1

GOLDEN2-LIKE1 IS SUFFICIENT BUT NOT NECESSARY FOR CHLOROPLAST BIOGENESIS IN MESOPHYLL CELLS OF C₄ GRASSES

Julia Lambret-Frotte[1,2], Georgia Smith[1] & Jane A. Langdale[1*]

[1] Department of Biology, University of Oxford, South Parks Road, OX1 3RB, Oxford, UK

[2] Current address: NIAB, Park Farm, Villa Road, Impington, CB24 9NZ, Cambridge, UK

[*] For correspondence: jane.langdale@biology.ox.ac.uk

ABSTRACT

Chloroplasts are the site of photosynthesis. In land plants, chloroplast biogenesis is regulated by a family of transcription factors named *GOLDEN2-like* (*GLK*). In C_4 grasses, it has been hypothesized that genome duplication events led to the sub-functionalization of *GLK* paralogs (*GLK1* and *GLK2*) to control chloroplast biogenesis in two distinct cell types: mesophyll and bundle sheath cells. Although previous characterization of *golden2* (*g2*) mutants in maize has demonstrated a role for *GLK2* paralogs in regulating chloroplast biogenesis in bundle sheath cells, the function of *GLK1* has remained elusive. Here we show that, contrary to expectations, *GLK1* is not required for chloroplast biogenesis in mesophyll cells of maize. Comparisons between maize and *Setaria viridis*, which represent two independent C_4 origins within the Poales, further show that the role of *GLK* paralogs in controlling chloroplast biogenesis in mesophyll and bundle sheath cells differs between species. Despite these differences, complementation analysis revealed that *GLK1* and *GLK2* genes from maize are both sufficient to restore functional chloroplast development in mesophyll and bundle sheath cells of *Setaria viridis* mutants. Collectively our results suggest an evolutionary trajectory in C_4 grasses whereby both orthologs retained the ability to trigger chloroplast biogenesis but *GLK2* adopted a more prominent developmental role, particularly in relation to chloroplast activation in bundle sheath cells.

KEY WORDS: C_4, chloroplasts, development, maize, green foxtail

INTRODUCTION

Chloroplasts are endosymbiotic organelles and are the site of photosynthetic reactions in a range of organisms from algae to land plants (Sagan 1967). Given the importance of chloroplast function for autotrophic growth, development of the organelle is a tightly controlled process that involves many levels of regulation (Waters and Langdale 2009). First described in the mid-1990s, the *golden2* (*g2*) mutant in *Zea mays* (maize) exhibits pale green leaves and impaired chloroplast development specifically in bundle sheath cells (Langdale and Kidner 1994, Hall *et al.* 1998). Subsequent characterization revealed that *ZmG2* encodes a transcription factor (Rossini *et al.* 2001), a founding member of the GARP transcription factor family (Riechmann *et al.* 2000). In most plant species examined, two *G2*-like (*GLK*) genes are present in the genome (Wang *et al.* 2013). *GLK* genes are defined by a characteristic helix-turn-helix DNA-binding domain related to MYB or MYB-like transcriptional factors, a conserved AREAEAA motif located immediately after the last helix and a GCT-box in the C-terminal region that is responsible for either homo- or heterodimerization (Rossini *et al.* 2001, Fitter *et al.* 2002, Safi *et al.* 2017). Functional characterization of *GLK* genes in maize and *Arabidopsis thaliana* (Arabidopsis) has shown that they are involved in the transcriptional activation of multiple photosynthesis-related genes including those encoding proteins of the light-harvesting complexes and enzymes of the chlorophyll biosynthesis pathway (Cribb *et al.* 2001, Waters *et al.* 2009). This pivotal role in chloroplast biogenesis has been associated with a myriad of physiological processes including stomatal movement, hormone signalling, defence responses and fruit ripening (Kobayashi *et al.* 2012, Nguyen *et al.* 2014, Li *et al.* 2020). The emerging picture is one where GLK proteins act as linchpins in any metabolic or developmental pathway that requires chloroplast function to be modulated.

During the evolution of land plants, the *GLK* gene family underwent multiple independent duplication events (Wang *et al.* 2013). In cases such as Arabidopsis where duplications likely occurred within species, the resulting paralogs are functionally redundant (Waters *et al.* 2009). However, duplication events in the monocots are more intriguing. The common ancestor of the Poales underwent a whole genome duplication that subdivided the *GLK* gene family into two sister clades: *GLK1* and *GLK2* (Wang *et al.* 2013). Notably, this event occurred before the many independent events that led to the evolution of C_4 photosynthesis within the Poales group (Christin and Osborne 2013). In species such as rice that perform C_3 photosynthesis, the *GLK* paralogs are both expressed in mesophyll cells and are functionally redundant as in Arabidopsis (Waters *et al.* 2009, Wang *et al.* 2013). Unlike C_3 photosynthesis, where carbon reactions occur predominantly in mesophyll cell chloroplasts, in C_4 plants carbon assimilation and fixation reactions are compartmentalized in the chloroplasts of mesophyll and bundle sheath cells, respectively (Langdale 2011). This biochemical compartmentalization is often accompanied by anatomical differences in chloroplast ultrastructure (Woo *et al.* 1970, Sage *et al.* 2011). In the C_4 species maize, *Setaria viridis* (setaria) and *Sorghum bicolor* (sorghum), *GLK1* is expressed preferentially in

mesophyll cells whereas *GLK2* transcripts accumulate preferentially in bundle sheath cells, with the

level of specificity differing between species such that *GLK2* transcripts accumulate to detectable levels

in mesophyll cells of setaria but not maize (Li *et al.* 2010, Chang *et al.* 2012, Wang *et al.* 2013, John *et*

al. 2014, Tausta *et al.* 2014). Importantly, localization of *ZmG2* transcripts in bundle sheath cells of

maize and defects in bundle sheath but not mesophyll chloroplast development in *g2* mutants

(Langdale and Kidner 1994, Cribb *et al.* 2001, Rossini *et al.* 2001) demonstrated that *GLK* paralogs in

C_4 grasses can control chloroplast development in a cell-specific manner.

GLK paralogs have been extensively characterized in Arabidopsis, however, little is known about gene

function in monocots, and in particular the function of *GLK1* genes has not been investigated. Because

ZmGLK1 is expressed in mesophyll cells of maize, and *ZmG2* is expressed and functions specifically

in bundle sheath cells, it has been hypothesized that *ZmGLK1* controls chloroplast biogenesis in

mesophyll cells (Tausta *et al.* 2014). Here we test this hypothesis by generating and characterizing loss-

of-function mutants in maize and setaria, two species that represent independent origins of C_4. Our

findings reveal that *ZmG2* and *SvGLK2* orthologs function in the cell-type(s) where the corresponding

transcripts accumulate but contrary to expectations, neither *ZmGLK1* nor *SvGLK1* are necessary for

chloroplast development in mesophyll cells.

RESULTS

***ZmGLK1* is not necessary for chloroplast biogenesis in mesophyll cells of maize**

To determine whether *ZmGLK1* regulates chloroplast biogenesis in maize, loss-of-function mutants

were generated using CRISPR/Cas9 (Figure 1A). Four independent mutant lines (referred to

collectively as *Zmglk1*) were obtained, all of which exhibited the same overall growth phenotype as

segregating wild-type siblings (Figure 1B, C; Figure S1). Ultrastructure analysis revealed that

chloroplasts in *Zmglk1* lines were similar to wild-type, with well-developed granal thylakoids in

mesophyll cells and agranal thylakoids in bundle sheath cells (Figure 1D-I; Figure S1). Total chlorophyll

levels were also similar in wild-type and *Zmglk1* mutants (Figure 1J). As such, we concluded that

ZmGLK1 is not necessary for thylakoid assembly or chlorophyll biosynthesis in bundle sheath or

mesophyll chloroplasts of maize.

An evaluation of chloroplast function in *Zmglk1* mutants was made by measuring photosystem II (PSII)

parameters. PSII is a large protein complex comprised of the antenna and chlorophyll molecules that

harvest light and together initiate the electron transfer reactions of photosynthesis (Nickelsen and

Rengstl 2013). In maize, PSII accumulates in the granal thylakoids of mesophyll cells and is largely

absent from the agranal bundle sheath cell chloroplasts (Woo *et al.* 1970). As such, PSII function is a

proxy for mesophyll cell chloroplast function. Fluorescence measurements were therefore taken in light

and dark-adapted plants to quantify PSII operating efficiency (PhiPSII) and maximum quantum

efficiency of PSII photochemistry (Fv/Fm), in both wild-type and *Zmglk1* mutant plants. Figures 1K-L show that PSII is operating normally in *Zmglk1* mutants, suggesting that *ZmGLK1* is not necessary for chloroplast biogenesis or function in maize mesophyll cells.

SvGLK1 is not necessary for chloroplast biogenesis in mesophyll cells of setaria

Maize and setaria have different origins of C_4 photosynthesis within the PACMAD clade (Li and Brutnell 2011) but like the orthologous *ZmGLK1*, *SvGLK1* transcripts accumulate preferentially in mesophyll cells (John *et al.* 2014). Phylogenetic analysis and amino acid sequence comparisons revealed that *SvGLK1* shares many structural features with *ZmGLK1* (Figure S2). To assess whether *SvGLK1* plays a role in chloroplast biogenesis, two independent loss-of-function mutants were generated in setaria using CRISPR/Cas9 (Figure 2A). As in maize, *Svglk1* mutants exhibited overall plant growth (Figure 2B-D) and chloroplast ultrastructure (Figure 2E-J) phenotypes that were similar to segregating wild-type siblings, with thylakoid membranes developing normally in both mesophyll and bundle sheath cells (Figure 2E-L). Furthermore, there was no significant difference in chlorophyll levels between *Svglk1* mutants and wild-type (Figure 2M), and electron flow through PSII was unperturbed (Figure 2N-O). Collectively, these results suggest that the preferential expression of *GLK1* genes in mesophyll cells of C_4 grass species does not reflect a requirement for gene function during chloroplast development and/or maintenance.

SvGLK2 regulates chloroplast development in both mesophyll and bundle sheath cells of setaria

Given that dimorphic chloroplast development is observed in bundle sheath and mesophyll cells of both maize and setaria, we next sought to determine whether *SvGLK2,* like its ortholog *ZmG2* (Figure S2)*,* regulates bundle sheath chloroplast development. To this end, CRISPR/Cas9 was used to generate loss-of-function *Svglk2* mutants (Figure 3A). In contrast to maize where *g2* mutants are weaker than wild-type but are viable and fertile, any *Svglk2* mutant plants that regenerated after transformation did not progress beyond the seedling stage before dying. The only viable line recovered from tissue culture was heterozygous for *SvGLK2* and this was used to isolate a null homozygous line via self-pollination. *Svglk2* 8G5 mutants, henceforth referred to as *Svglk2*, exhibited severe growth defects, had pale yellow leaves and failed to complete the life cycle (Figure 3B). Leaf cross sections showed misshapen and smaller chloroplasts in the bundle sheath cells of *Svglk2* mutants compared to wild-type (Figure 3C-D), and observations of chloroplast ultrastructure revealed impaired chloroplast anatomy with discontinuous thylakoids that occupied less of the organelle than in wild-type (Figure 3E-F), albeit a reduction that was not statistically significant (Figure 3I). More striking, given the bundle sheath specific perturbations observed in maize *g2* mutants, was the aberrant phenotype of mesophyll chloroplasts in *Svglk2* mutants. Compared to wild-type there was a significant reduction in thylakoid membrane coverage (15% less than wild-type) and a noticeable reduction in the size and frequency of grana (Figure 3G-H, J). As in bundle sheath cells, thylakoid membranes in mutant mesophyll chloroplasts

appeared fragmented (Figure 3H). The observed perturbations in bundle sheath and mesophyll cells
were accompanied by a reduction in leaf chlorophyll content of more than 80% (Figure 3K). These data
reveal a direct association between the presence of *SvGLK2* and *ZmG2* transcripts in a particular cell-
type and a requirement for gene function during chloroplast development, with the presence of *SvGLK2*
transcripts in mesophyll cells of setaria (albeit at lower levels than in bundle sheath cells) (John *et al.*
2014) reflecting a role in chloroplast biogenesis in both cell-types.

GLK2 transcript levels are not upregulated in *glk1* mutants of maize or setaria

Given the direct correspondence between *ZmG2* and *SvGLK2* transcript accumulation profiles and
necessity of gene function for chloroplast development, the absence of such correspondence for
ZmGLK1 and *SvGLK1* orthologs is intriguing. It is particularly notable because all *GLK* genes
characterized to date have a role in chloroplast biogenesis, albeit a redundant role alongside paralogs
in many cases (Fitter *et al.* 2002, Wang *et al.* 2013). Although *SvGLK2* may compensate for loss of
SvGLK1 function because *SvGLK2* transcripts comprise ~40% of total *GLK* transcript levels in
mesophyll cells of wild-type setaria, this is unlikely to be the case in maize where *ZmG2* transcripts only
comprise ~6% of total *GLK* transcript levels in mesophyll cells of wild-type plants (Figure S3). However,
loss of *ZmGLK1* function could lead to altered regulation of *ZmG2* and/or to the upregulation of genes
encoding the GATA transcription factors GNC and CGA1 which also regulate chloroplast development
(Chiang *et al.* 2012, Hudson *et al.* 2013) and have been shown to partially rescue loss of GLK function
in Arabidopsis (Zubo *et al.* 2018). To test this hypothesis, transcript levels of *ZmGLK1*, *ZmG2*, *ZmGNC*
and *ZmCGA1* were quantified in maize using the *Zmglk1* gene edited mutants generated above
alongside the previously reported transposon-induced *Zmg2-bsd1-s1* mutant (Cribb *et al.* 2001). In
each case the mutant allele produces transcripts that encode non-functional protein(s). Figure 4 reveals
that loss of *Zmglk1* function does not lead to consistently elevated levels of *ZmGLK1* (Figure 4A), *ZmG2*
(Figure 4B), *ZmGNC* (Figure 4C) or *ZmCGA1* (Figure 4D) transcripts. A small but significant increase
in *ZmG2*, *ZmGNC* and *ZmCGA1* transcript levels was seen in one of the four *Zmglk1* mutant lines but
that cannot explain how chloroplasts develop normally in all four lines. We thus conclude that
transcription of *ZmG2, ZmGNC* and *ZmCGA1* is not upregulated in mesophyll cells when *Zmglk1*
function is lost and thus that, unlike in all other contexts examined, chloroplasts in the mesophyll cells
of maize can develop normally in the absence of any GLK activity within that cell-type.

Whereas loss of ZmGLK1 function in maize does not impact on *ZmGLK1* or *ZmG2* transcript levels,
loss of ZmG2 function in bundle sheath cells leads to a statistically significant reduction in the
accumulation of both *ZmGLK1* and *ZmG2* transcripts (Figure 4A, B), and a similar situation is seen in
setaria where both *SvGLK1* and *SvGLK2* transcript levels are dramatically reduced in *Svglk2* mutants
(Figure 4E, F). This observation suggests that the ZmG2/SvGLK2 orthologs, either directly or indirectly,
promote the accumulation of both *GLK1* and *GLK2* transcripts in the respective species. In the case of

maize, the effect on *ZmGLK1* must be manifest non-cell-autonomously because *ZmG2* and *ZmGLK1*

transcripts accumulate in different cell-types. In setaria, however, because *SvGLK2* transcripts

accumulate in mesophyll cells, the effect on *SvGLK1* could be direct. Although reduced levels of

SvGLK1 transcripts in one of the two *Svglk1* mutant lines (Figure 4E) further suggest that *SvGLK1*

might also regulate its own transcript levels, the absence of such an effect in the other mutant line

makes it equally likely that the specific edits in line #2062 lead to transcript turnover, possibly via the

nonsense-mediated decay pathway. Collectively these data suggest that chloroplast development in

C$_4$ grasses is predominantly regulated by genes of the *GLK2* clade, with direct or indirect contributions

to mesophyll chloroplast biogenesis determined by the presence or absence of transcripts in that cell

type.

***ZmGLK1* and *ZmG2* are both sufficient to restore dimorphic chloroplast development in**

mesophyll and bundle sheath cells of *Svglk2* mutants

Our findings thus far show that *SvGLK2* and *ZmG2* play a more central role in controlling chloroplast

biogenesis in C$_4$ grasses than their respective *GLK1* paralogs. To evaluate whether this difference in

necessity also applies to sufficiency, *ZmGLK1* and *ZmG2* were expressed under the control of the

constitutive maize ubiquitin gene promoter (*ZmUBI*$_{pro}$) in the chloroplast-defective *Svglk2* mutant

background. Three independent lines were obtained for each construct and quantitative analysis of

transgene transcript levels revealed differences between lines that correlated directly with plant

phenotype (Figure 5A-D). For example, the weaker plants in line #17808-15254 accumulated around

half the level of *ZmGLK1* transcripts seen in healthier plants of the other two lines (Figure 5A, 5C).

Notably plants of all six lines were viable, healthy and fertile, suggesting complementation of the *Svglk2*

mutant phenotype (compare *Svglk2* plants in Figure 3B with Figure 5A, B).

Given that loss of *Svglk2* function leads to reduced levels of both *SvGLK1* and *SvGLK2* transcripts

(Figure 4E, F), we next sought to evaluate whether ZmGLK1 or ZmG2 could influence endogenous

SvGLK1 or *SvGLK2* expression. Figure 5E, F show that *SvGLK1* and *SvGLK2* transcripts were restored

to wild-type levels in *Svglk2* lines expressing either *ZmGLK1* or *ZmG2* transgenes. In fact, regardless

of the expression level of either *ZmGLK1* or *ZmG2*, all complemented lines exhibit the same wild-type-

like level of *SvGLK1* and *SvGLK2* expression (Figure 5C-F). Both ZmGLK1 and ZmG2 can thus activate

expression of *SvGLK1* and *SvGLK2*, providing further support for the existence of a regulatory network

between *GLK* genes. The functional significance of this observation must be considered in the context

of transgene expression driven by a constitutive promoter and gene-edited *Svglk2* transcripts encoding

non-functional proteins (and hence any effect on *SvGLK1* expression is a direct effect of the maize

transgene and not via *SvGLK2*). In this regard, we must deduce that the transgenic lines expressing

ZmGLK1 accumulate transcripts encoding functional ZmGLK1 and SvGLK1 proteins in both mesophyll

and bundle sheath cells, and those expressing *ZmG2* accumulate transcripts encoding functional
ZmG2 and SvGLK1 proteins, similarly in both cell types.
232
To determine whether *ZmGLK1* and/or *ZmG2* can rescue chloroplast ultrastructure defects in
mesophyll and/or bundle sheath cells of *Svglk2* mutants, confocal images and transmission electron
micrographs of transverse leaf sections from all six transgenic lines were examined. In all cases,
chloroplasts were restored to normal size (Figure 6A-C, Figure S4), with predominantly agranal
thylakoids in bundle sheath cell chloroplasts (Figure 6D-F, Figure S4) and granal thylakoids in
mesophyll chloroplasts (Figure 6G-I, Figure S4). Quantification revealed that with the exception of the
one line that had low levels of transgene expression (#17808-15254), thylakoid occupancy levels in
mesophyll (Figure 6J) and bundle sheath (Figure 6K) chloroplasts of transgenic lines were statistically
equivalent (or in one case higher) than in wild-type plants. Consistent with ectopic expression of *GLK*
genes, mature chloroplasts were also observed in the leaf vasculature of all complemented lines, more
with *ZmG2* than *ZmGLK1* (Figure 6A-C, Figure S4). Irrespective of which maize ortholog was
overexpressed, the total chlorophyll content in leaves of transgenic plants was restored to wild-type
levels (Figure 6L). Notwithstanding the fact that SvGLK1 may be functional in bundle sheath and
mesophyll cells of all transgenic plants, collectively these data demonstrate that ZmGLK1 and ZmG2
are functionally equivalent in the context of the C_4 setaria leaf and that the dimorphic identity of bundle
sheath and mesophyll cell chloroplasts develops regardless of which maize ortholog is overexpressed.
249

DISCUSSION

Prior to this study it was known that *GLK* genes are duplicated in many land plant species and that in
species that carry out C_3 photosynthesis the duplicate genes act redundantly to regulate chloroplast
biogenesis in all photosynthetic cells (Waters *et al.* 2008, Bravo-Garcia *et al.* 2009, Wang *et al.* 2013).
It had also been hypothesized that *GLK1* and *GLK2* genes regulate dimorphic chloroplast development
in the two photosynthetic cell-types of C_4 species, with each gene functioning in one of the two cell-
types (Rossini *et al.* 2001). This hypothesis was based on the observation that *GLK1* genes are
preferentially expressed in mesophyll cells of maize, sorghum and setaria whereas *GLK2/G2* genes are
preferentially expressed in bundle sheath cells (Li *et al.* 2010, Wang *et al.* 2013, John *et al.* 2014), and
loss of *Zmg2* function in maize perturbs bundle sheath but not mesophyll chloroplast development
(Cribb *et al.* 2001). Through the phenotypic characterization of *glk1* loss-of-function mutants in maize
and setaria (Figures 1, 2 & 4), *glk2* loss-of-function mutants in setaria (Figures 3 & 4), and *glk2* setaria
mutants complemented with maize *GLK1* and *GLK2* genes (Figure 5 & 6), we have revealed a more
complex situation both with respect to the developmental processes operating within individual C_4
species and to likely evolutionary trajectories.
265
266

GLK paralogs have equivalent functional capacity

GLK genes duplicated in a common ancestor of the Poales, before the speciation of maize, sorghum, setaria and rice, and before the evolution of C_4 photosynthesis (Wang *et al.* 2013). Given that *GLK* genes regulate chloroplast biogenesis in the moss *Physcomitrium patens* (Yasumura *et al.* 2005, Bravo-Garcia *et al.* 2009), which last shared a common ancestor with flowering plants over 450 million years ago (Kenrick and Crane 1997, Wickett *et al.* 2014), it is reasonable to assume that prior to this duplication the ancestral *GLK* gene played a role in chloroplast development. The duplication event, which must have occurred between 85-95 million years ago (Gallaher *et al.* 2022), provided an opportunity for neo- and/or sub-functionalization of gene activity and thus for divergent roles to evolve in different grass species. It has previously been suggested that the compartmentalized expression of *GLK1* and *GLK2* genes in mesophyll and bundle sheath cells of C_4 species reflects distinct roles in the biogenesis of granal versus agranal chloroplasts. This hypothesis was supported by the observation that *GLK* homologs are not differentially expressed in *Gynandropsis gynandra* (formerly known as *Cleome gynandra*), a C_4 dicot that exhibits granal thylakoids in chloroplasts of both mesophyll and bundle sheath cells (Marshall *et al.* 2007, Wang *et al.* 2013). However, data presented here show that appropriate cell-type specific thylakoid stacking is restored in mesophyll and bundle sheath cells of *Svglk2* mutants when either *ZmGLK1* or *ZmG2* is constitutively expressed (Figures 5 & 6). These results demonstrate that the maize paralogs have equivalent functional capacity. Furthermore, the data are consistent with a recent report which showed that the identity of genes activated by GLK transcription factors is determined by *cis* regulatory factors, such that a heterologous GLK protein will activate downstream targets that are normally activated by endogenous GLK proteins in the host species as opposed to targets that are activated by the heterologous protein in its species of origin (Tu *et al.* 2022). Together these observations suggest that the evolution of dimorphic chloroplasts in C_4 species was unlikely to have been mediated by sub-functionalization of GLK protein activity.

Given that *GLK* gene paralogs encode functionally equivalent proteins both in C_3 species where they act redundantly and in the C_4 species maize, the question arises of why the paralogs are expressed in a cell-type preferential manner in C_4 grass species (at least in the ones that have been examined). Estimates based on the topology of BEP (C_3 species only) and PACMAD (C_3 & C_4 species) clades in the Poales suggest that maize diverged from rice ~80 million years ago (Gallaher *et al.* 2022) and that within the PACMAD clade, maize and sorghum share a common history of C_4 evolution whereas independent events occurred in maize and setaria around ~45 million years apart (Gallaher *et al.* 2022). Both *GLK* gene paralogs are predominantly expressed in mesophyll cells of C_3 rice leaves, with *GLK1* transcripts accumulating at higher levels than *GLK2* (Hua *et al.* 2021), and gene function being redundant (Wang *et al.* 2013). Notably, both transcripts can also be detected in bundle sheath cells, at levels roughly equivalent to those of *GLK2* in mesophyll cells, suggesting that the few chloroplasts that develop in rice bundle sheath cells are also regulated by the redundant action of both genes. By

contrast, an element of sub-functionalization is apparent in both maize and setaria, with the *GLK2* paralog expressed preferentially in bundle sheath cells, almost exclusively so in maize. Given that the evolution of C_4 required activation of photosynthesis and a significant increase in chloroplast volume in bundle sheath cells, selection on the promoter of the *GLK2* ortholog to enable high levels of expression in that cell type may have been a critical step. Indeed, whether due to selection or drift, a phylogenetic signal of variation is evident between C_3 and C_4 species in the *GLK2* but not the *GLK1* promoter sequence (Figure S3). In this scenario, older C_4 lineages in the Poales would display more specific localization of *GLK2* transcripts in bundle sheath cells than younger lineages, a situation that is evident in comparisons between maize and setaria (Figure S3). However data from a number of different C_3, C_4 and intermediate species (Washburn *et al.* 2021) (Figure S3) invoke a more complex scenario because preferential expression of *GLK2* paralogs in bundle sheath cells is evident in species where decarboxylation reactions occur in the bundle sheath chloroplasts (NADP-ME type) but not in species where decarboxylation occurs in the mitochondria (NAD-ME type) or in the cytoplasm (PEP-CK type). This suggests that preferential localization of GLK2 in bundle sheath cells is associated with specific aspects of chloroplast function as opposed to general activation of photosynthesis.

Despite equivalent functional capacity, GLK2 plays a dominant role in both maize and setaria
The difference in levels of bundle sheath specificity of *GLK2* transcripts in maize versus setaria can partially explain the different phenotypes observed in loss-of-function mutants. In maize, where *ZmG2* is almost exclusively expressed in bundle sheath cells, *Zmg2* mutants exhibit perturbed chloroplast development only in that cell-type. By contrast, expression of *GLK2* in both mesophyll and bundle sheath cells of setaria is associated with defective chloroplast development in both cell-types (Figure 3). This observation was initially counter-intuitive because it would be expected that, as in C_3 plants, the high levels of *SvGLK1* transcripts in mesophyll cells would compensate for loss of SvGLK2 function. However, we observed that *SvGLK1* transcript accumulation was virtually abolished in *Svglk2* mutants and levels of the non-functional *Svglk2* transcripts were also reduced (Figure 4). As such, we conclude that SvGLK2 activates expression (or transcript stabilization) both of itself and of *SvGLK1,* and that the *Svglk2* mutant phenotype reflects loss of function of both genes. Collectively, these data suggest that SvGLK2 is the primary regulator of chloroplast development in both bundle sheath and mesophyll cells of setaria.

Whereas normal mesophyll chloroplast development in the absence of GLK1 function can be explained in setaria because SvGLK2 likely compensates for loss of GLK1 function, the mutant phenotype in maize was unexpected. In every land plant species examined to date, a role for *GLK* genes in chloroplast development has been revealed, albeit often a redundant role with paralogs operating in the same cell-type. Because ZmGLK1 is highly expressed in maize mesophyll cells and no other GLK-like transcripts appear as possible candidates for a redundant paralog, a previously unknown

mechanism must be operating in this cell-type. Although GLK gene function has previously been shown
to be cell-autonomous in Arabidopsis, it is possible that a non-cell-autonomous mechanism could
operate in maize. Indeed, there is some evidence of this because *ZmGLK1* transcript levels are reduced
when G2 function is lost in the bundle sheath (Figure 4). In such a scenario, loss of GLK1 function in
mesophyll cells would be compensated for by direct or indirect activity of G2 proteins in the bundle
sheath cells. Any such effect would only be revealed in double *Zmglk1;g2* mutants, which would have
defective chloroplasts in both cell-types. Regardless of whether ZmG2 compensates for loss of
ZmGLK1 function, the GLK2 paralog plays the dominant role in chloroplast biogenesis in maize, as in
setaria.
350

METHODS
Plant growth conditions
Maize and *S. viridis* seeds were germinated in hydrated vermiculite or on paper, respectively, and
placed in an Aralab Fitoclima D1200 PLH incubator under 12 h light/12 h dark cycle at an intensity of
320 µmol photon m^{-2} s^{-1}, with temperatures ranging from 31 °C during the day to 22 °C at night and
humidity between 50-60 %. About 7-10 days after sowing (DAS), seedlings were transferred to pots
containing John Innes no. 2 soil mixture and irrigated with water to keep the soil moist. Plants were
kept under these conditions until sampled for experimental analysis.
359

360 For genetic crosses, maize plants were transferred to larger pots with the same soil mixture
361 supplemented with Osmocote slow-release granules. Plants were grown in a greenhouse at the
362 Department of Biology at the University of Oxford, UK with 16 h light/8 h dark cycle, daytime
363 temperature 28 °C and night-time temperature 20 °C, with supplemental light provided when natural
364 light levels were below 120 µmol photon m^{-2} s^{-1}.
365

Construct design and assembly
ZmG2 and *ZmGLK1* orthologs were identified in *S. viridis* via phylogenetic reconstruction of the GARP
gene family (see Figure S2). Guide RNAs (gRNA) to drive CRISPR/Cas9 site-directed mutagenesis
were designed using the CRISPOR (Concordet and Haeussler 2018) online tool
(http://crispor.tefor.net/) to achieve high editing efficiency for the target genes *ZmGLK1*
(GRMZM2G026833), *SvGLK1* (Sevir.4G123200) and *SvGLK2* (Sevir.5G003100), and to minimize off-
targets. For maize, the designed gRNAs were sent to the Wisconsin Crop Innovation Centre (University
of Wisconsin) to be cloned and transformed using their standard pipeline for maize transformation. For
S. viridis, the gRNA sequences were synthesized as oligonucleotides with appropriate flanking sites for
Golden Gate cloning (Table S1; JLF_Ox43 and JLF_Ox44 for *SvGLK1*; JLF_Ox51 and JLF_Ox52 for
SvGLK2). Prior to cloning, the oligonucleotides were annealed and phosphorylated by incubating 10
µM of each with 1x Ligase Buffer (New England Biolabs) and 10 U of T4 Polynucleotide Kinase (New

England Biolabs) in a final reaction volume of 15 µl at 37°C for 1.5 hours. After that period, 2 µl of 0.5 M NaCl was added, and the reaction was placed in a water bath heated to 95 °C for 2 min. The reaction was cooled to room temperature before being diluted 100-fold and cloned downstream of the *TaU6* promoter (module pJG310) using the Golden Gate one-step one-pot protocol (Engler *et al.* 2009) with an appropriate restriction enzyme (*Esp3I*). This module was then combined with another level 1 module containing *TaCas9* downstream of the maize ubiquitin promoter (pJG471) into the final acceptor and binary vector pTC278. Both modules and the vector were kindly provided by Dr Dan Voytas (University of Minnesota).

To overexpress *ZmGLK1* and *ZmG2*, the coding sequences of the maize genes were domesticated to avoid recognition sites for the type II restriction enzymes used in Golden Gate assembly: *BsaI*, *BpiI*, *Esp3I* and *DraIII*. The final sequence was synthesized as a level 0 module with the appropriate SC flanking sites (Engler and Marillonnet 2013). *ZmGLK1* and *ZmG2* level 0 modules (EC17054 and EC17056) were cloned downstream of the maize ubiquitin promoter (*ZmUBI*$_{pro}$, EC15455) and upstream of a *nos* terminator (tNOS, EC41421) in level 1 backbone position forward 2 using the Golden Gate one-step one-pot protocol (Engler *et al.* 2009). The previously described *hygromycin phosphotransferase* (HygR) coding sequence cloned downstream of the rice actin promoter (*OsACT*$_{pro}$) was used as the selectable marker (Vlad *et al.* 2019). Level 1 modules were assembled into the binary vector pAGM4723 to obtain the constructs EC17808 and EC17810 (hereafter referred to as 17808 and 17810, respectively) depicted in Figure S4A and S4B.

Plant transformation

S. viridis accessions A10 and ME034V were used to obtain CRISPR/Cas9-driven mutations for *SvGLK1* and *SvGLK2*, respectively. *Agrobacterium tumefaciens* strain AGL1 harbouring the construct of interest was co-cultivated with callus generated from seeds of *S. viridis*. Callus induction, transformation and seedling regeneration were performed as described by VanEck and collaborators (Van Eck and Startwood 2015). Complementation lines were obtained by transforming constructs 17808 and 17810 into callus generated from a population of seeds that segregated for the gene-edited *Svglk2* allele but no longer segregated the original t-DNA which had been recombined out. The presence of t-DNA in regenerated seedlings was confirmed via duplex PCR (Table S1; JLF_Ox12 and JLF_Ox13 amplify the HygR gene; JLF_Ox14 and JLF_Ox15 amplify an endogenous gene used as a DNA quality control). The occurrence of gene editing by CRISPR/Cas9 was verified by PCR amplification of the region flanking the expected mutation site (Table S1; JLF_Ox45 and JLF_Ox46 for *SvGLK1*; JLF_Ox53 and JLF_Ox54 for *SvGLK2*) followed by endonuclease digestion with *MmeI* or *Tsp45I* for *SvGLK1* and *SvGLK2*, respectively, following the manufacturers' guidelines.

Z. mays Hi-II was transformed by the Wisconsin Crop Innovation Centre (University of Wisconsin)

according to their transformation pipeline. The presence of the t-DNA was assessed in T1 lines by PCR

(Table S1; JLF_Ox178 and JLF_Ox179) and *Zmglk1* mutants were identified by PCR amplification of

the region flanking the expected mutation site (Table S1; JLF_Ox39 and JLF_Ox175) followed by

endonuclease digestion with *HinfI* according to the manufacturer's instructions.

For all lines, mutated sequences were verified by Sanger sequencing (Source BioScience, UK).

Experimental analyses were performed on homozygous plants in the T2 or subsequent generations.

Confocal imaging

The mid-portion of fully expanded leaf 7 from 21 DAS *S. viridis* plants was fixed in 4 %

paraformaldehyde and 0.2 % glutaraldehyde in 25 mM phosphate buffer pH 7.2. After fixation, leaves

were stored in 0.2 M Na-EDTA pH 9.0. Free-hand cross sections were imaged using a Leica SP9

confocal microscope with an Argon laser set to excite chlorophyll at 663 nm wavelength and to emit

from 650-800 nm.

Ultrastructure analysis via transmission electron microscopy

Sample fixation and preparation were performed by the Electron Microscopy Facility team at the Sir

Willian Dunn School of Pathology (University of Oxford). Leaf discs of 2 mm diameter were collected

from leaf 5 of 40 DAS maize or 25 DAS *S. viridis* plants and placed into 3 % glutaraldehyde in 0.06 M

Sorenson's phosphate buffer pH 7.2. Samples were transferred to a Leica AMW microwave tissue

processor for fixation, osmification (with 1 % OsO_4), dehydration with an acetone series, and resin

infiltration (LRW Hard). The samples were then removed from the microwave and transferred to fresh

resin for four days, renewing the resin twice a day. The embedded samples were transferred into gelatin

capsules and filled with resin for polymerization at 60 °C for about 65 h. Sections of 150 nm were cut

using a Leica UC7 ultramicrotome, collected onto formvar-coated copper slot grids and post-stained

with lead citrate. Images were acquired on a Thermo Fischer Tecnai T12 TEM with a Gatan OneView

CMOS camera.

Thylakoid occupancy of chloroplasts was quantified using ImageJ. The thylakoid area was measured

by adjusting the image threshold to create a binary mask on which only the thylakoid membrane was

apparent. This area value was divided by the chloroplast planar area to give the thylakoid occupancy

level.

Chlorophyll extraction and quantification

Leaf discs were harvested from fully expanded leaf 5 at 30 DAS for maize and 21 DAS for *S. viridis*,

and immediately frozen in liquid nitrogen. For total chlorophyll extraction, finely powdered tissue was

resuspended in 80 % acetone buffered in 100 mM Tris-HCl pH 8.0 (Chazaux *et al.* 2022). Following a centrifugation step, the absorbance of each sample was read in technical triplicates in a FLUOstar Omega spectrometer (BMG Labtech) at 646 and 663 nm. Background absorbance was measured at 750 nm. Absorbance values recorded at 750 nm were subtracted from values recorded at 646 and 663 nm for normalization purposes. Technical replicates were averaged for each sample. Concentration of chlorophylls a + b were calculated in micrograms (μg) per leaf area using the following the equation: $(19.54 \times Abs_{646} + 8.29 \times Abs_{663}) \times$ dilution factor (Porra *et al.* 1989).

Fluorescence quantification of PSII

Photosystem II (PSII) fluorescence was measured using a LiCOR Li-6800 portable photosynthesis system. LiCOR chamber conditions were set to a flow intensity of 500 umol s^{-1}, delta pressure of 0.2 kPa, relative air humidity of 50 %, reference concentration of CO_2 at 400 umol mol^{-1} and temperature exchange at 25 °C. For quantification of PSII operating efficiency (PhiPSII), plants were adapted to light under the conditions described above for at least 1.5 h and actinic light intensity on the LiCOR chamber was set to 1000 μmol photon m^{-2} s^{-1}. After clamping, each leaf was adapted for a few minutes until stabilization was reached before logging the fluorescence value. Quantification of maximum quantum efficiency of PSII photochemistry (Fv/Fm) was performed on plants that were dark-adapted for at least 16 h. LiCOR actinic lights were off. In both cases, the fluorescence action log was set to FoFm (dark), FsFm' (light), rectangular flash type with red target at 8000 μmol photon m^{-2} s^{-1}, duration of 1000 ms, output rate of 100 Hz and margin of 5 points.

Gene expression analysis via qPCR

The mid-portion of fully expanded leaf 5 from at least three biological replicates of 30 DAS maize or 21 DAS *S. viridis* were harvested and immediately frozen in liquid nitrogen. Tissue was ground prior to total RNA extraction using a RNeasy Plant Mini Kit (Qiagen) according to the manufacturer's instructions. To avoid any DNA contamination, samples were treated with TURBO DNA-free DNase (Thermo Fischer Scientific) according to the manufacturer's guidelines. For first strand cDNA synthesis, 100 ng of total RNA was added to 1 μl of Maxima Enzyme Mix (Thermo Fischer Scientific) and 1x Reaction Mix to a final volume of 10 μL. The mixture was incubated at 25 °C for 10 min, followed by 50 °C for 15 min. Enzyme inactivation was carried out at 85 °C for 5 min.

Amplification reactions were performed in a StepOnePlus Real-Time PCR System (Thermo Fischer Scientific) using SYBRGreen to monitor dsDNA synthesis. The reaction mixtures contained 5 μL of diluted cDNA (1:25), 0.2 μM of each primer and 10 μL of SYBR™ Green PCR Master Mix (Thermo Fischer Scientific) in a total volume of 10 μL. The reaction cycles began with a 5 min denaturation step at 94 °C, followed by 40 amplification cycles of 15 s at 94 °C, 10 s at 60 °C, 15 s at 72 °C. After each cycle, the fluorescence was measured at 60 °C for 35 s. The melting curve was produced by a cycle of

95 °C for 15 s, 60 °C for 1 min, 95 °C for 30 s and finally 60 °C for 15 s. Every qPCR reaction was repeated three times to make technical replicates.

Transcript levels for multiple target genes were measured with primers designed to specifically amplify the gene of interest (Table S1). Two reference genes were used to normalize gene expression for maize (Lin *et al.* 2014) (Table S1; ZmCYP_F and ZmCYP_R, ZmEF1α_F and ZmEF1α_R) and *S. viridis* (Lambret-Frotté *et al.* 2015) (Table S1; JLF_Ox26 and JLF_Ox27, JLF_Ox28 and JLF_Ox29). Exported *Rn* values were used to calculate amplification efficiency for each primer pair and the Cq values for each qPCR reaction using the '*qpcR*' package from R (Ritz and Spiess 2008). Normalized relative expression of the target genes was calculated using the R package '*EasyqpcR*' (https://www.bioconductor.org/packages//2.12/bioc/html/EasyqpcR.html), according to the ΔΔCq model (Pfaffl 2001) based on the previously calculated Cq and amplification efficiencies.

Statistical analysis

Datasets were tested for Normal distributions using the Shapiro-Wilk test (Shapiro and Wilk 1965). For thylakoid coverage, chlorophyll concentration and PSII fluorescence, variance was calculated using a one-way ANOVA followed by a pairwise comparison analysis with Tukey's HSD test (Tukey 1949). For relative gene expression, statistical significance was calculated based on a Student's t-test, comparing the mean variance of each mutant line with their respective controls. All statistical analyses were performed using *scipy* v.1.5.0 library and plots were prepared using *matplotlib* v3.5.1 library; both from Python v.3.7.6, using Visual Studio Code as a code editor.

LIST OF SUPPLEMENTARY MATERIAL

Table S1. Oligonucleotides used for cloning, genotyping and qPCR.

Figure S1. Phenotypic characterization of three independent *Zmglk1* mutant lines.

Figure S2. Comparison of *GLK* sequences.

Figure S3. *GLK* transcript accumulation in bundle sheath and mesophyll cells.

Figure S4. Phenotypic characterization of two independent lines of *Svglk2* complemented with *ZmGLK1* or *ZmG2*.

AUTHOR CONTRIBUTIONS

J.L-F and J.A.L conceived and designed the experiments; G.S carried out setaria transformation, plant cultivation and genotyping; J.L-F carried out all of the remaining experiments and analysed the data. J.L-F and J.A.L wrote the manuscript.

ACKNOWLEDGEMENTS

The authors thank Dan Voytas for making available the CRISPR/Cas9 cloning vectors for gene editing in setaria; Raman Dhaliwal, Charlotte Melia and Errin Jonson for processing and imaging the TEM samples; Ray Collier, Mike Pettersen and the Wisconsin Crop Innovation Centre team for maize transformation; John Baker for plant photography; Roxaana Clayton, Julie Bull and Lizzie Jamison for technical support; Emma Raven for assisting with initial characterization of mutant lines; Jacob Washburn for providing TPM data for Figure S3D; Sophie Johnson, Chiara Perico, Daniela Vlad, Sovanna Tan, Thomas Hughes and Maricris Zaidem for discussion throughout the experimental work and during manuscript preparation.

COMPETING INTERESTS

No competing interests are declared.

FUNDING

This research was funded by a BBSRC sLoLa grant (BB/P003117/1), a Newton International Fellowship from The Royal Society to J.L-F (NF171598, 2018 – 2020) and by the Bill and Melinda Gates Foundation C_4 Rice grant awarded to the University of Oxford (2015-2019; OPP1129902).

REFERENCES

Bravo-Garcia, A., Yasumura, Y., and Langdale, J.A., 2009. Specialization of the Golden2-like regulatory pathway during land plant evolution. *New Phytologist*, 183 (1), 133–141.

Chang, Y.-M., Liu, W.-Y., Shih, A.C.-C., Shen, M.-N., Lu, C.-H., Lu, M.-Y.J., Yang, H.-W., Wang, T.-Y., Chen, S.C.-C., Chen, S.M., Li, W.-H., and Ku, M.S.B., 2012. Characterizing Regulatory and Functional Differentiation between Maize Mesophyll and Bundle Sheath Cells by Transcriptomic Analysis. *Plant Physiology*, 160 (1), 165–177.

Chazaux, M., Schiphorst, C., Lazzari, G., and Caffarri, S., 2022. Precise estimation of chlorophyll a, b and carotenoid content by deconvolution of the absorption spectrum and new simultaneous equations for Chl determination. *Plant Journal*, 109 (6), 1630–1648.

Chiang, Y.-H., Zubo, Y.O., Tapken, W., Kim, H.J., Lavanway, A.M., Howard, L., Pilon, M., Kieber, J.J., and Schaller, G.E., 2012. Functional Characterization of the GATA Transcription Factors GNC and CGA1 Reveals Their Key Role in Chloroplast Development, Growth, and Division in Arabidopsis. *Plant Physiology*, 160 (1), 332–348.

Christin, P.-A. and Osborne, C.P., 2013. The recurrent assembly of C4 photosynthesis, an evolutionary tale. *Photosynthesis research*, 117 (1–3), 163–75.

Concordet, J.-P. and Haeussler, M., 2018. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Research*, 46 (W1), W242–W245.

Cribb, L., Hall, L., and Langdale, J., 2001. Four Mutant Alleles Elucidate the Role of the G2 Protein in

the Development of C4 and C3 Photosynthesizing Maize Tissues. *Genetics*, 159, 787–797.

Van Eck, J. and Startwood, K., 2015. Setaria viridis. *In*: K. Wang, ed. *Agrobacterium Protocols*. Springer,
417.

Engler, C., Gruetzner, R., Kandzia, R., and Marillonnet, S., 2009. Golden gate shuffling: a one-pot DNA
shuffling method based on type IIs restriction enzymes. *PloS one*, 4 (5), e5553.

Engler, C. and Marillonnet, S., 2013. Combinatorial DNA assembly using golden gate cloning. *Methods*
in Molecular Biology, 1073, 141–156.

Fitter, D.W., Martin, D.J., Copley, M.J., Scotland, R.W., and Langdale, J.A., 2002. GLK gene pairs
regulate chloroplast development in diverse plant species. *Plant Journal*, 31 (6), 713–727.

Gallaher, T.J., Peterson, P.M., Soreng, R.J., Zuloaga, F.O., Li, D.-Z., Clark, L.G., Tyrrell, C.D., Welker,
C.A.D., Kellogg, E.A., and Teisher, J.K., 2022. Grasses through space and time: An overview of the
biogeographical and macroevolutionary history of Poaceae. *Journal of Systematics and Evolution*,
60 (3), 522–569.

Hall, L.N., Rossini, L., Cribb, L., and Langdale, J.A., 1998. GOLDEN 2: A novel transcriptional regulator
of cellular differentiation in the maize leaf. *Plant Cell*, 10 (6), 925–936.

Hua, L., Stevenson, S.R., Reyna-Llorens, I., Xiong, H., Kopriva, S., and Hibberd, J.M., 2021. The bundle
sheath of rice is conditioned to play an active role in water transport as well as sulfur assimilation
and jasmonic acid synthesis. *Plant Journal*, 107 (1), 268–286.

Hudson, D., Guevara, D.R., Hand, A.J., Xu, Z., Hao, L., Chen, X., Zhu, T., Bi, Y.-M., and Rothstein, S.J.,
2013. Rice Cytokinin GATA Transcription Factor1 Regulates Chloroplast Development and Plant
Architecture. *Plant Physiology*, 162 (1), 132–144.

John, C.R., Smith-Unna, R.D., Woodfield, H., Covshoff, S., and Hibberd, J.M., 2014. Evolutionary
convergence of cell-specific gene expression in independent lineages of C4 grasses. *Plant*
Physiology, 165 (1), 62–75.

Kenrick, P. and Crane, P.R., 1997. The origin and early evolution of plants on land. *Nature*, 389 (6646),
33–39.

Kobayashi, K., Baba, S., Obayashi, T., Sato, M., Toyooka, K., Keränen, M., Aro, E.M., Fukaki, H., Ohta,
H., Sugimoto, K., and Masudaa, T., 2012. Regulation of root greening by light and auxin/cytokinin
signaling in Arabidopsis. *Plant Cell*, 24 (3), 1081–1095.

Lambret-Frotté, J., Almeida, L.C.S. De, Moura, S.M. De, and Souza, F.L.F., 2015. Validating Internal
Control Genes for the Accurate Normalization of qPCR Expression Analysis of the Novel Model
Plant Setaria viridis, 1–22.

Langdale, J.A., 2011. C4 cycles: past, present, and future research on C4 photosynthesis. *The Plant*
Cell, 23 (11), 3879–92.

Langdale, J.A. and Kidner, C.A., 1994. Bundle sheath defective, a mutation that disrupts cellular
differentiation in maize leaves. *Development*, 120 (3), 673–681.

Li, P. and Brutnell, T.P., 2011. Setaria viridis and Setaria italica, model genetic systems for the Panicoid

grasses. *Journal of Experimental Botany*, 62 (9), 3031–7.

Li, P., Ponnala, L., Gandotra, N., Wang, L., Si, Y., Tausta, S.L., Kebrom, T.H., Provart, N., Patel, R., Myers, C.R., Reidel, E.J., Turgeon, R., Liu, P., Sun, Q., Nelson, T., and Brutnell, T.P., 2010. The developmental dynamics of the maize leaf transcriptome. *Nature Genetics*, 42 (12), 1060–7.

Li, X., Wang, P., Li, J., Wei, S., Yan, Y., Yang, J., Zhao, M., Langdale, J.A., and Zhou, W., 2020. Maize GOLDEN2-LIKE genes enhance biomass and grain yields in rice by improving photosynthesis and reducing photoinhibition. *Communications Biology*, 3 (1), 151.

Lin, Y., Zhang, C., Lan, H., Gao, S., Liu, H., Liu, J., Cao, M., Pan, G., Rong, T., and Zhang, S., 2014. Validation of potential reference genes for qPCR in maize across abiotic stresses, hormone treatments, and tissue types. *PloS one*, 9 (5), e95445.

Marshall, D.M., Muhaidat, R., Brown, N.J., Liu, Z., Stanley, S., Griffiths, H., Sage, R.F., and Hibberd, J.M., 2007. Cleome, a genus closely related to Arabidopsis, contains species spanning a developmental progression from C3 to C4 photosynthesis. *Plant Journal*, 51 (5), 886–896.

Nguyen, C. V., Vrebalov, J.T., Gapper, N.E., Zheng, Y., Zhong, S., Fei, Z., and Giovannoni, J.J., 2014. Tomato GOLDEN2-LIKE transcription factors reveal molecular gradients that function during fruit development and ripening. *Plant Cell*, 26 (2), 585–601.

Nickelsen, J. and Rengstl, B., 2013. Photosystem II Assembly: From Cyanobacteria to Plants. *Annual Review of Plant Biology*, 64 (1), 609–635.

Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research*, 29 (9), e45.

Porra, R.J., Thompson, W.A., and Kriedemann, P.E., 1989. Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 975 (3), 384–394.

Riechmann, J.L., Heard, J., Martin, G., Reuber, L., Jiang, C., Keddie, J., Adam, L., Pineda, O., Ratcliffe, O.J., Samaha, R.R., Creelman, R., Pilgrim, M., Broun, P., Zhang, J.Z., Ghandehari, D., Sherman, B.K., and Yu, G., 2000. Arabidopsis transcription factors: genome-wide comparative analysis among eukaryotes. *Science (New York, N.Y.)*, 290 (5499), 2105–10.

Ritz, C. and Spiess, A.-N., 2008. qpcR: an R package for sigmoidal model selection in quantitative real-time polymerase chain reaction analysis. *Bioinformatics*, 24 (13), 1549–1551.

Rossini, L., Cribb, L., Martin, D.J., and Langdale, J.A., 2001. The maize Golden2 gene defines a novel class of transcriptional regulators in plants. *Plant Cell*, 13 (5), 1231–1244.

Safi, A., Medici, A., Szponarski, W., Ruffel, S., Lacombe, B., and Krouk, G., 2017. The world according to GARP transcription factors. *Current Opinion in Plant Biology*, 39, 159–167.

Sagan, L., 1967. On the origin of mitosing cells. *Journal of Theoretical Biology*, 14 (3), 225-IN6.

Sage, R.F., Christin, P.-A., and Edwards, E.J., 2011. The C(4) plant lineages of planet Earth. *Journal of Experimental Botany*, 62 (9), 3155–69.

Shapiro, S.S. and Wilk, M.B., 1965. An analysis of variance test for normality (complete samples).
Biometrika, 52 (3–4), 591–611.

Tausta, L.S., Li, P., Si, Y., Gandotra, N., Liu, P., Sun, Q., Brutnell, T.P., and Nelson, T., 2014.
Developmental dynamics of Kranz cell transcriptional specificity in maize leaf reveals early onset
of C4-related processes. *Journal of Experimental Botany*, 65 (13), 3543–3555.

Tu, X., Ren, S., Shen, W., Li, J., Li, Y., Li, C., Li, Y., Zong, Z., Xie, W., Grierson, D., Fei, Z., Giovannoni,
J., Li, P., and Zhong, S., 2022. Limited conservation in cross-species comparison of GLK
transcription factor binding suggested wide-spread cistrome divergence. *Nature
Communications*, 13 (1), 7632.

Tukey, J.W., 1949. Comparing Individual Means in the Analysis of Variance. *Biometrics*, 5 (2), 99–114.

Vlad, D., Abu-Jamous, B., Wang, P., and Langdale, J.A., 2019. A modular steroid-inducible gene
expression system for use in rice. *BMC Plant Biology*, 19 (1), 426.

Wang, P., Fouracre, J., Kelly, S., Karki, S., Gowik, U., Aubry, S., Shaw, M.K., Westhoff, P., Slamet-Loedin,
I.H., Quick, W.P., Hibberd, J.M., and Langdale, J.A., 2013. Evolution of GOLDEN2-LIKE gene
function in C3 and C4 plants. *Planta*, 237 (2), 481–495.

Washburn, J.D., Strable, J., Dickinson, P., Kothapalli, S.S., Brose, J.M., Covshoff, S., Conant, G.C.,
Hibberd, J.M., and Pires, J.C., 2021. Distinct C4 sub-types and C3 bundle sheath isolation in the
Paniceae grasses. *Plant Direct*, 5 (12), 1–14.

Waters, M.T. and Langdale, J.A., 2009. The making of a chloroplast. *The EMBO Journal*, 28 (19), 2861–
2873.

Waters, M.T., Moylan, E.C., and Langdale, J.A., 2008. GLK transcription factors regulate chloroplast
development in a cell-autonomous manner. *Plant Journal*, 56 (3), 432–444.

Waters, M.T., Wang, P., Korkaric, M., Capper, R.G., Saunders, N.J., and Langdale, J.A., 2009. GLK
Transcription Factors Coordinate Expression of the Photosynthetic Apparatus in Arabidopsis.
Plant Cell, 21 (4), 1109–1128.

Wickett, N.J., Mirarab, S., Nguyen, N., Warnow, T., Carpenter, E., Matasci, N., Ayyampalayam, S.,
Barker, M.S., Burleigh, J.G., Gitzendanner, M.A., Ruhfel, B.R., Wafula, E., Der, J.P., Graham, S.W.,
Mathews, S., Melkonian, M., Soltis, D.E., Soltis, P.S., Miles, N.W., Rothfels, C.J., Pokorny, L., Shaw,
A.J., DeGironimo, L., Stevenson, D.W., Surek, B., Villarreal, J.C., Roure, B., Philippe, H.,
dePamphilis, C.W., Chen, T., Deyholos, M.K., Baucom, R.S., Kutchan, T.M., Augustin, M.M., Wang,
J., Zhang, Y., Tian, Z., Yan, Z., Wu, X., Sun, X., Wong, G.K.-S., and Leebens-Mack, J., 2014.
Phylotranscriptomic analysis of the origin and early diversification of land plants. *Proceedings of
the National Academy of Sciences*, 111 (45), E4859–E4868.

Woo, K.C., Anderson, J.M., Boardman, N.K., Downton, W.J.S., Osmond, C.B., and Thorne, S.W., 1970.
Deficient Photosystem II in Agranal Bundle Sheath Chloroplasts of C4 Plants. *Proceedings of the
National Academy of Sciences*, 67 (1), 18–25.

Yasumura, Y., Moylan, E.C., and Langdale, J.A., 2005. A Conserved Transcription Factor Mediates

Nuclear Control of Organelle Biogenesis in Anciently Diverged Land Plants. *Plant Cell*, 17 (7),

1894–1907.

Zubo, Y.O., Blakley, I.C., Franco-Zorrilla, J.M., Yamburenko, M. V, Solano, R., Kieber, J.J., Loraine, A.E.,

and Schaller, G.E., 2018. Coordination of Chloroplast Development through the Action of the GNC

and GLK Transcription Factor Families. *Plant Physiology*, 178 (1), 130–147.

    FIGURE LEGENDS

Figure 1. Chloroplast development is not perturbed in *Zmglk1* mutants of maize. **A)** Gene model

of *ZmGLK1* showing predicted exons in black boxes and the target site of CRISPR/Cas9 mediated

mutagenesis in yellow and red below. The table shows the sequence of the wild-type (WT) and mutated

alleles with the predicted protein length. The gRNA sequence is highlighted in yellow and the PAM

sequence in red. **B-C)** Whole plant phenotype of wild-type (B) and a representative *Zmglk1* mutant 30

days after sowing (C). Scale bars correspond to 10 cm. **D-G)** Electron micrographs of chloroplast

ultrastructure in bundle sheath (BSC; D-E) and mesophyll (MC; F-G) cells of WT (D, F) and a

representative *Zmglk1* mutant (E, G). Scale bars correspond to 2 µm. **H-L)** Measurements of thylakoid

occupancy in mesophyll cells (H), thylakoid occupancy in bundle sheath cells (I), total chlorophyll (J),

PSII operating efficiency (K) and maximum quantum efficiency of PSII photochemistry (L). The WT

segregant is depicted in blue and the four independent *Zmglk1* lines are depicted in green. Each dot

represents a biological replicate, the square is the average value for that line and the bars are the

associated standard deviations. A minimum of 5 biological replicates were evaluated for thylakoid

occupancy and chlorophyll quantification, and 3 biological replicates were assessed for PSII

fluorescence. Letters indicate statistically different groups in comparison to WT; n.s. – no significant

statistical support for a difference from WT (pvalue≤0.05, one-way ANOVA and Tukey's HSD).

Figure 2. Chloroplast development is not perturbed in *Svglk1* mutants of setaria. A) Gene model

of *SvGLK1* showing predicted exons in black boxes and the target site of CRISPR/Cas9 mediated

mutagenesis in yellow and red below. The table shows the sequence of the wild-type (WT) and mutated

alleles with the predicted protein length. The gRNA sequence is highlighted in yellow and the PAM

sequence in red. **B-D)** Whole plant phenotype of wild-type (B) and *Svglk1* mutants 20 days after sowing

(C-D). Scale bars correspond to 5 cm. **E-J)** Electron micrographs of chloroplast ultrastructure in bundle

sheath (BSC; E-G) and mesophyll (MC; H-J) cells of WT (E, H) and *Svglk1* mutants (F, G, I, J). Scale

bars correspond to 2 µm. **K-O)** Measurements of thylakoid occupancy in mesophyll cells (K), thylakoid

occupancy in bundle sheath cells (L), total chlorophyll (M), PSII operating efficiency (N) and maximum

quantum efficiency of PSII photochemistry (O). The WT segregant is depicted in blue and the two

independent *Svglk1* lines are depicted in green. Each dot represents a biological replicate, the square

is the average value for that line and the bars are the associated standard deviations. Twelve

chloroplasts were assessed for thylakoid occupancy quantification, and 5 biological replicates were

assessed for chlorophyll quantification and PSII fluorescence for each sample. n.s. - no significant

statistical support for a difference from WT (p-value≤0.05, one-way ANOVA and Tukey's HSD).

Figure 3. Chloroplast development is perturbed in both mesophyll and bundle sheath cells of

***Svglk2* mutants of setaria. A)** Gene model of *SvGLK2* showing predicted exons in black boxes and

the target site of CRISPR/Cas9 mediated mutagenesis in yellow and red below. The table shows the

sequence of the wild-type (WT) and mutated alleles with the predicted protein length. The gRNA sequence is highlighted in yellow and the PAM sequence in red. **B)** Whole plant phenotype of wild-type, a heterozygous and a homozygous *Svglk2* mutant 20 days after sowing. Scale bar corresponds to 5 cm. **C-D)** Confocal images of leaf cross sections showing overall chloroplast structure in WT (C) and the *Svglk2* mutant (D). Bundle sheath cells are outlined in white. Scale bars correspond to 50 μm. **E-H)** Electron micrographs of chloroplast ultrastructure in bundle sheath (BSC; E-F) and mesophyll (MC; G-H) cells of WT (E, G) and the *Svglk2* mutant (F, H). Yellow arrows point to fragmented membranes. Scale bars correspond to 2 μm. **I-K)** Measurements of thylakoid occupancy in bundle sheath cells (I), thylakoid occupancy in mesophyll cells (J) and total chlorophyll (K). The WT segregant is depicted in purple and the *Svglk2* line is depicted in pink. Each dot represents a biological replicate, the square is the average value for that line and the bars are the associated standard deviations. Twelve chloroplasts were assessed for thylakoid occupancy quantification, and 5 biological replicates were assessed for chlorophyll quantification and PSII fluorescence for each sample. Letters indicate statistically different groups in comparison to WT; n.s. - no significant statistical support for a difference from WT (pvalue≤0.05, one-way ANOVA and Tukey's HSD).

Figure 4. *GLK1* and *GLK2* transcript levels are reduced in *Svglk2* and *Zmg2* mutants of setaria and maize. A-H) Quantitative analysis of gene expression via qPCR in segregating wild-type (WT) siblings and *glk* mutants of maize (A-D) and setaria (E-F). Relative normalized expression of *ZmGLK1* (A), *ZmG2* (B), *ZmGNC* (C), *ZmCGA1* (D), *SvGLK1* (E) and *SvGLK2* (F) was calculated based on the expression of two reference genes for each sample. Each dot represents a biological replicate, the square is the mean value and the bars are the associated standard deviations. A minimum of 3 and 5 biological replicates were evaluated for maize and setaria samples, respectively. Statistical significance was calculated using a Student's t-test and pairwise comparison with the respective WT sample. Letters indicate statistically different groups in comparison to WT; n.s. - no significant statistical support for a difference from WT (p-value≤0.05).

Figure 5. *ZmGLK1* and *ZmG2* complement the *Svglk2* phenotype and activate transcription of the setaria homologs. **A-B)** Whole plant phenotype of *Svglk2* mutant overexpressing *ZmGLK1* (A) or *ZmG2* (B) 30 days after sowing. Scale bars correspond to 5 cm. **C-F)** Quantitative analysis of gene expression via qPCR of *ZmGLK1* (C), *ZmG2* (D), *SvGLK1* (E) and *SvGLK2* (F). Relative gene expression was calculated based on the expression of two reference genes for each sample. Each dot represents one of the 5 biological replicates evaluated, the square is the mean value and the bars are the associated standard deviations. Statistical significance was calculated using a Student's t-test and pairwise comparison with the respective wild-type (WT) sample. Letters indicate statistically different groups in comparison to WT and *Svglk2* (p-value≤0.05).

Figure 6. *ZmGLK1* and *ZmG2* are sufficient to induce dimorphic chloroplast development in both mesophyll and bundle sheath cells of setaria. **A-C)** Confocal images of leaf cross sections showing overall chloroplast morphology in the *Svglk2* mutant (A, same as Figure 3D), and representative lines expressing ZmGLK1 (B) or ZmG2 (C). Bundle sheath cells are outlined in white. Yellow arrows indicate ectopic chloroplasts developing in vascular cells. Scale bars correspond to 50 μm. **D-I)** Electron micrographs of chloroplast ultrastructure in bundle sheath (BSC; D-F) and mesophyll (MC; G-I) cells of the *Svglk2* mutant (D-G; same as Figure 3G, H) and one representative line overexpressing *ZmGLK1* (E-H) or ZmG2 (F-I). Scale bars correspond to 2 μm. **J-L)** Measurements of thylakoid occupancy in mesophyll cells (J), thylakoid occupancy in bundle sheath cells (K) and total chlorophyll (L). The wild-type (WT) segregant is depicted in purple, the *Svglk2* line in pink, the three independent lines overexpressing *ZmGLK1* in orange and those overexpressing *ZmG2* in yellow. Each dot represents a biological replicate, the square is the average value for that line and the bars are the associated standard deviations. A minimum of 10 chloroplasts were evaluated for thylakoid occupancy quantification and 5-10 biological replicates were used for chlorophyll quantification. Letters indicate statistically different groups in comparison to WT and *Svglk2* (p-value≤0.05, one-way ANOVA and Tukey's HSD).

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Phenotypic characterization of three independent *Zmglk1* mutant lines. A-C) Whole plant phenotype 30 days after sowing. Scale bars correspond to 10 cm. **D-F)** Transmission electron microscopy (TEM) images showing mesophyll chloroplast ultrastructure. Scale bars correspond to 2 µm, 5 µm and 2 µm, respectively. **G-I)** TEM images showing bundle sheath chloroplast ultrastructure. Scale bars correspond to 5 µm, 10 µm and 2 µm, respectively.

Figure S2. Comparison of GLK sequences. **A)** Phylogenetic reconstruction of the *GLK* gene family. Protein sequences of the extensively characterised GLK members ZmGLK1, ZmG2, OsGLK1 and OsGLK2 were used to create a HMMER profile that was used as a query to search for homologs in whole protein databases of 11 species (*Zea mays, Setaria viridis, S. italica, Panicum virgatum, Sorghum bicolor, Oryza sativa, Brachypodium distachyon, Vitis vinifera, Arabidopsis thaliana, Marchantia polymorpha* and *Physcomitrium patens*). The presence of the MYB domain in those sequences was confirmed using InterproScan, and only the sequences containing the conserved AREAEAA domain characteristic of the GLK subfamily were selected. The resulting 25 amino acid sequences were aligned using Clustal Omega. Phylogenetic reconstruction was performed using Bayesian inference on Mr. Bayes using Poisson with fixed substitution rates as a model. Two independent MCMC runs with 1,000,000 generations were performed and no further improvement of log likelihood of prior probabilities was observed. The tree was visualized on FigTree. Clade support on each node was assessed using Bayesian posterior probabilities. Members of the GLK1 subgroup are depicted in green and GLK2 in blue. **B)** Amino acid sequence alignment showing conservation between ZmGLK1, SvGLK1, ZmG2 and SvGLK2. Magenta bars show the HLH associated with MYB or MYB-related proteins; the AREAEAA conserved site is in blue and the last exon that contains the GCT-box domain is in dark green.

Figure S3. *GLK* transcript accumulation in bundle sheath and mesophyll cells. A-B) Gene expression in reads per kilobase million (RPKM) in mesophyll (MC) and bundle sheath cells (BSC) of maize – *ZmGLK1* and *ZmG2* (A) and setaria – *SvGLK1* and *SvGLK2* (B). Maize expression data were published by Chang et al. (2012) and setaria expression data were published by John et al. (2014). **C)** Phylogenetic reconstruction based on 1000 bp upstream of the ATG in *GLK1* and *GLK2* genes. Promoter sequences were aligned using MAFFT and phylogenetic reconstruction was performed using MEGA software v. 7.0.26 and Maximum Likelihood based on the Tajima-Nei model. Clade support is based on 1000 bootstrap replicates. **D)** Accumulation levels of *GLK1* and *GLK2* in transcripts per million (TPM) in three different species of Paniceae grasses (*Urochloa fusca*, *Panicum hallii* and *Digitaria californica*). All species carry out C_4 photosynthesis, but each uses a different decarboxylation pathway in the bundle sheath cells.

Figure S4. Phenotypic characterization of two independent lines of *Svglk2* complemented with *ZmGLK1* or *ZmG2*. A-B) Schematic of the constructs used to express *ZmGLK1* (A) or *ZmG2* (B) in the *Svglk2* background. *HygR* depicts the hygromycin phosphotransferase gene and $OsACT_{pro}$ and $ZmUBI_{pro}$ represent the constitutive rice actin and maize ubiquitin promoters, respectively. LB and RB refers to left and right borders. **C-J)** Transmission electron microscopy images showing mesophyll (MC) and bundle sheath (BSC) cell chloroplast ultrastructure in lines overexpressing *ZmGLK1* (C-F) or *ZmG2* (G-J). Scale bar sizes are stated on the figures. **K-N)** Confocal images of leaf cross sections from lines overexpressing *ZmGLK1* (K-L) or *ZmG2* (M-N) showing morphology of chloroplasts. Bundle sheath cells are outlined in white. Scale bars correspond to 50 µm. Yellow arrows indicate ectopic chloroplast formation in vascular cells.

Figure 1. Chloroplast development is not perturbed in *Zmglk1* mutants of maize. **A)** Gene model of *ZmGLK1* showing predicted exons in black boxes and the target site of CRISPR/Cas9 mediated mutagenesis in yellow and red below. The table shows the sequence of the wild-type (WT) and mutated alleles with the predicted protein length. The gRNA sequence is highlighted in yellow and the PAM sequence in red. **B-C)** Whole plant phenotype of wild-type (B) and a representative *Zmglk1* mutant 30 days after sowing (C). Scale bars correspond to 10 cm. **D-G)** Electron micrographs of chloroplast ultrastructure in bundle sheath (BSC; D-E) and mesophyll (MC; F-G) cells of WT (D, F) and a representative *Zmglk1* mutant (E, G). Scale bars correspond to 2 μm. **H-L)** Measurements of thylakoid occupancy in mesophyll cells (H), thylakoid occupancy in bundle sheath cells (I), total chlorophyll (J), PSII operating efficiency (K) and maximum quantum efficiency of PSII photochemistry (L). The WT segregant is depicted in blue and the four independent *Zmglk1* lines are depicted in green. Each dot represents a biological replicate, the square is the average value for that line and the bars are the associated standard deviations. A minimum of 5 biological replicates were evaluated for thylakoid occupancy and chlorophyll quantification, and 3 biological replicates were assessed for PSII fluorescence. Letters indicate statistically different groups in comparison to WT; n.s. – no significant statistical support for a difference from WT (pvalue≤0.05, one-way ANOVA and Tukey's HSD).

Figure 2. Chloroplast development is not perturbed in *Svglk1* mutants of setaria. A) Gene model of *SvGLK1* showing predicted exons in black boxes and the target site of CRISPR/Cas9 mediated mutagenesis in yellow and red below. The table shows the sequence of the wild-type (WT) and mutated alleles with the predicted protein length. The gRNA sequence is highlighted in yellow and the PAM sequence in red. **B-D)** Whole plant phenotype of wild-type (B) and *Svglk1* mutants 20 days after sowing (C-D). Scale bars correspond to 5 cm. **E-J)** Electron micrographs of chloroplast ultrastructure in bundle sheath (BSC; E-G) and mesophyll (MC; H-J) cells of WT (E, H) and *Svglk1* mutants (F, G, I, J). Scale bars correspond to 2 µm. **K-O)** Measurements of thylakoid occupancy in mesophyll cells (K), thylakoid occupancy in bundle sheath cells (L), total chlorophyll (M), PSII operating efficiency (N) and maximum quantum efficiency of PSII photochemistry (O). The WT segregant is depicted in blue and the two independent *Svglk1* lines are depicted in green. Each dot represents a biological replicate, the square is the average value for that line and the bars are the associated standard deviations. Twelve chloroplasts were assessed for thylakoid occupancy quantification, and 5 biological replicates were assessed for chlorophyll quantification and PSII fluorescence for each sample. n.s. - no significant statistical support for a difference from WT (p-value≤0.05, one-way ANOVA and Tukey's HSD).

Figure 3. Chloroplast development is perturbed in both mesophyll and bundle sheath cells of *Svglk2* mutants of setaria. A) Gene model of *SvGLK2* showing predicted exons in black boxes and the target site of CRISPR/Cas9 mediated mutagenesis in yellow and red below. The table shows the sequence of the wild-type (WT) and mutated alleles with the predicted protein length. The gRNA sequence is highlighted in yellow and the PAM sequence in red. **B)** Whole plant phenotype of wild-type, a heterozygous and a homozygous *Svglk2* mutant 20 days after sowing. Scale bar corresponds to 5 cm. **C-D)** Confocal images of leaf cross sections showing overall chloroplast structure in WT (C) and the *Svglk2* mutant (D). Bundle sheath cells are outlined in white. Scale bars correspond to 50 μm. **E-H)** Electron micrographs of chloroplast ultrastructure in bundle sheath (BSC; E-F) and mesophyll (MC; G-H) cells of WT (E, G) and the *Svglk2* mutant (F, H). Yellow arrows point to fragmented membranes. Scale bars correspond to 2 μm. **I-K)** Measurements of thylakoid occupancy in bundle sheath cells (I), thylakoid occupancy in mesophyll cells (J) and total chlorophyll (K). The WT segregant is depicted in purple and the *Svglk2* line is depicted in pink. Each dot represents a biological replicate, the square is the average value for that line and the bars are the associated standard deviations. Twelve chloroplasts were assessed for thylakoid occupancy quantification, and 5 biological replicates were assessed for chlorophyll quantification and PSII fluorescence for each sample. Letters indicate statistically different groups in comparison to WT; n.s. - no significant statistical support for a difference from WT (pvalue≤0.05, one-way ANOVA and Tukey's HSD).

Figure 4. *GLK1* **and** *GLK2* **transcript levels are reduced in** *Svglk2* **and** *Zmg2* **mutants of setaria and maize. A-H)** Quantitative analysis of gene expression via qPCR in segregating wild-type (WT) siblings and *glk* mutants of maize (A-D) and setaria (E-F). Relative normalized expression of *ZmGLK1* (A), *ZmG2* (B), *ZmGNC* (C), *ZmCGA1* (D), *SvGLK1* (E) and *SvGLK2* (F) was calculated based on the expression of two reference genes for each sample. Each dot represents a biological replicate, the square is the mean value and the bars are the associated standard deviations. A minimum of 3 and 5 biological replicates were evaluated for maize and setaria samples, respectively. Statistical significance was calculated using a Student's t-test and pairwise comparison with the respective WT sample. Letters indicate statistically different groups in comparison to WT; n.s. - no significant statistical support for a difference from WT (p-value≤0.05).

Figure 5. *ZmGLK1* **and** *ZmG2* **complement the** *Svglk2* **phenotype and activate transcription of the setaria homologs**. **A-B)** Whole plant phenotype of *Svglk2* mutant overexpressing *ZmGLK1* (A) or *ZmG2* (B) 30 days after sowing. Scale bars correspond to 5 cm. **C-F)** Quantitative analysis of gene expression via qPCR of *ZmGLK1* (C), *ZmG2* (D), *SvGLK1* (E) and *SvGLK2* (F). Relative gene expression was calculated based on the expression of two reference genes for each sample. Each dot represents one of the 5 biological replicates evaluated, the square is the mean value and the bars are the associated standard deviations. Statistical significance was calculated using a Student's t-test and pairwise comparison with the respective wild-type (WT) sample. Letters indicate statistically different groups in comparison to WT and *Svglk2* (p-value≤0.05).

Figure 6. *ZmGLK1* and *ZmG2* are sufficient to induce dimorphic chloroplast development in both mesophyll and bundle sheath cells of setaria. **A-C)** Confocal images of leaf cross sections showing overall chloroplast morphology in the *Svglk2* mutant (A, same as Figure 3D), and representative lines expressing ZmGLK1 (B) or ZmG2 (C). Bundle sheath cells are outlined in white. Yellow arrows indicate ectopic chloroplasts developing in vascular cells. Scale bars correspond to 50 µm. **D-I)** Electron micrographs of chloroplast ultrastructure in bundle sheath (BSC; D-F) and mesophyll (MC; G-I) cells of the *Svglk2* mutant (D-G; same as Figure 3G, H) and one representative line overexpressing *ZmGLK1* (E-H) or ZmG2 (F-I). Scale bars correspond to 2 µm. **J-L)** Measurements of thylakoid occupancy in mesophyll cells (J), thylakoid occupancy in bundle sheath cells (K) and total chlorophyll (L). The wild-type (WT) segregant is depicted in purple, the *Svglk2* line in pink, the three independent lines overexpressing *ZmGLK1* in orange and those overexpressing *ZmG2* in yellow. Each dot represents a biological replicate, the square is the average value for that line and the bars are the associated standard deviations. A minimum of 10 chloroplasts were evaluated for thylakoid occupancy quantification and 5-10 biological replicates were used for chlorophyll quantification. Letters indicate statistically different groups in comparison to WT and *Svglk2* (p-value≤0.05, one-way ANOVA and Tukey's HSD).