Hierarchical cortical and subcortical mechanisms underlying binocular rivalry

Chencan Qian[1,9], Zhiqiang Chen[1,2,3,9], Gilles de Hollander[5], Tomas Knapen[6,7,8], Zihao Zhang[1,4], Sheng He[1,2], Peng Zhang[1,2,4,*]

[1]State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese Academy of Sciences, Beijing 100101, China

[2]School of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

[3]Sino-Danish College, University of Chinese Academy of Sciences, Beijing 100049, China

[4]Institute of Artificial Intelligence, Hefei Comprehensive National Science Center, Hefei 230026, China

[5]Zurich Center for Neuroeconomics, Department of Economics, University of Zurich, Zurich, Switzerland

[6]Cognitive Psychology, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

[7]Spinoza Centre for Neuroimaging, Royal Academy of Sciences, Amsterdam, Netherlands

[8]Netherlands Institute for Neuroscience, Royal Academy of Sciences, Amsterdam, Netherlands

[9]C.C.Q. and Z.Q.C. contributed equally to this work.

*Address correspondence to Peng Zhang (zhangpeng@ibp.ac.cn)

Abstract

Conscious perception alternates between the two eyes' images during binocular rivalry. How hierarchical processes in our brain interact to resolve visual competition remains unclear. Here we investigated mesoscale neural circuitry of binocular rivalry in human cortical and subcortical areas using high-resolution functional MRI at 7 Tesla. Eye-specific rivalry modulation was strongest in the superficial layers of V1 ocular dominance columns (ODCs), and more synchronized in the superficial and deep layers. The intraparietal sulcus (IPS) generated stronger eye-specific modulation and increased feedback connectivity to the early visual cortex during binocular rivalry compared to stimulus replay. Although there was no evidence of eye-specific rivalry modulation in the lateral geniculate nucleus (LGN) of the thalamus, strong perceptual rivalry modulation can be found in its parvocellular (P) subdivision. Finally, IPS and ventral pulvinar showed robust perceptual rivalry modulation and increased connectivity to the early visual cortex. These findings demonstrate that local interocular competition arises from lateral mutual inhibition between V1 ODCs, and feedback signals from IPS to visual cortex and visual thalamus further resolve and synchronize visual competitions to generate conscious perception.

Highlights

- Eye rivalry is strongest in the superficial layers of V1 ODCs and more synchronized in superficial and deep layers
- IPS generates stronger eye-specific modulation and increased connectivity to V1 during rivalry compared to replay
- LGN shows no evidence of eye rivalry but strong perceptual rivalry in its P subdivision
- IPS and ventral pulvinar show robust perceptual rivalry and increased connectivity to the early visual cortex

Key words

Visual consciousness; Binocular rivalry; Cortical columns and layers; Subcortical nuclei; Feedforward; Feedback; Lateral inhibition; 7T high-resolution fMRI

Introduction

Two incompatible images presented to the two eyes compete for access to consciousness in alternation. This visual illusion, called binocular rivalry, is an ideal model to study how our brain resolves visual competition (R Blake & Logothetis, 2002; Dayan, 1998; Wilson, 2003), a key mechanism to generate conscious visual perception (Randolph Blake et al., 2014; Crick, 1996; Myerson et al., 1981). Although rivalry-related activity has been found in many brain areas (Jan Brascamp et al., 2018; Tong et al., 2006), how hierarchical neural processes in the brain interact to resolve visual competition remains unclear.

It has been proposed that binocular rivalry could arise from interocular competitions in the early visual areas (R Blake, 1989), either lateral mutual inhibition between adjacent ocular dominance columns (ODCs) in the primary visual cortex (V1), or interlaminar inhibition between adjacent ocular layers in the lateral geniculate nucleus (LGN) of the thalamus (Kacie Dougherty et al., 2018, 2021; Guillery & Colonnier, 1970). In support of this hypothesis, human fMRI studies found robust eye-specific rivalry modulations in the blindspot area (Tong & Engel, 2001) and ocular-biased voxels in V1 (Haynes et al., 2005), and even in the LGN (Haynes et al., 2005). However, since these early fMRI studies didn't resolve activity from V1 ODCs or LGN ocular layers, the neural mechanisms of interocular competition still lack concluding evidence. Different from the fMRI results, single-unit spiking activity showed no evidence of binocular rivalry in the LGN of alert monkeys (Lehky & Maunsell, 1996), and a weak effect in V1 (Leopold & Logothetis, 1996). Since BOLD signals can reflect synaptic input activity (Logothetis & Wandell, 2004), one possible explanation for the discrepancy between single-unit and fMRI results is that feedback modulations from higher-order brain areas drive rivalry-related activity in the early visual areas (de Jong et al., 2020; Maier et al., 2008). A potential role of eye-specific feedbacks in resolving interocular conflicts is supported by behavioral evidence that top-down attention can be eye-specific (Zhang et al., 2012), and by electrophysiology and neuroimaging evidence of eye-specific representations in extrastriate cortex (Burkhalter & Van Essen, 1986; Maunsell & Van Essen, 1983; Schwarzkopf et al., 2010; Zaretskaya et al., 2020). Therefore, it remains unclear whether interocular competition in binocular rivalry arises from interlaminar inhibition between ocular layers in the LGN, lateral inhibition between V1 ODCs, or is driven by eye-specific feedbacks from higher-order brain areas.

In addition to interocular competition, binocular rivalry could also involve pattern competition between stimulus representations at multiple levels of the visual hierarchy, and possibly attention and perceptual decision-making mechanisms in high-level brain areas. In this hierarchical whole-brain network, the role of frontoparietal areas is the most debated. Although a causal role of frontoparietal activity in generating perceptual transitions remains

controversial (J Brascamp et al., 2015; Lumer et al., 1998), converging evidence demonstrate that binocular rivalry requires top-down attention (J. W. Brascamp & Blake, 2012; Li et al., 2017; Zhang et al., 2011), suggesting a potential role of frontoparietal attention network in resolving visual competition. Moreover, it is unclear whether frontoparietal areas also represent perceptual state during rivalry. Another subcortical area that is rarely investigated but might play important roles in perceptual rivalry is the pulvinar of the thalamus. Interconnected with frontoparietal areas and the visual cortex, pulvinar may regulate information transfer between cortical areas and support cortical computations to resolve perceptual conflicts (Jaramillo et al., 2019; Saalmann et al., 2012; Wilke et al., 2009; Zhou et al., 2016). Finally, parallel visual pathways might be differentially involved in binocular rivalry. Although behavioral studies suggest that the parvocellular (P) pathway is more involved in rivalry than the magnocellular (M) pathway (He et al., 2005), there is no direct neuroimaging evidence supporting this hypothesis. To this date, the hierarchical whole-brain network of perceptual rivalry has not been clearly demonstrated.

Using high-resolution fMRI at 7 Tesla to measure mesoscale activity in the human brain, we investigated hierarchical neural mechanisms underlying binocular rivalry in cortical and subcortical areas. To reveal the neural circuitry of interocular competition, Experiment 1 and 2 studied eye-specific rivalry modulations in V1 ODCs at different cortical depth and LGN ocular layers, and also in the higher-order extrastriate and parietal cortex. To investigate the hierarchical whole-brain network of perceptual rivalry, Experiment 3 used M and P pathway-selective visual stimuli to study perceptual rivalry modulations over the whole brain.

Results

Fig. 1a shows the stimuli and procedures for Experiment 1 and 2. In the rivalry condition, a pair of red/green gratings in orthogonal orientations were dichoptically presented to the two eyes. Subjects reported their perception with button presses (red, green or mixed). In the replay condition, monocular images were presented in alternations with simulated transitions to match the temporal sequence of perception during binocular rivalry, and subjects reported their perception as in the rivalry condition. In localizer runs, a full contrast counterphase flickering checkerboard was alternatively presented to the two eyes to measure the ocular bias of voxels in V1 and LGN. Figure 1b shows three possible mechanisms that may support interocular competition in binocular rivalry: interlaminar inhibition between ocular layers of the LGN, lateral inhibition between V1 ODCs, or eye-specific feedback modulation of V1 and/or LGN activity. Although there is no evidence that corticogeniculate feedbacks can be eye-specific, we still included it as a possibility in the last hypothesis according to the (Haynes et al., 2005) study.

Figure 1. (a) Stimuli and procedures of Experiment 1 and 2. In rivalry runs, rotating red and green gratings in
orthogonal orientations were dichoptically presented to the two eyes. In replay runs, monocular stimuli were
alternatively presented to the two eyes to simulate the perception in the previous rivalry run. Subjects reported
their perception with button presses in binocular rivalry and simulated replay. In localizer runs, a high contrast
checkerboard flicker was monocularly delivered to the two eyes in alternation. **(b) Possible neural circuits of**
interocular competition in binocular rivalry. (1) Interlaminar mutual inhibition between LGN ocular layers. (2)
Lateral mutual inhibition between V1 ODCs. (3) Eye-specific feedback modulation from higher-order cortical
areas. Solid and dashed arrows indicate feedforward and feedback connections, respectively. Green dots connected
by solid green lines denote mutual inhibitions. Abbreviations: S (superficial), M (middle), D (deep), LE (left eye),
RE (right eye), ODC (ocular dominance column).

Based on known anatomical connections of the primate geniculostriate pathway
(Felleman & Van Essen, 1991), feedforward input from the LGN mainly terminates in the
middle layer (layer 4) of V1, cortico-cortical feedbacks target the superficial (layers 1/2/3)
and deep layers (layers 5/6), and lateral inhibitions between ODCs through horizontal
connections are most prominent in the superficial layers (layers 2/3) (Buzs et al., 2001; K
Dougherty et al., 2019; Gilbert & Wiesel, 1983; Sengpiel et al., 1995). If interocular
competition arises from interlaminar inhibition in the LGN, eye-specific rivalry modulation
should be strongest in V1 middle layer. Otherwise, if interocular competition arises from
lateral inhibition between V1 ODCs, eye-specific modulation should be strongest in the
superficial layers. Finally, if feedback processes are involved to resolve interocular
competition, eye-specific effect of rivalry should be stronger in the superficial and deep layers
compared to the middle layer.
Therefore, in Experiment 1, we tested the three hypotheses using cortical layer-
dependent fMRI at submillimeter resolution. T2*w BOLD signals from the early visual
cortex and parietal cortex were acquired with a gradient echo planar imaging (GE-EPI)
sequence at 0.8-mm isotropic resolution. In a representative subject (S06), we also acquired
T2w BOLD signals from V1 using a passband balanced steady-state free-precession (bSSFP)

sequence, with 3-mm thick slices at 0.5-mm in-plane resolution placed perpendicular to the

calcarine sulcus. In Experiment 2, we further investigated whether the LGN was involved in

eye-specific rivalry, as suggested by the first and possibly the last hypothesis, using a GE-EPI

sequence at 1.2-mm isotropic resolution covered the visual thalamus and early visual cortex.

Eye-specific rivalry modulation is strongest in the superficial depth of V1 ODCs

Figure 2. Eye-specific modulation of V1 ODCs at different cortical depths in Experiment 1. (a) OD patterns

of a representative subject (S01, the author P.Z.) on the inflated cortical surface of right hemisphere. The color bar

indicates the beta values for LE-RE contrast, thresholded at abs(t) > 2 (or $p < 0.05$ uncorrected), same for lower

panel in (d). The white lines delineate V1/V2, as well as the 5° eccentricity based on the Benson14 atlas (Noah C.

Benson et al., 2014). **(b)** Event-related timecourses of eye-specific modulations in ocular-biased voxels. Solid

(dashed) lines indicate responses for the preferred (non-preferred) perception, e.g., stimulus presented to the LE

(RE) for LE (RE)-biased voxels. The bars below indicate time points with a significant difference between

different perceptions (cluster-based permutation test, cluster defining threshold and cluster significance level, p <

0.05) **(c)** Comparison of the ODC map from the ocular-bias localizer and eye-specific response pattern during

binocular rivalry for S01. **(d)** Cortical depth map in the occipital lobe generated with an equivolume algorithm

overlaid on T1w image (upper) for the same subject; GLM beta map (LE-RE) within gray matter overlaid on mean

EPI image (lower). Green and purple lines indicate the pial and WM surfaces, respectively. **(e)** Eye-specific

response modulation peaked in the middle of cortical depth in the ocular-bias localizer. **(f)** Normalized eye-

specific modulation from different cortical depths in rivalry and replay conditions. Error bars indicate 95%

confidence interval from bootstrap. **(g)** The modulation ratio between the rivalry and replay conditions. Error bars

indicate 95% confidence interval from bootstrap. **(h)** Slice prescriptions for the 2D-bSSFP Experiment (0.5-mm

in-plane resolution, 3-mm thickness) in a representative subject (S06, the author C.Q.). From T2*w GRE images,

the line of Gennari is clearly visible in the middle layer of V1 gray matter. **(i)** The original bSSFP image. **(j)** ODCs

can be clearly identified on the cross section of calcarine sulcus (white arrow). **(k)** The V1 depth profile of eye-

specific modulation in the rivalry, replay and localizer conditions for T2w BOLD signals with bSSFP fMRI.

Figure 2 shows eye-specific modulations in V1 ODCs in Experiment 1. The

interdigitated patterns of V1 ODCs can be robustly resolved with 7T GE-BOLD fMRI at 0.8-

mm isotropic resolution (Fig. 2a for a representative subject S01, Fig. S1 for all subjects). The

orientations of the columns are roughly perpendicular to the V1/V2 boundary in its vicinity,

and highly reproducible across sessions on different days (Fig. S2, $r = 0.697$, $p < 0.001$,

Monte Carlo test). Event-related average of eye-specific modulations were time-locked to key

presses reflecting perceptual switches (Fig. 2b). From the time of perceptual switch, BOLD

signals increased when subjects perceived the preferred stimulus of the ocular-biased voxels

(LE/RE percept for LE/RE biased voxels), and decreased when the non-preferred stimulus

was perceived (RE/LE percept for LE/RE biased voxels). The modulation amplitude during

binocular rivalry was about 40% of that in the replay condition. The map of rivalry

modulation (difference between LE and RE percepts, 8-mm FWHM high-pass filtered)

matched well with the ODC map acquired from the localizer (Fig. 2c, $r = 0.475$, $p < 0.001$).

These results clearly demonstrate that eye-specific modulation of rivalry occurs at the level of

cortical columns.

Cortical depth was estimated for each voxel with an equivolume method (Waehnert et al.,

2014), based on manually edited cortical surface reconstructions (Fig. 2d upper). The

columnar structure of ODCs perpendicular to the cortical surface can be clearly seen (Fig. 2d

lower). The differential response between the left and right eye stimulation in the localizer

peaked in the middle depth of V1 (Fig. 2e), consistent with the fact that thalamocortical

projections terminate mainly at layer 4C. Thus, the OD column-specific response derived

from the differential of balanced responses to the LE and RE stimuli largely reduced the non-

specific signals in the superficial layers associated with the blooming effect of pial veins

(Moon et al., 2007; Uludag & Havlicek, 2021). We next investigated how this laminar-

columnar circuit reflects the endogenous mechanisms that gate perceptual awareness in

binocular rivalry. The normalized eye-specific modulations in the rivalry and replay

conditions were shown in Fig. 2f. Two-way repeated measures ANOVA showed a significant

interaction of cortical depth (superficial/middle/deep) and stimulus conditions (rivalry/replay):

$F(2,22) = 6.372$, $p = 0.013$, $\eta_p^2 = 0.367$. In the replay condition, the middle layer showed the

strongest effect of eye-specific modulation (main effect of depth, $F(2,22) = 15.290$, $p < 0.001$

Bonferroni corrected, $\eta_p^2 = 0.582$; deep vs. middle, $t(11) = -6.218$, $p < 0.001$, Cohen's $d =$

2.921; middle vs. superficial, $t(11) = 3.608$, $p = 0.004$, Cohen's $d = 1.600$), consistent with the

feedforward input from the LGN. During binocular rivalry, eye-specific modulation was more

biased to the superficial depth (main effect of depth, $F(2,22) = 12.859$, $p < 0.001$, $\eta_p^2 = 0.539$;

deep vs. middle, $t(11) = -3.486$, $p = 0.005$, Cohen's $d = 1.794$; middle vs. superficial, $t(11) = -$

1.617, $p = 0.134$, Cohen's $d = 0.665$),suggesting a major contribution of lateral mutual

inhibition between ODCs that is most prominent in V1 superficial layers. The laminar profiles

without normalization were qualitatively the same. To further reduce the influence of

superficial bias of BOLD signals and to directly reveal the difference in depth profile between

the two conditions, we calculated a modulation ratio by dividing the rivalry modulation by the

replay modulation. This rivalry/replay modulation ratio was strongest in the superficial layer

(Fig. 1g; main effect of depth, $F(2,22) = 8.118$, $p = 0.009$, $\eta_p^2 = 0.425$; deep vs. middle, $t(11)$

$= -2.403$, $p = 0.035$, Cohen's $d = 0.479$; middle vs. superficial, $t(11) = -2.551$, $p = 0.027$,

Cohen's $d = 0.578$).

One subject also performed multiple sessions of the same Experiment using a passband

bSSFP sequence. Compared to the T2*w GE-BOLD signals, T2w BOLD signals from bSSFP

fMRI are more sensitive to microvasculature activity in the gray matter, which is closer to the

site of neural activity. Previous studies show that T2w BOLD has higher spatial specificity to

reveal the laminar profile of cortical processing (Beckett et al., 2020; Liu et al., 2020; Olman

et al., 2012; Scheffler et al., 2018). Two coronal slices (0.5-mm in-plane resolution with 3-

mm slice thickness) were carefully prescribed to be perpendicular to the calcarine sulcus in

one hemisphere, where the ODCs went approximately parallel with the orientation of 'pencil'

voxels (Fig. 2h). One of the slices shows clear ODC patterns (Fig. 2j), highly reproducible

across sessions (Fig. S2). Eye-specific modulation peaked in the middle layer in the replay

and localizer conditions (Fig. 2k), but in the superficial layer during rivalry. This laminar

pattern is consistent with the GE-EPI data. This finding further supports that interocular

competition in binocular rivalry mainly arises from lateral mutual inhibitions between ODCs

in V1 superficial layers.

Eye-specific rivalry dynamics is more synchronized in V1 superficial and deep layers

Local interocular competition may result in different local winners and piecemeal perception.

It has been hypothesized that feedback signals from higher areas help synchronize and

stabilize local competitions into a globally coherent perceptual state over extended visual

field (Kovács et al., 1996; Tong et al., 2006). To test this hypothesis, we characterized the

synchrony of eye-specific modulations across V1 ODCs by calculating TR-by-TR Pearson

correlations between the ongoing V1 response pattern and the localizer-derived OD pattern

(Fig. 3a). More coherent perception would predict more synchronized OD dynamics with

larger correlation coefficients, quantified by the width of their distribution (O'Hashi et al.,

2018; Omer et al., 2018). As stimuli were fully coherent in the replay condition, this gives us

a benchmark against which to compare the rivalry response patterns at each cortical depth.

Figure 3. Pattern synchronization of eye-specific signals across ODCs. (a) TR-by-TR Pearson correlations

between the OD pattern from the localizer and the real-time V1 response pattern in typical runs of rivalry and

replay. **(b)** The distributions of pattern correlation (r) for different V1 layers in rivalry (solid) and replay (dashed).

Shaded area indicates 95% confidence intervals across subjects. **(c)** The ratio of r distribution widths between

rivalry and replay conditions across V1 layers.

As predicted, we found that the replay condition was associated with a larger distribution

width of correlation coefficients (Fig. 3b). The critical question is, how does the synchrony of

OD dynamics differ across cortical depth during rivalry? If pattern synchronization relies on

feedback signals, its signature would be more evident in the superficial and deep layers.

Indeed, V1 superficial and deep layers showed significantly larger normalized distribution

width than the middle layer (Fig. 3c; main effect of depth, $F(2,22) = 11.789$, $p < 0.001$, $\eta^2_p =$

0.517; deep vs. middle, $t(11) = 3.210$, $p = 0.008$, Cohen's $d = 0.690$; middle vs. superficial,

$t(11) = -4.789$, $p < 0.001$, Cohen's $d = 0.942$). To verify whether this actually reflected

difference in the temporal structure of the response or a mere SNR difference across cortical

depths, we performed a permutation analysis. A GLM with variable duration of perceptual

states was fitted to the vertex timeseries in the rivalry and replay conditions, and the residuals

were temporally permuted independently for each vertex before recombined with the fitted

timeseries. The permutation destroyed any synchronous fluctuation unmodeled in the GLM

without changing the overall SNR across layers, which we reasoned might reduce the

observed laminar difference. As expected, the difference between the deep and middle layers

was largely eliminated for the permuted data (Fig. S3; $t(11) = 0.389$, $p = 0.705$, $BF_{01} = 3.257$,

Cohen's $d = 0.038$) and the main effect of depth was no longer significant ($F(2,22) = 2.115$, p

$= 0.165$, $\eta_p^2 = 0.161$). These results suggest that feedbacks from IPS of the parietal cortex may

be involved to synchronize local interocular competitions in V1 into a coherent visual

representation.

IPS shows stronger eye-specific modulation and increased connectivity to the early visual
cortex during rivalry compared to replay

To identify the source of feedback modulations of rivalry dynamics in V1 ODCs, we further

investigated eye-specific modulations in higher-order cortical areas. Given the critical role of

attention in binocular rivalry and that top-down attention can be eye-specific (Zhang et al.,

2011, 2012), and especially a possible role of the intraparietal sulcus (IPS) region in bi-stable

perception as suggested by TMS studies (Carmel et al., 2010; Kanai et al., 2011; Zaretskaya

et al., 2010), we first examined whether IPS encoded the state of currently dominating eye,

and its relationship with rivalry modulations in V1. To exploit the sensitivity of multivariate

methods, we trained a support vector machine (SVM) to predict the currently stimulated eye

on a TR-by-TR basis using data from our ocular-bias localizer within retinotopic-atlas-

defined IPS (L. Wang et al., 2015). Voxels were selected based on their visual responsiveness

and ocular bias (see methods). The model was then used to predict which eye dominated at

each TR in the rivalry and replay conditions. The distance between each activation pattern

and the SVM decision boundary was used as a graded measure of the internal representation

reflecting eye dominance (Fig. 4a/b). Event-related averages around the time of perceptual

switches showed that, in IPS, significant eye-specific modulation was observed in binocular

rivalry but not in stimulus replay (Fig. 4c/d; cluster-based permutation test). The posterior

(pIPS, IPS0-2 in (L. Wang et al., 2015)) and anterior (aIPS, IPS3-5) portions of IPS showed

similar pattern (interaction not significant, $F(1,11) = 0.238$, $p = 0.635$, $\eta_p^2 = 0.021$). It suggests

that IPS might play a role in resolving interocular conflicts, e.g. by setting a competition bias

to one eye. Indeed, the modulation amplitudes of IPS significantly correlate with those of V1

only in the rivalry condition (Fig. 4e; rivalry $r = 0.778$, $p = 0.003$; replay $r = 0.409$, $p = 0.186$;

and their difference marginally significant, $p = 0.055$ assessed using bootstrap). Note that we

tried to avoid the mere influence of inter-subject variation in switch duration by taking into

account the duration of perceptual states in the GLM for estimating the modulation amplitude.

Figure 4. Eye-specific modulations in IPS and its relationship with the early visual cortex. (a) Schematic about the distance of activation pattern at each time point to the SVM decision boundary for LE vs. RE stimulation. Red (blue) dots denote the activation patterns of TRs when LE (RE) was stimulated. **(b)** The same distances replotted as a time course, i.e., a multivariate differential response. **(c)** Event-related average of the decoding-distance timecourse in IPS for LE (solid) and RE (dashed) percepts in rivalry (orange) and replay (green). **(d)** Differential waveforms between LE and RE events. The gray bars indicate a significant difference between rivalry and replay. Shaded area indicates SEM. Light and dark gray bars in (c) and (d) denote time points with significant difference from zero for corresponding conditions before and after multiple testing correction, respectively. **(e)** Inter-subject correlation between eye-specific modulations in IPS and V1. Shaded area indicates 95% confidence interval of the linear fit. **(f)** Changes in effective connectivity among V1, V2, and IPS during rivalry compared to replay. Numbers beside the connections denote the estimated modulatory effects in coupling strength, and the line thickness indicate the z value. In-significant connections ($Pp < 0.95$) were rendered as faint, dashed lines. Inset: DCM model specification for effective connectivity between IPS, V2 and V1. Green arrow: Eye-specific driving input for both rivalry and replay. Orange arrows: additional eye-specific driving inputs for rivalry only. Orange dots: eye-specific modulatory effects of rivalry.

Eye-specific rivalry modulation was also found in V2 of the extrastriate cortex (Fig. S4). To further investigate the causal relationship of eye-specific activity between V1, V2 and IPS,

we performed a dynamic causal modeling (DCM) analysis on the multivariate projected timeseries that best discriminated LE from RE perception based on SVM trained on localizer runs. In the full DCM model (Fig. 4f), V1 receives eye-specific driving inputs in both rivalry and replay conditions. Intrinsic or fixed connections were defined between and within cortical areas. The between-area connections as well as each node could be modulated or driven by an additional eye-specific input in rivalry but not in replay. This input thus captured the difference between the two conditions. The full DCM model was first estimated for each individual (Zeidman, Jafarian, Corbin, et al., 2019). For group-level analysis, we used Parametric Empirical Bayes (PEB), Bayesian model reduction, and Bayesian model average to make inference about the model parameters (Friston et al., 2016; Zeidman, Jafarian, Seghier, et al., 2019). Compared to stimulus replay, binocular rivalry significantly increased the feedback connectivity from IPS to V2 and from V2 to V1 (posterior probability (Pp) = 0.997 and 1.000, respectively), whereas decreased the feedforward connectivity from V1 to V2 (Pp = 0.999) and the driving input to V1 (Pp = 1.000). These findings support IPS as the source of feedback processes in resolving and synchronizing interocular competitions in V1 ODCs.

No evidence for eye-specific rivalry modulation in ocular-biased clusters of the LGN

In Experiment 2, to investigate whether the LGN was involved in interocular competition during binocular rivalry, we used a 1.2-mm isotropic GE-EPI sequence to study ocular-layer selective signals in the LGN. Stimuli and procedures were similar as in Experiment 1, fMRI slices were orientated to cover both LGN and V1. Robust ocular-biased patterns were clearly and consistently revealed in the LGN across sessions in separate days by the ocular-bias localizer (Fig. 5a/b for S05; r = 0.862, p < 0.001, Monte Carlo test, see Fig. S3 for the ocular-biased pattern of all participants). There are two ocular-biased clusters for each LGN: a ventromedial one biased to the ipsilateral eye, and a dorsolateral one biased to the contralateral eye, which replicated the findings of our recent study (Qian et al., 2020). Based on the simulation analysis of the previous study, these ocular-biased clusters were the result of BOLD blurring and fMRI down-sampling of the LGN laminar pattern (shown here in the lower right of Fig. 5a). The contralateral-eye biased cluster located at the anatomical boundary of the LGN is simply due to the blurring effect of BOLD point-spread-function. For more details about the simulation analysis, please refer to figure 1b and figure 3a in our previous study (Qian et al., 2020).

Figure 5. Eye-specific modulation in ocular-biased clusters of the LGN. (a, b) Highly reproducible ocular-biased clusters in the LGN of a representative subject (S05). The LE-RE beta maps were thresholded at p < 0.01 uncorrected. See Fig. S3 for ocular-biased patterns of the LGNs for all participants. The lower right panels of (a) show the simulation results of ocular-biased clusters, reproduced from figure 3a in (Qian et al., 2020). **(c)** Eye-specific modulation of BOLD signals in the ocular-biased clusters of the LGN. **(d)** Differential timecourses between preferred and non-preferred perceptual events in rivalry and replay conditions, averaged across LE- and RE-biased voxels. Horizontal bars in light and dark color denote time points with significant difference from zero for corresponding conditions before and after multiple testing correction, respectively.

To our surprise, although BOLD signals in ocular-biased clusters of the LGN showed a transient increase after perceptual switch (Fig. 5c, mean response to both preferred and non-preferred switches averaged between 2-6 s, $t(14) = 4.112$, $p = 0.001$, Cohen's $d = 1.062$), there was no significant eye-specific modulation during binocular rivalry (Fig. 5d; $t(14) = -0.619$, $p = 0.546$, $BF_{01} = 3.226$). As positive controls, the differential response to preferred- and non-preferred eye stimulation averaged between 4-12 s was as strong in LGN as in V1 in the replay condition, and V1 showed robust eye-specific modulations in both conditions (Fig. 5d). Although there was no evidence of eye-specific rivalry modulation in the ocular-biased clusters (Fig. S6). This negative effect is likely due to attentional suppression outside the stimulus region (Shmuel et al., 2002; Tootell et al., 1998). No significant eye-specific modulation was found in the pulvinar ($t(14) = 1.253$, $p = 0.231$, Cohen's $d = 0.323$).

Therefore, our 7T fMRI results suggest weak, if any, eye-specific rivalry modulation within the stimulus regions of the LGN. This finding is consistent with the single-unit study

of macaque LGN (Lehky & Maunsell, 1996), and also with the observation in Experiment 1

that eye-specific rivalry modulation peaked in V1 superficial layers. If interocular

competition of binocular rivalry was resolved in the LGN, one would expect a peak of eye-

specific modulation in the middle layer of V1 that receives thalamic input. Also, it is unlikely

that feedback processes in binocular rivalry can modulate ocular-layer selective activity in the

LGN. Nonetheless, it remains possible that LGN activity might be modulated by perception-

related feedback processes in a non-eye-specific manner (Wunderlich et al., 2005). To test

this hypothesis and to investigate the whole-brain network of perceptual rivalry, we did

Experiment 3.

Robust perceptual rivalry modulation in P subdivisions of the LGN and ventral pulvinar

Figure 6. Effect of perceptual rivalry between chromatic and achromatic gratings in the LGN and ventral

pulvinar. (a) The M-biased achromatic grating and P-biased chromatic grating in Experiment 3. **(b)** Upper: M-

and P-biased voxels in the LGN and vPulv of a representative subject (S02). These voxels were defined based on

their P-M response bias in the functional localizer and anatomical locations. Voxel colors indicate the P-M

response from the functional localizer. Lower: Group probability map of M and P ROIs in MNI space. Voxel

colors indicate the number of overlapped ROIs (Map threshold: LGNp=11, LGNm=5, Pulp=3, Pulm=3). Black

dotted lines indicate the anatomical boundaries of LGN and pulvinar. **(c, d)** Event-related BOLD timeseries in M-

and P-biased voxels of the LGN and ventral pulvinar. Shaded areas represent SEM across subjects. Light and dark

gray bars denote time points with significant difference from zero for corresponding conditions before and after

multiple testing correction, respectively.

In Experiment 3, we exploited the spatial segregation of magnocellular (M) and parvocellular

(P) pathways in subcortical (e.g., LGN, ventral pulvinar) and cortical (e.g, MT+, hV4) visual

areas as a novel tagging method to study the effect of perceptual rivalry across the whole

brain. According to the laminar organization of the LGN, M and P layers are located in the

ventral and dorsal portions of LGN, respectively. In the ventral pulvinar (vPul), the

parvocellular lateral portion reciprocally connects with the early visual cortex and ventral

visual stream, while the magnocellular medial portion receives input from the superior

colliculus (SC) and connects with dorsal visual stream such as area MT (Arcaro et al., 2015;

Bridge et al., 2016; Kaas & Lyon, 2007).

We designed M- and P-biased visual stimuli to preferentially activate the M and P visual

pathways (Derrington & Lennie, 1984; Wiesel & Hubel, 1966). The M stimulus was a low

spatial frequency (0.5 c.p.d.) achromatic grating (30% or 50% contrast), while the P stimulus

was a red/green equiluminant chromatic grating presented at high contrast (Fig. 6a). M and P

gratings were dichoptically presented in orthogonal orientations to the two eyes in binocular

rivalry, and presented to the two eyes in alternation in simulated replay. An independent

localizer was used to measure the M-P bias of voxels (see methods for details). In localizer

runs, the M-biased achromatic stimulus evoked stronger responses in the medial portion of

vPul and in the ventral LGN, while the P-biased chromatic stimulus more strongly activated

the lateral portion of vPul and the dorsal LGN (Fig. 6b), consistent with the known functional

organizations of these thalamic nuclei. The M and P stimuli also selectively activated the

dorsal and ventral cortical visual streams (Fig. S9a).

During binocular rivalry, BOLD signals in P-biased voxels of the LGN increased when

subjects reported seeing the chromatic grating, and decreased when the achromatic grating

took dominance (Fig. 6c; Fig. 7a, $t(12) = 2.993$, $p = 0.011$ Holm corrected, Cohen's $d =$

0.830). Similar modulation was also observed in the replay condition (Fig. 7a, $t(12) = 3.892$,

$p = 0.004$, Cohen's $d = 1.079$). For the M-biased voxels, there were marginally significant

stimulus-driven modulations in the replay condition (Fig. 7b, $t(12) = 1.460$, $p = 0.085$

uncorrected, Cohen's $d = 0.405$), but without significant perceptual modulation during

binocular rivalry (Fig. 6c, Fig. 7b). The different involvement of LGNp and LGNm in

perceptual rivalry were further supported by a significant interaction between condition and

pathway ($F(1,12) = 5.339$, $p = 0.039$, $\eta_p^2 = 0.308$). Interestingly, the rivalry/replay

modulation ratio in LGNp was significantly larger than V1p (condition*ROI interaction,

$F(1,12) = 7.518$, $p = 0.018$, $\eta_p^2 = 0.385$), whereas comparable with hV4 (interaction not

significant, $F(1,12) = 0.365$, $p = 0.557$, $\eta_p^2 = 0.030$), suggesting that the perceptual rivalry

modulation in LGNp might be related to feedback signals from higher-order brain areas.

BOLD signals in both P- and M-biased voxels of the ventral pulvinar showed significant

perceptual modulations during binocular rivalry (Fig. 6d; Fig.7a for Pulp, $t(12) = 3.347$, $p =$

0.009, Cohen's $d = 0.928$; Fig. 7b for Pulm, $t(12) = 2.079$, $p = 0.030$ uncorrected, Cohen's $d =$

0.576), whereas only marginally significant for stimulus-driven modulations of P-biased

voxels in simulated replay ($t(12) = 1.596$, $p = 0.068$ uncorrected, Cohen's $d = 0.443$). No

significant interaction was found between condition and pathway ($F(1,12) = 0.072$, $p = 0.793$,

$\eta_p^2 = 0.006$). Searchlight decoding also revealed significant activations in the ventral pulvinar

in both rivalry and replay conditions (Fig. S9c). We did not find significant perceptual state-

related modulation in the SC in either rivalry or replay (Fig. S8c), thus SC data was not

included in the following analysis.

Figure 7. Hierarchical whole-brain network of perceptual rivalry. (a-c) Normalized rivalry and replay
modulations in the P and M visual pathways and frontoparietal areas. **(d)** Effective connectivity between aIPS, V4,
ventral pulvinar and V1. **(e)** Hierarchical clustering of rivalry and replay modulations in cortical and subcortical
areas. **(f)** 2D visualization of normalized rivalry and replay modulations of all ROIs.

Perceptual rivalry modulation is weak or absent in early visual areas of the magnocellular

pathway

Robust perceptual rivalry modulation can be found in P-biased voxels of the early visual

cortex (V1p, V2p, V3p, hV4, Fig. 7a). Rivalry modulation was significantly smaller than

replay in V1p ($t(12) = -3.389$, $p = 0.032$, Holm corrected across ROIs, Cohen's $d = 0.602$),

and the ratio between the response amplitudes of the two conditions steadily went up along

the cortical hierarchy (slope = 0.137, [0.051,0.279], p = 0.001), consistent with previous

reports (Leopold & Logothetis, 1996; Mo et al., 2022). In contrast, early stages of the M

pathway showed significant modulations in replay (LGN: $t(12) = -3.180$, $p = 0.008$, Cohen's d

= 0.478; V1: $t(12) = -2.936$, $p = 0.012$, Cohen's $d = 0.896$), but not in rivalry (LGN: $t(12) = -$

0.195, $p = 0.576$, Cohen's $d = 0.054$; V1: $t(12) = -0.437$, $p = 0.665$, Cohen's $d = 0.121$). There

were also significantly stronger modulations in replay compared to rivalry (LGN: $t(12) = -$

3.180, $p = 0.040$, Cohen's $d = 0.478$; V1: $t(12) = -2.936$, $p = 0.049$, Cohen's $d = 0.896$). The

rivalry/replay modulation ratio also increased along the visual hierarchy (V1m, V2m, V3m,

MT+, Fig. 7b), but more than 3-fold faster than the P pathway (*slope* = 0.450, [0.196,1.051],

$p = 0.001$). Rivalry and replay modulations were comparable at high-level areas of the M and

P pathways (i.e. area MT+ and hV4). Therefore, perceptual rivalry modulation was strong in

the P pathway, but weak or absent in early stages of the M pathway.

Robust perceptual rivalry modulation in IPS and pSTS

We further investigated whether perception-state related information is represented in the

frontal and/or parietal association cortices. Since there is no known large-scale segregation of

M/P representations in frontoparietal areas, we again deployed multivariate methods to test if

perceptual content can be decoded TR-by-TR from these high-order brain areas. Regions with

significant visual response in the localizer runs were included (FEF, IFJ, TPJ, pSTS, aIPS,

pIPS, see Fig. S7 for ROI definition). To improve decoding performance in these high-level

brain areas, a cross-validated grid-search approach was used for feature selection based on

visual responsiveness and M/P bias in localizer runs (see methods). The anterior and posterior

portions of IPS were engaged differently in rivalry and replay (Fig. 7c, significant ROI

*condition interaction in normalized modulation, $F(1,12) = 8.385$, $p = 0.013$, $\eta_p^2 = 0.411$).

Modulation amplitudes were comparable between rivalry and replay in pIPS, whereas

marginally larger in rivalry compared to replay in aIPS ($t(12) = 2.057$, $p = 0.062$ uncorrected,

Cohen's $d = 0.367$; most significant around 8 s after perceptual switch, see Fig. S8b). Similar

results were also found in pSTS, showing significant modulation in rivalry ($t(12) = 3.680$, $p =$

0.016, Cohen's $d = 1.021$), but not in replay ($t(12) = 2.584$, $p = 0.076$, Cohen's $d = 0.717$).

These findings are consistent with the stronger eye-specific rivalry modulation in IPS found

in Experiment 1 (Fig. 4). No reliable above-chance decoding was observed in the frontal lobe.

Searchlight decoding also revealed significant activations in the parietal lobe during binocular

rivalry (Fig. S9b). Therefore, the perceptual states in binocular rivalry can be decoded from

parietal but not frontal activity in our data.

IPS and ventral pulvinar show stronger perception-state related connectivity to the early

visual cortex in rivalry compared to replay

To investigate the casual relationship of perceptual rivalry modulation in a whole-brain

network, we performed an effective connectivity analysis using DCM within a minimal

network containing multi-level visual areas (V1, V4), and high-order parietal cortex (aIPS)

and ventral pulvinar (Pul). Here we selected aIPS as the potential source of feedback signals

because it showed stronger modulation in rivalry compared to replay. Similar to the eye-

specific connectivity in Experiment 1, binocular rivalry showed an enhanced feedback

connectivity from aIPS to V4 ($Pp = 0.966$) and from V4 to V1 ($Pp = 1.000$) compared to

stimulus replay. In addition, connections from Pul to V1 and V4 were also significantly

stronger during rivalry compared to replay ($Pp = 0.995$ and 1.000, respectively). Feedforward

connections were significantly weaker during rivalry from V1 to V4 and Pul ($Pp = 1.000$ for

both V1-V4 and V1-Pul connections). These results suggest aIPS as the potential source of

feedback signals to the early visual cortex to help resolve conflicts, e.g., by setting a bias in

the competition. Meanwhile, pulvinar may regulate the feedback connectivity across the

hierarchy.

Hierarchical whole-brain network of perceptual rivalry

Finally, to further illustrate the relationship of perceptual rivalry in cortical and subcortical

areas, we performed a clustering analysis of rivalry and replay modulations across different

brain areas. Each ROI was represented as a vector comprising normalized modulation

amplitude during rivalry and replay for each subject, which captured both similarity in

modulation in different conditions and inter-subject correlation between regions. Hierarchical

clustering procedure based on Euclidean distance suggested the ROIs were best described as

3 groups (Fig. 7e), which was visualized by plotting the ROIs in a 2D-plane according to their

normalized modulation amplitude in rivalry and replay (Fig. 7f), along with eye-specific

modulation of LGN, V1, and IPS in Experiment 1 & 2. The first group, comprising mainly

the early stages of the M pathway (LGNm, V1m, V2m), was characterized by the near

absence of rivalry modulation and robust replay modulation. Thus, low-level areas of the M

pathway are stimulus driven but modulated little by perceptual rivalry. The second group,

including cortical areas of the P pathway and higher levels of the M pathway (V1p, V2p, V3p,

hV4, V3m, MT+), featured more or less similar modulation in both conditions. Perceptual

modulation in this group became increasingly indistinguishable with stimulus-driven response

along the visual hierarchy. The third group tended to show stronger modulation during

binocular rivalry, mainly composed of cortical and subcortical attention network (aIPS, pIPS,

pSTS, TPJ, FEF, IFJ, Pulp, Pulm). Rivalry modulations in these areas may partly reflect

processes engaged in resolving competition between conflicting representations. Interestingly,

the earliest stage of the P pathway (LGNp) also belonged to this group, presumably due to

perceptual feedback modulation. These analyses summarize the hierarchical whole-brain

network of perceptual rivalry, which may suggest distinct roles of sub-networks in

visuosensory processing, perceptual representation and conflict resolution.

Discussion

Here we studied hierarchical neural mechanisms of binocular rivalry in human cortical and

subcortical areas using high-resolution fMRI at 7 T. Experiment 1 and 2 investigated the

neural circuitry of interocular competition underlying binocular rivalry. Results show that

eye-specific rivalry modulation was strongest in the superficial layers of V1 ODCs, and more

synchronized in the superficial and deep layers. IPS generated stronger eye-specific

modulation and increased feedback connectivity to V1 during rivalry compared to replay.

Ocular-layer selective activity of the LGN showed no evidence of eye-specific rivalry

modulation. Experiment 3 used M- and P pathway-selective visual stimuli to further

investigate hierarchical neural processes of perceptual rivalry over the whole brain. Robust

perceptual rivalry modulation was found in P- and not M-biased voxels in the geniculostriate

pathway. IPS and ventral pulvinar showed robust perceptual modulation in rivalry and

increased connectivity to the early visual cortex compared to replay.

Interocular competition arises from lateral mutual inhibition between V1 ODCs

Horizontal connections between orientation-selective neurons are most prominent in layers

2/3 of V1 (Angelucci et al., 2017; Gilbert & Wiesel, 1983). Inhibitory large basket cells

spanning ODCs were found in layer 3 of cat V1 (Buzs et al., 2001). Dichoptic cross-

orientation suppression is strongest in the superficial layers of V1 in both anesthetized cats

and awake monkeys (M. A. Cox et al., 2019; Sengpiel et al., 1995). According to these

studies, our findings that eye-specific rivalry modulation was strongest in the superficial

layers of V1 ODCs (Fig. 2f) but was weak or absent in the LGN (Fig. 5d), clearly support that

local interocular competition in binocular rivalry arises from lateral mutual inhibition

between V1 ODCs. The eye-specific effect in V1 was unlikely driven by pulvino-cortical

feedback targeting the superficial layers (Shipp, 2003), since we found no eye-specific

modulation in the pulvinar. However, our data suggest that cortico-cortical feedbacks could

play roles to resolve and synchronize interocular competition. Eye-specific rivalry dynamics

was more coherent in V1 superficial and deep layers (Fig. 3c), and IPS showed stronger eye-

specific modulation and connectivity to the early visual cortex during rivalry compared to

replay (Fig. 4). These findings suggest that eye-specific feedback from IPS might help to

resolve and synchronize local interocular competitions in V1 into a coherent perceptual

representation. Our results also support that the discrepancy between single-unit and

LFP/fMRI results of perceptual suppression could be due to feedback modulation of synaptic

input (Maier et al., 2008).

With ocular-layer selective activity robustly resolved with 7T fMRI (Fig. 5a, Fig. S5),

we found no evidence of eye-specific rivalry modulation in the stimulus region of the LGN

(Fig. 5d). This finding is consistent with electrophysiological studies showing no effect of

binocular rivalry on the spike rate of LGN neurons in alert monkeys (Lehky & Maunsell,

1996). However, a 3T fMRI study found significant eye-specific effect of binocular rivalry in

ocular-biased voxels of the human LGN (Haynes et al., 2005). What might underlie this

discrepancy? In the previous study, a larger stimulus (bilaterally presented 120-deg wedges

subtending from 1.5 to 7.5 deg of eccentricity, compared to the 0.5 to 4.5 deg disc in the

current study) was used to map the ocular bias of voxels from more peripheral visual field.

Given that the temporal-nasal asymmetry of attention was more pronounced in the peripheral

visual field (Rafal et al., 1991) and that attention can strongly modulate LGN activity

(McAlonan et al., 2008; O'Connor et al., 2002), the eye-specific effect might be a result of

attentional modulation due to temporal-nasal asymmetry. Consistent with this hypothesis, we

also found an eye-specific suppression effect outside the ocular-bias clusters (Fig. S6), likely

due to attentional suppression outside of the stimulus region (Shmuel et al., 2002; Tootell et

al., 1998). Therefore, although binocular suppression exists in the primate LGN (Kacie

Dougherty et al., 2021; Schroeder et al., 1990), binocular rivalry does not strongly modulate

the ocular-layer selective activity.

Parvocellular feedback to the LGN may serve as a thalamic gatekeeper of perception-related

signals

Although we found no evidence of eye-specific rivalry modulation in the LGN, perceptual

rivalry between chromatic and achromatic gratings strongly modulated LGN activity in its

parvocellular subdivision (Fig. 6c). Modulation amplitudes were comparable between rivalry

and replay, similar to the higher-order visual areas such as V4 (Fig. 7a). Moreover,

hierarchical clustering analysis revealed closer relationship of rivalry modulations in the LGN

with those in high-order brain areas (Fig. 7e). Therefore, the effect of perceptual rivalry in the

LGN should be a result of feedback modulation, likely through pathway-specific

corticogeniculate connections (Briggs & Usrey, 2011). Since perceptual rivalry selectively

modulated P but not M responses, this effect cannot be explained by attentional modulation

that would influence both M and P activity (Schneider, 2011; Schneider & Kastner, 2009).

With exogenous flash suppression, Wilke and colleagues found an effect of perceptual

suppression on low-frequency LFP power in the LGN of alert monkeys (Wilke et al., 2009).

However, this effect might be related to the disengagement of attention upon target

disappearance. Another 3T fMRI study found a correlation of LGN activity with contrast perception during binocular rivalry between a pair of low and high contrast gratings (Wunderlich et al., 2005). However, this effect could also be explained by a multiplicative effect of attention on contrast responses. Also given the relatively low fMRI resolution (3-mm isotropic), LGN signals in this study might be influenced by those of the ventral lateral pulvinar due to partial volume effect. Therefore, our results are consistent with these previous studies that perceptual state can strongly modulate LGN activity, and further suggest that parvocellular feedback to the LGN might serve as a gating mechanism for conscious perception-related processing at the thalamic level.

Binocular rivalry mainly occurs in the parvocellular visual pathway

Stronger perceptual rivalry in the P than the M pathway (Fig. 7a/b) provides direct neuroimaging evidence that binocular rivalry is primarily a P-pathway phenomenon. This finding is consistent with psychophysics studies showing weak rivalry suppression of M-biased visual stimuli (He et al., 2005). The dissociation of M and P pathways in binocular rivalry can be understood in terms of their distinct functional roles and neurophysiological properties. The highly sensitive and transient nature of the M pathway supports fast detection of and immediate action to potentially important events, functions that may not require consciousness or the relatively slow feedback processes. In contrast, the function of the P pathway is to encode spatial details and color information for accurate object recognition, which may require an active inferential process with feedback modulation to refine information coding. Different involvement of parallel pathways in binocular rivalry could be also due to their differences in neural circuitries of binocular interactions. Recent studies show that binocular facilitation occurs at low contrast level (M-biased) and at an early stage of visual processing, while binocular suppression occurs at high contrast level (P-biased) and at a later stage (K Dougherty et al., 2019; Mitchell et al., 2022).

Perceptual state-related feedback from IPS resolves visual competition in the early visual cortex

Previous studies mainly focused on the role of frontoparietal attention network in perceptual transitions of bi-stable perception (Jan Brascamp et al., 2018). Due to technical limitations in resolving weak and possibly finescale perceptual representations in frontoparietal regions, whether frontoparietal activity represents perceptual state in bi-stable perception remains unclear. Our 7T fMRI results revealed robust perceptual state representations in IPS during binocular rivalry (Fig. 7c), and increased feedback connectivity to the visual cortex in rivalry compared to replay (Fig. 7d), suggesting that perceptual state-related feedback from IPS played roles to resolve visual competitions in the early visual cortex. Since we trained the

state classifier based on data from the functional localizer in which subjects performed a central fixation task, the state-related activity in IPS was unlikely due to active reports. In addition, IPS represented perceptual eye dominance in binocular rivalry but not stimulus eye-of-origin in simulated replay (Fig. 4c). Since subjects were not aware of the eye-of-origin information, the role of IPS should be resolving visual competitions in the early visual cortex rather than representing the content of visual consciousness. Given the critical role of IPS in attention, the current findings are also consistent with previous studies that binocular rivalry requires attention (J. W. Brascamp & Blake, 2012; Zhang et al., 2011), but not awareness of interocular conflict (Xu et al., 2016; Zou et al., 2016).

Ventral pulvinar regulates perceptual state-related feedback connectivity across hierarchy
The parallel pathway-selective visual stimuli selectively activated the lateral (P) and medial (M) subdivisions of ventral pulvinar (Fig. 6b). BOLD responses in both subregions significantly correlated with conscious perception during binocular rivalry (Fig. 6d, Fig. 7b). The stimulus-specific perceptual modulation cannot be explained by a non-specific effect of spatial attention. This poses an advantage over the study by Wilke et al., where target disappearance was induced by flash suppression, potentially inducing a non-specific effect of attention. However, it remains possible that perceptual modulations in the ventral pulvinar could be related to feature-based attention. Similar to IPS, pulvinar showed robust perceptual modulation during rivalry and stronger connectivity to the visual cortex in rivalry compared to replay. Therefore, our findings support a critical role of ventral pulvinar in generating conscious visual perception, which could be regulating the feedback connectivity from IPS to visual cortex and support cortical computations to resolve visual competition.

Conclusions

The current study revealed the most complete picture so far about how binocular rivalry is resolved in the human brain. Binocular rivalry arises from interocular competition between ocular dominance columns through lateral mutual inhibition in V1 superficial layers. Feedback modulations from IPS further resolve and synchronize visual competitions in the visual cortex into a coherent perceptual representation. The ventral pulvinar serves as the network hub regulating the feedback connectivity from IPS to visual cortex to resolve perceptual conflicts. Finally, parvocellular feedback to the LGN might serve as a gatekeeper of perception-related signals at the thalamic level. These findings elucidate the functional roles of major brain areas involved in binocular rivalry, and provide important insights about how hierarchical processes in our brain resolve visual competitions to generate conscious perception.

Methods and Materials

Participants

Sixteen healthy volunteers (seven females, age 22–40 years) participated in Experiment 1.
Three of them were excluded due to lack of clear OD pattern in V1, and one subject was
excluded due to excessive bias toward one eye. Fifteen subjects (seven females, age 22–39
years) participated in Experiment 2. Sixteen subjects (seven females, age 22–41 years)
participated in Experiment 3. One subject was excluded due to response box failure, and two
were excluded due to lack of significant M- or P-biased voxels in ventral pulvinar. All
observers had normal or corrected-to-normal vision and gave written informed consent.
Experimental protocols were approved by the Institutional Review Panel at the Institute of
Biophysics, Chinese Academy of Sciences.

Stimuli and procedures

Figure 1 shows the stimuli and procedures for Experiment 1 and 2. For the ocular-bias
localizer, to selectively activate V1 ODCs and LGN ocular layers, a high contrast
checkerboard (1 deg check size, ~9 deg in diameter) counterphase flickering at 8 Hz was
monocularly delivered to the left or right eye in alternating 24-s blocks. Two 24-s fixation
blocks were included at the beginning and the end of the localizer run. The checkerboard
slowly rotated in 3.75 degrees steps every second to reduce adaptation. Subjects viewed the
dichoptic stimuli in the scanner with prism glasses and a cardboard divider, and reported
occasional fixation-size changes. During binocular rivalry, red and green gratings (0.8
cycle/deg) in orthogonal orientations were dichoptically presented to the two eyes. The
gratings rotated at 0.67 round/s to prevent adaptation. The association between color and eye
swapped every run so that each eye was not bound to a particular color. Subjects continuously
reported whether they were seeing red, green, or a mixed percept using three buttons. In
replay runs, the perception and timing of the previous rivalry run were simulated with
physically alternating monocular stimulus. The transition was simulated as a blurred and
alpha blended boundary rotating and swiping across the grating, gradually revealing the
stimulus from the other eye. Each run for the rivalry or replay condition lasts 260 s. Each run
for the ocular-bias localizer lasted 340 s. Subjects scanned 4-6 localizer runs, 4 rivalry runs,
and 4 replay runs in a single session. For the bSSFP scans of S06 in Experiment 1, 6 localizer
and 6 rivalry runs were collected in a first session, followed by 6 localizer and 6 replay runs
in a second session on a different day.

In Experiment 3, achromatic and chromatic gratings were designed to preferentially
activate the M or P pathway while remaining nearly balanced during rivalry. In localizer runs,

the M-biased stimulus was a low contrast (30%, and for some subjects 50% if the dominant

duration for the M stimulus was too short during rivalry in pilot runs), luminance-defined

sinewave grating (0.5 cycles/deg), counterphase flickering at 10 Hz; the P-biased stimulus

was a red/green isoluminant chromatic grating (0.5 cycles/deg), counterphase flickering at 4

Hz. The isoluminance of the red/green colors and the gray in background was adjusted with a

minimal-flicker procedure. The stimulus was delivered in 16-s blocks, interleaved with 16-s

fixation period. In each block, one grating was monocularly presented to either left or right

eye, and its orientation changed 45 degrees every 2 s (counterclockwise or clockwise in

separate blocks). Each localizer run lasted 336 s. During binocular rivalry, the chromatic and

achromatic gratings were dichoptically presented in orthogonal orientations, continuously

rotating at 1 round/s without flickering. Subjects pressed one of three buttons to indicate

chromatic, achromatic or mixed perception, and the timings were used in replay runs to

simulate the perception during binocular rivalry. Each run for the rivalry or replay condition

lasted 300 s.

MRI data acquisition

MRI data were acquired with a 7T scanner (Siemens Magnetom) using a 32-channel receive

single-channel transmit head coil (NOVA medical), from the Beijing MRI Center for Brain

Research (BMCBR). A bite bar was used to reduce head motion. In Experiment 1, T2*w

BOLD signals from early visual cortex and parietal cortex were acquired with a 2D GE-EPI

sequence (0.8 mm isotropic, 31 oblique-coronal slices, FOV = 128×128 mm, TE = 23 ms, TR

= 2000 ms, nominal flip angle = 80°, bandwidth = 1157 Hz/pixel, partial Fourier = 6/8,

GRAPPA = 3). The author C.Q. was also scanned with a 2D passband bSSFP sequence to

acquire T2w BOLD signals (voxel size 0.5 × 0.5 × 1.5 mm, 2 oblique-coronal slices, FOV =

96 × 96 mm, volume acquisition time = 2400 ms for localizer and 1600 ms for rivalry/replay,

TR = 5.64 ms, TE = 2.82 ms, nominal flip angle = 29° or 30°, bandwidth = 521 Hz/pix,

GRAPPA = 0 for localizer and 2 for rivalry/replay). The same 2D GE-EPI sequence was used,

albeit with different parameters in Experiment 2 (1.2 mm isotropic voxels, 31 oblique-

transversal slices, FOV = 180×180 mm, TE = 22 ms, flip angle = 78°, bandwidth = 1587

Hz/pix, GRAPPA = 2) and Experiment 3 (1.5 mm isotropic voxels, 68 oblique-transversal

slices, FOV = 183×183 mm, TE = 21.6 ms, bandwidth = 1576 Hz/pix, GRAPPA = 2,

multiband = 2). EPI volumes with reversed phase encoding and readout directions were also

acquired for susceptibility distortion correction. For all Experiments, T1w anatomical

volumes were acquired using a MP2RAGE sequence (0.7 mm isotropic voxels, FOV =

224×224 mm, 256 sagittal slices, TE = 3.05 ms, TR = 4000 ms, TI1 = 750 ms, flip angle = 4°,

TI2 = 2500 ms, flip angle = 5°, bandwidth = 240 Hz/pix, partial Fourier = 7/8, GRAPPA = 3).

MRI data analysis

Preprocessing

MRI data were preprocessed using AFNI (R. W. Cox, 1996), FreeSurfer (version 6.0) (Fischl, 2012), ANTs (Avants et al., 2011), and the mripy package developed in our lab (https://github.com/herrlich10/mripy). EPI volumes were corrected for slice timing, susceptibility distortion (blip-up/down method), head motion (6 parameters rigid body), and rescaled to percent signal change before GLM analysis. To minimize the loss of spatial resolution, all spatial transformations were combined and applied in a single interpolation step (Sinc method), in which the data were also up-sampled by a factor of 2 (J. Wang et al., 2022). For the bSSFP data, motion correction was performed in-plane with three free parameters (in-plane rotation and translation) within the central image region that was free of aliasing. Susceptibility distortion correction was safely omitted due to the very low distortion of bSSFP images. A canonical HRF (BLOCK4 in AFNI) was used for both cortical and subcortical ROIs in GLM. The anatomical volume as well as the reconstructed surfaces were aligned to the mean of preprocessed EPI images.

To alleviate bias created by surface veins in laminar analysis, surface vertices with excessively high BOLD signal change (mean stimulus-driven response over 5% in localizer runs) or low EPI intensity (below 75% of the mean EPI intensity) were classified as veins and the corresponding column of voxels were excluded from further analysis. The column-wise voxel exclusion and ROI selection (see below) were used to balance the number of voxels from different depths in laminar analysis, aiming for a within-column comparison in activation profile in Experiment 1.

In Experiment 3, to increase the power for detecting P-biased clusters in subcortical regions where SNR was relatively low, within-ROI smoothing was performed within anatomical masks of the LGN and pulvinar (3dBlurInMask in AFNI, FWHM = 3 mm) before GLM analysis. M-biased voxels were defined with unsmoothed data because they were expected to locate in thin structures that were close to P-biased voxels, and might easily be contaminated or even overshadowed by their P neighbors.

Surface segmentation and depth estimation

The T1w MP2RAGE anatomical volume was segmented into white matter (WM), gray matter (GM), and cerebrospinal fluid (CSF) using FreeSurfer (version 6.0)'s automated procedure, using its high-resolution option (-*hires*). The results of initial segmentation were visually inspected and manually edited to eliminate dura matter, sinus, etc., ensuring correct GM boundaries. To match the up-sampled volume grid and to alleviate the vertex-missing problem during surface-volume projection, high density surface meshes (Polimeni et al., 2017)

were created by subdividing each triangular face into 4 smaller ones at the midpoint of each

edge and repeated twice (getting 16 small triangles).

The relative cortical depth for each voxel was estimated using the equivolume method

(Waehnert et al., 2014), which we implemented in mripy package. The neighborhood area for

a pair of nodes on the pial or smoothwm surface was approximated by summing up the area

of all triangular faces surrounding the vertex on the corresponding surface mesh. A set of

intermediate surfaces on specified equivolume depths were then generated according to

Equation 10 in (Waehnert et al., 2014). Finally, voxel depth was computed by interpolating

between two nearest equi-depth surfaces. The pial surface was defined to have a zero depth,

and the relative cortical depth for the boundary between white and gray matter was one. The

deep, middle, and superficial layers were defined to take up 30%, 35%, and 35% of the

cortical thickness, respectively (Balaram et al., 2014; de Sousa et al., 2010; Liu et al., 2020).

For the eye-specific pattern analysis in Fig. 3, we used 3dVol2Surf in AFNI when projecting

volume data onto the surface, which employed an equidistance algorithm. According to

previous studies (Liu et al., 2020; Renzo et al., 2021; Waehnert et al., 2014), equivolume and

equidistance estimates of cortical depth showed only mild differences in the final results.

ROI definition

In Experiment 1, V1 ROIs were manually drawn on the cortical surface to select vertices with

a clear and roughly balanced pattern of ODCs (see Fig. S1 for the OD pattern and ROIs of all

subjects). Vertices with significant ocular bias (LE-RE t > 2 for LE-biased columns, and t < -

2 for RE-biased columns) were then projected to the volume space to select voxels in a

column-wise manner. IPS was defined as the union of IPS0 to IPS5 in Wang15 atlas (L.

Wang et al., 2015), whose masks were generated using the neuropythy package (N C Benson

et al., 2018), whereas pIPS and aIPS was defined as IPS0-2 and IPS3-5, respectively.

In Experiment 2, two clusters of voxels with significant ocular-bias were identified for

each LGN (Fig. S5 shows the ocular-biased clusters of LGNs for all subjects). V1 voxels with

significant ocular bias and positive visual response were included for ROI analysis.

In Experiment 3, cortical ROIs were first defined as a coarse mask on the cortical surface,

then projected back to the native voxel space, and finally refined based on M-P bias and

stimulus responsiveness derived from GLM of the functional localizer runs. Surface masks

for the early visual cortices (V1, V2, V3, and hV4) were corresponding brain areas in the

Benson14 atlas (N C Benson et al., 2018; Noah C. Benson et al., 2014). The surface mask for

MT+ was manually drawn on the native mesh around the main M-biased cluster (1st-level M-

P contrast t > 2, or for some subjects as low as 1, FWHM = 3 mm smoothed on the surface)

within TO of the Benson14 atlas. After surface-to-volume projection, only voxels with

significant M-P bias and positive M/P response were kept as the M/P subdivision of these

early visual areas (for V1m, V2m, V3m, MT+: M-P t > 2 and M beta > 0; for V1p, V2p, V3p,

hV4: M-P t < -2 and P beta > 0). For the frontoparietal areas (IFJ, FEF, TPJ, pSTS, aIPS and

pIPS) where M-P bias were not obvious, surface masks were manually defined on the

standard surface (std141 in AFNI/SUMA) by encircling major clusters on the group-level t

map of M+P response (t > 0.5, FWHM = 4 mm, Fig. S7). A cross-validated feature-selection

procedure was then used for each individual to select the most relevant voxels that

discriminated M from P stimulation (see below). For subcortical areas (LGN, vPul, and SC),

group-level anatomical masks were first manually delineated on a symmetric T1w template in

MNI space (Pauli et al, 2018) by an experienced Experimenter, and then nonlinearly

transformed to each subject's native space using ANTs (Avants et al., 2011). The individual

masks were carefully inspected and adjusted based on T1w MP2RAGE images, and were

used as the anatomical reference for cluster selection and the mask for within-ROI smoothing

(3dBlurInMask in AFNI) to reduce signal crosstalk between neighboring nuclei. The LGN P-

biased voxels were selected from its dorsal part (M-P t < -2 and P > 0, 3-mm FWHM blur-in-

mask to increase SNR), and the M-biased voxels from the ventral part (M-P > 0 and M t > 1,

no smoothing to reduce the influence from P signals nearby). For the ventral pulvinar, P- and

M-biased voxels were selected from its lateral and medial portions, respectively, using similar

1st-level contrasts and thresholds as the LGN. The ROI for the SC was defined as visually

responsive voxels in the anatomical mask (M+P t > 1).

Univariate and multivariate differential response

To quantify eye-specific modulations, we computed the differential response between LE and

RE ODCs in assessing the laminar perceptual modulation and in the DCM analysis. This was

implemented by selecting V1 surface regions that contained visually balanced and

interdigitated OD stripes as ROI. For univariate analysis, estimating perceptual modulation

(see below) separately for LE and RE ODCs and taking their average, is equivalent to

computing perceptual modulation of the differential response:

$$(LE_{L\,event} - LE_{R\,event}) + (RE_{R\,event} - RE_{L\,event})$$
$$= (LE - RE)_{L\,event} - (LE - RE)_{R\,event}$$

The eye-specific differential response is essentially a special linear combination of all

voxels, where voxels belonging to LE ODCs are given a uniform weight of 1 while the RE

voxels are all given a weight of -1. Although faithfully reflecting the mean response

amplitude (e.g., across layers), this way of weight assignment may not be optimal for V1, or

even possible for higher-level areas for extracting eye-specific information. To increase

sensitivity, a better set of weights were obtained by training a linear classifier (we used linear

support vector machines from sklearn with default hyper-parameters) on ocular-bias localizer data to predict which eye was stimulated on a TR-by-TR basis, and the multivariate response patterns from other conditions can then be linearly projected using the optimal weights into a 1D timeseries that reflects the distance to the decision boundary at each moment. This is the multivariate differential response.

For Experiment 1, the multivariate method was used for IPS. Within atlas-defined aIPS or pIPS region, voxels with above-threshold visual response (omnibus $F > 1$, and L+R $t > 1$, but L+R beta < 5) and eye-bias (200 most biased voxels (2x up-sampled) in both ends of the L-R t distribution, and positive monocular response, e.g., L beta > 0 for LE-biased voxels) based on the GLM results of ocular-bias localizer were selected as features. Results were similar across a reasonable range of thresholds. The ideal response time course for the localizer was created by convolving the HRF ("GAM" with default parameters in AFNI) with the LE and RE block timings, and then taking their difference. Volumes at the flat part of the block responses (absolute value of the ideal response > 0.75 * maximum) were selected for training, whereas all volumes from the rivalry/replay runs were used at test time for generating the multivariate differential response. Each sample (feature vector) was normalized to have unitary Euclidean norm before train or test.

For Experiment 3, the multivariate method was used for frontoparietal areas and in the searchlight analysis. The models were trained on the localizer data to discriminate M from P stimulation. Since the format of M vs P representation might greatly differ across areas, the hyperparameters for feature selection were titrated for each ROI using grid search and cross validation. Data from rivalry or replay condition were split into two halves comprising disjoint runs. The space of possible feature selection hyperparameters was sampled by a 2D grid comprising the Cartesian product of 6 levels of visual responsiveness (from no constraint to GLM omnibus $F > 3$, M+P $t > 2$, M beta > 0, P beta > 0) and 8 levels of M-P bias (from no constraint to top 1% most biased voxels (2x upsampled) from both ends of the M-P t distribution). For each set of hyperparameters, a SVM was trained on all localizer data and an 1D discriminative projection of the rivalry or replay activation pattern at each time point was performed on the two halves separately. The models and the corresponding sets of features that resulted in significant perceptual modulation (see below) across subjects on the first half (the validation set) were chosen, and their results on the second half (the test set) were averaged and recorded, weighted by the effect size on the validation set. The roles of validation and test were then swapped and the average of the two test results was taken as the final result. The procedure was repeated for each ROI and each condition separately.

Perceptual modulation

Perceptual modulation, i.e., the difference in BOLD activity when subjects perceived one stimulus over the other, time-locked to button presses in either rivalry or replay condition, was estimated using several methods that led to similar results. To reduce the impact of HRF difference among brain areas (especially for subcortical nuclei), a model-free event-related average of BOLD signals was used in most cases. BOLD signals were averaged across voxels within each ROI (e.g., V1 LE-biased voxels or pulvinar P-biased voxels; for higher-level areas like IPS, multivariate differential response was used), linearly interpolated (0.1 s for Experiment 1/2 and 0.01 s for Experiment 3), smoothed with a 10-s hamming window (only for Experiment 3), and sorted into epochs time-aligned with button presses (LE/RE trials for Experiment 1/2 and M/P trials for Experiment 3). Only epochs with exclusive perception (non-mixed) for at least 4 s after and 2 s before the perceptual switch were analyzed. Button presses without a corresponding event in the paired rivalry/replay run were also discarded. Epochs were baseline corrected (subtracting the mean between -1 to 1 s; this was omitted for decoding time course in which case zero is a natural baseline) and averaged to get the event-related response. The modulation timecourse was obtained by subtracting responses between the L (M) and R (P) events. For eye-specific modulation, the results from LE- and RE-biased voxels were averaged. Finally, the mean value of the modulation timecourse between 4-12 s after the switch was taken as the estimated perceptual modulation. The map of rivalry modulation was computed on the surface in a similar way but by first projecting the BOLD timeseries onto the surface. The resulting map was high-pass filtered by subtracting the surface smoothed (8-mm FWHM) version. In Experiment 3, since some frontal areas exhibited large variability in the shape of the modulation time course, using a fixed time window for all ROIs to summarize the modulation amplitude seems suboptimal. Thus, we determined the window using a data-driven approach based on cross-validation. For each ROI, the mean modulation timecourse across subjects for one half of the data (see previous session) was smoothed, and the interval supporting the first positive peak (within 0-15 s) was taken as the time window, within which the mean response on the other half of the data could be recorded avoiding the risk of double-dipping.

To discount the influence of sluggish BOLD signal from previous trials, we estimated the perceptual modulation for V1 laminar analysis using a GLM-based method (3dDeconvolve with CSPLINzero model in AFNI), which may benefit from the high SNR in V1. BOLD timecourses from 0 to 24 s after each perceptual switch was modeled by 11 parameters separated by 2 s, with response at 0 and 24 s fixed to be zero. For each cortical layer, we selected voxels in a pre-defined range of equivolume depth (see above). Voxels

within each layer for each eye were pooled and the averaged timeseries was fed to the model.

The LE and RE events were modeled separately, and their results were differentiated

(preferred - non-preferred) and then averaged between LE- and RE-ODCs to get the

modulation curve. The mean response under the first positive peak was taken as the estimated

perceptual modulation.

To further discount the influence of perceptual duration in the V1-IPS correlation

analysis, the multivariate differential response for each ROI was modeled by a GLM with

variable length block (dmUBLOCK in AFNI). Besides run-wise baseline drift and head

motion regressors, only one perception-related regressor was included, in which LE- and RE-

dominant intervals were modeled as blocks of 1's and -1's before convolving with the HRF.

The resulting beta value was taken as the estimated perceptual modulation.

Laminar analysis of eye-specific modulation

To compare the shapes of laminar profiles in rivalry and replay conditions, eye-specific

modulations were normalized by dividing the sum of responses across cortical depth for each

subject (Fig. 2f). Since binocular rivalry and stimulus replay shared the same perception as

well as the laminar bias of BOLD signals, we further calculated the rivalry/replay modulation

ratio across cortical depth (Fig. 2g). To generate the continuous laminar profiles in Fig. 2e/2k,

the relative cortical depth from 0 to 1 was resampled into 31 points, and the mean response at

each depth was computed by averaging voxels near that depth with Gaussian weights (sigma

= 0.067 for EPI and 0.1 for bSSFP). Ocular modulations in the localizer data were defined as

the differential of LE and RE responses across cortical depth.

Pattern correlation

To quantify the synchrony of eye-specific rivalry dynamics in V1, we computed Pearson's

correlation coefficient between the moment-to-moment V1 response pattern in rivalry and

replay with the ODC pattern acquired from localizer. The preprocessed BOLD signal within

each layer was projected to the surface as the instantaneous activity pattern, and the ODC

pattern was computed by projecting the localizer LE-RE beta values within the gray matter to

the cortical surface. The volume-surface projection used median map-function. Perceptual

modulation was manifested both temporally and spatially (Figure 2b/c). By dividing the

standard deviation of the activity pattern that discounted differences in modulation amplitude

across layers, pattern correlation coefficient could indicate the synchrony of eye-specific

rivalry dynamics. If the ODCs for one eye become activated at slightly different times, or

their signals fluctuate asynchronously across the visual field, the correlation coefficient would

be closer to zero on average. Thus, the distribution width of all r values in the TR-by-TR

pattern correlation timecourse can be used as an index for the synchrony of eye-specific

dynamics. The distribution width was defined as two times the std, estimated separately for

each layer and condition. Since SNR also limits the maximally attainable pattern correlation,

which may vary across layers, we used the distribution width of synchronized stimulus-driven

response in replay as a benchmark with which to measure the synchrony of rivalry dynamics.

In the control analysis, we evaluated whether the SNR difference across cortical layer

alone could produce a similar laminar profile of pattern correlation. We first modeled the

synchronized BOLD responses in rivalry and replay conditions and in each layer using GLMs.

LE and RE dominant intervals were modeled as variable length blocks (dmUBLOCK in

AFNI) in two separate regressors. After model fitting, the residual timeseries were shuffled in

time independently for each vertex, which destroyed any unmodeled synchronous activity

fluctuation (e.g., at a particular time, vertices in LE ODCs of the left hemisphere may all

show higher activation than expected based on the global perceptual state due to slightly

earlier switch at this part of the visual field). The shuffled residuals were then added back to

the fitted timeseries, and the same pattern correlation analysis was repeated for the

recombined dataset.

Dynamic causal modeling

Effective connectivity of fMRI data was analyzed with the DCM module of SPM12 (Version

7771). In Experiment 1, multivariate decoding from V1, V2, and IPS were used as the VOI

timeseries. In Experiment 3, the differential timeseries between P- and M-biased voxels were

used as the VOI inputs of V1 and pulvinar, the difference between the mean responses of P-

biased voxels in hV4 and M-biased voxels in MT+ was used as data for the high-level visual

cortex (label here as hV4), and multivariate decoding timeseries was used as the IPS data.

The timeseries of rivalry and replay conditions were concatenated and modeled together. In

the eye-specific connectivity model for Experiment 1 (Fig. 4), there were two inputs: the eye-

of-origin of the currently perceived stimulus (high for LE and low for RE) was defined as the

driving input to V1 in both rivalry and replay conditions, and eye-specific driving inputs to all

brain areas (V1/V2/IPS) only in rivalry. Fixed connections were defined between and within

all brain areas, and the between-areas connections were allowed to be modulated by the 2nd

input during binocular rivalry. Both inputs were mean-centered. A bilinear, single state,

deterministic model with default parameters was used. At the first or individual level, the full

DCM for each subject was estimated using all data from rivalry and replay runs (Zeidman,

Jafarian, Corbin, et al., 2019). At the second or group level, we used the parametric empirical

Bayes method (Friston et al., 2016; Zeidman, Jafarian, Seghier, et al., 2019) to perform

Bayesian model reduction, Bayesian model average, and make inferences about the

connectivity strength. Both the C matrix for the second input and the B matrix were tested

and the averaged model for explaining the commonalities across subjects was shown. The z

values for the estimated parameters (e.g., changes in effective connectivity, i.e., the B matrix)
were computed by dividing their expectation (Ep) with the square root of the corresponding
diagonal elements in the covariance matrix (Cp). The stimulus-specific connectivity model
for Experiment 3 (Fig. 7) was defined and estimated analogously.

Hierarchical clustering

The hierarchical clustering analysis was performed using the dendrogram function from Scipy
with Euclidean distance and Ward's linkage. The data point for each ROI was a vector of
normalized rivalry and replay modulations across all subjects. The default threshold of 0.7
times maximum distance between clusters was used, and the resulting number of clusters was
determined from the dendrogram.

Statistical analysis

Statistical analyses were conducted using the Pingouin package (v0.5), JASP (v0.14), R
(v4.1), and home-built Python code (for permutation and bootstrap procedures). Cluster-based
permutation test (Maris et al., 2007; Nichols & Holmes, 2002) was used to test the difference
in timeseries correcting for multiple comparisons (see (Ge et al., 2020) for detailed
procedure). Perceptual modulation (as well as other bar-plots) was tested against zero using
one-tailed one-sample t-test and Holm correction for multiple comparisons (or otherwise
noted). Modulation amplitudes across conditions were tested using repeated measures
ANOVA followed by paired t-test with Holm correction (unless there were only three levels
in which case correction was unnecessary (Levin et al., 1994), or otherwise noted). The
laminar profiles of perceptual modulation in rivalry and replay were normalized (so that the
sum of all layers was one) before comparison. Similarly, data were normalized for each ROI
by dividing the L2-norm of rivalry and replay modulations to facilitate comparison across
ROIs in Figure 7. The normalization would not change the test results against zero or between
rivalry and replay conditions, because the modulation of each subject in each condition was
divided by the same value for a given ROI.

     To account for the correlation between voxels in accessing the between-session
consistency of V1 ODC maps and LGN ocular-biased clusters, the observed correlation
coefficients were compared with the null distribution generated by Monte Carlo simulation.
The spatial auto-correlation function within the ROI was first estimated from the localizer
GLM residual volumes (3dFWHMx in AFNI using three-parameter ACF model). 10000
simulated volumes (Gaussian random noise with specified spatial smoothness) were then
generated (3dClustSim in AFNI) as surrogate data, with which correlation coefficient under
null hypothesis was computed to get the null distribution. Finally, the observed statistic was
compared to the critical value of the null distribution for its significance.

1057

Acknowledgement

This study was supported by STI2030-Major Projects (2022ZD0211900, 2021ZD0204200), National Natural Science Foundation of China (31871107, 31930053), and Chinese Academy of Sciences (XDB32020200, KJZD-SW-L08,   CAS-NWO Program 153311KYSB20160030).

References

Angelucci, A., Bijanzadeh, M., Nurminen, L., Federer, F., Merlin, S., & Bressloff, P. C. (2017). Circuits and Mechanisms for Surround Modulation in Visual Cortex. *Annual Review of Neuroscience*, *40*, 425–451. https://doi.org/10.1146/annurev-neuro-072116-031418

Arcaro, M. J., Pinsk, M. A., & Kastner, S. (2015). The Anatomical and Functional Organization of the Human Visual Pulvinar. *Journal of Neuroscience*, *35*(27), 9848–9871. https://doi.org/10.1523/JNEUROSCI.1575-14.2015

Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A reproducible evaluation of ANTs similarity metric performance in brain image registration. *Neuroimage*, *54*(3), 2033–2044. https://doi.org/10.1016/j.neuroimage.2010.09.025

Balaram, P., Young, N. A., & Kaas, J. H. (2014). Histological features of layers and sublayers in cortical visual areas V1 and V2 of chimpanzees, macaque monkeys, and humans. *Eye Brain*, *2014*(6 Suppl 1), 5–18. https://doi.org/10.2147/EB.S51814

Beckett, A. J. S., Dadakova, T., Townsend, J., Huber, L., Park, S., & Feinberg, D. A. (2020). Comparison of BOLD and CBV using 3D EPI and 3D GRASE for cortical layer functional MRI at 7 T. *Magnetic Resonance in Medicine*, *84*(6), 3128–3145. https://doi.org/10.1002/mrm.28347

Benson, N C, Jamison, K. W., Arcaro, M. J., Vu, A. T., Glasser, M. F., Coalson, T. S., Van Essen, D. C., Yacoub, E., Ugurbil, K., Winawer, J., & Kay, K. (2018). The Human Connectome Project 7 Tesla retinotopy dataset: Description and population receptive field analysis. *J Vis*, *18*(13), 23. https://doi.org/10.1167/18.13.23

Benson, Noah C., Butt, O. H., Brainard, D. H., & Aguirre, G. K. (2014). Correction of Distortion in Flattened Representations of the Cortical Surface Allows Prediction of V1-V3 Functional Organization from Anatomy. *PLoS Computational Biology*, *10*(3). https://doi.org/10.1371/journal.pcbi.1003538

Blake, R. (1989). A neural theory of binocular rivalry. *Psychol Rev*, *96*(1), 145–167. http://www.ncbi.nlm.nih.gov/pubmed/2648445

Blake, R, & Logothetis, N. K. (2002). Visual competition. *Nature Reviews Neuroscience*, *3*(1),
13–23. https://doi.org/Doi 10.1038/Nrn701

Blake, Randolph, Brascamp, J., & Heeger, D. J. (2014). Can binocular rivalry reveal neural
correlates of consciousness? *Philosophical Transactions of the Royal Society B-*
Biological Sciences, *369*(1641). https://doi.org/ARTN 2013021110.1098/rstb.2013.0211

Brascamp, J. W., & Blake, R. (2012). Inattention abolishes binocular rivalry: perceptual
evidence. *Psychol Sci*, *23*(10), 1159–1167. https://doi.org/10.1177/0956797612440100

Brascamp, J, Blake, R., & Knapen, T. (2015). Negligible fronto-parietal BOLD activity
accompanying unreportable switches in bistable perception. *Nat Neurosci*, *18*(11),
1672–1678. https://doi.org/10.1038/nn.4130

Brascamp, Jan, Sterzer, P., Blake, R., & Knapen, T. (2018). Multistable Perception and the
Role of the Frontoparietal Cortex in Perceptual Inference. *Annual Review of Psychology*,
69(1), 77–103. https://doi.org/10.1146/annurev-psych-010417-085944

Bridge, H., Leopold, D. A., & Bourne, J. A. (2016). Adaptive Pulvinar Circuitry Supports
Visual Cognition. *Trends in Cognitive Sciences*, *20*(2), 146–157.
https://doi.org/10.1016/j.tics.2015.10.003

Briggs, F., & Usrey, W. M. (2011). Corticogeniculate feedback and visual processing in the
primate. *J Physiol*, *589*(Pt 1), 33–40. https://doi.org/10.1113/jphysiol.2010.193599

Burkhalter, A., & Van Essen, D. C. (1986). Processing of color, form and disparity
information in visual areas VP and V2 of ventral extrastriate cortex in the macaque
monkey. *Journal of Neuroscience*, *6*(8), 2327–2351.
https://doi.org/10.1523/jneurosci.06-08-02327.1986

Buzs, P., Eysel, U. T., Adorjn, P., & Kisvrday, Z. F. (2001). Axonal topography of cortical
basket cells in relation to orientation, direction, and ocular dominance maps. *Journal of*
Comparative Neurology, *437*(3), 259–285. https://doi.org/10.1002/cne.1282

Carmel, D., Walsh, V., Lavie, N., & Rees, G. (2010). Right parietal TMS shortens dominance
durations in binocular rivalry. *Curr Biol*, *20*(18), R799-800.
https://doi.org/10.1016/j.cub.2010.07.036

Cox, M. A., Dougherty, K., Westerberg, J. A., Schall, M. S., & Maier, A. (2019). *Temporal*
dynamics of binocular integration in primary visual cortex. *19*, 1–21.

Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic
resonance neuroimages. *Comput Biomed Res*, *29*(3), 162–173.
https://doi.org/10.1006/cbmr.1996.0014

Crick, F. (1996). Visual perception: rivalry and consciousness. *Nature*.

Dayan, P. (1998). A hierarchical model of binocular rivalry. *Neural Comput*, *10*(5), 1119–
1135. http://www.ncbi.nlm.nih.gov/pubmed/9654769

de Jong, M. C., Vansteensel, M. J., van Ee, R., Leijten, F. S. S., Ramsey, N. F., Dijkerman, H.

C., Dumoulin, S. O., & Knapen, T. (2020). Intracranial Recordings Reveal Unique

Shape and Timing of Responses in Human Visual Cortex during Illusory Visual Events.

Current Biology, *30*(16), 3089-3100.e4. https://doi.org/10.1016/j.cub.2020.05.082

de Sousa, A. A., Sherwood, C. C., Schleicher, A., Amunts, K., MacLeod, C. E., Hof, P. R., &

Zilles, K. (2010). Comparative Cytoarchitectural Analyses of Striate and Extrastriate

Areas in Hominoids. *Cerebral Cortex*, *20*(4), 966–981.

https://doi.org/10.1093/cercor/bhp158

Derrington, A. M., & Lennie, P. (1984). Spatial and temporal contrast sensitivities of

neurones in lateral geniculate nucleus of macaque. *J Physiol*, *357*, 219–240.

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita

tion&list_uids=6512690

Dougherty, K, Cox, M. A., Westerberg, J. A., & Maier, A. (2019). Binocular Modulation of

Monocular V1 Neurons. *Curr Biol*, *29*(3), 381-391 e4.

https://doi.org/10.1016/j.cub.2018.12.004

Dougherty, Kacie, Carlson, B. M., Cox, M. A., Westerberg, J. A., Zinke, W., Schmid, M. C.,

Martin, P. R., & Maier, A. (2021). Binocular suppression in the macaque lateral

geniculate nucleus reveals early competitive interactions between the eyes. *ENeuro*, *8*(2),

1–12. https://doi.org/10.1523/ENEURO.0364-20.2020

Dougherty, Kacie, Schmid, M. C., & Maier, A. (2018). Binocular response modulation in the

lateral geniculate nucleus. *J Comp Neurol*, *527*(3), 522–534.

https://doi.org/10.1002/cne.24417

Felleman, D. J., & Van Essen, D. C. (1991). Distributed hierarchical processing in the primate

cerebral cortex. *Cereb Cortex*, *1*(1), 1–47.

https://www.ncbi.nlm.nih.gov/pubmed/1822724

Fischl, B. (2012). FreeSurfer. *Neuroimage*, *62*(2), 774–781.

Friston, K. J., Litvak, V., Oswal, A., Razi, A., Stephan, K. E., van Wijk, B. C. M., Ziegler, G.,

& Zeidman, P. (2016). Bayesian model reduction and empirical Bayes for group (DCM)

studies. *Neuroimage*, *128*, 413–431. https://doi.org/10.1016/j.neuroimage.2015.11.015

Ge, Y., Zhou, H., Qian, C., Zhang, P., Wang, L., & He, S. (2020). Adaptation to feedback

representation of illusory orientation produced from flash grab effect. *Nature*

Communications, *11*(1). https://doi.org/10.1038/s41467-020-17786-1

Gilbert, C. D., & Wiesel, T. N. (1983). Clustered intrinsic connections in cat visual cortex.

Journal of Neuroscience, *3*(5), 1116–1133. https://doi.org/10.1523/jneurosci.03-05-

01116.1983

Guillery, R. W., & Colonnier, M. (1970). Synaptic patterns in the dorsal lateral geniculate

nucleus of the monkey. *Zeitschrift Für Zellforschung Und Mikroskopische Anatomie*,

103(1), 90–108. https://doi.org/10.1007/BF00335403

Haynes, J. D., Deichmann, R., & Rees, G. (2005). Eye-specific effects of binocular rivalry in
the human lateral geniculate nucleus. *Nature*, *438*(7067), 496–499.
https://doi.org/nature04169 [pii]10.1038/nature04169

He, S., Carlson, T., & Chen, X. (2005). Parallel Pathways and Temporal Dynamics in
Binocular Rivalry. *Binocular Rivalry*.

Jaramillo, J., Mejias, J. F., & Wang, X. J. (2019). Engagement of Pulvino-cortical
Feedforward and Feedback Pathways in Cognitive Computations. *Neuron*, *101*(2), 321-
+. https://doi.org/10.1016/j.neuron.2018.11.023

Kaas, J. H., & Lyon, D. C. (2007). Pulvinar contributions to the dorsal and ventral streams of
visual processing in primates. In *Brain Research Reviews* (Vol. 55, Issue 2 SPEC. ISS.,
pp. 285–296). https://doi.org/10.1016/j.brainresrev.2007.02.008

Kanai, R., Carmel, D., Bahrami, B., & Rees, G. (2011). Structural and functional
fractionation of right superior parietal cortex in bistable perception. *Curr Biol*, *21*(3),
R106-7. https://doi.org/10.1016/j.cub.2010.12.009

Kovács, I., Papathomas, T. V., Yang, M., & Fehér, Á. (1996). When the brain changes its
mind: Interocular grouping during binocular rivalry. *Proceedings of the National*
Academy of Sciences of the United States of America, *93*(26), 15508–15511.
https://doi.org/10.1073/pnas.93.26.15508

Lehky, S. R., & Maunsell, J. H. R. (1996). No binocular rivalry in the LGN of alert macaque
monkeys. *Vision Res*, *36*(9), 1225–1234. https://doi.org/Doi 10.1016/0042-
6989(95)00232-4

Leopold, D. A., & Logothetis, N. K. (1996). Activity changes in early visual cortex reflect
monkeys' percepts during binocular rivalry. *Nature*, *379*(6565), 549–553.
https://doi.org/10.1038/379549a0

Levin, J. R., Serlin, R. C., & Seaman, M. A. (1994). A controlled, powerful multiple-
comparison strategy for several situations. *Psychological Bulletin*, *115*(1), 153.

Li, H. H., Rankin, J., Rinzel, J., Carrasco, M., & Heeger, D. J. (2017). Attention model of
binocular rivalry. *Proc Natl Acad Sci U S A*, *114*(30), E6192–E6201.
https://doi.org/10.1073/pnas.1620475114

Liu, C., Guo, F., Qian, C., Zhang, Z., Sun, K., Wang, D. J. J., He, S., & Zhang, P. (2020).
Layer-dependent multiplicative effects of spatial attention on contrast responses in
human early visual cortex. *Progress in Neurobiology*, *207*, 101897.
https://doi.org/https://doi.org/10.1016/j.pneurobio.2020.101897

Logothetis, N. K., & Wandell, B. A. (2004). Interpreting the BOLD signal. *Annu Rev Physiol*,
66, 735–769. https://doi.org/10.1146/annurev.physiol.66.082602.092845

Lumer, E. D., Friston, K. J., & Rees, G. (1998). Neural correlates of perceptual rivalry in the
human brain. *Science*, *280*(5371), 1930–1934.

http://www.ncbi.nlm.nih.gov/pubmed/9632390

Maier, A., Wilke, M., Aura, C., Zhu, C., Ye, F. Q., & Leopold, D. A. (2008). Divergence of

fMRI and neural signals in V1 during perceptual suppression in the awake monkey. *Nat*

Neurosci, *11*(10), 1193–1200. https://doi.org/10.1038/nn.2173

Maris, E., Schoffelen, J.-M., & Fries, P. (2007). Nonparametric statistical testing of coherence

differences. *Journal of Neuroscience Methods*, *163*(1), 161–175.

Maunsell, J. H. R., & Van Essen, D. C. (1983). Functional properties of neurons in middle

temporal visual area of the macaque monkey. II. Binocular interactions and sensitivity to

binocular disparity. *Journal of Neurophysiology, 49*(5), 1148–1167.

https://doi.org/10.1152/jn.1983.49.5.1148

McAlonan, K., Cavanaugh, J., & Wurtz, R. H. (2008). Guarding the gateway to cortex with

attention in visual thalamus. *Nature*, *456*(7220), 391–394.

https://doi.org/10.1038/nature07382

Mitchell, B. A., Dougherty, K., Westerberg, J. A., Carlson, B. M., Daumail, L., Maier, A., &

Cox, M. A. (2022). Stimulating both eyes with matching stimuli enhances V1 responses.

IScience, *25*(5). https://doi.org/10.1016/j.isci.2022.104182

Mo, C., Lu, J., Shi, C., & Fang, F. (2022). Neural representations of competing stimuli along

the dorsal and ventral visual pathways during binocular rivalry. *Cerebral Cortex*,

bhac238. https://doi.org/10.1093/cercor/bhac238

Moon, C. H., Fukuda, M., Park, S. H., & Kim, S. G. (2007). Neural interpretation of blood

oxygenation level-dependent fMRI maps at submillimeter columnar resolution. *Journal*

of Neuroscience, *27*(26), 6892–6902. https://doi.org/10.1523/Jneurosci.0445-07.2007

Myerson, J., Miezin, F., & Allman, J. (1981). Binocular rivalry in macaque monkeys and

humans: a comparative study in perception. *Behav. Anal. Lett*, *1*, 149–159.

Nichols, T. E., & Holmes, A. P. (2002). Nonparametric permutation tests for functional

neuroimaging: A primer with examples. *Human Brain Mapping*, *15*(1), 1–25.

https://doi.org/DOI 10.1002/hbm.1058

O'Connor, D. H., Fukui, M. M., Pinsk, M. A., & Kastner, S. (2002). Attention modulates

responses in the human lateral geniculate nucleus. *Nat Neurosci*, *5*(11), 1203–1209.

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita

tion&list_uids=12379861

O'Hashi, K., Fekete, T., Deneux, T., Hildesheim, R., van Leeuwen, C., & Grinvald, A. (2018).

Interhemispheric Synchrony of Spontaneous Cortical States at the Cortical Column

Level. *Cerebral Cortex*, *28*(5), 1794–1807. https://doi.org/10.1093/cercor/bhx090

Olman, C. A., Harel, N., Feinberg, D. A., He, S., Zhang, P., Ugurbil, K., & Yacoub, E. (2012).

Layer-specific fMRI reflects different neuronal computations at different depths in

human V1. *PLoS One*, *7*(3), e32536. https://doi.org/10.1371/journal.pone.0032536

Omer, D. B., Fekete, T., Ulchin, Y., Hildesheim, R., & Grinvald, A. (2018). Dynamic Patterns of Spontaneous Ongoing Activity in the Visual Cortex of Anesthetized and Awake Monkeys are Different. *Cereb Cortex*. https://doi.org/10.1093/cercor/bhy099

Polimeni, J. R., Renvall, V., Zaretskaya, N., & Fischl, B. (2017). Analysis strategies for high-resolution UHF-fMRI data. *Neuroimage*. https://doi.org/10.1016/j.neuroimage.2017.04.053

Qian, Y., Zou, J., Zhang, Z., An, J., Zuo, Z., Zhuo, Y., Wang, D. J. J., & Zhang, P. (2020). Robust functional mapping of layer-selective responses in human lateral geniculate nucleus with high-resolution 7T fMRI. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1925). https://doi.org/10.1098/rspb.2020.0245

Rafal, R., Henik, A., & Smith, J. (1991). Extrageniculate contributions to reflex visual orienting in normal humans: a temporal hemifield advantage. *J Cogn Neurosci*, *3*(4), 322–328. https://doi.org/10.1162/jocn.1991.3.4.322

Renzo, L., Poser, B. A., Bandettini, P. A., Arora, K., Wagstyl, K., Cho, S., Goense, J., Nothnagel, N., Tyler, A., Hurk, J. Van Den, Müller, A. K., Reynolds, R. C., Glen, D. R., Goebel, R., & Faruk, O. (2021). NeuroImage LayNii : A software suite for layer-fMRI. *NeuroImage*, *237*(February), 118091. https://doi.org/10.1016/j.neuroimage.2021.118091

Saalmann, Y. B., Pinsk, M. A., Wang, L., Li, X., & Kastner, S. (2012). The Pulvinar Regulates Information Transmission Between Cortical Areas Based on Attention Demands. *Science*, *337*(6095), 753–756. https://doi.org/10.1126/science.1223082

Scheffler, K., Heule, R., M, G. B.-Y., Kardatzki, B., & Lohmann, G. (2018). The BOLD sensitivity of rapid steady-state sequences. *Magn Reson Med*. https://doi.org/10.1002/mrm.27585

Schneider, K. A. (2011). Subcortical mechanisms of feature-based attention. *Journal of Neuroscience*, *31*(23), 8643–8653. https://doi.org/10.1523/JNEUROSCI.6274-10.2011

Schneider, K. A., & Kastner, S. (2009). Effects of sustained spatial attention in the human lateral geniculate nucleus and superior colliculus. *Journal of Neuroscience*, *29*(6), 1784–1795. http://www.ncbi.nlm.nih.gov/pubmed/19211885

Schroeder, C. E., Tenke, C. E., Arezzo, J. C., & Vaughan, H. G. (1990). Binocularity in the Lateral Geniculate-Nucleus of the Alert Macaque. *Brain Research*, *521*(1–2), 303–310. https://doi.org/Doi 10.1016/0006-8993(90)91556-V

Schwarzkopf, D. S., Schindler, A., & Rees, G. (2010). Knowing with which eye we see: Utrocular discrimination and eye-specific signals in human visual cortex. *PLoS ONE*, *5*(10). https://doi.org/10.1371/journal.pone.0013775

Sengpiel, F., Blakemore, C., & Harrad, R. (1995). Interocular suppression in the primary visual cortex: a possible neural basis of binocular rivalry. *Vision Res*, *35*(2), 179–195. https://doi.org/0042-6989(94)00125-6 [pii]

Shipp, S. (2003). The functional logic of cortico-pulvinar connections. *Philos Trans R Soc*
Lond B Biol Sci, *358*(1438), 1605–1624. https://doi.org/10.1098/rstb.2002.1213

Shmuel, A., Yacoub, E., Pfeuffer, J., Van de Moortele, P. F., Adriany, G., Hu, X., & Ugurbil,
K. (2002). Sustained negative BOLD, blood flow and oxygen consumption response and
its coupling to the positive response in the human brain. *Neuron*, *36*(6), 1195–1210.
https://www.ncbi.nlm.nih.gov/pubmed/12495632

Tong, F., & Engel, S. A. (2001). Interocular rivalry revealed in the human cortical blind-spot
representation. *Nature*, *411*(6834), 195–199.
https://doi.org/10.1038/3507558335075583 [pii]

Tong, F., Meng, M., & Blake, R. (2006). Neural bases of binocular rivalry. *Trends Cogn Sci*,
10(11), 502–511. https://doi.org/S1364-6613(06)00240-3
[pii]10.1016/j.tics.2006.09.003

Tootell, R. B. H., Hadjikhani, N., Hall, E. K., Marrett, S., Vanduffel, W., Vaughan, J. T., &
Dale, A. M. (1998). The retinotopy of visual spatial attention. *Neuron*, *21*(6), 1409–
1422.

Uludag, K., & Havlicek, M. (2021). Determining laminar neuronal activity from BOLD fMRI
using a generative model. *Progress in Neurobiology*, *207*(March), 102055.
https://doi.org/https://doi.org/10.1016/j.pneurobio.2021.102055

Waehnert, M. D., Dinse, J., Weiss, M., Streicher, M. N., Waehnert, P., Geyer, S., Turner, R.,
& Bazin, P. L. (2014). Anatomically motivated modeling of cortical laminae.
Neuroimage, *93 Pt 2*, 210–220. https://doi.org/10.1016/j.neuroimage.2013.03.078

Wang, J., Nasr, S., Roe, A. W., & Polimeni, J. R. (2022). Critical factors in achieving
fine‐scale functional MRI : Removing sources of inadvertent spatial smoothing .
Human Brain Mapping, *November 2021*, 1–21. https://doi.org/10.1002/hbm.25867

Wang, L., Mruczek, R. E. B., Arcaro, M. J., & Kastner, S. (2015). Probabilistic maps of
visual topography in human cortex. *Cerebral Cortex*, *25*(10), 3911–3931.
https://doi.org/10.1093/cercor/bhu277

Wiesel, T. N., & Hubel, D. H. (1966). Spatial and chromatic interactions in the lateral
geniculate body of the rhesus monkey. *J Neurophysiol*, *29*(6), 1115–1156.
http://www.ncbi.nlm.nih.gov/pubmed/4961644

Wilke, M., Mueller, K. M., & Leopold, D. A. (2009). Neural activity in the visual thalamus
reflects perceptual suppression. *Proc Natl Acad Sci U S A*, *106*(23), 9465–9470.
https://doi.org/10.1073/pnas.0900714106

Wilson, H. R. (2003). Computational evidence for a rivalry hierarchy in vision. *Proc Natl*
Acad Sci U S A, *100*(24), 14499–14503. https://doi.org/10.1073/pnas.2333622100

Wunderlich, K., Schneider, K. A., & Kastner, S. (2005). Neural correlates of binocular rivalry

in the human lateral geniculate nucleus. *Nat Neurosci*, *8*(11), 1595–1602.

https://doi.org/nn1554 [pii]10.1038/nn1554

Xu, H., Han, C., Chen, M., Li, P., Zhu, S., Fang, Y., Hu, J., Ma, H., & Lu, H. D. (2016).

Rivalry-Like Neural Activity in Primary Visual Cortex in Anesthetized Monkeys.

Journal of Neuroscience, *36*(11), 3231–3242.

https://doi.org/10.1523/JNEUROSCI.3660-15.2016

Yu, Z., Guindani, M., Grieco, S. F., Chen, L., Holmes, T. C., & Xu, X. (2022). Beyond t test

and ANOVA: applications of mixed-effects models for more rigorous statistical analysis

in neuroscience research. *Neuron*, *110*(1), 21–35.

https://doi.org/10.1016/j.neuron.2021.10.030

Zaretskaya, N., Bause, J., Polimeni, J. R., Grassi, P. R., Scheffler, K., & Bartels, A. (2020).

Eye-selective fMRI activity in human primary visual cortex: Comparison between 3   T

and 9.4   T, and effects across cortical depth. *NeuroImage*, *220*(October 2019).

https://doi.org/10.1016/j.neuroimage.2020.117078

Zaretskaya, N., Thielscher, A., Logothetis, N. K., & Bartels, A. (2010). Disrupting parietal

function prolongs dominance durations in binocular rivalry. *Curr Biol*, *20*(23), 2106–

2111. https://doi.org/10.1016/j.cub.2010.10.046

Zeidman, P., Jafarian, A., Corbin, N., Seghier, M. L., Razi, A., Price, C. J., & Friston, K. J.

(2019). A guide to group effective connectivity analysis, part 1: First level analysis with

DCM for fMRI. *Neuroimage*, *200*, 174–190.

https://doi.org/10.1016/j.neuroimage.2019.06.031

Zeidman, P., Jafarian, A., Seghier, M. L., Litvak, V., Cagnan, H., Price, C. J., & Friston, K. J.

(2019). A guide to group effective connectivity analysis, part 2: Second level analysis

with PEB. *Neuroimage*, *200*, 12–25. https://doi.org/10.1016/j.neuroimage.2019.06.032

Zhang, P., Jamison, K., Engel, S., He, B., & He, S. (2011). Binocular rivalry requires visual

attention. *Neuron*, *71*(2), 362–369. https://doi.org/10.1016/j.neuron.2011.05.035 S0896-

6273(11)00492-2 [pii]

Zhang, P., Jiang, Y., & He, S. (2012). Voluntary attention modulates processing of eye-

specific visual information. *Psychol Sci*, *23*(3), 254–260.

https://doi.org/10.1177/0956797611424289 0956797611424289 [pii]

Zhou, H., Schafer, R. J., & Desimone, R. (2016). Pulvinar-Cortex Interactions in Vision and

Attention. *Neuron*, *89*(1), 209–220. https://doi.org/10.1016/j.neuron.2015.11.034

Zou, J., He, S., & Zhang, P. (2016). Binocular rivalry from invisible patterns. *Proceedings of*

the National Academy of Sciences of the United States of America, *113*(30).

https://doi.org/10.1073/pnas.1604816113

1349