Curvature coding in early visual system revealed by scale variance during adaptation to flashing circles

Irfa Nisar

Department of Electrical Engineering and Computer Science

Center for Vision Research, York University

James H. Elder

Department of Electrical Engineering and Computer Science

Center for Vision Research, York University

How is curvature coded in the early human visual system? This question is the essence of basic shape recognition and may explain why humans are so successful in recognizing objects under varying scale conditions (Biederman & Cooper, 1992; Lindeberg, 2013). The circle-polygon illusion produces polygonal percepts during adaptation when a static dark outline circle is pulsed at 2 Hz alternating with a gradient luminance circle. Both the radius and eccentricity of the stimulus were varied in a crossed design over 1-8 deg. Observers reported a circle or the polygon order and the strength of the percept. We test a lower level account that argues for curvature opponency between neurons against a higher level account that codes for whole shapes. This higher level account supports scale invariance, a property through which we recognize objects regardless of the object's size on the retina. We show the following: (1) Scale invariance is not obeyed during adaptation. The mean order of the perceived polygon increased with stimulus size and decreased with eccentricity. It did not remain constant for a stimulus size. This also demonstrates that the curvature coding occurs in the early visual system. (2) Linear regression analysis reveals that the cortical size of the stimulus is a better predictor of perceived polygon order. We quantify the relationship parametrically between cortical size and polygon order. Using integration and regression, we identify the region of the cortex, V1, where the shape, a regular ordered polygon, is being computationally constructed.

Keywords: shape adaptation, cortical adaptation

Introduction

Why is curvature perception important? Coding of curvature through the visual system influences how complex shapes are recognized in the world and why the human visual system is able to identify the same shape presented at varying scales (Biederman & Cooper, 1992; Lindeberg, 2013). The research objective is to answer the following questions: (a) Is curvature coded on a low level retinal plane or is it a product of a higher order cortical process? (b) Is the coding stimuli-scale variant or invariant: Does the size of

the stimuli change the number of sides of the polygon seen in the percept after adaptation? (c) Does the polygon order (shape) change with size and eccentricity? Can we quantify the relationship between the shape seen during in-place adaptation and the independent variables, size and eccentricity, at which the stimuli is presented?

Circle Polygon Illusion We adopt the circle-polygon illusion that was discovered by Khuu (Khuu, 2002) and use Sakurai's technique (Sakurai, 2014; Sakurai & Beaudot, 2015) to construct the phenomenon to understand how curvature is coded in the brain. Sakurai's illusion, builds on Khuu's circle-appearing-as-polygon illusion, by temporally pulsing a graded circle with its outline at 2 Hz. The circle-polygon illusion demonstrates that polygons are produced after prolonged adaptation to flashing circles. Although (Khuu, 2002; Sakurai, 2014; Sakurai & Beaudot, 2015) only report hexagon as the emergent shape, Ito in (Ito, 2012) presents an array of shapes (triangle,square,pentagon,hexagon,heptagon and circle) as the independent variable response type with several responses that are not hexagons. (Ito, 2012) shows that adaptation to a static and a rotating circle,either filled or unfilled, generates an afterimage of a hexagon. In (Sakurai, 2014; Sakurai & Beaudot, 2015), Sakurai show that if a circle and its inward gradation are alternated, we see a polygon shape perception of the circular stimuli itself. That shape is a hexagon.

Lower level or higher level? The early visual system is defined by the retina, and controlled through the receptive fields of regions V1 through V2 (Hubel & Wiesel, 1959; Hubel, Wiesel, & Stryker, 1978; Hegdé & Van Essen, 2000). The higher mid-level V4 neurons, with larger receptive fields, while showing preference for certain angles, show scale-invariance to different sized arc boundaries that subtend the same angle at the stimulus's center (Pasupathy & Connor, 2001, 2002; El-Shamayleh & Pasupathy, 2016a). We probe systematically across eccentricity and size to see if this scaling phenomenon holds and if not, can we deduce a relationship that explains the scaling. Ito (Ito, 2012) assert that the circle-polygon phenomenon is cortical because interocular transfers result in users reporting circles whereas each eye under monoptical conditions, when shown the same circular stimuli would produce a polygon. Sakurai in (Sakurai & Beaudot, 2015), build on Khuu's circle-appearing-as-polygon illusion and also cite a cortical process is responsible given that the adaptation occurs almost instantly when the stimuli pulsation is varied between 0.5 Hz to 8 Hz. Beyond the plausible observation that it is cortical, can we quantitatively state that polygons produced are determinant of the size and eccentricity as a known function? If the polygonal response to curvature was coded on the retinal plane or lower levels V1 and V2, it would be influenced by the neuronal responses to the shape as it appears on the retina. However, if it was coded higher up in the cortex, it might show more complex representations such as scale invariance with respect to size (Pasupathy & Connor, 2001, 2002; El-Shamayleh & Pasupathy, 2016a).

Scale invariance is a well known property of the higher human visual system, beyond V4 (Schwartz, Desimone, Albright, & Gross, 1983; Desimone, Albright, Gross, & Bruce, 1984; Logothetis, Pauls, & Poggio, 1995; Brincat & Connor, 2004; Liu, Agam, Madsen, & Kreiman, 2009; Rust & DiCarlo, 2010). Schwartz et. al and Logothetis et. al. Schwartz et. al (Schwartz et al., 1983) demonstrate invariance by recording from the inferotemporal (IT) region, a region that codes for global shape, by showing that position and size invariance is observed for stimuli constructed from Fourier descriptors. Neurons are selective for certain views and broadly tolerant to transformations such as rotation and scaling that is centered around the primary view (Logothetis et al., 1995).

As seen in (Sakurai, 2014; Sakurai & Beaudot, 2015), the circular stimuli, when alternated with its inward gradation, makes the cortical process adapt to the curvature and gives rise to symmetrical regular hexagons instead of an irregular organic shape. Gheorghiu

and Kingdom in (Gheorghiu & Kingdom, 2007, 2009, 2017) demonstrate that population neural code affects perception of local curvature through an AND-gate multiplication mechanism. However, none of the above experiments explain the order of the polygon as a function of the stimuli size. Neurophysiological studies conducted on non-primate data in (Das & Gilbert, 1995) show that receptive fields that fire for a similar range of orientations cluster together but long-range connections can sometimes influence the range of degree to which orientations are recognized. Why is the regularity of the afterimage, also called regular polygonalization, an important attribute in the afterimage? The presence of regularization itself suggests that retinal bleaching or retinal adaptation (Spillmann, 1994; Schrauf, Lingelbach, & Wist, 1997) is an unfavourable hypothesis since bleaching produces disappearances in features or production of new ones (opposite coloured fault lines) when intensity rods are fatigued (Spillmann, 1994; Schrauf et al., 1997). The disappearance occurs peripheral to fixation where fatiguing is pronounced and lacks the deliberate production of symmetry which could only be produced if rods or neurons were aligned in a corresponding symmetrical pattern. We test if a cortical process operating in the lower regions such as V1 utilizes long range connections in the production of polygonal shapes. In this paper, we compare a retinal and a competing cortical model to see which better explains the adapted polygon and if we can quantitatively define a relationship. The retinal model relates the retinal size of the stimulus to the order of the polygon. The second model, relies on cortical M-scaling and relates the cortical image of the stimulus to the order of the polygon.

The visual cortex is a large area with several sub-regions V1, V2, V3 and V4. Each region has properties that are specific to the region. For example, V1, scales linearly on the cortex while region V4 is known to show shape specific orientation preference. We try to answer the research question: which sub-region and its specific preference for orientation or scaling best explains the edge length being reported for each stimulus? There is a correspondence between neurons firing for a particular feature and its recorded response. We use this behavioural criterion as the dependent variable and the neuroscience theory to construct what the stimulus circumference as a shape looks like on the visual cortex.

Parametric cortical equation How can one quantify the stimuli's projection on the cortex? It is known that receptive field size increase towards the retina's periphery (Rovamo & Virsu, 1979). M is defined as the ratio between the cortical representation of the receptive field (in mm) and its size on the retina (degrees of visual angle). In (Daniel & Whitteridge, 1961), Daniel and Whitteridge show that an increment of 1 degree visual angle measured from the central fixation point corresponds to an arc length of M mm on the theoretical visual cortex.

Rovamo and Virsu in (Rovamo & Virsu, 1979) estimate M, for the principal meridians of the visual field in equation 1 and the accompanying table for coefficients. Here, M_o is $7.99 mm/°$, the magnification at the fovea. r is the eccentricity in degrees.

$$M(r) = (1 + ar + br^3)^{-1} M_o \tag{1}$$

Area where magnification is computed	a	b	Eccentricity range
Nasal	0.33	0.00007	$0 \leq E \leq 60 \deg$

If we integrate M, we obtain the distance on the cortex as a function of eccentricity. Our skeletal algorithm, see Algorithm 1,

computes the integral on the cortex and is explained in the Methods section.

Experiment 1

The experiment reports the polygon percept seen during adaptation as the circle radius and the eccentricity of the circle's placement varies.

Methods

In order to explain if the polygons observed above can be explained quantitatively by a retinal model or a higher level cortical model, we apply linear regression. We fit a first order linear model of the polygon order as a function of retinal size and another as a function of cortical circumference size. A retinal function samples the circular stimulus in 2D cartesian co-ordinates. The domain of the function is degree visual angle representing eccentricity. The cortical length can be computed from the points on the circle seen on the 2D retinal plane and applying the cortical M-scaling, that measures millimetres of cortex per degree of visual angle, as a function of eccentricity, on those sampled 2D points.

We shall hereby call the retinal circumference size, retinal length, and the cortical circumference size, cortical length. See Figure 11 (a,b,d).

The M-scaling factors in the cortical function were borrowed from (Rovamo & Virsu, 1979). This is a skeletal model with Rovamo and Virsu cortical magnification factors. The cortical magnification factors can be replaced by other magnification factors that perform hierarchical sampling, such as when the cortical representation in V4 is represented by how the region in V1 is sampled (Harvey & Dumoulin, 2011). Harvey and Dumoulin create a model that maps population receptive fields and cortical magnification factor as a function of the cortical magnification factor of the first field in the visual hierarchy, V1. Harvey and Dumoulin (Harvey & Dumoulin, 2011) show that the cortical magnification factor in area V4 is smaller than those in V2 and V3 which was in turn is less than V1. The cortical magnification factor for fields such as V2,V3 and V4 that sample from V1 is independently constructed. We, therefore, use V4's cortical magnification factor when considering if polygon order reported could be a function of cortical scaling in region V4, which is known for curvature representation.

In Algorithm 1, we approximate the cortical length formed by sampling the retinal stimuli's circumference at a fixed number of points and projecting that sampled length onto the cortex. The length formed on the cortex is scaled by the magnification factor, M, to give the cortical length.

We use boot strapping and use the coefficient of determination, R^2 to estimate the goodness of fit when fitting a linear regression function, polygon order as a function of the retinal and cortical length.

Observers

Nine graduate students (6 male, 3 female) at the university participated in the experiment. Four of the graduate students had prior experience in participating in psychophysical experiments. The remaining five students had limited or no experience related to

---

Algorithm 1: Cortical mapping function using a neurophysiological derived cortical magnification factor
Compute the cortical length using the M-scaling magnification factor for a circle of a given radius and presented at a particular eccentricity.

  ▷ radius: Radius of the circle   ▷ center: Center of the circle; The center, in cartesian coordinates, represents the eccentricity at which the circle is presented on the retinal 2D plane.

procedure COMPUTE(radius,center)

  Output Cortical length (in mm) as represented on the cortex region called V1.

  Initialization

  Set number of point samples on the circle, call it n

  unit sample length = 1/n

  Do:

  tangent=$2 * \pi * radius * (-sin(2\pi t), cos(2\pi t))$ ▷ Compute the tangent; t runs in unit sample length increments between 0 and 1

  speed = $\|tangent\|$ ▷ Compute the speed on the curve using the tangent

  M = $(1 + 0.33 * \text{eccentricity} + 0.00007 * \text{eccentricity}^3)^{-1} * M_o$ ▷ Cortical magnification factor for region V1. Use Rovamo and Virsu's value as an example.

  Cspeed=M*speed   ▷ Speed on the cortex

  len=Cspeed*unit sample length   ▷ Compute the cortical length by multiplying the M-scaling magnification factor with the speed over the sample length.

---

psychophysics. All the students had normal or corrected to normal vision. The experimental data and source code are stored at the Open Science Framework at the following link https://osf.io/x8n3p/.

Stimuli

The stimuli consisted of a circle with a black outline that alternated with another circle of the same size but with an inward gradation pattern. The black outlines were constructed of penwidth 5 using Psychtoolbox. See 1 (b). The circles were built of different sizes. The diameter of the circle was 2,4,8 or 16 °. See Figure 1 (a). The eccentricity at which the circle was presented was 1,2,4 and 8 °. See Figure 2. Randomized trials presented every combination of the circle radius and its eccentricity. Every unique combination of eccentricity and radius was presented ten times for a total of 160 trials. The circle was placed left of the central fixation cross half the time and on the right of the fixation cross half of the time. The stimuli was presented against a white background.

Apparatus

Stimuli were generated using the Psychtoolbox (Brainard, 1997) in Matlab on a Windows machine with a Quadro K2200 graphics card and a DELL P2715Q LCD screen. The stimuli were presented on the same machine in the lab. A chin rest was used to ensure stable head position. The monitor resolution was 3840 pixels by 2160 pixels with a refresh rate of 60 Hz. The distance of the observer from the center of the monitor screen was 60 cm. The background was white. The mean luminance of the monitor was 26 cd/m^2.

Psychtoolbox's *Screen* function was used to create stimuli. The distance of the observer from the center of the screen and the size of the screen, both in centimeters, were used to compute the angle in degrees subtended at the user's eye. Using the monitor's resolution

(a)

(b)

Figure 1: (a) Stimuli size: Circles,with inward gradation, whose size (diameter) subtends 2,4,8 and 16 degrees of visual angle respectively on the user's eye. The pictures are for illustration only and show the relative sizes of the stimuli. Stimuli do not subtend the exact degree on the reader's eye. (b) Flashing stimuli: Circle in a black outline and its inward luminance gradation. The outline and the inward gradation alternate at 2Hz. The circle is presented to the left of the fixation cross.

Figure 2: Presentation eccentricity: Circles,with inward gradation, presented at eccentricities of 1,2,4 and 8 degrees of visual angle respectively on the user's eye. The circle's size (diameter) is 4 degrees.

in pixels, the pixel per degree was computed. The *Screen* function was used to draw the appropriate stimuli using the pixel per degree information.

Procedure

Measurement Techniques

The testing took place in a darkened room. Observers used a standard mouse with a middle mouse button and a left button. The participants indicated the polygon to which they adapted, the polygon's orientation and its strength. Once the participant recorded a response, they pressed the middle mouse button on the mouse to move to the next stimuli presentation. The experiment was run in 5 blocks. After a block of 32 stimuli, a break was awarded to the observer. They could resume the next block of trial when ready. The left mouse button was used for all response recording tasks such as selecting the shape, its orientation and the strength of the percept.

Results

The results from the nine observers is shown in Figure 3. On average, most observers report more polygons than circles during the entire experiment. Polygons are reported 61.9 % of the time versus circles that are reported 38% of the time. This means that the formation of polygonal percepts is a phenomenon that occurs in the sampled population. A Student's t-test confirmed no significant difference in tendency to perceive polygons between displays shown to the right or left of observer's fixation (t(x) =T, p=.75). This rules out any effect of anisotropy in peripheral vision, asymmetric foveal confluence influencing shape formation and in the difference between the left and right visual field (Schwartz et al., 1983; Schira, Tyler, Breakspear, & Spehar, 2009). Anisotropy would distort the shape percept on the left and right. However, this test reveals that the emergent percept does not suffer from anisotropic shape distortions based on its presentation within the visual field.

Strength

The experiment measured the strength of a percept on a scale from 0 to 10. 0 meant that the observer did not see a polygon percept after adaptation. The percept in this case remained the same as the circular stimuli. A 10 on the scale indicated a crisp regular polygon. Figure 4 (a) shows that observers do not report a difference between the left and right visual receptive fields when adapting to the stimuli. The strongest effect is for small objects at large eccentricities. See Figure 6 (a).

Large objects at small eccentricities and small objects at large eccentricities produce large effects. Small objects at small eccentricities and large objects at large eccentricities produce smaller effects. This is demonstrated in the co-relation matrix in Figure 6 (a). The minimum contour eccentricity measures the eccentricity of the closest part of the contour from the fovea center or the fixation cross. Figure 6(b) shows that the further the contour is from the fovea, the stronger the effect.

The strength of the stimulus, at the middle of the Likert scale, with relative confidence. Using fitlme, we probed (a) if the users influence the strength of the percept (b) if size, as a fixed effect, influences the strength of the percept (c) if eccentricity, as a fixed effect, influences the strength of the percept (d) if both size and eccentricity, influences the strength of the percept (e) if both size, eccentricity and participant as a random effect influences the strength of the percept. We find the following: (a) Users do have an influence on the

(a)   (b)

Figure 3: (a) Proportion Polygon: Observers report seeing polygons after adaptation 61.94 % of the time and seeing the same circle as the stimulus 38.06% of the time across all the stimuli combinations and trials presented. Error bars indicate the standard error of mean (SEM) around the mean. (b) Probability distribution of the different polygon orders across all participants. Error bars indicate the standard error of mean (SEM) around the polygon order.

strength reported. A linear model with strength as the dependent variable and users as the independent variable shows users effect the strength (F =111.9 , p < 0.05). A one-way Anova on the strength reported by the users shows that users have a large effect $\eta = 0.2516$. (b) Size effects the strength reported $\eta = 1.1836\mathrm{e}\text{-}04$. (c) Eccentricity effects the strength reported $\eta=3.6048\mathrm{e}\text{-}145$ (d) Both size and eccentricity effects the size reported $\eta=0.0175$. (e) Size, eccentricity and users all effect the size reported $\eta = 0.26$. Size and eccentricity offer a better e

One may argue that the percepts are distinct and as there is very little inter subject variance and the percepts are reproducible across participants. Figure As circles are reported at a strength of 0, one can infer from the overall plateaued strength reported for polygons, that the circles are also

Size

The major effect here is stimuli size. Smaller circular stimuli generally show smaller order polygons. Bigger circular stimuli show higher order polygons. This is an interesting feature that indicates that rather than creating large edges between vertices, smaller, more numerous edges are created between numerous vertices.

Eccentricity

Eccentricity has a modest effect on the polygon order reported. Proximity to the fovea shows higher order polygons. This is a smaller effect. See Figure 7(a),(b)

Orientation

We compare observer responses to a uniform distribution of 15 angle increments starting from 0 to the first axis of symmetry for the polygon in question. For most polygon orders, a uniform distribution is observed across all orientations of the polygon. The triangle

Figure 4: (a) Strength Versus Side: Users do not report any significant differences between the left and right visual receptive fields. The same stimuli is presented on the left and the right. (b) The mean strength remains at 2.5 across almost all stimulus size and eccentric presentation. The only exception is observed with the smallest stimulus : size 2 deg that shows certain deviations across users. Users show a wide variation in the strength reported, indicating that there is a lack of uniform percept formation for the smallest stimulus sizes across users but becomes uniform as the stimulus becomes bigger. Error bars indicate the standard error of the mean centered around the mean. (c) Mean strength reported when polygon order indicated is selected as percept. Error bars indicate the standard error of mean around the mean. (d) Mean strength (Y-axis) reported by each participant (X-axis) across all trials of stimuli. The dash line represents the minimum and maximum strength reported. The bars represent standard deviation around the mean.

(a)   (b)

Figure 5: (a) Strength versus size: Users, on average, report the sharpest perception of the polygon when the radius of the polygon subtends 4 degrees of visual angle on the user's eye. (b) Strength versus eccentricity: Users report the sharpest polygon responses when the stimuli is presented at the largest degree (8 deg) of visual angle. Error bars indicate the standard error of mean interval around the mean.

(a)   (b)

Figure 6: (a) A co-relational matrix shows the strength as a function of size and eccentricity (b) Strength versus minimum contour eccentricity: The strength of the percept strengthens as we move away from the fovea. Strength values are means across all participants.

(a)

(b)

Figure 7: (a) Polygon order versus size: Larger order polygons were reported as the radius of the stimuli increased. (b) Polygon order versus eccentricity: In general, observers report lower order polygons at higher eccentricities, larger order polygons at middle eccentricities, followed by slightly lower order polygons at smaller eccentricities. Eccentricity has a modest effect on polygon order. All values reported are means across all participants. The median and mean are shown for eccentricity only since they are not as closely overlapping and show slight user differences.

(a)

(b)

Figure 8: (a) Co-relation matrix showing polygon order (mean) versus size and eccentricity. (b) Polygon order versus minimum contour eccentricity. On average, the minimum contour eccentricity from the fovea only modestly effects the polygon order. Means are over all participants.

showed a relatively high KL divergence value compared to other polygons. With the triangle, observers preferred the 45 ° orientation strongly. Figure 9 shows the responses recorded and the orientation selected for the response polygon.

Preference for contour orientation, where the contour is pointing in a particular angle and direction, is a known phenomenon in mid-level vision, specifically in region V4 (Pasupathy & Connor, 1999a, 2001, 2002). However, the responses recorded do not see any emergent internal polygon angle, after fatiguing of the circular contour, that is angled or oriented as part of its constituent contour shape as shown in (Pasupathy & Connor, 1999a, 2001, 2002). Pasupathy and Connor (Pasupathy & Connor, 1999b) show that there is a neural preference for angles oriented in the 135 ° to 180 ° direction.

Figure 9: Probability versus orientation : For each polygon that was selected by the user, we show the distribution of the orientations selected. The canonical form (0 deg orientation) is when the polygon is centered with its apex at the 12 o'clock position. The responses are computed over all users, shown as a probability of selecting that orientation over all the orientations selected per polygon. In Figure 3(b,c), the first shape in the orientation array is the canonical representation of the triangle and square respectively.

Model that explains the polygon order observed

Figure 10: KL divergence: KL divergence shows how much the orientation differs from a uniform distribution of orientations. When observers select a triangle as the in-place adaptation polygon, they mostly select the 45 degree orientations. Other shapes show a more uniform spread in the orientations selected.

Using regression, we build an equation for the polygon order as a function of the retinal circumference (Equation 2) and as a function of the cortical circumference (Equation 3).

$$y = 6.8995 + 0.0552x, \text{x is a point on the retinal circle} \tag{2}$$

$$y = 5.9936 + 0.0364x, \text{x is a point on the cortical circle} \tag{3}$$

The R^2 value explains how well the model explains the variability in the responses based on the variability in the predictor values. The cortical length's R^2 reports higher value than the R^2 for the retinal length measurements. We used bootstrapping on the observer's responses i.e. polygon order and computed the retinal function and cortical function R^2 values. Among the models that we studied - (a) the skeletal model with Rovamo and Virsu cortical magnification factor and (b) the Harvey model - we found R^2 values are the highest with the skeletal model using the Rovamo and Virsu cortical magnification factors. This model, therefore, becomes our model of choice in explaining the type of polygon seen during adaptation. The p-value measures what proportion of times R^2 is bigger for the retinal than the cortical function. The cortical function R^2 of 1 compared favourably over a retinal function R^2 value of 0. See Figure 12. This proves that our brain samples the curve at points along the circle's circumference which is better explained by a cortical length function.

As evident from regression, a better summary of the relationship between the polygon order seen during adaptation is provided by a cortical function than a retinal function in the brain's region V1 using Rovamo and Virsu's magnification factors. The cortical magnification factors used from lower level V1 (Rovamo and Virsu model) offers a better fit than the cortical magnification factors used from higher regions from the brain such as V4 (Harvey model).

Figure 11: (a) Retinal model: Polygon order regressed over retinal circumference. (b) Rovamo and Virsu cortical model: Polygon order regressed over retinal circumference using Rovamo and Virsu's cortical magnification factor (CMF) for region V1. (c) Harvey cortical model: Polygon order regressed over cortical circumference using Harvey's Cortical Magnification Factor for V4.

(a)

Figure 12: R-squared, from 10000 bootstrapped samples, for retinal circumference function and cortical circumference function. p-value measures by how much R^2 for the retinal circumference function is greater than R^2 for the cortical circumference function. Proportion times retinal is greater than cortical is 0 . Proportion times cortical is greater than retinal is 1 .

Discussion

Several studies have attempted to understand how the curve is mechanistically coded. Curvature built by a neural population is widely favoured because curvature appearance is explained by neurons tuned to different curvatures (Ben-Shahar & Zucker, 2004; Gheorghiu & Kingdom, 2017). Neural populations often define the shapes of afterimages, because it is often the aggregated inhibition responses, collectively that influence what is seen as an afterimage (Gheorghiu & Kingdom, 2017). In our experiment, we show that the unique formation of regular polygons during the adaptation process is a manifestation of the early cortical computation. The simplest mechanistic model that describes the polygon is defined by the equation given by the regression equation,

$$y = 5.9936 + 0.0364x, \text{x is a point on the cortical circle} \tag{4}$$

We make an important observation about scale invariance in this paper. As polygon order changes with changing size, it shows that scale invariance, a distinct property of the visual system, is not obeyed. Scale invariance is a feature of the higher order visual system. A low-level account explains an opponency between localized curvature-tuned neurons in early visual cortex, while the higher-level account assumes an opponency between neurons in higher visual areas coding whole shapes. Because the high-level account that would predict invariance to size and position fails, the low-level account that predicts that the order of the perceived polygon will increase with the size and decrease with the eccentricity of the stimulus, due to cortical magnification, becomes a plausible theory.

Future work can explore the stimuli as gradually increasing arcs of the circle to see how much of the local curvature i.e. arc

circumference is needed to code the polygon seen. El-Shamayleh et. al in (El-Shamayleh & Pasupathy, 2016b) show that scale invariance can be explained by neurons that respond to normalized curvature instead of absolute curvature. Normalized curvature is described as the rate of change in tangent angle per unit of angular length. Absolute curvature is described as the rate of change in tangent angle with respect to contour length. When the shape scales up or down, the normalized curvature remains the same while the absolute curvature changes. A large set of neurons, coding for normalized curvature, maintain selectivity for the shape across size changes whereas a smaller proportion show a shift in selectivity for shape with changing size. Neurons firing for normalized curvature describe orientation that scales with size whereas neurons firing for absolute curvature describe orientation at points described by linear distances. This raises interesting questions about the lower limits of normalized curvature coding by the population neurons and the upper limit of local curvature coding by the population neurons when polygon order changes with size. These limits are exposed during adaptation when the neurons are fatigued in upcoming future work. A limitation of the current experiment is the upper limit on the polygon order offered during shape selection. This will be addressed in future work that allows large polygon order shape selection.

Acknowledgements

The authors would like to thank Bob Ho for setting up the apparatus and Nicholas Baker for graciously reading this manuscript and offering valuable editorial suggestions.

References

Ben-Shahar, O., & Zucker, S. W. (2004). Sensitivity to curvatures in orientation-based texture segmentation. *Vision Research*.

Biederman, I., & Cooper, E. E. (1992). Size Invariance in Visual Object Priming. *Journal of Experimental Psychology: Human Perception and Performance*. Available from http://doi.org/10.1037/0096-1523.18.1.121 , doi:10.1037/0096-1523.18.1.121

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*. Available from http://doi.org/10.1163/156856897X00357

Brincat, S. L., & Connor, C. E. (2004). Underlying principles of visual shape selectivity in posterior inferotemporal cortex. *Nature Neuroscience*.

Daniel, P. M., & Whitteridge, D. (1961). The representation of the visual field on the cerebral cortex in monkeys. *The Journal of Physiology*, *159*(2), 203221.

Das, A., & Gilbert, C. D. (1995). Long-range horizontal connections and their role in cortical reorganization revealed by optical recording of cat primary visual cortex. *Nature*. , doi:10.1038/375780a0

Desimone, R., Albright, T. D., Gross, C. G., & Bruce, C. (1984). Stimulus-selective properties of inferior temporal neurons in the macaque. *Journal of Neuroscience*.

El-Shamayleh, Y., & Pasupathy, A. (2016a). Contour curvature as an invariant code for objects in visual area v4. *Journal of Neuroscience*, *36*(20), 5532–5543. Available from https://www.jneurosci.org/content/36/20/5532 ,

doi:10.1523/JNEUROSCI.4139-15.2016

El-Shamayleh, Y., & Pasupathy, A. (2016b). Contour curvature as an invariant code for objects in visual area V4. *Journal of Neuroscience*.

Gheorghiu, E., & Kingdom, F. A. (2007). The spatial feature underlying the shape-frequency and shape-amplitude after-effects. *Vision Research*.

Gheorghiu, E., & Kingdom, F. A. (2009). Multiplication in curvature processing. *Journal of Vision*.

Gheorghiu, E., & Kingdom, F. A. (2017). Dynamics of contextual modulation of perceived shape in human vision. *Scientific Reports*.

Harvey, B. M., & Dumoulin, S. O. (2011). The relationship between cortical magnification factor and population receptive field size in human visual cortex: Constancies in cortical architecture. *Journal of Neuroscience*. Available from http://doi.org/10.1523/JNEUROSCI.2572-11.2011

Hegdé, J., & Van Essen, D. C. (2000). Selectivity for complex shapes in primate visual area V2. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. Available from http://doi.org/10.1523/JNEUROSCI.20-05-j0001.2000

Hubel, D. H., & Wiesel, T. N. (1959). Receptive fields of single neurones in the cat's striate cortex. *The Journal of Physiology*. Available from http://doi.org/10.1113/jphysiol.1959.sp006308

Hubel, D. H., Wiesel, T. N., & Stryker, M. P. (1978). Anatomical demonstration of orientation columns in macaque monkey. *Journal of Comparative Neurology*. Available from http://doi.org/10.1002/cne.901770302

Ito, H. (2012). Cortical shape adaptation transforms a circle into a hexagon: A novel afterimage illusion. *Psychological Science*, *23*(2), 126–132. Available from http://doi.org/10.1177/0956797611422236

Khuu, B., McGraw. (2002). Misperceptions in the peripheral representation of curvature. *Perception*, *31*(1_suppl), 1-190. Available from https://doi.org/10.1177/03010066020310S101

Lindeberg, T. (2013). Invariance of visual operations at the level of receptive fields. *PLoS ONE*. Available from http://doi.org/10.1371/journal.pone.0066990

Liu, H., Agam, Y., Madsen, J. R., & Kreiman, G. (2009). Timing, Timing, Timing: Fast Decoding of Object Information from Intracranial Field Potentials in Human Visual Cortex. *Neuron*.

Logothetis, N. K., Pauls, J., & Poggio, T. (1995). Shape representation in the inferior temporal cortex of monkeys. *Current Biology*.

Pasupathy, A., & Connor, C. (1999b). Responses to contour features in macaque area V4. *Journal of Neurophysiology*, *82*(5), 2490-2502.

Pasupathy, A., & Connor, C. (2001, Nov). Shape representation in area V4: Position-specific tuning for boundary conformation. *Journal of Neurophysiology*, *86*(5), 2505-2519.

Pasupathy, A., & Connor, C. E. (1999a). Responses to contour features in macaque area V4. *Journal of Neurophysiology*, *82*(5), 2490-2502. Available from http://doi.org/10.1152/jn.1999.82.5.2490

Pasupathy, A., & Connor, C. E. (2002). Population coding of shape in area V4. *Nature Neuroscience*, *5*(12), 1332-1338. Available from

http://doi.org/10.1038/nn972

Rovamo, J., & Virsu, V. (1979). An estimation and application of the human cortical magnification factor. *Experimental Brain Research*.

Rust, N. C., & DiCarlo, J. J. (2010). Selectivity and tolerance ("invariance") both increase as visual information propagates from cortical area V4 to IT. *Journal of Neuroscience*.

Sakurai, K. (2014). Shape distortion illusion of circles without prolonged viewing. *Journal of vision*, *14 10*, 247. Available from http://doi.org/10.1167/14.10.247

Sakurai, K., & Beaudot, W. (2015). Shape distortion illusion of flashed circles can be induced by dichoptic stimulation. *Journal of Vision*. Available from http://doi.org/10.1167/15.12.528

Schira, M. M., Tyler, C. W., Breakspear, M., & Spehar, B. (2009). The foveal confluence in human visual cortex. *Journal of Neuroscience*. , doi:10.1523/JNEUROSCI.1760-09.2009

Schrauf, M., Lingelbach, B., & Wist, E. R. (1997). The scintillating grid illusion. *Vision Research*. , doi:10.1016/S0042-6989(96)00255-6

Schwartz, E. L., Desimone, R., Albright, T. D., & Gross, C. G. (1983). Shape recognition and inferior temporal neurons. *Proceedings of the National Academy of Sciences of the United States of America*.

Spillmann, L. (1994). *The Hermann grid illusion: a tool for studying human perspective field organization*.