Eco-evolutionary dynamics in two-species mutualistic systems: One-sided population decline triggers joint interaction disinvestment

Franz Weyerer[1]
Avril Weinbach[2]
Christiane Zarfl[1]
Korinna T. Allhoff[1,2]

1) Eberhard Karls Universität Tübingen, Tübingen, Germany
2) Universität Hohenheim, Stuttgart, Germany

Corresponding author: Dr. Korinna Theresa Allhoff, University of Hohenheim, Institute of Biology, FG Eco-Evolutionary Modelling (190m), Garbenstr. 30, 70599 Stuttgart, Germany. Email: korinna.allhoff@uni-hohenheim.de

ORCIDs: Allhoff (0000-0003-0164-7618); Zarfl (0000-0002-2044-1335); Weinbach (0000-0001-7598-1112)

Running title: Joint interaction disinvestment in mutualistic systems

Key words: Evolutionary rescue, evolutionary murder, coevolution, mutualism, insect decline, adaptive dynamics.

Total word count (excluding references): 6785

Number of figures and tables: 6 Figures and 1 Table

Items to be published as online appendix: Simulation code and summarising pdf document containing 4 additional figures.

Acknowledgments: We thank all members of the Eco-Evolutionary Modelling group at the University of Hohenheim for many inspiring discussions throughout the whole project as well as valuable feedback on the final manuscript. We furthermore thank Fynn Bachmann for assistance with programming.

Abstract:

The interplay between ecological and evolutionary dynamics can create feedback that reinforces external disturbances and potentially threatens species' coexistence.

40 For example, plants might invest less into attracting insect pollinators (decreased flower or nectar production) and more into independence (increased selfing or vegetative reproduction) when faced with pollinator decline. This adaptive response saves plant resources at the cost of further threatening the pollinator population. We ask under which conditions such self-reinforcing feedback occurs in two-species

45 mutualistic systems when considering one-sided population decline and whether it can be counteracted by self-dampening feedback if co-evolution of both interaction partners is considered. Based on a mathematical model and in line with previous studies, we find that the described pattern of accelerated population decline occurs for a wide range of parameter values if a concave allocation trade-off between

50 independent growth and interaction investment is assumed. The undisturbed population typically disinvests first, which then forces the declining population to also disinvest, in favour of other energy sources. However, a decelerated population decline can occur if the adaptation of the undisturbed partner is relatively slow compared to environmental decay, reducing the speed of its disinvestment, or if the

55 initial investment into the interaction was very high. Our results suggest that if actions are taken to save endangered populations, not only the evolution of the target species but also of their interaction partner, as well as the interaction between them should be considered.

1. **Introduction**

Insects experience stressors from many directions (Wagner et al., 2021), be it the fragmentation, degradation, and destruction of habitats (Fabienne Harris & Johnson, 2004), the enhanced use of pesticides (Hoffmann et al., 2010), invasive species (Madjidian et al., 2008), new diseases (Le Conte et al., 2010) or the effects of climate change in general. All these factors have led to a decline in the diversity and biomass of flying insects (Fox, 2013; Hallmann et al., 2017; Potts et al., 2010). This is not only a problem for the insect populations themselves but also for all species that interact with them, either antagonistically or mutualistically. Mutualistic interactions, which play a crucial role in sustaining ecosystem functioning and services (Reid et al., 2019), are of special importance in this context, given that they are less stable than antagonistic ones and more prone to be lost (Sachs & Simms, 2006; Toby Kiers et al., 2010).

The stressors mentioned above put insect populations under enormous pressure to adjust to new environmental conditions, e.g. via phenotypic plasticity or adaptation, which in turn forces their mutualistic interaction partners to adjust as well. In a worst-case scenario, the adaptation of one species might be too slow to catch up with their interaction partner and their dependence on each other could then lead to co-extinction. In other cases, the cost of maintaining the mutualistic interaction might simply increase under the new environmental conditions, meaning that the interaction itself could become subject to change in case the costs ultimately outweigh the benefits (Bronstein, 2001). This can lead to a mutualism loss, a partner switch, or even change the type of interaction from mutualism to antagonism (Toby Kiers et al., 2010).

Adaptive and evolutionary responses to external stressors are not limited to timescales of hundreds or thousands of years but can occur surprisingly fast (Cheptou et al., 2008; Hairston et al., 2005; Thompson, 1998). For example, Roels and Kelly (2011) showed that plants (*Mimulus guttatus*) increased their self autonomous reproduction in response to loss of pollinators (*Bombus impatiens*) within only a few generations. This trend is typically associated with smaller flowers, reduced flower longevity, and fewer rewards for the pollinator (Thomann et al., 2013). On the other hand, Gervasi and Schiestl (2017) showed that after

I

exposing plants (*Brassica rapa*) to pollinators (*Bombus terrestris)*, the plants evolved to higher investment in plant height and amount of volatile emission to attract the pollinator. Both plant responses potentially reinforce the initial change in pollinator abundance by decreasing or increasing food resources, creating so-called eco-evolutionary feedback (Pelletier et al., 2009; Post & Palkovacs, 2009).

Quantifying the impact of such eco-evolutionary feedback on system dynamics and species survival is very challenging, especially in systems that contain more than one population (Koch et al., 2014; N. Loeuille, 2019), but the mathematical framework of adaptive dynamics can help to gain valuable insights (Brännström et al., 2013; Dieckmann & Law, 1996; Ferriere & Legendre, 2013; Geritz et al., 1996; Govaert et al., 2019). For instance, Weinbach et al. (2022) used this method to investigate eco-evolutionary dynamics in a simple Lotka-Volterra mutualism system. They studied this system through the lens of plant-pollinator interactions. One of the interaction partners, which was interpreted as the pollinator, experienced a sudden decline in intrinsic growth due to external stressors. The other interaction partner, which was interpreted as a plant, was then able to react to changes in pollinator abundances by shifting its energy allocation. It could either invest into attractiveness for the pollinator, in order to maintain the interaction, or into its own intrinsic growth, in order to become more independent. The model suggests that plants tend to disinvest from the interaction in the face of pollinator decline, in line with the empirical evidence from Roels and Kelly (2011) mentioned above, if a concave trade-off function was assumed. This disinvestment could then further threaten the already declining pollinator population, resulting in self-reinforcing eco-evolutionary feedback. The authors hypothesised that plant disinvestment could even result in so-called eco-evolutionary murder (name suggested by Parvinen in 2005), meaning that the pollinator does not go extinct as a direct consequence of an external stressor but due to plant evolution.

It remains unclear, however, whether this key finding is robust to changes in the underlying model assumptions. This is in particular relevant when considering co-evolution of both interaction partners. We hypothesise that if pollinators were also allowed to re-adjust their energy investment in response to changing conditions, similar to the plants, then this might counteract the impact of external stressors. In consequence, we would observe self-dampening instead of self-reinforcing eco-evolutionary feedback and hence decelerated instead of accelerated pollinator

decline. Pollinator evolution could even enable so-called evolutionary rescue (S. M. Carlson et al., 2014; Ferriere & Legendre, 2013, 2013; Gomulkiewicz & Holt, 1995; Gonzalez et al., 2013), meaning that pollinators are able to survive
130   environmental conditions that would lead to extinction in the absence of evolution.

To test this hypothesis, we extended the model by Weinbach et al. to account for coevolution of both interaction partners. More specifically, we analysed the evolution of the investments in the mutualistic interaction of both species in response to the declining abundance of one interaction partner. By design, the
135   declining interaction partner sooner or later reached the extinction threshold. We asked under which conditions the self-reinforcing feedback found by Weinbach et al. (2022) dominates system dynamics, resulting in earlier extinction, and what conditions have to be met to observe the reversed outcome, that is delayed extinction. To address these questions, we varied ecological model parameters, in
140   particular interaction benefits and intraspecific competition, as well as the relative timescales of the environmental decay versus the evolutionary responses of both interaction partners.

2. __Model & Methods__

In the following, we first give an overview of the ecological processes considered in the model before explaining how population decline due to environmental decay is implemented. We then specify how evolutionary changes in energy investment are taken into account and how all three model components (ecology, environment and evolution) are linked within our numerical algorithm. A summary of all model parameters can be found in Table 1.

2.1 Ecological dynamics

As in Weinbach et al. (2022) the basic model consists of two interacting species. The model is relatively abstract and not system-specific, meaning that it could in principle describe any mutualistic system (e.g. plant-mycorrhizal, plant-ant, or ant-fungi systems, see Bronstein (1994). However, we refer to the two interaction partners as plant population P and pollinator population A [$\#indv\cdot m^{-2}$]. The population dynamics are given by a Lotka-Volterra-type model with two differential equations:

$$\frac{dP}{dt} = r_P(\alpha)P - c_P P^2 + \alpha\beta\gamma_A PA \tag{1a}$$

$$\frac{dA}{dt} = r_A(\beta)A - c_A A^2 + \alpha\beta\gamma_P PA \tag{1b}$$

The equations consist of three terms each. The first term describes the independent growth of the populations, where r_P and r_A represent the net growth coefficients in [1/time] of the plant and the pollinator, respectively. The plant coefficient includes processes like selfing or vegetative reproduction that allow for growth even in the absence of pollinators. The pollinator coefficient, on the other hand, includes not only energetic gain from alternative food sources, reproduction spots or shelter, but also mortality losses. It can become negative, if those losses outweigh the positive terms. Both coefficients are dependent on the corresponding interaction parameters α or β via trade-off functions, as explained in more detail further below.

The second term describes the intraspecific competition for both plants and pollinators, with coefficients c_P and c_A in [$m^2\cdot(\#indv\cdot time)^{-1}$], respectively. Competition is assumed to be at a constant level, meaning that evolution and also

environmental decay do not change the intensity of the competition, in line with the original model by Weinbach et al. (2022). The first two terms combined give logistic growth with carrying capacities r_i/c_i for the plant ($i=P$) and the pollinator ($i=A$).

175   The last term finally defines the mutualistic interaction between plants and pollinators. More precisely, γ_A (measured in [$m^2 \cdot (\#indv \cdot time)^{-1}$]) describes the interaction benefits for the plant (e.g. sexual reproduction due to the fertilisation or protection against herbivory provided by the mutualist interactor), while γ_P corresponds to the benefits for the pollinator (e.g. energetic gain due to nectar,

180   pollen, or other plant exudates provided by the plant). The parameters α and β (dimensionless) capture the investment of the plant and the pollinator into the interaction. On the plant side, this might describe the amount of nectar or size, shape, colour or number of flowers (Willmer, 2011). On the pollinator side, this could represent the pollinator's morphology, such as adapted mouthparts, or the

185   designated time to search for this particular plant. Note that without β the model simplifies into the original model version by Weinbach et al. (2022).

The basic ecological dynamics of the model have already been investigated by Weinbach et al. (2022), but the results have to be adjusted to the introduction of β, as explained in the following. A stable equilibrium for ecological coexistence can

190   be found at the crossing point of the non-trivial zero-growth isoclines (see online supplementary material A for more details). The corresponding equilibrium population densities can be calculated via the following equations:

$$A^* = \frac{\alpha \beta \gamma_P r_P + r_A c_P}{c_A c_P - \alpha^2 \beta^2 \gamma_A \gamma_P} \tag{2a}$$

$$P^* = \frac{\alpha \beta \gamma_A r_A + r_P c_A}{c_A c_P - \alpha^2 \beta^2 \gamma_A \gamma_P} \tag{2b}$$

The equilibrium point is only feasible for positive population densities. This condition is fulfilled, assuming that all parameters are positive, if the denominator

195   is positive:

$$c_A c_P > \alpha^2 \beta^2 \gamma_A \gamma_P \tag{3}$$

The above inequation can be interpreted as a balance between intraspecific competition (left) and interspecific mutualism (right). If equation (3) is not true, the self-regulatory effect of the intraspecific competition is too weak to compensate for the positive feedback that arises from the mutualistic interaction, resulting in uncounded population growth. Such unlimited growth is usually controlled by additional processes, such as predation or interspecific competition, which are not explicitly considered in the model. Instead, Weinbach et al. defined a critical level for plant interaction investment α_{cl}. The same is done here for the critical level of pollinator interaction investment β_{cl}:

$$\alpha_{cl} = \sqrt{\frac{c_A c_P}{\gamma_A \gamma_P \beta^2}}$$

(4a)

$$\beta_{cl} = \sqrt{\frac{c_A c_P}{\gamma_A \gamma_P \alpha^2}}$$

(4b)

Stable coexistence of both populations is thus only possible if $0 \leq \alpha \leq \alpha_{max} < \alpha_{cl}$ and $0 \leq \beta \leq \beta_{max} < \beta_{cl}$. Weinbach et al. set $\alpha_{max} = 0.8 * \alpha_{cl}$. To keep the existing symmetry between both populations, we define $\beta_{max} = 0.8 * \beta_{cl}$.

2.2 Environmental decay

We assume a steady decline in habitat quality, which directly affects the mortality of pollinators, so that the net growth rates of pollinators continuously decrease. We furthermore assume that this decrease can be described via the following equation:

$$r_A = r'_A - t \cdot \delta_{Env}$$

(5)

The parameters r'_A and δ_{Env} denote the initial growth rate at time t=0 and the change in pollinator mortality rate per unit time, respectively. Note that $r_A = r'_A$ in the absence of environmental decline, that is when $\delta_{Env} = 0$. In all other cases, we can expect that the habitat quality will eventually become so bad that pollinators are no longer able to survive. Simulations are stopped as soon as the pollinator population density falls below the extinction threshold ε.

`

2.3 Evolution of interaction investment

Both plants and pollinators are assumed to have a finite amount of energy that they can invest either into independence from each other (via the growth rates r_P or r'_A) or into the joint mutualistic interaction (via the investment into the interaction α or β). We use the same trade-off function that Weinbach et al. (2022) introduced for the plant also for the pollinator:

$$r_P = \left(1 - \left(\frac{\alpha}{\alpha_{max}}\right)^s\right)^{1/s} \tag{6a}$$

$$r'_A = \left(1 - \left(\frac{\beta}{\beta_{max}}\right)^s\right)^{1/s} \tag{6b}$$

The exponent s in these trade-off functions controls the trade-off shape. Weinbach et al. (2022) found that the plant evolution strongly depends on this parameter. Small values of s translate into a convex trade-off shape (see supplementary Fig. B1). Plants with such a trade-off tend to evolve towards the extreme cases, that is strategies where either attractiveness or independent growth is absent (either $\alpha=0$ and $r_P=1$ or $\alpha=1$ and $r_P=0$). A balanced strategy with intermediate investment into both attractiveness and independent growth can only be expected for relatively large values of s, which translate into a concave trade-off shape (see supplementary material B). Weinbach et al. (2022) showed that in these cases, the plant could indeed maintain both the mutualistic interaction and selfing or vegetative reproduction and that the pollinator could coexist with the plant for a large range of net growth rate coefficient values (r_A). In the following analysis, we therefore chose *s=3*.

Which trait values turn out to be optimal for the plant and its pollinator in a given situation obviously depends on plant and pollinator abundances, as well as on the strength of the mutualistic interaction. For example, with limited pollination service due to low pollinator densities, the plants might invest less into attractiveness (low value of α) and more into intrinsic growth (high value of r_P). Similarly, with low food availability due to low plant densities and/or low attractiveness, the pollinator might also invest less into the interaction (low value of β) and more into other energy sources (high value of r_A). In the face of declining pollinator abundances, both populations will consequently adapt by shifting their energy allocation, which would in turn change the strength of the mutualistic interaction and hence either accelerate or decelerate the initial pollinator decline.

2.3 Numerical simulation of eco-evolutionary dynamics

Our analysis of adaptive changes in energy allocation of plants and pollinators in response to pollinator decline is inspired by the mathematical framework of Adaptive Dynamics (see Brännström et al. (2013) for a review). However, with two
255 evolving traits, namely α and β, analytical calculations using Adaptive Dynamics are typically very complicated (see Dieckmann and Law (1996) for the general method and Kisdi (2006) for an application). Our investigation of the eco-evolutionary dynamics within our model was therefore exclusively based on numerical simulations. Simulations were programmed in Python 3.10. The
260 simulation code is available via as supplementary material.

Each simulation started with a single ancestor plant population and a single ancestor pollinator population at an eco-evolutionary equilibrium. We then performed recurrent "mutation events". During such a mutation event, we chose one of the populations as "resident" and simply added a new "mutant" population.
265 The initial population density of the mutant equalled the extinction threshold ε and was taken from the resident population. The traits of the mutant population were chosen from a normal distribution around the parent's traits. This normal distribution had a standard deviation of 1/2.5758, so that 99% of all randomly chosen numbers were in the interval [-1,1], and was multiplied with $\Delta\alpha$ and $\Delta\beta$,
270 respectively, so that $\Delta\alpha$ and $\Delta\beta$ effectively describe the maximum mutation distance between resident and mutant. The initial per-capita growth rate of the mutant was derived via the so-called "invasion fitness", that is the change in mutant density over time under the assumption that the mutant m is rare and that the resident r is at equilibrium densities P^*_r or A^*_r, respectively:

$$\frac{d P_m}{dt}\frac{1}{P_m}=r_{P_m}\big(\alpha_m\big)-c_P\big(P_m+P^*_r\big(\alpha_r\big)\big)+\alpha_m\beta_r\gamma_A A^*_r\big(\beta_r\big) \tag{7a}$$

$$\frac{d A_m}{dt}\frac{1}{A_m}=r_{A_m}\big(\beta_m\big)-c_A\big(A_m+A^*_r\big(\beta_r\big)\big)+\alpha_r\beta_m\gamma_P P^*_r\big(\alpha_r\big) \tag{7b}$$

275   The fate of the mutant population was determined by its traits, which affect the strength of the mutualistic interaction, as well as the state of the environmental decay. The mutant could (i) have a positive growth rate and eventually replace the resident population in case it was better adapted to the current situation, (ii) have a negative growth rate so it quickly went extinct or (iii) coexist with the resident. The

280   latter outcome is known as evolutionary branching and can, in principle, lead to the emergence of complex interaction networks (Brännström et al., 2011; Geritz et al., 1996; Nicolas Loeuille & Loreau, 2005). However, in our system we typically observed outcome (i) and (ii).

   We used two different implementations of the algorithm outlined above. Following

285   the core hypothesis of the Adaptive Dynamics framework, the first approach was based on a separation of ecological and evolutionary time scales. In particular, we assumed that mutation events are so rare that the population dynamics of the resident will always reach an ecological equilibrium before the next mutation event takes place. To shorten the simulation runtime, we therefore omitted the

290   calculation of the population dynamics. Instead, we instantly replaced the resident population with the mutant population if the initial per-capita growth rate of the mutant was positive (in line with outcome (i)). By contrast, mutants with negative growth rates were assumed to be not viable and simply removed from the system (in line with outcome (ii)). The code for this first model version thus includes

295   equations (2a) and (2b) for the equilibrium densities, equation (5) for the environmental decay, equations (6a) and (6b) for the trade-off functions, as well as equations (7a) and (7b) for the per-capita growth rates of new mutant populations, as summarised in Fig 1.

   In a second approach, we then relaxed the assumption of separate ecological and

300   evolutionary timescales. The corresponding algorithm is summarised in supplementary Fig. C1. Instead of instantly replacing the resident with the mutant population, as explained above, we simulated the explicit resident and mutant population dynamics over time. However, mutation events were still performed

with fixed probabilities q_A and q_P, respectively, as in the first approach, meaning that the establishment of the previous mutant was typically not completed when the next mutant was added. In consequence, we observed multiple mutant populations of plants and/or pollinators apparently coexisting over time. For computational reasons, at any given time we limited the number of plant and/or pollinator mutants to a maximum of 25 coexisting populations. Population dynamics were implemented using the following set of equations, with indices running over the multiple mutant populations, to accommodate a growing (or shrinking) number of populations:

$$\frac{dP_i}{dt} = r_{P_i}(\alpha_i)P_i - c_P P_i \sum_v P_v + \alpha_i \gamma_A P_i \sum_w \beta_w A_w \tag{8a}$$

$$\frac{dA_j}{dt} = r_{A_j}(\beta_j)A_j - c_A A_j \sum_w A_w + \beta_j \gamma_P A_j \sum_v \alpha_v P_v \tag{8b}$$

Simulations performed with the first approach were very fast at the cost of limited biological realism. Simulations based on the second approach, on the other hand, were more realistic, but also very time consuming, due to the high number of coupled equations. We therefore performed most of our analysis using only the first approach and then repeated only part of the analysis using the second one, as a robustness check. In both cases, we compared the extinction time points of pollinators with and without evolution for various parameter combinations, to analyse the impact of eco-evolutionary feedback on pollinator persistence.

`

Fig. 1: Overview of the model algorithm. The ecological part (in blue) calculates the ecological equilibrium densities, the environmental part (in green) implements the population decline via environmental decay and the evolutionary part (in orange) implements evolution for both the plant and the pollinator populations. Most simulations started at an eco-evolutionary equilibrium and were run with the environmental degradation being active. The end condition was then simply whether or not the pollinator was still alive. Note that the algorithm can also be run without environmental decline, with the goal to pinpoint the initial eco-evolutionary equilibrium. In these cases, the end condition was either a fixed time limit or whether the eco-evolution dynamics already reached an evolutionary endpoint that is both convergent stable and evolutionarily stable (a so-called CSS).

Table 1: Overview of model parameters. Base values correspond to the initial eco-evolutionary equilibrium used in reference simulation S1. Environmental decay rate refers to simulations with separated timescales.

Definition	Symbol	Unit	Base value	Analysed range
Ecological parameter				
Plant net growth coefficient	r_P	time^{-1}	0.66	0 – 1
Pollinator net growth coefficient	r_A	time^{-1}	0.66	0 – 1
Pollinator growth without env. decay	r'_A	time^{-1}	0.66	-∞ - 1
Plant intraspecific competition	c_P	m^2·(#$indv$·time)$^{-1}$	1.00	0.65–2.15
Pollinator intraspecific competition	c_A	m^2·(#$indv$·time)$^{-1}$	1.00	0.65–2.15
Plant interaction investment	α	1	0.72	0 – 0.80
Pollinator interaction investment	β	1	0.72	0 – 0.80
Plant energic gain	γ_A	m^2·(#$indv$·time)$^{-1}$	1.4	0 – 1.50
Pollinator energic gain	γ_P	m^2·(#$indv$·time)$^{-1}$	1.4	0 – 1.50
Environmental parameter				
Environmental decay constant	δ_{Env}	time^{-2}	0.001	0.1-0.00001
Extinction threshold	ε	#$indv$·m^{-2}	0.01	0.01
Evolutionary parameter				
Plant mutation probability	q_P	1	0	0 – 1
Pollinator mutation probability	q_A	1	0	0 – 1
Plant mutation maximum step size	$\Delta\alpha$	1	0.016	0.016
Pollinator mutation maximum step size	$\Delta\beta$	1	0.016	0.016
Trade-off shape	s	1	3	3

3. Results

3.1 Delayed pollinator extinction can occur for high interaction benefits, low competition and slow plant evolution

The pollinator population in our simulations is always doomed to go extinct, due to ever increasing mortality rates, see eq. (5), but the exact time point of extinction depends on the involved eco-evolutionary forces. We found that eco-evolutionary feedback could in principle delay pollinator extinction and thus help those pollinators to survive harsher conditions. This was in particular true for relatively high interaction benefits α and β and low competition c_P and c_A (blue area in Fig. 2). Most of the parameter combinations, however, resulted in self-reinforcing eco-evolutionary feedback that accelerated pollinator extinction (red area in Fig. 2). Note that extremely low interaction benefits and very intense competition led to a rather weak interaction between relatively small populations, which explains why in these cases the effect of eco-evolutionary feedback was less pronounced. Extremely high interaction benefits in combination with very low competition, on the other hand, led to a strong coupling of large populations, which pushed the systems towards unrealistic unbounded growth, as explained in the methods section (see eq. 3).

Not only the ecological parameters had an effect on the eco-evolutionary forces but also the relative speed of plant versus pollinator adaptation (Fig 3). We typically observed earlier pollinator extinction in the extreme case of only plant evolution ($q_A=0$, $q_P>0$, see x-axis in Fig. 3a-c). Delayed extinction was possible in the other extreme case, when plant evolution was switched off, in particular when the interaction benefit was sufficiently high ($q_P=0$, $q_A>0$, see y-axis in Fig. 3a-c). In the case of co-evolving interaction investments, we found that delayed extinction became more likely when plant evolution was decelerated or when pollinator evolution was accelerated.

The effect of evolutionary adaptation on pollinator survival strongly depends on how fast the evolutionary response is in comparison to the environmental decay. In general, we found that delayed pollinator extinction is more likely and/or more pronounced for faster environmental decay, especially if plant evolution is sufficiently slow (Fig 4). In particular, we found that delayed pollinator extinction is possible in a rapidly changing environment, even if pollinator evolution was switched off in the model structure (Fig. 4c).

Fig. 2: High interaction benefits and low competition can lead to delayed pollinator extinction. The colour code indicates the observed extinction time point for the given parameter combination relative to the extinction time point of the corresponding simulation without evolution in [%]: earlier extinction is shown in red, delayed extinction in blue. White areas indicate no effect of eco-evolutionary feedback on pollinator extinction. Mutation probabilities were $q_P = 0.1$ for the plants and $q_A = 0.5$ for the pollinators. All other parameters were chosen according to Table 1, if not indicated otherwise. The time series for the simulation highlighted as S1 can be found in Fig. 5.

Fig. 3: Delayed pollinator extinction is more likely with slow plant and fast pollinator evolution. The colour code indicates the observed extinction time point for the given parameter combination relative to the extinction time point of the corresponding simulation without evolution in [%], as in Fig 2. Parameter values were chosen according to Table 1, if not indicated otherwise. The time series for the simulations highlighted as S1-S4 can be found in Fig. 5.

Fig. 4: Delayed pollinator extinction is more likely with fast environmental decay. The colour code indicates the observed extinction time point for the given parameter combination relative to the extinction time point of the corresponding simulation without evolution in [%], as in Fig 2. Parameter values were chosen according to Table 1, if not indicated otherwise. The time series for the simulations highlighted as S2-S5 can be found in Fig. 5.

3.2 Plants disinvest first, pollinators follow

In our simulation, we always observed the same two processes, namely disinvestment of the plants and disinvestment of the pollinators (Fig. 5). Increasing pollinator mortality rates primarily affected the pollinators themselves and resulted in declining pollinator abundances (top row, yellow lines). The plant population then responded to such declining abundances by reducing its investment into the interaction (decreasing values of α, bottom row, green lines) in favour of intrinsic growth (increasing values of r_P, not shown). The plant disinvestment slowed down the plant decline that was triggered by the reduced interaction (top row, green line) at the cost of further threatening the pollinator population. The pollinator consequently responded to this double pressure of increasing mortality rates and decreasing interaction strength by also reducing its investment into the interaction (bottom row, yellow lines). This disinvestment then allowed the pollinator to partly compensate for increasing mortality rates (middle row), which in turn also slowed down pollinator decline (top row, yellow lines). The disinvestment of both interaction partners was typically rather slow in the beginning but accelerated in during the simulation (bottom row in Fig 5), which was partly due to the fact that both processes reinforce each other but also reflects the concave shape of the chosen trade-off functions (see eq. 6).

The first process (plant disinvestment) was in most cases harmful for the pollinator, as it weakened the mutualistic interaction, while the second process (pollinator disinvestment) typically helped them to survive longer. Whether or not the combined effect of both processes translated into earlier or delayed pollinator extinction thus strongly depended on their relative strength, which was modified via differences in the mutation frequency (see Fig. 3). In particular, we found that delayed extinction can turn into earlier extinction if plants evolve faster (compare S1 and S2 in Fig 5). However, also the initial situation at simulation start played a key role, which became apparent when analysing simulations with high interaction benefits and/or low competition (see Fig. 2 and 3). These simulations started with a very strong interaction between large populations, indicated via relatively high initial population densities P and A (see top panel for S3 in Fig 5) and high initial interaction investments α and β (bottom panel for S3). Plants then needed more time to actually disinvest, simply because they started at a higher level of

interaction investment α, so that the mutualistic interaction was maintained longer.

440 At the same time, the beneficial effect of pollinator disinvestment was particularly strong, due to the concave shape of the trade-off function (eq. 6): small mutation steps led to a substantial increase in intrinsic growth r_A as long as the interaction investment β was still high, which in turn directly compensated for increasing mortality rates. In consequence, we found that the beneficial effect of pollinator

445 disinvestment typically dominated when the initial interaction was very strong, resulting in delayed extinction, even if both interaction partners had the same mutation frequency ($q_A = q_P$ in S3).

The delayed extinction observed in S3 turned into earlier extinction after switching off pollinator evolution, indicating that plant evolution is indeed harmful for the

450 pollinator (S4). However, plant evolution could also be beneficial, in particular when the environmental decay was relatively fast (S5). This surprising result is again due to the shape of the concave trade-off function (eq. 5): small mutation steps led to a substantial increase in intrinsic growth r_P as long as the investment α was still high. This had a direct positive effect on plant abundance, which could

455 then indirectly benefit the pollinators as long as the mutualistic interaction was still strong. Note, however, that the positive effect of plant evolution was negligible compared to the negative effect of increased environmental decay, meaning that the pollinators still went extinct much earlier compared to scenarios with moderate or slow environmental decay.

Fig. 5: Overview of selected time series. Background colours highlight ecological (blue), environmental (green), and evolutionary (orange) processes. Vertical lines indicate pollinator extinction time points. Note that plant densities always approach 1, which equals the plant carrying capacity $1/c_P$ with $c_P=1$, that is the equilibrium density of the plant in the absence of pollinators. Dotted lines show reference simulation without evolution ($q_P=q_A=0$). Simulation S1 was run with interaction benefits $\gamma_P = \gamma_A = 1.4$ and mutation probabilities $q_P=0.1$ for the plant and $q_A=0.5$ for the pollinators. All other parameters were chosen according to Table 1. We then changed one parameter at a time, while keeping all other parameters constant. From S1 to S2: increase plant mutation probability ($q_P=0.5$ instead of 0.1). From S2 to S3: increase interaction benefit ($\gamma_P=\gamma_A=1.5$ instead of 1.4). From S3 to S4: switch off pollinator evolution ($q_A=0$ instead of 0.5). From S4 to S5: increase speed of environmental decay ($\Delta_{Env}=0.02$ instead of 0.001).

3.3 Robustness checks

We acknowledge that the model outcomes presented above are to some extent dependent on the specific trade-off function. In particular, the observed eco-evolutionary dynamics might accelerate or slow down when choosing a higher or lower value of the parameter s, which captures the trade-off shape. However, we

485   still find the same qualitative pattern of joint disinvestment if s is sufficiently large, meaning that the trade-off function is sufficiently concave. For more detailed information of this matter please refer to the methods section and to the online supplementary Fig. D1 and D2.

The results presented so far are all based on a simplified evolutionary algorithm

490   that allowed us to perform many simulations within a reasonable time span. The algorithm is based on the assumption that the ecological and evolutionary time scale are separate (see Methods section). As a final step, we relaxed this assumption by using a slower, but more realistic algorithm that captures overlapping timescales. The results obtained with both model versions were

495   generally consistent and differed only slightly in the exact parameter combinations for which delayed extinction could be observed (compare Fig 3 and 6). In particular, the critical value of the interaction benefit γ at which the system moved from delayed to earlier extinction tended to be higher in the simplified model version compared to the extended version. Also, the effect sizes varied, meaning

500   that the maximum possible delay of extinction was larger in the simplified model version. We explain these differences via a time lag between the 'optimal' evolutionary strategy of the plants, as predicted based on the assumption of separate time scales, and the evolutionary strategy observed during simulations with overlapping timescales. In the latter case, the plants simply did not have

505   enough time to fully adjust their energy allocation strategy to the current environmental condition, meaning that the mutualistic interaction was maintained longer, which in turn lessened the detrimental effect of plant evolution on pollinator survival.

Fig. 6: Heatmaps with overlapping timescales and intraspecific variability in resource allocation. The colour code indicates the observed extinction time point for the given parameter combination relative to the extinction time point of the corresponding simulation without evolution in [%], as in Fig 2. Environmental decay speed was $\delta_{Env}=0.0001$; Mutation events took place every 25 timesteps; Other parameter values were chosen according to Table 1, if not indicated otherwise.

4. **Discussion**

In this study, we investigated how eco-evolutionary dynamics within a mutualistic two-player system shape the system's response to one-sided population decline. Our model analysis revealed two important key insights. First, in all scenarios that were studied a joint disinvestment of both interaction partners in response to the decline of one interaction partner was found. Second, despite the generality of the first key finding, it still remains difficult to predict whether such joint disinvestment eventually translates into delayed or earlier pollinator extinction. This is due to a highly non-trivial interplay between the ecological, environmental and evolutionary processes considered in the model, as well as the relative speed of these processes. In summary, we found that delayed pollinator extinction can occur if the interaction benefits are sufficiently high, if intraspecific competition is sufficiently low and if plant evolution is relatively slow.

The observed joint disinvestment is surprising at first, because one might naively assume that at least one interaction partner would invest more into the interaction, not less, in order to maintain its benefits. However, the disinvestment can be easily explained as a direct consequence of the assumed energy allocation trade-off. This trade-off introduces a cost to the mutualistic interaction, meaning that maintaining interaction benefits goes hand in hand with losing other energy sources, such as alternative food sources, reproduction spots or shelter. These alternative resources become relatively more important as soon as the mutualistic interaction is weakened due to pollinator decline, leading to a shift in energy allocation in favour of those other resources, which in turn further weakens the interaction.

Those scenarios in which joint disinvestment led to delayed pollinator extinction provide potential for evolutionary rescue (see Gonzalez et al. (2013) or Bell (2013, 2017): If the pollinator mortality rate does not continue to increase forever, as in our study, but instead reaches at a value that is just high enough to push the pollinators below the extinction threshold in the absence of evolution, then those pollinators might still be able to survive when evolution is switched on, due to eco-evolutionary feedback. Note that in the past, evolutionary rescue theory has mostly been applied to single populations (see for example the summarising reviews of C. Carlson et al. (2022) and Bell (2017)), while our model demonstrates that the

550   concept of evolutionary rescue can be extended towards multi-species systems: The survival of a target species does not only depend on its own evolutionary trajectory, but also on adaptive responses of their interaction partners. This is in line with findings by Yamamichi and Miner (2015), who showed that a non-evolving predator population can be rescued from extinction by adaptive evolution

555   of its prey, resulting in indirect evolutionary rescue. It is also in line with recent work by (Yacine et al., 2021), who analysed body mass evolution within model food webs. Their simulations suggest that long-term biodiversity losses triggered by warming are considerably higher when evolution is slowed down or switched off completely, pointing towards evolutionary rescue at the community scale.

560   However, most scenarios that were studied here actually resulted in accelerated pollinator extinction, which provides potential for evolutionary murder instead of evolutionary rescue (as suggested by Parvinen (2005)): If pollinator mortality rates increase only until they reach a value that is still smaller than the critical value at which pollinators go extinct in the absence of evolution, then pollinator survival

565   would nevertheless be threatened when evolution is switched on, again due to eco-evolutionary feedback. These scenarios consequently challenge the naive assumption that evolution is generally beneficial for species survival and coexistence, in the sense that it will help ecosystems to adapt to external stressors. This insight is consistent with previous studies on evolutionary trapping

570   and suicide (see for example Ferriere and Legendre (2013) or Uchiumi et al. (2023)), evolutionary extinction debts (Norberg et al., 2012) and even eco-evolutionary tipping points (Dakos et al., 2019), including the original model version without pollinator evolution (Weinbach et al., 2022).

    We acknowledge that due to its simplicity, our model clearly does not capture the

575   full complexity of plant-pollinator interactions. This is in particular true for the linear functional response that we chose to represent the mutualistic interaction. However, the goal of this study was not to model a specific mutualistic system with all its typical characteristics. Instead, we focus on the impact of the eco-evolutionary feedback loop on the evolution of the investment into mutualism and

580   on pollinator persistence. We argue that whether or not this feedback loop is self-reinforcing or self-dampening depends on whether or not the biotic interaction is mutualistic or antagonistic, as well as on the strength of this interaction, but not on the exact shape of the underlying functional response. This is in line with a recent

review of models for pairwise mutualistic interactions, which showed that
population dynamics of mutualisms are qualitatively robust across derivations,
including levels of detail, types of benefit, and inspiring systems (Hale &
Valdovinos, 2021).

Because our model is quite abstract and general, it could, at least in principle,
describe any type of mutualistic interaction experiencing one-sided population
decline. However, we deliberately chose to study our model through the lens of
plant-pollinator systems, given that our results are of direct relevance for
understanding and predicting the consequences of global pollinator decline (Potts
et al., 2010). Our results offer two contrasting hypotheses for eco-evolutionary
responses of plant-pollinator interaction networks to insect decline. Both
hypotheses are based on the assumption that our model system represents a
small network module embedded into a larger network but differ in the
interpretation of plant and pollinator growth rates. If these growth rates are
interpreted as energy gain from interaction with alternative partners, that is with
the extended network in which the plant-pollinator module is embedded, then
interaction disinvestment directly translates into evolution towards generalism or
towards specialisation on alternative interaction partners. In consequence, we
might expect to see network rewiring, but not necessarily network collapse. On the
other hand, disinvestment can also be interpreted as evolution towards self-
reproduction, in particular on the plant side, with potentially detrimental long-term
consequences for the whole system due to reduced gene flow and inbreeding
depression (Lepers et al., 2014).

The goal of this study was to see if the reinforcing ecological feedback loop
observed between two mutualistic species still occurs when coevolution of both
mutualistic partners is considered. We saw here that evolution, via the joint
disinvestment of both partners, speeds up the decline of the already threatened
population. Species will evolve away from interacting with bad quality interactors in
the presence of energetically more efficient alternatives (Werner et al., 2018),
given that mutualism is always a service at a cost of other reproduction and
survival means (Bronstein, 2001). Our model is useful in identifying interactions
that might be too costly to maintain in future scenarios. Specialist species are
already known to be more prone to extinction (Biesmeijer et al., 2006), but
according to our model this is also the case for species who provide a bad

mutualistic service (e.g. not efficient pollinators) and suffer already from strong internal competition (e.g. competition for nesting sites in very artificial areas or for partners in very fragmented ones). Thus, if restoration attempts are made to rescue endangered mutualistic populations, the most fragile ones would be those populations with inefficient pollinators and/or high intrinsic competition.

However, because evolution reinforces any tendency, it could also accelerate any recovery that could be implemented with restoration attempts, which is motivating for conservation actions. According to our model, restoration should focus not only on the endangered species (e.g. by providing more food, nests and potential reproduction partners) but also on the species interactors. Bettering the plant's living conditions could also decrease its internal competition and improve its interaction gains, reducing the speed of disinvestment into the interaction or even reversing the evolutionary disinvestment once the energetic balance is again in favour of mutualism.

`

Statements and Declarations

Funding

The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

640 **Competing Interests**

The authors have no relevant financial or non-financial interests to disclose.

Ethics approval

Not applicable.

645 **Consent to participate**

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

650 Not applicable.

Code availability

The simulation code will be made available as supplementary material.

Author Contributions

KTA and FW designed the study, with minor contributions from AW and CZ. FW
655 carried out the numerical simulations and analysed the data. KTA took the lead in supervision and project organisation. All authors discussed the simulation results. FW and KTA wrote the first manuscript draft, with input from AW. All authors contributed critically to revisions. All authors read and approved the final manuscript.

References

Bell, G. (2013). Evolutionary rescue and the limits of adaptation. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *368*(1610), 20120080. https://doi.org/10.1098/rstb.2012.0080

Bell, G. (2017). Evolutionary Rescue. *Annual Review of Ecology, Evolution, and Systematics*, *48*(1), 605–627. https://doi.org/10.1146/annurev-ecolsys-110316-023011

Biesmeijer, J. C [J. C.], Roberts, S. P. M., Reemer, M., Ohlemüller, R., Edwards, M., Peeters, T., Schaffers, A. P., Potts, S. G [S. G.], Kleukers, R., Thomas, C. D., Settele, J., & Kunin, W. E [W. E.] (2006). Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science (New York, N.Y.)*, *313*(5785), 351–354. https://doi.org/10.1126/science.1127863

Brännström, Å., Johansson, J., & Festenberg, N. von (2013). The Hitchhiker's Guide to Adaptive Dynamics. *Games*, *4*(3), 304–328. https://doi.org/10.3390/g4030304

Brännström, Å., Loeuille, N [Nicolas], Loreau, M., & Dieckmann, U. (2011). Emergence and maintenance of biodiversity in an evolutionary food-web model. *Theoretical Ecology*, *4*(4), 467–478. https://doi.org/10.1007/s12080-010-0089-6

Bronstein, J. L. (1994). Our Current Understanding of Mutualism. *The Quarterly Review of Biology*, *69*(1), 31–51. https://doi.org/10.1086/418432

Bronstein, J. L. (2001). The Costs of Mutualism. *American Zoologist*, *41*(4), 825–839. https://doi.org/10.1093/icb/41.4.825

Carlson, C., Akçay, E., & Morsky, B. (2022). *The evolution of partner specificity in mutualisms* (Vol. 29). https://doi.org/10.1101/2021.12.10.472103

Carlson, S. M., Cunningham, C. J., & Westley, P. A. H. (2014). Evolutionary rescue in a changing world. *Trends in Ecology & Evolution*, *29*(9), 521–530. https://doi.org/10.1016/j.tree.2014.06.005

Cheptou, P.-O [P-O], Carrue, O., Rouifed, S., & Cantarel, A. (2008). Rapid evolution of seed dispersal in an urban environment in the weed Crepis sancta. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(10), 3796–3799. https://doi.org/10.1073/pnas.0708446105

Dakos, V., Matthews, B., Hendry, A. P [Andrew P.], Levine, J., Loeuille, N [Nicolas], Norberg, J., Nosil, P., Scheffer, M., & Meester, L. de (2019). Ecosystem tipping points in an evolving world. *Nature Ecology & Evolution*, *3*(3), 355–362. https://doi.org/10.1038/s41559-019-0797-2

Dieckmann, U., & Law, R. (1996). The dynamical theory of coevolution: A derivation from stochastic ecological processes. *Journal of Mathematical Biology.* Advance online publication. https://doi.org/10.1007/BF02409751

Fabienne Harris, L., & Johnson, S. D. (2004). The consequences of habitat fragmentation for plant–pollinator mutualisms. *International Journal of Tropical Insect Science*, *24*(01), none. https://doi.org/10.1079/IJT20049

705 Ferriere, R., & Legendre, S. (2013). Eco-evolutionary feedbacks, adaptive dynamics and evolutionary rescue theory. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *368*(1610), 20120081. https://doi.org/10.1098/rstb.2012.0081

Fox, R. (2013). The decline of moths in Great Britain: a review of possible causes. *Insect Conservation and Diversity*, *6*(1), 5–19.
710 https://doi.org/10.1111/j.1752-4598.2012.00186.x

Geritz, S., Kisdi, É [É.], Meszéna, G., & Metz, J. (1996). Evolutionarily singular strategies and the adaptive growth and branching of the evolutionary tree. *Evolutionary Ecology*, *12*(1), 35–57. https://doi.org/10.1023/A:1006554906681

715 Gervasi, D. D. L., & Schiestl, F. P. (2017). Real-time divergent evolution in plants driven by pollinators. *Nature Communications*, *8*, 14691. https://doi.org/10.1038/ncomms14691

Gomulkiewicz, R., & Holt, R. D. (1995). When does Evolution by Natural Selection Prevent Extinction? *Evolution*, *49*(1), 201. https://doi.org/10.2307/2410305

720 Gonzalez, A., Ronce, O., Ferriere, R., & Hochberg, M. E. (2013). Evolutionary rescue: An emerging focus at the intersection between ecology and evolution. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *368*(1610), 20120404. https://doi.org/10.1098/rstb.2012.0404

725 Govaert, L., Fronhofer, E. A., Lion, S., Eizaguirre, C., Bonte, D., Egas, M., Hendry, A. P [Andrew P.], Brito Martins, A. de, Melián, C. J., Raeymaekers, J. A. M., Ratikainen, I. I., Saether, B.-E., Schweitzer, J. A., & Matthews, B. (2019). Eco-evolutionary feedbacks—Theoretical models and perspectives. *Functional Ecology*, *33*(1), 13–30.
730 https://doi.org/10.1111/1365-2435.13241

Hairston, N. G., Ellner, S. P., Geber, M. A., Yoshida, T., & Fox, J. A. (2005). Rapid evolution and the convergence of ecological and evolutionary time. *Ecology Letters*, *8*(10), 1114–1127. https://doi.org/10.1111/j.1461-0248.2005.00812.x

735 Hale, K. R. S., & Valdovinos, F. S. (2021). Ecological theory of mutualism: Robust patterns of stability and thresholds in two-species population models. *Ecology and Evolution*, *11*(24), 17651–17671. https://doi.org/10.1002/ece3.8453

Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H.,
740 Stenmans, W., Müller, A., Sumser, H., Hörren, T., Goulson, D., & Kroon, H. de (2017). More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PloS One*, *12*(10), e0185809. https://doi.org/10.1371/journal.pone.0185809

Hoffmann, M., Hilton-Taylor, C., Angulo, A., Böhm, M., Brooks, T. M.,
745 Butchart, S. H. M., Carpenter, K. E., Chanson, J., Collen, B., Cox, N. A., Darwall, W. R. T., Dulvy, N. K., Harrison, L. R., Katariya, V., Pollock, C. M., Quader, S., Richman, N. I., Rodrigues, A. S. L., Tognelli, M. F., . . . Stuart, S. N. (2010). The impact of conservation on the status of the world's vertebrates. *Science (New York, N.Y.)*, *330*(6010), 1503–1509.
750 https://doi.org/10.1126/science.1194442

Kisdi, É [Éva] (2006). Trade-off geometries and the adaptive of two co-evolving species. *Evolutionary Ecology Research*(8), 959–973.

Koch, H., Frickel, J., Valiadi, M., & Becks, L. (2014). Why rapid, adaptive evolution matters for community dynamics. *Frontiers in Ecology and Evolution*, *2*. https://doi.org/10.3389/fevo.2014.00017

Le Conte, Y., Ellis, M., & Ritter, W. (2010). Varroa mites and honey bee health: can Varroa explain part of the colony losses? *Apidologie*, *41*(3), 353–363. https://doi.org/10.1051/apido/2010017

Lepers, C., Dufay, M., & Billiard, S. (2014). How does pollination mutualism affect the evolution of prior self-fertilization? A model. *Evolution; International Journal of Organic Evolution*, *68*(12), 3581–3598. https://doi.org/10.1111/evo.12533

Loeuille, N [N.] (2019). Eco-evolutionary dynamics in a disturbed world: implications for the maintenance of ecological networks, [version 1; peer review: 2 approved]. F1000Research 2019, 8(F1000 Faculty Rev):97.

Loeuille, N [Nicolas], & Loreau, M. (2005). Evolutionary emergence of size-structured food webs. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(16), 5761–5766. https://doi.org/10.1073/pnas.0408424102

Madjidian, J. A., Morales, C. L., & Smith, H. G. (2008). Displacement of a native by an alien bumblebee: Lower pollinator efficiency overcome by overwhelmingly higher visitation frequency. *Oecologia*, *156*(4), 835–845. https://doi.org/10.1007/s00442-008-1039-5

Norberg, J., Urban, M. C., Vellend, M., Klausmeier, C. A., & Loeuille, N [Nicolas] (2012). Eco-evolutionary responses of biodiversity to climate change. *Nature Climate Change*, *2*(10), 747–751. https://doi.org/10.1038/nclimate1588

Parvinen, K. (2005). Evolutionary suicide. *Acta Biotheoretica*, *53*(3), 241–264. https://doi.org/10.1007/s10441-005-2531-5

Pelletier, F., Garant, D., & Hendry, A. P [A. P.] (2009). Eco-evolutionary dynamics. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *364*(1523), 1483–1489. https://doi.org/10.1098/rstb.2009.0027

Post, D. M., & Palkovacs, E. P. (2009). Eco-evolutionary feedbacks in community and ecosystem ecology: Interactions between the ecological theatre and the evolutionary play. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *364*(1523), 1629–1640. https://doi.org/10.1098/rstb.2009.0012

Potts, S. G [Simon G.], Biesmeijer, J. C [Jacobus C.], Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E [William E.] (2010). Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology & Evolution*, *25*(6), 345–353. https://doi.org/10.1016/j.tree.2010.01.007

Reid, A. J., Carlson, A. K., Creed, I. F., Eliason, E. J., Gell, P. A., Johnson, P. T. J., Kidd, K. A., MacCormack, T. J., Olden, J. D., Ormerod, S. J., Smol, J. P., Taylor, W. W., Tockner, K., Vermaire, J. C., Dudgeon, D., & Cooke, S. J. (2019). Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biological Reviews of the Cambridge Philosophical Society*, *94*(3), 849–873. https://doi.org/10.1111/brv.12480

800   Roels, S. A. B., & Kelly, J. K. (2011). Rapid evolution caused by pollinator loss in Mimulus guttatus. *Evolution; International Journal of Organic Evolution*, *65*(9), 2541–2552. https://doi.org/10.1111/j.1558-5646.2011.01326.x

Sachs, J. L., & Simms, E. L. (2006). Pathways to mutualism breakdown. *Trends in Ecology & Evolution*, *21*(10), 585–592.
805   https://doi.org/10.1016/j.tree.2006.06.018

Thomann, M., Imbert, E., Devaux, C., & Cheptou, P.-O [Pierre-Olivier] (2013). Flowering plants under global pollinator decline. *Trends in Plant Science*, *18*(7), 353–359. https://doi.org/10.1016/j.tplants.2013.04.002

Thompson, J. N. (1998). Rapid evolution as an ecological process. *Trends in
810   Ecology & Evolution*, *13*(8), 329–332. https://doi.org/10.1016/s0169-5347(98)01378-0

Toby Kiers, E., Palmer, T. M., Ives, A. R., Bruno, J. F., & Bronstein, J. L. (2010). Mutualisms in a changing world: An evolutionary perspective. *Ecology Letters*, *13*(12), 1459–1474. https://doi.org/10.1111/j.1461-
815   0248.2010.01538.x

Uchiumi, Y., Sato, M., & Sasaki, A. (2023). Evolutionary double suicide in symbiotic systems. *Ecology Letters*, *26*(1), 87–98. https://doi.org/10.1111/ele.14136

Wagner, D. L., Grames, E. M., Forister, M. L., Berenbaum, M. R., & Stopak, D.
820   (2021). Insect decline in the Anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(2). https://doi.org/10.1073/pnas.2023989118

Weinbach, A., Loeuille, N [Nicolas], & Rohr, R. P. (2022). Eco-evolutionary dynamics further weakens mutualistic interaction and coexistence under
825   population decline. *Evolutionary Ecology*, *10*, 351. https://doi.org/10.1007/s10682-022-10176-7

Werner, G. D. A., Cornelissen, J. H. C., Cornwell, W. K., Soudzilovskaia, N. A., Kattge, J., West, S. A., & Kiers, E. T. (2018). Symbiont switching and alternative resource acquisition strategies drive mutualism breakdown.
830   *Proceedings of the National Academy of Sciences of the United States of America*, *115*(20), 5229–5234. https://doi.org/10.1073/pnas.1721629115

Willmer, P. (2011). *Pollination and Floral Ecology*. Princeton University Press. https://doi.org/10.1515/9781400838943

Yacine, Y., Allhoff, K. T., Weinbach, A., & Loeuille, N [Nicolas] (2021). Collapse
835   and rescue of evolutionary food webs under global warming. *The Journal of Animal Ecology*, *90*(3), 710–722. https://doi.org/10.1111/1365-2656.13405

Yamamichi, M., & Miner, B. E. (2015). Indirect evolutionary rescue: Prey adapts, predator avoids extinction. *Evolutionary Applications*, *8*(8), 787–795. https://doi.org/10.1111/eva.12295

840