Neurovascular Coupling is Required for Amygdala Neuronal Activation and Associated Emotional Behaviors in Mice, Depending on NMDA Receptor Subunit 2D-expressing Cerebral Arteriolar Smooth Muscle Cells

Running title: Neurovascular Coupling Regulates Amygdala-dependent Emotion Homeostasis

Jiayu Ruan[1,2,3], Yiyi Zhang[2,3], Dongdong Zhang[2,3], Huiqi Xie[2,3], Xinghua Quan[2,3], Wentao Wang[2,3], Xu Hu[2,3], Yanmei Tao[5,6], Bingrui Zhao[2,3], Danping Lu[2,3], Jie-Min Jia[2,3,4*]

Affiliations

1. College of Life Sciences, Zhejiang University, Hangzhou, Zhejiang 310058, China.

2. Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou 310024, China.

3. Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou 310024, China.

4. Laboratory of Neurovascular Biology, Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou 310024, China.

5. Key Lab of Organ Development and Regeneration of Zhejiang Province, Institute of Life Sciences, College of Life and Environmental Sciences, Hangzhou Normal University, Hangzhou 311121, China.

6. Department of Physiology, Medical College, Southeast University, Nanjing 210009, China.

*Correspondence: Jie-Min Jia (jiajiemin@westlake.edu.cn)

Abstract

Proper brain function depends on intact neurovascular coupling (NVC). Through NVC, local cerebral blood flow (CBF) is rapidly increased in activated cerebral regions to meet the corresponding demands of neural activities. NVC has been believed to support advanced cerebral functions such as complex emotional homeostasis. However, few studies provide causal link evidence with specific molecular and cellular identities. Here, using RNAscope, cell-type specific conditional knockout (cKO) perturbation in BL6/C57 mice, laser speckle contrast imaging, and mouse behavior assays, we demonstrated that the N-methyl-D-aspartic acid (NMDA) receptor noncore subunit NR2D is only expressed in cerebral arteriole smooth muscle cells (aSMCs) but not in peripheral aSMCs. Both aSMC-NR2D cKO and aSMC-NR2D+NMDA receptor subunit 1 (NR1) double cKO mice displayed a severe decline in functional hyperemia induced by whisker stimulation. In addition, the NR2D pathway was independent of previously demonstrated COX-2-PGE2-dependent mechanism and high-fat diet (HFD)-induced impairments in NVC. Caldesmon is a potential downstream protein in the NR2D pathway for NVC regulation. Notably, we demonstrated that NVC deficits caused abnormalities in complex emotional behaviors, such as anxiety-like behaviors (exhibiting increased defecation and urine), reduced curiosity and decreased fear instinct (displaying less fear of aggressive CD1 mice). Furthermore, fear-response gene c-FOS expression in the amygdala neurons was attenuated in aSMC-NR2D knockout mice after response to acute stresses. In contrast, its expression level remained unchanged in the non-fear-related cerebral regions such as the piriform cortex and hippocampus. These findings demonstrated that inadequate functional hyperemia impaired cerebral complex emotional functions and suggested a co-existence of anxiety and fearlessness in single individuals.

Introduction

Cerebral blood flow (CBF) provides logistic support for maintaining the homeostasis of the cerebral microenvironment by delivering oxygen and nutrients and clearing carbon dioxide and metabolic wastes as promptly and adequately as possible (1). CBF impacts the cerebral microenvironment by priming extracellular ion homeostasis, which could influence neuronal excitability (2). The emerging views are that moderate CBF dysregulation, not to such an extent as ischemia, leads to chronic cerebral dysfunction and disease (3). Cerebrovascular dysregulation becomes a feature not only of cerebrovascular pathologies but also of complex cognitive (4) and emotional disorders (5). Regionally rapid functional hyperemia induced by local neural activities, termed neurovascular coupling (NVC), is a functional interplay between the central nervous system

(CNS) and cerebrovascular systems by signaling neural messages, resulting in cerebral arterial smooth muscle cells (aSMCs) relaxation and hence vasodilation, leading the CBF increasing in regions of neural activating (6). NVC is a universally conserved biological process among mammalians that can effectively and spatiotemporally regulate cerebral blood supply distribution (7). The dynamics of CBF are widely assumed to represent neuronal activity in functional neuroimaging, which is the only non-invasive approach to measuring the cerebral cognitive and emotional status of humans. One of the known applications of NVC is functional magnetic resonance imaging (fMRI) (8).

With fMRI, high levels of the blood oxygenation level-dependent (BOLD) signal mirrored with an increased NVC are often repeatedly observed in some classic cerebral regions of patients with emotional disorders such as major depressive disorder (MDD), general anxiety disorder (GAD), and fear-related post-traumatic stress disorder (PTSD). For instance, MDD patients commonly manifest robust and persistent high BOLD signals in the lateral habenula (LHb), where excessive neural excitements were verified in many rodent animal models of depression (9). Besides, almost all PTSD patients and fear-stressed humans exhibit high BOLD signals in the amygdala (10, 11). Notably, induced by acute fear stress (e.g., instinct fear of a predator) or emotional disorders, the correlation between physiological or pathological neural excitement and an increased BOLD signal in the amygdala is an evolutionally conserved observation, crossing humans (12, 13), nonhuman primates (14), and rodents (15). However, the physiological roles of NVC in complex high-level cerebral functions (namely, how vital the increased amount of CBF beyond the resting CBF level is for advanced cerebral functions) remain largely unknown despite decades of extensive investigations on the underlying mechanisms of NVC.

Specifically, in emotional disorder studies, little insight has been drawn into how the increased magnitude of CBF over the resting level is involved in neuronal excitement in the amygdala and fear perception. On the other dimension, recent findings point out that in the *in vivo* experiments of rats, fluoxetine (a selective serotonin reuptake inhibitor) for rats has been proven to cause cerebral arterioles vasodilation (16) and reverse the deficit in escape latency in the learned helplessness model of the emotional disorder (17). It is well established that except neurons, aSMCs express a high level of serotonin transporter (18) and receptor subtypes (19, 20). These two facts together implicate that blood flow alteration caused by aSMCs may be involved in reversing emotional status. The consequence of ignoring the contributions of NVC to emotional disorders leads to stagnation in studies.

Compared with the study of the correlation between NVC and emotional disorders, the assessment of NVC is

more mature. The extent of NVC can be quantified by measuring CBF variation and vascular diameters during

the functional hyperemia in the barrel cortex of mice induced by the whisker stimulation, using laser doppler

imaging (21), laser speckle imaging (22), and two-photon Imaging (23). Previous studies have revealed

multiple signalling pathways mediating NVC: neurons or endothelial cells releasing nitric oxide (NO) to cerebral

arterioles SMCs (aSMCs) cause vasodilation (24-26); astrocyte-secreted small vasoactive molecules cause

the constriction of aSMCs (27-29). Although extensive manipulations have been established to enhance or

inhibit NVC, most depend on pharmacology or null mutations, which lack cellular specificity. Therefore, no

clear conclusion can be yielded by these means because both the neural circuit *per se* and the cerebrovascular

system are being affected, which confounds the contributions from vascular cell-dependent NVC inhibition.

Only the conditional knock-out genetic perturbation strategy developed during the recent decade can achieve

the goal of cell specificity (23, 30, 31). Most recent findings indicate that the N-methyl-D-aspartic acid (NMDA)

receptor expressed in aSMCs controls NVC robustly, revealed by cell-type specific genetic perturbation.

However, which noncore subunit of the NMDA receptor works with the obligatory NR1 core subunit in aSMCs

to mediate NVC remained unknown. The NMDA receptor has long been involved in emotional disorders. For

instance, the extinction of conditioned fear is mediated by NMDA receptors in the amygdala (32, 33); in

emotional disorders patients, it can be found that NMDA receptors decreased in the prefrontal cortex (34);

NR2D (one of a noncore subunit of NMDA receptor) total knock-out mice exhibited anxiety-like and depression

behavior phenotypes (35). Since the total knock-out strategy and widespread nature of drug interruption lose

the cellular specificity manipulation, the contribution and cellular identity of NMDA receptors in fear-like and

anxiety-like behavior presentation are unclear.

Here, we first reported that NR2D is a unique noncore subunit of the NMDA receptor expressed in cerebral

aSMCs. In whisker stimulation-mediated functional hyperemia, conditional knockout (cKO) of NR2D in aSMCs

of mice resulted in reduced functional hyperemia. The NVC regulatory pathway involving NR2D was similar to

that mediated by NR1 but independent of the COX-2 and high-fat diet (HFD) pathways. Caldesmon is a

potential downstream protein in the NR2D pathway for NVC regulation. Notably, behaviors that represent fear

status were significantly diminished in aSMC-specific NR2D cKO mice. Furthermore, neuronal c-Fos

expressions in the amygdala, which usually represents neurons activated in response to fear stimuli, were

robustly attenuated in aSMC-specific NR2D cKO mice. More interestingly, aSMC-specific NR2D cKO mice

exhibited a higher level of anxiety-like behavior by producing more defecation, similar to that in anxiety patients (36). Based on the above evidence, being less fearful and having a higher anxiety level can coexist in the same individuals. Our findings may shed light onto a new angle of emotional disorder treatment development.

Materials and methods

Animal and animal care

All mice procedures complied with the Institutional Animal Care and Use Committee (IACUC) guidelines of the School of Life Sciences, Westlake University. All mice housed in Westlake University Laboratory Animal Resource Center with standard animal rooms provided a 12-h light/dark cycle at 25°C. No more than six mice were fed in each cage, and standard chow and water were provided to the mice *ad libitum*. All the genetic information of mice is in Table 1. All mice were B6 background. Unless otherwise specified, the age of all male adult mice was from 2-month-old to 3-month-old.

Cell culture

P0 mice anesthetized with ice. Surgical tweezers were used to quickly extract pial membrane or cortex from the brains of mice in DMEM (Gibico, Cat# 11965092, USA). Then, 0.25% Trypsin (Thermo Scientific, Cat# 25200072, USA) was used to digest pia or cortex for 15min at 37℃ in the incubator. After that, cells were added with FBS (Gibico, Cat# 10099, USA) and filtered by a 40μm filter (Falcon, Cat# 352340, CHN). Finally, cells were centrifuged 1,600*g at 4℃ for 5min and resuspended by SMCM (ScienCell, Cat# 1101, USA) in T25 flasks (Corning, Cat# 430639, USA).

TMX administration

1. Tamoxifen for mice

Tamoxifen (MCE, Cat# HY-13757A, USA) was orally administrated to mice. Tamoxifen was dissolved in corn oil (Aladdin, Cat# C116023-500ml, CHN), and the dose for each mouse per day was 0.2ml (10mg/ml). In almost all cases, for adult mice, tamoxifen was orally administered to mice from postnatal day 60 to postnatal day 63 (P60 to P63). Otherwise, for the P5 conditional knock-out group and its littermate control group, infant mice got tamoxifen through breast milk. On the fifth day after the birth of mice (P5), the mother of infant mice was orally administrated with tamoxifen for four days (from P5 to P8). In particular cases, tamoxifen administration time refers to the results and figures of the corresponding experiments.

2.   Tamoxifen for cell culture (include cell sorting)

When cells reached approximately 80% full of the bottom surface of T25 flasks, tamoxifen was administrated to cells. tamoxifen was dissolved in DMSO (Solarbio, Cat# D8371, CHN), and the final concentration of tamoxifen for SMCM was 5µM. After four-day culture, EGFP or Tdtomato in cells were expressed. Cells were sorted by MA-900 (Sony, FCCF-MBR, JPN) to obtain EGFP-positive or Tdtomato-positive cells, and these cells were pure sorted cerebral arterioles smooth muscle cells (aSMCs). Sorted cells were cultured in SMCM for further assay. The purity of cells sorted by the same method had been verified in the paper published by our laboratory (37).

RNA scope *in situ* hybridization

Horizontal sections (15µm) of tamoxifen administrated SMAcre^{ER+}:Ai14 mice brains were obtained by CM1950 (Lecia, Histo-CM1950, DEU). All RNA scope *in situ* hybridisation productions were obtained from ACDbio, including RNAcope Multiplex Fluorescent Reagent Kit v2 (ACDbio, Cat# 323100, USA), *Grin2d* probe (ACDbio, Cat# 425951-C3, USA) and *Grin1* probe (ACDbio, Cat# 431611, USA). All operations were carried out following the reagent instructions of ACDbio company and a published study that used RNA scope *in situ* hybridization (38).

RT-PCR

aSMCs from triple transgenic mouse lines, SMAcre^{ER+}:*grin2d*$^{f/f}$:Ai47 (conditional knock-out group) and SMAcre^{ER+}:*grin2d*$^{+/+}$:Ai47 (littermate control group), were used as a source for RNA extraction. Briefly, after obtaining pure sorted EGFP/Tdtomato-positive aSMCs from pia (pial aSMCs) and cortex (parenchymal aSMCs) of the conditional knock-out group and its littermate control group, cells were cultured in 6cm dishes (Corning, Cat# 430166, USA) with SMCM. After cells reached around 90% density, Trizol (Sangon Biotec, Cat# B511311-0100, CHN) was used for the RNA extraction. RNA from P0 SMAcre^{ER+}:*grin2d*$^{+/+}$:Ai47 cortex/cerebellum was used as a positive control. After the P0 mice were anesthetized with ice, the cerebellum of the mice was directly taken out, and TRIzol was used for RNA extraction.

After RNA extraction, all RNA samples were reverse transcription to cDNA (50ng/µl) using HiScript III 1st Strand cDNA Synthesis Kit (Vazyme, Cat# R312-02, CHN). For the PCR amplification system, each reaction contained 10µl of Green Taq Mix (Vazyme, Cat# P131-AA, CHN), 2µl of cDNA, 2µl of primer pairs (10µM), and

6μl of RNAase-free water. Except the amplification cycle of *18s* was 20, the amplification cycle of other primers

was 35. Agarose gel electrophoresis used 2% agarose gel (BIOWESTE, Cat# 111860, ESP) and ran at 220V

for 30min. RT-PCR gel images were obtained by Tanon imager (Tanon, 1600 Geil imager, CHN). All operations

were carried out in strict accordance with the reagent instructions. A list of primer pairs is in Table 2.

Western-bolt

For tissue sample collection, tissue samples were ground together with RIPA (Thermo Scientific, Cat# 89901,

USA) on ice for 30min. For cell samples, after cells reached around 90% density, the RIPA was used for protein

extraction on ice for 30min. Protease Inhibitor Cocktail (CWBIO, Cat# CW22008, CHN) was used to prevent

protein enzymolysis during the protein extraction. After that, samples were centrifuged 12,000*g at 4℃ for

5min, and supernatant were collected. Concentrations of proteins were measured by Micro BCA Protein Assay

Kit (CWBIO, Cat# CW2011S, CHN), and after the concentration measurement, all samples were boiled with

SDS-PAGE loading buffer (CWBIO, Cat# CW0027S, CHN) for 10min at 100℃. 8% SDS-polyacrylamide

electrophoresis gels were used for gel running, and a protein marker (GenStar, Cat# M221-05, CHN) was used

to identify the protein size. After 90min gels running at 120V (80V for the first 30min), gels were transferred to

0.45μm PVDF membranes (Millipore, Cat# IPVH00010, USA) at 400mA for 120min. After transfer, PVDF

membranes were blocked by 5% non-fat milk in 1X TBST buffer for 60min at room temperature with a mild

horizontal shake.

After the block, for each particular assay, the first antibody diluted with 5% non-fat milk in 1X TBST buffer was

used to incubate the membrane overnight at 4℃ with a mild horizontal shake. After the first antibody incubation,

the membrane was washed by 1X TBST 10min (3 replicates). The corresponding second HRP-conjugated

antibody diluted with 5% non-fat milk in 1X TBST buffer was used to incubate the membrane 45min at room

temperature with a mild horizontal shake. After the second antibody incubation, the membrane was washed

by 1X TBST 10min (3 replicates). ECL kits (Abbkine, Cat# BMU102-CN, CHN) and gel image system (GE,

GEL Imaging System AI680RGB v.2.0, USA) were used for exposure photography. All operations were carried

out in strict accordance with the reagent instructions. A list of antibodies is in Table 3.

Immunofluorescence staining

For tissue Immunofluorescence (IF) staining, briefly, after anesthesia with 1% pentobarbital sodium, the mice

were perfused with 1X PBS (5min) and 4%PFA in 1X PBS (5min) through the left ventricle to empty blood and

fix tissue. Afterwards, tissue was extracted and fixed by 4%PFA in 1X PBS overnight at 4℃ with a mild horizontal shake. The next day, for MCA branches IF staining, MCA branches were carefully extracted from the brain using surgical tweezers. Then, treated isolated MCA branches by 0.5% Triton X-100 (Diamond, Cat# A110694-0500, CHN) in 1X PBS for 30min at room temperature, and then isolated MCA branches were washed with 1X PBS for 10min (3 replicates). Next, 5% BSA (BBI, Cat# A600332-0100, CHN) in 1X PBS was used to block isolated MCA branches for 30min at room temperature. After the block, isolated MCA branches were rewashed as described before. For other tissue like brain and peripheral organs, all organs were sliced into 50μm (Leica, Histo-VT1200S, DEU) after fixed by 4%PFA in 1X PBS overnight at 4℃ with a mild horizontal shake. The subsequent process was similar to that of isolated MCA branches.

For cultured cell immunofluorescence staining. After cells reached around 90% density, cells were passed by into 12-well plates (Corning, Cat# 3515, USA) with glass slices (30k cells per well). After cells in 12-well plates reached around 70% density, cells in glass slices were fixed by 4%PFA in 1X PBS for 10min at room temperature. Washed cell slices with 1X PBS 10min (3 replicates), treated cells with 0.3% Triton X-100 in 1X PBS for 10min at room temperature, and then washed again with 1X PBS 10min (3 replicates). Next, 5% BSA in 1X PBS was used to block cell slices for 30min at room temperature. After the block, cell slices were rewashed as described above.

In the staining process, the first antibody diluted with 1% BSA+0.1% Triton X-100 in 1X PBS was used to incubate samples overnight at 4℃. After the first antibody incubation, samples were washed with 1X PBS 10min (3 replicates). The corresponding second fluorescence-conjugated antibody diluted with 1% BSA+0.1% Triton X-100 in 1X PBS was used to incubate samples 120min at room temperature. After the second antibody incubation, samples were washed with 1X PBS 10min (3 replicates). After that, 1000X diluted Hoechst (AbMole, Cat# M5112, CHN) was used for nucleus staining. The operation was slightly different for the co-staining of NR2D and NR1 in MCA. After incubating with the first antibody of NR2D, Goat Anti-Mouse IgG (H+L) was used as the second antibody for 120min incubating at room temperature. Then, Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 647, was used as the third antibody to incubate at room temperature for 120min. After incubating the third antibody, 5% BSA in 1X PBS was used to block isolated MCA branches for 30min at room temperature. Then, the first antibody of NR1 was incubated overnight at 4℃. After the first antibody of NR1 incubation, samples were incubated with Goat anti-Mouse IgG (H+L)

Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488 at room temperature for 120min. 1X PBS was used for washing after each step (10min, 3 replicates). Tissue slices and cell culture slices were covered by a glass slice with the mounting medium (SouthernBiotech, 0100-01, USA). Zeiss LSM800 (Zeiss, LSM800, DEU) microscopy was used for photography. All operations were carried out in strict accordance with the reagent instructions. A list of antibodies is in Table 3.

Whisker stimulation

The whisker was trimmed to only remain lane three and lane four on one side of the face, and the scalp was exposed on the same day. The mice did not perform any operation the day before the assay to reduce potential interference (Particular set-up diagrams refers to the results and figures of the corresponding experiments). On the day of the assay, mice were anesthesia with 1% pentobarbital sodium, and stable fixated on the platform. The electro-motor-driven brush stimulated the whisker longitudinally for 1 minute at a frequency of 4Hz. The principle of laser speckle contrast imaging has been described in the previous study (39), and the operation referred to the published study (40). Laser Speckle Image System (LSIS, RWD, RFSLI ZW/RFSLI III, CHN) was used to monitor the CBF variation in the region of interest (ROI) of the opposite barrel cortex of mice. The recording parameters were: 512*512 resolution, 10FPS, and 2min recording (the 30s before stimulation, 1min during stimulation, and 30s after stimulation).

Cisternal injection and whisker stimulation

All operations were followed by study ***Cannula Implantation into the Cisterna Magna of Rodents*** *(https://doi.org/10.3791/57378)*. NS-398 (MCE, Cat# HY-13913, USA) was mixed with FITC (10mg/ml, Sigma, FD40S-1G, USA) for injection, and the final concentration of NS-398 in the CSF was 100µM. 10min after the injection, remove the scalp, and the FITC signal was detected by microscope (Zeiss, Axio Zoom v16, DEU). 20min after the injection, whisker stimulation (parameters were similar to that described above in "Whisker stimulation" part) was performed for successfully injected mice.

Physical examination of mice and the HFD challenge experiment

Electronic scales were used for weighing mice and high-fat food (SYSEbBio, Cat# D12492, CHN). The body fat/lean/total water content of mice was measured with EchoMRI™-100H (Echo MRI, Cat# 100H, USA). The blood pressure of mice was measured by Kent Noninvasive blood pressure measurement system for animals/Biopac physiological signal analytical system (Kent/Biopac, Cat#, USA). HFD challenge kept for 8-

weeks. The food consumption and the body fat/lean/total water content of mice were monitored during the HFD challenge. Whisker stimulation was performed after the HFD challenge (parameters were similar to that described above in "Whisker stimulation" part).

Immunoprecipitation assay (include lentivirus infection)

Tdtomato-positive pure aSMCs from P0 pial culture of SMAcre[ER+]:Ai14 mice were collected for lentivirus infection and further immunoprecipitation (IP) assay. Cells were cultured in 6cm dishes with SMCM. After cells reached around 90% density, add lentivirus (Lenti-EF1-EGFP-CMV-*grin2d*-3XFlag-WPRE) into the culture and keep for one week (virus titer: $2.6*10^9$). Infected cells were green. MA-900 sorted out EGFP and Tdtomato double-positive cells, and these cells were lentivirus-infected aSMCs. These cells were then cultured in 10cm dishes. After cells reached around 90% density, cells were collected by RIPA reagent. A Protein G Magnetic beads kit (Epizyme, Cat# YJ002, CHN) was used for the immunoprecipitation assay. NR2D antibody and IgG control antibody were mixed with Protein G Magnetic beads respectively to pull down NR2D and its related protein. Pull-down products were verified by western-bolt and analyzed through mass spectrometry. All operations were carried out in strict accordance with the reagent instructions. Data analysis and inclusion criteria methods are described in "Statistical analysis software and code" part. A list of antibodies is in Table3.

Open field test and open field-like test

The setup of open field test (OFT) referred to published studies (41-44). Each mouse was placed in the central area of a 40*40 cm box for the OFT. The camera recorded their path, fecal/urine production and movement parameters within 15 minutes. The setup of open field-like test (OFLT) referred to published studies (45). Each mouse was placed in the central area of a 23*23 cm box for the OFLT. An iron mesh wall replaced one wall of the box. Sketch paper was put on the bottom of the box to collect urine. A transparent acrylic plate covered the top of the box. Symmetric space is on the opposite side of the iron mesh wall. The camera recorded their path, fecal/urine production and movement parameters within 20 minutes. During the analysis, the box was divided into three parts (left area, central area, and right area) to analyze the preference of mice for iron mesh wall.

CD-1 fear assay

The setup of CD-1 fear assay referred to published papers (45-48). CD-1 mice that were never exposed to wild-type (WT, SMAcre[ER-]:*Grin2d*[f/f]) and cKO (SMAcre[ER+]:*Grin2d*[f/f]) mice were used for the assay. Before the assay, simultaneously put a single WT or cKO mouse and a CD-1 mouse into a cage. The cage contains new

padding and a metal lid on the top. One side of the metal lid is used for placing food and water bottles, so the height is lower than another side. After that, the camera was used to record all behaviors of C57 mice (WT or cKO mouse) and CD-1 mice, including but not limited to fighting behaviors within 10 minutes. After the assay, CD-1 and C57 mice were separated again. C57 mice were fed alone in a clean new cage for 90 minutes and then used for brain extraction to detect neural c-Fos expression in the amygdala, S1BF, Pir cortex, and hippocampus.

Elevated plus-maze test

The setup of elevated plus-maze (EPM) referred to published studies (38, 49). The EPM comprises two open arms (65×5 cm) and two close arms (65×5 cm). Each mouse was placed in the center, facing one of the two open arms, and allowed to explore for 5 minutes. Time spent in open arms, center of arms, and close arms were recorded.

Statistical analysis software and code

All the analysis of fluorescence intensity and grey value used Zeiss Zen Lite (v2.3) and ImageJ (v1.53u). All whisker stimulation original data were measured by LSIS-self software (v5.0). Visualization of CBF variation, CBF plateau mean value, CBF plateau maximum value, CBF "Time to plateau", CBF increase value (K-slope), data inclusion criteria before bioinformatics analysis of mass spectrometry data, and OFLT heat map were calculated using self-compiled MATLAB codes. Besides, The Veen map was made by Hello jvenn! (http://jvenn.toulouse.inra.fr/app/example.html). KEGG and GO analysis referred to DAVID Bioinformatics Resources (https://david.ncifcrf.gov/home.jsp) and Kyoto Encyclopedia of Genes and Genomes (https://www.genome.jp/kegg/). The figure of KEGG pathway map made by self-compiled R code. All codes were available at https://github.com/JialabEleven/Neurovascular_coupling. All the original behavioral data of OFLT were recorded using any-maze (v4.73). OFT and EPM original behavioral data were recorded using Noldus. All statistical data were processed by Numbers (v12.2), and all figures were made by Prism (v9.4.1).

Brief description of codes

1. **Data and code availability.** All image processing, analysis, and visualization, as detailed below, were performed in ImageJ (v1.53u), R studio and MATLAB (MathWorks). All quantitative analyses were done in a blinded manner. The codes were divided into four parts, visualization of cerebral blood flow variation, analysis of cerebral blood variation periods, data pre-analysis of mass spectrum, and the heat map of mice

tracking of OFLT. All codes can be found at https://github.com/JialabEleven/Neurovascular_coupling.

2. Visualization of the CBF change collected by LSIS. Using custom-written scripts, the LSIS data were

analyzed in MATLAB (version R2021a; MathWorks). It generates relative perfusion values (arbitrary units),

and the CBF was expressed as a percentage change over baseline values in whisker stimulation

experiments. For the whisker stimulation experiments, the imaging sampling frequency is 10Hz for a total

of 120 seconds, and the first frame per second of the image (512 pixels × 512 pixels) was taken as the

analysis object. A 30-second-long period (typically 30 images) was averaged and considered as the

baseline image, and which grey value of every pixel was recorded at the beginning of the imaging session.

Then every whisker stimulation image (60 seconds) and after-stimulation image (30 seconds) were

normalized to the baseline image. To demonstrate the differences in CBF variation between

SMAcre^{ER-}:$Grin2d^{f/f}$ mice and SMAcre^{ER+}:$Grin2d^{f/f}$ mice, the ROI of barrel cortex opposite to the whisker

stimulation side was chosen, and the diagram of CBF change over baseline of the ROI was presented in

Figure 2B.

3. Analysis of CBF periods. Acquired laser speckle data of relative perfusion values were processed in

MATLAB using custom-written scripts for the automated detection and analysis of perfusion change

parameters. To find the endpoint of the rising period of CBF value under whisker stimulation, the local

maximum intensity threshold was determined visually for each image and the cubic interpolation calculated

the time to 20% and 80% of the local maximum. Within the threshold range, the inflection point where the

value rises rapidly to a stable level was selected as the end point to reach the blood flow plateau. The

beginning of the whisker stimulation (30 seconds) to the time of the selected local maximum was the rising

period, and the local maximum to the end of the whisker stimulation (90 seconds) was the plateau period.

Their kinetics of them were characterized as rise20-80 time in the rising period (from 20% to 80% of local

maximum amplitude), K-slope (slope of rising speed) in the rising period, and maximum and mean value

in the plateau period, as shown in Figure 2C-F, Figure 3B/E-G, Figure 4F/H-J, Figure S3, and Figure S4.

4. Mass spectrum analysis for IP products of NR2D. Acquired mass spectrum data was analyzed in

MATLAB and R studio using a custom algorithm. Firstly, the data is filtered through the rules set by

MATLAB as below, the proteins needed to appear at least once in the three IP experiments and the

peptide-spectrum match (PSM) value was higher than ten. Furthermore, the proteins would be preserved

for subsequent analysis if the PSM value of this protein existed in the NR2D-IP group and not in the IgG

control group. If the PSM value of proteins exists in both groups, the proteins would be preserved if the

PSM value of NR2D-IP groups was greater than twice that of the corresponding IgG control group. The

selected mass spectrum data was further analyzed by R studio to achieve KEGG pathway enrichment, as

shown in Figure 5.

5. Heat map of mice tracking. Acquired Open Field-Like Test data of mice activity was analyzed using a

custom algorithm in MATLAB. To compare the location preference in an OFT between SMAcre^{ER-}:*Grin2d*$^{f/f}$

mice and SMAcre^{ER+}:*Grin2d*$^{f/f}$ mice, the track map of mice was transformed into heat map which employed

the Gaussian blur operation, as shown in Figure 6B.

Results

Expression of NR2D in cerebral aSMCs

We first confirmed the expression of NR2D/*Grin2d* (the gene name of NR2D) in cerebral aSMCs through a

series of measurement approaches. First, we used RT–PCR to detect the gene expression of all NMDA

receptors in aSMCs collected from the brain parenchyma (Figure 1A/B). The functional NMDA receptor

comprises two core subunits of NR1 and two noncore subunits, NR2/3 (50). The PCR results showed that

among the seven known NMDA receptors, only the gene of the core subunit, *Grin1*, and the gene of the

noncore subunit, *Grin2d*, were expressed in cerebral aSMCs. This result suggests that the NMDA receptor in

cerebral aSMCs may comprise NR1 (encoded by *Grin1*) and NR2D (encoded by *Grin2d*) only. In addition, we

also measured the expression of *Grin2d* in cerebral aSMCs *in situ* by RNAscope. We first tested the specificity

of the RNAscope probe (Figure S1A). According to previous reports, with increasing age, the expression of

Grin2d and its encoded protein NR2D in the cerebellum is significantly reduced, while that of *Grin1* and its

encoded protein NR1 increased (51, 52). We used RNAscope probes of *Grin1* and *Grin2d* to measure the

gene expression of *Grin1* and *Grin2d* in the cerebellum of young (postnatal day 14, P14) and adult (3-month-

old) mice. In the cerebellum of young mice, *Grin2d* was expressed at high levels, but *Grin1* was expressed at

low levels. In contrast, in adult mice, *Grin2d* was expressed at low levels, but *Grin1* was expressed at high

levels. This trend is consistent with previous reports. Next, we measured the gene expression of *Grin2d* in the cerebral arterioles *in situ* in brain slices from adult SMACre[ER+]:Ai14 mice (Figure 1C). One week after oral administration of tamoxifen, the aSMCs of mice specifically expressed tdTomato fluorescent protein. We used the expression of this red fluorescent protein to confirm the location of cerebral aSMCs. The results of RNAscope showed that although the density of the *Grin2d* probe signal was less than that of neurons, gene expression of *Grin2d* occurred in cerebral aSMCs. This result supported the PT-PCR data and further confirmed the expression of *Grin2d* in cerebral aSMCs.

Next, we introduced the Cre-loxp system to knockout NR2D in aSMCs. We tested whether Cre protein was expressed in neurons and endothelial cells, leading to the leakage of Cre protein to non-aSMCs in the CNS. The IF results demonstrated negligible tdTomato signal in NeuN-positive cells and CD-31 positive cells in brain slices of SMACre[ER+]:Ai14 mice (Figure S1B). Then, due to the extensive-expression of Cre protein in all aSMCs of SMACre[ER+] mice, we used SMACre[ER+]:Ai14 mice to detect the colocalization of NR2D protein and tdTomato protein in the peripheral organs of mice after the administration of tamoxifen. The immunofluorescence (IF) staining showed no colocalization of NR2D protein and tdTomato protein in the heart, liver, spleen, lung, kidney or small intestine (Figure S1C). The above evidence showed that even though SMACre[ER+] mice could express Cre protein in SMCs of other peripheral organs outside the CNS, the expression of NR2D in cerebral aSMCs was specific.

After validating the reliability of SMACre[ER+] mice, we also validated the reliability of *Grin2d*-flox mice. Nestin-Cre[+]:*Grin2d*[f/f] mice were obtained by mating Nestin-Cre[+] mice with *Grin2d*[f/f] mice. By verifying the expression of NR2D protein in the cerebellum of P0 Nestin-Cre[+]:*Grin2d*[f/f] mice through western-blot (WB), we determined that, compared with Nestin-Cre[-]:*Grin2d*[f/f] mice, the expression of NR2D in the cerebellum was significantly reduced in Nestin-Cre[+]:*Grin2d*[f/f] mice, to levels even lower than those in the cerebellum of adult Nestin-Cre[-]:*Grin2d*[f/f] mice (Figure 1D/H). Moreover, we used SMACre[ER+]:*Grin2d*[f/f] mice to verify the expression of NR2D in the whole cerebellum at P10. On the fifth day after the birth of mice, the infant mothers received oral administration of tamoxifen for four days (from P5 to P8 of infant mice). Since this strain of mice had NR2D knocked out only in aSMCs, the expression of NR2D in other non-aSMCs was not affected. In addition, because cerebral aSMCs accounted for only approximately 3‰ of the total number of cells in the brain, the loss of NR2D expression in cerebral aSMCs did not significantly impact the expression of NR2D in the whole

cerebellum. The WB results showed that there was no significant difference in the expression of NR2D in the

cerebellum between the WT(SMACre^{ER-}:*Grin2d*$^{f/f}$) mice and cKO (SMACre^{ER+}:*Grin2d*$^{f/f}$) mice (Figure 1E/I).

Based on the above results, we mated SMACre^{ER+}:*Grin2d*$^{f/f}$ mice with SMACre^{ER+}:Ai47 mice to generate triple-

positive SMACre^{ER+}:*Grin2d*$^{f/f}$:Ai47 mice to obtain pure pial aSMCs. Then, we used RT–PCR, WB, and IF to

measure *Grin2d*/NR2D expression (Figure 1F/J, Figure 1G/K, and Figure 1L/M/O). The results confirmed the

expression of *Grin2d* in cerebral aSMCs and showed the expression level of *Grin2d*/NR2D in cKO

(SMACre^{ER+}:*Grin2d*$^{f/f}$) mice was significantly lower than that in WT (SMACre^{ER-}:*Grin2d*$^{f/f}$) mice. In the *ex vivo*

experiment, we extracted the middle cerebral artery (MCA) from the cortex surface of adult mice to determine

whether there was NR2D protein expression. The expression of NR2D protein was detected in the aSMCs of

the MCA in WT mice, and the expression of NR2D in the MCA of SMACre^{ER+}:*Grin2d*$^{f/f}$ mice decreased

significantly. The above results again confirm that the expression of the *Grin2d* gene and the NR2D protein in

mouse cerebral aSMCs and the Cre-loxp system was a suitable genetic tool to significantly reduce the

expression of *Grin2d* and the encoded NR2D protein in mouse cerebral aSMCs. Finally, we verified the effect

of aSMCs-NR2D cKO on the expression of biomarkers in aSMCs by WB and IF. The results showed that the

expression of both smoothelin (Figure 1N) and SMA (Figure 1P) did not significantly change with the cKO of

NR2D in aSMCs.

Conditional knock-out of NR2D in aSMCs leads to a decrease in functional hyperemia

We tested the functional hyperemia capacity with the conventional whisker stimulation assay. We set two time

points (P5 and P60) for tamoxifen administration to knockout *Grin2d* in aSMCs in SMACre^{ER+}:*Grin2d*$^{f/f}$ mice to

determine the contribution of NR2D to functional hyperemia. Moreover, we also used platelet-derived growth

factor receptor beta (Pdgfrβ)$^{ER+}$:*Grin2d*$^{f/f}$ mice to knockout *Grin2d* at P60 to validate the contribution of NR2D

in all mural cells. All mice underwent whisker stimulation at the same age (P65) (Figure 2A). The laser speckle

image system (LSIS) was used to monitor blood volume variation during the whisker stimulation assay (Figure

2B). The results showed that compared with WT mice, all cKO mice exhibited dysfunctional hyperemia during

the whisker stimulation period, regardless of whether cKO NR2D mice were SMACre^{ER+} at P5/P60 or Pdgfrβ^{ER+}

at P60 (Figure 2C). Specifically, during the period in which whisker stimulation-induced changes in CBF had

reached a plateau, the mean CBF changes of P5 SMACre^{ER+}:*Grin2d*$^{f/f}$ mice, P60 SMACre^{ER+}:*Grin2d*$^{f/f}$ mice,

and P60 Pdgfrβ^{ER+}:*Grin2d*$^{f/f}$ mice were 4.88%, 4.75%, and 3.14%, respectively. Compared with WT mice

(14.13%), the CBF changes in cKO mice were attenuated by at least 65%, a significant difference. In addition,

there were no significant differences in the mean CBF changes among all cKO groups. The maximum CBF changes during the whisker stimulation exhibited a similar trend as the mean CBF changes in the four groups. The maximum CBF changes of WT mice, P5 SMACre^{ER+}:*Grin2d*$^{f/f}$ mice, P60 SMACre^{ER+}:*Grin2d*$^{f/f}$ mice, and P60 Pdgfrβ^{ER+}:*Grin2d*$^{f/f}$ mice were 18.38%, 8.23%, 7.34% and 5.31%, respectively. The WT group significantly differed from all three cKO groups. Moreover, the maximum CBF increase rate of P60 Pdgfrβ^{ER+}:*Grin2d*$^{f/f}$ mice was significantly lower than that of P5 and P60 SMACre^{ER+}:*Grin2d*$^{f/f}$ mice (Figure 2D). Next, we quantified the CBF change from the beginning of whisker stimulation to the time the CBF variations began to stabilize (plateau). We defined the latency between whisker stimulation and the start of the plateau period as the "time to plateau". This period reflected how long the whisker stimulation-induced CBF change took to reach a plateau. Next, we used the "K-slope" to describe the rate of slope increase of CBF during the "time to plateau", reflecting the initial rate of CBF change. Time to plateau did not significantly differ among all four groups (Figure 2E); however, due to different mean and maximum CBF changes during the plateau period, the K-slope of the WT group was significantly higher than that of all cKO groups. In addition, the K-slopes of the three cKO groups did not significantly differ (Figure 2F). In summary, cKO of NR2D in mural cells led to dysfunctional hyperemia during the whisker stimulation assay. No differences in parameters were observed between the P5 SMACre^{ER+}:*Grin2d*$^{f/f}$ mice and the P60 SMACre^{ER+}:*Grin2d*$^{f/f}$ mice. Only the maximum CBF change in P60 Pdgfrβ^{ER+}:*Grin2d*$^{f/f}$ mice was lower than that in the two SMACre^{ER+}:*Grin2d*$^{f/f}$ groups. There is no significant difference between aSMCs-NR2D cKO during development and adulthood, suggesting that the original effect of functional hyperemia is aSMCs.

As each mouse underwent at least two rounds of whisker stimulation during the whisker stimulation assay, we wondered whether high-frequency whisker stimulation led to desensitization or hypersensitization of CBF change. We used SMACre^{ER-}:*Grin2d*$^{f/f}$ mice to validate whether long-term continuous whisker stimulation led to desensitization or hypersensitization of functional hyperemia (Figure S3). We collected data on twenty-one consecutive days of whisker stimulation (ten times per day, each time lasting 1 minute). Comparison of the data obtained before and after the long-term stimulation revealed no significant difference in any parameter (mean CBF change, maximum CBF change, time to plateau, or K-slope). Therefore, the results suggest that long-term continuous whisker stimulation did not induce desensitization or hypersensitization of the CBF change.

aSMC-expressed NR2D and NR1 may work together to regulate functional hyperemia

Previous reports have suggested that the NR1 core subunits of NMDA receptors are always present in

functional NMDA receptors in neurons. To verify whether NMDA receptors containing NR2D and NR1 subunits

are present in cerebral aSMCs and whether these subunits work together, we measured the CBF change

induced by whisker stimulation in NR1 plus NR2D double-knockout mice (SMACre^{ER+}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$ mice)

before and after tamoxifen administration (Figure 3A). If the NR1 and NR2D subunits in aSMCs are similar to

those in neurons and form complexes to work together, the CBF decrease caused by a double knockout of

these two genes should be similar to that caused by a single knockout of NR2D or NR1 alone. In contrast, if

the CBF decrease differs, these two proteins work independently. First, we performed the whisker stimulation

assay before and after tamoxifen administration in WT (SMACre^{ER-}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$) mice to confirm whether

this experimental procedure plus the administration of tamoxifen affected functional hyperemia. All parameters

measured (the mean CBF change, maximum CBF change, time to plateau, and K-slope) showed no significant

differences in WT (SMACre^{ER-}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$) mice before and after tamoxifen administration (Figure S4).

This means that the experimental operation itself could not affect functional hyperemia. Then, we administered

the whisker stimulation assay to SMACre^{ER+}:*Grin2d*$^{f/f}$ mice and SMACre^{ER+}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$ mice before and

after tamoxifen administration. Before tamoxifen administration, SMACre^{ER+}:*Grin2d*$^{f/f}$ mice and

SMACre^{ER+}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$ mice showed normal functional hyperemia. However, after tamoxifen

administration, both groups of mice exhibited a decreased CBF change during the whisker stimulation plateau

period (Figure 3B). Specifically, during the plateau period, the mean CBF change (9.98% and 11.55%) and

maximum CBF change (14.07% and 16.07%) before tamoxifen administration in SMACre^{ER+}:*Grin2d*$^{f/f}$ mice

and SMACre^{ER+}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$ mice, respectively, showed no significant differences. However, after

tamoxifen administration, these two groups of mice showed significantly decreased mean values (3.42% and

3.85%, respectively) and maximum CBF change (6.01% and 7.05%, respectively) during the plateau period.

There was no significant difference between SMACre^{ER+}:*Grin2d*$^{f/f}$ mice and SMACre^{ER+}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$ mice

after tamoxifen administration (Figure 3E). Regarding time to plateau, SMACre^{ER+}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$ mice had a

longer time to plateau after tamoxifen administration than SMACre^{ER+}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$ mice before tamoxifen

administration and SMACre^{ER+}:*Grin2d*$^{f/f}$ mice after tamoxifen administration (Figure 3F). Before the tamoxifen

administration, the K-slopes of the SMACre^{ER+}:*Grin2d*$^{f/f}$ mice and SMACre^{ER+}:*Grin1*$^{f/f}$:*Grin2d*$^{f/f}$ mice showed no

significant difference; similar results were observed after tamoxifen administration. Nevertheless, the K-slope

showed significant differences from before to after tamoxifen administration in each group (Figure 3G). Except

for the functional assay, IF staining demonstrated the colocalization of the NR1 and NR2D subunits in aSMCs

of the MCA (Figure 3C/D). In summary, based on the structural and functional evidence shown above, cKO of

NR2D and NR2D plus NR1 in aSMCs of mice led to similar patterns of functional hyperemia, which highly

suggests that the NR2D and NR1 subunits work together as a complex to form functional NMDA receptors in

cerebral aSMCs to regulate functional hyperemia. Hence, in later experiments, we used *Grin2d* knockouts to

yield similar results (knockout of NMDA receptors) in aSMCs.

aSMC-expressed NR2D works independently of COX-2 and HFD pathways to regulate functional hyperemia

Previous studies demonstrated that COX-2 is a molecule involved in NVC regulation. When COX-2 is knocked

out or inhibited, functional hyperemia is attenuated by approximately 50%. However, total knockout of COX-2

in mice can lead to severe peripheral defects, such as kidney dysfunction (53). Thus, NVC impairment may be

partially due to other global side effects. We wanted to study the similarities and differences between the

aSMCs-NR2D-related pathway and the reported COX-2-related pathway through more local and precise

intervention in the NVC regulation. We administered NS-398, a specific inhibitor of COX-2, locally through the

cistern magnum to inhibit the activity of COX-2 (Figure 4A). By injecting fluorescein isothiocyanate (FITC) at

the same time, we could track the diffusion of NS-398. We selected only mice with successful cistern magnum

administration for the further whisker stimulation assay. The successful injection of NS-398 and FITC clearly

labelled the Virchow-Robin spaces of arterioles (Figure 4B). Through analysis of the whisker stimulation data,

we found that administration of NS-398 to WT (SMACre^{ER-}:*Grin2d*$^{f/f}$) mice significantly decreased the mean

CBF change during the plateau period by approximately 42% (from 13.70% to 7.91%), which is consistent with

the currently known reports. Moreover, in cKO mice administered NS-398, the mean CBF change was further

attenuated to 2.99%, significantly lower than that of the cKO (SMACre^{ER+}:*Grin2d*$^{f/f}$) group without NS-398

(5.96%). A similar trend was observed in the maximum CBF change during the plateau period: the maximum

CBF change of the WT group without NS-398 administration was 20.09%, significantly decreasing to 11.58%%

after NS-398 administration. In the cKO group, the maximum CBF changes without and with NS-398 were

9.48% and 6.88%, respectively, which significantly differed (Figure 4F/H). However, regarding time to plateau,

no significant differences were found among the four groups of mice (Figure 4I). Regarding K-slopes, the WT

group without NS-398 had significantly higher values than the other groups (Figure 4J). Interestingly, through

IF and RT–PCR, we found that *Ptgs2* (gene name of COX-2) and COX-2 were expressed not only in neurons

but also in cerebral aSMCs (Figure 4C/D). Moreover, the RT–PCR results of *Ptgs2* expression in adult cortex

lysates were even lower than those in cultured cerebral aSMCs (Figure 4G). This evidence suggests that COX-

2 expressed in cerebral aSMCs may also contribute to the COX-2 pathway to regulate CBF. The conclusions

drawn from previous studies are not comprehensive. In addition, WB did not detect COX-2 expression in the immunoprecipitated (IP) products of NR2D (Figure 4E). The above evidence suggests that COX-2 and the NR2D subunit in cerebral aSMCs may independently regulate functional hyperemia.

Moreover, we also administered a high-fat diet (HFD) to WT (SMACre^{ER-}:$Grin2d^{f/f}$) and cKO (SMACre^{ER+}:$Grin2d^{f/f}$) mice because a previous study reported a relationship between HFD challenge and functional hyperemia (54). We monitored the food consumption of mice during the HFD challenge (Figure S2G) and the fat/lean/total water content of mice (Figure S2H). These parameters confirmed the successful establishment of the HFD model. After HFD administration, mice underwent whisker stimulation (Figure 4F). During the plateau period, the mean CBF change of HFD_SMACre^{ER-}:$Grin2d^{f/f}$ mice was 5.00%, and the mean CBF change of HFD_SMACre^{ER+}:$Grin2d^{f/f}$ mice was 1.33%, which significantly differed. A similar trend was observed in maximum CBF change; in HFD_WT mice, it was 7.62%, and in HFD_cKO mice, it was 3.03% (Figure 4H). The time to the plateau of HFD_WT and HFD_cKO mice after the HFD challenge was not significantly different (Figure 4I). The K-slope of HFD_cKO mice was significantly lower than that of HFD_WT mice, which indicated that HFD and aSMC-NR2D cKO have a superimposed effect on functional hyperemia. Compared with WT and cKO mice in the control groups, after the HFD challenge, all parameters significantly decreased in both WT and cKO mice (Figure 4J). The above evidence suggests that HFD and the NR2D subunit in cerebral aSMCs may independently regulate functional hyperemia.

Impact of NR2D cKO in aSMCs on physical parameters

In addition to functional experiments, we also physical monitored SMACre^{ER-}:$Grin2d^{f/f}$ and SMACre^{ER+}:$Grin2d^{f/f}$ mice. We tracked the weight of P5 tamoxifen-administered SMACre^{ER+}:$Grin2d^{f/f}$ (P5) mice and p60 SMACre^{ER+}:$Grin2d^{f/f}$ tamoxifen-administered (P60) mice. There was a significant difference in body weight between the P5 group during part of the period from P38 to P60. Specifically, the weight of cKO (SMACre^{ER+}:$Grin2d^{f/f}$) mice was lower than that of their WT (SMACre^{ER-}:$Grin2d^{f/f}$) littermates (controls) (Figure S2A/B). However, in the P60 group, even at P92, WT and cKO mice exhibited no significant difference in body weight (Figure S2C/D). Moreover, at P65, we also monitored the blood pressure of mice. The results showed no significant differences among WT mice, SMACre^{ER+}:$Grin2d^{f/f}$ mice, and SMACre^{ER+}:$Grin1^{f/f}$:$Grin2d^{f/f}$ mice before/after tamoxifen administration in terms of diastolic blood pressure, systolic blood pressure, and mean arterial pressure (Figure S2E). In addition, we monitored the weight of mice before the whisker stimulation

assay at P65 by sampling survey and found no significant difference between the SMACre^{ER-}:*Grin2d*$^{f/f}$ group

and the SMACre^{ER+}:*Grin2d*$^{f/f}$ group (Figure S2F).

Caldesmon may be the key downstream molecule of NR2D-regulated functional hyperemia in cerebral aSMCs

We wanted to discover the critical downstream molecule of the NR2D subunit that regulates functional hyperemia. We first used the lentivirus to overexpress NR2D in cultured SMCs. After NR2D was fully overexpressed, NR2D and the proteins interacting with NR2D were pulled down by immunoprecipitation (IP). Next, we drew a Venn diagram after mass spectrometry analysis of the products from three independent IPs; we found that 75 proteins were detected in all three independent IP experiments (Figure 5A). We selected the proteins that met the following criteria: appeared at least once in the three IP experiments, had a PSM value higher than ten, and were observed in only the NR2D-IP groups or existed in both NR2D-IP groups and immunoglobulin (IgG) control groups, but the PSM values in NR2D-IP groups were more than twice that in the IgG control groups. A total of 916 proteins were selected for Kyoto Encyclopedia of Genes and Genomes (KEGG) and Gene Ontology (GO) analyses. According to the KEGG pathway analysis, the signalling pathway related to vascular SMC contraction was enriched in the IP products of NR2D (Figure 5B). In this pathway, we found that caldesmon was detected in all three independent IP experiments. Caldesmon is critical for cell contraction and dilation. We further verified the reliability of bioinformatics analysis data through WB. In WB, we first assessed the expression of NR2D in IP products. Then, we assessed caldesmon expression in IP products by WB. Caldesmon was detected in the "Input" and the "NR2D-IP" lanes (Figure 5C). Three types of GO analysis of the 916 selected proteins were shown in Figure 5D-F. In short, by analyzing bioinformatics data and performing WB, we found that caldesmon may be the critical downstream molecule that regulates functional hyperemia by altering the contractional and relaxational homeostasis of aSMCs.

Conditional knock-out of NR2D in aSMCs leads mice to express a unique anxiety-like behavior and reduced curiosity

We designed a new open field test (OFT), the open field-like test (OFLT), to explore whether the defect in functional hyperemia induced by NR2D cKO in aSMCs affected a series of emotional functions in mice. In the initial experimental proposal, we aimed to use the OFLT to simultaneously detect anxiety, curiosity, and instinctive fear. In the experiment, we placed WT (SMACre^{ER-}:*Grin2d*$^{f/f}$) mice and cKO (SMACre^{ER+}:*Grin2d*$^{f/f}$) mice in the left part of the box, with the middle of the box blocked with an iron mesh wall (Figure 6A). This area

was similar to the classic OFT, which is used to observe the anxiety-like behavior of mice. A mirrored, symmetrical area was on the opposite side of the iron mesh wall. A CD-1 mouse was placed behind the iron mesh wall. Because CD-1 mice are aggressive, they usually fight with C57 mice when placed them together. Thus, CD-1 mice could generate fear memories in C57 mice. C57 mice were predicted to exhibit spontaneous curiosity and exploration of the iron mesh wall and areas behind it. However, the fear of the CD-1 mouse was predicted to motivate the C57 mouse to avoid the iron mesh wall as much as possible, creating a behavioral conflict between curiosity and fear. However, the pilot experiment did not detect fear behavior from cKO mice in the OFLT. This finding raised doubts about whether the fear intensity generated by the OFLT was sufficient for C57 mice, which would make the OFLT unable to detect fear. Therefore, in the formal experiment, we changed the experimental procedure to assess fear behavior separately.

During the formal behavioral test, SMACre^{ER-}:*Grin2d*$^{f/f}$ and SMACre^{ER+}:*Grin2d*$^{f/f}$ mice were placed in the left area of the box, which was lined with sketch paper to absorb urine produced by mice to enable urine weighing. The top of the box was covered with a transparent acrylic plate (Figure 6A). After a 20-minute behavioral test, we analyzed some parameters. First, we analyzed the distance travelled by mice. WT (SMACre^{ER-}:*Grin2d*$^{f/f}$) mice were more curious about the iron mesh wall and were more willing to stay near the iron mesh wall for exploration. However, cKO (SMACre^{ER+}:*Grin2d*$^{f/f}$) mice seemed less interested in the iron mesh wall. The heatmap showed that the distance travelled by cKO mice was increased compared with that by WT mice, but that exploration near the iron mesh wall was reduced in cKO mice (Figure 6B). Specifically, during the 20-minute behavioral test, the average distance travelled by cKO mice was 30.24 m, significantly higher than that of WT mice (22.98 m) (Figure 6C). In addition, the mean urine weight of cKO mice was 247.81 mg, significantly higher than that of WT mice (78.06 mg) (Figure 6D). The fecal results were similar to those of urine volume, as the number of feces produced was significantly higher in the cKO group (4.9) than in the WT group (2.8) (Figure 6E). In addition, we assessed the preference of the mice for the iron mesh wall and found that the proportion of the time that cKO mice stayed near the iron mesh wall (on the right side of the left half of the whole box) was significantly lower than that of WT mice (60.75% vs. 70.60%). Instead, cKO mice spent more time in the central area of the box than WT mice (19.96% vs. 14.53%) (Figure 6F). Finally, we analyzed the ratio of time spent in the right and left areas (Figure 6G) and the ratio of time spent in the right and all the remaining areas (Figure 6H) in WT and cKO mice. Both of these parameters significantly decreased. Based on the above results, cKO mice exhibited unconventional urine-specific and fecal-specific anxiety-like behaviors and showed reduced curiosity toward the iron mesh wall and the area behind it.

Then, we used a 40-cm square box to conduct the traditional OFT (Figure S5A) to verify our conclusion obtained by the OFLT. In the 15-minute OFT, we found no significant differences between WT (SMACre^{ER-}:*Grin2d*$^{f/f}$) mice and cKO (SMACre^{ER+}:*Grin2d*$^{f/f}$) mice in terms of time spent in the center area (Figure S5B), time spent in the periphery (Figure S5C) and the total distance travelled (Figure S5D). Surprisingly, the probabilities of defecation (Figure S5E) and urination (Figure S5F) were higher in cKO mice than in WT mice. In addition, the number of feces produced by the cKO group was much greater than that produced by the WT group (Figure 6E, 3.38 vs. 0.79). Moreover, regarding the latency to fecal production, the WT group produced the first feces at 517.40s on average, while the cKO group produced the first feces at 285.67s on average, nearly half the latency of the WT group (Figure S5G). In summary, the fecal production latency of cKO mice was shorter than that of WT mice, and the number of feces produced was greater than that of WT mice. The increase in urine and fecal volume in the OFT was consistent with that in the OFLT. Urination and defecation-specific anxiety were conservative in two independent experiments.

Conditional knock-out of NR2D in aSMCs leads mice to exhibit less fear in the CD-1 fear assay

Due to the lack of fear responses in the OFLT, we conducted an acute fear-induction experiment to test whether NVC regulates the fear response. Glutamatergic neurons are present in the amygdala (49), but whether NR2D expressed in cerebral aSMCs in the amygdala are regulated by local glutamatergic neurons and thus influence the fear response is unknown. First, we demonstrated the presence of arterioles and capillaries in the amygdala (Figure S6A), which is the structural fundamental of the behavioral test. Then, we placed CD-1 mice and WT (SMACre^{ER-}:*Grin2d*$^{f/f}$) or cKO (SMACre^{ER+}:*Grin2d*$^{f/f}$) mice together in a cage to directly induce acute fear in WT or cKO mice (Figure 7A). The amygdala is involved in regulating fear memory. Therefore, in addition to recording behavioral parameters, we also performed IF staining of c-Fos in the amygdala of WT and cKO mice to detect the activation of neurons in this area. In this assay, we first exposed WT or cKO mice to CD-1 mice for 10 minutes and recorded their fighting and social behaviors with video cameras. Then, brain extraction of C57 mice was performed 90 minutes later to detect the activation of neurons in the amygdala through c-Fos IF staining.

Behavioral data showed no difference between the WT and cKO groups in the amount (number of bouts) of active sniffing (Figure 7B). Nevertheless, the latency to the first fight in the WT group was significantly shorter than that in the cKO group (58.13 s vs. 201.3 s, Figure 7C). The number of fights in the WT group (4.63 on

average) was also significantly higher than that in the cKO group (1.9 on average) (Figure 7D). In addition, regarding the space use of mice (i.e., close proximity), cKO mice were more willing to share a small space with CD-1 mice than WT mice (161.3 s of cKO mice vs. 88.63 s of WT mice, Figure 7E). The total duration of fighting of the WT group (25.25 s) was greater than that of the cKO group (10.40 s) (Figure 7F). In addition, we conducted a deeper analysis of the fight data of WT and cKO mice. In the WT group, 7/37 (18.92%) fights were initiated by WT C57 mice, and CD-1 mice initiated 30/37 (81.08%) fights. In the cKO group, 5/19 (26.32%) fights were initiated by cKO C57 mice, and 14/19 (73.68%) fights were initiated by CD-1 mice (Figure 7G). In terms of flighted after fights, in 36/37 fights (97.3%), WT mice lost and fled; only one CD-1 mouse lost. However, in the cKO group, cKO C57 mice lost and fled in 7/19 (36.94%) fights, and CD-1 mice lost and fled in 12/19 (63.16%) fights (Figure 7H).

The IF results showed that significantly more amygdala neurons were activated in WT mice than in cKO mice ($88.19/mm^2$ vs. $25.99/mm^2$, Figure 7I). Additionally, the activation of neurons in the barrel cortex (S1BF area) in the WT group ($219.18/mm^2$) was significantly higher than in the cKO group ($109.59/mm^2$) (Figure 7J). S1BF is also the location of the ROI used for CBF detection by whisker stimulation. This result indicates that CBF and neuronal activity decrease in this region is conservative across experiments. However, in cerebral regions unrelated to fear or social interaction, such as the piriform (Pir) cortex and hippocampus, there was no significant difference in the level of neuronal activation between WT and cKO mice (Figure 7K/L). We additionally established control groups without CD-1 mouse exposure (Figure S6 B-E). All statistical data are presented in Figure 7M. Based on the above data, cKO mice exhibited less fear than WT mice in the CD1-1 fear assay, as seen most obviously in the reduced activation of amygdala neurons and the reduced probability of fleeing a fight.

No significant behavioral differences in the elevated plus-maze (EPM) test between SMACre^{ER-}:*Grin2d*$^{f/f}$ and SMACre^{ER+}:*Grin2d*$^{f/f}$ mice

Based on the OFT, OFLT and CD-1 fear assay results, we concluded that SMACre^{ER+}:*Grin2d*$^{f/f}$ mice exhibit anxiety-like and fear-attenuated behaviors. Therefore, we additionally performed an elevated plus-maze (EPM) test. The EPM test evaluates mouse anxiety as a trade-off between curiosity (exploring open arms) and fear (staying in closed arms). In a 5-minute EPM test, we found that the behavior of WT (SMACre^{ER-}:*Grin2d*$^{f/f}$) and cKO (SMACre^{ER+}:*Grin2d*$^{f/f}$) mice were similar. There were no significant differences between WT mice and cKO mice in terms of distance travelled, time spent in the central area, time spent in the closed arms, time

699 spent in the open arms, the proportion of time in the open/closed arms, and proportion of time in the

700 center/periphery (Figure S7). This may be due to the behavior of cKO mice, which both exhibited anxiety-like

701 and fear-attenuated behaviors, which may have led to no net difference in preference for open or closed arms,

702 leading the behavior of cKO mice to be similar to that of WT mice.

Discussion

Unlike previous studies, we abandoned the use of drugs and null mice to study the molecular function of NR2D

but adopt Cre-dependent transgenic mice, which makes our conclusion more cell-specific. According to the

current data, the maximum CBF change during the plateau stage in Pdgfrβ^{ER+}:*Grin2d*$^{f/f}$ mice was lower than

that in SMACre^{ER+}:*Grin2d*$^{f/f}$ mice (Figure 2D), which suggests that there may be low levels of NR2D expression

in pericytes and/or venous SMCs. Subsequent studies should investigate the expression and role of NR2D in

these cells.

Conditional knock-out of NR2D in aSMCs caused an approximately two-thirds decrease in the CBF variation

during the whisker stimulation (Figure 2C). Compared with an approximately 50% decrease in the CBF

variation in the COX-2 total knock-out model (55), it can be inferred that NR2D on aSMCs contributes more to

NVC regulation than COX-2 in all organs. The drawback of COX-2 total knock is the loss of cell specificity and

systemic side effects. Our study found that COX-2 was expressed in not only neurons but also cerebral aSMCs.

Therefore, COX-2 total knock-out also impacts the function of COX-2 in aSMCs. Previous studies found that

COX-2 is expressed only in neurons in the brain (55-58). Thus, these conclusions may not be entirely correct.

Besides, previous studies found that after whisker stimulation, c-Fos-positive cells were enriched in layer IV of

the cortex, but the distribution of COX-2-positive cells was enriched in layers II/III of the cortex (59, 60). Hence,

it would be inaccurate to attribute COX-2-induced regulation of functional hyperemia to neurons alone. Hence,

in subsequent experiments, we plan to explore the contribution of COX-2 expressed in aSMCs to cerebral

functional hyperemia in SMACre^{ER+}:*Ptgs2*$^{f/f}$ mice.

The "time to plateau" and "K-slope" parameters innovative reflect the impact of a particular pathway in two

periods of functional hyperemia. The first period encompasses increases in hyperemia (how quickly CBF

reaches a plateau), and the next reflects hyperemia maintenance (how much CBF increased). If a particular

pathway impacts only the first period, interfering with this pathway will increase the time to plateau and

decrease the K-slope. If a particular pathway impacts only the second period, interfering with this pathway will

decrease the time to the plateau without impacting the K-slope (61). This study is the first to distinguish these

two periods by using parameters "time to plateau" and "K-slope". For example, we can consider data shown

in Figure 2, compared with SMACre^{ER-}:*Grin2d*$^{f/f}$ mice, SMACre^{ER+}:*Grin2d*$^{f/f}$ mice did not differ in the time to

plateau but had a lower K-slope, which means that NR2D was involved in both the increasing and maintenance

periods of hyperemia. Therefore, NR2D could influence two potential pathways in cerebral aSMCs to regulate

functional hyperemia. If NR2D contributed only to glutamate signalling, SMACre^{ER+}:*Grin2d*$^{f/f}$ mice would exhibit

decreased K-slope and increased time to plateau, with the CBF change equal to that of WT mice. If NR2D

contributed only to the maintenance of cellular dilation, SMACre^{ER+}:*Grin2d*$^{f/f}$ mice would exhibit no changes in

K-slope and decreased time to plateau, with the CBF change lower than that in WT mice. However,

SMACre^{ER+}:*Grin2d*$^{f/f}$ mice did not differ in time to plateau, although the K-slope decreased. Hence, based on

these parameters, we speculate that NR2D in cerebral aSMCs participates in glutamate signalling and the

maintenance of cellular dilation. These two parameters make it possible to accurately analyze the increase of

CBF in the future.

We identified caldesmon, a critical downstream protein of NR2D, as regulating CBF in aSMCs through IP data.

Previous reports have revealed that nonphosphorylated caldesmon binds with actin and myosin to prevent

their interaction, thereby inhibiting the contraction of the cell skeleton (62-65). Through phosphorylation of

MAPK (66) and PAK (67) or binding with calcium-activated CaM (68), caldesmon is removed from the binding

site of actin and myosin, leading to the contraction of the cytoskeleton. Therefore, if the knockout of NR2D in

aSMCs affects the phosphorylation of caldesmon or calcium activation of CaM, it is likely to lead to abnormal

cell contraction and relaxation. In addition, caldesmon and arp2/3 are antagonistic, and increased caldesmon

expression will affect the expression of arp2/3 (69, 70). Arp2/3 is responsible for the assembly and extension

of the actin skeleton to maintain an angle of 70° between two actin filaments. If the loss of NR2D in aSMCs

leads to the abnormal expression of caldesmon, it is likely to further lead to the abnormal expression of arp2/3,

which affects the assembly of actin and the angle of actin skeletons, thus affecting the normal contraction and

relaxation of cells. Therefore, future experiments should focus on manipulate caldesmon in aSMCs-NR2D cKO

mice to further study the role of the NR2D-caldesmon pathway in regulating cerebral aSMC contraction and

relaxation.

The change in emotional behavior is the most important discovery of this study. We first link the correlation of

NVC and emotional disorders in one study by the transgenic strategy. Compared with the extensive behavioral

changes produced by the traditional anxiety-like model, SMACre^{ER+}:*Grin2d*$^{f/f}$ mice showed an increase in the production of urine and defecation, which may indicate that the cKO of NR2D in aSMCs specifically affected the cerebral regions of anxiety related to urination and defecation in mice. Clinical studies have confirmed that patients with anxiety show a significant increase in defecation (36), which is consistent with our report. The correlation between anxiety and increased defecation is conservative across species. This precise effect may find a more specific target for treating this type of anxiety and avoiding the side effects caused by the widespread use of drugs.

In addition, the fecal production of both SMACre^{ER-}:*Grin2d*$^{f/f}$ and SMACre^{ER+}:*Grin2d*$^{f/f}$ mice was higher in the OFLT than in the OFT. On the one hand, the OFLT lasted for 20 minutes, while the OFT lasted for 15 minutes. On the other hand, the OFLT box (26 × 26 cm) was smaller than the OFT box (40 × 40 cm). The overall anxiety level of mice may be higher in a narrower space. Based on physical monitoring during the HFD challenge, food consumption and changes in fat content of SMACre^{ER-}:*Grin2d*$^{f/f}$ mice and SMACre^{ER+}:*Grin2d*$^{f/f}$ mice were similar (Figure S2 G/H). These findings indicated that the increase in defecation of SMACre^{ER+}:*Grin2d*$^{f/f}$ mice in the OFT and OFLT may not be due to differences in food consumption.

The increase of CBF in the amygdala during fear is conservative across species. A previous study on depressed adolescents showed a decreased CBF baseline in the amygdala (71). Hence, a stable CBF supply in the amygdala is critical for emotional homeostasis. We hypothesize that the "CBF-Neuron-Behavior" axis is how NVC regulates the cerebral complex emotional behaviors. Our manipulation only influenced the expression of NR2D in aSMCs (CBF level), and this manipulation decreased c-Fos expression in neurons of the amygdala (neuronal level). Finally, it caused the fear attenuation of mice (behavioral level). Therefore, this study is the first to show the high correlation between NVC regulation and advanced cerebral functions. Besides, what makes us curious is the cellular identity of neurons not activated in SMACre^{ER+}:*Grin2d*$^{f/f}$ mice after the acute fear stimulation. So, we expected to explore the identity of these groups of neurons, and these neurons may both blood supply-sensitive and participate in the regulation of fear.

In the EPM test, mice can explore open arms. However, their preference for dark areas and fear of heights leads them to fear open arms. The conflict between exploration (curiosity) and avoidance (fear) can thus be assessed. In our study, SMACre^{ER+}:*Grin2d*$^{f/f}$ mice showed anxiety-like and fear-attenuated behaviors, leading to contradictory behavior. Because of their increased anxiety, SMACre^{ER+}:*Grin2d*$^{f/f}$ mice did not want to explore

open arms. However, because these mice felt less fear of heights (fear-attenuated behavior), they did not fear open arms. Thus, there was no difference between WT and SMACre^{ER+}:*Grin2d*$^{f/f}$ mice regarding exploring open arms and closed arms in the EPM test (Figure S7). Because SMACre^{ER+}:*Grin2d*$^{f/f}$ mice exhibited both anxiety-like and fear-attenuated behaviors, there was no net change in behavior. The results of the EPM test further confirm that SMACre^{ER+}:*Grin2d*$^{f/f}$ mice exhibited increased anxiety-like behaviors and attenuated fear simultaneously.

Encouragingly, we only intervened in the expression of NR2D in cerebral aSMCs, which only occupied 0.3% of total cells in the mouse brain, enough to change the CBF variation and advanced emotional behaviors of mice. This phenomenon suggested that continually decreased CBF supply during neuron activation is a vital trigger factor for advanced behavioral dysfunction. As the absence of NR2D in cerebral aSMCs leads to abnormalities in higher-order cognitive functions, NR2D is a promising target for treating anxiety-related and amygdala-related mental disorders or diseases. This study provides new strategies and ideas for treating psychiatric diseases and can inform drug development by interfering with molecules expressed in cerebral aSMCs to treat mental disorders.

Acknowledgments

We thank Yanmei Tao for sharing Nestin-Cre mice. We thank Bo Zhou for sharing SMAcre^{ER} mice. We thank Zilong Qiu for sharing Ai47 mice. We thank Xiang Yu for sharing PdgfrβCre^{ER} mice. We thank Yangang Sun for sharing *Grin1* mice. We thank Yajie Liu from Westlake University Laboratory Animal Resource Center for guiding behaviors assays. We thank Yuxiao Niu from Yanxiao Zhang lab of Westlake University for drawing KEGG pathway map.

Fundings

J.‑M.J. acknowledges supports from Westlake University startup funding, the Westlake Education Foundation, the Zhejiang Province Natural Science Foundation (Project # 2022XHSJJ004), the Westlake Laboratory of Life Sciences and Biomedicine (Project# 202109013).

Conflict of Interest

821   We declare that we do not have any financial or personal relationships with other people or organizations,

822   which may affect our work inappropriately, and there are no professional or other personal interests of any

823   nature or kind in any products, services and/or companies. These interests may be interpreted as affecting or

824   reviewing the positions in the following manuscripts.

References

1.   Attwell D, Buchan AM, Charpak S, Lauritzen M, Macvicar BA, Newman EA. Glial and neuronal control

of brain blood flow. Nature. 2010;468(7321):232-43.

2.   Lauritzen M, Dreier JP, Fabricius M, Hartings JA, Graf R, Strong AJ. Clinical relevance of cortical

spreading depression in neurological disorders: migraine, malignant stroke, subarachnoid and intracranial

hemorrhage, and traumatic brain injury. J Cereb Blood Flow Metab. 2011;31(1):17-35.

3.   Girouard H, Iadecola C. Neurovascular coupling in the normal brain and in hypertension, stroke, and

Alzheimer disease. J Appl Physiol (1985). 2006;100(1):328-35.

4.   Ruitenberg A, den Heijer T, Bakker SL, van Swieten JC, Koudstaal PJ, Hofman A, et al. Cerebral

hypoperfusion and clinical onset of dementia: the Rotterdam Study. Ann Neurol. 2005;57(6):789-94.

5.   Pico-Perez M, Radua J, Steward T, Menchon JM, Soriano-Mas C. Emotion regulation in mood and

anxiety disorders: A meta-analysis of fMRI cognitive reappraisal studies. Prog Neuropsychopharmacol Biol

Psychiatry. 2017;79(Pt B):96-104.

6.   Iadecola C. Neurovascular regulation in the normal brain and in Alzheimer's disease. Nat Rev

Neurosci. 2004;5(5):347-60.

7.   Drew PJ. Neurovascular coupling: motive unknown. Trends Neurosci. 2022;45(11):809-19.

8.   Hillman EM. Coupling mechanism and significance of the BOLD signal: a status report. Annu Rev

Neurosci. 2014;37:161-81.

9.   Clemm von Hohenberg C, Weber-Fahr W, Lebhardt P, Ravi N, Braun U, Gass N, et al. Lateral

habenula perturbation reduces default-mode network connectivity in a rat model of depression. Transl

Psychiatry. 2018;8(1):68.

10.   Kamiya K, Abe O. Imaging of Posttraumatic Stress Disorder. Neuroimaging Clin N Am.

2020;30(1):115-23.

11.   Ahmed-Leitao F, Spies G, van den Heuvel L, Seedat S. Hippocampal and amygdala volumes in adults

with posttraumatic stress disorder secondary to childhood abuse or maltreatment: A systematic review.

Psychiatry Res Neuroimaging. 2016;256:33-43.

12.   Pitman RK, Rasmusson AM, Koenen KC, Shin LM, Orr SP, Gilbertson MW, et al. Biological studies of

post-traumatic stress disorder. Nat Rev Neurosci. 2012;13(11):769-87.

13.   Whalen PJ, Kagan J, Cook RG, Davis FC, Kim H, Polis S, et al. Human amygdala responsivity to

masked fearful eye whites. Science. 2004;306(5704):2061.

14.   Hadj-Bouziane F, Bell AH, Knusten TA, Ungerleider LG, Tootell RB. Perception of emotional

expressions is independent of face selectivity in monkey inferior temporal cortex. Proc Natl Acad Sci U S A.

2008;105(14):5591-6.

15.   Harris AP, Lennen RJ, Marshall I, Jansen MA, Pernet CR, Brydges NM, et al. Imaging learned fear

circuitry in awake mice using fMRI. Eur J Neurosci. 2015;42(5):2125-34.

16.   Ofek K, Schoknecht K, Melamed-Book N, Heinemann U, Friedman A, Soreq H. Fluoxetine induces

vasodilatation of cerebral arterioles by co-modulating NO/muscarinic signalling. J Cell Mol Med.

2012;16(11):2736-44.

17.   Malberg JE, Duman RS. Cell proliferation in adult hippocampus is decreased by inescapable stress:

reversal by fluoxetine treatment. Neuropsychopharmacology. 2003;28(9):1562-71.

18.   Li RW, Yang C, Kwan YW, Chan SW, Lee SM, Leung GP. Involvement of organic cation transporter-

3 and plasma membrane monoamine transporter in serotonin uptake in human brain vascular smooth muscle

cells. Front Pharmacol. 2013;4:14.

19.   Maddahi A, Edvinsson L. Enhanced expressions of microvascular smooth muscle receptors after focal

cerebral ischemia occur via the MAPK MEK/ERK pathway. BMC Neurosci. 2008;9:85.

20.   Cao L, Xu CB, Zhang Y, Cao YX, Edvinsson L. Secondhand cigarette smoke exposure causes

upregulation of cerebrovascular 5-HT(1) (B) receptors via the Raf/ERK/MAPK pathway in rats. Acta Physiol

(Oxf). 2013;207(1):183-93.

21.   Li W, Prakash R, Chawla D, Du W, Didion SP, Filosa JA, et al. Early effects of high-fat diet on

neurovascular function and focal ischemic brain injury. Am J Physiol Regul Integr Comp Physiol.

2013;304(11):R1001-8.

22.   Remzso G, Nemeth J, Toth-Szuki V, Varga V, Kovacs V, Domoki F. NMDA attenuates the

neurovascular response to hypercapnia in the neonatal cerebral cortex. Sci Rep. 2019;9(1):18900.

23.   Chow BW, Nunez V, Kaplan L, Granger AJ, Bistrong K, Zucker HL, et al. Caveolae in CNS arterioles

mediate neurovascular coupling. Nature. 2020;579(7797):106-10.

24.   Offenhauser N, Thomsen K, Caesar K, Lauritzen M. Activity-induced tissue oxygenation changes in

rat cerebellar cortex: interplay of postsynaptic activation and blood flow. J Physiol. 2005;565(Pt 1):279-94.

25.   St Lawrence KS, Ye FQ, Lewis BK, Frank JA, McLaughlin AC. Measuring the effects of indomethacin

on changes in cerebral oxidative metabolism and cerebral blood flow during sensorimotor activation. Magn

Reson Med. 2003;50(1):99-106.

26.   Busija DW, Bari F, Domoki F, Louis T. Mechanisms involved in the cerebrovascular dilator effects of

N-methyl-d-aspartate in cerebral cortex. Brain Res Rev. 2007;56(1):89-100.

27.   Roman RJ. P-450 metabolites of arachidonic acid in the control of cardiovascular function. Physiol

Rev. 2002;82(1):131-85.

28.   Alkayed NJ, Narayanan J, Gebremedhin D, Medhora M, Roman RJ, Harder DR. Molecular

characterization of an arachidonic acid epoxygenase in rat brain astrocytes. Stroke. 1996;27(5):971-9.

29.   Amruthesh SC, Boerschel MF, McKinney JS, Willoughby KA, Ellis EF. Metabolism of arachidonic acid

to epoxyeicosatrienoic acids, hydroxyeicosatetraenoic acids, and prostaglandins in cultured rat hippocampal

astrocytes. J Neurochem. 1993;61(1):150-9.

30.   Hogan-Cann AD, Lu P, Anderson CM. Endothelial NMDA receptors mediate activity-dependent brain

hemodynamic responses in mice. Proc Natl Acad Sci U S A. 2019;116(21):10229-31.

31.   Lu L, Hogan-Cann AD, Globa AK, Lu P, Nagy JI, Bamji SX, et al. Astrocytes drive cortical vasodilatory

signaling by activating endothelial NMDA receptors. J Cereb Blood Flow Metab. 2019;39(3):481-96.

32.   Davis M. NMDA receptors and fear extinction: implications for cognitive behavioral therapy. Dialogues

Clin Neurosci. 2011;13(4):463-74.

33.   Shultz B, Farkash A, Collins B, Mohammadmirzaei N, Knox D. Fear learning-induced changes in

AMPAR and NMDAR expression in the fear circuit. Learn Mem. 2022;29(3):83-92.

34.   Feyissa AM, Chandran A, Stockmeier CA, Karolewicz B. Reduced levels of NR2A and NR2B subunits

of NMDA receptor and PSD-95 in the prefrontal cortex in major depression. Prog Neuropsychopharmacol Biol

Psychiatry. 2009;33(1):70-5.

35.   Yamamoto H, Kamegaya E, Hagino Y, Takamatsu Y, Sawada W, Matsuzawa M, et al. Loss of GluN2D

subunit results in social recognition deficit, social stress, 5-HT(2C) receptor dysfunction, and anhedonia in

mice. Neuropharmacology. 2017;112(Pt A):188-97.

36.   Gorard DA, Gomborone JE, Libby GW, Farthing MJ. Intestinal transit in anxiety and depression. Gut.
1996;39(4):551-5.

37.   Li X, Zhou L, Zhang X, Jin Y, Zhao B, Zhang D, et al. Proteins secreted by brain arteriolar smooth
muscle cells are instructive for neural development. Mol Brain. 2022;15(1):97.

38.   Yu XD, Zhu Y, Sun QX, Deng F, Wan J, Zheng D, et al. Distinct serotonergic pathways to the amygdala
underlie separate behavioral features of anxiety. Nat Neurosci. 2022;25(12):1651-63.

39.   Boas DA, Dunn AK. Laser speckle contrast imaging in biomedical optics. J Biomed Opt.
2010;15(1):011109.

40.   Jin Y, Shi P, Wang Y, Li J, Zhang J, Zhao X, et al. Precise control of embolic stroke with magnetized
red blood cells in mice. Commun Biol. 2022;5(1):136.

41.   Liu CY, Chen JB, Liu YY, Zhou XM, Zhang M, Jiang YM, et al. Saikosaponin D exerts antidepressant
effect by regulating Homer1-mGluR5 and mTOR signaling in a rat model of chronic unpredictable mild stress.
Chin Med. 2022;17(1):60.

42.   Guo Z, Li S, Wu J, Zhu X, Zhang Y. Maternal Deprivation Increased Vulnerability to Depression in
Adult Rats Through DRD2 Promoter Methylation in the Ventral Tegmental Area. Front Psychiatry.
2022;13:827667.

43.   Kim S, Hong KB, Kim S, Suh HJ, Jo K. Creatine and taurine mixtures alleviate depressive-like
behaviour in Drosophila melanogaster and mice via regulating Akt and ERK/BDNF pathways. Sci Rep.
2020;10(1):11370.

44.   Lapmanee S, Charoenphandhu J, Teerapornpuntakit J, Krishnamra N, Charoenphandhu N.
Agomelatine, venlafaxine, and running exercise effectively prevent anxiety- and depression-like behaviors and
memory impairment in restraint stressed rats. PLoS One. 2017;12(11):e0187671.

45.   Borelli KG, Blanchard DC, Javier LK, Defensor EB, Brandao ML, Blanchard RJ. Neural correlates of
scent marking behavior in C57BL/6J mice: detection and recognition of a social stimulus. Neuroscience.
2009;162(4):914-23.

46.   Arakawa H, Arakawa K, Blanchard DC, Blanchard RJ. Scent marking behavior in male C57BL/6J mice:
sexual and developmental determination. Behav Brain Res. 2007;182(1):73-9.

47.   Arakawa H, Arakawa K, Blanchard DC, Blanchard RJ. A new test paradigm for social recognition
evidenced by urinary scent marking behavior in C57BL/6J mice. Behav Brain Res. 2008;190(1):97-104.

48.   McQuaid RJ, Audet MC, Anisman H. Environmental enrichment in male CD-1 mice promotes

aggressive behaviors and elevated corticosterone and brain norepinephrine activity in response to a mild

stressor. Stress. 2012;15(3):354-60.

49.   Liu J, Li D, Huang J, Cao J, Cai G, Guo Y, et al. Glutamatergic Neurons in the Amygdala Are Involved

in Paclitaxel-Induced Pain and Anxiety. Front Psychiatry. 2022;13:869544.

50.   Traynelis SF, Wollmuth LP, McBain CJ, Menniti FS, Vance KM, Ogden KK, et al. Glutamate receptor

ion channels: structure, regulation, and function. Pharmacol Rev. 2010;62(3):405-96.

51.   Monyer H, Burnashev N, Laurie DJ, Sakmann B, Seeburg PH. Developmental and regional expression

in the rat brain and functional properties of four NMDA receptors. Neuron. 1994;12(3):529-40.

52.   Laurie DJ, Bartke I, Schoepfer R, Naujoks K, Seeburg PH. Regional, developmental and interspecies

expression of the four NMDAR2 subunits, examined using monoclonal antibodies. Brain Res Mol Brain Res.

1997;51(1-2):23-32.

53.   Slattery P, Frolich S, Goren I, Nusing RM. Salt supplementation ameliorates developmental kidney

defects in COX-2(-/-) mice. Am J Physiol Renal Physiol. 2017;312(6):F1044-F55.

54.   Huber G, Ogrodnik M, Wenzel J, Stolting I, Huber L, Will O, et al. Telmisartan prevents high-fat diet-

induced neurovascular impairments and reduces anxiety-like behavior. J Cereb Blood Flow Metab.

2021;41(9):2356-69.

55.   Niwa K, Araki E, Morham SG, Ross ME, Iadecola C. Cyclooxygenase-2 contributes to functional

hyperemia in whisker-barrel cortex. J Neurosci. 2000;20(2):763-70.

56.   Yamagata K, Andreasson KI, Kaufmann WE, Barnes CA, Worley PF. Expression of a mitogen-

inducible cyclooxygenase in brain neurons: regulation by synaptic activity and glucocorticoids. Neuron.

1993;11(2):371-86.

57.   Breder CD, Dewitt D, Kraig RP. Characterization of inducible cyclooxygenase in rat brain. J Comp

Neurol. 1995;355(2):296-315.

58.   Kaufmann WE, Worley PF, Pegg J, Bremer M, Isakson P. COX-2, a synaptically induced enzyme, is

expressed by excitatory neurons at postsynaptic sites in rat cerebral cortex. Proc Natl Acad Sci U S A.

1996;93(6):2317-21.

59.   Lacroix A, Toussay X, Anenberg E, Lecrux C, Ferreiros N, Karagiannis A, et al. COX-2-Derived

Prostaglandin E2 Produced by Pyramidal Neurons Contributes to Neurovascular Coupling in the Rodent

Cerebral Cortex. J Neurosci. 2015;35(34):11791-810.

60.   Lecrux C, Toussay X, Kocharyan A, Fernandes P, Neupane S, Levesque M, et al. Pyramidal neurons

are "neurogenic hubs" in the neurovascular coupling response to whisker stimulation. J Neurosci.

2011;31(27):9836-47.

61.   Koizumi K, Hattori Y, Ahn SJ, Buendia I, Ciacciarelli A, Uekawa K, et al. Apoepsilon4 disrupts

neurovascular regulation and undermines white matter integrity and cognitive function. Nat Commun.

2018;9(1):3816.

62.   Kordowska J, Huang R, Wang CL. Phosphorylation of caldesmon during smooth muscle contraction

and cell migration or proliferation. J Biomed Sci. 2006;13(2):159-72.

63.   Wang CL. Caldesmon and smooth-muscle regulation. Cell Biochem Biophys. 2001;35(3):275-88.

64.   Hai CM, Gu Z. Caldesmon phosphorylation in actin cytoskeletal remodeling. Eur J Cell Biol. 2006;85(3-

4):305-9.

65.   Wang CL. Caldesmon and the regulation of cytoskeletal functions. Adv Exp Med Biol. 2008;644:250-

72.

66.   Childs TJ, Watson MH, Sanghera JS, Campbell DL, Pelech SL, Mak AS. Phosphorylation of smooth

muscle caldesmon by mitogen-activated protein (MAP) kinase and expression of MAP kinase in differentiated

smooth muscle cells. J Biol Chem. 1992;267(32):22853-9.

67.   Foster DB, Shen LH, Kelly J, Thibault P, Van Eyk JE, Mak AS. Phosphorylation of caldesmon by p21-

activated kinase. Implications for the Ca(2+) sensitivity of smooth muscle contraction. J Biol Chem.

2000;275(3):1959-65.

68.   Janovick JA, Natarajan K, Longo F, Conn PM. Caldesmon: a bifunctional (calmodulin and actin)

binding protein which regulates stimulated gonadotropin release. Endocrinology. 1991;129(1):68-74.

69.   Yamakita Y, Oosawa F, Yamashiro S, Matsumura F. Caldesmon inhibits Arp2/3-mediated actin

nucleation. J Biol Chem. 2003;278(20):17937-44.

70.   Morita T, Mayanagi T, Yoshio T, Sobue K. Changes in the balance between caldesmon regulated by

p21-activated kinases and the Arp2/3 complex govern podosome formation. J Biol Chem. 2007;282(11):8454-

63.

71.   Ho TC, Wu J, Shin DD, Liu TT, Tapert SF, Yang G, et al. Altered cerebral perfusion in executive,

affective, and motor networks during adolescent depression. J Am Acad Child Adolesc Psychiatry.

2013;52(10):1076-91 e2.

Figure legends

	Main Figures

Figure 1. Expression of *Grin2d*/NR2D in cerebral aSMCs. A) Expression of all NMDAR receptors in mRNA

level detected by RT-PCR. B) Statistical data of Figure 1A. Each dot indicates an independent experiment,

and the data are mean ±s.e.m. C) RNA scope results of *Grin2d* expression in cerebral aSMCs. Red is the

signal of aSMCs. Green is the signal of the *Grin2d* probe. Blue is the signal of the nucleus. Histogram of the

red, green, and blue fluorescent intensities of the *Grin2d* signal inside the nucleus of smooth muscle cells

pointed out by the red arrow. D) WB of NR2D expression in P0 Nestin-cre$^-$:*Grin2d*$^{f/f}$ cerebellum, Adult Nestin-

cre$^-$:*Grin2d*$^{f/f}$ cerebellum, and P0 Nestin-cre$^+$:*Grin2d*$^{f/f}$ cerebellum. E) WB of NR2D expression in P10

SMAcre^{ER-}:*Grin2d*$^{f/f}$ cerebellum and P10 SMAcre^{ER+}:*Grin2d*$^{f/f}$ cerebellum. F) RT-PCR of *Grin2d* expression in

SMAcre^{ER+}:*Grin2d*$^{+/+}$:Ai47 sorted pial aSMCs and SMAcre^{ER+}:*Grin2d*$^{f/f}$:Ai47 sorted pial aSMCs. G) WB of

NR2D expression in SMAcre^{ER+}:*Grin2d*$^{+/+}$:Ai47 sorted pial aSMCs and SMAcre^{ER+}:*Grin2d*$^{f/f}$:Ai47 sorted pial

aSMCs. H-K), Statistical data of Figure 1D-G. Each dot indicates an independent experiment, and the data

are mean ±s.e.m. L) IF of NR2D expression in MCA of SMAcre^{ER-}:*Grin2d*$^{f/f}$ and SMAcre^{ER+}:*Grin2d*$^{f/f}$ (Top two

rows). Red is the signal of smoothelin. Green is the signal of NR2D. Blue is the signal of the nucleus. The

bottom row is the IF of NR2D in SMAcre^{ER+}:*Grin2d*$^{+/+}$:Ai47 sorted pial aSMCs and SMAcre^{ER+}:*Grin2d*$^{f/f}$:Ai47

sorted pial aSMCs. Red is the signal of NR2D. White is the signal of smooth muscle cells. M) Statistical data

of NR2D expression in MCA of Figure 1L. Each dot indicates a smooth muscle cell. Data are mean ±s.e.m.

N) Statistical data of Smoothelin expression in MCA of Figure 1L. Each dot indicates a smooth muscle cell.

Data are mean ±s.e.m. O) Statistical data of NR2D expression in cultured aSMCs of Figure 1L. Each dot

indicates an independent experiment, and the data are mean ±s.e.m. P) Statistical data of SMA expression

in cultured aSMCs through WB. Each dot indicates an independent experiment, and the data are mean ±

s.e.m. RFI, Relative Fluorescence Intensity. T-test used for statistical analysis. **** means $P<0.0001$.

1021

Figure 2. Cerebral blood flow variation of SMAcre^{ER+}/Pdgfrβ^{ER+}:*Grin2d*$^{f/f}$ mice during the whisker stimulation.

A) Schematic diagram of experimental timeline and operations. B) Schematic diagram of CBF increases in the

ROI before, during, and after the whisker stimulation of SMAcre^{ER-}:*Grin2d*$^{f/f}$ mice and SMAcre^{ER+}:*Grin2d*$^{f/f}$ mice.

In the right area, the redder the colour is, the more CBF increases, and the bluer the colour is, the more CBF

decreases. C) CBF variation before, during, and after the whisker stimulation. The solid line indicates the mean

value, and the dotted line indicates ±s.e.m. D) The mean and maximum value of CBF variation during the

plateau period. E) The value "Time to plateau" of mice. F) The value "K-slope" of mice. In Figure 2, Each group

contains six mice, except the P60 TMX_ Pdgfrβ^{ER+}:*Grin2d*$^{f/f}$ group contains five mice. Each mouse was

stimulated at least twice. Each dot indicates the mean value obtained from one mouse, and the data are mean ±s.e.m. T-test used for statistical analysis. **** means P<0.0001.

Figure 3. Cerebral blood flow variation of SMAcre[ER+]:*Grin2d*[f/f] mice and SMAcre[ER+]:*Grin1*[f/f]:*Grin2d*[f/f] mice during the whisker stimulation. A) Schematic diagram of experimental timeline and operations. B) CBF variation before, during, and after the whisker stimulation. The solid line indicates the mean value, and the dotted line indicates ±s.e.m. C) The IF of colocalization of NR2D and NR1 in MCA. Blue is the signal of smooth muscle cells. Green is the signal of NR2D. Red is the signal of NR1. White is the signal of nucleus. D) Statistical data of Figure 3C. Each dot indicates an independent experiment. Data are mean ±s.e.m. E) The mean and maximum value of CBF variation during the plateau period. F) The value "Time to plateau" of mice. G) The value "K-slope" of mice. In Figure 3, each group contains four mice. Each mouse was stimulated at least twice. Each dot indicates the mean value obtained from one mouse, and the data are mean ±s.e.m. T-test used for statistical analysis. **** means P<0.0001.

Figure 4. Cerebral blood flow variation of SMAcre[ER-]:*Grin2d*[f/f] mice and SMAcre[ER+]:*Grin2d*[f/f] mice with COX-2 inhibition and HFD challenge during the whisker stimulation. A) Schematic diagram of experimental timeline and operations. B) Schematic diagram of the validation of cistern magnum injection. NS-398 mixed with FITC. The green signal of FITC can be detected in the Virchow-Robin spaces of cerebral blood vessels. C) RT-PCR of *Ptgs2* expression in adult cortex and sorted SMAcre[ER+]:Ai47 parenchymal aSMCs. D) IF of COX-2 in cerebral smooth muscle cells in the cortex of mice. The red signal is smooth muscle cells, green is the signal of COX-2, and white is the signal of the nucleus. The Arrow shows the signal of COX-2 expressed inside the signal of smooth muscle cells. E) WB of COX-2 expression validation in the IP products. F) CBF variation before, during, and after the whisker stimulation. The solid line indicates the mean value, and the dotted line indicates ± s.e.m. G) Statistical data of Figure 4C. H) The mean and maximum value of CBF variation during the plateau period. I) The value "Time to plateau" of mice. J) The value "K-slope" of mice. In Figure 4, each group contains three mice. Each mouse was stimulated at least twice. Each dot indicates an independent experiment, and the data are mean ±s.e.m. T-test used for statistical analysis. **** means P<0.0001.

Figure 5. Downstream signal pathways analysis of NR2D in aSMCs. A) Veen diagram of NR2D-related proteins obtained from three independent mass spectrum data. B) KEGG pathway analysis of 915 proteins selected from three independent mass spectrum data. C) WB of NR2D and caldesmon expression validation

in the IP products. D-F) GO analysis of 915 proteins selected from three independent mass spectrum data. They are GO_BP, GO_CC, and GO_MF, respectively.

Figure 6. The open field-like test (OFLT) of SMAcre^{ER-}:*Grin2d*$^{f/f}$ (WT) mice and SMAcre^{ER+}:*Grin2d*$^{f/f}$ (cKO) mice. A) Schematic diagram of the box used for the OFLT. B) Schematic diagram of the heat map of WT and cKO mice in the box during 20min. C) Running distance of WT and cKo mice. D) Urine volume of WT and cKo mice. E) Fecal particles of WT and cKo mice in OFT and OFLT. F) Box area preference of WT and cKo mice. G) Ratio of time spent in the right and left area of WT and cKO mice. Data presented in log10 format. H) Ratio of time spent in the right and the rest of area of WT and cKO mice. Data presented in log10 format. T-test used for statistical analysis. Each dot indicates one mouse, and the data are mean ±s.e.m.

Figure 7. CD-1 fear assay of SMAcre^{ER-}:*Grin2d*$^{f/f}$ (WT) mice and SMAcre^{ER+}:*Grin2d*$^{f/f}$ (cKO) mice. A) Schematic diagram of the cage for CD-1 fear assay. B) The value "Number of active sniffing" of mice. C) The value "First fight latency" of mice. D) The value "Number of fights" of mice. E) The value "Two mice in tiny space together" of mice. F) The value "Total fighting duration" of mice. G) The percentage of "Who triggered the fight" of mice. H) The percentage of "Who flighted after the fight" of mice. I-L) IF of c-Fos expression in the amygdala, S1BF, pir cortex, and hippocampus of WT and cKO mice. Red, the signal of NeuN. Green, the signal of c-Fos. M) Statistical data of Figure 7I-L, including the data of controls (Figure S6B-E). Each dot indicates one mouse, and the data are mean ±s.e.m. T-test used for statistical analysis. **** means P<0.0001.

Supplementary Figures

Figure S1. RNA scope probe specificity, SMAcreER specificity, and NR2D expression specificity validation. A) RNA scope of *Grin2d* and *Grin1* in mice cerebellum. Red is the signal of the *Grin1* probe. Green is the signal of the *Grin2d* probe. Blue is the signal of the nucleus. B) IF of NeuN in the brain slices of SMAcre^{ER+}:Ai14 mice. Red is the signal of the smooth muscle cell. Green is the signal of CD-31. Blue is the signal of NeuN, and the white signal is for the nucleus. In the statistical data, each dot indicates an ROI of analysis. Three independent experiments were performed, and the data are mean ±s.e.m. C) IF of NR2D expression in heart, liver, spleen, lung, kidney, and small intestine of SMAcre^{ER+}:Ai14 mice. Red is the signal of the smooth muscle cell. Green is the signal of NR2D. Blue is the signal of the nucleus. The first row is zoom-out images of six organs, the second row is zoom-in images of white circles in the first row, and the third row is histograms of

the red and green fluorescent intensities of the indicated sites in the second row. RFI, Relative Fluorescence Intensity. T-test used for statistical analysis. **** means P<0.001.

Figure S2. Physical parameters of SMAcre[ER-]:*Grin2d*[f/f] mice and SMAcre[ER+]:*Grin2d*[f/f] mice. A) The absolute weight change of SMAcre[ER-]:*Grin2d*[f/f] mice and SMAcre[ER+]:*Grin2d*[f/f] mice. Tamoxifen was administrated at P5. The solid line indicates the mean value, and the dotted line indicates ±s.e.m. B) The relative weight change in Figure S2A. The solid line indicates the mean value, and the dotted line indicates ±s.e.m. C) The absolute weight change of SMAcre[ER-]:*Grin2d*[f/f] mice and SMAcre[ER+]:*Grin2d*[f/f] mice. Tamoxifen was administrated at P60. The solid line indicates the mean value, and the dotted line indicates ±s.e.m. D) The relative weight change in Figure S2C. The solid line indicates the mean value, and the dotted line indicates ±s.e.m. E) The blood pressure of mice. F) The body weight of SMAcre[ER-]:*Grin2d*[f/f] mice and SMAcre[ER+]:*Grin2d*[f/f] mice before the whisker stimulation at P65. G) High-fat food consumption of SMAcre[ER-]:*Grin2d*[f/f] mice and SMAcre[ER+]:*Grin2d*[f/f] mice. H) Fat/lean/total water of SMAcre[ER-]:*Grin2d*[f/f] mice and SMAcre[ER+]:*Grin2d*[f/f] mice under HFD challenge. Each dot indicates one mouse, and the data are mean ±s.e.m. T-test used for statistical analysis. * means P=0.01-0.05, and ** means P<0.01. **** means P<0.0001.

Figure S3. Cerebral blood flow variation of SMAcre[ER-]:*Grin2d*[f/f] mice in the whisker stimulation before and after long-term stimulation. A) CBF variation before, during, and after the whisker stimulation. The solid line indicates the mean value, and the dotted line indicates ±s.e.m. B) The mean and maximum value of CBF variation during the plateau period. C) The value "Time to plateau" of mice. D) The value "K-slope" of mice. In figure S3, four SMAcre[ER-]:*Grin2d*[f/f] mice were used in the experiment. T-test used for statistical analysis. Each dot indicates the mean value obtained from one mouse, and the data are mean ±s.e.m.

Figure S4. Cerebral blood flow variation of SMAcre[ER-]:*Grin2d*[f/f] mice in the whisker stimulation before and after Tamoxifen administration. A) CBF variation before, during, and after the whisker stimulation. The solid line indicates the mean value, and the dotted line indicates ±s.e.m. B) The mean and maximum value of CBF variation during the plateau period. C) The value "Time to plateau" of mice. D) The value "K-slope" of mice. In figure S4, five SMAcre[ER-]:*Grin2d*[f/f] mice were used in the experiment. T-test used for statistical analysis. Each dot indicates the mean value obtained from one mouse, and the data are mean ±s.e.m.

Figure S5. The open field test (OFT) of SMAcre^{ER-}:*Grin2d*$^{f/f}$ (WT) mice and SMAcre^{ER+}:*Grin2d*$^{f/f}$ (cKO) mice. A) Schematic diagram of the box used for the Open Field Test. B) Time of WT and cKO mice spent in the central area of the box. C) Time of WT and cKO mice spent in the non-central area of the box. D) Running distance of WT and cKO mice spent in the center of the box. E) Rate of fecal production for WT and cKO mice. F) Rate of urine production for WT and cKO mice. G) Fecal latency for WT and cKO mice. T-test used for statistical analysis. Each dot indicates one mouse, and the data are mean ±s.e.m.

Figure S6. Blood vessel distribution in the amygdala and negative control of CD-1 fear assay. A) The blood vessel distribution in the amygdala. Red is the signal of mural cells. Green is the signal of NeuN. Blue is the signal of nucleus. B-E) The negative control IF staining of CD-1 fear assay. They are amygdala, pir cortex, S1BF, and hippocampus.

Figure S7. Elevated plus maze (EPM) test of SMAcre^{ER-}:*Grin2d*$^{f/f}$ (WT) mice and SMAcre^{ER+}:*Grin2d*$^{f/f}$ (cKO) mice. A) Schematic diagram and representative diagram of WT and cKO mice in EPM, B) Running distance of WT and cKO mice spent in the EPM. C) Time percentage of WT and cKO mice spent in the central area of the EPM. D) Time percentage of WT and cKO mice spent in close arms of the EPM E) Time percentage of WT and cKO mice spent in open arms of the EPM. F) Ratio of time spent in open and closed arms of WT and cKO mice. Data presented as - log10 format. G) Ratio of time spent in the central and non-central area of WT and cKO mice. Data presented in - log10 format. T-test used for statistical analysis. Each dot indicates one mouse, and the data are mean ±s.e.m.

 Tables

 Table 1. List of Mice Genetics

Common name	Strain name	Resource
Nestin-cre	B6.Cg-Tg(Nes-cre)1Kln/J	Yanmei Tao, Hangzhou Normal University/ Southeast University
Grin2d	B6/JGpt-Grin2d^{em1Cflox}/Gpt	GemPharmatech
SMAcre^{ER}	Tg(Acta2-cre/ERT2)12Pcn	Bo Zhou, ShanghaiInstitutes for Biological Sciences, Chinese Academy of Sciences
Ai47	Gt(ROSA)26Sor$^{tm47(CAG-EGFP*)Hze}$	Zilong Qiu, Chinese Academy of Sciences Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences.
PdgfrβCre^{ER}	Tg(Pdgfrb-cre/ERT2)6096Rha	Xiang Yu, Peking University
Grin1	*Grin1*^{tm2Stl}	Yangang Sun, Chinese Academy of Sciences Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences
Ai14	B6.Cg-Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J	Woo-Ping Ge, Chinese Institute of Brain Research (CIBR)

Table 2. List of Primer Pairs (Some primers referred to the published PrimerBank database)

Name		Sequence
Grin1	F	AAGAACCTGCAGCAGTACCA
	R	GCAGCAGGACTCATCAGTGT
Grin2a	F	TCAGGGACTTTGGAAAACGCT
	R	AAGTCATGCAGGGTTGGTTG
Grin2b	F	ACTAACTATCAATGAAGAACGGT
	R	CCAGACCCCAGAGTAACCA
Grin2c	F	TGTGGGCCTTCTTCGCTGTCATCT
	R	TGCCATTAGGTACCGTGCCAAAAC
Grin2d	F	TGGAGGAGTACGACTGGACAT
	R	CGCACTGACACTACGGAGC
Grin3a	F	CTGCACACCAGTCAGAGGTT
	R	TTCGTTGGTTGTCGTGACTC
Grin3b	F	TGGGGGTCTACCACTTGTCA
	R	TGGATCTTCTGGCTCGTGTG
Ptgs2	F	TTCAACACACTCTATCACTGGC
	R	AGAAGCGTTTGCGGTACTCAT
18s	F	CGGACACGGACAGGATTGACA
	R	CCAGACAAATCGCTCCACCAACTA

 Table 3. List of Antibodies

Name	Host	Cat#	Dilution	Company
Anti-NMDAR2D Antibody	Mouse	MAB5578	1:400 (IF) 1:2,500 (WB) 1:1,000 (IP)	Merck
Anti-Smoothelin antibody	Rabbit	ab219652	1:400 (IF)	abcam
α-Smooth Muscle Actin (ACTA2) Rabbit mAb	Rabbit	A17910	1:2,500 (WB)	ABclonal
Anti-NeuN purified Antibody	Guinea Pig	ABN90P	1:800 (IF)	Merck
Anti-NMDAR1 Antibody, clone 54.1	Mouse	MAB363	1:400 (IF)	Merck
COX2/PTGS2 Rabbit pAb	Rabbit	A1253	1:400 (IF)	ABclonal
c-Fos antibody	Rabbit	226 003	1:1,000 (IF)	SYSY
Recombinant Alexa Fluor® 488 Anti-Caldesmon/CDM antibody	Rabbit	ab208116	1:400 (IF) 1:2,500 (WB)	abcam
Mice IgG	Mouse	A7028	1:1,000 (IP)	Beyotime
Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488	Goat	A-11001	1:1,000 (IF)	Invitrogen
Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 546	Goat	A-11010	1:1,000 (IF)	Invitrogen
Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488	Goat	A-11034	1:1,000 (IF)	Invitrogen
Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 647	Donkey	A-21447	1:1,000 (IF)	Invitrogen
Goat Anti-Guinea pig IgG H&L (Alexa Fluor® 488)	Goat	ab150185	1:1,000 (IF)	abcam
Goat Anti-Guinea pig IgG H&L (Alexa Fluor® 555)	Goat	ab150186	1:1,000 (IF)	abcam
Goat Anti-Guinea pig IgG H&L (Alexa Fluor® 647)	Goat	Ab150187	1:1000 (IF)	abcam
β-Tubulin Mouse mAb	Mouse	AC021	1:5,000 (WB)	ABclonal
Goat Anti-Mouse IgG, HRP Conjugated	Goat	CW0102	1:10,000 (WB)	CWBIO
Goat Anti-Rabbit IgG, HRP Conjugated	Goat	CW0103	1:10,000 (WB)	CWBIO
Goat Anti-Mouse IgG (H+L)	Goat	AS071	1:500 (IF)	ABclonal
DDDDK-Tag Rabbit mAb	Rabbit	AE063	1:5000 (WB)	ABclonal
BD Pharmingen™ Purified Rat Anti-Mouse CD31	Rat	557355	1:400 (IF)	BD
Alexa Fluor® 488 AffiniPure Donkey Anti-Rat IgG (H+L)	Donkeny	712-545-453	1:500 (IF)	Jackson ImmunoResearch

Data availability

All codes involved in this study are available at https://github.com/JialabEleven/Neurovascular_coupling with citation. Another data is available from authors upon reasonable request.

Author contributions

Jie-Min Jia conceived the project, designed the experiments, and supervised this study. Jiayu Ruan did all main experiments, obtained all main original data, and wrote the manuscript. Yiyi Zhang wrote main MATLAB code. Dongdong Zhang provided primer pairs of *Grin1* and provided critical suggestions. Huiqi Xie participated in blood pressure measurement, long-term whisker stimulation, and provided suggestion for manuscript writing. Xinghua Quan participated in all behavioral tests and IF staining of blood vessel distribution in the Amygdala. Wentao Wang participated in the writing of MATLAB code. Xu Hu provided primer pairs of *Grin2a*, *Grin2b*, *Grin2c*, *Grin3a*, *Grin3b*, and *18s*. Yanmei Tao provided Nestin-cre mice. Bingrui Zhao participated in genotyping of all mice and cell sorting. Danping Lu participated in brain slice acquisition.

Figure 1

A

Grin1 *Grin2a* *Grin2b* *Grin2c* *Grin2d* *Grin3a* *Grin3b*

1 2 1 2 1 2 1 2 1 2 1 2 1 2

NMDARs

18s

1: P0 cerebellum
2: SMAcre^(ER+):Ai47 sorted parenchymal aSMCs

C

SMA:Ai14
Grin2d
DAPI

SMCs of arterioles

Neuron-like cells

Grin2d

Tdtomato
Grin2d
DAPI

Grin2d signal

D

NR2D

Tubulin

E

NR2D

Tubulin

F

Grin2d

18s

G

NR2D

Tubulin

1: P0 Nestin-cre^(-):*Grin2d*^(f/f) cerebellum
2: Adult Nestin-cre^(-):*Grin2d*^(f/f) cerebellum
3: P0 Nestin-cre^(+):*Grin2d*^(f/f) cerebellum

4: P10 SMAcre^(ER-):*Grin2d*^(f/f) cerebellum
5: P10 SMAcre^(ER+):*Grin2d*^(f/f) cerebellum

6: SMAcre^(ER+):*Grin2d*^(+/+):Ai47 sorted pial aSMCs
7: SMAcre^(ER+):*Grin2d*^(f/f):Ai47 sorted pial aSMCs

L

8_SMAcre^(ER-):*Grin2d*^(f/f)   9_SMAcre^(ER+):*Grin2d*^(f/f)

Smoothelin
NR2D
DAPI

NR2D

10_SMAcre^(ER+):*Grin2d*^(+/+):Ai47   11_SMAcre^(ER+):*Grin2d*^(f/f):Ai47

EGFP
NR2D

Figure 2

A

TMX administration
(P5 to P8)
(P5_cKO group & littermate control)

Whisker trimming, scalp open
(P63)

Whisker stimulation 4Hz, 1min
(P65)

TMX administration
(P60 to P63)
(P60_cKO group & littermate control)

Rest
(P64)

B

LSIS image diagram

(△f/f)*100% >0%
(△f/f)*100% <0%

	Before stimulation	During stimulation	After stimulation
WT			
P5_cKO			
P60_cKO			

C

Whisker stimulation
(30s-90s)

CBF Variation (%)

— SMAcreER⁻/PdgfrβER⁻:*Grin2d^{f/f}*
— P5 TMX_SMAcre^{ER+}:*Grin2d^{f/f}*
— P60 TMX_SMAcre^{ER+}:*Grin2d^{f/f}*
— P60 TMX_Pdgfrβcre^{ER+}:*Grin2d^{f/f}*

Seconds

D

Plateau mean

Plateau Max

****
****
****
****
****
0.0342
0.0413

CBF Variation(%)

E

Seconds

F

K_{Slope}

0.0001
0.0001
0.0001

Figure 4

Figure 5

Figure 6

Figure 7