Neurovascular Coupling in the Basolateral Amygdala Modulates Negative Emotions

Jiayu Ruan[1, 2, 3, 5], Xinghua Quan[3], Huiqi Xie[3], Yiyi Zhang[3], Dongdong Zhang[2], Wentao Wang[3], Bingrui Zhao[3], Danping Lu[2], Yuxiao Niu[6], Mengyuan Diao[7], Wei Hu[7], Jie-Min Jia[2, 3, 4, 5*]

Affiliations

[1]College of Life Sciences, Zhejiang University, Hangzhou, Zhejiang, China.

[2]Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, China.

[3]Laboratory of Neurovascular Biology, School of Life Sciences, Westlake University, Hangzhou, China.

[4]State Key Laboratory of Gene Expression, School of Life Sciences, Westlake University, Hangzhou, Zhejiang, China.

[5]Laboratory of Neurovascular Biology, Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, China.

[6]Laboratory of Computational and Functional Genomics, School of Life Sciences, Westlake University, Hangzhou, China

[7]Department of Critical Care Medicine, Affiliated Hangzhou First People's Hospital, School of Medicine, Westlake University, Hangzhou, China.

*Correspondence: jiajiemin@westlake.edu.cn

ABSTRACT

Emotion induces changes in regional cerebral blood flow (CBF), a manifestation

of neurovascular coupling (NVC), yet whether NVC might feed back to

modulate emotion actively remains unexplored. Here, we demonstrate that

NVC actively and bidirectionally modulates stress-induced negative emotions.

We established bidirectional manipulations of NVC in freely moving mice by

employing integrated pharmacological, genetic, and arteriolar optogenetic

approaches. Our results showed that both systemic and basolateral amygdala

(BLA) region-specific NVC deficiencies heightened emotional responses when

mice transitioned from a safe, familiar environment to anxiogenic environments,

and local restoration of NVC in the BLA normalized these responses.

Mechanistically, NVC dysfunction impairs the capacity of BLA neuronal scaling

during state transitions, manifesting as a characteristic biphasic pattern of c-

Fos topology. Namely, the NVC-deficient animal aberrantly adopts high-stress

configurations under mild stress but regresses to low-stress templates during

high-demand survival threats, thereby compromising defensive sustainability.

Notably, the genetic NVC-enhancement model counteracts NVC impairments

caused by chronic stress, thereby alleviating stress-driven emotional distress.

These findings established NVC in the BLA as an allostatic program that fine-

tunes neural circuit activity for emotional responses, with implications for

understanding and treating emotional disorders.

INTRODUCTION

Emotion is a complex, integrative response to changes in internal and external environments.[1] Emotional flexibility—the capacity to dynamically regulate and adapt these responses—is fundamental for mental health and survival.[2] Conversely, deficits in this adaptive capacity result in emotional rigidity, a hallmark of dysregulation strongly associated with psychiatric disorders such as depression and post-traumatic stress disorder (PTSD).[3, 4] Therefore, delineating the molecular and cellular mechanisms that govern the balance between emotional flexibility and rigidity is critical for elucidating the biological basis of resilience.[5, 6]

Neurovascular coupling (NVC), the core physiological foundation of blood oxygen level-dependent (BOLD) functional MRI (fMRI) signals, has long been categorized as a passive neural proxy.[7-10] Clinical observations in the basolateral amygdala (BLA)—a pivotal hub for processing negative affect—reveal pervasive neurovascular dysregulation across diverse psychiatric landscapes.[11] Specifically, exaggerated BOLD responses occur in post-traumatic stress disorder (PTSD), bipolar disorder, high body mass index (BMI), and early-life stress-exposed cohorts;[8, 12-15] In contrast, autosomal recessive Urbach-Wiethe disease, stemming from *ECM1* (coding extracellular matrix protein 1) gene mutations, is marked by a lack of hemodynamic responsiveness: Such patients exhibit BLA vascular perturbations leading to localized calcification, alongside a selective deficit in fear perception when confronted

with threat.[16, 17]

Consistent with this clinical phenotype, NVC impairments are recapitulated

in animal models, where acute, chronic, and neuroendocrine stressors

significantly impair NVC fidelity within stress-sensitive circuits.[18-20] Yet, a

fundamental question of causality remains: do these deficits merely reflect

secondary consequences of neuronal dysfunction, or do they actively drive

emotional pathology?

Emerging evidence indicates that vascular-derived cues actively modulate

diverse neural processes. For instance, nitric oxide generated during NVC

promotes adult hippocampal neurogenesis,[21] while vascular-derived nitric oxide

depolarizes the membrane potential of the optic nerve.[22] In the hypothalamus,

salt-loading-induced vasoconstriction during NVC exerts positive feedback to

boost local parvalbumin interneuron activity.[23] Additionally, whole-cell patch

recordings in brain slices demonstrate that altered parenchymal arteriolar tone

modulates nearby neuronal firing–decreasing pyramidal neuron activity and

increasing interneuron activity, with effects depending on the mechanosensitive

cation channel.[24] These observations imply that NVC possesses an intrinsic

capacity, likely via chemical and/or mechanical cues, to actively regulate neural

circuits, extending beyond its traditional role in metabolic support.[25]

However, its potential as a real-time gatekeeper of emotional circuitry

remains unexplored. To address this gap, we investigated NVC within the BLA

as a model system. We hypothesize that BLA NVC functions as a critical

regulatory checkpoint that calibrates the intensity of negative emotions, thereby

maintaining emotional flexibility.[26]

In this study, employing a multidimensional emotion assessment

framework combined with cell-type-specific genetic manipulations and

arteriolar optogenetics, we demonstrate that NVC in the BLA actively and

bidirectionally modulates emotional reactivity. Unlike its non-contribution in the

somatosensory barrel cortex (S1BF), we reveal that neurovascular decoupling

in the BLA precipitates emotional rigidity—a maladaptive state defined by a

collapse in the dynamic range of emotional responses—manifesting as a

distinctive biphasic dysregulation where physiological and behavioral outputs

are pathologically exaggerated under mild stress, yet paradoxically blunted

under intense stress. Mechanistically, we demonstrate that this behavioral

rigidity is underpinned by corresponding biphasic shifts in BLA neural activity

across graded stressors. Notably, the enhanced NVC model confers emotional

resilience against chronic stress by normalizing BLA neural circuit activity.

These findings establish NVC as an intrinsic regulatory gain controller in

emotional processing and suggest that targeting circuit-specific neurovascular

communication offers a novel therapeutic strategy for psychiatric dysregulation.

RESULTS

Validation of a quantitative behavioral framework linking graded BLA activation to escalating spatial stress levels

To dissect the neurovascular mechanisms actively modulating emotion, we first established a robust, quantifiable behavioral paradigm that scales linearly with stress intensity. We exposed male age-matched wild-type (*WT*) mice to contexts for 15 min with graded anxiogenic potential: the home cage (HC; minimal stress), the 40 cm, 35 cm, and 10 cm open field tests (OFTs; mild-to-moderate stress), and acute restraint stress (ARS; intense stress) (Fig. 1a).[27-31] We verified BLA neuronal activation 90 min post-assay by assessing c-Fos expression (Fig. 1b). Immunofluorescence mapping revealed that c-Fos intensity increased in a stress-dependent manner (Fig. 1c). Crucially, this graded BLA activation was mirrored by a stepwise increase in defecation bouts (Fig. 1d), showing a robust linear correlation with BLA c-Fos intensity ($R = 0.8160$, $P < 0.0001$; Fig. 1e). Furthermore, we quantified serum corticosterone levels, the primary stress-responsive glucocorticoid in rodents, in a parallel cohort of mice at 5 min post-assay (Fig.1a). Serum corticosterone concentrations exhibited a stress-dependent trend mirroring that of BLA neural c-Fos (Fig. 1f). Likewise, defecation bouts also showed a strong positive correlation with serum corticosterone concentrations ($R = 0.7765$, $P = 0.0004$; Fig. 1g). Collectively, these results establish defecation as a reliable, robust, and non-invasive marker that accurately tracks systemic and spatially graded

stress levels in free-moving live mice.[31-33]

Reducing the arena size from 40 cm to 35 cm prominently increased

defecation bouts (Fig. 1d). Although the shift in linear dimensions from 40 cm

to 35 cm appears modest, this modification represents a ~33% reduction in total

cubic volume, significantly intensifying the degree of spatial confinement.[34] This

compressed environment provoked a more potent stress state than the 40 cm

OFT, characterized by a multi-dimensional escalation in BLA c-Fos density (Fig.

1c), serum corticosterone levels (Fig. 1f), locomotor activity (Fig. 1h), coupled

with reductions in classic readouts of exploration, specifically center time (Fig.

1i) and unsupported rearing (Fig. 1j). Spatial mapping further revealed that

more than 90% of defecation behavior in the 40 cm OFT occurred in the

peripheral zone, consistent with thigmotaxis behavior (Fig. 1k, l).[27, 35] Together,

converging behavioral, neural, and hormonal readouts confirm that the 35 cm

OFT elicits a stronger stress response than the standard 40 cm OFT.[36]

To determine whether BLA neuronal activity is necessary for this response,

we chemogenetically silenced BLA excitatory neurons using the inhibitory

DREADD hM4Di (Fig. 1m).[37] In mice expressing hM4Di in BLA excitatory

neurons treated with the agonist C21 effectively blunted stress-induced BLA c-

Fos expression in ARS (Fig. 1n, o). Intriguingly, this also suppressed defecation

bouts in both the 40 cm OFT and ARS contexts compared to vehicle-treated

control mice (Fig. 1p). Furthermore, BLA silencing significantly reduced the

defecation rate (the incidence of mice that defecated) and prolonged the

defecation latency (the time elapsed before the production of the first fecal

pellet) (Fig. 1q) in the 40 cm OFT. Interestingly, neither center preference nor

locomotor activity was altered in the 40 cm OFT following chemogenetic

inhibition of the BLA (Fig. 1r-s). Control mice expressing EGFP alone (sham

group) exhibited no alterations in defecation metrics (Fig. 1q, t).

Collectively, these data establish a validated, BLA-dependent quantitative

framework—the "negative emotion indices"—encompassing BLA neural

activation, multi-dimensional defecation metrics (bouts, rate, latency),

unsupported rearing, and serum corticosterone concentration to dynamically

assess emotional states.

A genetic model of cerebral SMC-specific GluN2D deficiency exhibits BLA

neurovascular decoupling

Equipped with this multi-dimensional assessment framework, we next sought

to identify the specific molecular substrates within the BLA neurovascular unit

(NVU) that govern these emotional responses. We previously identified that the

core N-methyl-D-aspartate (NMDA) receptor subunit GluN1 (encoded by *Grin1*)

in smooth muscle cells (SMCs) is essential for NVC at the neuro-smooth

muscle junction (NsMJ; Fig. 2a),[38] but the other regulatory NMDA subunits in

SMCs remained to be characterized. To address this, we screened mRNA

expression of all NMDA receptor subunits in cultured SMCs sorted from

postnatal day 0 (P0) brain parenchymal tissue and identified *Grin2d* (encoding

GluN2D) as the sole regulatory subunit detectably expressed (Fig. 2b). High-

resolution RNAscope further confirmed that *Grin2d* colocalized specifically with

SMC markers in cerebral arterioles with high colocalization coefficients

(visualized via *SMA-creER:Ai14* reporter mice; Fig. 2c). To rigorously validate

the specificity of our RNAscope probes, we leveraged the well-documented

developmental divergence of NMDA subunits in the cerebellum as an internal

biological control. We exploited the distinct trajectories where *Grin1* expression

increases from P0 to P60, while *Grin2d* expression follows a reciprocal

decline.[39, 40] Our probes accurately recapitulated these age-dependent patterns,

thereby confirming their high target specificity (Supplementary information, Fig.

S1a, b). This biological validation was further extended to the protein level; our

anti-GluN2D antibody successfully distinguished the differential GluN2D protein

abundance between P0 and P60 stages in cerebellum tissue, mirroring the

transcriptomic data (Supplementary information, Fig. S1c).

Using this validated antibody, immunofluorescence analysis of isolated

middle cerebral arteries (MCA) (Fig. 2d, e) and cultured SMCs sorted from P0

brain parenchymal tissue (Supplementary information, Fig. S1d) confirmed the

specific ablation of GluN2D in the cerebral SMCs in *SMA-creER:Grin2d^{f/f}*

(*Grin2d*-cKO) mice. Furthermore, Western blot analysis of whole-cortical

lysates revealed that *Grin2d*-cKO did not affect GluN2D expression in other

non-SMC populations (Supplementary information, Fig. S1e), confirming the

cell-type specificity of our genetic model. To assess the global impact of this

genetic depletion, we examined GluN2D expression in the SMCs of peripheral

organs, including the heart, liver, spleen, lung, kidney, intestine, and bladder

(Supplementary information, Fig. S2a). Notably, GluN2D expression seemed

restricted to cerebral SMCs and absent in these peripheral tissues' SMCs,

suggesting that our *Grin2d*-cKO model exerts a brain-restricted impact

(Supplementary information, Fig. S2b, c). Furthermore, we systematically

verified the physiological status of *Grin2d*-cKO mice; no differences were

observed in systemic metabolic homeostasis (Supplementary information, Fig.

S3a-g), blood pressure (Supplementary information, Fig. S3h), gastrointestinal

function (Supplementary information, Fig. S3i-n), and blood-brain barrier (BBB)

integrity[41, 42] (Supplementary information, Fig. S3o-p) compared to littermate

controls. Finally, we assessed key contractile proteins, finding that α-SMA and

smoothelin expression remained unaltered (Supplementary information, Fig.

S4). These data demonstrate that the basic metabolic functions,

gastrointestinal system, and basic contractile structural integrity of SMCs are

preserved in *Grin2d*-cKO mice.

To assess the functional impact of SMC-specific *Grin2d* deletion and

facilitate a direct, one-to-one comparison with our previously reported *Grin1*-

cKO model, we employed the same classical paradigm: measuring whisker

stimulation-evoked hemodynamic responses in the somatosensory barrel

cortex (S1BF) (Fig. 2f).[38] Consistently, *Grin2d*-cKO mice exhibited a ~66%

attenuation in functional hyperemia upon whisker stimulation (Fig. 2g),

confirming that GluN2D is functionally comparable to GluN1 in regulating

cerebral NVC. Control experiments confirmed that tamoxifen administration

alone in *Grin2d^{f/f}* mice did not alter NVC responses (Supplementary information,

Fig. S5).

Having established the NVC deficit in the cortex, we sought to determine

whether this NVC dysfunction extends to the BLA. To this end, we developed a

de novo system based on previous strategies[43, 44] to simultaneously record

neuronal activity and blood supply within the BLA of freely moving mice, thereby

enabling direct assessment of NVC. First, we mapped the BLA arteriolar

topology by superimposing 3D-reconstructed arterioles in the tissue-cleared

brain on a schematic cerebral vasculature. This analysis identified that the

corticoamygdaloid artery (coamg) penetrates the BLA, making it suitable for

photometric monitoring (Supplementary Information, Fig. S6, Video S1).

Second, we validated the fidelity of our cerebral blood volume (CBV) recording

method: fiber photometry-based blood plasma fluorescence intensity recording

locally in the BLA showed a robust linear correlation with orbital Rhodamine

injection concentration (Supplementary information, Fig. S7a, b; $R = 0.8946$, P

< 0.0001). Signal stability was achieved 15 min post-injection (Supplementary

information, Fig. S7c), and this time point was designated as the initiation time

for the behavior assay.

Using this optimized paradigm, after tamoxifen administration, we injected

AAV2/9-hSyn-GCaMP8f into the BLA and implanted optic fibers at the injection site two weeks before the recordings. After three days of habituation (one trial per day for a continuous 3-day period), we monitored NVC during the 40 cm OFT (Fig. 2h-k). Consistent with our earlier c-Fos data (Fig. 1c), the 40 cm OFT elicited robust BLA Ca^{2+} transients that were temporally matched by CBV elevations, indicating tight local NVC in littermate control mice (Fig. 2i-k). In contrast, *Grin2d-cKO* mice exhibited a dissociated neurovascular phenotype, where augmented BLA neuronal Ca^{2+} signals were paradoxically paired with a blunted hemodynamic response (Fig. 2i-k). This inverse pattern—hyperactive neurons coupled with hypoperfused blood supply—was further corroborated by increased c-Fos expression in the BLA 90 min post-assay (Fig. 2l, m). Furthermore, we confirmed that the observed attenuation in functional hyperemia was not attributable to differences in basal perfusion, locomotor activity, or spatial preference within the 40 cm OFT (Supplementary information, Fig. S8)

Collectively, these findings demonstrate robust NVC within the BLA during 40 cm OFT exposure and validate that SMC-specific *Grin2d*-cKO mice exhibit an NVC deficit in this region.

BLA-target pharmacological restoration of NVC reversed emotional hyper-reactivity in *Grin2d-cKO* mice under mild stress

Building on the BLA's established role as one of the central hubs for stress

responses, we next utilized our negative emotion indices (Fig. 1) to evaluate

whether the BLA neurovascular decoupling observed in *Grin2d*-cKO mice (Fig.

2) results in behavioral and physiological impairments. Notably, we observed

that *Grin2d*-cKO mice exhibited a ~12-fold increase in defecation bouts during

the first 5 min of the 40 cm OFT, a trend sustained throughout the session (Fig.

3a). This phenomenon was not observed in the home cage context

(Supplementary information, Fig. S3c-e, n). Littermate control mice (*Grin2d^{f/f}*)

displayed a graded increase in defecation bouts in response to escalating

spatial stress, consistent with the pattern observed in *WT* mice (Figs. 1d and

3b). In contrast, *Grin2d*-cKO mice failed to show this graded response; instead,

they exhibited an abrupt rise to a maximal plateau level of defecation bouts, a

response typically elicited by intense ARS stress (Fig. 3b). This hyper-reactivity

was further reflected in an increased defecation rate, reduced latency (Fig. 3c),

a more than 3-fold reduction in unsupported rearing, and a ~1.3-fold increase

in serum corticosterone levels in the 40 cm OFT (Fig. 3d). Importantly, basal

stress levels in the home cage remained indistinguishable between genotypes

(Supplementary information, Fig. S9a). Altogether, *Grin2d*-cKO mice displayed

a unified shift across multiple-dimensional stress metrics, indicating a robust,

exaggerated negative emotional response to mild stressors, such as those

encountered in OFTs.

Given that *Grin2d*-cKO mice showed NVC dysfunction beyond the BLA, to

determine the contribution of BLA local NVC in hyper-reactivity, we employed

administration of HET0016, a selective inhibitor of CYP450ω-hydroxylases that

suppresses the synthesis of 20-HETE (a potent vasoconstrictor), thereby

facilitating vasodilation to compensate for the NVC deficit.[45, 46]

To strictly validate the pharmacological efficacy of HET0016 before its site-

specific delivery to the BLA, we first replicated the systemic administration

protocol described in our previous study.[38] Systemic HET0016 administration

dose-dependently reduced ARS-induced defecation bouts (Supplementary

Information, Fig. S10a). This behavioral rescue provided a clear rationale for

selecting 1 mg/kg as the optimal dose for subsequent investigations. Under this

condition, we confirmed that 1 mg/kg HET0016 produced a ~3-fold increase in

whisker-evoked functional hyperemia amplitude in the S1BF compared with

saline controls (Supplementary Information, Fig. S10b, c)

Notably, HET0016 effectively rescued the hyper-reactive phenotype in

Grin2d-cKO mice, both in systemic (Fig. 3e-j) and local (Fig. 3k-p)

administrations in the 40 cm OFT. Both administration routes effectively

normalized BLA neural activation and multi-dimensional defecation metrics—

including bouts, rate, and latency—to littermate control levels (Fig. 3f-i and l-o).

Importantly, this vascular intervention also significantly normalized unsupported

rearing bouts and reduced serum corticosterone concentrations (Fig. 3j and p)

in the 40 cm OFT. In contrast, in the home cage, neither systemic nor BLA local

HET0016 impacted serum corticosterone levels (Supplementary information,

Fig. S9b, c).

Collectively, local pharmacological compensation in the BLA sufficiently

normalized the hyper-reactivity features in the *Grin2d*-cKO genetic model,

despite NVC deficits in other brain regions, thereby establishing that intact NVC

in the BLA is essential for maintaining appropriate emotional responses to mild

stress.

We observed consistent behavioral phenotypes in the 40 cm OFT across

various mechanistically distinct models of NVC dysfunction, each targeting

different nodes of the NVU (Supplementary information, Fig. S11, S12).

Specifically, either perturbing upstream neural signaling (*Emx1-cre:Ptgs2^{f/f}*)

(Supplementary information Fig. S11a-h, and S12o-p)[47-49] or disrupting the

neurovascular interface–achieved via pan-mural cell-specific *Grin2d*-cKO

(*Pdgfrβ-creER:Grin2d^{f/f}*) or SMC-specific *Grin1-Grin2d* double-cKO (*SMA-*

creER:Grin1^{f/f}:Grin2d^{f/f}) (Supplementary information Fig. S11i-r and S12o-p)–

consistently recapitulated the NVC deficits and behavioral phenotypes

observed in the primary *Grin2d*-cKO mice in the 40 cm OFT (Fig. 2, 3).

To ultimately disturb contractile protein machinery within SMC, we targeted

the *Acta2* gene (encoding α-SMA).[50, 51] This model (*Pdgfrβ-creER:Acta2^{f/f}* mice,

shorten as *Acta2*-cKO) indistinguishably mirrored the hyper-reactive

phenotypes of the signaling-deficient models in the 40 cm OFT (Supplementary

information, Fig. S11s-v, x-aii, and S12o, p). Surprisingly, a high-fat diet

(HFD)—known to diminish NVC in both animals and humans via multiple

factors[14, 52]—produced an identical phenotype of *Grin2d*-cKO mice in the 40 cm

OFT (Supplementary information, Fig. S12a-d, g-h, j-n, o, and p). These

findings indicate that this emotional hyper-reactivity was not restricted to the

single *Grin2d*-cKO model but represented a universal consequence of

neurovascular dysfunction within the 40 cm OFT, reinforcing the universality of

NVC-driven emotional dysregulation.

Although local HET0016 delivery to the BLA successfully restored

emotional metrics in other models (Fig. 3k-p, Supplementary information, Fig.

S12i-n) in the 40 cm OFT, it failed to produce similar effects in *Acta2*-cKO mice

(Supplementary information, Fig. S11w-aii) in the same context. This indicates

that once the downstream contractile apparatus is disrupted, upstream

regulators are unable to exert their regulatory functions. Finally, we confirmed

that these convergent effects did not involve BBB leakage (Supplementary

information, Fig. S3o, p; S12e, f).[53] Together, these findings establish that NVC

integrity per se is indispensable for emotional homeostasis.

Caldesmon acts as the downstream effector of GluN2D in NVC thereby

modulating emotional reactivity

Given that direct disruption of the vascular contractile apparatus (*Acta2*-cKO)

phenocopies the emotional hyper-reactivity of *Grin2d*-cKO mice (Fig. 3;

Supplementary information, Fig. S11; Fig. S12), we hypothesized that GluN2D

couples to the SMC cytoskeletal machinery. To this end, we performed co-

immunoprecipitation (Co-IP) followed by mass spectrometry to map the

GluN2D interactome in vascular SMCs. This unbiased screen identified

caldesmon (encoded by *Cald1*), an actin-binding protein critical for regulating

actomyosin contractility, as a top candidate enriched in *Vascular Smooth*

Muscle Contraction pathway (Fig. 4a).[54, 55] This physical association between

GluN2D and caldesmon was further confirmed by Western blotting of the

immunoprecipitated complex (Fig. 4b).

Caldesmon serves as a "blocker" on smooth muscle cells' contraction; its

disruption is predicted to impair cross-bridge cycling and thereby compromise

NVC. To test this, we employed siRNA-mediated knockdown (KD). Unlike

GluN2D, caldesmon expression was restricted exclusively to vascular mural

cells (Fig. 4c–e),[56] allowing for vascular-specific manipulation without the risk

of direct neuronal off-target effects. We validated the efficacy and spatial

specificity of this approach: both whole-brain and local BLA siRNA delivery

robustly reduced caldesmon expression in background-matched groups (both

established in *Grin2d^{f/f}* mice) (Fig. 4f-i). Crucially, gene silencing was spatially

restricted to the injection site with anatomical precision: local BLA *Cald1* KD did

not impact caldesmon levels in the adjacent Ventromedial Hypothalamus (VMH)

(Fig. 4g–i), nor did S1BF-restricted *Cald1* KD alter expression in the BLA

(Supplementary information, Fig. S13).

Functionally, local S1BF *Cald1* KD—but not local BLA *Cald1* KD—

attenuated whisker-evoked functional hyperemia in the barrel cortex (Fig. 4j, k),

confirming *Cald1* KD as an effective tool for disrupting NVC with regional

precision. We then focused on the emotional circuit. Behaviorally, both whole-

brain and local BLA *Cald1* KD recapitulated the hyper-reactive phenotype of

Grin2d-cKO mice in the 40 cm OFT: *Cald1* KD mice exhibited exaggerated

negative emotional responses, characterized by BLA neuronal hyperactivation

(c-Fos), reduced unsupported rearing, elevated serum corticosterone

concentrations, and intensified defecation dynamics (increased bouts/rate,

decreased latency) (Fig. 4l–q). In contrast, local S1BF *Cald1* KD did not alter

the above parameters, confirming regional specificity (Supplementary

information, Fig. S14).

To elucidate the functional relationship between GluN2D and caldesmon,

we applied the local BLA *Cald1* KD approach in *Grin2d*-cKO mice. This

epistasis analysis showed no additive phenotype (Fig. 4l-q), suggesting that

caldesmon likely functions downstream of GluN2D within a hierarchical

vascular signaling cascade. Together, these results identify caldesmon as a

critical effector linking GluN2D to SMC contractility and demonstrate that

localized NVC deficits in the BLA–despite intact NVC in other brain regions–are

sufficient to impair emotional responses to mild stress.

BLA arteriolar SMCs' optogenetic suppression of NVC precipitates negative emotional states under conditions of mild and negligible stress

To achieve acute suppression of NVC within a millisecond timescale–unattainable through genetic or pharmacological manipulations–we employed an SMC-optogenetic approach to reversibly "clamp" vascular diameter.[57]

We generated SMC-specific ChR2-expressing mice (*SMA-creER:Ai32;* Fig. 5a) and optimized a fiber-optic setup to locally activate BLA SMCs while simultaneously monitoring regional CBV (Fig. 5b). In the home cage under minimal external stress, we optimized a protocol that reliably induced a consistent ~3% CBV decrease (Fig. 5c red line). This hemodynamic manipulation was rapid and reversible (Fig. 5d).

Consistent with earlier data obtained from *Grin2d*$^{f/f}$ mice (blue line in Fig. 2i, blue line), the stress during the 40 cm OFT elicited functional hyperemia in an independent cohort of *SMA-creER:Ai47* (EYFP) mice (Fig. 5e-g, green line). Optogenetic clamping of BLA arterioles partially blocked this hemodynamic response (Fig. 5e–g, red line). Furthermore, power spectral density analysis of the hemodynamic traces revealed a significant reduction in blood flow power across low-frequency bands—typically associated with vascular vasomotion—during optogenetic stimulation (Supplementary information, Fig. S15). This suppression of spectral energy confirms that optogenetic clamping effectively stabilized vascular diameter.

This acute NVC suppression precipitated an emotional shift: ChR2-EYFP

mice exhibited an exacerbation of negative emotional responses, characterized

by markedly increased BLA neural activation (Fig. 5h, i), reduced unsupported

rearing (Fig. 5j), and elevated serum corticosterone concentrations (Fig. 5k),

and shifted defecation metrics (Fig. 5l, m) in the 40 cm OFT. These alterations

were consistent with the *Grin2d*-cKO phenotype (Fig. 3a-d) in the 40 cm OFT.

Locomotor activity and center preference remained unaffected, ruling out motor

confounds (Supplementary information, Fig. S16a, b) in the 40 cm OFT. This

optogenetically driven, temporally controlled suppression of NVC reveals that

preventing the stress-induced vascular surge removes a critical physiological

component, thereby amplifying negative emotion.

Next, we investigated whether BLA-targeted vascular clamp, in the

absence of actual external stress, is sufficient to drive stress-like behaviors. We

applied the same optogenetic stimulation protocol (~3% CBV decrease) to mice

in the safe home cage environment for 15 min (Fig. 5n, o). Strikingly, this

hemodynamic mismatch alone was sufficient to induce a stress-like state: While

basal defecation was negligible in the home cage (0.23 pellets/15 min),

optogenetic stimulation induced a ~6.4-fold cumulative CBV change (Fig. 5o)

coupled with a ~10-fold surge in defecation bouts (Fig. 5p). These changes

were associated with increased BLA neural activation and elevated serum

corticosterone levels (Fig. 5q, r).

Crucially, we verified that this optogenetic manipulation did not induce

hypoxic injury. Unlike the Stroke Induced by Magnetic ParticLEs (SIMPLE)

stroke model[58-60] which triggers robust HIF-1α upregulation, our optogenetic

protocol elicited no obvious HIF-1α response after optogenetic stimulation in

the home cage (Supplementary information, Fig. S16c, d). This confirms that

the observed behavioral shifts stem from functional neurovascular modulation,

not ischemic stroke or tissue hypoxia.

437 NVC dysfunction conversely exhibited hyporeactivity under high-

438 intensity stress

Our findings from the standard 40 cm OFT established that NVC dysfunction

precipitates a hyper-reactive, anxiety-like state under mild stress (Fig. 3). We

next characterized how the NVC-deficient BLA responds to imminent, life-

threatening high-intensity stress.[61] To this end, we employed two paradigms: a

looming assay[62] and a virtual reality (VR$_{eagle}$) simulation of a diving eagle[63], both

of which mimic an approaching aerial predator.

Prior to looming stimulation, basal locomotor activity was comparable

between groups (Fig. 6a, blue region; Supplementary information, Fig. S17a).

Upon stimulus, both groups exhibited immediate and robust freezing (Fig. 6a,

yellow region), confirming that NVC dysfunction did not impair sensory

detection, threat evaluation, or the initiation of defensive strategies.[64]

However, a striking divergence emerged during the maintenance phase:

while control mice sustained a prolonged freezing state, *Grin2d*-cKO mice

exhibited a "premature exit" phenotype (Fig. 6b). Velocity heatmaps confirmed

that *Grin2d*-cKO mice resumed locomotor activity significantly earlier than

controls (Fig. 6b, c). This behavioral collapse was mirrored by an inappropriate

increase in unsupported rearing bouts—a high-risk behavior exposing the

animal to recurring predation—immediately following stimulus termination (Fig.

6d). This dissociation (intact initiation but shortened maintenance) in free-

moving mice underscores the critical role of NVC in sustaining defensive states.

The second life-threatening paradigm, virtual reality (VR$_{eagle}$) eagle,

allowed for head-fixed monitoring of pupil diameter in awake mice—a robust

index of autonomic arousal[65]—alongside locomotor activity on a float ball (Fig.

6e). Baseline metrics (pupil diameter and locomotor activity) were

comparable between genotypes (Supplementary information, Fig. S17b, c).

Upon exposure to the eagle call (0–1.5 s, Figure 6f blue region) and visual

dive (1.5–3 s, Figure 6f yellow region), both groups exhibited robust defensive

engagement, characterized by immediate freezing and sharp pupil dilation

sustained throughout the stimulus (0–3 s, Fig. 6f; Supplementary information,

Fig. S17d and Video S2). This confirms that sensory processing and fear

initiation remain preserved in *Grin2d*-cKO mice, consistent with observations

made in the looming assay.

However, a striking divergence emerged during the "defensive offset

phase", defined as the interval from stimulus termination to the intersection

of pupil diameter and locomotion speed in tracing plots for the recovery period

(Fig. 6f, double-arrows lines). While littermate control mice maintained high

arousal, *Grin2d*-cKO mice exhibited a rapid collapse of the defensive state

(Fig. 6g). Moreover, the recovery time for pupil diameter back to baseline was

reduced by ~59% in *Grin2d*-cKO mice, consistent with the accelerated

recovery time of locomotion speed observed in the looming assay (Fig. 6h).

Collectively, these two independent paradigms demonstrate that NVC-

deficient mice cannot physiologically sustain high-load fear states.

To establish an intermediate stress condition between the OFTs and a

life-threatening predator scenario (looming and VR_{eagle}), we modified the

standard OFT to create a complex OFT (c-OFT). This modification involved

the addition of a metal fence barrier that separated an aversive start zone

(white floor) from a safe zone (black floor) (Fig. 6i).[29] c-Fos mapping of

littermate controls revealed that c-OFT paradigm indeed represented an

intermediate level of BLA activation, which fell between the mild stress elicited

by the OFTs and the intense stress induced by predator simulation (Fig. 6j,

blue line). In contrast, c-Fos mapping in *Grin2d*-cKO mice demonstrated

exaggerated neural activation under mild stress yet blunted activation under

intense stress, matching their corresponding behavioral phenotypes (Fig. 6j,

magenta line).

Notably, under the intermediate c-OFT condition, one-dimensional

analysis of c-Fos density failed to correlate with the observed heighted

defecation bouts in *Grin2d*-cKO mice (Fig. 6k). This discrepancy prompted us

to extend beyond simple intensity quantification and conduct further analyses

of the topological distribution patterns of c-Fos positive neurons within the

BLA (Fig. 6l-m; supplementary information, S18).

We therefore decoded c-Fos topology using the structural similarity index

measure (SSIM) and unbiased clustering to dissect BLA ensemble

configurations (Fig. 6l, m). In littermate controls, BLA topology exhibited

dynamic "stress scaling": activation patterns evolved distinctly from minimal

(Level 0) to high-threat states (Level 3), forming a precise structural hierarchy

(Fig. 6l). In contrast, *Grin2d*-cKO ensembles displayed topological

disorganization and a lack of hierarchy (Fig. 6m).

Mapping *Grin2d*-cKO neural SSIM signatures against the structural

hierarchy of littermate control mice revealed a striking inversion of emotional

scaling (Supplementary information, Fig. S18b-g): Under mild stress (OFTs),

Grin2d-cKO patterns aberrantly clustered with high-threat control templates

(Levels 2 and 3), providing a topological explanation for their hyper-reactivity

(Supplementary information, Fig. S18d, e). Conversely, under high load

(looming, VR_{eagle}), these patterns failed to scale appropriately and instead

regressed to low-stress templates (Level 1), underpinning their functional

collapse (Supplementary information, Fig. S18f, g). Specifically, hierarchical

clustering revealed that the *Grin2d*-cKO topological signature under c-OFT

aberrantly converged with that of control mice under ARS, rather than their

stage-matched controls. This pathological shift was absent in the home cage,

where topologies of both genotypes remained tightly clustered

(Supplementary information, Fig. S18b, c). These findings reveal that BLA

NVC deficiency decouples behavior from c-Fos intensity by inducing a

nuanced reconfiguration of BLA neural topology.

Collectively, these data reveal a stress load-dependent biphasic pattern

(Supplementary information, Fig. S18h): NVC dysfunction precipitates

topological hyper-reactivity under low stress load (OFTs) but leads to hypo-

sustainability under high stress load (looming and VR$_{eagle}$), demonstrating

that functional NVC is indispensable for the topological scaling of neural

networks during state transitions.

Genetic NVC-enhancement model of *Glrb*-cKO mice

Our preceding findings established that the cation-permeable NMDA receptor

(GluN2D) acts as the essential "accelerator" for NVC. Its deficiency disrupts

NVC, locking the brain into a state of behavioral rigidity—manifested as hyper-

reactivity under mild stress yet functional collapse under high load contexts.

Guided by the fundamental principle of electrochemical balance, we

hypothesized that if cation influx drives NVC, a converse mechanism mediated

by anion channels might functionally restrict hemodynamic gain. Consequently,

inhibiting such anionic "brakes" could theoretically potentiate NVC and confer

emotional resilience.

To identify these regulators, we screened transcriptomic databases for

anion channels enriched in mural cells.[56] This unbiased screen identified *Glrb*

(encoding the chloride-permeable glycine receptor subunit beta, GlyRβ) as the

top candidate (Supplementary information, Fig. S19)[56]. RNAscope and

immunofluorescence mapping revealed a distinct spatial segregation: *Glrb*

(GlyRβ) was predominantly expressed in venous SMCs (vSMCs) (Fig. 7a–d).

We validated this expression pattern using *Pdgfrβ-CreER:Glrb^{f/f}* (*Glrb*-cKO)

mice, which exhibited a significant reduction in mural *Glrb* (GlyRβ) levels (Fig.

7c, e, f). To determine whether these venous receptors are functionally

innervated, we examined their molecular and ultrastructural organization:

Molecularly, GlyRβ puncta closely colocalized with gephyrin (Fig. 7g), a

scaffolding protein essential for clustering inhibitory receptors,[66] suggesting the

presence of postsynaptic-like machinery; Structurally, confocal imaging and

serial electron microscopy (EM) revealed frequent direct physical contacts

between neuronal axons and venular SMCs (Fig. 7h, i). Collectively, these

molecular and ultrastructural features provide the morphological substrate for

potential neuro-venular communication sites. Unlike the NVC deficit caused by

Grin2d ablation, *Glrb* deletion resulted in a "super-coupled" state: *Glrb*-cKO

mice exhibited significantly enhanced functional hyperemia upon whisker

stimulation (Fig. 7j-m). To pinpoint the cellular origin of this enhancement, we

performed a genetic dissection. First, deletion restricted to arteriolar SMCs

(driven by *SMA-creER*) failed to recapitulate the phenotype (Supplementary information, Fig. S20), ruling out an arteriolar origin. Second, given that GlyRβ is selectively enriched in venular SMCs—and considering the established view that capillaries nearly lack the contractile machinery for active dilation[50]—the enhanced NVC in *Pdgfrβ* mutants is primarily attributable to venular disinhibition. Consequently, the enhanced NVC in *Glrb* mutants suggests that venular GlyRβ may physiologically constrain the magnitude of hemodynamic responses.

Given that pharmacological vasodilation (HET0016) successfully rescued the *Grin2d*-cKO phenotype, we hypothesized that genetic removal of *Glrb* could serve as a genetic rescue strategy to counterbalance the *Grin2d* deficit. To test this genetic disinhibition hypothesis, we generated double-knockout mice (*Pdgfrβ-CreER:Grin2d^{f/f}:Glrb^{f/f}*; Fig. 7k). Remarkably, the concomitant deletion of *Glrb* effectively reversed the NVC deficiency triggered by *Grin2d* loss. Whisker stimulation in *Pdgfrβ-CreER:Grin2d^{f/f}:Glrb^{f/f}* mice elicited functional hyperemia responses comparable to littermate controls (Fig. 7l, m). This result implied that *Glrb*-cKO mimics the effect of HET0016. This finding establishes a genetic paradigm of hemodynamic rescue, demonstrating that venular disinhibition can effectively counterbalance the loss of the arteriolar accelerator. We next asked the fundamental translational question: Does this enhancement of neurovascular coupling confer behavioral resilience against stress?

NVC enhancement confers resilience against both acute and chronic stress

Given that *Glrb* deletion unlocks a "super-coupled" hemodynamic state (Fig. 7), we first asked whether this vascular enhancement translates into immediate behavioral resilience against acute stress. We subjected mice to ARS. While littermate controls exhibited characteristic stress-induced physiological and behavioral shifts, *Glrb*-cKO mice appeared remarkably stress-buffered. They displayed significantly attenuated BLA neuronal activation (Fig. 8a, b) and blunted serum corticosterone concentrations (Fig. 8c) in ARS. Behaviorally, this physiological resilience was mirrored by stabilized defecation dynamics, with *Glrb*-cKO mice exhibiting significantly fewer defecation bouts in ARS (Fig. 8d).

The observation that *Glrb* removal in mural cells confers stress resilience prompted us to investigate whether this NVC-enhancement model could counteract the deficits caused by *Grin2d*-cKO. We found that *Pdgfrβ-CreER:Grin2d$^{f/f}$:Glrb$^{f/f}$* mice effectively neutralized the core hyper-anxious phenotype observed in *Grin2d-cKO* mutants (Fig. 8d–j). Specifically, *Pdgfrβ-CreER:Grin2d$^{f/f}$:Glrb$^{f/f}$* mice exhibited defecation metrics comparable to NVC-intact littermate controls in ARS and 40 cm OFT (Fig. 8d-f), accompanied by attenuated BLA neural activation (Fig. 8g, h) and a significant increase in unsupported rearing bouts (Fig. 8i) in the 40 cm OFT. Collectively, the observation that *Pdgfrβ-CreER:Grin2d$^{f/f}$:Glrb$^{f/f}$* mice closely resemble NVC-intact mice indicates that emotional reactivity is dynamically calibrated by the

precise homeostatic balance between opposing vasoconstrictive and vasodilatory signaling pathways within the BLA.

To determine whether the stress resilience observed in systemic mural cell-specific *Glrb*-cKO mice is driven by localized NVC within the BLA, we targeted the disruption of BLA NVC via local delivery of *Cald1* siRNA. In intense ARS, local BLA NVC disruption abolished the resilience-conferring effects typically seen in the *Glrb*-cKO mice, effectively reverting neural, behavioral, and hormonal stress responses to littermate control levels (Supplementary information, Fig. S21a-d). Furthermore, laser speckle contrast imaging (LSCI) revealed that whole-brain knockdown of *Cald1* via cisterna magna delivery did not alter basal CBF at the cortical surface (Supplementary information, Fig. S21e). This indicates that *Cald1* deficiency did not compromise basal vascular integrity, but specifically impairs the vascular response to neuronal activation.

While *Glrb* deletion effectively buffers the immediate impact of acute stress, a more rigorous test of neurovascular integrity is whether this protection can sustain itself against chronic adversity. We thus escalated our paradigm to chronic restraint stress (CRS, 3 hours/day for 28 days) (Fig. 8k).[19, 67] We first validated the efficacy of the CRS model in littermate controls, which exhibited a comprehensive state of emotional and physiological exhaustion. Specifically, CRS-treated littermate control mice exhibited significantly attenuated functional hyperemia in S1BF during whisker stimulation (Fig. 7l, m).[19] Concurrently, these mice developed a distinct constipation-like phenotype, characterized by

progressive reductions in both defecation frequency and Bristol scores over the 28-day stress period (Fig. 8l, m). This was accompanied by classical anxiety- and depression-like behaviors, including significantly reduced time in the open arms of the elevated plus maze (EPM; Supplementary information, Fig. S22a), heightened neural activation in the lateral habenula (LHb; Supplementary information, Fig. S22b, c)—a critical node in depression-related circuitry[68]—and prolonged immobility in the forced swim test after the CRS modelling (FST; Fig. 8n). Furthermore, we verified that CRS exposure did not compromise BBB integrity (Supplementary information, Fig. S23). This confirms that the observed impairments in NVC and subsequent emotional and behavioral dysregulation after CRS were not confounded by non-specific vascular leakage or barrier disruption.[42, 69]

Under this sustained adversity, *Glrb*-cKO mice displayed profound resilience. Specifically, they preserved intact whisker-evoked functional hyperemia and remained resistant to the stress-induced constipation-like phenotype, maintaining normal defecation dynamics throughout the 28-day CRS modelling (Fig. 7l, m and Fig. 8l, m). Furthermore, *Glrb*-cKO mice displayed high mobility in the FST, independent of CRS exposure (Fig. 8n). At the cellular and endocrine levels, *Glrb*-cKO mice were protected from the maladaptive sensitization of stress circuits, showing no significant increase in baseline BLA neural activity and serum corticosterone concentrations compared with CRS-treated littermate controls at the end of CRS (Fig. 8o, p).

Importantly, *Glrb*-cKO mice exhibited normal locomotion and spatial

preference in the 40 cm OFT (Supplementary information, Fig. S22d, e), ruling

out hyperactivity or altered exploratory drive as confounding factors for the

observed behavioral improvements. Finally, it is important to note that *Glrb*-cKO

mediated protection may brain-region specific. While emotional and

neurovascular homeostasis was preserved, *Glrb*-cKO mice still experienced

stress-induced weight loss comparable to CRS-treated littermate controls (Fig.

8q). This suggested that while enhanced NVC effectively shields the brain's

emotional circuits, it did not negate the systemic physiological costs of chronic

stress.

DISCUSSION

In this study, we demonstrate that bidirectional control of NVC enables

bidirectional modulations of the negative emotion states. Far from being a

passive metabolic read-out of neuronal activity, NVC within the BLA–but not in

the primary somatosensory cortex–operates as an indispensable active

regulator of the affective negative emotion network. By combining genetic,

pharmacological, and optogenetic approaches, our findings reveal that the

vascular interface acts as a gatekeeper, tuning stress signals to either be

amplified into anxiety-like under mild stress or dampened into resilience to

strong stress. These findings redefine the BLA NVC as a proactive component

of affective circuitry.

More than a century after the first measurements of regional CBF laid the

foundation for modern NVC research, the field has come to a new door–what

next?[70] We knock on this door by beginning with addressing the role of BLA

NVC in emotion modulation, offering a conceptual framework for exploring the

regulatory capacity of NVC across functionally distinct brain regions and their

respective outputs.

During optogenetic manipulations, the ~3% hemodynamic changes we

detected (Fig. 5n) is highly consistent with recent work by Hauglund et al.[57]

showing that analogous ~3% CBV changes sufficiently drive cerebrospinal fluid

flow. This concordance across two separate research groups verifies that subtle,

persistent shifts in basal blood volume can be sufficient to elicit notable

functional effects, for instance, elevated neural activity in the BLA (Fig. 5h, i, q,

r left).

Our arteriolar optogenetic data provide genuine *in vivo* validation that the

neurovascular unit (NVU) acts as a critical regulator of proximal neural activity

(Fig. 5e-m). Even under the minimal-stress home cage context with negligible

external stress, arteriole contraction potently activated neighboring neurons

(Fig.5q, r left), defining an *in vivo* form of vasculo-neuronal coupling (VNC) that

mirrors earlier *ex vivo* observations.[24] These findings imply an implication that

NVC and VNC operate as two arms of a fast, closed loop, enabling real-time,

bidirectional crosstalk between vascular and neuronal compartments.

NVC is a widely recognized, evolutionarily conserved biological process.[71]

The "premature exit" observed in NVC-deficient mice under predator-like stress

(Fig. 6) suggests that NVC is innately encoded to optimize the kinetics

underlying the predatory fear response cycle. As a survival-critical program, this

regulatory property may explain the evolutionary conservation of NVC across

species.

The specific mechanism by which $Glrb$ deficiency enhances NVC warrants

further investigation. In mural cells, the intracellular chloride concentration (Cl^-

$_i$) is typically higher than the extracellular concentration (Cl^-_e), creating an

electrochemical gradient where the opening of chloride-permeable glycine

receptors leads to Cl^- outflow—a sharp contrast to the Cl^- influx observed in

mature neurons.[72-75] This efflux is predicted to depolarize mural cells and

promote contraction; thus, the ablation of $Glrb$ likely diminishes the baseline

contractile ability of mural cells. Given that $Glrb$ (GlyRβ) is selectively enriched

in venular SMCs (vSMCs) (Fig. 7a–d) and that its deletion in arteriolar SMCs

(aSMCs) fails to recapitulate the "super-coupled" phenotype (Supplementary

information, Fig. S20a, b), we propose that $Glrb$ (GlyRβ) functions as a

physiological constraint on venular conductance. Upon its removal, neural

inputs—such as those during whisker stimulation—elicit a significantly

amplified NVC response (Fig. 7l, m).

While *Grin2d* and *Glrb* appear to exert their primary effects on distinct vascular segments—the arteriolar and venular compartments, respectively—their mutual counteraction suggests a complex hemodynamic synergy (Fig. 7l, m). One plausible interpretation is that vSMCs also possess a regulatory weight within the arteriole-capillary-venule network that is functionally comparable to that of aSMCs. These findings suggest that NVC operates as an integrated network property in the vasculature system, where enhanced venular hemodynamics functionally offsets arteriolar deficits to maintain an apparent partial equilibrium (Figures 7 and 8).

At the molecular level, we propose that GluN2D activates a signaling cascade in arteriolar SMCs—specifically, a GluN2D–Ca^{2+}-bound calmodulin–caldesmon–α-SMA axis. Caldesmon typically binds to α-SMA filaments and sterically blocks their interaction with myosin, thereby suppressing cross-bridge formation.[55] Ca^{2+}-bound calmodulin, generated following Ca^{2+} entry through GluN2D-containing NMDARs, is known to relieve this inhibition.[55] Consequently, GluN2D activation is predicted to release the caldesmon "brake", modulating α-SMA–myosin cross-bridge cycling to increase CBF. Although this pathway warrants further direct validation, our prediction is supported by genetic epistasis analysis: *Cald1* knockdown effectively phenocopies the *Grin2d*-cKO phenotypes (Fig. 4l-q), and the lack of an additive effect when combining these two perturbations (*Grin2d*-cKO+local BLA *Cald1* KD) suggests that they potentially operate within the same axis (Fig. 4l-q).

This study has two primary limitations. First, the current lack of a venular

SMCs-specific *CreER* line necessitated the use of the pan-mural *Pdgfrb-CreER*

driver to perturb *Glrb* in arteriolar SMCs, pericytes, and venular SMCs. Second,

we utilized male cohorts to minimize hormonal variability; future investigations

involving female mice are warranted to establish the sex-dependent

generalizability of these neurovascular mechanisms.

In conclusion, we identified that NVC functions as an active physiological

gatekeeper rather than a passive support system. This study establishes NVC

as a key target for basic research into neural circuits and mood disorders.

MATERIALS AND METHODS

Animal care and subject selection

All animal procedures complied with the Institutional Animal Care and Use

Committee (IACUC) guidelines of the School of Life Sciences, Westlake

University (Protocol 24013JJM). Mice were housed in the Westlake University

Laboratory Animal Resource Centre under strictly controlled conditions: a 12-h

light/dark cycle at 25 °C, with maximum six mice per individually ventilated cage,

and *ad libitum* access to standard chow and water. To exclude potential

confounding effects from the estrous cycle on emotional behavior, we

exclusively used age-matched male C57BL/6J mice (60–80 days old at the

onset of experiments).[30]

Rigor and bias control for behavior-related assay

We implemented rigorous protocols to minimize behavioral variability arising from habituation or handling. To ensure stable baseline conditions, animals were first acclimated to the facility for at least two weeks and then habituated to the testing room environment for at least 90 min prior to each specific assay. All handling and behavioral assays were performed by a single trained experimenter who was strictly blinded to genotype and group assignments; blinding was maintained by a separate researcher who managed genotyping results. Furthermore, testing was confined to a fixed daily time window (13:00–19:00). To prevent exposure to uncontrolled olfactory or auditory cues, testing areas were strictly controlled for light, noise, and odor: apparatuses were thoroughly cleaned with 70% ethanol and ventilated between trials, and animals were transported in standardized containers. Finally, animals were randomly distributed across experimental groups from different cages to control for potential cage effects.[30, 76, 77]

Home cage non-invasive physiological monitoring

To assess physiological parameters with minimal disturbance, mice were monitored in their home cages. Blood pressure was measured using a non-invasive tail-cuff system (Kent/Biopa), and the occupation of fat was measured

using EchoMRI^{TM}-100H (EchoMRI) during the fixed experimental time window

(13:00–19:00); crucially, mice were habituated to the restrainer for three

consecutive days prior to recording to minimize stress artifacts.

For baseline metabolic and excretory profiling, mice were housed in groups

of four. Over three consecutive 3-day cycles, animals were provided with

standardized quantities of food (60 g), water (340 g), and bedding (100 g). At

the end of each cycle, body weight, ingestive behaviors (food and water intake),

fecal output, and urination volume (derived from bedding weight gain) were

recorded. For home-cage baseline assessments where individual excreta could

not be definitively assigned in group-housed conditions, data were averaged

per mouse within each cage. To ensure statistical rigor and avoid

pseudoreplication, the cage mean was utilized as the independent

experimental unit (N = 4 cages). To facilitate cross-group comparisons, weight

data were normalized to the littermate control baseline. Specifically, the value

of each sample was divided by the mean value of the control group (defined as

100%). To record defecation bouts, mice were individually housed in new home

cages for 3 days of habituation. Existing fecal pellets were cleared 15 min prior

to the assay, and the number of defecation bouts was recorded during the

subsequent 15-min interval. Following monitoring, mice were randomly

assigned to terminal procedures, including gastrointestinal inspection, retro-

orbital blood collection for ELISA, blood-brain barrier (BBB) integrity

assessment, and brain extraction for c-Fos immunofluorescence mapping. All

assessors remained blinded to group identity throughout data collection and

analysis.

Behavioral assays

We performed all behavioral assays using a rigorous double-blind protocol. One

trained experimenter conducted the assays, while a second, independent

experimenter performed the video scoring and data analysis. To eliminate

olfactory cues and ensure consistency, we cleaned all apparatus with 70%

ethanol and fully aerated them between subjects.

Open field tests (OFTs)

To assess emotional responses under graded stress intensities, the OFT was

conducted in square arenas of three specific dimensions: standard (40 × 40

cm), intermediate (35 × 35 cm), and constricted (10 × 10 cm). Mice were placed

in the center of the arena, and behavior was recorded for 15 min using an

overhead camera. Unsupported rearing—defined as standing on hindlegs

without contacting the arena walls—was quantified during the first 5 min.

Defecation metrics (bouts, rate, and latency), running distance, and area

preference were measured across the entire 15-min session. For zonal analysis,

the arena floor was digitally divided into 25 equal squares (5 × 5 grid); the center

zone was defined as the inner 9 squares, excluding the 16 peripheral squares

adjacent to the walls.

After testing, mice were transferred to clean, individual cages. Retro-orbital

blood samples for ELISA analysis were collected 5 min post-assay. Mice

designated for immunofluorescence were perfused 90 min after the assay to

capture peak c-Fos expression. For linear correlation analyses, behavioral and

physiological data were paired from the same individual subjects.

To adhere to the 3Rs principles (Replacement, Reduction, and Refinement)

and minimize animal usage, 40 cm OFT data (including whisker stimulation

data) for *SMA-creER:Grin1f/f* mice were obtained through the retrospective re-

analysis of unpublished datasets independently generated in our laboratory.

This approach enhances the reproducibility of our findings, as these data were

collected by a different experimenter at a distinct time point using identical

experimental parameters. Due to the nature of the original experimental

recordings, which were restricted to behavioral videos and whisker-evoked

hemodynamic measurements, c-Fos and serum corticosterone data were not

available for the *SMA-creER:Grin1f/f* line.

For fiber photometry and optogenetic-based OFT, only 40 cm OFT used.

Tamoxifen induction 21 days prior to formal testing. For real-time NVC

monitoring, *AAV2/9-hSyn-GCaMP8f* was stereotaxically injected into the BLA

(AP: −1.6 mm; ML: ±3.45 mm; DV: −4.00 mm), and optic fibers were implanted

immediately above the injection site (DV: −3.95 mm). Following a two-week

recovery and three days of fiber-connection habituation, mice received a retro-

orbital injection of Rhodamine B (Sigma, R9379; 100 µl, 5 mg/ml for formal

assay) 15 min prior to the 40 cm OFT. The experimental timeline comprised

three sequential recording phases: pre-test (home cage), test (40 cm OFT), and

post-test (home cage).

To rigorously isolate genuine functional signals from hemodynamic and

motion-related artifacts, we employed a multi-wavelength fiber photometry

system (Thinkertech) for all fiber photometry assays referred published

methods.[78, 79] Fluorescence signals were acquired via 473 nm excitation for

GCaMP8f (Ca^{2+}-dependent) and 580 nm for Rhodamine (blood volume). A 405

nm channel was utilized as an isosbestic reference; at this wavelength,

GCaMP8f fluorescence remains Ca^{2+}-insensitive, serving as a precise proxy for

non-specific absorption and motion artifacts that affect both the green and red

emission spectra. To decontaminate the functional data, the 405 nm reference

trace was independently fitted to the 473 nm and 580 nm raw signals using

least-squares linear regression. These fitted components were then subtracted

from their respective raw traces. This dual-correction strategy effectively

neutralized common-mode noise and ensured that the reported $\triangle$F/F for both

GCaMP8f and Rhodamine reflects true physiological dynamics—neuronal

activity and functional blood volume changes, respectively—undistorted by

physical movement or non-functional optical fluctuations. These data

processing and normalization protocols were applied consistently to

optogenetic experiments to ensure cross-experimental comparability (details

and formula shown below).

For optogenetic manipulation of vascular contractility,[43, 44, 57] optic fibers

were implanted in *SMA-creER:Ai32* (ChR2-EYFP) and control *SMA-*

creER:Ai47 (EYFP) mice using the coordinates described above (without

AAV2/9-hSyn-GCaMP8f injection). Mice received tamoxifen induction 21 days

prior to testing. Following habituation and Rhodamine injection, blue light pulses

(473 nm) were delivered to induce optogenetic stimulation while simultaneously

monitoring cerebral blood volume (CBV) via Rhodamine fluorescence (580 nm).

To capture the full hemodynamic response profile, recordings were segmented

into three phases: pre-stimulation baseline, optical stimulation, and post-

stimulation recovery. Optogenetic stimulation cycle consisted of an 85 ms

stimulation pulse followed by a 5 ms interval and a 5 ms recording window (10

Hz, 0.939 mW laser power).

As for isosbestic correction for hemodynamic artifacts, similar to BLA

photometry recording, signal fluctuations at the 405 nm isosbestic point are

calcium-independent and primarily reflect changes in light scattering and

absorption by dynamic hemoglobin concentration changes (functional

hyperemia). We applied a linear regression correction algorithm. As mentioned

above, the 405 nm signal was fitted to the 473 nm signal, and the predicted

hemodynamic/motion component was subtracted. This correction confirms that

the CBV recording during the optogenetic stimulation represents a genuine

hemodynamics.

Two distinct normalization strategies were applied to quantify CBV

dynamics:

1.   Static Baseline Comparison (Cross-group): To allow for cross-

group comparisons of basal vascular hemodynamics, data were normalized

to the mean raw value of the control group. Specifically, the raw value of

each individual sample was divided by the mean raw value of the littermate

control group (defined as 100%). This approach standardizes the readout,

ensuring that reported values reflect specific genotypic differences in

perfusion capacity relative to a physiological baseline.

2.   Dynamic Response Quantification (Time-series): For time-series

recordings of CBV (as well as Ca^{2+} signal recording and Laser Speckle

Contrast Imaging recording), a temporal normalization strategy was applied.

The baseline signal ($F_{baseline}$) was defined as the mean value recorded

during the defined pre-stimulation period (e.g., the 30-second baseline

window). The relative change at each time point (F_t) was calculated as a

percentage of its own baseline using the formula:

900             $\triangle CBV\ (\%)\ or\ \triangle Ca^{2+}\ (\%) = (F_t - F_{baseline}) / F_{baseline} \times 100\%$

This self-normalization approach isolates the evoked hemodynamic

response from background variations. This peak-aligned averaging strategy

ensures that the quantification of the metabolic plateau is temporally

synchronized to the individual hemodynamic kinetics of each animal.

Fourier transform plots were generated in MATLAB, with frequency and

power spectral density represented on the X and Y axes, respectively. The

analysis was conducted using the acquisition parameters specified in the 40

cm OFT optogenetic stimulation section. To resolve the spectral characteristics

of the BLA hemodynamic response, Fast Fourier Transform (FFT)[80-82] was

applied to the CBV time series. For visualization, the low-frequency range (0–

1 Hz) is presented in the representative spectral panel. For statistical

quantification, the Area Under the Curve (AUC) was calculated specifically for

the ultra-low frequency band (0–0.3 Hz) to represent oscillatory power across

experimental groups.[83, 84] The relevant codes can be found at the GitHub

website:

https://github.com/JialabEleven/Functional_hyperemia_regulates_emotion-

master/tree/master/Part7_Frequency_analysis.

Home cage optogenetic stimulation and recording

To determine if BLA vasoconstriction is sufficient to induce negative emotion in

a non-aversive environment, simultaneous optogenetics and photometry were

performed in the home cage. *SMA-creER:Ai32* (ChR2-EYFP) and control *SMA-*

creER:Ai47 (EYFP) mice received tamoxifen induction 21 days prior to testing.

Optic fibers were implanted above the BLA (AP: −1.6 mm; ML: ±3.45 mm; DV:

−3.95 mm) two weeks before the assay, and mice were habituated to the recording tether for three days. The experimental procedures for optogenetic stimulation in the home cage were identical to those conducted in the 40 cm OFT, with the exception of the testing environment.

On the test day, Rhodamine B (100 μL, 5 mg/mL) was administered via retro-orbital injection 15 min prior to recording. The session comprised three 15-min epochs: baseline, stimulation, and recovery. To ensure signal fidelity and prevent spectral crosstalk, an optimized interleaved illumination sequence was employed: (1) 405 nm for isosbestic control; (2) 580 nm for continuous CBV recording; and (3) 473 nm for optogenetic stimulation (delivered exclusively during the second epoch). The duty cycle consisted of an 85 ms stimulation pulse followed by a 5 ms interval and a 5 ms recording window (10 Hz, 0.939 mW laser power). To standardize baseline conditions, all existing fecal pellets were removed immediately before recording, and fresh fecal output was quantified at the end of the session. Data analysis and processing for optogenetics in the home cage is similar to that in the 40 cm OFT.

Tube stress assay

Restraint stress was induced using modified 50 mL conical tubes (Corning, Cat# 430290).[85, 86] To prevent hypoxia and hyperthermia, tubes were equipped with multiple ventilation apertures drilled into the cap and body, ensuring

adequate airflow while strictly restricting physical movement.

For the acute restraint stress (ARS) paradigm, mice were restrained for a

single 15-min session, and fecal pellet output was quantified immediately upon

release. To interrogate the causal role of BLA activity in stress responses, we

utilized a chemogenetic BLA neural silencing strategy. *AAV2/5-mCaMKIIa-*

hM4D(Gi)-EGFP-ER2-WPRE-pA (inhibitory DREADD) or the control virus

AAV2/5-mCaMKIIa-EGFP-ER2-WPRE-pA was bilaterally injected into the BLA

(AP: −1.6 mm; ML: ±3.45 mm; DV: −4.0 mm). Following a two-week recovery

period, mice were subjected to behavioral testing (including the 40 cm OFT

described above). To ensure peak DREADD activation during the stressor, the

agonist Compound 21 (C21; MCE, Cat# HY-100234; 0.1 mg/kg, i.p.) was

administered 45 min prior to initiating restraint. Following ARS, mice were

individually housed in clean cages. Retro-orbital blood samples for ELISA

analysis were collected 5 min post-assay. To capture stress-induced neuronal

activation, mice were perfused 90 min post-assay for c-Fos

immunofluorescence mapping.

For the chronic restraint stress (CRS) paradigm, mice were subjected to daily

restraint sessions for 28 consecutive days. To prevent habituation to the

stressor, both the duration (randomized between 120 and 180 min) and the

daily onset time (within the light phase) were varied. The physiological impact

of chronic stress was assessed weekly (at the end of each 7-day block) by

recording fecal output and stool consistency (Bristol Score)[87] during a 60-min

monitoring period in a clean home cage. Bristol scores analysis was conducted

by an independent researcher blinded to the mouse genotypes and treatment

groups. Mice designated for Elevated Plus-Maze (EPM) Test, Forced Swim Test

(FST), Blood-brain-barrier (BBB) integrity assessment, ELISA inspection, and

immunofluorescence analysis were sacrificed 24 hours after the final CRS

session.

Complex-open field test

To simulate an environment of high visual contrast and frustrated escape, the

complex OFT (c-OFT) was conducted in a specialized chamber divided into two

compartments: a high-illumination white zone (aversive) and a low-illumination

black zone.[29, 88] Crucially, these zones were separated by a barred metal

partition that permitted visual contact with the safe environment but precluded

physical entry. Mice were placed in the center of the white compartment at trial

onset, and behavioral responses to this context of unattainable safety were

continuously recorded for 20 min. Defecation bouts inspection after the assay.

To capture stress-induced neuronal activation, mice were perfused 90 min post-

assay for c-Fos immunofluorescence mapping.

Looming assay

Innate defensive behaviors against aerial predators were assessed by adapting

established protocols,[62] with specific optimizations implemented to standardize

threat perception. Mice were first acclimated to the arena for 10 min to stabilize

baseline anxiety levels. To ensure that defensive responses (freezing or flight)

were distinct from baseline immobility or thigmotaxis, a state-dependent

triggering protocol was employed. Unlike continuous stimulation paradigms, the

looming stimulus—a rapidly expanding overhead black disk—was manually

triggered only when the mouse met specific behavioral criteria: spontaneous

locomotion (speed 13–15 cm/s) and positioning within the arena center (away

from the walls). Behavioral responses were recorded using a high-resolution

overhead camera. Each mouse underwent a single stimulation trial to prevent

habituation. Immediately following the assay, mice were housed individually in

clean cages for 90 min prior to perfusion for c-Fos analysis.

1001

Virtual reality eagle (VR$_{eagle}$) assay

To simulate an aerial predator attack within a highly controlled, multimodal

environment, we employed a custom-built virtual reality (VR) flight simulation

system. Assay design modified from previous work.[62, 63] The apparatus

consisted of an air-supported spherical treadmill (float ball) allowing

unrestricted locomotion, surrounded by a panoramic display system to create

an immersive visual field. Two weeks prior to testing, mice were implanted with

titanium headplates. To strictly decouple restraint stress from predator-induced

responses, animals underwent a rigorous daily habituation protocol (one week) to the head-fixation apparatus and the floating ball mechanism until they demonstrated stable voluntary locomotion. On the testing day, mice were head-fixed and allowed a 5-min adaptation period within a neutral visual landscape. To ensure the capture of active defensive responses, a state-dependent triggering protocol was enforced: the predator stimulus was triggered only after the mouse initiated spontaneous locomotion following the adaptation phase. The stimulus consisted of a high-fidelity eagle avatar diving rapidly from the dorsal to the proximal visual field (3 s duration), temporally synchronized with an eagle call (audiovisual integration, ~75 dB at first 1.5 s). Pupil diameter was tracked via an infrared camera, and locomotion speed was recorded via rotary encoders sampling the spherical treadmill rotation. Data acquisition continued for 2 min post-stimulation to assess recovery kinetics. Each mouse underwent a single stimulation trial to prevent habituation. Immediately following the assay, mice were housed individually in clean cages for 90 min prior to perfusion for c-Fos analysis.

Elevated Plus-Maze (EPM) Test

To assess anxiety-like behavior, the EPM was conducted using an apparatus elevated 50 cm above the floor, consisting of two open arms (65 × 5 cm) and two closed arms (65 × 5 cm) intersecting at a central platform.[89, 90] At the trial

1031   onset, mice were placed in the center zone facing an open arm and allowed to

1032   explore freely for 5 min. The duration of time spent in the open arms was

1033   recorded.

1034

Forced Swim Test (FST)

To evaluate behavioral despair, the FST was performed in a transparent glass

cylinder filled with water (25 °C) to a depth sufficient to preclude caudal

support.[91] Behavior was recorded for a 5-min session. The duration of

immobility—defined as floating with the absence of active struggling, exhibiting

only minimal movements necessary to maintain the head above water—was

quantified. Immediately post-assay, mice were dried and returned to warmed

home cages.

1043

Enzyme-linked immunosorbent assay (ELISA)

To assess physiological stress levels, blood samples were allowed to clot at

controlled room temperature (22–25 °C) for 2 hours. Serum was isolated by

centrifugation at 3,000 rpm for 20 min at 4 °C, followed by a secondary

clarification spin at 8,000 rpm for 3 min to remove residual debris.

Corticosterone concentrations were quantified using a commercial ELISA kit

(Absin, Cat# abs554164-96T) strictly following the manufacturer's protocol.

Optical density (OD) was measured using a Cytation 5 microplate reader

(Agilent, Model MS-SP104).

1053

Stereotaxic viral transduction

Mice were anesthetized with isoflurane and secured in a stereotaxic frame. To

minimize bias, surgical procedures were performed by investigators who were

blinded to the group allocations. For manipulations targeting the BLA, viral

vectors were microinjected at the following coordinates relative to Bregma: AP:

−1.6 mm; ML: ±3.45 mm; DV: −4.0 mm. For chemogenetic inhibition, mice

received bilateral injections (150 nL per side) of the neural inhibitory DREADD

virus *AAV2/5-mCaMKIIa-hM4D(Gi)-EGFP-ER2-WPRE-pA* (2.3×10^{12} VG/mL)

or the control virus *AAV2/5-mCaMKIIa-EGFP-ER2-WPRE-pA* (2.3×10^{12}

VG/mL).[37] For neuronal calcium recording: The calcium indicator *AAV2/9-hSyn-*

jGCaMP8f-WPRE-pA (2.0×10^{12} VG/mL) was injected bilaterally (150 nL per

side) using the same coordinates. To label glycinergic neurons in the medullary

reticular formation, *rAAV2/4-Glyt2-EGFP-WPRE-hGHpolyA* (5×10^{12} VG/mL)

was injected bilaterally (500 nL per side) at coordinates: AP: −3.5 mm; ML:

±0.84 mm; DV: −2.8 mm.[92]

Following surgery, mice were allowed to recover for at least two weeks to

ensure optimal viral expression and physiological recovery prior to behavioral

testing or brain collection.

1072

Immunofluorescence staining and microscopy

To preserve tissue integrity and antigenicity, mice were deeply anesthetized with 1% pentobarbital sodium (80 mg/kg, i.p.) and transcardially perfused with ice-cold PBS (5 min) followed by 4% paraformaldehyde (PFA) in PBS (5 min). Brains and peripheral organs were harvested, post-fixed in 4% PFA for 6 hours at 4 °C, and sectioned at 50 μm using a vibratome (Leica Histo-VT1200S). Cultured cells were washed with PBS and subsequently fixed in 4% PFA for 10 min at room temperature.

For immunolabeling, free-floating sections were permeabilized and blocked in PBS containing 1% Triton X-100 and 0.25% normal donkey serum (NDS) for 60 min at room temperature. Following three washes in PBS, sections were incubated overnight at 4 °C with primary antibodies in a buffer containing 0.1% Triton X-100. After extensive washing, sections were incubated with species-specific fluorescence-conjugated secondary antibodies (in 0.1% Triton X-100) for 2 hours at room temperature. Nuclei were counterstained with DAPI (1:1000; Beyotime, Cat# C1002). Sections were mounted using anti-fade medium (SouthernBiotech, Cat# 0100-01) and imaged using a Zeiss LSM800 confocal microscope.

To explicitly quantify the spatial distribution of proteins relative to the vasculature (e.g., GlyRβ distribution), orthogonal linescan profiling was performed. Regions of interests (ROIs) were selected where vessels appeared

in longitudinal cross-section (brain slice) or appropriate places in the boundary

of cultured Human Umbilical Vein Smooth Muscle Cells. Line profiles (width: 5

pixels) were drawn to capture the fluorescence intensity gradient. Besides, to

validate the robustness of these spatial associations and control for selection

bias, Pearson's Correlation Coefficient (PCC) was calculated across the entire

ROI using the ImageJ Coloc 2 plugin.[93] This global metric confirmed the degree

of pixel-by-pixel co-localization between the target protein and the vascular

marker, independent of specific linescan trajectories.

As for cellular quantification and normalization, distinct quantification and

normalization strategies were applied based on the specific biological metrics:

1.   Cellular Intensity (e.g., c-Fos): The total number of NeuN-positive

nuclei was identified within the defined ROI. Preliminary analysis confirmed

that the density of NeuN-positive neurons was consistent across groups,

validating the use of area-based normalization. Activation intensity was

calculated by dividing the positive neural count by the total area of the ROI,

expressed as c-Fos positive neurons per mm^2. Additionally, to facilitate

comparison with alternative quantification methodologies, the proportion of

c-Fos-positive neurons relative to the total neuronal population within the

BLA and LHb was also calculated. These supplementary data are provided

in the supplementary information (Supplementary information, Fig. S24).

2.   Relative Protein Expression (e.g., GluN2D, HIF-1α, and IgG): Mean

Fluorescence Intensity (MFI) was measured specifically within the somatic

ROIs of NeuN-positive neurons. To facilitate cross-group comparisons, data

were normalized to the littermate control baseline. Specifically, the MFI of

each sample was divided by the mean MFI of the littermate control group

(defined as 100%).

3.   Signal colocalization (e.g., COX-2): Background fluorescence in

the COX-2 channel (546 nm) was first subtracted across the field of view.

The colocalization between the COX-2 signal and the NeuN signal (488 nm)

within NeuN-positive regions was then quantified using the Pearson

correlation coefficient (PCC, Coloc 2 plugin in ImageJ software).

1125

Pharmacological interventions

To induce Cre-mediated recombination, Tamoxifen (MCE, Cat# HY-13757A)

was dissolved in corn oil (Aladdin, Cat# C116023) at a concentration of 10

mg/mL. The solution was administered via oral gavage (200 µL daily). Dosing

schedules were tailored to specific genotypes: *Acta2*-cKO mice were treated

from postnatal day 21 (P21) to P24, whereas all other transgenic lines received

treatment from P60 to P63, unless otherwise specified.

For acute pharmacological manipulations, two distinct delivery routes were

employed:

1.   Systemic Administration: Drugs were delivered via intraperitoneal

(i.p.) injection at a standardized volume of 200 µL per mouse. Behavioral

assays commenced 15 min post-infusion to target the peak efficacy window.

2.   Local BLA injection: To target the BLA specifically, a stainless-steel

guide cannula (RWD, Cat# 800-00245-00) was stereotaxically implanted at

coordinates AP: −1.6 mm, ML: ±3.45 mm, DV: −4.0 mm. Drugs were

microinfused (1 µL total volume) at a controlled rate of 250 nL/min using an

automated syringe pump. HET0016 (10 µM) was microinfused locally into

the BLA. Behavioral assays commenced 5 min post-infusion to target the

peak efficacy window.

1145

Cell isolation and culture

To characterize SMC-specific molecular mechanisms, primary cultures were

established by the SMCs sorted from P0 brain parenchymal tissue of *SMA-*

creER:Ai47 mice. Cortical parenchymal tissue was dissected following

hypothermia-induced anesthesia and digested in 0.25% trypsin (Thermo

Scientific, Cat# 25200072) for 15 min at 37 °C. Digestion was terminated with

fetal bovine serum (FBS; Gibco, Cat# 10099). The suspension was passed

through a 40 µm cell strainer (Falcon, Cat# 352340) and centrifuged at 1,600 ×

g for 5 min at 4 °C. Pelleted cells were resuspended in Smooth Muscle Cell

Medium (SMCM; ScienCell, Cat# 1101) and plated in T25 flasks (Corning, Cat#

430639).

To purify SMC lineages, primary cultures (~80%) were treated with

Tamoxifen (5 µM in DMSO; Solarbio, Cat# D8371) to induce EGFP expression

specifically in α-SMA-positive cells. Four days post-induction, EGFP-positive

cells were purified using an MA-900 cell sorter (Sony, FCCF-MBR). Sorted cells

were maintained in SMCM for downstream assays. The purity of SMCs isolated

via this protocol has been previously validated.[94]

To capture the *in vivo* transcriptional profile of mural cells (e.g., validation

of *Glrb* expression), freshly isolated cells were used rather than cultured

populations. Pdgfrβ-positve cells were sorted directly from dissociated whole-

brain single-cell suspensions for P60 mice using an anti-Pdgfrβ antibody and

immediately lysed in TRIzol reagent to minimize transcriptional alterations. For

the assay in Fig. 7g, Human Umbilical Vein Smooth Muscle Cells (HU-vSMC;

ScienCell, Cat# 8020) were cultured under standard conditions recommended

by the manufacturer.

1171

Semi-quantitative reverse transcription PCR (RT-PCR)

Total RNA was extracted from two distinct sources using TRIzol reagent

(Sangon Biotec, Cat# B511311-0100): (1) Cultured Cells: Sorted EGFP-positive

SMCs harvested at ~90% confluence; and (2) Brain Tissue: Cortices or

cerebella dissected from P60/P0 mice under 1% pentobarbital sodium (80

mg/kg, i.p.)-induced (P60) or hypothermia-induced (P0) anesthesia.

First-strand cDNA was synthesized (50 ng/µL) using the HiScript III 1st

Strand cDNA Synthesis Kit (Vazyme, Cat# R312-02). For transcript abundance

analysis, semi-quantitative PCR amplification was performed using Green Taq

Mix (Vazyme, Cat# P131-AA). The reaction mixture contained 10 µL Taq Mix, 2

µL cDNA, 2 µL primer pairs (10 µM), and 6 µL RNase-free water. PCR

amplicons were resolved on 2% agarose gels (BIOWEST, Cat# 111860) via

electrophoresis (220 V for 30 min) and imaged using a Tanon 1600 system.

To quantify relative mRNA abundance, the mean grey value of

electrophoretogram bands was quantified using ImageJ software: Band

intensities were first normalized to the corresponding internal loading control

(e.g., *18s*) to correct for variations in lane-to-lane loading. Then, to determine

relative expression levels across groups, these normalized values were

subsequently divided by the mean value of the littermate control group (defined

as 100%).

RNAscope *in situ* hybridization

To validate cell-type-specific gene expression, 15 µm horizontal brain sections

were prepared using a Leica CM1950 cryostat. Genetically modified lines were

utilized to map specific targets: *SMA-creER:Ai14* mice for *Grin2d* expression in

arteriolar SMCs, and *Pdgfrβ-creER:Ai14* mice for *Glrb* in mural cells; *WT* mice

were used for probe specificity validation (*Grin1* and *Grin2d*). Multiplex

fluorescent *in situ* hybridization was performed using the RNAscope Multiplex

Fluorescent Reagent Kit v2 (ACDbio, Cat# 323100) strictly adhering to the

published works and manufacturer's guidelines.[38, 89] Specific probes included

Grin2d (Cat# 425951-C3), *Grin1* (Cat# 431611), and *Glrb* (Cat# 1225261-C1).

Spatial quantification, including orthogonal linescan profiling and co-localization

assessment, was performed identically to the immunofluorescence protocol

described above. Briefly, regions of interests (ROIs) were selected where

vessels appeared in longitudinal cross-section. Line profiles (width: 5 pixels)

were drawn to capture the fluorescence intensity gradient. Besides, to validate

the robustness of these spatial associations and control for selection bias,

Pearson's Correlation Coefficient (PCC) was calculated across the entire ROI

using the ImageJ Coloc 2 plugin.

1211

Western blot

Protein extraction was performed on two distinct sample types: (1) Brain Tissue:

Dissected regions were homogenized in ice-cold RIPA buffer (Thermo Scientific,

Cat# 89901) supplemented with a protease inhibitor cocktail (CWBIO, Cat#

CW22008); and (2) Cultured Cells: Cells were lysed in the same buffer at ~90%

confluence. Lysates were centrifuged at 12,000 × g for 5 min at 4 °C to remove

insoluble debris. Protein concentrations were determined using a Micro BCA

Protein Assay Kit (CWBIO, Cat# CW2011S). Samples were denatured by

heating at 100 °C for 10 min in SDS-PAGE loading buffer. Proteins were

resolved on 8% SDS-polyacrylamide gels and transferred onto 0.45 μm PVDF

membranes (Millipore, Cat# IPVH00010). Following blocking with 5% non-fat

milk in TBST, membranes were incubated overnight at 4 °C with primary

antibodies, washed, and subsequently incubated with HRP-conjugated

secondary antibodies for 45 min at room temperature. Immunoreactive bands

were visualized using ECL reagents (Abbkine, Cat# BMU102-CN) and captured

on a GE AI680RGB imaging system.

To quantify relative protein abundance, the mean grey value of

immunoreactive bands was quantified using ImageJ software. A rigorous dual-

normalization strategy was applied identically to the RT-PCR protocol: Band

intensities were first normalized to the corresponding internal loading control

(e.g., GAPDH) to correct for variations in lane-to-lane loading. Then, to

determine relative expression levels across groups, these normalized values

were subsequently divided by the mean value of the littermate control group

(defined as 100%).

1236

1237 **siRNA administration**

1238 To achieve broad cerebrovascular knockdown; siRNA was delivered via

1239 injection into the cisterna magna following established protocols.[38] A mixture

1240 containing 0.8 nmol siRNA and FITC-dextran (10 mg/mL; Sigma) was prepared

1241 to allow for visual tracking of the distribution (20 μL total volume, infused 1

ul/min). 10 min post-injection, successful delivery was rigorously verified by visualizing perivascular FITC fluorescence through the open scalp using a Zeiss Axio Zoom V16 microscope. Crucially, only mice exhibiting confirmed widespread perivascular perfusion were included in downstream analyses. Functional outcomes (Whisker Stimulation and 40 cm OFT) were assessed 48 hours post-injection to align with the window of peak knockdown efficiency.

For local BLA and S1BF knockdown, siRNA solution (1 nM, 150 nL) was microinjected bilaterally directly into the BLA at coordinates: AP: −1.55 mm; ML: ±3.40 mm; DV: −4.0 mm (BLA)/ DV: −0.5 mm (S1BF). The 40 cm OFT and ARS was conducted 48 hours post-injection.

Whisker stimulation and Laser Speckle Contrast Imaging (LSCI)

To assess functional hyperemia, mice were anaesthetized with 1% sodium pentobarbital (80 mg/kg, i.p.). Assay design referred published paper.[38, 48, 60] Following anesthesia, the head was secured in a stereotaxic frame, and the scalp was retracted to expose the skull. We stimulated whiskers longitudinally using a motor-driven brush (12 Hz, 1 min) and recorded CBF dynamics in the contralateral barrel cortex using a laser speckle contrast imaging system (RFLSI ZW/RFSLI III, RWD Life Science).[95] Images were acquired at a resolution of 512 x 512 pixels with an exposure time 5 ms, optimized for 10 frames per second (fps) sampling.

The LSCI system provides data in "Perfusion Units (PU)", which represent

a relative blood flow index derived from the spatial blurring of laser speckle

patterns. This index is related to the product of average speed and

concentration of moving red blood cells in the tissue sample volume. While

these raw PU values were used to compare basal perfusion levels between

groups, functional hyperemia data were normalized to correct for inter-

individual variations in cranial optical properties. Specifically, the CBF response

was calculated as the percentage change relative to baseline:

1271 $\triangle CBF\ (\%) = (CBF_t - CBF_{baseline})\ /\ CBF_{baseline} \times 100\%$

Where CBF_t is defined as the perfusion value of each frame; $CBF_{baseline}$ is

defined as the mean perfusion value during the 30-second pre-stimulation

period.

To characterize the temporal dynamics of functional hyperemia, the peak

response time was identified using a custom MATLAB algorithm (Signal

Processing Toolbox). Specifically, local maxima in the $\triangle CBF$ time-series were

detected using the findpeaks function, with empirical constraints applied to

reject signal noise (minimum peak height threshold > 2.2%; minimum peak

separation > 50 frames). The global maximum detected within the stimulation

window was defined as the peak functional hyperemia. To quantify the

magnitude of the sustained vascular response, we calculated the mean $\triangle CBF$

over a period from the identified peak latency to 90 seconds (which marks the

conclusion of the whisker stimulation).

Tissue clearing and 3D vascular reconstruction

To visualize the 3D vascular architecture, mouse brains were washed in PBS (3 × 2 hours) on a shaker (40 rpm) at room temperature. Tissue clearing was performed using a commercial kit (Nuohai, NH-CR-210701), adhering strictly to the manufacturer's protocol. Cleared whole-brain samples were imaged using light-sheet fluorescence microscopy (LSFM; Nuohai, LS18) with a 561 nm excitation laser. For structural analysis, raw datasets were stitched using LS18, software and downsampled (0.25 ×) in ImageJ.

The datasets, comprising 1,500 sagittal slices were registered to the Allen Mouse Brain Common Coordinate Framework (CCFv3) using the BIRDS plugin. Specifically, nine reference slices were selected, and their corresponding atlas indices were manually defined to guide the registration algorithm. The automated registration for the entire dataset was then executed based on these reference points. To ensure spatial accuracy, each slice was monitored during the process, followed by randomized spot-checks post-registration. Any misalignments were resolved through manual adjustment or by re-running the registration pipeline. Based on this refined registration, Imaris Surface modules were generated for specific brain regions (eg., BLA). Vascular structures were segmented based on fluorescence signal intensity, and vascular volume was quantified using Imaris software.

High-Fat Diet (HFD) challenge

To evaluate NVC integrity under clinically relevant metabolic stress, a diet-induced obesity model was employed. Mice were randomized into HFD (SYSE, Cat# PD6001; 60% kcal from fat) or normal fat diet (NFD) control groups for four weeks. Body weight and food intake were monitored weekly. Physiological impact was assessed by quantifying home-cage fecal output as described above. Following the 4-week modelling, mice were subjected to the 40 cm OFT, and whisker stimulation assays to assess emotional and hemodynamic function, respectively. For ELISA analysis, serum was collected 5 min after the 40 cm OFT. For c-Fos immunofluorescence (IF) staining, brain tissues were harvested 90 min post- 40 cm OFT.

Immunoprecipitation (IP) assay

To map the GluN2D interactome, cultured SMCs sorted from P0 *SMA-creER:Ai14* mice brain parenchymal tissue (via tdTomato-based sorting) were collected and subsequently expanded in SMCM. At ~90% confluence, cells were transduced with lentivirus encoding Grin2d-3XFlag (*Lenti-EF1-EGFP-CMV-Grin2d-3XFlag-WPRE*, 5×10^{12} VG/mL). One-week post-transduction, infected cells (EGFP-tdTomato double positive) were purified using an MA-900 sorter and expanded in 10 cm dishes. Upon reaching 90% confluence, cells

were lysed in RIPA buffer. Lysates were incubated with Anti-GluN2D or IgG control antibodies conjugated to Protein G Magnetic Beads (Epizyme, Cat# YJ002) to pull down the protein complex. Eluates were analyzed via Western blot and mass spectrometry.

Blood-Brain Barrier (BBB) permeability assay

To evaluate BBB integrity, the Evans Blue dye extravasation assay was performed.[96] Mice were injected retro-orbitally with 2% Evans Blue (Merck, Cat# E2129; dissolved in saline; 4 μL/g). 45 min post-injection, mice were anesthetized, and whole brains were harvested. Macroscopic dye leakage was first assessed qualitatively using a Zeiss Axio Zoom V16 microscope. For quantitative assessment, brains were weighed and homogenized in 1 mL of 50% trichloroacetic acid (Merck, Cat# 190780). Following centrifugation at 10,000 × g for 20 min at 4 °C, the supernatant was collected, and Evans Blue concentration was quantified by measuring absorbance at 620 nm. To detect microscopic leakage, immunofluorescence staining for Immunoglobulin G (IgG) was performed as described in the Immunofluorescence Staining section. We utilized brain sections from mice subjected to non-perfusion as positive controls—defining their mean fluorescence intensity as the 100% benchmark—to quantify the IgG intensity.

Colorectal endoscopy

To assess stress-induced gastrointestinal pathology, representative images of the distal colon were captured using the ENDOQ colonoscopy system. To ensure unbiased evaluation, a gastroenterologist blinded to the experimental groups scored mucosal morphology and signs of inflammation based on the endoscopic images.

Hematoxylin and Eosin (H&E) staining

For histological analysis, mice were anesthetized and perfused as described above. Harvested tissues were post-fixed in 4% PFA for 6 hours at 4 °C. Following standard paraffin embedding, 8 µm sections were cut (Histo. Microtome, Leica FC-LRM), and H&E staining was performed to visualize gross morphological structures.

Topological characterization of BLA activation patterns

To objectively classify emotional states based on the spatial configuration of neuronal ensembles, a topology-based estimation framework was developed, integrating the Structural Similarity Index (SSIM) with unsupervised hierarchical clustering.

To transition from discrete cellular signals to a continuous representation

of regional activation, raw c-Fos immunofluorescence staining images were

preprocessed in MATLAB. Images were converted to greyscale, inverted, and

normalized to a [0, 1] intensity scale. A multi-scale Gaussian smoothing protocol

($\sigma 1$ = 6 pixels, $\sigma 2$ = 12 pixels) was applied to generate a continuous spatial

intensity map representing the macro-scale topography of neuronal activation

within the BLA. This step effectively minimizes high-frequency noise from

individual cell variations while preserving the global spatial structure.

To quantify the topological similarity between experimental conditions,

pairwise SSIM values were computed between the activation intensity maps.

Unlike traditional metrics that rely on absolute cell counts or pixel-by-pixel

subtraction, SSIM assesses the preservation of structural information by

integrating comparisons of luminance (*l*), contrast (*c*), and structure (*s*). The

similarity index between image *x* and *y* is calculated as:

1381
$$\text{SSIM}(x, y) = [l(x, y)]^{\alpha} \cdot [c(x, y)]^{\beta} \cdot [s(x, y)]^{\gamma}$$

This metric provides a robust measure of spatial fidelity, converting image-

based topographical correlations into a numerical similarity matrix.

The resulting SSIM similarity matrix was utilized for cluster analysis in R.

We defined the topological distance (*D*) between any two conditions as *D* = 1 -

SSIM. Based on this distance matrix, unsupervised hierarchical clustering was

performed using Ward's minimum variance method (method = "ward.D2"). This

algorithm minimizes total within-cluster variance to construct a dendrogram,

effectively grouping behavioral contexts into distinct clusters based purely on topological proximity. These naturally emerging clusters were then mapped to discrete stress Levels (Levels 0–3), defining the subject's state based on the intrinsic topology of their neuronal response rather than arbitrary thresholds.

To ensure rigor, a double-blind workflow was implemented: image processing and algorithmic analysis were conducted by two independent analysts, distinct from the personnel performing wet-lab experiments. Statistical analyses were performed using GraphPad Prism (v9.4.1) and custom MATLAB scripts. All source code is available at website in the Github shown below: https://github.com/JialabEleven/Functional_hyperemia_regulates_emotion-master.

Serial Block-Face Scanning Electron Microscopy (SBF-SEM)

To visualize the ultrastructure of the neurovascular interface, we prepared specimens consistent with our previous protocols.[38] Briefly, we identified ascending venules using X-ray microscopy (Zeiss Xradia 520 Versa) at 4 × magnification (80 kV). We mounted the specimen on an aluminum pin using conductive epoxy (Chemtronics, CW2400) and trimmed it into a 600 × 600 × 700 μm^3 block using a glass knife and a diamond trimming knife (DiATOME, Trim 90). Prior to imaging, we sputter-coated the sample with gold–palladium (Leica EM ACE200). We acquired images using a Carl Zeiss Gemini 300

1410   scanning electron microscope equipped with a Gatan 3View ultramicrotome

1411   and a BSE detector. Imaging parameters were set to 2 kV acceleration voltage

1412   under high vacuum (~1.8×10^{-3} mBar) with focal charge compensation. We

1413   performed serial sectioning at a thickness of 50–75 nm with a cutting speed of

1414   0.1 mm/s.

1415

Stroke modelling

To validate the absence of hypoxic injury in our optogenetic model, we

generated a positive control for hypoxia using the Stroke Induced by Magnetic

ParticLEs (SIMPLE) method.[58-60] Briefly, we injected magnetic microparticles

into the bloodstream and positioned a magnet over the middle cerebral artery

(MCA) to induce targeted occlusion. This permanent ischemia triggers robust

upregulation of Hypoxia-inducible factor 1-alpha (HIF-1α) within 24 hours. We

utilized brain sections from mice subjected to the SIMPLE stroke model as

positive controls—defining their mean fluorescence intensity in the stroke

infarct area as the 100% benchmark—to quantify HIF-1α expression.

1426

Software, code, and data availability

Fluorescence images were processed using Zeiss Zen Lite (v3.4) and ImageJ

(v1.53u). For fiber photometry, signal demodulation and linear regression

correction were performed using the system's built-in proprietary software

(Thinkertech). Whisker stimulation data were acquired and analyzed using the

RWD LSCI software (v5.0). Trajectory tracking for the OFTs, complex OFT, and

EPM was performed using the Noldus EthoVision XT system (v18), which was

also utilized to generate locomotion heatmaps. For the Looming and VR_{eagle}

assays, tracking was conducted using custom scripts (Bayonne). Physiological

and Topology Analysis: Custom MATLAB scripts were developed to analyze

CBF peak kinetics, map aggregate defecation coordinates, and generate neural

activation heatmaps for the topology-based estimation framework. KEGG

pathway enrichment analysis and SSIM-based hierarchical clustering were

executed using custom R algorithms. Statistical analyses were conducted using

GraphPad Prism (v9.4.1) and MATLAB. Figures were assembled using Adobe

Illustrator 2024 (v28.3). Outliers were identified and excluded using the

Interquartile Range (IQR) method (values falling outside the range of Q1 − 1.5

× IQR to Q3 + 1.5 × IQR).[97] Data are expressed as mean ± standard error of

the mean (SEM). The normality of data distribution was assessed using the

Shapiro-Wilk test. Specific statistical tests employed for each experiment are

detailed in the corresponding figure legends. Exact *P* values are reported to

four decimal places. A *P* value < 0.05 was considered statistically significant.

Statistical analyses were performed using GraphPad Prism (v9.4.1). A

comprehensive data resource checklist, detailing figure attributions and sample

sources, is provided in Supplementary Table S2 for reference. Furthermore, to

ensure maximum transparency and statistical robustness, the percentage of c-

Fos positive neurons in the BLA across all experimental cohorts is quantified and presented in Supplementary Figure S24. All custom MATLAB and R code generated for the topology-based estimation framework and hemodynamic analysis is available at the GitHub website for free searching and use: https://github.com/JialabEleven/Functional_hyperemia_regulates_emotion-master.

REFERENCES

1 Roelofs K, Bramson B, Toni I. A neurocognitive theory of flexible emotion control: The role of the lateral frontal pole in emotion regulation. *Ann N Y Acad Sci* 2023; **1525**:28-40.

2 Bortolon C, Chen S, Bonanno GA. Components of emotion regulation flexibility and psychosis: The association between psychosis-proneness and context sensitivity. *Br J Clin Psychol* 2023; **62**:82-95.

3 D'Agostino A, Covanti S, Rossi Monti M, Starcevic V. Reconsidering Emotion Dysregulation. *Psychiatr Q* 2017; **88**:807-825.

4 Easdale-Cheele T, Parlatini V, Cortese S, Bellato A. A Narrative Review of the Efficacy of Interventions for Emotional Dysregulation, and Underlying Bio-Psycho-Social Factors. *Brain Sci* 2024; **14**.

5 De Prisco M, Oliva V, Fico G *et al.* Emotion dysregulation in bipolar disorder compared to other mental illnesses: a systematic review and meta-analysis.

Psychol Med 2023; **53**:7484-7503.

Geng H, Xu P, Aleman A, Qin S, Luo YJ. Dynamic Organization of Large-scale Functional Brain Networks Supports Interactions Between Emotion and Executive Control. *Neurosci Bull* 2024; **40**:981-991.

Szaflarski JP, Allendorfer JB, Heyse H, Mendoza L, Szaflarski BA, Cohen N. Functional MRI of facial emotion processing in left temporal lobe epilepsy. *Epilepsy Behav* 2014; **32**:92-99.

Etkin A, Wager TD. Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 2007; **164**:1476-1488.

Mathias EJ, Kenny A, Plank MJ, David T. Integrated models of neurovascular coupling and BOLD signals: Responses for varying neural activations. *Neuroimage* 2018; **174**:69-86.

Iadecola C. The Neurovascular Unit Coming of Age: A Journey through Neurovascular Coupling in Health and Disease. *Neuron* 2017; **96**:17-42.

Garfinkel SN, Abelson JL, King AP *et al.* Impaired contextual modulation of memories in PTSD: an fMRI and psychophysiological study of extinction retention and fear renewal. *J Neurosci* 2014; **34**:13435-13443.

Delvecchio G, Fossati P, Boyer P *et al.* Common and distinct neural correlates of emotional processing in Bipolar Disorder and Major Depressive Disorder: a voxel-based meta-analysis of functional magnetic resonance imaging studies. *Eur Neuropsychopharmacol* 2012; **22**:100-113.

13 Fan Y, Herrera-Melendez AL, Pestke K *et al.* Early life stress modulates

amygdala-prefrontal functional connectivity: implications for oxytocin effects.

Hum Brain Mapp 2014; **35**:5328-5339.

14 Yang S, Webb AJS. Reduced neurovascular coupling is associated with

increased cardiovascular risk without established cerebrovascular disease: A

cross-sectional analysis in UK biobank. *J Cereb Blood Flow Metab*

2024:271678x241302172.

15 Kunimatsu A, Yasaka K, Akai H, Kunimatsu N, Abe O. MRI findings in

posttraumatic stress disorder. *J Magn Reson Imaging* 2020; **52**:380-396.

16 Terburg D, Morgan BE, Montoya ER *et al.* Hypervigilance for fear after

basolateral amygdala damage in humans. *Transl Psychiatry* 2012; **2**:e115.

17 Feinstein JS, Adolphs R, Damasio A, Tranel D. The human amygdala and

the induction and experience of fear. *Curr Biol* 2011; **21**:34-38.

18 Han K, Lee M, Lim HK *et al.* Excitation-Inhibition Imbalance Leads to

Alteration of Neuronal Coherence and Neurovascular Coupling under Acute

Stress. *J Neurosci* 2020; **40**:9148-9162.

19 Lee S, Kang BM, Shin MK *et al.* Chronic Stress Decreases Cerebrovascular

Responses During Rat Hindlimb Electrical Stimulation. *Front Neurosci* 2015;

9:462.

20 Johnson AC, Uhlig F, Einwag Z, Cataldo N, Erdos B. The neuroendocrine

stress response impairs hippocampal vascular function and memory in male

and female rats. *Neurobiol Dis* 2022; **168**:105717.

21 Shen J, Wang D, Wang X *et al.* Neurovascular Coupling in the Dentate Gyrus

Regulates Adult Hippocampal Neurogenesis. *Neuron* 2019; **103**:878-890 e873.

22 Garthwaite G, Bartus K, Malcolm D *et al.* Signaling from blood vessels to

CNS axons through nitric oxide. *J Neurosci* 2006; **26**:7730-7740.

23 Roy RK, Althammer F, Seymour AJ *et al.* Inverse neurovascular coupling

contributes to positive feedback excitation of vasopressin neurons during a

systemic homeostatic challenge. *Cell Rep* 2021; **37**:109925.

24 Kim KJ, Ramiro Diaz J, Iddings JA, Filosa JA. Vasculo-Neuronal Coupling:

Retrograde Vascular Communication to Brain Neurons. *J Neurosci* 2016;

36:12624-12639.

25 Drew PJ. Neurovascular coupling: motive unknown. *Trends Neurosci* 2022;

45:809-819.

26 Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion

perception II: Implications for major psychiatric disorders. *Biol Psychiatry* 2003;

54:515-528.

27 Hale MW, Hay-Schmidt A, Mikkelsen JD, Poulsen B, Shekhar A, Lowry CA.

Exposure to an open-field arena increases c-Fos expression in a distributed

anxiety-related system projecting to the basolateral amygdaloid complex.

Neuroscience 2008; **155**:659-672.

28 Kovacs LA, Schiessl JA, Nafz AE, Csernus V, Gaszner B. Both Basal and

Acute Restraint Stress-Induced c-Fos Expression Is Influenced by Age in the

Extended Amygdala and Brainstem Stress Centers in Male Rats. *Front Aging*

Neurosci 2018; **10**:248.

Shoji H, Miyakawa T. Effects of test experience, closed-arm wall color, and illumination level on behavior and plasma corticosterone response in an elevated plus maze in male C57BL/6J mice: a challenge against conventional interpretation of the test. *Mol Brain* 2021; **14**:34.

Sturman O, Germain PL, Bohacek J. Exploratory rearing: a context- and stress-sensitive behavior recorded in the open-field test. *Stress* 2018; **21**:443-452.

Kakol-Palm D, Brusberg M, Sand E *et al.* Role of tachykinin NK(1) and NK(2) receptors in colonic sensitivity and stress-induced defecation in gerbils. *Eur J Pharmacol* 2008; **582**:123-131.

Taguchi R, Shikata K, Furuya Y *et al.* Selective corticotropin-releasing factor 1 receptor antagonist E2508 reduces restraint stress-induced defecation and visceral pain in rat models. *Psychoneuroendocrinology* 2017; **75**:110-115.

Monarca RI, Silva RFB, Gabriel SI, Cerveira AM, von Merten S. The Presence of a Shelter in an Open Field Test Has Differential Effects on the Behavior and Stress Response of Two Mouse Species. *J Exp Zool A Ecol Integr Physiol* 2025; **343**:480-492.

An W, Zhang Y, Zhou A, Hu Y. Detrimental Effects of Restricted Cage Size on Reproductive Performance, Exploration Ability, and Anxiety but Not Working Memory of Kunming Mice. *Front Behav Neurosci* 2021; **15**:651782.

Powell WH, Annett LE, Depoortere R, Newman-Tancredi A, Iravani MM. The

selective 5-HT(1A) receptor agonist NLX-112 displays anxiolytic-like activity in mice. *Naunyn Schmiedebergs Arch Pharmacol* 2022; **395**:149-157.

Chen K, Gupta R, Martin-Avila A, Cui M, Xie Z, Yang G. Anesthesia-induced hippocampal-cortical hyperactivity and tau hyperphosphorylation impair remote memory retrieval in Alzheimer's disease. *Alzheimers Dement* 2024; **20**:494-510.

Khalil V, Faress I, Mermet-Joret N, Kerwin P, Yonehara K, Nabavi S. Subcortico-amygdala pathway processes innate and learned threats. *Elife* 2023; **12**.

Zhang D, Ruan J, Peng S *et al.* Synaptic-like transmission between neural axons and arteriolar smooth muscle cells drives cerebral neurovascular coupling. *Nat Neurosci* 2024; **27**:232-248.

Monyer H, Burnashev N, Laurie DJ, Sakmann B, Seeburg PH. Developmental and regional expression in the rat brain and functional properties of four NMDA receptors. *Neuron* 1994; **12**:529-540.

Laurie DJ, Bartke I, Schoepfer R, Naujoks K, Seeburg PH. Regional, developmental and interspecies expression of the four NMDAR2 subunits, examined using monoclonal antibodies. *Brain Res Mol Brain Res* 1997; **51**:23-32.

Qiu YM, Zhang CL, Chen AQ *et al.* Immune Cells in the BBB Disruption After Acute Ischemic Stroke: Targets for Immune Therapy? *Front Immunol* 2021; **12**:678744.

Dion-Albert L, Dudek KA, Russo SJ, Campbell M, Menard C. Neurovascular

adaptations modulating cognition, mood, and stress responses. *Trends Neurosci* 2023; **46**:276-292.

Oyarzabal EA, Hsu LM, Das M *et al.* Chemogenetic stimulation of tonic locus coeruleus activity strengthens the default mode network. *Sci Adv* 2022; **8**:eabm9898.

Chao TH, Zhang WT, Hsu LM, Cerri DH, Wang TW, Shih YI. Computing hemodynamic response functions from concurrent spectral fiber-photometry and fMRI data. *Neurophotonics* 2022; **9**:032205.

Liu X, Li C, Falck JR, Roman RJ, Harder DR, Koehler RC. Interaction of nitric oxide, 20-HETE, and EETs during functional hyperemia in whisker barrel cortex. *Am J Physiol Heart Circ Physiol* 2008; **295**:H619-631.

Renic M, Klaus JA, Omura T *et al.* Effect of 20-HETE inhibition on infarct volume and cerebral blood flow after transient middle cerebral artery occlusion. *J Cereb Blood Flow Metab* 2009; **29**:629-639.

Lacroix A, Toussay X, Anenberg E *et al.* COX-2-Derived Prostaglandin E2 Produced by Pyramidal Neurons Contributes to Neurovascular Coupling in the Rodent Cerebral Cortex. *J Neurosci* 2015; **35**:11791-11810.

Lecrux C, Toussay X, Kocharyan A *et al.* Pyramidal neurons are "neurogenic hubs" in the neurovascular coupling response to whisker stimulation. *J Neurosci* 2011; **31**:9836-9847.

Niwa K, Araki E, Morham SG, Ross ME, Iadecola C. Cyclooxygenase-2 contributes to functional hyperemia in whisker-barrel cortex. *J Neurosci* 2000;

20:763-770.

Hill RA, Tong L, Yuan P, Murikinati S, Gupta S, Grutzendler J. Regional Blood Flow in the Normal and Ischemic Brain Is Controlled by Arteriolar Smooth Muscle Cell Contractility and Not by Capillary Pericytes. *Neuron* 2015; **87**:95-110.

Heissler SM, Chinthalapudi K. Structural and functional mechanisms of actin isoforms. *FEBS J* 2025; **292**:468-482.

Li W, Prakash R, Chawla D *et al.* Early effects of high-fat diet on neurovascular function and focal ischemic brain injury. *Am J Physiol Regul Integr Comp Physiol* 2013; **304**:R1001-1008.

Chang HC, Tai YT, Cherng YG *et al.* Resveratrol attenuates high-fat diet-induced disruption of the blood-brain barrier and protects brain neurons from apoptotic insults. *J Agric Food Chem* 2014; **62**:3466-3475.

Foster DB, Shen LH, Kelly J, Thibault P, Van Eyk JE, Mak AS. Phosphorylation of caldesmon by p21-activated kinase. Implications for the Ca(2+) sensitivity of smooth muscle contraction. *J Biol Chem* 2000; **275**:1959-1965.

Yao YB, Xiao CF, Lu JG, Wang C. Caldesmon: Biochemical and Clinical Implications in Cancer. *Front Cell Dev Biol* 2021; **9**:634759.

Vanlandewijck M, He L, Mae MA *et al.* A molecular atlas of cell types and zonation in the brain vasculature. *Nature* 2018; **554**:475-480.

Hauglund NL, Andersen M, Tokarska K *et al.* Norepinephrine-mediated slow

vasomotion drives glymphatic clearance during sleep. *Cell* 2025; **188**:606-622 e617.

Lan Y, Zhang X, Liu S *et al.* Fate mapping of Spp1 expression reveals age-dependent plasticity of disease-associated microglia-like cells after brain injury. *Immunity* 2024; **57**:349-363 e349.

Jia JM, Chowdary PD, Gao X *et al.* Control of cerebral ischemia with magnetic nanoparticles. *Nat Methods* 2017; **14**:160-166.

Jin Y, Shi P, Wang Y *et al.* Precise control of embolic stroke with magnetized red blood cells in mice. *Commun Biol* 2022; **5**:136.

Hu Y, Chen Z, Huang L *et al.* A translational study on looming-evoked defensive response and the underlying subcortical pathway in autism. *Sci Rep* 2017; **7**:14755.

Shang C, Chen Z, Liu A *et al.* Divergent midbrain circuits orchestrate escape and freezing responses to looming stimuli in mice. *Nat Commun* 2018; **9**:1232.

Wang L, Chen IZ, Lin D. Collateral pathways from the ventromedial hypothalamus mediate defensive behaviors. *Neuron* 2015; **85**:1344-1358.

Silva BA, Gross CT, Graff J. The neural circuits of innate fear: detection, integration, action, and memorization. *Learn Mem* 2016; **23**:544-555.

Clewett D, Gasser C, Davachi L. Pupil-linked arousal signals track the temporal organization of events in memory. *Nat Commun* 2020; **11**:4007.

Grunewald N, Jan A, Salvatico C *et al.* Sequences Flanking the Gephyrin-Binding Site of GlyRbeta Tune Receptor Stabilization at Synapses. *eNeuro*

2018; **5**.

Chiba S, Numakawa T, Ninomiya M, Richards MC, Wakabayashi C, Kunugi H. Chronic restraint stress causes anxiety- and depression-like behaviors, downregulates glucocorticoid receptor expression, and attenuates glutamate release induced by brain-derived neurotrophic factor in the prefrontal cortex. *Prog Neuropsychopharmacol Biol Psychiatry* 2012; **39**:112-119.

Cui Y, Yang Y, Ni Z *et al.* Astroglial Kir4.1 in the lateral habenula drives neuronal bursts in depression. *Nature* 2018; **554**:323-327.

Lee S, Kang BM, Kim JH *et al.* Real-time in vivo two-photon imaging study reveals decreased cerebro-vascular volume and increased blood-brain barrier permeability in chronically stressed mice. *Sci Rep* 2018; **8**:13064.

Masamoto K, Hirase H, Yamada K, Kanno I. Neurovascular coupling-What next? *Prog Brain Res* 2016; **225**:269-272.

Malezieux M, Klein AS, Gogolla N. Neural Circuits for Emotion. *Annu Rev Neurosci* 2023; **46**:211-231.

Yamasaki E, Thakore P, Ali S *et al.* Impaired intracellular Ca(2+) signaling contributes to age-related cerebral small vessel disease in Col4a1 mutant mice. *Sci Signal* 2023; **16**:eadi3966.

Liu Z, Le Q, Lv Y *et al.* A distinct D1-MSN subpopulation down-regulates dopamine to promote negative emotional state. *Cell Res* 2022; **32**:139-156.

Chipperfield AR, Harper AA. Chloride in smooth muscle. *Prog Biophys Mol Biol* 2000; **74**:175-221.

75 Ben-Ari Y. Excitatory actions of gaba during development: the nature of the

nurture. *Nat Rev Neurosci* 2002; **3**:728-739.

76 Runegaard AH, Sorensen AT, Fitzpatrick CM *et al.* Locomotor- and Reward-

Enhancing Effects of Cocaine Are Differentially Regulated by Chemogenetic

Stimulation of Gi-Signaling in Dopaminergic Neurons. *eNeuro* 2018; **5**.

77 Seibenhener ML, Wooten MC. Use of the Open Field Maze to measure

locomotor and anxiety-like behavior in mice. *J Vis Exp* 2015:e52434.

78 Ma Y, Shaik MA, Kim SH *et al.* Correction to 'Wide-field optical mapping of

neural activity and brain haemodynamics: considerations and novel

approaches'. *Philos Trans R Soc Lond B Biol Sci* 2017; **372**.

79 Lerner TN, Shilyansky C, Davidson TJ *et al.* Intact-Brain Analyses Reveal

Distinct Information Carried by SNc Dopamine Subcircuits. *Cell* 2015; **162**:635-

647.

80 Li J, Zhang Y, Zhang D *et al.* Ca(2+) oscillation in vascular smooth muscle

cells control myogenic spontaneous vasomotion and counteract post-ischemic

no-reflow. *Commun Biol* 2024; **7**:332.

81 Zhang YY, Li JZ, Wang WT, Xie HQ, Ruan JY, Jia JM. Vasomotion delineates

cerebral vascular dynamic features and participates in the homeostatic cerebral

blood flow regulation. *Sci Rep* 2025; **15**:36210.

82 Zhang YY, Li JZ, Xie HQ *et al.* High-resolution vasomotion analysis reveals

novel arteriole physiological features and progressive modulation of cerebral

vascular networks by stroke. *J Cereb Blood Flow Metab* 2024; **44**:1330-1348.

83 Stefanovska A, Bracic M, Kvernmo HD. Wavelet analysis of oscillations in

the peripheral blood circulation measured by laser Doppler technique. *IEEE*

Trans Biomed Eng 1999; **46**:1230-1239.

84 Kleinfeld D, Mitra PP. Spectral methods for functional brain imaging. *Cold*

Spring Harb Protoc 2014; **2014**:248-262.

85 Seo JS, Park JY, Choi J *et al.* NADPH oxidase mediates depressive behavior

induced by chronic stress in mice. *J Neurosci* 2012; **32**:9690-9699.

86 Chen M, Ma S, Liu H *et al.* Brain region-specific action of ketamine as a

rapid antidepressant. *Science* 2024; **385**:eado7010.

87 Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller

RC. Functional bowel disorders. *Gastroenterology* 2006; **130**:1480-1491.

88 Arakawa H, Arakawa K, Blanchard DC, Blanchard RJ. Scent marking

behavior in male C57BL/6J mice: sexual and developmental determination.

Behav Brain Res 2007; **182**:73-79.

89 Yu XD, Zhu Y, Sun QX *et al.* Distinct serotonergic pathways to the amygdala

underlie separate behavioral features of anxiety. *Nat Neurosci* 2022; **25**:1651-

1663.

90 Liu J, Li D, Huang J *et al.* Glutamatergic Neurons in the Amygdala Are

Involved in Paclitaxel-Induced Pain and Anxiety. *Front Psychiatry* 2022;

13:869544.

91 Warden MR, Selimbeyoglu A, Mirzabekov JJ *et al.* A prefrontal cortex-

brainstem neuronal projection that controls response to behavioural challenge.

Nature 2012; **492**:428-432.

92 Rampon C, Luppi PH, Fort P, Peyron C, Jouvet M. Distribution of glycine-

immunoreactive cell bodies and fibers in the rat brain. *Neuroscience* 1996;

75:737-755.

93 Dunn KW, Kamocka MM, McDonald JH. A practical guide to evaluating

colocalization in biological microscopy. *Am J Physiol Cell Physiol* 2011;

300:C723-742.

94 Li X, Zhou L, Zhang X *et al.* Proteins secreted by brain arteriolar smooth

muscle cells are instructive for neural development. *Mol Brain* 2022; **15**:97.

95 Ferris HR, Stine NC, Hill-Eubanks DC, Nelson MT, Wellman GC, Koide M.

Epidermal Growth Factor Receptors in Vascular Endothelial Cells Contribute to

Functional Hyperemia in the Brain. *Int J Mol Sci* 2023; **24**.

96 Hou W, Yao J, Liu J *et al.* USP14 inhibition promotes recovery by protecting

BBB integrity and attenuating neuroinflammation in MCAO mice. *CNS Neurosci*

Ther 2023; **29**:3612-3623.

97 Villani AP, Rozieres A, Bensaid B *et al.* Massive clonal expansion of

polycytotoxic skin and blood CD8(+) T cells in patients with toxic epidermal

necrolysis. *Sci Adv* 2021; **7**.

1734

1735   **ACKNOWLEDGMENTS**

1736   We sincerely thank Prof. Bo Li for his invaluable advice and for editing the

1737   manuscript. We are grateful to Prof. Hailan Hu for providing the *Ai32* mice, and

to Prof. Yanmei Tao and Dr. Xu Hu for sharing the *Nestin-Cre* and *Emx1-Cre* lines. We also thank Xiaoxuan Zhang for providing stroke-modeling mice; Xuan Li from Prof. Bo Li's lab for guidance on fiber photometry in the amygdala; and Wuyang Zhang for assisting with data assessment during the revision. We thank Dr. Zhu Zhu for providing critical discussion during the revision. We thank rotation/visiting students Jiaxing Tang, Zeyu Meng, Jiaqi Tu, and Yiran Ge for double-bind analysis of behavioral data. furthermore, we thank the following staff at Westlake University for their technical support: Yi Zhu for sharing psychological insights that facilitated our data interpretation; Yajie Liu at the Laboratory Animal Resource Center (LARC) for guidance on behavioral assays; Dr. Jiongfang Xie at the Microscopy Core Facility for imaging assistance; Dr. Mingzhu Fan at the Mass Spectrometry & Metabolomics Core Facility (MSMCF) for mass spectrometry data collection; and Yanan Tang at the Flow Cytometry Core Facility (FCCF) for assistance with cell sorting.

This work was supported by Westlake Laboratory of Life Sciences and Biomedicine (Project No. 202309002), the National Natural Science Foundation of China (project No. 32571142 and 32170961), State Key Laboratory of Gene Expression (Grant No. 2025ZY01117), Hangzhou Leading Innovation Team (TD2024001), the Joint Fund of the Zhejiang Provincial Natural Science Foundation of China (LHZQN25H160001), the "Pioneer" and "Leading Goose" R&D Program of Zhejiang (grant 2024SSYS0031), and the partially supported from the Construction Fund of Key Medical Disciplines of Hangzhou

1760 (2025HZZD04) from H.W.

1761

1762 **AUTHOR CONTRIBUTIONS**

1782 double-blind genotyping management of mice and cell sorting. D.P.L. helped

1783 the mouse brain section after 35 cm OFT. Y.X.N illustrated the KEGG maps.

1784

1785 **CONFLICT OF INTERESTS**

1786 The authors declare no competing interests.

1787

 FIGURE LEGENDS

 Fig.1 Validation of defecation bouts as a robust quantitative index of

 negative emotional states. a Experimental design for home cage (HC), open

 field tests (OFTs), and acute restraint stress (ARS) assays. **b, c** Representative

 images (**b**) and quantification (**c**) of BLA neuronal activation across different

 stress paradigms shown in (**a**). One-way ANOVA followed by Tukey's multiple

 comparisons test ($P < 0.0001$). **d** Quantification of defecation bouts under the

 paradigms shown in (**a**). Kruskal-Wallis test followed by Dunn's multiple

 comparisons test ($P < 0.0001$). **e** Linear correlation between BLA c-Fos-positive

 neuron counts and defecation bouts. Data were pooled from (**c**) and (**d**). Dotted

 lines are SEM. **f, g** Serum corticosterone concentrations (**f**) under the

 paradigms shown in (**a**) and their linear correlation with defecation bouts (**g**).

 Data were pooled in (**g**) from (**d**) and (**f**). One-way ANOVA followed by Tukey's

 multiple comparisons test ($P < 0.0001$). Dotted lines are SEM in (**g**). **h–j** Total

 running distance (**h**), time spent in the center zone (**i**), and unsupported rearing

 bouts (**j**) in 40 cm and 35 cm OFT. Student's *t*-test (**h**, **i**) and Mann-Whitney test

 (**j**). **k** Representative heatmap of zone preference for a single mouse and the

 spatial distribution of defecation locations from 15 mice in the 40 cm OFT. Each

 dot indicates one defecation location. **l** Statistical quantification of the spatial

 distribution data in (**k**). **m** Experimental schematic of chemogenetic inhibition of

 BLA neurons using *AAV2/5-hSyn-hM4Di-EGFP*. **n, o** Representative images (**n**)

 and quantification (**o**) of BLA neuronal activation under chemogenetic inhibition

in ARS. Student's *t*-test. **p** Quantification of defecation bouts in 40 cm OFT and

ARS following BLA chemogenetic inhibition. Mann-Whitney test. **q** Defecation

rate and defecation latency in the 40 cm OFT. Sample sizes (*N*) are indicated

in the panels. Gehan-Breslow-Wilcoxon test. **r, s** Time spent in the center zone

(**r**) and total running distance (**s**) following chemogenetic inhibition in the 40 cm

OFT. Student's *t*-test. **t** Defecation bouts in the 40 cm OFT and ARS following

BLA injection of the control virus (EGFP). Mann-Whitney test. Except (**k**), each

dot represents one mouse. **** indicates $P < 0.0001$. Data are presented as

mean ± SEM.

Fig.2 GluN2D in arteriolar SMCs is essential for NVC in S1BF and BLA. a

Schematic representation of the neuro-smooth muscle junction (NsMJ) within

the neurovascular unit (NVU). **b** Electrophoretogram (top) and quantification

(bottom) of mRNA levels for NMDA receptor subunits via RT-PCR in cultured

SMCs sorted from P0 brain parenchymal tissue. P0 cerebellum lysates used as

benchmark. Arrows indicate bands' locations. **c** Validation of *Grin2d* expression

via RNAscope. Top: representative image from the cortex of brain slice (yellow

dotted line indicates the arteriolar SMC boundary). Middle: representative

fluorescence intensity linescan of the region indicated by white arrows. Bottom:

Pearson's correlation coefficient analysis. Each dot represents one linescan

region from one mouse. **d, e** Representative images (**d**) and quantification (**e**)

of GluN2D immunofluorescence (IF) staining in the middle cerebral artery

(MCA). Each dot represents one analysis region of MCA from a single mouse. Student's *t*-test. **f-g** Experimental schematic (**f**) and whisker-evoked functional hyperemia in the S1BF (**g**). Panel (**g**) left: time-series traces of functional hyperemia; shaded areas represent SEM across mice. Panel (**g**) right: quantification of the functional hyperemia plateau phase. One-way ANOVA followed by Tukey's multiple comparisons test ($P < 0.0001$). Floxed control mice group including comparable 8 *Grin2d^{f/f}* mice and 7 *Grin1^{f/f}* mice. **h** Schematic for concurrent recording of CBV (Through Rhodamine) and neuronal Ca^{2+} (Through *AAV2/9-hSyn-GCaMP8f*) signals in the BLA. **i, j** Time-series recordings of BLA CBV (**i**) and Ca^{2+} signals (**j**) during the 15-min (900 s) 40 cm OFT session. Shaded areas represent SEM across mice. **k** Area under curve (AUC) quantification of the first 5 minutes of CBV and Ca^{2+} responses from data in (**i**) and (**j**). Student's *t*-test. **l, m** Representative images (**l**) and quantification (**m**) of BLA neuronal activation following the 40 cm OFT. Student's *t*-test. Except (**c** and **e**), each dot represents one mouse. **** indicates $P < 0.0001$. Data are presented as mean ± SEM. See also Supplementary information, Figs. S1–S8.

Fig.3 Systemic SMC-specific GluN2D deficiency-driven negative emotionality is pharmacologically rescued via BLA local intervention. a Defecation bout in the 40 cm OFT shown in 5-minute intervals. Mann-Whitney test. **b** Comparison of defecation bouts across HC (Home cage), OFTs, and ARS paradigms. Kruskal-Wallis test followed by Dunn's multiple comparisons

test (P = 0.0365). **c** Defecation rate and defecation latency in the OFTs. Sample

sizes (N) are indicated in the panels. Gehan-Breslow-Wilcoxon test. **d**

Unsupported rearing bouts (left) and serum corticosterone concentrations (right)

in the 40 cm OFT. Mann-Whitney test (left) and Student's t-test (right). **e**

Experimental schematic of systemic HET0016 administration. **f, g**

Representative images (**f**) and quantification (**g**) of BLA neuronal activation

following systemic HET0016 treatment in the 40 cm OFT. Two-way ANOVA

followed by Tukey's multiple comparisons test (P = 0.0104). **h–j** Quantification

of defecation bouts (**h**), defecation rate and latency (**i**), and unsupported rearing

bouts and serum corticosterone concentrations (**j**) in the 40 cm OFT following

systemic HET0016 administration. Two-way ANOVA on rank-transformed data

followed by Sidak's multiple comparisons test (**h**, P < 0.0001; **j left**, P = 0.00003),

Gehan-Breslow-Wilcoxon test (**i**), and Two-way ANOVA followed by Tukey's

multiple comparisons test (**j right**, P = 0.0212). Sample sizes (N) are indicated

in the panel (**i**). **k** Experimental schematic of local BLA microinjection of

HET0016. **l, m** Representative images (**l**) and quantification (**m**) of BLA

neuronal activation following local BLA HET0016 treatment in the 40 cm OFT.

Two-way ANOVA followed by Tukey's multiple comparisons test (P < 0.0001).

n–p Quantification of defecation bouts (**n**), defecation rate and latency (**o**), and

unsupported rearing bouts and serum corticosterone concentrations (**p**) in the

40 cm OFT following local BLA HET0016 administration. Two-way ANOVA on

rank-transformed data followed by Sidak's multiple comparisons test (**n**, P <

0.0001, **p left**, P = 0.0020), Gehan-Breslow-Wilcoxon test (**o**), and Two-way

ANOVA followed by Tukey's multiple comparisons test (**p right,** P = 0.0190).

Sample sizes (N) are indicated in the panel (**o**). Each dot represents one mouse.

**** indicates P < 0.0001. Data are presented as mean ± SEM. Detailed

statistical data for panel (**c**, **i**, and **o**) are shown in the supplementary table S1,

sub-table 1-3. See also Supplementary information, Figs. S9–S12.

Fig.4 Caldesmon interacts with GluN2D in the SMCs to regulate neurovascular coupling and further modulate emotional states. a KEGG pathway analysis of GluN2D-interacting proteins identified via proteomic analysis from cultured SMCs sorted from P0 brain parenchymal tissue. **b** Immunoprecipitation validation of the interaction between GluN2D and caldesmon from cultured SMCs sorted from P0 brain parenchymal tissue. **c** Experimental schematic for whole-brain (left) and local BLA (right) *Cald1* knockdown (KD) using siRNA. **d, e** Representative images (**d**) and quantification (**e**) of caldesmon immunofluorescence (IF) staining. Student's *t*-test. **f** Electrophoretogram (top) and quantification (bottom) of *Cald1* mRNA expression in the cortex lysates following whole-brain *Cald1* siRNA injection. Student's *t*-test. **g, h** Representative immunofluorescence (IF) images of caldesmon in the BLA (**g**) and ventromedial hypothalamus (VMH, **h**) of mice received local BLA *Cald1* siRNA injection. **i** Quantification of caldesmon expression levels from the experiments shown in (**g, h**). Student's *t*-test. **j, k**

Functional hyperemia in the S1BF during whisker stimulation (**j**) and quantification of the plateau phase (**k**). Shaded areas represent SEM across mice. One-way ANOVA followed by Tukey's multiple comparisons test (P = 0.0041). Floxed control mice group including comparable 3 mice with cisterna magna scramble siRNA injection and 3 mice with local BLA scramble siRNA injection. **l, m** Representative images (**l**) and quantification (**m**) of BLA neuronal activation in the 40 cm OFT. One-way ANOVA followed by Tukey's multiple comparisons test (P < 0.0001). **n** Unsupported rearing bouts in the 40 cm OFT. Kruskal-Wallis test followed by Dunn's multiple comparisons test (P = 0.0001). **o** Serum corticosterone concentrations in the 40 cm OFT. One-way ANOVA followed by Tukey's multiple comparisons test (P = 0.0013). **p, q** Defecation bouts (**p**), defecation rate, and latency (**q**) in the 40 cm OFT. Kruskal-Wallis test followed by Dunn's multiple comparisons test (**p**, P < 0.0001) and Gehan-Breslow-Wilcoxon test (**q**). Sample sizes (N) are indicated between panels (**p** and **q**). Each dot represents one mouse. **** indicates P < 0.0001. Data are presented as mean ± SEM. Detailed statistical data for panel (**q**) are shown in the supplementary table S1, sub-table 4. See also Supplementary information, Figs. S13–S14.

Fig.5 Optogenetic vascular clamping in the BLA is sufficient to drive negative emotionality. a, b Experimental schematic showing the fiber optic implantation site (**a**) and the setup for optogenetic stimulation in a freely moving

mouse within the home cage (**b**). **c, d** Representative CBV traces showing optogenetically induced CBV changes under various stimulation protocols (**c**) and the corresponding quantification (**d**). Each dot represents the area under curve (AUC) change from one mouse during the stimulation period. Shaded areas represent SEM across mice. One-way ANOVA followed by Tukey's multiple comparisons test ($P < 0.0001$). **e–g** Experimental schematic (**e**), representative CBV traces (**f**), and quantification (**g**) in the 40 cm OFT under optogenetic vascular clamping. Each dot represents the AUC change during the optogenetic stimulation from one mouse in (**g**). Shaded areas represent SEM across mice in (**f**). Dotted lines in (**f**) indicate $X = 0$ and $Y = 0$. Student's t-test. **h, i** Representative images (**h**) and quantification (**i**) of BLA neuronal activation following optogenetic stimulation in the 40 cm OFT. Student's t-test. **j–m,** Quantification of unsupported rearing bouts (**j**), serum corticosterone concentrations (**k**), defecation bouts (**l**), and defecation rate and latency (**m**) in the 40 cm OFT during optogenetic stimulation. Coloured dotted lines in (**m**) indicate the median defecation latency for each group. Mann-Whitney test (**j, l**), Student's t-test (**k**), and Gehan-Breslow-Wilcoxon test (**m**). Sample sizes (N) are indicated in the panel (**m**). **n, o** Experimental schematic of 15-min home cage optogenetic stimulation (**n, top**), corresponding BLA CBV response (**n, bottom**), and quantification (**o**). Each dot represents the area under curve (AUC) change during the optogenetic stimulation from one mouse in the home cage. Shaded areas represent SEM across mice. Student's t-test. **p**

Quantification of defecation bouts during home cage optogenetic stimulation. Colored numbers indicate the mean defecation bouts for each group. Kruskal-Wallis test followed by Dunn's multiple comparisons test (P < 0.0001). **q, r** Representative images (**q**) and quantification (**r**) of BLA neuronal activation (**r left**) and serum corticosterone concentrations (**r right**) in the home cage optogenetic stimulation. Student's *t*-test. Each dot represents one mouse. **** indicates P < 0.0001. Data are presented as mean ± SEM. See also Supplementary information, Figs. S15–S16.

Fig.6 Compromised NVC impairs defensive sustainability and alters BLA neural topological scaling. a, b Time-series plots (**a**) and heatmaps (**b**) profiling locomotion speed dynamics before (**a, blue region**) and during (**a, yellow region**) looming stimulation. shaded areas represent SEM across mice in (**a**). The yellow dotted box in (**b**) indicates the 3-second looming duration. **c, d** Quantification of the accumulated recovery latency from freezing (**c**) and unsupported rearing bouts 5 min before and after looming stimulation (**d**). N = 15 mice per group. Kolmogorov-smirnov test (**c**) and Two-way ANOVA on rank-transformed data followed by Sidak's multiple comparisons test (**d**, P < 0.0001). **e** Experimental schematic of the virtual reality facility for visual threat stimulation. **f** Time-series traces showing normalized pupil diameter and locomotion speed during (0–3 s) and after (3–6 s) VR$_{eagle}$ stimulation. The blue region (first 1.5 s) indicates the eagle call, and the yellow region (1.5–3.0 s) indicates the eagle

diving phase. Representative images of the VR$_{eagle}$ stimulus and the mouse eye with a clearly visualized pupil are shown on the bottom. Shaded areas represent SEM across mice. **g, h** Quantification of defensive offset phase (**g**) dynamics and recovery latencies for pupil diameter and locomotion (**h**) from the data in (**b** and **f**). N = 7 mice per group in VR$_{eagle}$ assay, and N = 15 mice per group in looming assay. Student's t-test. **i** Experimental schematic of the complex open field test (c-OFT) arena. **j** Quantification of BLA neuronal activation in contexts of home cage, 40 cm OFT, 35 cm OFT, c-OFT, Looming, and VR$_{eagle}$ assays. Data of 40 cm OFT groups are shared with **Fig. 2m**. Student's t-test. **k** Defecation bouts of mice in the c-OFT. Mann-Whitney test. **l, m** Structural Similarity Index (SSIM)-based unbiased classification of BLA c-Fos topological patterns across seven standard stress contexts in littermate control mice (**l**) and *Grin2d*-cKO mice (**m**). Heatmaps indicate the modified c-Fos intensity in the BLA (processing details can be seen in the supplementary information, **Figure S18a**). **** indicates $P < 0.0001$. Data are presented as mean ± SEM. See also Supplementary information, Figs. S17–S18.

Fig.7 Mural cell-specific GlyRβ deficiency establishes an enhancement model for NVC. a, b Representative RNAscope images (**a**) and quantification (**b**) of *Glrb* mRNA expression via RNAscope in arteriolar SMCs (aSMCs) and venous SMCs (vSMCs) from brain slices. Each dot represents one dot per mural cell; N = 4 mice per group. Mann-Whitney test. **c–e** Representative

images of GlyRβ immunofluorescence from brain slices (**c**) and quantification

comparing arterioles vs. venules in littermate control mice (**d**) and venules

between littermate control and *Glrb*-cKO mice (**e**). In panel (**d**), each dot

represents the number of clusters per 10 µm vessel; N = 4 mice per group. In

panel (**e**), each dot represents one analysis region of vascular mural cells from

a single mouse. Mann-Whitney test (**d**) and Student's *t*-test (**e**). **f**

Electrophoretogram (top) and quantification (bottom) of *Glrb* mRNA expression

in acute sorted Pdgfrβ-positive parenchymal mural cells compared with P60

medullary reticular formation lysates (as positive control). Each dot represents

one mouse. One-way ANOVA followed by Tukey's multiple comparisons test (P

< 0.0001). **g** Co-localization of GlyRβ with gephyrin in cultured human umbilical

venous SMCs (HU-vSMCs). The panel on the right shows the Pearson's

correlation coefficient analysis. Each dot represents the mean correlation

coefficient of ten cells from an independent biological replicate. **h** Co-

localization of GlyRβ with glycinergic axon terminals in the cortex of brain slice.

The orange dotted box indicates the magnified area. Top right: Pearson's

correlation coefficient analysis. Bottom right: Fluorescence intensity linescan

across the region indicated by the white arrow. Each dot represents one

linescan region from one mouse. **i** Sequential electron microscopy (EM)

sections of an ascending venule in the mouse cerebral cortex. White arrows

indicate axonal terminals structural contacts with venous SMCs. EC:

endothelial cell; SMC: smooth muscle cell. **j, k** Experimental schematics of the

2008 whisker stimulation paradigm (**j**) and the transgenic mouse mating strategy (**k**).

2009 **l, m** Functional hyperemia in the S1BF during whisker stimulation (**l**) and

2010 quantification of the corresponding plateau phase (**m**). Shaded areas represent

2011 SEM across mice in (**l**). Each dot represents one mouse. One-way ANOVA

2012 followed by Tukey's multiple comparisons test (P < 0.0001). Control data

2013 (*Grin2d^{f/f}* and *Pdgfrβ-creER:Grin2d^{f/f}*) are shared with **Fig. S11j**. Floxed control

2014 mice group including comparable 3 *Grin2d^{f/f}* mice, 3 *Glrb^{f/f}* mice, and 3

2015 *Grin2d^{f/f}:Glrb^{f/f}* mice. **** indicates P < 0.0001. Data are presented as mean ±

2016 SEM. See also Supplementary information, Figs. S19–S20.

2017

2018 **Fig.8 An enhancement model of NVC promotes resilience against acute**

2019 **and chronic emotional distress. a, b** Representative images (**a**) and

2020 quantification (**b**) of BLA neuronal activation in ARS. Student's *t*-test. **c** Serum

2021 corticosterone concentrations in ARS. Student's *t*-test. **d-f** Defecation bouts in

2022 ARS (**d**) and 40 cm OFT (**e**), and defecation rate and latency in the 40 cm OFT

2023 (**f**). Kruskal-Wallis test followed by Dunn's multiple comparisons test (P <

2024 0.0001 for **d** and P = 0.0014 for **e**). Sample sizes (N) are indicated in the panel

2025 (**f**). Gehan-Breslow-Wilcoxon test for (**f**). Floxed control data in (**e** and **f**) (14

2026 *Grin2d^{f/f}* mice and 17 *Pdgfrβ-creER:Grin2d^{f/f}* mice) are shared with **Fig. 3b, c**

2027 and **S12o**, **p**. Floxed control group in (**e**, **f**) contains comparable 14 *Grin2d^{f/f}*

2028 mice, 5 *Glrb^{f/f}* mice, and 5 *Grin2d^{f/f}:Glrb^{f/f}* mice. **g, h** Representative images (**g**)

2029 and quantification (**h**) of BLA neuronal activation in the 40 cm OFT. Student's *t*-

test. **i, j** Unsupported rearing bouts (**i**) and serum corticosterone concentrations (**j**) in the 40 cm OFT. Mann-Whitney test (**i**) and Student's *t*-test (**j**). **k** Experimental schematic of the chronic restraint stress (CRS) paradigm. **l, m** Representative fecal appearance (**l**) and statistical analysis of Bristol scores and defecation bouts during the CRS session (**m**). All fecal scoring was performed by an independent observer blinded to the experimental groups. *N* = 15 mice per group. Two-way ANOVA on rank-transformed data followed by Sidak's multiple comparisons test ($P < 0.0001$). Bristol s.: Bristol score. **n** Immobility time in the forced swim test (FST) after CRS modelling. One-way ANOVA followed by Tukey's multiple comparisons test ($P < 0.0001$). **o, p** Representative images (**o**) and quantification (**p**) of BLA neuronal activation (**p left**) and serum corticosterone concentrations (**p right**) after CRS. Quantification was performed by an independent observer blinded to the experimental groups. One-way ANOVA followed by Tukey's multiple comparisons test ($P < 0.0001$ for **p left**; $P = 0.0400$ for **p right**). **q** Body weight changes during the CRS paradigm. *N* = 15 mice per group. Student's *t*-test. Each dot represents one mouse. **** indicates $P < 0.0001$. Data are presented as mean ± SEM. Detailed statistical data for panel (**f**) are shown in the supplementary table S1, sub-table 5. Naïve indicates mice without CRS modelling as controls. See also Supplementary information, Figs. S21–S23.

Supplementary information

 Supplementary figure legends

 Fig.S1 Validation of successful GluN2D conditional Knock-out. a, b

 Representative images (**a**) and quantification (**b**) of *Grin1* and *Grin2d* mRNA

 expression via RNAscope in the cerebellum. Each dot represents one region of

 interest (ROI) from a single brain slice per mouse. Student's *t*-test. **c–e**

 Representative immunoblot (top) and quantification (bottom) of GluN2D protein

 levels in P0/P60 cerebellum lysates (**c**), cultured SMCs sorted from P0

 parenchymal tissue (**d**), and P60 cortex lysates(**e**). Each dot represents one

 mouse. One-way ANOVA followed by Tukey's multiple comparisons test for (**c,**

 $P < 0.0001$), and Student's *t*-test for (**d**, **e**). **** indicates $P < 0.0001$. Data are

 presented as mean ± SEM.

 Fig.S2 Validation of GluN2D expression in peripheral organs' SMCs. a

 Representative low-magnification images of tissue sections from the heart, liver,

 spleen, lung, kidney, small intestine, and bladder of *SMA-creER:Ai14* mice (top).

 Yellow dotted boxes indicate the magnified areas shown below. White arrows

 indicate ROIs for fluorescence intensity linescan analysis. **b** Fluorescence

 intensity distribution (linescan) across the ROIs indicated by white arrows in (**a**).

 c Quantification of GluN2D immunofluorescence (IF) staining intensity of

 GluN2D in SMCs across tissues. Each dot represents one mouse. One-way

 ANOVA followed by Tukey's multiple comparisons test ($P < 0.0001$). ****

2073 indicates *P* < 0.0001. Data are presented as mean ± SEM.

2074

Fig.S3 Verification of mouse metabolic status, gut health, and blood-brain

barrier integrity. a–h Quantification of food consumption (**a**), water intake (**b**),

number of fecal pellets (**c**), daily fecal weight (**d**), weight per pellet (**e**), urine

output (**f**), relative body weight changes (**g**), and blood pressure (**h**) measured

in the home cage. For (**a–f**), each dot represents the average value per cage

(4 mice per cage). For (**g, h**), each dot represents an individual mouse. Mann-

Whitney test (**c**) and Student's *t*-test (except **c**). **i** Representative images of

intestinal morphology (appearance and length) and H&E staining of intestinal

sections. **j–n** Quantification of data from (**i**), including mice body weight (**j**),

intestinal length (**k**), intestinal length-to-body weight ratio (**l**), villus length (**m**),

and total 24-hour defecation count (**n**) for mice in the home cage. Each dot

represents one mouse. Student's *t*-test. **o, p** Representative images of the brain

following retro-orbital Evans Blue injection (**o**) and the corresponding

quantification (**p**). Each dot represents one mouse. One-way ANOVA followed

by Tukey's multiple comparisons test (*P* = 0.7654). Data are presented as mean

± SEM.

2091

2092 **Fig.S4 Validation of SMC marker proteins in *Grin2d*-cKO mice. a**

2093 Representative immunoblot (top) and quantification (bottom) of α-SMA

2094 expression. Each dot represents one mouse. Student's *t*-test. **b** Quantification

2095 of Smoothelin immunofluorescence (IF) staining in the middle cerebral artery

2096 (MCA), corresponding to the representative images in **Fig. 2d**. Each dot

2097 represents one analysis region of MCA from a single mouse. *N* = 4 mice per

2098 group. Student's *t*-test. Data are presented as mean ± SEM.

2099

2100 **Fig.S5 The effect of tamoxifen on NVC. a** Whisker-evoked functional

2101 hyperemia in the S1BF of littermate control mice treated with tamoxifen.

2102 Shaded areas represent SEM across mice. **b** Quantification of the functional

2103 hyperemia plateau phase from the data in (**a**). Each dot represents one mouse.

2104 Student's *t*-test. Data are presented as mean ± SEM.

2105

Fig.S6 3D-Reconstruction of BLA arteriolar topology through correlative

integration of the Allen Brain Atlas with tissue-cleared brains from *SMA-*

***creER:Ai14* mouse. a** Ventral view of the mouse BLA. Schematic diagram (left)

was adapted from *The Mouse Nervous System* (p. 460, Fig. 14.1), and 3D-

reconstruciton (right) was generated in Imaris using the Allen Mouse Brain

Common Coordinate Framework plugin. BLA is denoted translucent grey. Blue

denotes arteriolar SMCs. Colored lines represent distinct arteriolar segments.

b Coronal view of the mouse BLA along with arterioles inside. Colored lines

indicate arterioles in basal amygdala anterior part (BAa), basal amygdala

ventral part (BAv), and lateral amygdala (LA).

2116

Fig.S7 Calibration and validation of plasma Rhodamine fluorescence

intensity. a Real-time fluorescence intensity measurement following retro-

orbital injection of various Rhodamine concentrations. **b** Linear correlation

between Rhodamine concentration and fluorescence intensity in (**a**). Each dot

represents the average intensity from 120–150 s post-injection for a specific

concentration. **c** Real-time fluorescence intensity measurements following

retro-orbital injection of 5 mg/mL Rhodamine, showing signal stability from 900–

3600 s post-injection.

2125

2126 **Fig.S8 Baseline CBV and OFT performance in *Grin2d*-cKO mice. a** Baseline

2127 CBV in the BLA during the home cage stage. Each dot represents one mouse.

2128 Student's *t*-test. **b, c** Running distance (**b**) and time spent in the center zone (**c**)

2129 in the 40 cm OFT. Each dot represents one mouse. Student's *t*-test. Data are

2130 presented as mean ± SEM.

2131

2132 **Fig.S9 Serum corticosterone concentrations under baseline conditions.**

2133 **a–c** Serum corticosterone concentrations in mice at the home cage condition

2134 (**a**), following i.p. injection of HET0016 (**b**), and following local BLA

2135 microinjection of HET0016 (**c**). Each dot represents one mouse. Student's *t*-

2136 test. Data are presented as mean ± SEM.

2137

Fig.S10 Dose-dependent and functional validation of HET0016. a Effects of

different HET0016 concentrations on defecation bouts in ARS. Each dot

represents one mouse. Kruskal-Wallis test followed by Dunn's multiple

comparisons test (P = 0.0005). **b, c** Functional hyperemia in the S1BF during

whisker stimulation (**b**) and quantification of the corresponding plateau phase

(**c**). Shaded areas represent SEM across mice. Each dot represents one mouse.

Paired Student's t-test. Data are presented as mean ± SEM.

2145

Fig.S11 NVC dysfunction across multiple models drives negative

emotionality. a–h Validation of the *Emx1-cre:Ptgs2* model. Experimental

schematic of mating strategy (**a**). Representative images (**b**) and quantification

of neuronal COX-2 (encoded by *Ptgs2*) expression (**c**). Functional hyperemia

in the S1BF (**d, e**), BLA neuronal activation (**f, g**), unsupported rearing bouts (**h,**

left), and serum corticosterone (**h, right**) in the 40 cm OFT. **i–r** Validation of the

mural cell-based NMDA conditional knock out models. Mating strategy (**i**),

functional hyperemia (**j**), BLA neuronal activation (**k, l**, and **o, p**). Unsupported

rearing (**m, q**) and serum corticosterone (**n, r**) in the 40 cm OFT. **s–v** Validation

of the *Acta2*-cKO model. Mating strategy (**s**), α-SMA immunoblot quantification

(**t**), and functional hyperemia in the S1BF (**u, v**). **w–aii** Effects of local BLA

HET0016 injection in *Acta2*-cKO mice. Experimental design (**w**), BLA neuronal

activation (**x, y**), serum corticosterone concentrations (**z**), unsupported rearing

bouts (**ai, left**), and defecation bouts (**ai, right**), and defecation rate and latency

in the 40 cm OFT (**aii**) with local BLA HET0016 injection. Each dot represents

one mouse. Shaded areas represent SEM across mice in panel (**d**, **j**, and **u**).

Floxed control mice group in (**j**) including comparable 8 *Grin2d^{f/f}* mice, 7 *Grin1^{f/f}*

mice, and 3 *Grin2d^{ff}: Grin1^{f/f}* mice. Control data (*Grin2d^{f/f}*, *Grin1^{f/f}*, *SMA-*

creER:Grin2d^{f/f}, and *SMA-creER:Grin1^{f/f}*) are shared with **Fig. 2g**. Student's *t*-

test (**c**, **e**, **g**, **h right**, **l**, **n**, **p**, **r**, **t**, and **v**). Mann-Whitney test (**h left**, **m**, and **q**).

Two-way ANOVA followed by Tukey's multiple comparisons test for (**y,** $P =$

0.6564; **z**, $P = 0.7144$). Two-way ANOVA on rank-transformed data followed by

Sidak's multiple comparisons test (**ai left**, $P = 0.4442$; **ai right**, $P = 0.7932$).

One-way ANOVA followed by Tukey's multiple comparisons test (**j**, $P < 0.0001$).

Gehan-Breslow-Wilcoxon test (**aii**). Sample sizes (*N*) are indicated in the panel

(**aii**). **** indicates $P < 0.0001$. Data are presented as mean ± SEM.

2172

Fig.S12 HFD-induced NVC dysfunction and its impact on emotional

distress. a–d High-fat diet (HFD) model validation. Schematic (**a**), body weight

changes (**b**), body composition (fat and lean mass) changes (**c**), and food

consumption (**d**) of mice with HFD or normal-fat diet (NFD). Student's *t*-test. *N*

= 16 mice per group for **b** and **d**, two-way ANOVA followed by Tukey's multiple

comparisons test ($P < 0.0001$). **e-f** Assessment of BBB integrity via Evans Blue

injection (**e**) and quantification (**f**). student's *t*-test. **g, h** Functional hyperemia in

the S1BF (**g**) and quantification of the plateau phase (**h**). shaded areas

represent SEM across mice in (**g**). Student's *t*-test. **i–n** Effects of local BLA

HET0016 injection in HFD mice. Experimental design (**i**), BLA neural activation

(**j, k**), defecation bouts (**l**), defecation rate and latency (**m**), and unsupported

rearing (**n**) of HFD mice in the 40 cm OFT with local BLA HET0016 injection.

Two-way ANOVA followed by Tukey's multiple comparisons test for (**k**), P =

0.0001. Two-way ANOVA on rank-transformed data followed by Sidak's multiple

comparisons test (**l** and **n**), P < 0.0001 for (**l**), and P = 0.0009 for (**n**). Gehan-

Breslow-Wilcoxon test (**m**). Sample sizes (N) are indicated in the panel (**m**). **o,**

p Defecation bouts (**o**), and defecation rate and latency (**p**) in the 40 cm OFT

and ARS (panel **o** only). Kruskal-Wallis test followed by Dunn's multiple

comparisons test for (**o**) (P < 0.0001). Gehan-Breslow-Wilcoxon test for (**p**).

Sample sizes (N) are indicated in the panels (**o** and **p**). Control data

(*Grin2d^{f/f}*and *SMA-creER:Grin2d^{f/ff}*) are shared with **Fig. 3a-c**. Detailed

statistical data for panel (**p**) are shown in the supplementary table S1, sub-table

6. Dotted line in (**p**) used to separate two cluster of distribution pattern between

NVC-deficiency groups and their littermate controls. Each dot represents one

mouse. **** indicates P < 0.0001. Data are presented as mean ± SEM.

Fig.S13 Specificity of S1BF *Cald1* knockdown. a Schematic showing the

siRNA injection site in the S1BF and its spatial relationship with the BLA (orange

2201 dotted line). **b–d** Representative images of caldesmon IF stainningin the S1BF

2202 (**b**) and BLA (**c**) following local S1BF *Cald1* siRNA injection, with quantification

2203 in (**d**). Each dot represents one mouse. Two-way ANOVA followed by Tukey's

2204 multiple comparisons test ($P < 0.0001$). **** indicates $P < 0.0001$. Data are

2205 presented as mean ± SEM.

2206

Fig.S14 Impact of S1BF *Cald1* knockdown on emotional responses. a, b

Defecation bouts (**a**), and defecation rate and latency (**b**) in the 40 cm OFT

following S1BF *Cald1* KD in the 40 cm OFT. Each dot represents one mouse.

Mann-Whitney test in (**a**). Sample sizes (*N*) are indicated in the panel (**b**).

Gehan-Breslow-Wilcoxon test in (**b**). **c, d** Representative images (**c**) and

quantification (**d**) of BLA neuronal activation in the 40 cm OFT. Each dot

represents one mouse. Student's *t*-test. **e, f** Unsupported rearing bouts (**e**) and

serum corticosterone concentrations (**f**) in the 40 cm OFT. Each dot represents

one mouse. Mann-Whitney test in (**e**), and student's *t*-test in (**f**). Data are

presented as mean ± SEM.

2217

Fig.S15 Spectral energy analysis of BLA blood flow signals. a, b

Representative Fourier plot of BLA CBV signals from 0–1 Hz (**a**) and

quantification of ultra-low-frequency oscillations (0–0.3 Hz) (**b**) in the 40 cm OFT.

Each dot represents one mouse. One-way ANOVA followed by Tukey's multiple

comparisons test ($P < 0.0001$). **** indicates $P < 0.0001$. Data are presented

as mean ± SEM.

2224

Fig.S16 Supplementary data for BLA optogenetic stimulation. a, b Running

distance (**a**) and time spent in the center zone (**b**) during optogenetic stimulation

in the 40 cm OFT. Each dot represents one mouse. Student's *t*-test. **c, d**

Representative images (**c**) and quantification (**d**) of Hif-1α immunofluorescence

in the BLA during the optogenetic stimulation in the home cage. Stroke model

tissue serves as a positive control. Each dot represents one mouse. One-way

ANOVA followed by Tukey's multiple comparisons test ($P < 0.0001$). ****

indicates $P < 0.0001$. Data are presented as mean ± SEM.

2233

2234 **Fig.S17 Supplementary locomotion speed and pupil diameter in looming**

2235 **and VR$_{eagle}$ assays. a–d** Quantification of pre-stimulation locomotion speed in

2236 looming (**a**) and VR$_{eagle}$ (**b**) assays, and pupil diameter before (**c**) and at the

2237 conclusion (**d**) of VR$_{eagle}$ stimulation. Each dot represents one mouse. Student's

2238 *t*-test. Data are presented as mean ± SEM.

2239

2240 **Fig.S18 Methodological framework for topological clustering and**

2241 **clustering coordinates of *Grin2d*-cKO mice. a** Schematic workflow for

2242 clustering data processing and generation. **b–g**, Clustering coordinates of

Grin2d-cKO mice mapped against the unbiased classification library of littermate controls across individual contexts. **h** the summary of the load-dependent biphasic pattern. Quantification data shared with **Fig. 6j**, and heatmap shared with **Figure. 6l, m**. Each 50% transparent dot represents one mouse. The non-transparent dot represents the mean value. Student's *t*-test. Data are presented as mean ± SEM.

Fig.S19 *Glrb* expression profile in the vascular transcriptome database. Transcriptional levels of *Glrb* in venous SMCs (vSMCs), arteriolar SMCs (aaSMCs), and arterial SMCs (aSMCs). Data are presented as mean ± SEM.

Fig.S20 Functional and behavioral validation of SMC-specific *Glrb*-cKO mice. a, b Functional hyperemia in the S1BF (**a**) and quantification of the plateau phase (**b**). **c** Defecation bouts of mice during the ARS. Each dot represents one mouse. Shaded areas represent SEM across mice in (**a**). Student's *t*-test (**b**) and Mann-Whitney test (**c**). Data are presented as mean ± SEM.

Fig.S21 Local BLA *Cald1* knockdown occludes the NVC enhancement in *Glrb*-cKO mice. a, b Representative images (**a**) and quantification (**b**) of BLA neuronal activation of *Glrb*-cKO mice with local BLA *Cald1* KD in ARS.

Student's *t*-test. **c, d** Defecation bouts (**c**) and serum corticosterone (**d**) in *Glrb*-cKO mice with local BLA *Cald1* KD in ARS. Mann-Whitney test (**c**) and Student's *t*-test (**d**). **e** Baseline cerebral perfusion analysis before and after *Cald1* siRNA cisterna magna injection in the *Glrb*-cKO mice. Paired Student's *t*-test. Each dot represents one mouse. Data are presented as mean ± SEM.

Fig.S22 CRS modelling and *Glrb*-cKO mice naïve behaviours in the 40 cm OFT. a Time spent in the open arms of the elevated plus maze after CRS modelling (EPM, left) and fecal quality (Bristol score, right, data similar to that in **Fig. 8m**) at day-28 of CRS. Student's *t*-test (left) and Mann-Whitney test (right). **b, c** Representative images (**b**) and quantification (**c**) of neuronal activation in the lateral habenula (LHb) after CRS modelling. Student's *t*-test. **d, e** Total running distance (**d**) and time spent in the center zone (**e**) in the 40 cm OFT. Naïve indicates mice without CRS modelling as controls. Each dot represents one mouse. Student's *t*-test. **** indicates $P < 0.0001$. Data are presented as mean ± SEM.

Fig.S23 Assessment of BBB integrity following CRS. Representative images (including non-perfusion positive control) and quantification of IgG fluorescence in the S1BF and BLA after CRS modelling. Each dot represents the average IgG level per mouse, calculated from three ROIs in both the cortex

2285 and the BLA. One-way ANOVA followed by Tukey's multiple comparisons test

2286 ($P < 0.0001$). **** indicates $P < 0.0001$. Data are presented as mean ± SEM.

2287

Fig.S24 Statistical quantification of the percentage of c-Fos positive

neurons in the BLA. (a–u) Re-quantification of c-Fos positive neuron ratios

(normalized to total NeuN positive neurons in the BLA) corresponding to the

original intensity-based data presented in Figure 1c (**a**), Figure 1e (**b**), Figure

1o (**c**), Figure 2m (**d**), Figure 3g (**e**), Figure 3m (**f**), Figure S11g (**g**), Figure S11l

(**h**), Figure S11p (**i**), Figure S11y (**j**), Figure S12k (**k**), Figure 4m (**l**), Figure S14d

(**m**), Figure 5i (**n**), Figure 5r left (**o**), Figure 6j (**p**), Figure 8b (**q**), Figure 8h (**r**),

Figure 8p left (**s**), Figure S21b (**t**), and Figure S22c (**u**), respectively.

2296

2297 *Supplementary tables*

2298 **Supplementary Table S1. P_{value} of Defecation rate-latency curve**

2299 Sub table 1 for **Fig. 3c**

P	1	2	3	4	5	6
1	1	0.0031	0.0019	****	****	****
2		1	0.1934	0.3581	0.1384	0.1189
3			1	0.8685	0.3193	0.3458
4				1	0.1585	0.1083

5					1	0.9278
6						1

1. OFT_40 *Grin2d*^{f/f}

2. OFT_35 *Grin2d*^{f/f}

3. OFT_10 *Grin2d*^{f/f}

4. OFT_40 *Grin2d*-cKO

5. OFT_40 *Grin2d*-cKO

6. OFT_40 *Grin2d*-cKO

**** indicates $P < 0.0001$

2307

2308 Sub table 2 for **Fig. 3i**

P	1	2	3	4
1	1	0.1650	****	0.6470
2		1	****	0.2934
3			1	****
4				1

2309 1. *Grin2d*^{f/f}+Saline

2310 2. *Grin2d*^{f/f}+HET0016

2311 3. *Grin2d*-cKO+Saline

2312 4. *Grin2d*-cKO+HET0016

2313 **** indicates $P < 0.0001$

2314

2315 Sub table 3 for **Fig. 3o**

P	1	2	3	4
1	1	0.0679	0.0003	0.2123
2		1	****	0.3173
3			1	****
4				1

2316 5. *Grin2d^{f/f}*+Saline

2317 6. *Grin2d^{f/f}*+HET0016

2318 7. *Grin2d*-cKO+Saline

2319 8. *Grin2d*-cKO+HET0016

2320 **** indicates $P < 0.0001$

2321

2322 Sub table 4 for **Fig. 4q**

P	1	2	3	4	5
1	1	****	****	0.0009	****
2		1	0.5745	0.8881	0.9999
3			1	0.99990	0.6926
4				1	0.6353
5					1

1. Scramble (Whole-brain+BLA)

2. *Cald1* KD (Whole-brain)

3. *Cald1* KD (BLA)

4. Scramble (BLA)+*Grin2d*-cKO

5. *Cald1* KD (BLA)+*Grin2d*-cKO

**** indicates $P < 0.0001$

2329

2330   Sub table 5 for **Fig. 8f**

P	1	2	3	4
1	1	0.0012	0.6285	0.4703
2		1	0.0035	0.0244
3			1	0.2637
4				1

2331   1. Floxed control mice

2332   2. *Pdgfrβ-creER:Grin2d^{f/f}*

2333   3. *Pdgfrβ-creER:Glrb^{f/f}*

2334   4. *Pdgfrβ-creER:Grin2d^{f/f}:Glrb^{f/f}*

2335   **** indicates $P < 0.0001$

2336

2337

2338

2339 Sub table 6 for **Fig. S12p**

P	1	2	3	4	5	6	7	8	9	10	11	12
1	1	****	0.0002	0.0823	0.0005	0.8480	****	****	0.4906	****	0.5680	****
2		1	0.7333	0.0092	0.7209	0.0002	0.0786	0.2073	0.0043	0.4279	0.0010	0.2679
3			1	0.0047	0.7747	0.0001	0.2580	0.3542	0.0028	0.7549	0.0010	0.4218
4				1	0.0130	0.1038	****	0.0001	0.2961	0.0009	0.0548	0.0005
5					1	0.0004	0.2596	0.3170	0.0078	0.9480	0.0022	0.5377
6						1	****	****	0.5970	****	0.4793	****
7							1	0.4699	****	0.2935	0.0003	0.8118
8								1				
9								****	1	0.0008	0.2695	0.0005
10								0.3463		1	0.0003	0.5583
11								****			1	0.0005
12								0.9373				1

1. *Grin2d^{f/f}*

2. *SMA-creER:Grin2d^{f/f}*

3. *Pdgfrβ-creER:Grin2d^{f/f}*

4. *Grin1^{f/f}*

5. *SMA-creER:Grin1^{f/f}*

6. *Ptgs2^{f/f}*

7. *Emx1-cre:Ptgs2^{f/f}*

8. *SMA-creER:Grin1^{f/f}:Grin2d^{f/f}*

9. *Acta2*$^{f/f}$

10. *Pdgfrβ-creER:Acta2*$^{f/f}$

11. NFD

12. HFD

**** indicates $P < 0.0001$

Supplementary Table S2. Sample source checklist

No.	Location	Sample type	Source
1	Fig. 1b	Immunofluorescence staining	Brain slice
2	Fig. 1n	Immunofluorescence staining	Brain slice
3	Fig. 2b	Semi-quantified PCR	Sorted cultured cell
4	Fig. 2c	RNAscope	Brain slice
5	Fig. 2d	Immunofluorescence staining	Isolated MCA
6	Fig. 2l	Immunofluorescence staining	Brain slice
7	Fig. S1a	RNAscope	Brain slice
8	Fig. S1c	Western blot	Cerebellum lysates
9	Fig. S1d	Western blot	Sorted cultured cell
10	Fig. S1e	Western blot	Cortex lysates
11	Fig. S2a	Immunofluorescence staining	Tissue slice
12	Fig. S3i	Hematoxylin and eosin staining	Intestine slice
13	Fig. S3o	Absorbance measurement	Brain lysates

14	Fig. S4a	Western blot	Sorted culture cell
15	Fig. S4b	Immunofluorescence staining	Isolated MCA
16	Fig. 3f	Immunofluorescence staining	Brain slice
17	Fig. 3l	Immunofluorescence staining	Brain slice
18	Fig. S11b	Immunofluorescence staining	Brain slice
19	Fig. S11f	Immunofluorescence staining	Brain slice
20	Fig. S11k	Immunofluorescence staining	Brain slice
21	Fig. S11o	Immunofluorescence staining	Brain slice
22	Fig. S11t	Western blot	Cortex lysates
23	Fig. S11x	Immunofluorescence staining	Brain slice
24	Fig. S12e	Absorbance measurement	Brain lysates
25	Fig. S12j	Immunofluorescence staining	Brain slice
26	Fig. 4b	Western blot	Sorted culture cell
27	Fig. 4d	Immunofluorescence staining	Brain slice
28	Fig. 4f	Semi-quantified PCR	Cortex lysate
29	Fig. 4g	Immunofluorescence staining	Brain slice
30	Fig. 4h	Immunofluorescence staining	Brain slice
31	Fig. 4l	Immunofluorescence staining	Brain slice
32	Fig. S13b	Immunofluorescence staining	Brain slice
33	Fig. S13c	Immunofluorescence staining	Brain slice
34	Fig. S14c	Immunofluorescence staining	Brain slice
35	Fig. 5a	Immunofluorescence staining	Brain slice

36	Fig. 5h	Immunofluorescence staining	Brain slice
37	Fig. 5q	Immunofluorescence staining	Brain slice
38	Fig. S16c	Immunofluorescence staining	Brain slice
39	Fig. 7a	RNAscope	Brain slice
40	Fig. 7c	Immunofluorescence staining	Brain slice
41	Fig. 7f	Semi-quantified PCR	Acute sorted cell
42	Fig. 7g	Immunofluorescence staining	Cultured cell line
43	Fig. 7h	Immunofluorescence staining	Brain slice
44	Fig. 7i	Electron microscopy image	Brain slice
45	Fig. 8a	Immunofluorescence staining	Brain slice
46	Fig. 8g	Immunofluorescence staining	Brain slice
47	Fig. 8o	Immunofluorescence staining	Brain slice
48	Fig. S21a	Immunofluorescence staining	Brain slice
49	Fig. S22b	Immunofluorescence staining	Brain slice
50	Fig. S23	Immunofluorescence staining	Brain slice

2355

2356 **Supplementary Table S3. Mice information**

Abbreviation	Full name	Source
Ai32	*B6;129S-Gt(ROSA)26Sortm32(CAG-COP4*H134R/EYFP)Hze/J*	*Hailan Hu, Zhejiang University*

Nestin-cre	*B6.Cg-Tg(Nes-cre)1Kln/J*	*Yanmei Tao and Xu Hu, Hangzhou Normal University/ Southeast University*
Emx1-cre	*B6.129S2-Emx1tm1(cre)Krj/J*	*Yanmei Tao and Xu Hu, Hangzhou Normal University/ Southeast University*
SMA-creER	*Tg(Acta2-cre/ERT2)12Pcn*	*Bo Zhou, ShanghaiInstitutes for Biological Sciences, Chinese Academy of Sciences*
Pdgfrβ-creER	*Tg(Pdgfrb-cre/ERT2)6096Rha*	*Xiang Yu, Peking University*
Ai47	*Gt(ROSA)26Sortm47(CAG-EGFP*)Hze*	*Zilong Qiu, Chinese Academy of Sciences Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences.*

Ai14	*B6.Cg-Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J*	*Woo-Ping Ge, Chinese Institute of Brain Research (CIBR)*
Grin2d	*B6/JGpt-Grin2dem1Cflox/Gpt*	*GemPharmatech*
Grin1	*Grin1tm2Stl*	*Yangang Sun, Chinese Academy of Sciences Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences*
Ptgs2	*C57BL/6JCya-Ptgs2em1flox/Cya*	*Cyagen*
Acta2	*C57BL/6JGpt-Acta2em1Cflox/Gpt*	*GemPharmatech*
Glrb	*C57BL/6JGpt-Glrbem1Cflox/Gpt*	*GemPharmatech*

2357

2358

2359

Supplementary Table S4. Antibodies information

Name	Host	Cat#	Dilution	Company
Anti-NMDAR2D Antibody	Mouse	MAB5578	1:400 (IF) 1:2,500 (WB) 1:1,000 (IP)	Merck
Anti-Smoothelin antibody	Rabbit	ab219652	1:400 (IF)	abcam
α-Smooth Muscle Actin (ACTA2) Rabbit mAb	Rabbit	A17910	1:2,500 (WB)	ABclonal
ACE2 Recombinant Rabbit Monoclonal Antibody (SN0754)	Rabbit	MA5-32307	1:400 (IF)	Invitrogen
Gephyrin Polyclonal Antibody	Rabbit	PA5-29036	1:400 (IF)	Invitrogen
BD Pharmingen™ Purified Rat Anti-Mouse CD31	Rat	557355	1:200 (IF)	BD
Anti-NeuN purified Antibody	Guinea Pig	ABN90P	1:800 (IF)	Merck
COX2/PTGS2 Rabbit pAb	Rabbit	A1253	1:400 (IF)	ABclonal
c-Fos antibody	Rabbit	226 003	1:1,000 (IF)	SYSY

Recombinant Alexa Fluor® 488 Anti-Caldesmon/CDM antibody	Rabbit	ab208116	1:400 (IF) 1:2,500 (WB)	abcam
iFluor™ 488 Conjugated Goat anti-mouse IgG polyclonal Antibody	Goat	HA1125	1:400 (IF)	HUABIO
Mouse Aminopeptidase N/CD13 Antibody	Goat	AF2335	1:200 (IF)	R&D
CD140b (PDGFRB) Monoclonal Antibody (APB5), APC, eBioscience™	Rat	17-1402-82	1:50 (FCM)	Invitrogen
Mice IgG	Mouse	A7028	1:1,000 (IP)	Beyotime
Donkey anti-Mouse IgG (H+L) ReadyProbes™ Secondary Antibody, Alexa Fluor™ 488	Donkey	R37114	1:1,000 (IF)	Invitrogen
Donkey anti-Rabbit IgG (H+L) ReadyProbes™ Secondary Antibody, Alexa Fluor™ 488	Donkey	R37118	1:1,000 (IF)	Invitrogen
Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 546	Donkey	A10040	1:1,000 (IF)	Invitrogen

Donkey anti-Rat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488	Donkey	A21208	1:1,000 (IF)	Invitrogen
Donkey anti-Rat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 555	Donkey	A78945	1:1,000 (IF)	Invitrogen
Donkey anti-Rat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 647	Donkey	A48272	1:1,000 (IF)	Invitrogen
Alexa Fluor® 647 AffiniPure™ Donkey Anti-Guinea Pig IgG (H+L)	Donkey	706-605-148	1:1,000 (IF)	Jackson ImmunoResearch
β-Tubulin Mouse mAb	Mouse	AC021	1:5,000 (WB)	ABclonal
Goat Anti-Mouse IgG, HRP Conjugated	Goat	CW0102	1:10,000 (WB)	CWBIO
Goat Anti-Rabbit IgG, HRP Conjugated	Goat	CW0103	1:10,000 (WB)	CWBIO
DDDDK-Tag Rabbit mAb	Rabbit	AE063	1:400 (IF)	ABclonal

| Hif-1α | | Rabbit | ER1802-41 | 1:100 (IF) | HUABIO |

2361

Supplementary Table S5. Primer pairs information

Name		Sequence
Grin1	F	AAGAACCTGCAGCAGTACCA
	R	GCAGCAGGACTCATCAGTGT
Grin2a	F	TCAGGGACTTTGGAAAACGCT
	R	AAGTCATGCAGGGTTGGTTG
Grin2b	F	ACTAACTATCAATGAAGAACGGT
	R	CCAGACCCCAGAGTAACCA
Grin2c	F	TGTGGGCCTTCTTCGCTGTCATCT
	R	TGCCATTAGGTACCGTGCCAAAAC
Grin2d	F	TGGAGGAGTACGACTGGACAT
	R	CGCACTGACACTACGGAGC
Grin3a	F	CTGCACACCAGTCAGAGGTT
	R	TTCGTTGGTTGTCGTGACTC

Grin3b	F	TGGGGGTCTACCACTTGTCA
	R	TGGATCTTCTGGCTCGTGTG
Glrb	F	TTCAAAGGCATTCCTGTTGATGT
	R	TGAGCCTCTGAAGTCACTAGG
Cald1	F	GCCGTTCAAGTGCTTCACTC
	R	TGTCAATCTTGGAGACTACTGCT
18s	F	CGGACACGGACAGGATTGACA
	R	CCAGACAAATCGCTCCACCAACTA
Cald1 (siRNA)	F	GUUUGAUCCGACCAUAACAGA
	R	UGUUAUGGUCGGAUCAAACUC

2363

2364 *Supplementary Video*

2365 **Supplementary Video S1. Topological distribution of arterioles in the BLA.**

2366

2367 **Supplementary Video S2. Pupil diameter changes during the stimulation**

2368 **in the VR$_{eagle}$ stimulation assay.**

a KEGG Pathway Enrichment_Top10

b

c siRNA administration
Whole-brain   Local BLA

d Caldesmon DAPI — 1 mm

e

f Whole-brain injection
Cald1 18s
0.0271

g BLA with local BLA injection
Scramble   *Cald1* KD
Caldesmon NeuN DAPI — 400 μm

h VMH with local BLA injection
Scramble   *Cald1* KD
Caldesmon NeuN DAPI — 400 μm

i S: Scramble
C: *Cald1* KD
From the same mouse

j

k

l Local BLA injection
NeuN c-Fos DAPI — 200 μm
Scramble+*Grin2d*[f/f]
Cald1 KD+*Grin2d*[f/f]
Scramble+*Grin2d*-cKO
Cald1 KD+*Grin2d*-cKO

m

n

o Local BLA injection

p

q

a SMA-creER:Ai32 ChR2-EYFP NeuN DAPI ChR2-EYFP 200 μm

b Stimulation (473 nm) CBV recording HC free-moving mouse BLA recording

c △CBV (%) Optogenetic stimulation Seconds
- ChR2-No stimulation
- ChR2-85ms-5ms-5ms-10Hz-0.302mW
- ChR2-5ms-5ms-5ms-10Hz-0.939mW
- ChR2-85ms-5ms-5ms-10Hz-0.939mW
- EYFP-No stimulation
- EYFP-85ms-5ms-5ms-10Hz-0.302mW
- EYFP-5ms-5ms-5ms-10Hz-0.939mW
- EYFP-85ms-5ms-5ms-10Hz-0.939mW

d Stimulation period AUC (au.) 0.0077 0.0403 ns ns ns ns

e Stimulation (473 nm) CBV recording 40 cm OFT free-moving mouse BLA recording 40 cm OFT

f △CBV (%) 40 cm OFT + Optogenetic stimualtion Seconds EYFP ChR2-EYFP

g AUC (a.u.x10³) 0.0043 EYFP ChR2-EYFP

h EYFP ChR2-EYFP EYFP c-Fos DAPI ChR2-EYFP c-Fos DAPI 200 μm

i c-Fos⁺ neurons in BLA per mm² **** EYFP ChR2-EYFP

j Unsupported rearing bouts 0.0157 EYFP ChR2-EYFP

k Corticosterone (ng/ml) 0.0473 EYFP ChR2-EYFP

l Defecation bouts 0.0338 EYFP ChR2-EYFP

m Defecation rate (%) 144.5 s 565 s 0.0007 Defecation latency (s) EYFP (N = 12) EYFP-ChR2 (N = 12)

n Fecal collection period 15 min 15 min 15 min Recording Recording + Optogenetic stimulation Recording Optogenetic stimulation △CBV (%) Seconds EYFP ChR2-EYFP

o EYFP ChR2-EYFP AUC (a.u.) ns 0.0004 ns Before During After

p Defecation bouts (stimulation period) 0.0052 0.0061 0.23 2.00 4.30 EYFP_HC ChR2-EYFP_HC Control mice_ARS

q EYFP ChR2-EYFP EYFP c-Fos DAPI ChR2-EYFP c-Fos DAPI 200 μm

r c-Fos⁺ neurons in BLA per mm² Corticosterone (ng/ml) **** 0.0487 EYFP ChR2-EYFP

a — Locomotion speed (cm/s); *Grin2d*[f/f], *Grin2d*-cKO; ns; Spontaneous locomotion, Stress-induced locomotion; Seconds (-3 to 3)

b — 60 45 30 15 0 (cm/s); Mouse number; *Grin2d*[f/f], *Grin2d*-cKO; -6s 0s 3s 24s

c — Accumulated recovery latency (s); *Grin2d*[f/f], *Grin2d*-cKO; 0.0320; Accumulated mice amount

d — Unsupported rearing bouts; ns; ****; ns; *Grin2d*[f/f] (Before/After); *Grin2d*-cKO (Before/After)

e — VR Screen; Float ball

f — Eagle call, Eagle diving; Pupil diameter, Speed; *Grin2d*-cKO, *Grin2d*[f/f]; Normalized (%); Seconds (0 to 6)

g — Defensive offset phase (s); 0.0249; *Grin2d*[f/f], *Grin2d*-cKO

h — VR[eagle]_pupil diameter recovery latency (s); Looming_locomotion speed recovery latency (s); 0.0468; 0.0344; 12.41; 5.01; 59.61%; 13.13; 5.38; 59.05%; VR[eagle], Looming; *Grin2d*[f/f], *Grin2d*-cKO

i — Complex OFT; Metal fence; Black floor; 21 cm; White floor; 46 cm

j — c-Fos[+] neurons in BLA per mm²; ns; 0.0016; 0.0278; ns; 0.0122; 0.0123; Home cage, OFT_40, OFT_35, Complex OFT, Looming, VR[eagle]; *Grin2d*[f/f], *Grin2d*-cKO

k — Complex OFT; Defecation bouts; 0.0002; *Grin2d*[f/f], *Grin2d*-cKO

l — Stress level: Level_0, Level_1, Level_2, Level_3; Level_0 Home cage; Level_1 OFT_40, OFT_35; Level_2 ARS, Complex OFT; Level_3 Looming, VR[eagle]; 1 High, 0.5, 0 Low; *Grin2d*[f/f]

m — *Grin2d*[f/f]; *SMA-creER:Grin2d*[f/f] (*Grin2d*-cKO); Home cage, VR[eagle], OFT_40, Looming, Complex OFT, OFT_35, ARS; Level_0, Level_3, Level_1, Level_3, Level_2, Level_1, Level_2; 200 μm

a *Pdgfrβ-creER:Ai14*

Glrb mRNA tdTomato / *Glrb* mRNA — 10 μm

c *Pdgfrβ-creER:Ai14* / *Pdgfrβ-creER:Ai14:Glrb^{f/f}*

GlyRβ tdTomato / GlyRβ — 40 μm

b *Glrb* signal dots per cell — vSMCs aSMCs — ****

d GlyRβ clusters per 10 μm — Venules Arterioles — 0.0007

e GlyRβ expression level (%) — 0.0045 — *Glrb^{f/f}* / *Glrb^{f/f}*-cKO

f *Glrb* expression level (%) — *Glrb* 18s — Adult cortex — **** —
● Adult cerebral cortex
● Sorted mural cells from *Glrb^{f/f}*
● Sorted mural cells from *Glrb^{f/f}*-cKO

g Gephyrin GlyRβ ACE2 DAPI — Pearson's correlation — Gephyrin GLRB ACE2 DAPI — 50 μm — 2 μm

h GlyT2-EGFP GlyRβ CD13 DAPI — 5 μm / 10 μm — Pearson's correlation — Intensity (a.u) — CD13 GlyRβ EGFP — Distance (μm)

i **1** SMC EC Lumen / **2** SMC EC Lumen / **3** SMC EC Lumen / **4** SMC EC Lumen — 1 μm

j

k *Pdgfrβ-creER:Grin2d^{f/f}* × *Glrb^{f/f}* — NVC-deficiency × NVC-enhancement → *Pdgfrβ-creER:Grin2d^{f/f}:Glrb^{f/f}* NVC-neutrality

l △CBF (%) — Whisker stimulation — Seconds —
● Floxed control
● *Glrb^{f/f}*-cKO
● *Pdgfrβ-creER:Grin2d^{f/f}*
● *Pdgfrβ-creER:Grin2d^{f/f}:Glrb^{f/f}*
● CRS_*Glrb^{f/f}*
● CRS_*Glrb^{f/f}*-cKO

m △CBF (%) — ns — **** — 0.0128 — 0.0014 — 0.0239 — ns

a

tdTomato GluN2D DAPI

b

c

a

α-SMA expression level in SMCs (%)

ns

Grin2d^f/f
Grin2d^f/f-cKO

b

Smoothelin expression level in MCA (%)

ns

Grin2d^f/f
Grin2d^f/f-cKO

a

b

a

Ventral view

Anterior cerebral artery (ACA)

Middle cerebral artery (MCA)

Circle of Willis

Corticoamygdaloid artery (Coamg)

Internal carotid artery (ICA)

Ventral view
A single hemisphere

ACA

MCA

ICA

Coamg

SMA-creER:Ai14

b

Coronal view (from the anterior side)

LA

BAa

BAv

a Naive home cage
b HET0016 i.p injection home cage
c HET0016 BLA local injection home cage

a

i.p injection

b

c

a

OFT_40

b

OFT_40

Vanlandewijck et al. Nature (2018)