Neurovascular coupling conversely modulates emotion

Jiayu Ruan[1,2,3,4], Xinghua Quan[2,3,4], Yiyi Zhang[2,3,4], Huiqi Xie[2,3,4], Dongdong Zhang[2,3,4], Wentao Wang[2,3,4], Bingrui Zhao[2,3,4], Danping Lu[2,3,4], Jie-Min Jia[2,3,4*]

Affiliations

[1] College of Life Sciences, Zhejiang University, Hangzhou, Zhejiang, China.

[2] Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, China.

[3] Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, China.

[4] Laboratory of Neurovascular Biology, Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, China.

*Correspondence: jiajiemin@westlake.edu.cn

Summary

Emotion induces changes in regional cerebral blood flow (rCBF), a manifestation of neurovascular coupling (NVC). However, whether rCBF changes during NVC might, in turn, influence emotion is unclear. Here, by establishing arteriolar-optogenetic control of rCBF in free-moving mice, we demonstrate rCBF changes themselves induce emotional changes, and NVC actively modulates emotion. NVC deficiency impairs emotional flexibility, leading to contextually inappropriate emotional rigidity. Conversely, enhancing NVC counteracts this rigidity. To categorize inappropriate responses, we developed a predictive platform based on neural-activity patterns of the basolateral amygdala (BLA) in the mice exposed to a range of contexts. This platform revealed that NVC deficiency leads to BLA activations in a bipolar form, whereby predicting context-dependent hyper-reactivity and hypo-reactivity. Notably, this prediction was verified by actual physiological and behavioral responses. Mechanistically, neuronal mechanosensory receptor Piezo2 is essential for NVC-modulated emotion flexibility. Our findings establish NVC as an allostatic program for shaping emotion flexibility.

Introduction

Emotion changes over time, a key indicator of emotion flexibility.[1] Such flexibility involves the regulatory ability to adapt and respond to emotional experiences in a contextually appropriate manner.[2] Emotion regulation is of crucial significance for mental health and well-being.[3-5] Impairment in emotional flexibility can lead to emotion rigidity, which is one of the dimensions of emotion dysregulation and has recently been highlighted as a key contributor to psychiatric conditions.[2,6] Emotion rigidity manifests in two primary ways: restricted emotion ranges (reduced emodiversity) and context-inappropriate emotional responses.[2,7] Understanding the underlying molecular and cellular mechanisms of emotional flexibility and rigidity holds critical implications for elucidating basic biological explanations for resilience in the face of potential psychological trauma.[8,9]

Neuroimaging techniques, such as functional magnetic resonance imaging (fMRI), are instrumental not only in detecting emotion dysregulation but also in shedding light on the underlying neural mechanisms.[10,11] For instance, they have shown that hyperactivity or hypoactivity in the basolateral amygdala (BLA) can contribute to psychiatric conditions associated with emotion rigidity.[6,12] These neural activities are reflected indirectly through hemodynamic changes during neurovascular coupling (NVC), which forms the physiological basis for functional brain imaging.[13,14] Consequently, changes in

rCBF during NVC are often regarded as passive indicators.[15] Whether and

how they might actively regulate neural circuit functions is unclear *in vivo*.

Alternatively, despite over a century of research, the definitive functions of

NVC remain elusive.[16,17] This study seeks to explore this open question.

Recent proposals suggest NVC's involvement in modulating cognition and

neuropathologies[14,16,18] but the definitive role of local NVC in the BLA in

influencing emotion flexibility and rigidity is poorly understood. To investigate

this, two critical requirements must be met: first, achieving BLA-specific control

of NVC in freely moving animals; second, selectively blocking the

communication between neural and vascular systems without directly

disrupting either system. Thus far, to our knowledge, no experimental

paradigm meets these criteria, underscoring the urgent need to advance our

understanding of NVC's functional roles.

Arteriolar smooth muscle cells (aSMCs) are a critical component in the

neurovascular unit (NVU), effectively governing NVC,[19,20] and its contractility

can be successfully controlled by optogenetics on the surface of the cortical

cortex of head-fixed mice.[21,22] These reports have promised the feasibility of

stimulation of aSMC-covered arterioles in the BLA of free-moving mice by

combining with the currently well-developed fiber photometry technique.

Moreover, such photometry can simultaneously record hemodynamics in

deep-brain regions following intravenous injection of fluorescent dye.[23] This

paradigm would allow simultaneous control and monitoring of rCBF during

NVC in freely behaving mice, meeting the requirements to address the central

question of this study.

In parallel, independent mouse models are necessary to ascertain the

consistency of the effects derived from the aforementioned optical control of

NVC. For instance, a mouse model based on gene perturbations is essential.

The genetic model should precisely block neurovascular communication,

avoiding directly interrupting neural circuits and vascular integrity. However,

because conventional pharmacology and null knockout models lack necessary

cell-type specificity or directly interrupt neural circuitry,[24-27] meeting these

requirements has seemed challenging. Although recently developed

cell-type-specific NVC-deficient model may be able to address the central

question,[28,29] their effects on modulating emotion have yet to be explored.

More critically, these perturbations are global and cannot achieve

brain-region-specific NVC manipulations, thus introducing ambiguity.

Here, we established the paradigm for simultaneous monitoring and optical

controlling NVC in the BLA of free-moving mice. This paradigm, together with

three more fundamentally distinct NVC-deficient models, all commonly indicate

NVC deficiency impairs emotion flexibility. They exhibited one hallmark of

emotion rigidity (context-inappropriate emotional hyperactivity or hyporeactivity,

a bipolar-like manifestation). But they did not display the other hallmark

(reduction of emodiversity). Notably, enhancing NVC helped mice gain

resilience to emotion rigidity development following chronic stress.

Mechanistically, NVC deficiency caused hyperactivity or hypoactivity of BLA

neurons in a context-dependent manner, reminisced with similar

neuropathologies in the BLA of patients with psychiatric disorders.[6,12]

Furthermore, vasomechanics during NVC seemed to be critical. Overall, this

study further clarified the relationship between NVC and emotion, deepening

our understanding of their reciprocal interactions and suggesting new insights

into how to counteract the progression of emotion rigidity and intervene in

emotional dysregulation.

Results

Developing an assay to simultaneously record the BLA's rCBF, control it,

and monitor emotion-representing behaviors.

If rCBF changes have active roles, artificial manipulation of rCBF should be

able to actively change mouse emotion status, even in the innocuous context

of the home cage. To explore this, we first established an assay that integrated

measurement and manipulation of BLA's rCBF in free-moving mice, alongside

setting appropriate behavioral emotion indexes (EIs) from a range of potential

EIs (Figure 1A).

We began this study with defecation behavior for the following two

reasons. First, numerous studies in human research have established a

relationship between emotion change and defecation behavior.[30] Second, in

animal studies, the defecation behavior has long been used as one of the EIs

to reflect emotional changes induced by stress.[31-33]

Our findings further supported these conclusions by showing that BLA

activity was required for defecation behavior induced by acute tube stress

(Figure 1B-1C). Such stress induced every single mouse to defecate with no

exception (saline group in Figure 1C right). The BLA inactivation by

chemogenetics stopped three out of ten mice's defecation and reduced

average fecal production (Figure 1B and 1C). We additionally found that the

regular open field test (OFT) with minor stress intensity also induced

defecation behavior, similarly requiring BLA's activity (Figure 1D). These

results indicated that defecation behavior is an effective emotion representing

index, shortened as EI_Defecation here, which was associated with BLA

activity.

Moreover, we found that EI_Defecation exhibited a spatial preference

similar to the frequently reported EI of time spent in the center area of the OFT,

shortened as EI_Time-in-center (Figure 1E and 1F). Additionally,

EI_Defecation showed a stronger preference for the peripheral area (Figure

1F). Thus, we decided to use EI_Defecation as the primary EI, combined with

other EIs, for subsequent investigations.

After settling down the primary EI, we embarked on rCBF recording and

manipulations as designed in Figure 1A. First, we successfully recorded the

BLA's rCBF using fiber photometry recording of fluorescent signals in blood

plasma following venous injection of Rhodamine B (Figure 1G). We detected

the increasing signals along multiple injections of higher doses, demonstrating

effective recording of rCBF (Figure 1G-H. See Methods).

We next effectively suppressed rCBF in the BLA where arteriolar cells

expressed light-sensitive channelrhodopsin2 (ChR2) in *SMA-creER:Ai32* mice

(Figure 1A, 1I and Figure S1A). The specificity of this Cre-driver line to aSMCs

was validated previously (Figure S1B).[19] After exploring a series of protocols,

we finally identified one that can reliably and repeatedly suppress rCBF (Figure

1J shown in red line. See Methods). Once the light was on, rCBF became ~3%

instantly lower than baseline; it immediately returned to baseline once the light

was off (Figure 1J and 1K). These results indicated that we can reversibly

control rCBF in the BLA. All these data thus demonstrated a comprehensive

assay of measuring the BLA activity-associated behavioral EIs while

simultaneously and artificially changing rCBF in the BLA.

Optogenetic reduction of the BLA's rCBF likely de novo implants emotion

Next, we performed arteriolar optogenetics in the home cages, a context with

minimal external stress. We particularly chose a stable period to perform

arteriole optogenetics (pointed by arrows in Figure 1L). Within the 15 minutes

of light-On, the total amount of cerebral blood supply to the BLA was

profoundly reduced in *SMACreER:Ai32* mice compared to that of the ChR2

negative group (ChR2⁻, *SMACreER:Ai47* mice) that received all the same

handlings (Figure 1M and 1N).

To assess the EI_Defecation, all the previously produced feces in the

home cage were first removed before a total of 45 minutes observation period

(Figure 1M above). We defined the defecation rate as the percentage of mice

that produce at least one fecal pellet during a specified observation period.

Within this 45-minute observation period, ChR2$^-$ mice only exhibited a

defecation rate of 63.3% (9 out of 14 mice, Figure 1O left). In contrast, arteriole

optogenetics induced every single *SMA-creER:Ai32* mouse to defecate with no

exception, namely, a 100% defecation rate (Figure 1O middle), comparable to

the results of the tube stress assay (Figure 1O right).

Notably, ChR2$^+$ mice produced 1.14 fecal particles on average in 15

minutes, which was 3.7 times greater than that of ChR2$^-$ mice, achieving levels

not previously observed in the home cage environment (Figure 1P).

Unequivocally, this acute, distal stimulation in the BLA avoided any direct

stimulations on the local gastrointestinal tract that promotes bowel movement.

Thus, these calculations of EI_Defecation suggested that the artificial changes

in the BLA's rCBF under the context of home cage *de novo* implant emotion

represented by EI_Defecation.

This conclusion was further consolidated by the finding that artificial

changes in rCBF caused a robust increase in the BLA's neural activity (Figure

1Q and 1R). This result echoed the previous conclusion that the EI_Defecation

can reflect BLA activity-associated emotion changes (Figure 1B-1D).

Altogether, these findings demonstrated a cause-effect relationship between

rCBF change and emotion change.

NVC deficiency caused by arteriolar optogenetics promotes more rapid

initiation of defecation, a sign of hyper-reactivity.

Although OFT is known to increase the BLA's neural activity represented by

c-Fos expression level,[34] whether OFT can induce NVC, showing rCBF

increase in the BLA, has not been directly determined. The deep-brain optical

fiber photometry revealed rCBF increases once mice entered the open field

box, indicating that NVC occurred in the BLA (Figure 1S-T).

Optical suppression of such NVC in *SMA-creER:Ai32* mice resulted in a

hyperreactive-like response, as evidenced by a higher defecation rate, a trend

of producing more feces (Figure 1U-1W). In contrast, the total running distance

and EI_Time-in-center remained unchanged (Figure 1X and 1Y). Notably, NVC

deficiency caused profoundly shortened defecation latency (Figure 1W), which

was defined as the time from a mouse's introduction to a new environment of

OFT until the first fecal pellet is produced. The finding of faster latency

indicated that artificial NVC deficiency caused a more rapid initiation of

defecation, suggesting a hyper-reactive-like status. Altogether, these data

indicated that artificial suppression of NVC can induce a context-inappropriate

emotional response, a sign of impairment of emotional flexibility.

208 *Distinct models of NVC deficiency commonly eventuated a*

209 *hyper-reactivity to OFT with mild stress*

210 Recognizing that arteriolar optogenetics may inadvertently introduce unknown

211 side effects, we established three more types of NVC-deficient models. Each

212 type is based on different foundational principles (cell-type-specific genetic

213 knockout (cKO), local siRNA, and high-fat diet). If they all commonly induce

214 hyper-reactivity-like behavior, it would further strengthen the conclusion that

215 NVC deficiency impairs emotional flexibility.

216 Regarding the first cKO-type model, based on the previous finding of

217 neural–arteriolar smooth muscle cell junction (NsMJ, Figure 2A),[19] we

218 established four genetic tools (*SMA-creER:Grin1^f/f^*, *SMA-creER:Grin2d^f/f^*,

219 *SMA-creER:Grin1^f/f^:Grin2d^f/f^*, and *Pdgfrβ-creER:Grin2d^f/f^*), which are maximally

220 devoid of the direct disturbances of neural circuit or vascular integrity.[19]

221 Meanwhile, to have comparators, three more NVC-deficient lines were

222 generated by directly targeting neurons (*Emx1-cre:Ptgs2^f/f^*) and vascular

223 contractile proteins (*Pdgfrβ-creER:Acta2^f/f^*). *SMA-creER:Ptgs2^f/f^* was used as a

224 potential negative control. *Emx1-cre* was used to target nearly all excitatory

225 neurons;[35] *Ptgs2* encodes prostaglandin-endoperoxide synthase 2, also

226 known as cyclooxygenase-2 (COX2), producing potent vasodilators.[24]

227 *Pdgfrβ-creER* was used to target all mural cells that encompass SMCs and

228 capillary pericytes;[19] *Acta2* encodes smooth muscle alpha-actin (α-SMA), a

229 contractile protein critical for arteriolar contractility.[20]

Our prior work targeted the NMDAR core subunit GluN1 in aSMCs,[19] and

in this study, we identified *Grin2d* (Gene name of GluN2D) as the only

detectable non-core NMDAR subunit in aSMCs (Figure 2B and 2C). First,

RNAscope confirmed *Grin2d* presence in aSMCs (Figure 2D), and the probe's

specificity was first verified in cerebellum tissues (Figure S2A and S2B).[36] Next,

after the anit-GluN2D antibody was strictly verified (Figure S2C),[37] we found

that GluN2D protein was present in aSMCs of littermates but absent in those of

SMA-creER:Grin2d^{f/f} mice (Figure 2E-2G). Neither neural disruptions (Figure

2H) nor changes in contractile proteins (α-SMA and smoothelin) were

observed (Figure S2D and S2E). Also, we found GluN2D undetectable in

peripheral SMCs, which minimized systemic side effect concerns (Figure S3).

In addition, we also excluded tamoxifen administration's possible side effects

on NVC (Figure S4). All these examinations suggested that we neither directly

altered neural circuits nor caused any functional defects in myogenic tone.

Similarly, the effectiveness of *Ptgs2* and *Acta2* perturbation was carefully

verified (Figure S5). Functionally, we evaluated all these seven genetic cKO

lines by combining laser spectral contrast imaging (LSCI) measurements with

the traditional whisker stimulation paradigm.[38] Except for *SMA-creER:Ptgs2^{f/f}*

mice, the other six genetic cKO lines all effectively suppress NVC in the barrel

cortex (Figure S6).

We then chose *SMA-creER:Grin2d^{f/f}* as the exemplary line to further

determine whether BLA's NVC can also be suppressed (Figure 2I and 2J). The

basal levels of rCBF during the home cage between the two genotypes were

comparable (Figure 2I inserted panel). These observations suggested that the

genetic cKO lines may only compromise the amplitude of NVC without

affecting basal CBF supply to the BLA. We thus inferred that all these genetic

cKO lines may globally affect NVC, beyond BLA and barrel cortex.

Surprisingly, the EI_Defecation increased in whichever NVC-deficit lines,

indicating that they responded more intensely than typical to OFT (Figure 2K).

First, qualitative analysis revealed that all defecation rates were nearly twice

as high in all NVC deficient lines than in control mice (87.5%-100% vs.

50%-64.29%, percentiles in Figure 2K). In addition, the urination rate was

generally consistent with the defecation rate (Figure S7). Second, consistently,

quantitative analysis revealed that all NVC-deficient lines produced

approximately three times more fecal particles than controls on average

(Figure 2K). Collectively, these findings revealed a general negative correlation

between the NVC and EI_Defecation: the decreased NVC, the increased

EI_Defecation (Figure 2L).

Among these lines, the EI_Defecation observed in *Emx1-crePtgs2$^{f/f}$* and

Pdgfrb-creER:Acta2$^{f/f}$ lines even approached levels akin to those induced by

the tube stress assay (Figure 2K). We characterized this increased defecation

as a type of hyper-reactivity. Altogether, these genetic cKO lines of NVC

deficiency manifested a hyper-reactivity as observed optical suppression of

NVC in *SMACreER:Ai32* mice (Figure 1U-1W).

We took *SMA-creER:Grin2d$^{f/f}$* line as an exemplary line for further detailed

examinations. Similar to *SMA-creER:Ai32* mice (Figure 1W), we found a

profoundly shorter defecation latency (Figure 2M). Some *SMA-creER:Grin2d$^{f/f}$*

individuals immediately defecated once entering the box, which was never

observed in controls. In the 15-minute OFT test, *SMA-creER:Grin2d$^{f/f}$* mice

exhibited the strongest EI_Defecation response in the first 5 minutes,

producing nearly 12 times more feces than controls (Figure 2N, also see

vertical dashed lines in Figure 2I). This response decreased over the next two

5-minute periods, with fecal output 3.5 times higher in the second period and

3.0 times higher in the third period (Figure 2N). Again, there was no

difference in the total running distance and EI_Time-in-center (Figure S8),

consistent with the results from the optically stimulated ChR2$^+$ mice (Figure 1X

and 1Y).

Intriguingly, there was a neat, negative correlation between the sizes of

the OFT and the EI_Defecation of mice with normal NVC (Figure 2O and 2P).

Littermate control mice showed a gradual increase in EI_Defecation in OFT

boxes with decreasing floor areas of 40*40 cm², 35*35 cm², and 10*10 cm²

(Figure 2O and 2P). In great contrast, NVC-deficient mice exhibited an abrupt

increase in EI_Defecation rather than a gradual style. This finding indicated the

ability of mice to modulate the intensity of emotional responses in a

context-dependent manner. The abrupt plateau again indicated that NVC

deficiency led to context-inappropriate emotional responses, a status of

impaired emotional flexibility. (Figure 2P right).

Local NVC deficiency modeled by siRNA targeting aSMC-specific protein

In addition to the arteriolar optogenetics, we achieved another local model of

NVC deficiency. It is unavoidable to directly interrupt neurons if we choose

siRNA targeting GluN2D or GluN1, because they are also expressed in

neurons.[39] Therefore, we switched to determine whether there are molecular

targets that are not only GluN2D binding proteins but also exclusively

expressed in aSMC. We identified one protein called caldesmon[40] functionally

interacting with GluN2D to mediate NVC (Figure 2Q and 2R). We applied the

proteomic analysis for the protein complex that was pulled down by using the

anti-GluN2D antibody to incubate with the primary aSMC cell lysates. The

KEGG analysis of this complex revealed an enriched pathway of vascular

smooth muscle contraction pathway (Figure 2Q).

After their interaction was further verified by Western blot and

immunoprecipitation (Figure 2R), we designed and evaluated the effective

siRNAs targeting *Cald* (gene name of caldesmon. Figure 2S). Cisterna magna

injection of *Cald*-siRNAs indeed effectively suppressed NVC in the barrel

cortex following whisker stimulation (Figure 2T and Figure S9A). Thus, we

applied intracranial stereotactic injection of a strictly controlled amount of

siRNA (1 nM/100 nl) to achieve BLA-specific vascular perturbation so that

even the nearby regions were not affected (Figure 2U and 2V). For example,

ventromedial nucleus of the hypothalamus was not affected (Figure S9B-S9D).

More importantly, knocking down *Cald* in the BLA did not functionally affect the

NVC in the barrel cortex (Figure 2T). This would allow BLA-specific NVC

manipulation, which the Cre-lox system-mediated genetic model cannot

achieve.

Notably, local NVC deficiency in the BLA, despite intact NVC in the rest of

the brain regions, resulted in a 100% defecation rate, three times higher than

that of scrambled siRNA-injected mice (Figure 2K). Furthermore, a

combination of the local model and the genetic cKO model failed to increase

EI_Defecation further, suggesting that caldesmon acted in the same pathway

with GluN2D (Figure S9E and S9F).

Collectively, these results confirmed that both models of BLA-specific

NVC deficiency (optogenetic and siRNA-mediated), together with the global

cell-type-specific genetic cKO model, commonly induced emotional

hyper-reactivity (Figures 1U-1W and Figure 2K).

Obese mice induced by a high-fat diet show inappropriately intense

reactions to mild stress when blood-brain barrier (BBB) is still intact

We next extended to the obesity model induced by the high-fat diet (HFD),

which is known to induce NVC deficiency (Figure S10A and 10B).[41] Although

HFD-induced obesity involves intricate, multifaceted, and nonspecific effects, it

can serve as a fundamentally distinct way to interrupt NVC compared to the

above approaches. Importantly, obese humans with body mass index (BMI) >

30 are believed to have NVC deficiency.[42]

Notably, HFD-treated mice exhibited high EI_Defecation despite that they

consumed less amount of food than controls and produced fewer fecal pellets

in their home cage (Figure 2K, 2K inserted panel, and Figure S10C-S10E).

Even though they only received the mild stress resulting from OFT, their

EI_Defecation was even approaching the degree of tube stress with a 100%

defecation rate and an average above four fecal pellets (Figure 2K). These

data strongly demonstrated that the increased EI_Defecation was not due to

increased food intake but a context-dependent emotional response.

Moreover, we excluded the possibility that this emotional change is

attributed to BBB leakage (Figure S10F and S10G), despite BBB leakage

being frequently observed in HFD treatments and typically associated with

emotional changes.[18,43] This exclusion was based on the observations that

both NVC and emotion changes occurred in the fourth week of HFD when BBB

appeared intact (Figure 2K, Figure S10F, and 10G). This timeline aligned with

the commonly reported BBB leakage in the eighth week.[43]

In summary, we established four types of fundamentally distinct models of

NVC deficiency (BLA-targeting arteriole optogenetics, global genetic cKOs,

local siRNA, and HFD). In general, the overall physiological features, including

gastrointestinal (GI) system and BBB integrities, were unchanged, except for

body weight and fat content in the HFD group (Figure S10C and S10D). These

indicated such emotional change was not likely due to systemic physiological

side effects across these models (Figure S11).

Even though these four types of models might have entrained their own

side effects, their common effects on incurring hyper-reactivity to mild stress,

showing impaired emotional flexibility. This common effect suggested a

potentially universal link between NVC and emotional regulation. NVC may

serve as a necessary allostatic program for the brain to avoid generating an

overwhelmed emotional response to as mild stress.

NVC deficiency results in BLA hyperactivation, which is responsible for

emotional hyper-reactivity.

Next, we asked about the molecular and cellular mechanisms underpinning

NVC-deficiency-induced hyper-reactivity. We hypothesized that such

hyper-reactivity was due to BLA hyperactivation in NVC-deficient mice

because of the following two findings. First, local rCBF changes sufficiently

activated BLA neurons (Figure 1M, 1N, 1Q, and 1R). Second, BLA neural

activity was required for stress-induced defecation behavior (Figure 1B-1D). In

wild-type rodents, BLA's c-Fos is known to increase upon OFT,[34] so we

predicted more c-Fos increase in NVC-deficient mice than in littermates. We

examined two models, *SMA-creER:Grin2d*$^{f/f}$ and local *Cald* siRNA. Indeed,

both models exhibited hyperactivated BLA upon OFT (Figure 3A and 3B).

To further consolidate the above findings, we employed real-time

recording of rCBF and neural Ca^{2+} in the BLA. We transduced the BLA

excitatory neurons with *AAV-CaMKIIa-Gcamp8s*, which expressed a protein

probe to detect Ca^{2+} changes. We observed elevated neuronal Ca^{2+} signals in

the littermates with healthy NVC (Figure 3C and 3D), consistent with the c-Fos

results (Figure 3A and 3B).[34] Excitingly, we indeed observed higher Ca^{2+}

signals in the NVC-deficient mice, directly demonstrating hyperactivation,

aligning with c-Fos results (Figure 3A-3D). Altogether, these findings

demonstrated that the lower the capacity of NVC, the higher BLA activity.

Lastly, we chemogenetically inactivated the BLA of NVC-deficient mice

(*SMA-creER:Grin2d^{f/f}*). Indeed, deprivation of such hyperactivation stopped

the hyper-reactivity represented by the EI_Defecation (Figure 3E). This

provided a neural mechanism explaining how NVC deficiency led to

hyper-reactive emotional responses.

Neuronal Piezo2 perceives NVC deficits and is partially responsible for

the hyper-reactivity

During NVC, arterioles dilate and exert mechanical forces on surrounding

neurons. Removing NMDAR from aSMCs reduces arteriolar dilation [19]and thus

may reduce mechanical forces. If neurons possess mechanosensitive proteins

that detect these forces, depleting these proteins could mimic NVC deficits in

wild-type mice. This is because neurons would lose their ability to sense

arteriolar dilation, even though the dilation occurs normally.

406   As transcriptomics research profiled,[44] BLA neurons express the

407 mechanosensory protein Piezo2. We knocked it down via virus injection into

408 the BLA of *wild-type* mice (Figure 3F and Figure S12). Notably,

409 Piezo2-knockdown (KD) partially recapitulated the key features of NVC

410 deficient *SMA-creER:Grin2d$^{f/f}$* mice: hyperactivated of BLA (Figure 3G and 3H)

411 and partial increase in EI_Defecation (Figure 3I). These findings indicated that

412 Piezo2-KD mice presented slight hyper-reactivity upon OFT. This suggested

413 that Piezo2 in BLA excitatory neurons is partially responsible for

414 NVC-dependent emotion regulation.

416 ***Create a predictive platform to categorize the exaggerated emotional***

417 ***reponses mismatched with the actual stress level.***

418 To classify the extent of exaggerated emotional responses caused by NVC

419 deficiency, we developed a predictive platform that can define a spectrum of

420 stress levels. We created a library of BLA sections from seven groups of mice;

421 each group underwent distinct contexts that may differently activate BLA

422 (Figure 4A top, and Figure S13). This library encompassed a broad spectrum

423 of stress levels, ranging from nearly no stress (home cage) to strong stress

424 (predator-like threats. Figure 4A top panel). The latter were mimicked by

425 traditional looming assay and the newly established virtual reality of eagle

426 diving (VR$_{eagle}$). By using this library, we calculated a prediction index (PI)

427 based on the degree of similarity of the BLA neural activation patterns across

different contexts (Figure 4A top). The similarity calculation integrated two

dimensions of the BLA neural activity: anatomical distribution and their

intensity of c-Fos signals (Figure S13. See Methods).

An image analysis specialist performed unbiased classification for this

library by using this PI (See Methods). This specialist then revealed four stress

levels denoted as levels 0 to 3 (Figure 4A top). As expected, the home-cage

context generated the lowest stress (Level_0), and the predator-like-threat

context generated the highest stress (Level_3) (Figure 4A top). OFT is known

to cause mild stress and was positioned at Level_1, just right next to the

home-cage context. Moreover, the two contexts of OFT_40*40 and

OFT_35*35, known to be very similar to each other, were unbiasedly classified

together, self-reflecting the PI reliability (Figure 4A top).

To further test the PI's discriminative capacity, we generated a complex

version of the OFT (c-OFT) that can generate stronger stress. We introduced a

natural insecurity element associated with a bright-white floor. Prey C57/B6

mice are known to feel more stressed on white floors and more secure on

black floors.[45] We divided a chamber with a metal fence, with a white-floored

left and black-floored right sides (Figure S14A). Each mouse was placed on

the white side, allowing it to see but not cross into the darker side. The PI

ranked the context of c-OFT at Level_2, demonstrating its discriminative ability

because it distinguished the c-OFT one level higher than regular OFT and one

level lower than predator threats (Figure 4A top).

Interestingly, the tube-stress context was positioned at Level_2, lower

than predator threats. This could be possible because we did not force mice

but waited until they spontaneously entered the tube, and we only restrained

them for 15 minutes, substantially shorter than the typical ranges of up to 3

hours of restraint   time often reported.[46] This appropriate positioning of tube

stress again demonstrated the reliability of the PI of the prediction platform.

Regarding the highest Level_3, we updated the looming system (Figure

S14B) to the VR_{eagle} system (Figure S14C), which simulates an eagle-diving

scenario through a video played in a ballroom, creating a more realistic

environment. Their stress levels are theoretically the closest. Indeed, they

were ranked together based on the PI (Figure 4A top). Altogether, the PI can

appropriately rank the seven contexts in an order aligned with the known

stress levels (Figure 4A top and Figure S13).

By establishing this predictive platform, we revealed a regularity in the

relationship between stress levels and the patterns of BLA activity in mice with

normal NVC (Figure 4A top panel). Profoundly, NVC deficiency disrupted this

regularity, represented by the distorted c-Fos patterns shown in the exemplary

genetic cKO line of *SMA-creER:Grin2d^{f/f}* mice, which underwent the same

seven contexts (Figure 4A bottom). Only the home-cage context remained

unchanged between the two genotypes (Figure 4A bottom and Figure 4B),

aligned with previous findings of comparable basal characterizations (Figure 2I

inserted panel and Figure S11).

Once again, the PI was even able to reveal unbiasedly that the *Cald*-KD

and *SMA-creER:Grin2d^{f/f}* were the two most similar groups, which was exactly

as expected since caldesmon was first identified as working on the same

pathway with GluN2D (Figure 2Q, 2R, Figure S9E, and Figure S9F).

Additionally, this indicated that despite locally or globally impaired NVC, they

similarly hyper-activated the BLA neurons.

After the above validations, we proceeded to classify the mismatched

stress levels that NVC-deficient mice were likely to 'feel'. In the prediction

platform, the PI ranked "cKO_OFT_40" at Level_2 despite these mice

experiencing actual stress of Level_1, indicating hyper-reactivity, aligning with

observed behavioral responses (Figure 2K and Figure 4C). It appeared that

when *SMA-creER:Grin2d^{f/f}* mice entered a regular OFT, they hyper-reacted as

if they had entered an environment with the stress level of c-OFT or tube

stress. This prediction was consistent with previous actual EI_Defecation

observations (Figure 2K). Similarly, the local *Cald* KD treated group also had

consistency between our predictions and observed behaviors (Figure 2K and

Figure 4D). Thus, our predictions and actual observations commonly showed

that NVC-deficient mice hyper-responded to a mild context as if they

encountered more intense stress.

Intriguingly, arteriole optogenetics prompted a dramatic jump in the stress

levels by two levels: from the Level_0 to Level_2 (Figure 4E). Given that c-OFT

and tube stress at Level_2 both induce complex and mixed emotions, it is

possible that we optically implanted a mixed artificial emotion into the

free-behaving mice (Figure 1U-1Y).

Surprisingly, the PI predicted that *SMA-creER:Grin2d$^{f/f}$* mice would

hypo-respond to looming and VR$_{eagle}$, a stark and paradoxical contrast to the

predicted and behaviorally validated hyper-reactivity to OFT (Figure 2K and

2L). Instead of the expected further increase in stress levels, the

SMA-creER:Grin2d$^{f/f}$ mice showed a decrease, stepping down from Level_3

(red) to Level_2 (yellow) (Figures 4F, 4G, and S15A). This unexpected

prediction required actual physiological and behavioral validations.

NVC deficiency led to hypo-reactivity to innate fear threats

The complete sequence of an innate fear response in prey includes five steps,

integrating both cognitive and behavioral components.[47] Cognitively, it includes

predator recognition (Step_1), threat assessment (Step_2), and

decision-making on flight or alternative defenses (Step_3). Behaviorally, it

involves executing the chosen action (Step_4), relaxing defense, and returning

to a normal state once the threat diminishes (Step_5).[47] The last step matters

for the survival of individuals and species in a wild world because premature

relaxation may make them more susceptible to surprise attacks.[48]

Recognizing that a single EI_Defecation is insufficient to capture all these

dynamics occurring in these five steps (Figure S15B), we similarly

incorporated additional behavior-based EIs, such as changes in pupil size and

locomotion speed. While EI_Defecation remained a robust indicator of innate

fear, it failed to differentiate between the two genotypes as in the OFT context

(Figure S15B and 15C).

In the looming assay, mice would not receive looming stimulation until

they reached a locomotion speed of 13-15 cm/s, with no difference in basal

locomotion between the two genotypes (Figure 5A blue-shadow and Figure

S16A). The looming consistently induced instant typical freezing behavior in

both genotypes with speed reduced to 1-2 cm/s (freeze status) (Figure 5A

flesh-shadow). This implied that NVC-deficient mice had a normal innate fear

response from Step_1 to Step_4 (Figure 5A), which appeared contradictory to

the prediction (Figure 4F).

We continued to determine the Step_5 of relaxing defensive status. We

generated a heatmap that profiled locomotion speed dynamics until 21

seconds after threats because this time window would be long enough to cover

the most reported full duration of defensive states in *wild-type* mice.[49,50] This

heatmap revealed that NVC deficiency led to premature relaxation of defense

(Figure 5B). We found a substantially shorter period required for returning to

normal status (Figure 5B and 5C) as well as more rearing behaviors

representing a more relaxed state (Figure 5D). This finding of Step_5 was

consistent with the prediction (Figure 4F).

To assess the cognitive recognition of threats (Step_2), we upgraded the

looming threat to the VR$_{eagle}$ system, encompassing both visual and auditory

threats. We examined locomotion speed and pupil size, an involuntarily

controlled EI.[49,51] We recorded these two EIs simultaneously from awake,

head-fixed mice with semi-free moving on a floating ball (Figure S14C and

Supplementary Video 1). The two baselines of locomotion and pupil size

between the two genotypes were not different (Figure S16B and S16C).

Similarly, stimulation commenced only after the mice began moving.

When exposed to a distant-sounding eagle screech recording, both genotypes

displayed a similar transient decrease in locomotion speed without changes in

pupil size (Figure 5E, dashed and solid lines). These observations suggest that

the mice perceived the distant screech as a potential threat, yet not imminent

enough to cause harm. This supports the notion that Step 2, threat

assessment, was comparably executed by both genotypes.

Upon the eagle's dive following the screech, both genotypic mice

exhibited fear responses, evidenced by a similar increase in pupil size and

concurrent freezing behavior (arrows in Figure 5E and Figure S16D). This

response marked a second decrease in locomotion speed, deeper and more

prolonged than the initial reduction caused by the screech (Figure 5E). These

observations confirm normal visual processing and threat recognition from

Step 1 to Step 4.

Thus, we again investigated the last step of relaxing defense (Step_5).

The maximum of two independent EIs (largest pupil size and slowest

locomotion) took place almost at the same time point (Figure 5E, arrows). After

this time point, Step_5 began as pupil size started to decline, and locomotion

speed started to rise. We observed that Step_5 had a fast initial phase. We

defined this initial phase from the endpoint of the eagle's dive (brown line) to

the time point when the two dashed and solid curves came across, indicated

by magenta and blue vertical lines (Figure 5E). Profoundly, NVC deficiency led

to a shorter initial phase of Step_5 (Figure 5E and 5F), indicating potential

premature defense relaxation.

The two independent assays of looming and VR_{eagle} commonly revealed

that mice with normal NVC took an average of 12 seconds to fully relax from a

defensive state back to normal status (Figure 5G, blue dots). To our great

surprise, these two assays indicated almost identical time reductions between

the two genotypic mice (59.05% in looming and 59.61% in VR_{eagle}, Figure 5G).

These data immediately suggested that NVC deficiency prompted premature

relaxation by more than two times faster. This might be caused by lower BLA

activation. Therefore, these actual physiological and behavioral evaluations

validated the prediction of hypo-reactivity to predator-like threats (Figures 4G

and 5A-5G).

To exclude the effect of predator-type dependency, we designed a

paradigm–letting mice watch a video of an approaching cat, an independent

type of predator (Cat_{video}, Figure 5H-5L, and Figure S14D). First, we recorded

the behavioral responses of awake, head-fixed mice on a treadmill repeatedly

stimulated by Cat_{video} with three trials (Figure 5H and Supplementary Video 2).

As expected, normal mice showed typical defensive behaviors such as retreat

and tail rattling (Figure 5I-5L), demonstrating that Cat_{video} was an effective

stimulus. Notably, most NVC-deficient mice exhibited no retreat behavior at all

(Figure 5J and 5K). Even among the rest of the small subset of this group, the

intensity of defensive behaviors was also remarkably reduced (Figure 5H, and

5J-5L). These behavioral characterizations suggest that the found

hypo-defensiveness was not predator-type-dependent (Figure 5E-5L).

Altogether, these actual behavioral evaluations corroborated the prediction.

This consensus between the predicted and observatory results showed that

NVC was required for shaping contextually appropriate emotional responses–

emotional flexibility–by preventing the bipolar form of hyper- and

hypo-responsiveness.

NVC-deficient mice are predicted to hypo-react to real social stress,

verified by actual behavioral evaluation.

In addition to the above virtual and designated threats, we challenged the

prediction platform with a substantially more complex context, which is more

dynamically interactive and real mouse-mouse social stress. We decided to

generate social stress by cohabitation with selected socially aggressive CD1

mice in a fresh home cage (Figure 6A). In a single-time cohabitation for a brief

10 minutes, we observed an average of 20 seconds of fighting, accounting for

only 4% of the total time (Figure 6B). This was substantially shorter than

reported, which may be due to using fresh home cages, new to both mice.[52]

The home cage was separated into small and large subregions, but mice were

still allowed to freely inter-entrance between them (Figure 6A).

In the prediction platform, the PI positioned control mice experienced this

social stress at Level_1 (green), more intense than in a home cage but less

than tube stress (Figure 6C). Intriguingly, this was different in

SMA-creER:Grin2d^{f/f} mice (Figure 6D). They were predicted as if they

perceived a stress level of Level_0 (blue), like in a home cage. This predicted

a hypo-responsiveness to real social stress, which was again subsequently

validated by actual behavioral evaluations.

Due to fear, although both two genotypes exhibited a natural tendency to

avoid staying in the same smaller subregion with CD1, *SMA-creER:Grin2d^{f/f}*

mice showed less avoidance to CD1 than littermate controls (50 sec vs 100

sec out of a total of 600 seconds, Figure 6E). More strikingly, 98% of control

mice actively fled away from CD1 after fights, and so did only 37% of

NVC-deficient mice (Figure 6F). Consistent with the mice flee rate, we

normalized it to total fighting events. There were 37 and 19 fights between the

8 controls with CD1 and 10 NVC-deficient mice with CD1, respectively, among

which CD1 actively triggered the most fights (Figure 6G). NVC-deficient mice

exhibited a significantly smaller percentage of fleeing rate than controls (Figure

6G). The sniffing behaviors did not differ between the two genotypes,

indicating comparable basal exploratory ability (Figure 6H).

The consensus between the predicted results (Figure 6C and 6D) and

actual behavioral verifications (Figure 6E-6G) strongly supported the

prediction platform as reliable and the conclusion that NVC can modulate

emotional status.

NVC-enhancement resists emotion rigidity caused by chronic restraint

stress

Chronic restraint stress (CRS) is known to impair NVC and lead to emotion

rigidity, like anxiety- and depression-like status.[53,54] If NVC is truly capable of

modulating emotion, NVC-enhancement should be able to influence the

development of emotion dysregulation following CRS.

Next, we embarked on developing an NVC-enhancement model, as no

such model currently exists to our knowledge. Given that disrupting a cation

channel, such as the NMDA receptor (NMDAR), suppresses NVC (Figure 2I),

[19] we hypothesized that disrupting an opposite anion channel might enhance

NVC. Our experimental results confirmed this hypothesis (Figure 7).

Previous transcriptomic studies showed that perivascular mural cells

express ligand-gating chloride channels, for instance, glycine receptor subunit

b (GlyRβ, encoded by gene *Glrb*).[55] It seemed highly expressed in venular

SMCs (vSMCs, Figure S17A).[55] We confirmed *Glrb* puncta and GlyRβ protein

clusters with more frequency found in vSMCs than aSMCs (Figure 7A-7D). We

conditionally   knocked   out   *Glrb*   specifically   in   mural   cells   using

Pdgfrβ-creER:Glrb^{f/f} (Figure 7C, 7E, and 7F) and further identified GlyRβ

colocalizing with gephyrin *in vitro* (Figure 7G), a scaffold protein often

segregated in inhibitory postsynaptic sites,[56] suggesting that GlyRβ could be

clustered on the postjunctional membrane of glycinergic-NsMJ. This

speculation was further supported by the finding that frequent direct contact of

neural axons onto venular SMCs by both confocal laser scanning microscope

(Figure 7H) and electron microscope (Figure 7I).

Notably, the whisker stimulation assay revealed that *Pdgfrβ-creER:Glrb^{f/f}*

mice indeed enhanced NVC, approximately 37.25% higher than averaging

from littermate controls (Figure 7J and 7K). Together, these data suggested

that functional glycinergic-NsMJ suppressed physiological NVC, so removing it

would lead to enhanced NVC.

We hypothesized that if *Pdgfrβ-creER:Glrb^{f/f}* mice with NVC enhanced

can counteract the NVC deficiency caused by *Grin2d-cKO,* they may also be

able to counteract hyper-reactive behaviors. Indeed, double knockout mice not

only resisted hemodynamic attenuation during NVC (Figure 7J and 7K) but

also resisted the hyper-responsiveness by comparing the two groups of

Pdgfrβ-creER:Grin2d^{f/f} single cKO mice and *Pdgfrβ-creER:Grin2d^{f/f}:Glrb^{f/f}*

double cKO mice upon regular OFT (Figure 2K and Figure 7L). These

observations led us to test further whether *Pdgfrβ-creER:Glrb^{f/f}* is able to

counteract emotion rigidity, such as depressive-like emotional distress caused

by CRS. We applied CRS to *Pdgfrβ-creER:Glrb^{f/f}* mice to test this hypothesis.

After the *wild-type* mice were forced to stay in the tube for 2 hours every

day for 28 consecutive days, they showed higher anxiety-like status, reflected

by a shorter period of staying in the open arm of the elevated maze (Figure 7M

left). In addition, they produced stools with changed quality after CRS (Figure

7M right), confirming CRS induction was effective in this study.[30]

Notably, *Pdgfrb-creER:Glrb^{f/f}* mice completely resisted the changes in

defecation behaviors and the quality of stool (Figure 7N-7P). They appeared to

be even as normal as sham mice never received CRS (Figure 7N-7P).

Alternatively, the forced swimming test (FST) was believed to reflect the

degree of desperation in depression-like mice according to their duration of

immobilization in the water. More strikingly, FST revealed a complete

resistance to the development of depression-like status (Figure 7Q). The

constant movements of *Pdgfrb-creER:Glrb^{f/f}* mice in water were unlikely due to

the increased basal level of locomotion because they exhibited normal

locomotion in the OFT (Figure S18). In contrast, *Pdgfrb-creER:Glrb^{f/f}* mice

exhibited the lightest body weight over the course of CRS (Figure S17B),

suggesting that NVC-enhancement only partially resisted emotional distress.

CRS is also known to impair BBB, which has been frequently associated

with later emotional distress.[18,46,57] An alternative possibility could be that

Pdgfrb-creER:Glrb^{f/f} mice may be through protecting BBB leakage to resist the

effects of CRS. Because we restrained mice for a 4-week period shorter than

reported[57] and also because we observed no IgG leakage in parenchyma, we

conclude that BBB seemed intact under the current experimental settings

(Figure S17C and S17D).

All these data emphasized that NVC may be a future therapeutic target for

psychiatric disorders that are associated with emotion rigidity. Taking all NVC

manipulations through this study together, we assert that NVC is a built-in

program to modulate emotion.

Discussion

This study identified modification of emotion as one of the functions of NVC,

directly addressing the longstanding question regarding the definitive function

of NVC.[16-18] Unexpectedly, NVC actively engaged in the central nervous

system computation for programming innate fear emotional response, a

defensive system that promotes survival (Figures 4-5). NVC actively shaped

emotional flexibility (Figures 2, 4-5), a factor crucial for mental health and

well-being. For example, in high-stress situations such as emergency

response or combat, the ability to regulate emotions can significantly impact

decision-making and survival.[58] These findings may provide an explanation for

the evolutionary conservation of NVC across species, as appropriate emotion

regulation can be a matter of life or death in the animal kingdom and, at times,

in human society as well. More importantly, this study offers a fundamental

perspective on emotion regulation programs, promising a new avenue of

intervention in psychopathological progression by enhancing NVC, thus

complementing current strategies.

Neurovascular coupling—What next ?

Although enormous amounts of academic effects have been put into

measuring human's rCBF and understanding NVC mechanisms by Mosso,

Roy, and Sherrington since the late 19th century, [59,60] the NVC research has

come to a new door—what next? By knocking on this door, this study provides

an unexpected renewal for this topic using state-of-the-art technologies that

combine arteriolar optogenetics, real-time recordings of rCBF and neural Ca^{2+}

in the deep brain regions of free-moving animals, cell-type-specific gene

perturbations, and local vascular siRNA.

By exploring the BLA-emotion axis as an exemplary case, we propose that

NVC can reversely regulate brain functions. It merits further investigations of

this concept beyond emotion regulation in other brain regions where NVC also

occurs. For instance, robust NVC occurs in the anterior cingulate cortex (ACC)

when the ACC is involved in many brain functions and disease conditions,

such as the affection and cognitive aspects of pain and social exclusion,[61] the

transitions between wakefulness and sleep and its relations with insomnia

disorder,[62] and psychogenic erectile dysfunction (PED).[63,64] Then, whether

NVC has regulatory roles in the pathophysiology of chronic pain, insomnia,

and PED merits further investigation. This study opens the next door to the

NVC research field.

Defecation behavior is a robust and sensitive EI

The amygdala is known to project to Barrington directly, a defecation and

urination center located in the brainstem of mice.[65,66] Therefore, we

extrapolated that BLA may send emotion-related neural signals to Barrington's

nucleus, resulting in defecation. Suppression of BLA neurons by chemogenetic

inhibition suggested the functional relationship between BLA and defecation

(Figure 1C and 1D).

All seven contexts in this study can readily cause robust defecation

behavior compared to the home cage (Figure 2K, 2L, and Figure S15B),

mirrored with robust increases in BLA's c-Fos while with diverse patterns

(Figure 4A and Figure S15A). This diversity indicates that although different

contexts engaged distinct neuron subgroups, they all can eventually converge

to defecation behavior (Figure 2K). This may explain why the EI_Defecation is

sensitive EI while with limited discriminative ability (Figure 2K and Figure

S15B). This may also explain why defecation behavior can be commonly

induced despite those opposite changes in rCBF following OFT and optical

stimulation, resulting in distinct c-Fos patterns (Figure S15B).

We propose that the EI_Defecation can be used for fast and/or pilot

screening readouts because it is an assessment with robustness, simplicity,

and objectivity. The analysis of defecation locations in the OFT (Figure 1E and

1F) highlights the potency of using the EI_Defecation has been overlooked.

We recommend a broader scope in the mood research field to include the

EI_Defecation. For example, in depression research, recently, the two classic

tests of tail suspension and forced swimming have recently been under heated

debates.[67,68] The findings following chronic restrain stress in this study

implicate that the EI_Defecation might be a robust, alternative evaluation of

depressed status.

In addition, the EI_Defecation was prone to differentiate the impact of

NVC under the contexts of mild stress but failed to do so under the contexts of

strong stress (Figure S15B). However, this could be due to this study's limited

diversity of contexts or due to the investigation scope of this study. Multiple

physiological and behavioral emotion indexes (e.g. pupil size, freeze/flight) are

necessary, depending on specific context and scientific questions.

Neurovascular calculations should be considered in the basic

psychiatric study.

This study opens up a new field of how NVC actively participates and helps in

fine calculations during neural computations, highlighting not merely at the

synaptic level[69] but also at the neurovascular level. The neurovascular

computation seemed intrinsically programmed to optimize the kinetics of a

single cycle of fear response to predators, specifically determining the timing of

the transition into a defensive state and the subsequent relaxation from that

state. This neurovascular calculation should be taken into account by the basic

affective study of emotion flexibility.

This study emphasizes the critical role of bidirectional neurovascular

communications in a functional entity. Such bidirectional communications may

provide a unique biological insight into the next-generation algorithm

development for artificial intelligence if future robots need to include

multi-hierarchical modulatory modes of emotion designs.[70]

Neurovascular coupling (NVC) and vasculo-neuronal coupling (VNC)

function in a loop manner that is interlinked

NVC could, conversely, immediately send various types of signals back to

neurons when vasodilation is induced by neural activity. Among these signals,

a mechanic signal attributed to vasodilation could be one of them. This

backward vascular-neural talk is referred to as vasculo-neuronal coupling

(NVC). [71-73] We believe that the neurons that cause NVC may or may not be

the same group of neurons that perceive vasomechanics. If they are two

separate groups, future work could focus on whether the two groups also

communicate with each other. Thus, the NVC–VNC loop, sequentially

programmed, eventually satisfies the bidirectional communications between

the nervous and cerebrovascular systems.

NVC is generally thought to oversupply blood flow where needed.[16] How

neurons sense and respond to vasodilation during NVC has emerged as a new

research field.[71,73] This study broke the NVC-VNC loop by blocking VNC part

(Figure 3G-3I). Despite blocking VNC by knocking down neuronal Piezo2-KD

in *wild-type* mice, a similar trend was found in EI_Defecation as if in

NVC-deficient mice (Figure 2K and Figure 3I). This suggested that this loop

functioned as an interlinked program. Future investigation is needed to

determine the neuron subgroups that control vasodilation and sense

vasodilation.

Enhancing NVC may provide a new intervention

To our knowledge, this study, for the first time, developed a genetic model that

enhanced NVC: *Pdgfrβ-creER:Glrb$^{f/f}$*. This is likely achieved by blocking

glycinergic NsMJ, meaning glycinergic axon innervating vessels. During NVC,

there are believed to be negative and positive factors for regulating

vasodiameter.[74] We extrapolate glycinergic NsMJ as a negative factor. In

Pdgfrβ-creER:Glrb$^{f/f}$ mice, such a negative factor was removed, so the NVC

was enhanced. Nevertheless, the detailed molecular mechanism underlying

such potential negative regulation needs to be elucidated.

Previous work reported that glycinergic axon terminals can be observed

both in BLA and involved in depression regulation.[75,76] Enhanced NVC in

Pdgfrβ-creER:Glrb$^{f/f}$ mice may prevent hyper- or hypo-neural activations in the

BLA so that it can completely counteract emotion rigidity caused by chronic

stress. If the mural cell-specific virus is established, interruptions of GlyRβ will

have implications for developing new therapeutic approaches for psychiatric

conditions.

In addition, the vascular-cell transcriptomic study revealed that GlyRβ

transcripts are more expressed in mural cells of venules compared to capillary

pericytes and aSMCs.[55] This study also opens a new avenue for studying the

contribution of venules in brain functions.

In summary, this study is based on comprehensive models to expand our

understanding of the relationship between NVC and emotional regulation,

highlighting NVC actively participating in programming emotional

flexibility. The study also agrees with a growing consensus that the function of

NVC is far beyond the regulation of CBF and is involved in regulating cognition

and emotion at different scales.

Acknowledgments

We thank Y. T. and X. H. for sharing *Nestin-cre* and *Emx1-cre* mice. We thank

Y. Z. for sharing psychological knowledge to facilitate our understanding of

data. We thank Y. N. for the KEGG map drawing. We thank Westlake

University Laboratory Animal Resource Centre for guiding behaviour assays.

We thank the Microscopy Core Facility of Westlake University for imaging

guidance. We thank the Mass Spectrometry & Metabolomics Core Facility

(MSMCF) of Westlake University for mass data collection. We thank the Flow

Cytometry Core Facility (FCCF) for cell sorting assistance. J.-M.J.

acknowledges the support from Westlake Laboratory of Life Sciences and

Biomedicine (Project No. 202309002), Westlake University startup funding, the

Westlake Education Foundation, Zhejiang, Key R&D Program of Zhejiang

(2024SSYS0031), and the National Natural Science Foundation of China

(project No. 32170961).

Author contributions

J.-M.J. conceived the project, designed the experiments, and supervised this

study. J.R. did primary experiments, obtained and organized the majority of

original data, and wrote the manuscript draft. X.Q. participated in behavioural

assay design and operations, siRNA cisterna magna injection, gut health

inspection assay, BBB integrity inspection, caldesmon expression inspection,

cluster analysis of similarity index, and wrote Mann-Whitney U test R code.

Y.Z.wrote the main MATLAB code for PI. H.X. participated in the looming

assay and VR$_{eagle}$ assay, as well as in blood pressure measurement and

long-term whisker stimulation. D.Z. provided primer pairs of *Grin1*, raw data of

electron microscope, and *SMA-creER:Grin2d$^{f/f}$* OFT_40*40 data. W.W.

contributed to the writing of MATLAB code with Y.Z.. B.Z. participated in the

genotyping of mice and cell sorting. D.L. helped the mouse brain section.

Declaration of interests

The authors declare no competing interests.

Supplemental information

Supplemental figures: Figures S1-S18

Supplemental tables: Tables S1-S3

Supplemental videos: Videos S1-S2

Figure legends

Figure 1. Artificial reduction of rCBF *de novo* initiates emotion. (A) The schematic diagram of the potential route linking neural signals initiated by arteriolar optogenetics in the BLA and ended in defecation behavior. (B-C) c-Fos$^+$ neuron density and fecal pellet quantification after experiencing 15-min tube stress in the two groups of mice with IP injection of saline or C21. Both groups received *AAV2/5-hM4D (Gi)-EGFP* virus injection in their BLA. One dot represents one mouse. (D) Fecal pellet quantification for two groups of mice received the same treatment as C, except for the experience of 15-min OFT instead of tube stress (N =9 mice for each group). (E) The spatial map of defecation locations in the OFT from a total of 27 mice. (F) The percentage of defecation and traveling time behavior occurred in the center versus peripheral

areas in the OFT box (N =27 mice). (G) Optical photometry recording of

fluorescent intensity in the BLA of mouse received multiple orbital injections of

four doses of Rhodamine B. (H) Linear relationship between Rhodamine B

concentration and plasma fluorescence intensity in G. (I) The timeline of the

experiment of arteriolar optogenetic control and recording rCBF in the BLA. (J)

Temporal dynamics of rCBF changes before, during, and after arteriolar

optogenetics with four different stimulation protocols (see Methods). (K) The

quantified area under the curve (AUC) for J. Each dot represents one mouse.

(L) The fluorescence decay curve in the BLA of awake mice after receiving

orbital Rhodamine B injection. (M) Quantification of rCBF changes in the BLA

across three consecutive 15-minute recordings. Arteriolar optogenetics was

applied during the second 15-minute recording. mice. (N) The quantification of

AUC for M. N =8 *SMACreER:Ai47* (ChR2$^-$) mice, N =7, *SMACreER:Ai32*

(ChR2$^+$) mice. (O-P) Defecation behavior quantifications upon optical and tube

stress stimulations. The defecation rates of mice within a 45-minute period in

the home cage or tube stress assay in O. The fecal production every 15 min in

P. N =14, 7, and 10 mice for each group, respectively. (Q) Representative

image of co-staining of c-Fos and NeuN in the BLA from ChR2$^-$ or ChR2$^+$ mice.

(R) The quantification for Q. (S) Real-time rCBF changes in the BLA of mice

before and during OFT in S. N =4 mice. (T) The quantification for S. (U-Y)

Quantifications of behaviors of ChR2$^-$ or ChR2$^+$ mice while receiving optical

stimulation during OFT. Quantifications of defecation rate, feical production

and defecation latency in U-W. Quantifications of the total running distance

and the time spent in the center of the OFT box in X and Y.   N =4 mice for

each group. Data are presented as mean ± SEM. See also Figure S1, Table

S1 and S2. Student's t-test was used for differential analysis of all data, except

for fecal particles, using the Mann-Whitney U test. ****P<0.0001.

Figure 2. Establishment of various genetic lines of NVC deficiency and

evaluations of their emotional responsiveness (A) The schematic of NsMJ

in NVU. (B) mRNA expressions of NMDAR subunits in cerebral parenchymal

aSMCs. (C) The quantification for B. N =3 independent experiments. (D)

RNAscope of *Grin2d* transcripts (green dots) in *SMA-creER:Ai14* mouse brain

section. The white arrow points to *Grin2d*mRNA present in aSMC (red), the

relative location shown in the inserted histogram. (E-F) GluN2D and

Smoothelin protein expressions in arterioles. The quantification in F (N =11-12

aSMCs from 4 arterioles of 4 mice for each genotype). (G) GluN2D

expressions in the sorted aSMCs (EGFP$^+$) from *SMA-creER:Ai47* and

SMA-creER:Ai47:Grin2d$^{f/f}$ mice. N =3 biological replicates. (H) GluN2D

expressions in the cerebral tissue. N =3 biological replicates. (I) rCBF changes

in the BLA before and during OFT. The vertical black dash line marks OFT

begin. The vertical colour dash lines mark defecation time point. Inserted,

basal levels of rCBF at home cage. N =4 mice for each group. Figure 1S and

Figure 2I shared the same raw data for *Grin2d$^{f/f}$* mice. (J) Quantification of the

1st 5-min rCBF area under the curve (AUC) for I. (K) Total fecal production

within a 15-min OFT or tube stress, and 24 hrs home cage. Underlined

numbers mean defecation rate (%), under which the numbers mean fecal

particles averaged from total mice for each group. The mouse number (N) for

each group is shown along the genotype in the X axis. Inserted panel is the

food consumption of control/HFD mice in 24 hrs (N =16 mice for each group,

each dot presents the mean value of one cage contains 4 mice). (L) Heatmap

of the relationship between the extent of NVC and EI_defecation. (M)

Cumulative defecation latency from 14 mice for each genotype subjected to

the OFT. (N) Fecal production within every 5 min during the 15-minute OFT. N

=14 and 16 for each genotype). (O) Defecation rates after OFTs with different

floor areas and after tube test. (P) Fecal particles for D. Raw data of *Grin2d$^{f/f}$*

and *SMA-creER:Grin2d$^{f/f}$* was shared in K-P. N =12-19 mice shown in O.

(Q)Top 10 KEGG pathways of the Glu2D-binding protein complex. N =3

experiments. (R) Western blottings of GluN2D and Caldesmon using samples

pulled down by anti-GluN2D. N =3 independent experiments. (S) *Cald* mRNA

expression in the cortex lysis with cisterna magna injection of *Cald* siRNA or

scramble, N =3 biological replicates. (T) LSCI imaging of CBF changes of mice

with the same treatment as in U before receiving whisker stimulation. Insert,

quantification for mean changes in CBF. N =5-6 mice for each group. (U) The

immunofluorescence staining of Caldesmon of mice received local BLA siRNA

injection. (V) Quantifications for U and for Figure S9A. Dots represent mouse

numbers. Student's t-test was used for differential analysis of all data, except

fecal particles used the Mann-Whitney U test. Data are presented as mean ±

SEM. See also Table S1-S3 and Figures S2-S11. ****$P<0.0001$.

Figure3.   Neural   mechanism   for   NVC   deficiency   induced

hyper-responsiveness. (A) Representative images of c-Fos immunostaining

in the BLA mice received OFT. (B) Quantifications of c-Fos$^+$ neuron density for

A. Each dot represents one mouse. (C) Ca^{2+} changes in BLA before and

during OFT. (D) Quantification for C. (E) The fecal production of

SMA-creER:Grin2d$^{f/f}$ mice with hM4D (Gi) expressed in the BLA excitatory

neurons. N =8 mice received IP injection of saline, and N =7 mice received

C21. (F) Diagram of *Piezo2* KD virus infection range in BLA. (G-I) Before and

during OFT, real-time neural activity recordings in the BLA where stereotactic

injection with scramble or *Piezo2* siRNA as well as quantification of defecation

behaviors in OFT. Ca^{2+} changes in G and H. Fecal production in I. N =8 mice

for each group. Student's t-test was used for differential analysis of all data,

except fecal particles used the Mann-Whitney U test. Data are presented as

mean ± SEM. See also Table S1, S2, and Figure S12. ****$P<0.0001$.

Figure 4. Prediction platform for estimating false emotions. (A) Prediction

index-based unbiased classification of stress levels for seven contexts.   (B-G)

The predicted position of every single treatment in the unbiased classification

system. The stress levels are denoted as Level_0 (blue), Level_1 (green),

Level_2 (yellow), and Level_3 (brown). See also Table S1, Figure S13 and

Figure S14.

Figure 5. NVC deficiency prompted an early relaxation of defense upon

predator-like attack. (A-D) Looming assay. Temporal profile of locomotion

speed dynamics of the two genotypic mice before (cyan) and during (yellow)

stimulation in A. Heatmap that profiled the locomotion speed dynamics for a

total of 30 seconds in B. The yellow box marks the 3-sec looming duration.

Cumulative time used for relaxation from a freezing status following looming

stimulation in C. Rearing index of each group in D. N =15 mice for each group.

(E-G) VR_{eagle} stimulation. Temporal changes of the normalized pupil size and

locomotion speed during and after VR_{eagle} stimulation in E. Eagle screech only

present during the first 1.5s (blue) and eagle started to dive at 1.5 second.

Representative images of VReagle video and mouse eye with pupil clear

visualized. The statistical data of the initial phase of step_5 for E in F. The

statistical data of pupil size full-recovery latency and looming locomotion

full-recovery latency in G. For VR_{eagle}, N =7 mice for each group; for Looming,

N =15 mice for each group. (H-L) Cat_{video} assay. The locomotion pattern during

the Cat_{video} assay in H. Black bar marks retreat behavior; magenta lines

indicate tail rattling. Two vertical dash lines highlight the time points when the

cat appeared and pounced. Mouse numbering is on the left. Quantification of

retreat behaviors of *Grin2d^{f/f}* mice before and after cat appearance in I. Retreat

rate of two groups upon in Cat$_{video}$ stimulation in J. Retreat rate referred to the

fraction of mice that exhibited at least one retreat behavior of total mice. Dark

green means retreat exhibited; light green means no retreat. Cumulative

retreat time curve in K. Quantification of tail rattling times in L. N =12 mice for

each group. Student's t-test was used for differential analysis of all data. Data

are presented as mean ± SEM. See also Table S1, Figure S14-S16, and

Supplementary Video 1 and 2.

1008

Figure 6. NVC-deficient mice are predicted to hypo-respond to real social

stress. (A) Picture of a home cage from a side view. (B) The total fighting time.

(C-D) The predicted positions of the two genotypic miceexperience social

stress. (E) The total time of staying with CD1 mouse in the small space. (F-G)

The flee rate of *Grin2d^{f/f}* and *SMA-creER:Grin2d^{f/f}* mice (counted as per mouse)

after fighting in F, and the normalized flee rate (counted as per fight event)

after each fighting in G. (H) The sniffing times. N =8 and 10 mice for *Grin2d^{f/f}*

and *SMA-creER:Grin2d^{f/f}*). Student's t-test was used for differential analysis of

all data, except fecal particles used the Mann-Whitney U test. Data are

presented as mean ± SEM. See also Table S1.

1019

1020 **Figure 7. The NVC-enhancement model resisted emotion rigidity**

1021 **following chronic restraint stress.** (A)RNAscope of *Glrb* transcripts in brain

slices of *Pdgfrβ-creER:Ai14* mice. (B) Quantification of mRNA signal dots in

aSMCs and vSMCs for A. N =41 aSMCs and N =32 vSMCs from a total of 3

brain sections of each mouse. N =4 mice for each group. (C) GlyRβ protein

expression in arterioles and venules from two genotypic mice. (D)

Quantification of GlyRβ clusters in aSMCs and vSMCs in C. N =14 arterioles,

N =19 venules. n=6 mice for each group. (E) Quantification of GlyRβ

immunosignal intensity for C (n=6 mice for each group). (F) Semi-quantitative

analysis of *Glrb* mRNA by RT-PCR for the sorted PDGFRβ^{+} cells in the

cerebral cortex. (G) Co-immunostaining of GlyRβ and Gephyrin in HUvSMC

cell line. The white arrow marks their colocalizations. Insert, the statistical

co-localization analysis (N =4 cells). (H) Representative image of glycinergic

NsMJ in a venule. Insert, histogram analysis of the location indicated by the

white arrow. (I) Serial SEM sections of the ascending venule in mouse cerebral

cortex. The white arrows point to axon's boutons. (J) LSCI imaging of CBF

changes in the barrel cortex upon whisker stimulation. (K) The mean changes

in CBF for J. N =5-6 for each group. (L) The fecal production of mice in the

OFT and tube stress assay. Defecation rate (underline) and mean fecal

production. N =10-15 mice for each group. Figure 7J-7L used the raw data.

Data of of *Pdgfrβ-creER:Grin2d$^{f/f}$* mice from Figure 2K and Figure S6A. (M)

Quantification of time spent in open arms of EPM and the Modified Bristol

Score on the last day of CRS. N =6 control mice, N =13 mice for CRS group).

(N) Representative images of stools with different Modified Bristol Scores.

(O-P) Progressive changes following CRS. Dynamic changes in Modified

Bristol scores in O. Fecal particle number produced within a period of 60 min in

P. N =6 mice for $Glrb^{f/f}$_Sham group, N =14 mice for $Glrb^{f/f}$ group received CRS,

N =7 mice for $Pdgfr\beta$-$creER$:$Glrb^{f/f}$ group received CRS. The evaluation was

performed before the assay and once a week during CRS. (Q) Quantification

of immobility time in the 5-minute FST. N =7 mice for $Glrb^{f/f}$ group, N =4 mice

for $Pdgfr\beta$-$creER$:$Glrb^{f/f}$ group. ****P<0.0001. Student's t-test was used for

differential analysis of all data, except fecal particles used the Mann-Whitney U

test. Data are presented as mean ± SEM. See also Tables S1-S3, Figure S17

and Figure S18. ****P<0.0001.

1054

1055 **Methods**

1056

Animal and animal care

All mouse procedures complied with the Institutional Animal Care and Use

Committee (IACUC) guidelines of the School of Life Sciences, Westlake

University. All mice were housed in the Westlake University Laboratory Animal

Resource Centre with standard animal facilities, which provided a 12-hour

light/dark cycle at 25 °C. No more than six mice were fed in each cage, and

standard chow/water was provided to the mice *ad libitum*. All the genetic

information of mice is in Table S1.

1065

Tube stress assay (ARS & CRS)

A 50ml tube was used to restrain the mice[46,77] to perform acute restraint stress (ARS) and chronic restraint stress (CRS). The tube cover and wall will be drilled to ensure normal breathing. For ARS, the mice were restrained in the tube for 15 min, and then the fecal production was measured. For the ARS plus chemogenetic manipulation assay, the C21 intraperitoneal injection was 45 min before the tube stress assay. After the assay, mice were fed alone in a clean new cage for 90 min and then used for brain extraction to detect c-Fos expression. For CRS, the mice were restrained in the tube for 120 min/day, constant for 28 days, and fecal production and Modified Bristol Score (Add "0" to indicate no defecation) was measured at the end of each seven days within 60 minutes in a home cage box.

Immunofluorescence staining

For Immunofluorescence (IF) staining, after anesthesia with 1% pentobarbital sodium, the mice were perfused with 1X PBS (5 min) and 4% PFA in 1X PBS (5 min) through the left ventricle to empty blood and fix tissue. Afterwards, tissue was extracted and fixed by 4% PFA in 1X PBS for 6 hours at 4 °C. For samples like the brain and peripheral organs, samples were sliced into 50 μm (Leica, Histo-VT1200S, DEU). Then, treated samples with 1% Triton X-100 (Diamond, Cat# A110694-0500, CHN) and 0.25% NDS (Jackson Resources, Cat# 017-000-121, USA) in 1X PBS for 60 min at room temperature. After that,

samples were rewashed three times with 1X PBS.

In the staining process, the primary antibody diluted with 0.1% Triton

X-100 in 1X PBS was used to incubate samples overnight at 4 °C. After the

incubation, samples were washed with 1X PBS 10 min (3 replicates). The

corresponding secondary fluorescence-conjugated antibody diluted with 0.1%

Triton X-100 in 1X PBS was used to incubate samples for 120 min at room

temperature. After the incubation, samples were washed with 1X PBS for 10

min (3 replicates). After that, 1000X diluted DAPI (Beyotime, Cat# C1002,

CHN) was used for nucleus staining. Finally, samples were covered by a glass

slice with the mounting medium (SouthernBiotech, 0100-01, USA). Zeiss

LSM800 (Zeiss, LSM800, DEU) microscopy was used for photography. A list of

antibodies is in Table S2.

1100

Open field test

The setup of open field test (OFT) referred to published papers.[34,78] Each

mouse was placed in the central area of a 40*40 cm, 35*35 cm, or 10*10 cm

box for the OFT. The camera recording was 15 min. For the chemogenetic

manipulation assay, the C21 intraperitoneal injection was 45 min before the

OFT. After the assay, mice were fed alone in a clean new cage for 90 min and

then used for brain extraction to detect c-Fos expression. For the BLA CBF

and calcium signal recording, optic fibers were implanted at BLA (AP: -1.6 mm;

ML: +/-3.45 mm; DV: -4.0 mm) with the injection of hSyn-Gcamp-8f virus

(*SMA-creER:Ai32* mice do not need virus injection. Detailed information about

virus mentioned later) two weeks before the assay. Before the formal assay,

three days of adaptation were performed. On the day of the assay, orbital

injection of Rhodamine B 100 µl 5 mg/ml 15 minutes before the recording. 405

nm light was used for bassline control, 488 nm light recording for the Gcamp-8f

signal (For *SMA-creER:Ai32* mice, using 473 nm light for optogenetic

stimulation), and 580 nm light recording for the Rhodamine B

signal. Recording sustained 15 (For CBF and Ca^{2+} signal recoding)/45 minutes

(For *SMA-creER:Ai32* mice).

1119

1120 **Tamoxifen administration for mice**

1121 Tamoxifen (MCE, Cat# HY-13757A, USA) was dissolved in corn oil (Aladdin,

1122 Cat# C116023-500ml, CHN) and orally administered to mice (10 mg/ml, 200 µl

1123 per mouse per day). For *Acta2* cKO, tamoxifen was given from P21 to P24.

1124 Otherwise, tamoxifen was orally administered to mice from P60 to P63.

1125

Optogenetic stimulation and recording

Optogenetic stimulation referred published papers.[23,79] *SMA-creER:Ai32* and

SMA-creER:Ai47 mice were used for optogenetic stimulation. Tamoxifen was

administered to mice 18 days before tha assay. Then, optic fibers were

implanted at BLA (AP: -1.6 mm; ML: +/-3.45 mm; DV: -4.0 mm) two weeks

before the assay. Before the formal assay, three days of adaptation were

performed. On the day of the assay, orbital injection of Rhodamine B 100 µl 5

mg/ml (Sigma, Cat# R9379-1G, USA) 15 minutes before the recording. 405

nm light was used for bassline control, 580 nm light recording for Rhodamine B

signal for 45 min, and 473 nm light for optogenetic stimulation was performed

at the 2^{nd} 15 min of the 45 min recording. The parameters of the stimulation

protocol included the stimulation period, interval period, recording period, total

frequency, and laser power. The optimal protocol shown in Figure 1J is 85 ms

stimulation, 5 ms interval, 5 ms recording, 10 Hz frequency, 0.939 mW laser

power. During the 45-minute recording, mice always stayed in their home cage,

but all fecal particles were removed before the assay. After 45 min recording,

fecal particles were recorded.

1143

Cell culture

SMA-creER:Ai47 P0 mice were anesthetized with ice. Surgical tweezers were

used to quickly extract the cortex and then transfer it into the 0.25% Trypsin

(Thermo Scientific, Cat# 25200072, USA) for 15 min at 37 °C. Next, FBS

(Gibico, Cat# 10099, USA) was added to block digestion. Then, cells were

filted by a 40 µm filter (Falcon, Cat# 352340, CHN). Finally, cells were

centrifuged 1,600*g at 4 °C for 5 min and resuspended by SMCM (ScienCell,

Cat# 1101, USA) in T25 flasks (Corning, Cat# 430639, USA). When cells

reached approximately 80% full of the bottom surface of T25 flasks, tamoxifen

was administrated to cells. Tamoxifen was dissolved in DMSO (Solarbio, Cat#

D8371, CHN), and the final concentration of tamoxifen was 5 μM. After a

four-day culture, EGFP in cells were expressed. Cells were sorted by MA-900

(Sony, FCCF-MBR, JPN) to obtain EGFP$^+$ cells. Sorted cells were cultured in

SMCM for further assay. The purity of cells sorted by the same method was

verified in a paper published by our laboratory.[19] For data in Figure 7F,

samples were directly sorted from Pdgfrβ$^+$ cells in brain lysis by using

Anti-Pdgfrβ antibody (without cell culture). Then, Pdgfrβ$^+$ cells were quickly

resuspended by Trizol (Sangon Biotec, Cat# B511311-0100, CHN) for RNA

extraction. A list of antibodies is in Table S2. In Figure 7G, Human Umbilical

Vein Smooth Muscle Cells (HU-vSMC, Sciencell, Cat# 8020, USA) were used

for cell culture.

1165

RT-PCR

For cultured cells, after obtaining pure sorted EGFP$^+$ cells from the cortex,

cells were cultured in 6 cm dishes (Corning, Cat# 430166, USA) with SMCM.

After cells reached around 90% density, Trizol was used for the RNA extraction.

For brain tissue, after mice were anesthetized with ice, the cortex or

cerebellum of the mice was directly taken out, and TRIzol was used for RNA

extraction. After extraction, all RNA samples were reverse transcribed to cDNA

(50 ng/μl) using HiScript III 1st Strand cDNA Synthesis Kit (Vazyme, Cat#

R312-02, CHN). For the Real-Time Polymerase Chain Reaction (RT-PCR),

each reaction contained 10 μl of Green Taq Mix (Vazyme, Cat# P131-AA,

CHN), 2 µl of cDNA, 2 µl of primer pairs (10 µM), and 6 µl of RNAase-free

water. Agarose gel electrophoresis used 2 % agarose gel (BIOWESTE, Cat#

111860, ESP) and ran at 220 V for 30 min. RT-PCR gel images were obtained

from a Tanon imager (Tanon, 1600 Geil imager, CHN). A list of primer pairs is

in Table S3.

1181

RNAscope in situ hybridisation

Horizontal sections (15 µm) of tamoxifen administrated *SMA-creER:Ai14* (for

Grin2d), *Pdgfrβ-creER:Ai14* (for *Glrb*) and C57/B6 *wild-type* (for *Grin2d* and

Grin1 probe validation) mice brains were obtained by CM1950 (Lecia,

Histo-CM1950, DEU). All RNA scope *in situ* hybridisation productions were

obtained from ACDbio, including RNAscope Multiplex Fluorescent Reagent Kit

v2 (ACDbio, Cat# 323100, USA), *Grin2d* probe (ACDbio, Cat# 425951-C3,

USA), *Grin1* probe (ACDbio, Cat# 431611, USA), and *Glrb* probe (ACDbio,

Cat# 1225261-C1, USA). All operations were carried out following the reagent

instructions of ACDbio company and the published paper.[19,80]

1192

Western-bolt

For tissue sample collection, samples were ground with RIPA (Thermo

Scientific, Cat# 89901, USA) in the tube on ice for 30 min. For cells, after they

reached around 90% density, the RIPA was used for protein extraction on ice

for 30 min. Protease Inhibitor Cocktail (CWBIO, Cat# CW22008, CHN) was

added into RIPA to prevent protease. After that, samples were centrifuged at

12,000*g at 4 °C for 5 min, and the supernatant was collected. Concentrations

of proteins were measured by Micro BCA Protein Assay kit (CWBIO, Cat#

CW2011S, CHN), and all samples were boiled with SDS-PAGE loading buffer

(CWBIO, Cat# CW0027S, CHN) for 10 min at 100 °C. 8% SDS-polyacrylamide

electrophoresis gels were used for gel running. After 90 min, gels running at

120 V (80 V for the first 30 min), gels were transferred to 0.45 µm PVDF

membranes (Millipore, Cat# IPVH00010, USA) at 400 mA for 120 min. After

transfer, PVDF membranes were blocked by 5% non-fat milk in 1X TBST

buffer for 60 min at room temperature with a mild horizontal shake.

After the block, the primary antibody diluted with 5% non-fat milk in 1X

TBST buffer was used to incubate the membrane overnight at 4 °C with a mild

horizontal shake. After the incubation, the membrane was washed by 1X TBST

10min (3 replicates). The corresponding secondary HRP-conjugated antibody

diluted with 5% non-fat milk in 1X TBST buffer was used to incubate the

membrane for 45 minutes at room temperature with a mild horizontal shake.

After the incubation, the membrane was washed with 1X TBST for 10 min (3

replicates). ECL kits (Abbkine, Cat# BMU102-CN, CHN) and gel image system

(GE, GEL Imaging System AI680RGB v.2.0, USA) were used for exposure

photography. A list of antibodies is in Table S2.

1218

Whisker stimulation

Mice were anaesthesia with 1% pentobarbital sodium and open the scalp, then

mice stable fixated on the holder. The electro-motor-driven brush stimulated

the whisker longitudinally for 1 minute at a frequency of 12 Hz. The principle of

laser speckle contrast imaging has been described in the previous paper,[24]

and the operation referred to in the published paper.[38,81,82] Laser Speckle

Image System (RWD, RFSLI ZW/RFSLI III, CHN) was used to record the CBF

change in the ROI of the opposite barrel cortex of mice. The recording

parameters were 512*512 resolution, 10 FPS, and 2-minute recording.

1228

Immunoprecipitation assay

tdTomato[+] sorted aSMCs from P0 *SMA-creER:Ai14* mice were used for

lentivirus infection and further immunoprecipitation (IP) assay. Sorted cells

were cultured in 6 cm dishes with SMCM. After cells reach around 90% density,

add lentivirus (Lenti-EF1-EGFP-CMV-grin2d-3XFlag-WPRE) into the culture

and keep for one week to over-express GluN2D. MA-900 sorted out EGFP and

tdTomato double-positive cells, which were lentivirus-infected aSMCs. These

cells were then cultured in 10 cm dishes. After cells reached around 90%

density, cells were collected by RIPA reagent. The Protein G Magnetic beads

kit (Epizyme, Cat# YJ002, CHN) was used for the immunoprecipitation assay.

GluN2D and IgG control antibodies were mixed with Protein G Magnetic beads

to pull down GluN2D and its related protein. Pull-down products were verified

by western-bolt and analyzed through mass spectrometry. A list of antibodies

is in Table S3.

1243

siRNA cisternal magna/BLA local injection

Cisterna magna injection protocol referred to Cannula Implantation into the

cisterna magna of Rodents (https://doi.org/10.3791/57378) and paper

published by our laboratory.[19] 1 nM siRNA was mixed with FITC (10 mg/ml,

Sigma, FD40S-1G, USA) for injection (20 µl in total). The scalp was opened 10

min after the injection, and the microscope was used to detect the FITC signal

(Zeiss, Axio Zoom v16, DEU) in perivascular spaces. Only successfully

injected mice remain. Whisker stimulation was performed 48 hours after siRNA

injection, followed by OFT_40*40. As for BLA local siRNA injection, 150nl of

1nM siRNA was injected in two sides of BLA (AP: -1.55 mm; ML: +/-3.40 mm;

DV: -4.00 mm), and OFT_40*40 was performed 48 hours after siRNA injection.

Brain slice collection was performed 90 minutes after OFT for IF staining.

Sequence of siRNA can be found in the Table S3.

1257

High-fat diet challenge assay

Set up of High-fat diet challenge referred to published papers.[41,43,83] Mice were

given high-fat food (SYSE, Cat# PD6001, CHN) for four weeks. The weight of

mice and food consumption were measured before the assay and at the end of

each week. The home cage fecal production was measured using the same

method mentioned in the "Home cage monitor assay" part. OFT_40*40 assay

and whisker stimulation assay were performed after four weeks of HFD

challenge.

1266

Home cage monitor assay

The blood pressure of mice was measured by Kent Noninvasive blood

pressure measurement system for animals (Kent/Biopac, USA) before the

mice were put into the home cage. Each home cage only contains four mice. In

each cycle (three days), 60 g of food, 340 g of water (including bottle weight),

and 100 g of padding were provided. Weight of mice, food/water consumption,

fecal particles, and urination amount (counted by increased padding weight)

were measured by calculating the mean data. Three cycles of monitoring were

performed for each home cage. After the assay, mice were fed alone in a clean

new cage for 90 min and then used for brain extraction to detect c-Fos

expression. The intestine system inspection was performed after three cycles

of monitor.

1279

BBB leakage inspection (evens blue and IgG)

Assay setup referred to published papers.[43,84] 2% evens blue (Merck,

Cat#E2129, USA) dissolved in saline was used for orbital injection (4 ul/g).

After 45 min, the mouse was anesthetized, and the brain was extracted for

appearance inspection for BBB leakage using a microscope (Zeiss, Axio Zoom

v16, DEU). The brain was weighted and then homogenized using 1 ml of 50%

trichloroacetic acid (Merck, Cat#190780, USA). Then, samples were

centrifuged at 10,000*g at 4 °C for 20 min, supernatants were collected, and

absorbance detection at 620 nm was performed. For IgG inspection methods,

see the *Immunofluorescence staining* part above.

1290

1291 ***Colorectal inspection***

1292 The ENDOQ Colonoscopy system took representative images of colorectal

1293 tissue. A professional physician performed scoring of the endoscopic images in

1294 a double-blind manner to ensure unbiased assessment.

1295

Hematoxylin and eosin staining

For hematoxylin and eosin (HE) staining, after anesthesia with 1%

pentobarbital sodium, the mice were perfused with 1X PBS (5 min) and 4%

PFA in 1X PBS (5 min) through the left ventricle to empty blood and fix tissue.

Afterward, tissue was extracted and fixed by 4% PFA in 1X PBS for 6 hours at

4 °C. After a paraffin block, tissues were sliced into 8 µm sections and

examined following HE staining.

1303

In vivo virus injection

rAAV2/4-Glyt2-EGFP-WPRE-hGHpolyA (5E+12VG/ml) was used to label

glycinergic neurons. Two weeks before the assay, bilateral medullary reticular

formation injection (500 nl on each side, AP: -3.5 mm; ML: +/-0.84 mm; DV:

-2.8   mm)   was   administered.[75]

AAV2/5-mCaMKIIa-hM4D(Gi)-EGFP-ER2-WPRE-pA(2.30E+12VG/ml)   and

AAV2/5-mCaMKIIa-EGFP-ER2-WPRE-pA(2.30E+12VG/ml) were used for the

chemogenetical manipulation. Two weeks before the assay, bilateral amygdala

injection (150 nl on each side, AP: -1.6 mm; ML: +/-3.45 mm; DV: -4.1 mm)

was administered.[85] AAV2/9-hSyn-jGCaMP8f-WPRE-pA was used for the

neural calcium measurement. Two weeks before the assay, bilateral amygdala

injection (150 nl on each side, AP: -1.6 mm; ML: +/-3.45 mm; DV: -4.1 mm)

was   administered.   rAAV2/9-CamKIIa-3xflag-5'miR-30a-shRNA

(piezo2)-3'-miR30a-WPREs and rAAV2/9-CamKIIa-3xflag-5'miR-30a-shRNA

(scramble)-3'-miR30a-WPREs were used for the BLA local *Piezo2* KD. Two

weeks before the assay, bilateral amygdala injection (100 nl Piezo-2 virus +

100 nl jGCaMP8f virus) on each side, AP: -1.6 mm; ML: +/-3.45 mm; DV: -4.1

mm) was administered.

1322

Prediction Index (PI)

The Prediction Index (PI) is based on the Structural Similarity Index (SSIM)

and the cluster analysis of SSIM (See Statistical analysis software and code

part for details). In brief, translating BLA's c-Fos positive cells location and

density with different contexts into SSIM. Then, the cluster analysis was used

to classify the SSIM of different contexts to build a library of PI. To predict a

subject's "real" stress level, the SSIM of this subject will be clustered together

with the PI library to locate the subject's stress level.

1331

Complex-Open field test (c-OFT)

The complex-open field test (c-OFT) setup referred to a published paper.[45,86]

Each mouse was placed in the central area of the left part of the c-OFT box.

One side of the c-OFT box has a white bottom (the mouse is put on this side)

and a black bottom on the other. The camera recording was 20 min. For the

analysis, the box was divided into three parts (left area, central area, and right

area) to analyze the preference of mice for metal fence (right area close to the

metal fence). After the assay, mice were fed alone in a clean new cage for 90

minutes and then used for brain extraction to detect c-Fos expression.

1341

Looming assay

The assay was carried out in a 35 cm*35 cm OFT box. The settings of the

assay were based on the published paper.[49] Modifications include:

Modifications include:

• mice will be put in the OFT box to adapt for 10 minutes before starting

the assay;

• we only use a single camera at the top of the box to record;

• we stimulate the mice when they move and are not close to the box

wall rather than continuously stimulating them.

Each mouse performs three trials of stimulations. After the assay, mice

were fed alone in a clean new cage for 90 min and then used for brain

extraction to detect c-Fos expression.

1354

VR_{eagle} assay

Assay set up referred published paper.[49] The mouse head was fixed with a

fixator two weeks before the assay. The mice received daily adaptive training

on the VR system one week before the assay. On the day of the assay, the

mice were fixed on the VR system and given 5 min of adaptation. Unless

stimulated, the screen in front of the mouse will always play the regular

landscape video. The stimulation of the eagle lasts for three seconds. The

eagle dives down from far to near and plays the voice of the eagle at the

beginning of the stimulation. There were five trials of stimulation in total, and

the criterion for the first stimulation was that the mice moved for the first time

after 5 min of adaptation. The criteria for subsequent stimulation were that the

last stimulation ended for at least 2 min, and mice resumed movement. The

assay ended 2 min after the last stimulation. The pupil changes

(Supplementary Video 1) and mice's locomotion speed were recorded

simultaneously. After the assay, mice were fed alone in a clean new cage for

90 min and then used for brain extraction to detect c-Fos expression.

1371

Cat_{video} stimulation assay

The assay was carried out in a 40*40 cm OFT box. The mouse head was fixed

with a fixator and trained to fit the treadmill two weeks before the assay. On the

day of the assay, the mice were anesthetized with isoflurane and fixed on the

treadmill. After the mouse wakes up, play the video of cat hunting on the iPad

Air (Apple. Cat# 3rd generation, USA) in front of the mouse (Supplementary

Video 2). The stimulation video played three consecutive times, each

stimulation lasting 30 seconds. A camera directly above was used to record the

assay.

1381

Home cage social assay

The setup of the Home cage social assay referred to published papers.[52,86-89]

CD1 mice never exposed to $Grin2d^{f/f}$ or $SMA\text{-}creER\text{:}Grin2d^{f/f}$ mice were used

for the assay. Before the assay, simultaneously put a single $Grin2d^{f/f}$ or

$SMA\text{-}creER\text{:}Grin2d^{f/f}$ mouse and a CD1 mouse into a cage. The cage contains

new padding and a metal lid on the top. One side of the metal lid is used to

place food and water bottles, so the height is lower than the other. A camera

was used to record two mice's behaviours for 10 min. After the assay, the mice

were separated again. $Grin2d^{f/f}$ or $SMA\text{-}creER\text{:}Grin2d^{f/f}$ mice were fed alone in

a clean new cage for 90 min and then used for brain extraction to detect c-Fos

expression.

1393

1394   **Elevated plus-maze test**

1395   The setup of elevated plus-maze (EPM) referred to published papers.[80,90] The

1396 EPM comprises two open arms (65*5 cm) and two close arms (65*5 cm). Each

1397 mouse was placed in the centre, facing one of the two open arms, and allowed

1398 to explore for 5 min. Time spent in open arms was recorded.

1399

Serial block-face scanning electron microscopy image

The specimen was obtained in the same batch as the previously published

paper from our laboratory.[19] The details in the "SBF-SEM imaging" section of

the methods in that paper. In brief, the ascending venule was identified using

X-ray microscopy (Zeiss, XRM, Xradia 520 Versa, DEU) at ×4 and 80 kV

magnification. The specimen adhered to an aluminium metal pin using

conductive epoxy (Chemtronics, CW2400, USA). Then, it was further precisely

trimmed into a block of 600*600*700 μm^3 using a glass knife and a

right-angled diamond knife (DiATOME, Trim 90, CHE) after coating with

gold-palladium (Leica, EM ACE200, DEU) and a final trim using a 3View

ultramicrotome (Gatan, USA). Image acquisition was performed using a BSE

detector on a Carl Zeiss Gemini 300 scanning electron microscope at 2 kV

voltage and approximately 1.8*10–3 mBar in high-vacuum mode with focus

charge compensation. The specimens were sectioned at a 50–75 nm

thickness and a cutting speed of 0.1 mm/s.

1415

1416 ***Force Swimming Test (FST)***

1417 The Force Swimming Test (FST) referred to a published paper.[91] In brief, mice

1418 were put into the glass water tank with 25 °C clear water. Measuring the

1419 swimming time of mice in 5 minutes and then putting the mouse back in the

1420 home cage.

1421

Statistical analysis software and code

All the image processing and analysis of fluorescence intensity used Zeiss Zen

Lite (v3.4) and ImageJ (v1.53u). All whisker stimulation original data were

measured by RWD LSIS-self software (v5.0). All the original behavioral data

for OFT/c-OFT/EPM were recorded using Noldus. Data recording and analysis

of Looming and VR$_{eagle}$ assay was performed using self-compiled codes

provided by Bayonne company. All statistical data were processed by

Numbers (v12.2). Unless otherwise specified, Student's t-test was used for

differential analysis on all data that conform to a normal distribution.

Non-continuous data that does not conform to a normal distribution, such as

fecal particles, signal dots in vessels, and rearing index were analyzed using

the Mann-Whitney U test. All figures were made by Prism (v9.4.1) and Adobe

Illustrator 2024 (v28.3). CBF peak mean value, heat maps, mass spectrometry

data inclusion criteria, cross-correlation, and prediction index (based on the

Structural Similarity Index, SSIM) were analyzed by self-compiled MATLAB

codes. Besides, the KEGG pathways, cluster analysis of SSIM, and

Mann-Whitney U test were analyzed using self-compiled R code. All available

codes   can   be   found   at

https://github.com/JialabEleven/Functional_hyperemia_regulates_emotion.

1441

1442   **References**

1443

1.   Bortolon, C., Chen, S., and Bonanno, G.A. (2023). Components of

emotion regulation flexibility and psychosis: The association between

psychosis-proneness and context sensitivity. Br J Clin Psychol *62*,

82-95. 10.1111/bjc.12395.

2.   D'Agostino, A., Covanti, S., Rossi Monti, M., and Starcevic, V. (2017).

Reconsidering Emotion Dysregulation. Psychiatr Q *88*, 807-825.

10.1007/s11126-017-9499-6.

3.   Roelofs, K., Bramson, B., and Toni, I. (2023). A neurocognitive theory of

flexible emotion control: The role of the lateral frontal pole in emotion

regulation. Ann N Y Acad Sci *1525*, 28-40. 10.1111/nyas.15003.

4.   McCracken, L.M. (2024). Psychological Flexibility, Chronic Pain, and

Health.   Annu   Rev   Psychol   *75*,   601-624.

10.1146/annurev-psych-020223-124335.

5.   Addolorato, G., Mirijello, A., D'Angelo, C., Leggio, L., Ferrulli, A.,

Abenavoli, L., Vonghia, L., Cardone, S., Leso, V., Cossari, A., et al.

(2008). State and trait anxiety and depression in patients affected by

gastrointestinal diseases: psychometric evaluation of 1641 patients

referred to an internal medicine outpatient setting. Int J Clin Pract *62*,

1063-1069. 10.1111/j.1742-1241.2008.01763.x.

6.   Easdale-Cheele, T., Parlatini, V., Cortese, S., and Bellato, A. (2024). A

Narrative Review of the Efficacy of Interventions for Emotional

Dysregulation, and Underlying Bio-Psycho-Social Factors. Brain Sci *14*.

10.3390/brainsci14050453.

7.   Thompson, R.A. (2011). Methods and measures in developmental

emotions research: some assembly required. J Exp Child Psychol *110*,

275-285. 10.1016/j.jecp.2011.04.007.

8.   De Prisco, M., Oliva, V., Fico, G., Radua, J., Grande, I., Roberto, N.,

Anmella, G., Hidalgo-Mazzei, D., Fornaro, M., de Bartolomeis, A., et al.

(2023). Emotion dysregulation in bipolar disorder compared to other

mental illnesses: a systematic review and meta-analysis. Psychol Med

53, 7484-7503. 10.1017/S003329172300243X.

9.   Geng, H., Xu, P., Aleman, A., Qin, S., and Luo, Y.J. (2024). Dynamic

Organization of Large-scale Functional Brain Networks Supports

Interactions Between Emotion and Executive Control. Neurosci Bull *40*,

981-991. 10.1007/s12264-023-01168-w.

10.   Szaflarski, J.P., Allendorfer, J.B., Heyse, H., Mendoza, L., Szaflarski,

B.A., and Cohen, N. (2014). Functional MRI of facial emotion

processing in left temporal lobe epilepsy. Epilepsy Behav *32*, 92-99.

10.1016/j.yebeh.2014.01.012.

11.   Etkin, A., and Wager, T.D. (2007). Functional neuroimaging of anxiety: a

meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. Am J Psychiatry *164*, 1476-1488. 10.1176/appi.ajp.2007.07030504.

12. Phillips, M.L., Drevets, W.C., Rauch, S.L., and Lane, R. (2003). Neurobiology of emotion perception II: Implications for major psychiatric disorders. Biol Psychiatry *54*, 515-528. 10.1016/s0006-3223(03)00171-9.

13. Mathias, E.J., Kenny, A., Plank, M.J., and David, T. (2018). Integrated models of neurovascular coupling and BOLD signals: Responses for varying neural activations. Neuroimage *174*, 69-86. 10.1016/j.neuroimage.2018.03.010.

14. Iadecola, C. (2017). The Neurovascular Unit Coming of Age: A Journey through Neurovascular Coupling in Health and Disease. Neuron *96*, 17-42. 10.1016/j.neuron.2017.07.030.

15. Pico-Perez, M., Radua, J., Steward, T., Menchon, J.M., and Soriano-Mas, C. (2017). Emotion regulation in mood and anxiety disorders: A meta-analysis of fMRI cognitive reappraisal studies. Prog Neuropsychopharmacol Biol Psychiatry *79*, 96-104. 10.1016/j.pnpbp.2017.06.001.

16. Drew, P.J. (2022). Neurovascular coupling: motive unknown. Trends Neurosci *45*, 809-819. 10.1016/j.tins.2022.08.004.

17. Masamoto, K., Hirase, H., Yamada, K., and Kanno, I. (2016).

Neurovascular coupling-What next? Prog Brain Res *225*, 269-272.

10.1016/bs.pbr.2016.03.007.

18.   Dion-Albert, L., Dudek, K.A., Russo, S.J., Campbell, M., and Menard, C.

(2023). Neurovascular adaptations modulating cognition, mood, and

stress responses. Trends Neurosci *46*, 276-292.

10.1016/j.tins.2023.01.005.

19.   Zhang, D., Ruan, J., Peng, S., Li, J., Hu, X., Zhang, Y., Zhang, T., Ge, Y.,

Zhu, Z., Xiao, X., et al. (2024). Synaptic-like transmission between

neural axons and arteriolar smooth muscle cells drives cerebral

neurovascular coupling. Nat Neurosci *27*, 232-248.

10.1038/s41593-023-01515-0.

20.   Hill, R.A., Tong, L., Yuan, P., Murikinati, S., Gupta, S., and Grutzendler,

J. (2015). Regional Blood Flow in the Normal and Ischemic Brain Is

Controlled by Arteriolar Smooth Muscle Cell Contractility and Not by

Capillary Pericytes. Neuron *87*, 95-110. 10.1016/j.neuron.2015.06.001.

21.   Holstein-Rønsbo, S., Gan, Y., Giannetto, M.J., Rasmussen, M.K.,

Sigurdsson, B., Beinlich, F.R.M., Rose, L., Untiet, V., Hablitz, L.M.,

Kelley, D.H., and Nedergaard, M. (2023). Glymphatic influx and

clearance are accelerated by neurovascular coupling. Nat Neurosci *26*,

1042-1053. 10.1038/s41593-023-01327-2.

22.   Hartmann, D.A., Berthiaume, A.A., Grant, R.I., Harrill, S.A., Koski, T.,

Tieu, T., McDowell, K.P., Faino, A.V., Kelly, A.L., and Shih, A.Y. (2021).

Brain capillary pericytes exert a substantial but slow influence on blood

flow. Nat Neurosci *24*, 633-645. 10.1038/s41593-020-00793-2.

23. Chao, T.H., Zhang, W.T., Hsu, L.M., Cerri, D.H., Wang, T.W., and Shih,

Y.I. (2022). Computing hemodynamic response functions from

concurrent spectral fiber-photometry and fMRI data. Neurophotonics *9*,

032205. 10.1117/1.NPh.9.3.032205.

24. Niwa, K., Araki, E., Morham, S.G., Ross, M.E., and Iadecola, C. (2000).

Cyclooxygenase-2 contributes to functional hyperemia in whisker-barrel

cortex. J Neurosci *20*, 763-770.

10.1523/JNEUROSCI.20-02-00763.2000.

25. Yang, G., Zhang, Y., Ross, M.E., and Iadecola, C. (2003). Attenuation of

activity-induced increases in cerebellar blood flow in mice lacking

neuronal nitric oxide synthase. Am J Physiol Heart Circ Physiol *285*,

H298-304. 10.1152/ajpheart.00043.2003.

26. Ando, K., Tong, L., Peng, D., Vazquez-Liebanas, E., Chiyoda, H., He, L.,

Liu, J., Kawakami, K., Mochizuki, N., Fukuhara, S., et al. (2022).

KCNJ8/ABCC9-containing K-ATP channel modulates brain vascular

smooth muscle development and neurovascular coupling. Dev Cell *57*,

1383-1399 e1387. 10.1016/j.devcel.2022.04.019.

27. Shen, J., Wang, D., Wang, X., Gupta, S., Ayloo, B., Wu, S., Prasad, P.,

Xiong, Q., Xia, J., and Ge, S. (2019). Neurovascular Coupling in the

Dentate Gyrus Regulates Adult Hippocampal Neurogenesis. Neuron

103, 878-890 e873. 10.1016/j.neuron.2019.05.045.

28.   Hogan-Cann, A.D., Lu, P., and Anderson, C.M. (2019). Endothelial

NMDA receptors mediate activity-dependent brain hemodynamic

responses in mice. Proc Natl Acad Sci U S A *116*, 10229-10231.

10.1073/pnas.1902647116.

29.   Chow, B.W., Nunez, V., Kaplan, L., Granger, A.J., Bistrong, K., Zucker,

H.L., Kumar, P., Sabatini, B.L., and Gu, C. (2020). Caveolae in CNS

arterioles mediate neurovascular coupling. Nature *579*, 106-110.

10.1038/s41586-020-2026-1.

30.   Lee, C., Doo, E., Choi, J.M., Jang, S.H., Ryu, H.S., Lee, J.Y., Oh, J.H.,

Park, J.H., Kim, Y.S., Brain-Gut Axis Research Group of Korean Society

of, N., and Motility (2017). The Increased Level of Depression and

Anxiety in Irritable Bowel Syndrome Patients Compared with Healthy

Controls: Systematic Review and Meta-analysis. J Neurogastroenterol

Motil *23*, 349-362. 10.5056/jnm16220.

31.   Sarro, E.C., Sullivan, R.M., and Barr, G. (2014). Unpredictable neonatal

stress enhances adult anxiety and alters amygdala gene expression

related to serotonin and GABA. Neuroscience *258*, 147-161.

10.1016/j.neuroscience.2013.10.064.

32.   Suda, K., Setoyama, H., Nanno, M., Matsumoto, S., and Kawai, M.

(2013). Involvement of parasympathetic pelvic efferent pathway in

psychological stress-induced defecation. World J Gastroenterol *19*,

1572   1200-1209. 10.3748/wjg.v19.i8.1200.

1573 33. Gorard, D.A., Gomborone, J.E., Libby, G.W., and Farthing, M.J. (1996).

1574   Intestinal transit in anxiety and depression. Gut *39*, 551-555.

1575   10.1136/gut.39.4.551.

1576 34. Hale, M.W., Hay-Schmidt, A., Mikkelsen, J.D., Poulsen, B., Shekhar, A.,

1577   and Lowry, C.A. (2008). Exposure to an open-field arena increases

1578   c-Fos expression in a distributed anxiety-related system projecting to

1579   the basolateral amygdaloid complex. Neuroscience *155*, 659-672.

1580   10.1016/j.neuroscience.2008.05.054.

1581 35. Gorski, J.A., Talley, T., Qiu, M., Puelles, L., Rubenstein, J.L., and Jones,

1582   K.R. (2002). Cortical excitatory neurons and glia, but not GABAergic

1583   neurons, are produced in the Emx1-expressing lineage. J Neurosci *22*,

1584   6309-6314. 10.1523/JNEUROSCI.22-15-06309.2002.

1585 36. Monyer, H., Burnashev, N., Laurie, D.J., Sakmann, B., and Seeburg,

1586   P.H. (1994). Developmental and regional expression in the rat brain and

1587   functional properties of four NMDA receptors. Neuron *12*, 529-540.

1588   10.1016/0896-6273(94)90210-0.

1589 37. Laurie, D.J., Bartke, I., Schoepfer, R., Naujoks, K., and Seeburg, P.H.

1590   (1997). Regional, developmental and interspecies expression of the

1591   four NMDAR2 subunits, examined using monoclonal antibodies. Brain

1592   Res Mol Brain Res *51*, 23-32. 10.1016/s0169-328x(97)00206-4.

1593 38. Boas, D.A., and Dunn, A.K. (2010). Laser speckle contrast imaging in

biomedical optics. J Biomed Opt *15*, 011109. 10.1117/1.3285504.

39. Zhang, Y., Chen, K., Sloan, S.A., Bennett, M.L., Scholze, A.R., O'Keeffe,

S., Phatnani, H.P., Guarnieri, P., Caneda, C., Ruderisch, N., et al.

(2014). An RNA-sequencing transcriptome and splicing database of glia,

neurons, and vascular cells of the cerebral cortex. J Neurosci *34*,

11929-11947. 10.1523/JNEUROSCI.1860-14.2014.

40. Foster, D.B., Shen, L.H., Kelly, J., Thibault, P., Van Eyk, J.E., and Mak,

A.S. (2000). Phosphorylation of caldesmon by p21-activated kinase.

Implications for the Ca(2+) sensitivity of smooth muscle contraction. J

Biol Chem *275*, 1959-1965. 10.1074/jbc.275.3.1959.

41. Huber, G., Ogrodnik, M., Wenzel, J., Stolting, I., Huber, L., Will, O.,

Peschke, E., Matschl, U., Hovener, J.B., Schwaninger, M., et al. (2021).

Telmisartan prevents high-fat diet-induced neurovascular impairments

and reduces anxiety-like behavior. J Cereb Blood Flow Metab *41*,

2356-2369. 10.1177/0271678X211003497.

42. Yang, S., and Webb, A.J.S. (2024). Reduced neurovascular coupling is

associated with increased cardiovascular risk without established

cerebrovascular disease: A cross-sectional analysis in UK biobank. J

Cereb   Blood   Flow   Metab,   271678x241302172.

10.1177/0271678x241302172.

43. Chang, H.C., Tai, Y.T., Cherng, Y.G., Lin, J.W., Liu, S.H., Chen, T.L., and

Chen, R.M. (2014). Resveratrol attenuates high-fat diet-induced

disruption of the blood-brain barrier and protects brain neurons from

apoptotic insults. J Agric Food Chem *62*, 3466-3475.

10.1021/jf403286w.

44.   Hochgerner, H., Singh, S., Tibi, M., Lin, Z., Skarbianskis, N., Admati, I.,

Ophir, O., Reinhardt, N., Netser, S., Wagner, S., and Zeisel, A. (2023).

Neuronal types in the mouse amygdala and their transcriptional

response to fear conditioning. Nat Neurosci *26*, 2237-2249.

10.1038/s41593-023-01469-3.

45.   Shoji, H., and Miyakawa, T. (2021). Effects of test experience,

closed-arm wall color, and illumination level on behavior and plasma

corticosterone response in an elevated plus maze in male C57BL/6J

mice: a challenge against conventional interpretation of the test. Mol

Brain *14*, 34. 10.1186/s13041-020-00721-2.

46.   Chen, M., Ma, S., Liu, H., Dong, Y., Tang, J., Ni, Z., Tan, Y., Duan, C., Li,

H., Huang, H., et al. (2024). Brain region-specific action of ketamine as

a rapid antidepressant. Science *385*, eado7010.

10.1126/science.ado7010.

47.   Silva, B.A., Gross, C.T., and Graff, J. (2016). The neural circuits of

innate fear: detection, integration, action, and memorization. Learn

Mem *23*, 544-555. 10.1101/lm.042812.116.

48.   Apfelbach, R., Blanchard, C.D., Blanchard, R.J., Hayes, R.A., and

McGregor, I.S. (2005). The effects of predator odors in mammalian prey

species: a review of field and laboratory studies. Neurosci Biobehav Rev *29*, 1123-1144. 10.1016/j.neubiorev.2005.05.005.

49. Shang, C., Chen, Z., Liu, A., Li, Y., Zhang, J., Qu, B., Yan, F., Zhang, Y., Liu, W., Liu, Z., et al. (2018). Divergent midbrain circuits orchestrate escape and freezing responses to looming stimuli in mice. Nat Commun *9*, 1232. 10.1038/s41467-018-03580-7.

50. Lee, T., Weinberg-Wolf, H., Zapadka, T.E., Rudenko, A., Demb, J.B., and Kim, I.J. (2024). Specific retinal neurons regulate context-dependent defensive responses to visual threat. PNAS Nexus *3*, pgae423. 10.1093/pnasnexus/pgae423.

51. Wang, L., Chen, I.Z., and Lin, D. (2015). Collateral pathways from the ventromedial hypothalamus mediate defensive behaviors. Neuron *85*, 1344-1358. 10.1016/j.neuron.2014.12.025.

52. Bigi, S., Maestripieri, D., Aloe, L., and Alleva, E. (1992). NGF decreases isolation-induced aggressive behavior, while increasing adrenal volume, in adult male mice. Physiol Behav *51*, 337-343. 10.1016/0031-9384(92)90150-z.

53. Lee, S., Kang, B.M., Shin, M.K., Min, J., Heo, C., Lee, Y., Baeg, E., and Suh, M. (2015). Chronic Stress Decreases Cerebrovascular Responses During Rat Hindlimb Electrical Stimulation. Front Neurosci *9*, 462. 10.3389/fnins.2015.00462.

54. Chiba, S., Numakawa, T., Ninomiya, M., Richards, M.C., Wakabayashi,

C., and Kunugi, H. (2012). Chronic restraint stress causes anxiety- and

depression-like behaviors, downregulates glucocorticoid receptor

expression, and attenuates glutamate release induced by brain-derived

neurotrophic factor in the prefrontal cortex. Prog

Neuropsychopharmacol Biol Psychiatry *39*, 112-119.

10.1016/j.pnpbp.2012.05.018.

55.   Vanlandewijck, M., He, L., Mae, M.A., Andrae, J., Ando, K., Del Gaudio,

F., Nahar, K., Lebouvier, T., Lavina, B., Gouveia, L., et al. (2018). A

molecular atlas of cell types and zonation in the brain vasculature.

Nature *554*, 475-480. 10.1038/nature25739.

56.   Froehner, S.C. (1993). Regulation of ion channel distribution at

synapses. Annu Rev Neurosci *16*, 347-368.

10.1146/annurev.ne.16.030193.002023.

57.   Lee, S., Kang, B.M., Kim, J.H., Min, J., Kim, H.S., Ryu, H., Park, H., Bae,

S., Oh, D., Choi, M., and Suh, M. (2018). Real-time in vivo two-photon

imaging study reveals decreased cerebro-vascular volume and

increased blood-brain barrier permeability in chronically stressed mice.

Sci Rep *8*, 13064. 10.1038/s41598-018-30875-y.

58.   Liu, Y., Wu, Y., and Yang, Q. (2024). The impact of stress on the risk

decision-making process. Psychophysiology *61*, e14595.

10.1111/psyp.14595.

59.   Sandrone, S., Bacigaluppi, M., Galloni, M.R., Cappa, S.F., Moro, A.,

Catani, M., Filippi, M., Monti, M.M., Perani, D., and Martino, G. (2014).

Weighing brain activity with the balance: Angelo Mosso's original

manuscripts come to light. Brain *137*, 621-633. 10.1093/brain/awt091.

60. Roy, C.S., and Sherrington, C.S. (1890). On the Regulation of the

Blood-supply of the Brain. J Physiol *11*, 85-158 117.

10.1113/jphysiol.1890.sp000321.

61. Eisenberger, N.I., Lieberman, M.D., and Williams, K.D. (2003). Does

rejection hurt? An FMRI study of social exclusion. Science *302*, 290-292.

10.1126/science.1089134.

62. Guo, Y., Zou, G., Shao, Y., Chen, J., Li, Y., Liu, J., Yao, P., Zhou, S., Xu,

J., Hu, S., et al. (2023). Increased connectivity of the anterior cingulate

cortex is associated with the tendency to awakening during N2 sleep in

patients with insomnia disorder. Sleep *46*. 10.1093/sleep/zsac290.

63. Cera, N., Di Pierro, E.D., Ferretti, A., Tartaro, A., Romani, G.L., and

Perrucci, M.G. (2014). Brain networks during free viewing of complex

erotic movie: new insights on psychogenic erectile dysfunction. PLoS

One *9*, e105336. 10.1371/journal.pone.0105336.

64. Brom, M., Both, S., Laan, E., Everaerd, W., and Spinhoven, P. (2014).

The role of conditioning, learning and dopamine in sexual behavior: a

narrative review of animal and human studies. Neurosci Biobehav Rev

38, 38-59. 10.1016/j.neubiorev.2013.10.014.

65. Hou, X.H., Hyun, M., Taranda, J., Huang, K.W., Todd, E., Feng, D.,

Atwater, E., Croney, D., Zeidel, M.L., Osten, P., and Sabatini, B.L.

(2016). Central Control Circuit for Context-Dependent Micturition. Cell

167, 73-86.e12. 10.1016/j.cell.2016.08.073.

66. Tateno, F., Sakakibara, R., Kishi, M., Tsuyusaki, Y., Furukawa, R.,

Yoshimatsu, Y., and Suzuki, Y. (2012). Brainstem Stroke and Increased

Anal Tone. Low Urin Tract Symptoms *4*, 161-163.

10.1111/j.1757-5672.2012.00143.x.

67. Trunnell, E.R., Baines, J., Farghali, S., Jackson, T., Jayne, K., Smith, R.,

and Stibbe, T. (2024). The need for guidance in antidepressant drug

development: Revisiting the role of the forced swim test and tail

suspension test. Regulatory Toxicology and Pharmacology *151*, 105666.

https://doi.org/10.1016/j.yrtph.2024.105666.

68. Nestler, E.J., and Hyman, S.E. (2010). Animal models of

neuropsychiatric disorders. Nat Neurosci *13*, 1161-1169.

10.1038/nn.2647.

69. Hassabis, D., Kumaran, D., Summerfield, C., and Botvinick, M. (2017).

Neuroscience-Inspired Artificial Intelligence. Neuron *95*, 245-258.

10.1016/j.neuron.2017.06.011.

70. Park, S., and Whang, M. (2022). Empathy in Human-Robot Interaction:

Designing for Social Robots. Int J Environ Res Public Health *19*.

10.3390/ijerph19031889.

71. Kim, K.J., Ramiro Diaz, J., Iddings, J.A., and Filosa, J.A. (2016).

Vasculo-Neuronal Coupling: Retrograde Vascular Communication to Brain Neurons. J Neurosci *36*, 12624-12639. 10.1523/jneurosci.1300-16.2016.

72. Moore, C.I., and Cao, R. (2008). The hemo-neural hypothesis: on the role of blood flow in information processing. J Neurophysiol *99*, 2035-2047. 10.1152/jn.01366.2006.

73. Jammal Salameh, L., Bitzenhofer, S.H., Hanganu-Opatz, I.L., Dutschmann, M., and Egger, V. (2024). Blood pressure pulsations modulate central neuronal activity via mechanosensitive ion channels. Science *383*, eadk8511. 10.1126/science.adk8511.

74. Attwell, D., Buchan, A.M., Charpak, S., Lauritzen, M., Macvicar, B.A., and Newman, E.A. (2010). Glial and neuronal control of brain blood flow. Nature *468*, 232-243. 10.1038/nature09613.

75. Zeilhofer, H.U., Studler, B., Arabadzisz, D., Schweizer, C., Ahmadi, S., Layh, B., Bösl, M.R., and Fritschy, J.M. (2005). Glycinergic neurons expressing enhanced green fluorescent protein in bacterial artificial chromosome transgenic mice. J Comp Neurol *482*, 123-141. 10.1002/cne.20349.

76. Rampon, C., Luppi, P.H., Fort, P., Peyron, C., and Jouvet, M. (1996). Distribution of glycine-immunoreactive cell bodies and fibers in the rat brain. Neuroscience *75*, 737-755. 10.1016/0306-4522(96)00278-3.

77. Seo, J.S., Park, J.Y., Choi, J., Kim, T.K., Shin, J.H., Lee, J.K., and Han,

P.L. (2012). NADPH oxidase mediates depressive behavior induced by

chronic stress in mice. J Neurosci *32*, 9690-9699.

10.1523/JNEUROSCI.0794-12.2012.

78. Sturman, O., Germain, P.L., and Bohacek, J. (2018). Exploratory

rearing: a context- and stress-sensitive behavior recorded in the

open-field test. Stress *21*, 443-452. 10.1080/10253890.2018.1438405.

79. Oyarzabal, E.A., Hsu, L.M., Das, M., Chao, T.H., Zhou, J., Song, S.,

Zhang, W., Smith, K.G., Sciolino, N.R., Evsyukova, I.Y., et al. (2022).

Chemogenetic stimulation of tonic locus coeruleus activity strengthens

the default mode network. Sci Adv *8*, eabm9898.

10.1126/sciadv.abm9898.

80. Yu, X.D., Zhu, Y., Sun, Q.X., Deng, F., Wan, J., Zheng, D., Gong, W.,

Xie, S.Z., Shen, C.J., Fu, J.Y., et al. (2022). Distinct serotonergic

pathways to the amygdala underlie separate behavioral features of

anxiety. Nat Neurosci *25*, 1651-1663. 10.1038/s41593-022-01200-8.

81. Jin, Y., Shi, P., Wang, Y., Li, J., Zhang, J., Zhao, X., Ge, Y., Huang, Y.,

Guo, M., Wang, F., et al. (2022). Precise control of embolic stroke with

magnetized red blood cells in mice. Commun Biol *5*, 136.

10.1038/s42003-022-03082-9.

82. Li, J., Zhang, Y., Zhang, D., Wang, W., Xie, H., Ruan, J., Jin, Y., Li, T., Li,

X., Zhao, B., et al. (2024). Ca(2+) oscillation in vascular smooth muscle

cells control myogenic spontaneous vasomotion and counteract

post-ischemic no-reflow. Commun Biol *7*, 332.

10.1038/s42003-024-06010-1.

83. Li, W., Prakash, R., Chawla, D., Du, W., Didion, S.P., Filosa, J.A., Zhang,

Q., Brann, D.W., Lima, V.V., Tostes, R.C., and Ergul, A. (2013). Early

effects of high-fat diet on neurovascular function and focal ischemic

brain injury. Am J Physiol Regul Integr Comp Physiol *304*, R1001-1008.

10.1152/ajpregu.00523.2012.

84. Hou, W., Yao, J., Liu, J., Lin, X., Wei, J., Yin, X., Huang, H., Chen, X.,

Yang, G.Y., and He, X. (2023). USP14 inhibition promotes recovery by

protecting BBB integrity and attenuating neuroinflammation in MCAO

mice. CNS Neurosci Ther *29*, 3612-3623. 10.1111/cns.14292.

85. Khalil, V., Faress, I., Mermet-Joret, N., Kerwin, P., Yonehara, K., and

Nabavi, S. (2023). Subcortico-amygdala pathway processes innate and

learned threats. Elife *12*. 10.7554/eLife.85459.

86. Arakawa, H., Arakawa, K., Blanchard, D.C., and Blanchard, R.J. (2007).

Scent marking behavior in male C57BL/6J mice: sexual and

developmental determination. Behav Brain Res *182*, 73-79.

10.1016/j.bbr.2007.05.007.

87. Arakawa, H., Arakawa, K., Blanchard, D.C., and Blanchard, R.J. (2008).

A new test paradigm for social recognition evidenced by urinary scent

marking behavior in C57BL/6J mice. Behav Brain Res *190*, 97-104.

10.1016/j.bbr.2008.02.009.

88.   Borelli, K.G., Blanchard, D.C., Javier, L.K., Defensor, E.B., Brandao,

M.L., and Blanchard, R.J. (2009). Neural correlates of scent marking

behavior in C57BL/6J mice: detection and recognition of a social

stimulus. Neuroscience *162*, 914-923.

10.1016/j.neuroscience.2009.05.047.

89.   McQuaid, R.J., Audet, M.C., and Anisman, H. (2012). Environmental

enrichment in male CD-1 mice promotes aggressive behaviors and

elevated corticosterone and brain norepinephrine activity in response to

a mild stressor. Stress *15*, 354-360. 10.3109/10253890.2011.623249.

90.   Liu, J., Li, D., Huang, J., Cao, J., Cai, G., Guo, Y., Wang, G., Zhao, S.,

Wang, X., and Wu, S. (2022). Glutamatergic Neurons in the Amygdala

Are Involved in Paclitaxel-Induced Pain and Anxiety. Front Psychiatry

13, 869544. 10.3389/fpsyt.2022.869544.

91.   Warden, M.R., Selimbeyoglu, A., Mirzabekov, J.J., Lo, M., Thompson,

K.R., Kim, S.Y., Adhikari, A., Tye, K.M., Frank, L.M., and Deisseroth, K.

(2012). A prefrontal cortex-brainstem neuronal projection that controls

response to behavioural challenge. Nature *492*, 428-432.

10.1038/nature11617.

1810

A Neurovascular interface NsMJ — Cation/Anion Receptors, Bouton, Glu/Gly, SMC, Neuron, Lumen

B 1: P0 cerebellum; 2: Parenchymal aSMCs sorted from adult brains — NMDAR subunits: Grin1, Grin2a, Grin2b, Grin2c, Grin2d, Grin3a, Grin3b; 18s

C mRNA fold change NMDAR subunits — P0 cerebellum, Parenchymal aSMCs

D RNAscope_SMA-creER:Ai14 — tdTomato, Grin2d, DAPI; 10 µm

E Grin2d^f/f, SMA-creER:Grin2d^f/f — Smoothelin, GluN2D, DAPI; 10 µm

F Relative GluN2D protein level — Grin2d^f/f, SMA-creER:Grin2d^f/f; ****

G Relative GluN2D protein level — GluN2D, Tubulin; 0.0288

H Relative GluN2D protein level (Brain) — GluN2D, Tubulin; ns

I ΔCBF% — Grin2d^f/f, SMA-creER:Grin2d^f/f; rCBF at home cage (au.); ns

J AUC (au.) — Grin2d^f/f, SMA-creER:Grin2d^f/f; 0.0170

K Tube stress & OFT — Control, Global Genetic, BLA siRNA, Diet, Home cage

L rCBF (Δ%), Defecation (Δ%) — Increase, Decrease, Unchange

M Accumulated defecation latency sum (s) — 10687s, 4913s; ****

N OFT fecal particles — 11.81 folds, 3.50 folds, 3.00 folds

O Defecation rate (%) — Yes/No

P Fecal particles — Grin2d^f/f, SMA-creER:Grin2d^f/f

Q KEGG Pathway Enrichment_Top10 — Vascular smooth muscle contraction

R Input, IgG, GluN2D_IP — GluN2D, Caldesmon

S Cald relative expression level — Scramble, Cald KD; 0.0271

T rCBF Variation (%) — Scramble siRNA, Cald KD(Global), Cald KD (BLA)

U Scramble, Cald KD — Caldesmon, NeuN, DAPI; 200 µm

V Relative caldesmon expression level — Scramble, Cald KD; Global 0.0010, BLA local 0.0005

A

Manipulations
Grin2d^{f/f}
SMA-creER:Grin2d^{f/f}
Cald siRNA

Stress level
Level_0
Level_1
Level_2
Level_3

Home cage | OFT_40*40 | OFT_35*35 | c-OFT | Tube stress | Looming | VR_eagle

Unbiased prediction index (PI) classification

Home cage | VR_eagle | OFT_40*40 | WT_OFT_40*40 (BLA local *Cald* siRNA) | Looming | c-OFT | OFT_35*35 | Tube stress

200 µm

B

WT_HC | cKO_HC | WT_OFT_40 | WT_OFT_35 | WT_c-OFT | WT_TS | WT_Looming | WT_VR

Level_0 | Level_1 | Level_2 | Level_3

C

WT_HC | WT_OFT_40 | WT_OFT_35 | WT_c-OFT | WT_TS | cKO_OFT_40 | WT_Looming | WT_VR

Level_0 | Level_1 | Level_2 | Level_3

D

WT_HC | WT_OFT_40 | WT_OFT_35 | WT_C-OFT | *Cald* siRNA | WT_TS | WT_Looming | WT_VR

Level_0 | Level_1 | Level_2 | Level_3

E

WT_HC | WT_OFT_40 | WT_OFT_35 | WT_C-OFT | Ai32 (ChR2) | WT_TS | WT_Looming | WT_VR

Level_0 | Level_1 | Level_2 | Level_3

F

WT_HC | WT_OFT_40 | WT_OFT_35 | WT_C-OFT | cKO_looming | WT_TS | WT_Looming | WT_VR

Level_0 | Level_1 | Level_2 | Level_3

G

WT_HC | WT_OFT_40 | WT_OFT_35 | WT_C-OFT | cKO_VR | WT_TS | WT_Looming | WT_VR

Level_0 | Level_1 | Level_2 | Level_3

A

Small space:3072cm³   Large space: 4032cm³

B

0.0045

Total fighting (s)

- Grin2d^f/f
- SMA-creER:Grin2d^f/f

C

| WT_HC | WT_OFT_40 | WT_HC_Social | WT_OFT_35 | WT_c-OFT | WT_TS | WT_Looming | WT_VR |

Level_0   Level_1   Level_2   Level_3

D

| WT_HC | cKO_HC_Social | WT_OFT_40 | WT_OFT_35 | WT_c-OFT | WT_TS | WT_Looming | WT_VR |

Level_0   Level_1   Level_2   Level_3

E

0.0314

Time in lower side (s)

- Grin2d^f/f
- SMA-creER:Grin2d^f/f

F

Flee rate (%)

- Controls or NVC-deficit
- CD1

N=7/8   N=3/10

G

Rate (%)

37 events   19 events   37 events   19 events

N=7/37   N=5/19   N=36/37   N=7/19

Fight trigger   Flee

- Grin2d^f/f
- SMA-creER:Grin2d^f/f

H

ns

Sniffing (s)

- Grin2d^f/f
- SMA-creER:Grin2d^f/f