Neuron-vascular Coupling in the Amygdala Control Behavioral State

of Stress-induced Defecation and Innate Fear Retention

Jiayu Ruan[1,2,3,4], Huiqi Xie[1,2,3,4], Xinghua Quan[2,3,4], Yiyi Zhang[1,2,3,4], Dongdong Zhang[2,3,4],

Wentao Wang[2,3,4], Jie-Min Jia[1,2,3,4, *]

Affiliations

1. College of Life Sciences, Zhejiang University, Hangzhou, Zhejiang 310058, China.

2. Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province,

School of Life Sciences, Westlake University, Hangzhou 310024, China.

3. Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou 310024, China.

4. Laboratory of Neurovascular Biology, Institute of Basic Medical Sciences, Westlake Institute

for Advanced Study, Hangzhou 310024, China.

*Correspondence: Jie-Min Jia (jiajiemin@westlake.edu.cn)

Abstract

Proper cerebral function depends on intact neurovascular coupling (NVC), a conservative biological process in mammals that can effectively and spatiotemporally regulate cerebral blood distribution. The amygdala is a critical cerebral region for regulating emotion and stress-related behaviors. However, how NVC participates in advanced cerebral function and the consequences of NVC dysfunction in the amygdala remain unclear. The lack of cellular-specific transgenic models and non-specific drug administration mainly limited previous studies on NVC. Moreover, almost all papers take the function of hyperemia as the readout of NVC, but the advanced significance of functional hyperemia still needs to be discovered. Hence, the findings could have been more robust. In this study, we broadly used cellular-specific transgenic models to manipulate NVC and combined them with the chemogenetic strategy to reveal NVC-related cerebral advanced function. This study confirmed that GLRB loss in mural cells leads to NVC enhancement, the first report of a transgenic NVC enhancement model. Also, we are the first to use the transgenic model to clarify that COX-2 in neurons contributes to NVC regulation. The major work of this study is using arterial smooth muscle cells (aSMCs) NR2D conditional knockout model as a presentative NVC dysfunction model to find that NVC dysfunction in the amygdala context-dependently leads to shorter innate fear retention and increased stress-induced defecation. Through chemogenetic inhibition of neurons in the amygdala can correct increased stress-induced defecation in NVC dysfunction mice. Besides, during the behavioral assay, we first report the correlation pattern between pupil size and locomotion during the virtual predator stimulation. Last, we found that caldesmon is the downstream protein of NR2D in aSMCs to regulate NVC, and the interaction site of NR2D and caldesmon may become a drug target for NVC-initiated mental disorders.

Introduction

Neurovascular coupling (NVC), where corresponding changes in cerebral blood flow accompany changes in neural activity, is a universally conserved biological process among mammalians that can spatiotemporally regulate cerebral blood supply distribution(1). The amygdala is a crucial cerebral region that regulates emotional and stress-related behaviors cross species (2-5). One of the known applications of NVC is functional magnetic resonance

imaging (fMRI) (6), in which high levels of the blood oxygenation level-dependent (BOLD) signal mirrored with an increase in NVC are often repeatedly observed in some classic cerebral regions of patients with emotional disorders (7) such as in the amygdala of the fear-related post-traumatic stress disorder (PTSD) patients (8, 9).

The innate fear response is a complex emotional behavioral that helps organisms protect themselves from potential threats (10, 11). The amygdala is responsible for responding to potential threats in the environment, and it does so by activating a cascade of physiological and behavioral responses like fight or flight to avoid damage (12, 13). However, when the amygdala activation is abnormal, the excessive or weakened response to threat is harmful. Besides the innate fear response, several studies have explored the relationship between stress-induced defecation and the amygdala (14, 15). In the animal model, repeated OFT (16, 17) and wrap restraint (18) increased stress-induced defecation, and another study directly points out that restraint stress causes neuron activation in the amygdala (19). Moreover, a study examined the relationship between stress and bowel function in patients with irritable bowel syndrome (IBS), and researchers found that patients with IBS had increased activity in the amygdala during resting-state(20, 21).

The above evidence indicates that amygdala activation correlates with innate fear response and stress-induced defecation, but causality is unclear. Moreover, current studies usually use the variation of functional hyperemia as a readout of NVC, but how functional hyperemia controls advanced cerebral function is unknown. For instance, the impact of NVC dysfunction in the amygdala on innate fear response and stress-induced defecation remains unclear.

To answer these questions, we validate a series of cellular-specific transgenic models that can alter the extent of NVC. A representative example is NR2D, one of the non-core subunits of the NMDA receptor. Lack of NR2D in arterial smooth muscle cells (aSMCs) leads to attenuating phase NVC dysfunction and causes shorter innate fear retention time and increased stress-induced defecation. The abnormal increased stress-induced defecation was corrected in the NVC dysfunction mice by inhibiting the amygdala neurons by the chemogenetic strategy.

Besides, mural cell-specific expressed caldesmon was the downstream protein of NR2D in

aSMCs for NVC regulation, and the interaction site of NR2D and caldesmon may be a

potentially specific drug target for NVC-initiated mental disorders.

Materials and methods

Animal and animal care

All mice procedures complied with the Institutional Animal Care and Use Committee (IACUC)

guidelines of the School of Life Sciences, Westlake University. All mice housed in Westlake

University Laboratory Animal Resource Centre with standard animal rooms provided a 12 hours

light/dark cycle at 25°C. No more than six mice were fed in each cage, and standard chow and

water were provided to the mice *ad libitum*. All transgenic mice were listed in Table 1.

Cell culture

Surgical tweezers were used to quickly extract cortex of P0 anesthetized mice, and tissue was

put into 10ml 0.25% Trypsin (Thermo Scientific, Cat# 25200072, USA) to digest tissue for 15min

at 37°C. After that, the solution was added with 1ml FBS (Gibico, Cat# 10099, USA) and filtered

by a 40μm filter (Falcon, Cat# 352340, CHN). Finally, centrifuged 1,600*g at 4℃ for 5min and

cells were resuspended by 5ml SMCM (ScienCell, Cat# 1101, USA) in T25 flasks (Corning,

Cat# 430639, USA).

TMX administration

1.   Tamoxifen for mice

Tamoxifen (MCE, Cat# HY-13757A, USA) was dissolved in corn oil (Aladdin, Cat# C116023-

500ml, CHN) and orally administrated to mice (10mg/ml, 200ul per mice per day). Postnatal

day 5 (P5) *grin2d* conditional knockout infant mice got tamoxifen through breast milk. On the

fifth day after the birth of infant mice, their mother was orally administrated with tamoxifen for

four days. As for conditional knockout *acta2*, Tamoxifen was given from P21 to P24. Otherwise,

tamoxifen was orally administered from P60 to P63.

2.   Tamoxifen for cell culture (include cell sorting)

When cells reached approximately 80% full of the bottom surface of T25 flasks, tamoxifen was

dissolved in DMSO (Solarbio, Cat# D8371, CHN) to add into culture medium (final

concentration 5µM). After four-day culture, EGFP (Ai47) or Tdtomato (Ai14) in cells were

expressed. Cells were sorted by MA-900 (Sony, FCCF-MBR, JPN) to obtain EGFP-positive or

Tdtomato-positive cells. Sorted cells were cultured in SMCM for further assay. The purity of

cells sorted by the same method had been verified in the published paper (22).

RNA scope in situ hybridisation

Horizontal sections (15µm) of tamoxifen administrated *SMA-creER:Ai14* (for *grin2d*) or *Pdgfrβ-*

creER:Ai14 (for *glrb*) mice brain slices were obtained by CM1950 (Lecia, Histo-CM1950, DEU).

All RNA scope *in situ* hybridisation productions were obtained from ACDbio, including RNAcope

Multiplex Fluorescent Reagent Kit v2 (ACDbio, Cat# 323100, USA), *grin2d* probe (ACDbio,

Cat# 425951-C3, USA), *grin1* probe (ACDbio, Cat# 431611, USA), and *glrb* probe (ACDbio,

Cat# 1225261-C1, USA). All operation followed the reagent instructions and the published

paper (23).

RT-PCR

After obtaining sorted EGFP/Tdtomato-positive aSMCs, cells were cultured in 6cm dishes

(Corning, Cat# 430166, USA) with SMCM. After cells reached around 90% density, Trizol

(Sangon Biotec, Cat# B511311-0100, CHN) was used for the RNA extraction. For brain tissue,

after mice were anesthetized, the cortex or cerebellum of the mice was directly taken out, and

TRIzol was used for RNA extraction.

After RNA extraction, concentrations were measured by NanoDrop™ One (Thermo Scientific,

Cat# 840-317400. USA), and all samples were reverse transcription to cDNA (50ng/µl) using

HiScript III 1st Strand cDNA Synthesis Kit (Vazyme, Cat# R312-02, CHN). For the Real Time

Polymerase Chain Reaction (RT-PCR), each reaction contained 10µl of Green Taq Mix

(Vazyme, Cat# P131-AA, CHN), 2µl of cDNA, 2µl of primer pairs (10µM), and 6µl of RNAase-

free water. Agarose gel electrophoresis used 2% agarose gel (BIOWESTE, Cat# 111860, ESP),

100bp maker (Tsingke, Cat#: TSJ100-100, CHN), and ran at 220V for 30min. RT-PCR gel

images were obtained by Tanon imager (Tanon, 1600 Geil imager, CHN). Primer pairs listed in

Table 2.

Western-bolt

For tissue sample collection, samples were ground together with RIPA (Thermo Scientific, Cat#

89901, USA) on ice for 30min. For cell samples, after cells reached around 90% density, the

RIPA was used for protein extraction on ice for 30min. Protease Inhibitor Cocktail (CWBIO,

Cat# CW22008, CHN) was added into RIPA to prevent protein enzymolysis. Then, samples

were centrifuged 12,000*g at 4℃ for 5min, and supernatant were collected. Concentrations of

proteins were measured by Micro BCA Protein Assay kit (CWBIO, Cat# CW2011S, CHN), and

all samples were boiled with SDS-PAGE loading buffer (CWBIO, Cat# CW0027S, CHN) for

10min at 100℃. 8% SDS-polyacrylamide electrophoresis gels were used for gel running, and a

protein marker (GenStar, Cat# M221-05, CHN) was used to identify the protein size. After 90min

gels running at 120V (80V for the first 30min), gels were transferred to 0.45μm PVDF

membranes (Millipore, Cat# IPVH00010, USA) at 400mA for 120min. After transfer, PVDF

membranes were blocked by 5% non-fat milk in 1X TBST buffer for 60min at room temperature

with a mild horizontal shake.

After the block, the first antibody diluted with 5% non-fat milk in 1X TBST buffer was used to

incubate the membrane overnight at 4℃ with a mild horizontal shake. After the first antibody

incubation, the membrane was washed by 1X TBST 10min (3 replicates). The corresponding

second HRP-conjugated antibody diluted with 5% non-fat milk in 1X TBST buffer was used to

incubate the membrane 45min at room temperature with a mild horizontal shake. After the

second antibody incubation, the membrane was washed by 1X TBST 10min (3 replicates). ECL

kits (Abbkine, Cat# BMU102-CN, CHN) and gel image system (GE, GEL Imaging System

AI680RGB v.2.0, USA) were used for exposure photography. Antibodies listed in Table 3.

Immunofluorescence staining (IF)

For tissue Immunofluorescence (IF) staining, briefly, after anesthesia, the mice were perfused

with cold 1X PBS (10ml) and cold 4%PFA in 1X PBS (10ml) through the left ventricle to empty

blood and fix tissue. Afterwards, tissue was extracted and fixed by 4% PFA in 1X PBS 6 hours

at 4℃. For MCA branches IF staining, after the fixation, MCA branches were carefully extracted

from the brain using surgical tweezers. Then, treated isolated MCA branches by 1% Triton X-

100 (Diamond, Cat# A110694-0500, CHN) with 2.5% Normal Goat Serum (Jackson

ImmunoResearch, Cat# 005-000-121, USA) in 1X PBS for one hour at room temperature. Then,

isolated MCA branches were rewashed three times by 1XPBS. For other tissue like brain and

peripheral organs, all organs were sliced into 50μm (Leica, Histo-VT1200S, DEU) after the

fixation, then the subsequent process was similar to that of isolated MCA branches.

For cultured cell immunofluorescence staining. After cells reached around 90% density, cells

were passed by into 12-well plates (Corning, Cat# 3515, USA) with glass slices (30k cells per

well). After cells in 12-well plates reached around 70% density, cells in glass slices were fixed

by 4%PFA in 1X PBS for 10min at room temperature. Washed cell slices with 1X PBS 10min

(3 replicates), treated cells with 0.3% Triton X-100 in 1X PBS for 10min at room temperature,

and then washed again with 1X PBS 10min (3 replicates). Next, 5% BSA in 1X PBS was used

to block cell slices for 30min at room temperature. After the block, cell slices were rewashed as

described above.

In the staining process, the first antibody diluted with 1% BSA+0.1% Triton X-100 in 1X PBS

was used to incubate samples overnight at 4℃. After the first antibody incubation, samples

were washed with 1X PBS 10min (3 replicates). The corresponding donkey-source second

fluorescence-conjugated antibody diluted with 1% BSA+0.1% Triton X-100 in 1X PBS was used

to incubate samples 120min at room temperature. After the second antibody incubation,

samples were washed with 1X PBS 10min (3 replicates). After that, 1000X diluted Hoechst

(AbMole, Cat# M5112, CHN) was used for nucleus staining (5min). Finally, tissue slices and

cell culture slices were covered by a glass slice with the mounting medium (SouthernBiotech,

0100-01, USA). Zeiss LSM800 (Zeiss, LSM800, DEU) microscopy was used for photography.

Antibodies listed in Table 3.

Whisker stimulation

The whisker was trimmed to only remain lane three and lane four on one side of the face. On

the day of the assay, mice were anesthesia with 1% pentobarbital sodium, and stable fixated

on the platform. The electro-motor-driven brush stimulated the whisker for 1 minute at 12Hz.

The principle of laser speckle contrast imaging has been described in the previous paper (24),

and the operation referred to the published paper (25). Laser Speckle Image System (RWD,

RFSLI ZW/RFSLI III, CHN) was used to monitor the CBF variation in the ROI of the opposite

barrel cortex of mice. The recording parameters were: 512*512 resolution, 10FPS, and 2min

recording.

Vasomotion measurement

Two weeks before the assay, a head fixator was placed on the head of the mouse, and a glass

cranial window replaced the 3-mm round skull above the cortex. On the day of the assay,

isoflurane (RWD, cat# R510-22-10, CHN) was used to anesthetize mice and RhoB dextran

(sigma, cat#r9379, 10mg/ml) was injected into the orbit. The mice were then fixed under a two-

photon microscope and recorded while awake. The image and analysis strategy were

consistent with our pre-print paper (doi: https://doi.org/10.1101/2023.11.05.565680).

Immunoprecipitation assay

Sorted Tdtomato-positive aSMCs were used for lentivirus infection and further

immunoprecipitation (IP) assay. Cells were cultured in 6cm dishes with SMCM. After cells

reached around 90% density, add lentivirus (Lenti-EF1-EGFP-CMV-grin2d-3XFlag-WPRE) into

the culture and keep for one week (virus titer: 2.6*109) for cells to over express NR2D. MA-900

sorted out EGFP and Tdtomato double-positive cells, and these cells were lentivirus-infected.

These cells were then cultured in 10cm dishes. After cells reached around 90% density, cells

were collected by RIPA reagent. The Protein G Magnetic beads kit (Epizyme, Cat# YJ002, CHN)

was used for the immunoprecipitation assay. NR2D antibody and IgG control antibody were

mixed with Protein G Magnetic beads respectively to pull down NR2D and its related protein.

Pull-down products were verified by western-bolt and analyzed through mass spectrometry.

Antibodies listed in Table3.

cald siRNA interference

All operations were followed by paper *Cannula Implantation into the Cisterna Magna of Rodents*

(https://doi.org/10.3791/57378). cald siRNA (1ul, 1nM) was mixed with Fluorescein

Isothiocyanate (FITC) (10mg/ml, Sigma, FD40S-1G, USA) for injection (20μl/10min). Ten

minutes after the injection, the scalp was removed, and the microscope was used to detected

the FITC signal (Zeiss, Axio Zoom v16, DEU). Only successfully injected mice remain. The

whisker stimulation was performed 48 hours after the injection, and brain was extracted for

Immunofluorescence staining and RT-PCR after the whisker stimulation.

Looming assay

This assay was carried out in a 35cm*35cm OFT box. Setting of the assay referred to the

published paper (26). Modifications include: 1. mice will be put in the OFT box to adapt for 10

minutes before starting the assay. 2. We only use a single camera at the top of the box to record.

3. We stimulate the mice when they move and are not close to the box wall rather than

continuously stimulating them. Each mouse performs three stimulations. 4. Locomotion speed

index before (LSI_{before}) calculation used the mean of locomotion from -30s to -10s and the mean

of locomotion from 0s to 3s; LSI_{after} calculation used the mean of locomotion from 0s to 3s and

the mean of locomotion from 3s to 30s. After the assay, mice were fed alone in a clean new

home cage for 90 minutes and then brain was extraction to detect c-fos expression in the

amygdala.

VR assay

This assay referred to a published paper (27). Two weeks before the assay, the mouse head

was fixed with a fixator. One week before the assay, the mice received 10 minutes of adaptive

training on the vital reality (VR) system every day (refers Figure 3B). On the day of the assay,

the mice were fixed on the VR system and given 5 minutes of adaptation. The screen in front

of the mouse will always play the landscape video unless it is stimulated. The stimulation video

was 3 seconds with the eagle swooped down, and the voice of the eagle played at the beginning

of the stimulation. There were 5 times of stimulation in total, and the criterion for the first

stimulation was that the mice moved for the first time after 5 minutes of adaptation. The criteria

for subsequent stimulation were that the last stimulation ended for at least 2 minutes and the

mice resumed movement. The assay stopped 2min after the last stimulation. The pupil size

variation, locomotion speed, defecation and urination of mice were recorded during the whole

assay. After the assay, mice were fed alone in a clean new home cage for 90 minutes and then

brain was extraction to detect c-fos expression in the amygdala.

Cat video stimulation assay

The assay was carried out in a 40cm*40cm OFT box. The mouse head was fixed with a fixator

on the head two weeks before the assay. One week before the assay, the mice received 10

minutes of adaptive training on the treadmill. On the day of the assay, the mice were fixed on

the treadmill, and played the video of cat hunting on the iPad Air (Apple, Cat#: 3rd generation,

USA) in front of the mouse (16cm) three trails for stimulation (30s for each trail). One trail

contains 10s empty, and 20s cat appearance (cat ambush at 26s). The camera above the box

was used for recording in the assay.

CD-1 fear assay

The setup of CD-1 fear assay referred to published papers(28-31). Aggressive CD-1 mice that

were never exposed to NVC dysfunction and control mice (described as B6 mice below) were

used for the assay. Before the assay, simultaneously put a single B6 mouse and a CD-1 mouse

into a home cage. The cage contains new padding and a metal lid on the top. One side of the

metal lid is used for placing food and water bottles, so the height is lower than another side.

After that, the camera was used to record all behaviours of B6 mice and CD-1 mice in 10

minutes. After the assay, CD-1 and B6mice were separated again. After the assay, B6 mice

were fed alone in a clean new home cage for 90 minutes and then brain was extraction to detect

c-fos expression in the amygdala.

Home cage physical examination

The blood pressure of mice was measured by Kent Noninvasive blood pressure measurement

system for animals/Biopac physiological signal analytical system (Kent/Biopac, Cat#, USA)

before the mice were put into the home cage. Each home cage only contains four mice. In each

cycle (three days), 60g of food, 340g of water (including bottle weight), and 100g of padding were provided. Food/water consumption, weight of mice, fecal particles, and urination amount (counted by increased padding weight) were measured at the end of each cycle. Three cycles of monitoring were performed for each home cage.

Chemogenetic manipulation

2.88^{E+12}VG/ml AAV2/9-mCaMKIIa-hM3D(Gq)-mCherry-ER2-WPRE-pA (Shanghai Taitool Bioscience Co., Ltd, CHN) was used for chemogenetic manipulation. Bilateral amygdala injection (150nl each side, AP: -1.6mm; ML: +/-3.45mm; DV: -4.1mm) was performed two weeks before the assay. Intraperitoneal injection of DREADD agonist 21 (C21, MCE, Cat# HY-100234, USA) was used to inhibit neurons in the amygdala. The injection strategy referred a published paper (32).

Open field test

The setup of open field test (OFT) referred to published papers (33-36). Each mouse was placed in the central area of a 40*40 cm or 35*35 cm box for the OFT. The camera recording was 15 minutes. For chemogenetic manipulation assay, only use 40*40 cm box for OFT, and C21 injection was 45min before the OFT.

Open field-like test

The setup of open field-like test (OFLT) referred to published paper (29). Each mouse was placed in the central area of a 23*23 cm box for the OFLT. An iron mesh wall replaced one wall of the box. Sketch paper was put on the bottom of the box to collect urine. Symmetric space is on the opposite side of the iron mesh wall. The camera recording was 20 minutes. For the analysis, the box was divided into three parts (left area, central area, and right area) to analyze the preference of mice for iron mesh wall (right area close to the iron mesh wall).

Tube stress

The tube stress assay refers to published papers (37-39). 50ml tube were used to restrain the mice. The tube cap and tube surface will be drilled to ensure normal breathing of mice. Mice

were restrained in the tube for 15min, and then calculate the fecal production. For

chemogenetic manipulation assay, The tube stress assay was performed 45min after C21

injection.

Mice quarantine assay

Mice were intraperitoneal C21 injection at the beginning of the assay, and then each mouse

was isolated into a clean cage without water, food, and padding supply for 1 hour. Defecation

and urination results were checked at the end of assay.

Electrophysiology

Parameter settings of the Electrophysiology followed the published paper (40). Mice were

anaesthetized and then perfused with 10ml cold artificial cerebrospinal fluid (aSCF). Then, the

brain was extracted and put into chilled aCSF. 300μm Brain slices were prepared in coronal

sections (Leica, Histo-VT1200S, DEU). Then, brain slices were transferred to aCSF at 34 °C

for 30 minutes and then recovered to room temperature for one hour. All solutions were

oxygenated with 95% O2 + 5% CO2. For *in vitro* electrophysiological recording of the amygdala

neurons, currents were measured under whole-cell patch-clamp recordings using pipettes with

a typical resistance of 5–6 MΩ filled with an internal solution. Cells were visualized by

microscope. The recording system includes an amplifier (MultiClamp 700B) driven by

Axon™pCLAMP™ (v.10.7) and a digital-to-analog converter (Digidata 1550B). The resting

membrane potential (RMP) was determined during the silent period of neuronal spontaneous

activity. Then, continuously increasing stimulation current was given from -20pA to 300pA

(lasting one second), and the number of action potentials (AP) was recorded in different levels

of current stimulation.

Statistical analysis software and code

All the analysis of fluorescence intensity and grey value used Zeiss Zen Lite (v2.3) and ImageJ

(v1.53u). All whisker stimulation original data were measured by LSIS-self software (v5.0).

Parameters "Time to peak", "Peak mean", "Peak maximum", "OFLT heat map", "Z-score cross

area" and data inclusion criteria before bioinformatics analysis of mass spectrometry data were

calculated using self-compiled MATLAB codes. KEGG and GO analysis referred to DAVID Bioinformatics Resources (https://david.ncifcrf.gov/home.jsp) and Kyoto Encyclopedia of Genes and Genomes (https://www.genome.jp/kegg/). The c-fos expression distribution map and KEGG pathway made by self-compiled R code. All the original behavioral data of OFLT were recorded using any-maze (v4.73). OFT original behavioral data were recorded using Noldus. The implementation of Looming stimulation assay and VR stimulation assay including raw data collection were assisted by BAYONE Co., Ltd. All statistical data were processed by Numbers (v12.2), and all figures were made by Prism (v9.4.1).

Results

Neurotransmitter receptors loss in mural cells lead NVC dysfunction

Under physiological conditions, sustained and correct neuronal activation in cerebral regions, including the amygdala, requires normal functioning of NVC (41-43). Therefore, we first plan to validate the ability of some *de novo* and classic potential candidates on NVC and then to see the impact of NVC dysfunction on particular advanced functions of neural circuits.

Our previous study reported that NR1 (encoded by gene *grin1*), the core subunit of NMDA receptor encoded by gene *grin1* in aSMCs, regulates NVC, but the contribution of another non-core subunit is still unclear. Hence, we first detected the expression level of all NMDA receptors in cerebral aSMCs to map the expression pattern of NMDA receptors. Among the seven known NMDA receptors, we found that apart from NR1, only the non-core subunit NR2D (encoded by gene *grin2d*) can be detected (Figure 1A/B). This result suggests that the NMDA receptor in cerebral aSMCs may only be composed of NR1 and NR2D.

Based on the above results, the loss of NR2D in aSMCs may share the same consequence of NR1 loss. Therefore, we further validated the expression of *grin2d*/NR2D in cerebral smooth muscle cells and other organs (Figure 1C/D, Figure S1E/H). Then, we confirmed the genetic tool (cre-loxp system) was specific and efficient in manipulating the gene expression (Figure 1D/E/F/G/H/I/K, Figure S1F/G), and conditional knockout *grin2d* in aSMCs did not impact the maker protein expression of SMCs (Figure 1J/L).

Afterwards, we conducted the whisker stimulation to evaluate the NVC reflected by the ability

of functional hyperemia in our transgenic mice (Figure 1M). First, we confirmed that multiple

consecutive whisker stimulations did not impact the data on repeatability and reliability (Figure

S1A/C), and Tamoxifen administration did not impact functional hyperemia (Figure S1B/D).

Based on the above, data showed that specific *grin2d* loss in aSMCs (driven by SMA-creER)

can attenuate functional hyperemia, and this reduction is independent of the time of knocking

out *grin2d*, which indicates that the functional compensation from other non-core subunits of

NMDA receptors in aSMCs may not exist (Figure 1N/O). In addition, *grin2d* loss throughout all

mural cells (driven by Pdgfrβ-creER) rather than limited to aSMCs revealed a similar

aggravated inhibition of functional hyperemia, suggesting that mural cells other than aSMCs

may not express *grin2d*. Knocking out *grin2d* alone in aSMCs is enough to produce significant

functional hyperemia inhibition (Figure 1N/O). Finally, the double knockout of *grin1* and *grin2d*

in aSMCs is similar to those of a single knockout of *grin1* or *grin2d*, suggesting that *grin1* and

grin2d in aSMCs regulate the functional hyperemia in the same pathway (Figure 1N/O).

To verify the downstream pathway of NR2D regulating neurovascular coupling in aSMCs, we

identified the proteins interacting with NR2D by immunoprecipitation. The Vascular smooth

muscle contraction pathway, identified through KEGG Pathway Enrichment, contains

caldesmon and smooth muscle actin (SMA) (Figure S2A/G). SMA is the final executive protein

responsible for cell contraction and relaxation (44). Non-phosphorylated caldesmon binds with

actin and myosin to prevent their interaction, thereby inhibiting the contraction of the cell

skeleton (45-48). Through phosphorylation of MAPK (49) and PAK (50) or binding with calcium-

activated CaM (51), caldesmon is removed from the binding site of actin and myosin, leading

to the contraction of the cytoskeleton. First, we conditional knockout *acta2* (gene name of SMA)

in mural cells (Figure S2E) and found that functional hyperemia was inhibited (Figure S2H/I).

These results suggest that SMA is involved in regulating NVC. Directly affecting contraction

and relaxation machines of mural cells could impact NVC. As for caldesmon, this protein almost

mural cells specific in the brain. So, we block the expression of caldesmon in the whole brain

by injecting *cald* (gene name of caldesmon) siRNA into the cisterna magna. After 48 hours of

siRNA interference, we found that the expression of caldesmon decreased (Figure S2B/C/D/F)

and functional hyperemia in mice was inhibited at the same time (Figure S2H/I), suggesting

that caldesmon may participate in the regulation of nNVC as the downstream of NR2D. In

conclusion, caldesmon and SMA potential are downstream proteins of NR2D in aSMCs.

In addition to the NMDA receptor, we verified the contribution of GLRB and COX-2 in NVC.

Firstly, we found that GLRB (encoded by gene *glrb*), the beta subunit of the glycine receptor,

was also expressed in cerebral mural cells. The expression of *glrb*/GLRB is relatively

concentrated in venous smooth muscle cells (vSMCs) and less in aSMCs and pericytes (Figure

1P/R/S). Functional hyperemia can be enhanced when *glrb* lost in all mural cells (Figure 1N/O).

This phenomenon was the first reported example of NVC enhancement by the trangenetic

model rather than chemical drugs. Besides, we also found that *glrb* loss in mural cells impacts

the vasomotion (Figure S1I). Although the vasomotion frequency did not vary in veins, the

vasomotion amplitude increased. However, only conditional knockout *glrb* in aSMCs did not

impact the frequency and amplitude of venous vasomotion (Figure S1J/K). These data indicate

that *glrb* expressed on vSMCs was critical in inhibiting functional hyperemia and venous

vasomotion amplitude. This suggests that vasomotion variations may correlate with NVC

variations, but the causality is unknown.

As for COX-2 (encoded by the gene *ptgs2*), the previous view was that COX-2 mainly

expressed in neurons, and total COX-2 knockout or drug inhibition can inhibit functional

hyperemia by approximately 50% (52). However, we also found that COX-2 was also expressed

in cerebral aSMCs (Figure 1Q/T). To clarify the contribution of COX-2 in different types of cells,

we conditional knock out *ptgs2* in neurons (driven by Emx1-cre) and aSMCs (driven by SMA-

creER). Data showed that specific knockout of *ptgs2* in aSMCs did not affect functional

hyperemia. However, in neurons, it reduced by approximately 50%, indicating that COX-2

expressed in aSMCs was not involved in regulating functional hyperemia, and the significant

contribution came from COX-2 in neurons (Figure 1N/O).

NVC dysfunction mice had shorter innate fear retention during the looming stimulation assay

Looming stimulation assay is a classic model that reflects the innate fear response of mice, so

we chose this classic model to detect the innate fear response ability of NVC dysfunction mice

(Figure 2A). We characterized the locomotion of mice in the 30s before and after stimulation

(Figure 2B). First of all, we detected no difference in the locomotion between NVC dysfunction

mice and control mice during the non-stimulation period (-30s to -10s), which indicated no

difference in the motor function of mice (Figure 2C). The difference in the possible locomotion

data after stimulation did not come from the motor dysfunction of mice. After that, we evaluated

the latency of the locomotion of mice return to the mean locomotion before the stimulation (-

30s to -10s). We found that the latency of NVC dysfunction mice was shorter, which was more

evident in the first two trials (Figure 2D). After that, we calculated the LSI of three trials (Figure

2E) and found that the LSI of NVC dysfunction mice after stimulation was significantly higher

than that of control mice (Figure 2F). In addition, the Area Under Curve (AUC) of locomotion

between 6s to 24s in NVC dysfunction mice was larger than that of control mice in the first trial

(Figure 2G). The above data showed that NVC dysfunction had shorter innate fear retention.

Finally, we detected the activation of neurons in the amygdala of mice after the assay, and

found that the activation of neurons in NVC dysfunction mice decreased significantly (Figure

2H/I). This evidence suggests that the low-level activation of amygdala neurons was correlated

to the shorter innate fear retention.

NVC dysfunction mice had shorter innate fear retention during the VR stimulation assay

To further simulate the real natural predators attack, we further used the VR system (Figure

3A/B) to give mice more real threaten to induce more real innate fear behaviour of mice. We

analysed the data for 30s before and after each stimulation (Figure 3C). We first analysed the

changes in pupil diameter and locomotion speed of mice from -30s to -10s in 5 trials. We found

no difference in these two parameters between the NVC dysfunction mice and control mice

(Figure 3E/F). The above evidence indicates that NVC dysfunction mice did not exhibit pre-

existing abnormalities in pupil and locomotion before stimulation.

Afterwards, we used the Z-score to compare the pattern of mouse pupil diameter and

locomotion of mice to analyze the correlation between these two diameters in 5 trials. In the

range of -3s to 0s, due to the onset of movement, the pupil diameter increases synchronously

with the increase of locomotion speed, showing a positive correlation consistent with previous

papers. In the range of 0s to 3s, the pupil diameter of mice will rapidly expand due to the

stimulation. However, mice will also exhibit a freezing pattern, causing a rapid decrease in

locomotion speed, resulting in a negative correlation between pupil diameter and locomotion

speed. This pattern shift remained consistent across five trials (except for trial_4 in the NVC

dysfunction group, where there was no difference). In summary, a positive correlation exists

between pupil diameter and locomotion speed during normal movement in mice, and a negative

correlation between them during mice were shocked (Figure 3D). This assay was the first one

to characterize the correlation between pupil size and locomotion speed in mice during the

virtual predator stimulation.

Next, we analysed the amplitude of pupil diameter changes in 5 trials, and there was no

difference in the increase in pupil diameter between the NVC dysfunction mice and the control

mice during the stimulation period (0s-3s) (Figure 3G). However, after the stimulation (4s-10s),

the NVC dysfunction group mice had a faster rate of pupil shrinkage, especially in trial_2, where

the pupil diameter of the NVC dysfunction mice decreased significantly than that of the control

mice (Figure 3H). In addition, we also analysed the changes in the locomotion speed of mice

during the stimulation period (0s-3s). In trail_1, the NVC dysfunction mice showed a lower

locomotion speed decrease than the control mice (Figure 3I). After the VR stimulation assay,

we analysed the activation of amygdala neurons in mice and found that the activation of

amygdala neurons was lower in NVC dysfunction mice (Figure 3K/L). Based on the above

results, it is suggested that the NVC dysfunction group had a shorter fear retention time and

weaker instinctive fear response when the virtual predator attacked due to the less neuron

activation in the amygdala.

Finally, we analysed the area enclosed by the z-score curve of mouse pupil diameter and

locomotion speed during and after stimulation to reflect the adaptability of mice to continuous

stimulation. We found that the area decreased with increasing repeats of stimulation in both

groups of mice, indicating that the mice showed adaptability to continuous non-invasive

stimulation (Figure 3J/M).

NVC dysfunction mice had weaker innate fear response during the cat video stimulation

As a supplement to the looming and VR assay, to increase dimensions to verify whether the

NVC dysfunction in the amygdala affects innate fear behaviours, we used the cat video as a

visual trigger to stimulate the innate fear behaviour of mice. We used the retreat pattern (move

backward) and tail rattling to evaluate the ability of innate fear response.

Firstly, we mapped the standing and retreating patterns of mice on the treadmill (Figure S3A).

Firstly, we counted the number of tail rattling, which was positively correlated with the degree

of fear level. It can be seen that the number of tail rattling in the NVC dysfunction mice was

significantly lower than that in the control mice in the first trial (Figure S3B/C). Then, the number

of tail rattling decreased in the following two trails, showing an adaptation trend. As for the

retreat, it can be seen that compared with the control mice, NVC dysfunction mice had

significantly less time to retreat at the stage of cat appearance and cat ambush, which was

reflected in the probability and total time to retreat. However, with the increase of trials, the

response of mice to the video weakened, showing a trend of adaptation (Figure S3D). The

above data showed that NVC dysfunction mice had shorter innate fear retention, indicating that

NVC dysfunction can affect the innate fear response in mice.

NVC dysfunction mice had weaker innate fear response during the CD-1 fight assay

In the above assay, we administered non-contact stimulation to mice to evaluate innate fear

response. Next, we placed the NVC dysfunction mice in a cage with aggressive CD-1 mice for

10 minutes to observe the innate fear response in contact stimulation assay. We first map the

fighting patterns of B6 mice with CD-1 mice in the cage (Figure S3E). Then, we analyzed the

frequency of sniffing (Figure S3F). Results showed that the number of sniffing these between

NVC dysfunction mice and control mice had no differences, indicating that the information input

received by the two groups of mice was consistent. Therefore, the possible behavioural

differences in the subsequent parameters do not originate from the information input bias due

to NVC dysfunction.

Then, we found that compared with the control mice, the NVC dysfunction mice showed a

significant reduction in the fight latency (Figure S3G), number of fights (Figure S3H), and total

duration (Figure S3I). On the contrary, the NVC dysfunction mice spent longer in a narrow space

than CD-1 mice (Figure S3J/O). In addition, we also analyzed the triggers and losers of each

fight. Whether in the NVC dysfunction mice or the control mice, the main initiators of fights were

CD-1 mice, but the rate of fights triggered by NVC dysfunction mice mildly increased (Figure

S3K). As for fighting losers, we used flight behaviour as the index to identify the losers. CD-1

mice were nearly not defeated in the control mice group, but the probability of CD-1 defeat

significantly increased in the NVC dysfunction group (Figure S3L). After the CD-1 fight assay,

we measured the neuron activation in the amygdala and other cerebral regions (Figure S3M/N).

Results showed that the activation of amygdala neurons in the NVC dysfunction mice was lower

than in the control mice. Based on the above results, it can be concluded that in the contacted

fight assay, NVC dysfunction mice showed similar shorter innate fear retention to that in the

non-contact stimulation assay.

NVC dysfunction altered stress-induced defecation

We set up a series of assays from low-stress to high-stress levels to verify whether the NVC

dysfunction will impact stress-induced defecation (Figure 4A). In the home cage with the lowest

stress, there was no difference in the physiological and metabolic parameters between NVC

dysfunction mice and the control mice, such as food (Figure 4C) /water (Figure S4A) intake,

body weight (Figure S4D), blood pressure (Figure S4E), defecation (Figure 4B/D/E, and Figure

S4B), and urination (Figure S4C). In addition, we also measured the intestinal system and found

no difference in the length of intestines (Figure S4F), body weight (Figure S4G), and the rate

between length and weight (Figure S4H) between the NVC dysfunction mice and the control

mice. Also, intestinal bleeding cannot be detected in NVC dysfunction mice. After the home

cage assay, brain was extracted from the mouse for the detection of neuron activation in the

amygdala, and no obvious differences can be seen between NVC dysfunction mice and contril

mice (Figure S4U).

In the electrophysiological recording of brain slices, the amygdala neurons of NVC dysfunction

mice had similar minimum action potential activation current (Figure S4J), and resting-stage

membrane potential (Figure S4K) were similar to that of control mice, and the ability of BLA

neurons to generate action potentials during the different current stimulation was also normal

(Figure S4L/M). These results confirm that the basic physical parameters and function of

amygdala neurons were normal in NVC dysfunction mice.

In the Open Field Test (OFT), the fecal production of mice in the control group did not increase

significantly at the 40cm OFT box compared with that in the home cage (Figure 4B). However,

the defecation increased significantly after changing the OFT box to 35cm (Figure 4B),

suggesting that space may induce stress in mice. Hence, subsequent assays were to increase

the pressure by reducing the space. However, compared with the control group, the defecation

of NVC dysfunction mice was significantly increased in two sizes of OFT boxes (Figure 4B/F/G),

and NVC dysfunction mice produced more feces at each five minutes of OFT than that of control

mice in the OFT (Figure S4I), which suggests that NVC dysfunction mice were more sensitive

to increased stress. Furthermore, combining results obtained in the home cage and

electrophysiological recording, it can be seen that increased stress-induced defecation in NVC

dysfunction mice was context-dependent rather than the consequence of continuous defecation

increase. Besides, we also found no difference in the running distance (Figure 4H) and area

preference (Figure 4I) of the NVC dysfunction mice in the OFT—only the defecation-related

parameters were altered, which showed that NVC dysfunction did not entirely change mice

behaviors.

In addition, we also compared the stress-induced defecation in the OFT (40cm) in various NVC

dysfunction mice (Figure S4N). In general, the number of fecal particles negatively correlated

with the extent of functional hyperemia; that is, the decrease of functional hyperemia leads to

an increase in defecation and vice versa. Importantly, *grin1* and *grin2d* conditional knock mice

showed a similar increase in defecation, which is another dimension to indicate that they work

in the same pathway. Besides, conditional knockout *ptgs2* in aSMCs did not affect functional

hyperemia and defecation, while the conditional knockout *ptgs2* by Emx1-cre inhibited

functional hyperemia and increased defecation. Moreover, because the fecal production of mice

in the OFT is relatively low, *glrb* conditional knockout mice only exhibit defecation reduction in

the tube stress, which provides more space-induced stress to induce more stress-induced

defecation than the OFT. Based on the above, the increase in defecation may not be related to

a particular gene but to the alteration of NVC.

In the Open Field-like Test (OFLT), we modified the existing OFT paradigm to increase the

space-induced stress by narrowing the box size. The results showed that NVC dysfunction mice

produced more fecal particles than control mice (Figure 4B). Similarly, urination also increased

in the NVC dysfunction mice (Figure 4J, and Figure S4Q), and the increase near the iron mesh

wall was more prominent (Figure 4K, and Figure S4O), which was synchronous with the

increased preference of the right side of the box (Figure S4P/R/S/T), indicating that the

unknown space may attract mice and provide stress to mice at the same time. Due to this

phenomenon, the running distance of NVC dysfunction mice increased in the OFLT (Figure

S4R).

In the tube stress assay, mice were restrained in the tube, and there was no difference in

defecation between the NVC dysfunction mice and the control mice (Figure 4B). Still, the

defecation of the two groups increased significantly compared with that in the home cage and

the OFT, which showed that fecal production reached the maximum in this assay and more

stress can no longer make mice produce more feces. As mentioned above, conditional

knockout *glrb* in mural cells lead defecation decreased in the tube stress (Figure S4N).

To reveal the mechanism for the increase in defecation, we analyzed the activation of neurons

in different cerebral regions after the OFT. We found that neuron activations of NVC dysfunction

mice were almost attenuated or unchanged (Figure 4L/M/N). Only a few cerebral regions

showed increased neuron activation, including the amygdala (Figure S4V), which strongly

indicates that NVC dysfunction in the amygdala leads to abnormal neuron activation, which

may correlate with abnormal stress-induced defecation. Vascular anatomical evidence (Figure

S4W), including BOLD signal evidence, proved that the amygdala has abundant vascular

distribution, which is the structural cornerstone for NVC to affect the excitability of amygdala

neurons. Further analyses confirmed that neurons activated by the OFT in the BLA were mainly

excitatory neurons, and approximately 60% were excitatory neurons in CeA (Figure 4O). These

proportions did not change in the NVC dysfunction mice, indicating that the defecation increase

in the NVC dysfunction group was not due to extra circuit activation. However, the activation

extent of the same circuits was different.

The ultimate executor of stress-induced defecation is undoubtedly neurons. Therefore, if the

abnormal activation of amygdala neurons leads to increased stress-induced defecation in NVC

dysfunction mice, the artificial reduction of neuron activation may correct the increase in

abnormal feces. So, we used the chemogenetic method to inhibit the activation of amygdala

neurons (Figure 4P) and found that the defecation of mice in quarantine assay (Figure 4Q),

OFT (Figure 4R), and tube stress assay (Figure 4S) were significantly reduced. This decrease

in defecation was also observed in the NVC dysfunction mice, suggesting that NVC dysfunction

in the amygdala specifically leads to abnormally increased activation of excitatory neurons to

enhance the stress-induced defecation, and this phenomenon could be corrected by directly

inhibiting the neuron activation in the amygdala.

Discussion

In this study, we found three proteins (NR2D, GLRB, COX-2) involved in the regulation of NVC,

in which the loss of GLRB in mural cells leads to the enhancement of NVC, which is the first

reported transgenic model that can enhance NVC. Previous studies only reported that NVC

was enhanced by drug inhibition. In addition, we also clarified the cellular identity of COX-2

regulating NVC. COX-2 is expressed in neurons, glial cells, and cerebral aSMCs. Previous

studies have used total knockout mice to study the effect of COX-2 on NVC, but this model

lacks cell specificity and produces serious side effects (53). Similarly, the assay of drug

inhibition (like NS-398) on COX-2 cannot achieve cell specificity (52). In this study, we used

Emx1-cre and SMA-creER mice, respectively, to distinguish the contribution of COX-2 in

neurons and SMCs to NVC by transgenic strategies. Our conclusion confirmed that COX-2

expressed on neurons mainly mediated the regulation of NVC but not COX-2 in SMCs.

We selected NR2D conditional knockout mice as the representative NVC dysfunction model for

further study. We found that the double knockout of NR1 and NR2D in SMCs would not further

aggravate the effect of NVC. Besides, NR1 and NR2D single knockout mice showed similar

increased stress-induced defecation. This evidence suggests that the non-core NMDA receptor

subunit NR2D worked together with NMDA receptor core subunit NR1 in the same pathway.

We have further broadened the field of NVC research because this study first reported the effect

of NVC on advanced cerebral function, like stress-induced defecation and innate fear response.

Previous NVC studies, whether using laser Doppler imaging (54), laser speckle imaging (55)

or two-photon imaging (56), were limited to detecting functional hyperemia itself but not the

consequence of abnormal functional hyperemia. These assays cannot reflect the changes in

higher-level cerebral function during NVC dysfunction. This study found that NVC-attenuated

mice showed abnormally increased stress-induced defecation and shorter innate fear retention

under different stimulation models. It should be noted that there is no significant difference

between NVC dysfunction mice and control mice in the home cage without stimulation, which

indicates that NVC dysfunction is not obvious in the resting state. Only after stimulation can

NVC dysfunction mice show abnormal advanced function, which may also be the difficulty in

detecting NVC-related advanced function. The data from the electrophysiology confirmed that

the function of neurons was normal in NVC dysfunction mice, and only in the activation degree

to particular stimulation different from that of normal mice, which indicated that the cause of

some diseases, such as IBS, might not be the lesion of neurons but the lesion of NVC of these

neurons.

Therefore, NVC may become a potential candidate for mental diseases in the future, and NR2D

expressed on cerebral aSMCs is an ideal target. We found that only cerebral aSMCs expressed

NR2D in vascular SMCs. In addition, with the increase of age, the expression of NR2D in the

whole brain decreases, which makes the side effects of drugs targeting NR2D in SMCs

relatively low. More importantly, we found that caldesmon was downstream of NR2D by

immunoprecipitation and functional hyperemia detection, while caldesmon is only expressed in

mural cells in the brain. Suppose the drug candidate is designed to strengthen/block the

interaction between NR2D and caldesmon, in that case, the specificity of candidates will be significantly improved because the interaction between NR2D and caldesmon exists only in cerebral aSMCs. Hence, by targeting the interaction site of NR2D and caldesmon, the candidate compound can correct the mental disorders initiated by NVC.

Acknowledgement

We thank Yanmei Tao for sharing *Nestin-cre* mice and *Emx1-cre* mice. We thank Bo Zhou for sharing *SMA-creER* mice. We thank Zilong Qiu for sharing Ai47 mice. We thank Xiang Yu for sharing *PdgfrβCreER* mice. We thank Yangang Sun for sharing *grin1* mice. We thank Woo-Ping Ge for sharing Ai14 mice. We thank Yuxiao Niu for drawing KEGG pathway map. We thank Jishuai Wang for electrophysiology recording. We thank Bingrui Zhao for cell sorting. We thank Xu Hu for primer sequences sharing of NMDA receptors.

Conflict of Interest

We declare that we do not have any financial or personal relationships with other people or organizations, which may affect our work inappropriately, and there are no professional or other personal interests of any nature or kind in any products, services and/or companies. These interests may be interpreted as affecting or reviewing the positions in the following manuscripts.

References

1. Drew PJ. Neurovascular coupling: motive unknown. Trends Neurosci. 2022;45(11):809-19.
2. Pitman RK, Rasmusson AM, Koenen KC, Shin LM, Orr SP, Gilbertson MW, et al. Biological studies of post-traumatic stress disorder. Nat Rev Neurosci. 2012;13(11):769-87.
3. Whalen PJ, Kagan J, Cook RG, Davis FC, Kim H, Polis S, et al. Human amygdala responsivity to masked fearful eye whites. Science. 2004;306(5704):2061.

4. Hadj-Bouziane F, Bell AH, Knusten TA, Ungerleider LG, Tootell RB. Perception of emotional expressions is independent of face selectivity in monkey inferior temporal cortex. Proc Natl Acad Sci U S A. 2008;105(14):5591-6.

5. Harris AP, Lennen RJ, Marshall I, Jansen MA, Pernet CR, Brydges NM, et al. Imaging learned fear circuitry in awake mice using fMRI. Eur J Neurosci. 2015;42(5):2125-34.

6. Lacroix A, Toussay X, Anenberg E, Lecrux C, Ferreiros N, Karagiannis A, et al. COX-2-Derived Prostaglandin E2 Produced by Pyramidal Neurons Contributes to Neurovascular Coupling in the Rodent Cerebral Cortex. J Neurosci. 2015;35(34):11791-810.

7. Clemm von Hohenberg C, Weber-Fahr W, Lebhardt P, Ravi N, Braun U, Gass N, et al. Lateral habenula perturbation reduces default-mode network connectivity in a rat model of depression. Transl Psychiatry. 2018;8(1):68.

8. Ahmed-Leitao F, Spies G, van den Heuvel L, Seedat S. Hippocampal and amygdala volumes in adults with posttraumatic stress disorder secondary to childhood abuse or maltreatment: A systematic review. Psychiatry Res Neuroimaging. 2016;256:33-43.

9. Kamiya K, Abe O. Imaging of Posttraumatic Stress Disorder. Neuroimaging Clin N Am. 2020;30(1):115-23.

10. Coryell MW, Ziemann AE, Westmoreland PJ, Haenfler JM, Kurjakovic Z, Zha XM, et al. Targeting ASIC1a reduces innate fear and alters neuronal activity in the fear circuit. Biol Psychiatry. 2007;62(10):1140-8.

11. Misslin R. The defense system of fear: behavior and neurocircuitry. Neurophysiol Clin. 2003;33(2):55-66.

12. Davis M. NMDA receptors and fear extinction: implications for cognitive behavioral therapy. Dialogues Clin Neurosci. 2011;13(4):463-74.

13. Shultz B, Farkash A, Collins B, Mohammadmirzaei N, Knox D. Fear learning-induced changes in AMPAR and NMDAR expression in the fear circuit. Learn Mem. 2022;29(3):83-92.

14. Taguchi R, Shikata K, Furuya Y, Hirakawa T, Ino M, Shin K, et al. Selective corticotropin-releasing factor 1 receptor antagonist E2508 reduces restraint stress-induced defecation and visceral pain in rat models. Psychoneuroendocrinology. 2017;75:110-5.

15. Kakol-Palm D, Brusberg M, Sand E, Larsson H, Martinez V, Johansson A, et al. Role of tachykinin NK(1) and NK(2) receptors in colonic sensitivity and stress-induced defecation in

gerbils. Eur J Pharmacol. 2008;582(1-3):123-31.

16. Candland DK, Nagy ZM. The open field: some comparative data. Ann N Y Acad Sci.

1969;159(3):831-51.

17. Archer J. Tests for emotionality in rats and mice: a review. Anim Behav. 1973;21(2):205-

35.

18. Suda K, Setoyama H, Nanno M, Matsumoto S, Kawai M. Involvement of parasympathetic

pelvic efferent pathway in psychological stress-induced defecation. World J Gastroenterol.

2013;19(8):1200-9.

19. Morgan A, Gaulden A, Altemus M, Williford K, Centanni S, Winder D, et al.

Cyclooxygenase-2 inhibition prevents stress induced amygdala activation and anxiety-like

behavior. Brain Behav Immun. 2020;89:513-7.

20. Qi R, Liu C, Weng Y, Xu Q, Chen L, Wang F, et al. Disturbed Interhemispheric Functional

Connectivity Rather than Structural Connectivity in Irritable Bowel Syndrome. Front Mol

Neurosci. 2016;9:141.

21. Yu Z, Liu LY, Lai YY, Tian ZL, Yang L, Zhang Q, et al. Altered Resting Brain Functions in

Patients With Irritable Bowel Syndrome: A Systematic Review. Front Hum Neurosci.

2022;16:851586.

22. Li X, Zhou L, Zhang X, Jin Y, Zhao B, Zhang D, et al. Proteins secreted by brain arteriolar

smooth muscle cells are instructive for neural development. Mol Brain. 2022;15(1):97.

23. Yu XD, Zhu Y, Sun QX, Deng F, Wan J, Zheng D, et al. Distinct serotonergic pathways to

the amygdala underlie separate behavioral features of anxiety. Nat Neurosci.

2022;25(12):1651-63.

24. Boas DA, Dunn AK. Laser speckle contrast imaging in biomedical optics. J Biomed Opt.

2010;15(1):011109.

25. Jin Y, Shi P, Wang Y, Li J, Zhang J, Zhao X, et al. Precise control of embolic stroke with

magnetized red blood cells in mice. Commun Biol. 2022;5(1):136.

26. Shang C, Chen Z, Liu A, Li Y, Zhang J, Qu B, et al. Divergent midbrain circuits orchestrate

escape and freezing responses to looming stimuli in mice. Nat Commun. 2018;9(1):1232.

27. Wang L, Chen IZ, Lin D. Collateral pathways from the ventromedial hypothalamus mediate

defensive behaviors. Neuron. 2015;85(6):1344-58.

28. Borelli KG, Blanchard DC, Javier LK, Defensor EB, Brandao ML, Blanchard RJ. Neural

correlates of scent marking behavior in C57BL/6J mice: detection and recognition of a social

stimulus. Neuroscience. 2009;162(4):914-23.

29. Arakawa H, Arakawa K, Blanchard DC, Blanchard RJ. Scent marking behavior in male

C57BL/6J mice: sexual and developmental determination. Behav Brain Res. 2007;182(1):73-9.

30. Arakawa H, Arakawa K, Blanchard DC, Blanchard RJ. A new test paradigm for social

recognition evidenced by urinary scent marking behavior in C57BL/6J mice. Behav Brain Res.

2008;190(1):97-104.

31. McQuaid RJ, Audet MC, Anisman H. Environmental enrichment in male CD-1 mice

promotes aggressive behaviors and elevated corticosterone and brain norepinephrine activity

in response to a mild stressor. Stress. 2012;15(3):354-60.

32. Khalil V, Faress I, Mermet-Joret N, Kerwin P, Yonehara K, Nabavi S. Subcortico-amygdala

pathway processes innate and learned threats. Elife. 2023;12.

33. Liu CY, Chen JB, Liu YY, Zhou XM, Zhang M, Jiang YM, et al. Saikosaponin D exerts

antidepressant effect by regulating Homer1-mGluR5 and mTOR signaling in a rat model of

chronic unpredictable mild stress. Chin Med. 2022;17(1):60.

34. Guo Z, Li S, Wu J, Zhu X, Zhang Y. Maternal Deprivation Increased Vulnerability to

Depression in Adult Rats Through DRD2 Promoter Methylation in the Ventral Tegmental Area.

Front Psychiatry. 2022;13:827667.

35. Kim S, Hong KB, Kim S, Suh HJ, Jo K. Creatine and taurine mixtures alleviate depressive-

like behaviour in Drosophila melanogaster and mice via regulating Akt and ERK/BDNF

pathways. Sci Rep. 2020;10(1):11370.

36. Lapmanee S, Charoenphandhu J, Teerapornpuntakit J, Krishnamra N, Charoenphandhu

N. Agomelatine, venlafaxine, and running exercise effectively prevent anxiety- and depression-

like behaviors and memory impairment in restraint stressed rats. PLoS One.

2017;12(11):e0187671.

37. Seo JS, Wei J, Qin L, Kim Y, Yan Z, Greengard P. Cellular and molecular basis for stress-

induced depression. Mol Psychiatry. 2017;22(10):1440-7.

38. Oh YS, Gao P, Lee KW, Ceglia I, Seo JS, Zhang X, et al. SMARCA3, a chromatin-

remodeling factor, is required for p11-dependent antidepressant action. Cell. 2013;152(4):831-

43.

39.   Seo JS, Park JY, Choi J, Kim TK, Shin JH, Lee JK, et al. NADPH oxidase mediates

depressive behavior induced by chronic stress in mice. J Neurosci. 2012;32(28):9690-9.

40.   Cui Y, Yang Y, Ni Z, Dong Y, Cai G, Foncelle A, et al. Astroglial Kir4.1 in the lateral habenula

drives neuronal bursts in depression. Nature. 2018;554(7692):323-7.

41.   Lauritzen M, Dreier JP, Fabricius M, Hartings JA, Graf R, Strong AJ. Clinical relevance of

cortical   spreading   depression   in   neurological   disorders: migraine,   malignant   stroke,

subarachnoid and intracranial hemorrhage, and traumatic brain injury. J Cereb Blood Flow

Metab. 2011;31(1):17-35.

42.   Ruitenberg A, den Heijer T, Bakker SL, van Swieten JC, Koudstaal PJ, Hofman A, et al.

Cerebral hypoperfusion and clinical onset of dementia: the Rotterdam Study. Ann Neurol.

2005;57(6):789-94.

43.   Pico-Perez M, Radua J, Steward T, Menchon JM, Soriano-Mas C. Emotion regulation in

mood and anxiety disorders: A meta-analysis of fMRI cognitive reappraisal studies. Prog

Neuropsychopharmacol Biol Psychiatry. 2017;79(Pt B):96-104.

44.   Arnoldi R, Hiltbrunner A, Dugina V, Tille JC, Chaponnier C. Smooth muscle actin isoforms:

a tug of war between contraction and compliance. Eur J Cell Biol. 2013;92(6-7):187-200.

45.   Kordowska J, Huang R, Wang CL. Phosphorylation of caldesmon during smooth muscle

contraction and cell migration or proliferation. J Biomed Sci. 2006;13(2):159-72.

46.   Wang   CL.   Caldesmon   and   smooth-muscle   regulation.   Cell   Biochem   Biophys.

2001;35(3):275-88.

47.   Hai CM, Gu Z. Caldesmon phosphorylation in actin cytoskeletal remodeling. Eur J Cell Biol.

2006;85(3-4):305-9.

48.   Wang CL. Caldesmon and the regulation of cytoskeletal functions. Adv Exp Med Biol.

2008;644:250-72.

49.   Childs TJ, Watson MH, Sanghera JS, Campbell DL, Pelech SL, Mak AS. Phosphorylation

of smooth muscle caldesmon by mitogen-activated protein (MAP) kinase and expression of

MAP kinase in differentiated smooth muscle cells. J Biol Chem. 1992;267(32):22853-9.

50.   Foster DB, Shen LH, Kelly J, Thibault P, Van Eyk JE, Mak AS. Phosphorylation of

caldesmon by p21-activated kinase. Implications for the Ca(2+) sensitivity of smooth muscle

contraction. J Biol Chem. 2000;275(3):1959-65.

51. Janovick JA, Natarajan K, Longo F, Conn PM. Caldesmon: a bifunctional (calmodulin and

actin) binding protein which regulates stimulated gonadotropin release. Endocrinology.

1991;129(1):68-74.

52. Niwa K, Araki E, Morham SG, Ross ME, Iadecola C. Cyclooxygenase-2 contributes to

functional hyperemia in whisker-barrel cortex. J Neurosci. 2000;20(2):763-70.

53. Slattery P, Frolich S, Goren I, Nusing RM. Salt supplementation ameliorates

developmental kidney defects in COX-2(-/-) mice. Am J Physiol Renal Physiol.

2017;312(6):F1044-F55.

54. Li W, Prakash R, Chawla D, Du W, Didion SP, Filosa JA, et al. Early effects of high-fat diet

on neurovascular function and focal ischemic brain injury. Am J Physiol Regul Integr Comp

Physiol. 2013;304(11):R1001-8.

55. Remzso G, Nemeth J, Toth-Szuki V, Varga V, Kovacs V, Domoki F. NMDA attenuates the

neurovascular response to hypercapnia in the neonatal cerebral cortex. Sci Rep.

2019;9(1):18900.

56. Chow BW, Nunez V, Kaplan L, Granger AJ, Bistrong K, Zucker HL, et al. Caveolae in CNS

arterioles mediate neurovascular coupling. Nature. 2020;579(7797):106-10.

Figure legends

Figure 1. NR2D, GLRB, and COX-2 expression patterns and functional hyperemia ability

measurement of corresponding transgenic mice. A) Expression of all NMDA receptors in mRNA

level detected by RT-PCR. B) Statistical results of Figure 1A. Each dot indicates an independent

experiment, and the data are mean ± s.e.m. C) RNA scope results of *grin2d* expression in

cerebral aSMCs. Red arrow pointed out *grin2d* signal in aSMCs. The inserted histogram shows

the intensity of fluorescent signal. D) The WB and statistical results of NR2D in cerebellum of

P0 *grin2d f/f* mice, P60 grin2d f/f mice, and P0 *Emx1-cre:grin2d f/f* mice. Each dot indicates an

independent experiment, and the data are mean ± s.e.m. E) The WB and statistical results of

NR2D in cerebellum of P10 *grin2d f/f* mice, and P10 *SMA-creER:grin2d f/f* mice. Each dot

indicates an independent experiment, and the data are mean ± s.e.m. F) The RT-PCR and

statistical results of *grin2d* in cerebral aSMCs from *grin2d f/f mice*, and *SMA-creER:grin2d f/f*

mice. Each dot indicates an independent experiment, and the data are mean ± s.e.m. G) The WB and statistical results of NR2D in cerebral aSMCs from *grin2d f/f* mice, and *SMA-creER:grin2d f/f* mice. Each dot indicates an independent experiment, and the data are mean ± s.e.m. H) The top row: The IF results of NR2D in isolated MCA of *grin2d f/f mice* (left), and *SMA-creER:grin2d f/f mice* (right). The bottom row: The IF results of NR2D in cultured aSMCs from *grin2d f/f mice* (left), and SMA-creER:grin2d f/f mice (right). I) The statistical results of NR2D expression of the top raw in Figure 1H. Each dot indicates an independent experiment, and the data are mean ± s.e.m. J) The statistical results of smoothelin expression of the top raw in Figure 1H. Each dot indicates an independent experiment, and the data are mean ± s.e.m. K) The statistical results of NR2D expression of the bottom raw in Figure 1H. Each dot indicates an independent experiment, and the data are mean ± s.e.m. L) The WB and statistical results of SMA in the cortex of P60 *grin2d f/f* mice, P60 *SMA-creER:grin2d f/f* mice. Each dot indicates an independent experiment, and the data are mean ± s.e.m. M) Schematic diagram of blood flow increases in the ROI before, during, and after the whisker stimulation. N) CBF variation before, during, and after the whisker stimulation. The solid line indicates the mean value, and the dotted line indicates ± s.e.m. O) Statistical data of Figure 1N, including how long the CBF variation reaches the peak period and the mean CBF variation during the peak period. Each dot indicates a mean data from one mouse. Data are mean ± s.e.m. P) RNA scope results of *glrb* expression in cerebral mural cells (left), and GLRB expression in vSMCs. Q) The RT-PCR results of ptgs2 in cortex of P60 mice and aSMCs of cultured aSMCs (left). The IF results of COX-2 expression in cerebral aSMCs (right). R) Statistical results of RNA-scope in Figure 1P. Brain slices from 3 mice were analysed, and the data are mean ± s.e.m. S) Statistical results of IF in Figure 1P. Brain slices from 3 mice were analysed, and the data are mean ± s.e.m. T) Statistical results of RT-PCR in Figure 1Q. Each dot indicates a mean data from one mouse, and the data are mean ± s.e.m.

Figure 2. NVC dysfunction mice showed shorter innate fear retention in the looming assay. A) Schematic diagram of assay setting. B) Locomotion of mice around the stimulation in three trails (-30s to 30s). The solid line indicates the mean value, and the dotted line indicates ±

s.e.m. C) Locomotion of mice from -30s to -10s in three trails. Each dot indicates one mouse, and the data are mean ± s.e.m. D) Locomotion latency of mice, which means that how long the locomotion of the mouse return to the average locomotion from -30s to -10s. Each dot indicates one mouse, and the data are mean ± s.e.m. E) LSI of each mouse in three trails (from left to right). Each dot indicates one mouse.   F) Statistical results of LSI$_{after}$ of data in Figure 2E. Each dot indicates one mouse, and the data are mean ± s.e.m. G) Statistical analysis of the locomotion AUC in Figure 2B from 6s to 24s. Each dot indicates one mouse, and the data are mean ± s.e.m. H) Representative c-fos IF image in the amygdala. I) Statistical data of c-fos activation in Figure 2H. Each dot indicates one mouse, and the data are mean ± s.e.m.

Figure 3. NVC dysfunction mice showed shorter innate fear retention in the VR assay. A) Schematic diagram of assay setting. B) The structure of the VR system. C) Z-score of locomotion and pupil size of mice around the stimulation in five trails (-30s to 30s).   D) The correlation between locomotion and pupil size from -3s to 0s and 0s to 3s. Each dot indicates one mouse, and the data are mean ± s.e.m. E) The pupil size of mice from -30s to -10s. Each dot indicates one mouse, and the data are mean ± s.e.m. F) The locomotion of mice from -30s to -10s. Each dot indicates one mouse, and the data are mean ± s.e.m. G) Statistical analysis of the pupil size AUC in Figure 3C from 0s to 3s. Each dot indicates one mouse, and the data are mean ± s.e.m. H) Statistical analysis of the pupil size AUC in Figure 3C from 3s to 10s. Each dot indicates one mouse, and the data are mean ± s.e.m. I) Statistical analysis of the speed AUC in Figure 3C from 0s to 3s. Each dot indicates one mouse, and the data are mean ± s.e.m. J) Statistical analysis of the curve enclosed area of pupil size and locomotion in Figure 3C from the first cross during the stimulation. Each dot indicates one mouse, and the data are mean ± s.e.m. K) Representative c-fos IF image in the amygdala. L) Statistical data of c-fos activation in Figure 3K. Each dot indicates one mouse, and the data are mean ± s.e.m. M) The presentative diagram of curve enclosed area of pupil size and locomotion shown in orange.

Figure 4. NVC dysfunction mice showed increased stress-induced defecation. A) Schematic diagram of assay setting from low stress to high stress. B) Fecal production in five assays

normalized in 15min. Each dot indicates one mouse, and the data are mean ± s.e.m. C) The

average food consumption of mice in the home cage. Each dot indicates one cage, and the

data are mean ± s.e.m. D) The average water consumption of mice in the home cage. Each dot

indicates one cage, and the data are mean ± s.e.m. E) The average fecal production number

of mice in the home cage. Each dot indicates one cage, and the data are mean ± s.e.m. F) The

defecation rate of mice in the OFT. Each dot indicates one mouse, and the data are mean ±

s.e.m. G) The defecation number of mice in the OFT. Each dot indicates one mouse, and the

data are mean ± s.e.m. H) The time of mice stay in the side of the box in the OFT. Each dot

indicates one mouse, and the data are mean ± s.e.m. I) The running distance of mice in the

OFT. Each dot indicates one mouse, and the data are mean ± s.e.m. J) The urination weight of

mice in the OFLT. Each dot indicates one mouse, and the data are mean ± s.e.m. K) The

urination mark size of mice in the OFLT. Each dot indicates one mouse, and the data are mean

± s.e.m. L) Representative c-fos IF image in the amygdala. M) Statistical data of BLA c-fos

activation in Figure 4L. N) Statistical data of CeA c-fos activation in Figure 4L. O) Statistical

percentage of BLA /CeA c-fos and camKII double positive cells in all c-fos positive cells in Figure

4L. P) Schematic diagram of AAV2/5-mCaMKIIa-hM4D(Gi)-EGFP-ER2-WPRE-pA injection in

the amygdala (left). The c-fos activation pattern in the tube stress assay when I.P saline

(middle). The c-fos activation pattern in the tube stress assay when I.P C21 (right). Q) The fecal

production of amygdala-infected mice with/without C21 in the quarantine assay. Each dot

indicates one mouse, and the data are mean ± s.e.m. R) The fecal production of amygdala-

infected mice with/without C21 in the OFT. Each dot indicates one mouse, and the data are

mean ± s.e.m. S) The fecal production of amygdala-infected mice with/without C21 in the tube

stress assay. Each dot indicates one mouse, and the data are mean ± s.e.m.

Figure S1. Supplementary data of Figure1. A) CBF variation before, during, and after the

whisker stimulation. The solid line indicates the mean value, and the dotted line indicates ±

s.e.m. B) CBF variation before, during, and after the whisker stimulation with or without

tamoxifen administration. The solid line indicates the mean value, and the dotted line indicates

± s.e.m. C) Statistical data of Figure S1A, including how long the CBF variation reaches the

peak period and the mean CBF variation during the peak period. Each dot indicates a mean data from one mouse. Data are mean ± s.e.m. D) Statistical data of Figure S1A, including how long the CBF variation reaches the peak period and the mean CBF variation during the peak period. Each dot indicates a mean data from one mouse. Data are mean ± s.e.m. E) RNA-scope data of grin1 and grin2d in cerebellum. F) IF of NeuN in the brains of *SMA-creER:Ai14* mice. G) Statistical data in Figure S1F, and each dot indicates an data from one mouse, and the data are mean ± s.e.m. H) IF of NR2D expression in heart, liver, spleen, lung, kidney, and small intestine of *SMA-creER:Ai14* mice. The first row is zoom-out images of six organs, the second row is zoom-in images of white circles in the first row, and the third row is histograms of the red and green fluorescent intensities of the indicated sites in the second row.

Figure S2. Downstream signal pathways analysis of NR2D in aSMCs. A) WB results of caldesmon detection in the IP product of NR2D. B) RT-PCR results of cald siRNA in whole brain lysis. C) Statistics analyze of the bottom row in Figure S2A. Each dot indicates one mouse, and the data are mean ± s.e.m. D) Statistics analyze of Figure S2F. Each dot indicate one mouse, and the data are mean ± s.e.m. E) WB results of SMA expression in P60 *acta2 f/f* mice, and p*dgfrβ-creER:acta2 f/f* mice. Each dot indicates one mouse, and the data are mean ± s.e.m. F) IF results of caldesmon siRNA in the S1BF. G) KEGG pathway analysis of 915 proteins selected from three independent mass spectrum data. H) CBF variation before, during, and after the whisker stimulation. The solid line indicates the mean value, and the dotted line indicates ± s.e.m. I) Statistical data of Figure S2H, including how long the CBF variation reaches the peak period and the mean CBF variation during the peak period. Each dot indicates a mean data from one mouse. Data are mean ± s.e.m.

Figure S3. NVC dysfunction mice showed attenuated innate fear resource in the cat video stimulation assay and CD-1 fight assay. A) The locomotion pattern of mice in the treadmill of three trials. B) Tail rattling times of mice. Each dot indicates one mouse, and the data are mean ± s.e.m. C) Tail rattling rate of mice. D) The retreat times of mice. Each dot indicates one mouse, and the data are mean ± s.e.m. E) The fight pattern of mice in the cage. F) Sniffing times of

mice. Each dot indicates one mouse, and the data are mean ± s.e.m. G) Fight latency of mice.

Each dot indicates one mouse, and the data are mean ± s.e.m. H) Number of fights happened

during the assay. Each dot indicates one mouse, and the data are mean ± s.e.m. I) Total fighting

times of mice. Each dot indicates one mouse, and the data are mean ± s.e.m. J) Time of B6

mice together with CD-1 mice in the narrow space. Each dot indicates one mouse, and the data

are mean ± s.e.m. K) Fighting trigger rate of mice. L) Fighting defeat (flight) rate of mice. M)

Statistical data of c-fos activation in Figure S3N. Each dot indicates one mouse, and the data

are mean ± s.e.m. N) Representative c-fos IF image in the amygdala. O) Example of fighting

cages.

Figure S4. Supplementary data of stress-induced defecation assay. A) The mean water

consumption of mice in the home cage. Each dot indicates one cage, and the data are mean ±

s.e.m. B) The mean fecal weight of mice in the home cage. Each dot indicates one cage, and

the data are mean ± s.e.m. C) The mean urine production of mice. Each dot indicates one cage,

and the data are mean ± s.e.m. D) The weight variation of mice during the home cage assay.

Each dot indicates one cage, and the data are mean ± s.e.m. E) The blood pressure of mice.

Each dot indicates one mouse, and the data are mean ± s.e.m. F) The weight of mice for

intestine analysis. Each dot indicates one mouse, and the data are mean ± s.e.m. G) The length

of intestine of mice. Each dot indicates one mouse, and the data are mean ± s.e.m. H) The ratio

of length and weight. Original data from Figure S4F/G. Each dot indicates one mouse, and the

data are mean ± s.e.m. I)The defecation pattern of mice in OFT of every 5 minutes. Each dot

indicates one mouse, and the data are mean ± s.e.m. J) The minimum current stimulation that

can trigger AP. Each dot indicates one cells. Brain slices from 5 control mice and 5 NVC

dysfunction mice. The data are mean ± s.e.m. K) The RMP of each neuron. Each dot indicates

one cells. Brain slices from 5 control mice and 5 NVC dysfunction mice. The data are mean ±

s.e.m. L) the current stimulation pattern. M) The number of APs in different current stimulation.

Each dot indicates one cells. Brain slices from 5 control mice and 5 NVC dysfunction mice. The

data are mean ± s.e.m. N) Fecal production in different transgenic models in 15min OFT. Each

dot indicates one mouse, and the data are mean ± s.e.m. O) Urine map of mice in the OFLT.

P) The presentative heat map of mice trajectory in the OFLT. Q) The urine rate of mice in the OFLT. R) The running distance of mice in the OFLT. Each dot indicates one mouse, and the data are mean ± s.e.m. S) The time of mice speed distribution of mice in the OFLT. Each dot indicates one mouse, and the data are mean ± s.e.m. T) The Log10 ratio between time in right area and the rest of area. Each dot indicates one mouse, and the data are mean ± s.e.m. U) Representative c-fos IF image in the amygdala after the home cage assay. V) The analysis of c-fos activation after the OFT. Each dot indicates the average c-fos expression level of three mice, and the data are mean ± s.e.m. W) The vasculature system distribution in the amygdala. Red signals present blood vessels.

Tables

Table 1. List of Mice Genetics

Common name	Strain name	Resource
Nestin-cre	B6.Cg-Tg(Nes-cre)1Kln/J	Yanmei Tao, Hangzhou Normal University/ Southeast University
Emx1-cre	B6.129S2-Emx1^{tm1(cre)Krj}/J	Yanmei Tao, Hangzhou Normal University/ Southeast University
grin2d	B6/JGpt-Grin2d^{em1Cflox}/Gpt	GemPharmatech
SMA-creER	Tg(Acta2-cre/ERT2)12Pcn	Bo Zhou, ShanghaiInstitutes for Biological Sciences, Chinese Academy of Sciences
Ai47	Gt(ROSA)26Sor^{tm47(CAG-EGFP*)Hze}	Zilong Qiu, Chinese Academy of Sciences Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences.
Pdgfrβ-creER	Tg(Pdgfrb-cre/ERT2)6096Rha	Xiang Yu, Peking University
grin1	*Grin1*^{tm2Stl}	Yangang Sun, Chinese Academy of Sciences Center for Excellence in Brain Science and Intelligence

		Technology, Chinese Academy of Sciences
Ai14	B6.Cg-Gt(ROSA)26Sor^{tm14(CAG-tdTomato)Hze}/J	Woo-Ping Ge, Chinese Institute of Brain Research (CIBR)
glrb	C57BL/6JGpt-Glrb^{em1Cflox}/Gpt	GemPharmatech
acta2	C57BL/6JGpt-Acta2^{em1Cflox}/Gpt	GemPharmatech

1034

Table 2. List of Primer Pairs (Some primers referred to the published PrimerBank database)

Name		Sequence
grin1	F	AAGAACCTGCAGCAGTACCA
	R	GCAGCAGGACTCATCAGTGT
grin2a	F	TCAGGGACTTTGGAAAACGCT
	R	AAGTCATGCAGGGTTGGTTG
grin2b	F	ACTAACTATCAATGAAGAACGGT
	R	CCAGACCCCAGAGTAACCA
grin2c	F	TGTGGGCCTTCTTCGCTGTCATCT
	R	TGCCATTAGGTACCGTGCCAAAAC

grin2d	F	TGGAGGAGTACGACTGGACAT
	R	CGCACTGACACTACGGAGC
grin3a	F	CTGCACACCAGTCAGAGGTT
	R	TTCGTTGGTTGTCGTGACTC
grin3b	F	TGGGGGTCTACCACTTGTCA
	R	TGGATCTTCTGGCTCGTGTG
ptgs2	F	TTCAACACACTCTATCACTGGC
	R	AGAAGCGTTTGCGGTACTCAT
18s	F	CGGACACGGACAGGATTGACA
	R	CCAGACAAATCGCTCCACCAACTA
cald (siRNA)	S	GUCAAGAGCUUCUUGGAUA
	AS	UAUCCAAGAAGCUCUUGAC

1037

1038 **Table 3. List of Antibodies**

Name	Host	Cat#	Dilution	Company
Anti-NMDAR2D Antibody	Mouse	MAB5578	1:400 (IF) 1:2,500 (WB) 1:1,000 (IP)	Merck

Anti-Smoothelin antibody	Rabbit	ab219652	1:400 (IF)	abcam
α-Smooth Muscle Actin (ACTA2) Rabbit mAb	Rabbit	A17910	1:2,500 (WB)	ABclonal
Anti-NeuN purified Antibody	Guinea Pig	ABN90P	1:800 (IF)	Merck
COX2/PTGS2 Rabbit pAb	Rabbit	A1253	1:400 (IF)	ABclonal
Anti-GLRB antibody	Mouse	EM1710-32	1:400 (IF)	HUABIO
c-Fos antibody	Rabbit	226 003	1:1,000 (IF)	SYSY
Recombinant Alexa Fluor® 488 Anti-Caldesmon/CDM antibody	Rabbit	ab208116	1:400 (IF) 1:2,500 (WB)	abcam
Mice IgG	Mouse	A7028	1:1,000 (IP)	Beyotime
Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488	Goat	A-11001	1:1,000 (IF)	Invitrogen
Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 546	Goat	A-11010	1:1,000 (IF)	Invitrogen
Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488	Goat	A-11034	1:1,000 (IF)	Invitrogen
Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 647	Donkey	A-21447	1:1,000 (IF)	Invitrogen
Goat Anti-Guinea pig IgG H&L (Alexa Fluor® 488)	Goat	ab150185	1:1,000 (IF)	abcam

Goat Anti-Guinea pig IgG H&L (Alexa Fluor® 555)	Goat	ab150186	1:1,000 (IF)	abcam
β-Tubulin Mouse mAb	Mouse	AC021	1:5,000 (WB)	ABclonal
Goat Anti-Mouse IgG, HRP Conjugated	Goat	CW0102	1:10,000 (WB)	CWBIO
Goat Anti-Rabbit IgG, HRP Conjugated	Goat	CW0103	1:10,000 (WB)	CWBIO

1039

Data availability

1041 The code is available at https://github.com/JialabEleven/ with citation. Another data is available

1042 from authors upon reasonable request.

1043

Author contributions

Jie-Min Jia conceived the project, designed the experiments, and supervised this study. Jiayu

Ruan designed mice management, operated all major assay, obtained all major original data,

and wrote the manuscript. Huiqi Xie participated in blood pressure measurement, long-term

whisker stimulation, and behavioral assay. Xinghua Quan made *cald* siRNA interference assay,

and participated behavioral assay. Yiyi Zhang wrote main MATLAB code. Dongdong Zhang

provided primer pairs of *grin1* and provided suggestions. Wentao Wang participated in the

writing of MATLAB code.

Figure1

Figure2

Figure3

Figure4

Figure S1

Figure S2

A

	Input	IgG	NR2D_IP
NR2D_Flag			
Caldesmon			

B

Scramble control   Caldesmon siRNA

131bp — *cald*
98bp — *18s*

C

Relative expression level
0.0271
Scramble siRNA   *cald* siRNA

D

Relative expression level
0.0010
Scramble siRNA   *cald* siRNA

E

SMA
Tubulin

Relative expression level
0.0319
acta2 f/f   *Pdgfrβ-creER:acta2 f/f*

F

Scramble siRNA   Caldesmon siRNA

200μm   200μm   200μm   200μm
200μm   200μm   200μm   200μm

S1BF cortex Caldesmon CD31 DAPI

G

KEGG Pathway Enrichment

count: 20, 40, 60
padj: 0.8, 0.6, 0.4, 0.2

Pvalue

H

Stimulation (12Hz)

CBF Variation (%)
Seconds
30   60   90   120

— Littermate control mice
— *cald* siRNA
— *Pdgfrβ-creER:acta2 f/f*

I

CBF Variation (%) / Seconds
0.0161 0.0144   ns   ns

Peak mean (%)   Time to peak (s)

● Peak mean (%)
● *cald siRNA*
● *Pdgfrβ-creER:acta2 f/f*

Figure S3

Figure S4