Neurovascular Coupling in the Basolateral Amygdala Modulates Negative Emotions

Jiayu Ruan[1,2,3,4], Xinghua Quan[2,3,4], Yiyi Zhang[2,3,4], Huiqi Xie[2,3,4], Dongdong Zhang[2,3,4], Wentao Wang[2,3,4], Bingrui Zhao[2,3,4], Danping Lu[2,3,4], Jie-Min Jia[2,3,4*]

Affiliations

[1] College of Life Sciences, Zhejiang University, Hangzhou, Zhejiang, China.

[2] Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, China.

[3] Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, China.

[4] Laboratory of Neurovascular Biology, Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, China.

*Correspondence: jiajiemin@westlake.edu.cn

ABSTRACT

Emotion induces changes in regional cerebral blood flow (rCBF), a manifestation of neurovascular coupling (NVC), yet whether NVC might feedback to modulate emotion actively remains unexplored. Here, we demonstrated that NVC actively modulates stress-induced negative emotions. We established bidirectional manipulations of NVC in freely moving mice by employing integrated pharmacological, genetic, and arteriolar optogenetic approaches. Our results showed that both systemic and basolateral amygdala (BLA) regional specific NVC deficiencies heightened emotional responses when mice transitioned from a home cage to anxiogenic environments, and local restoration of NVC in the BLA normalized these responses. Furthermore, by combining BLA neuronal activity mapping with behavioral measures of responses to various stressors, we revealed that the spatial patterns of BLA neuronal activation predicted emotional response magnitudes, with hyper- and hypoactive BLA predicting heightened and blunted behavioral responses, respectively, in the NVC-deficient mice. Finally, enhancing NVC, achieved through either pharmacological or genetic manipulations, promoted resilience to chronic-stress-induced NVC impairments and alleviated emotional distress. These findings established NVC as an allostatic program for fine-tuning emotional responses, with implications for understanding and treating emotional disorders.

INTRODUCTION

Emotion is a complex, integrative response to changes in internal and external environments.[1] Emotional flexibility—the capacity to dynamically regulate and adapt emotional responses—is fundamental for mental health and, in some contexts, survival.[2] Deficits in this flexibility can lead to emotional rigidity, a hallmark of emotional dysregulation, strongly associated with psychiatric disorders such as Post-Traumatic Stress Disorder and depression.[3, 4] Therefore, understanding the molecular and cellular mechanisms that modulate emotional flexibility and rigidity holds critical implications for elucidating resilience to psychological trauma.[5, 6]

Regional neuronal activity, for example, elicited by emotional challenge, tightly couples with regional cerebral blood flow (rCBF) changes–a phenomenon known as neurovascular coupling (NVC)–which underlies the blood-oxygen-level-dependent (BOLD) signals widely used in functional imaging.[7-10] Traditionally, NVC has been widely viewed as a passive proxy for local neuronal activity.[11] Yet despite decades of investigation into the biophysical mechanisms of NVC, its active role—if any—in modulating neuronal circuit function and behavior remains unresolved. Recent studies suggest that NVC in the hippocampus may influence neurogenesis, hinting at a broader regulatory function beyond its hemodynamic role.[12]

Particularly in emotional processing circuits, does the cerebrovascular

network in emotional centers, such as the basolateral amygdala (BLA), simply

mirror changes in neuronal firing during stress, or can it actively gate or tune

emotional reactivity? BLA has long been recognized as a central hub for

processing negative emotions, with altered neuronal activity—either

hyperactivity or hypoactivity—linked to emotional dysfunction.[4, 13] The role of

NVC in the BLA in modulating negative emotions, a yet-underexplored

regulatory element, represents the focal point of our study.

In this study, using pharmacological, genetic, and arteriolar optogenetic

approaches, we demonstrated that NVC in the BLA plays a causal role in

modulating stress-induced emotional responses. In contrast, NVC in the

somatosensory barrel cortex (S1BF) appeared to be functionally dispensable

in this context. Our findings revealed that NVC dysfunction leads to context-

inappropriate emotional behaviors—characterized by exaggerated or blunted

reactions—reminiscent of bipolar-like emotional responses in emotional rigidity.

Conversely, enhancing BLA NVC mitigated negative emotional responses.

These results established BLA NVC as an active modulatory mechanism in

emotional processing and suggested that targeting NVC may offer novel

strategies for treating emotional dysregulation in psychiatric disorders.

RESULTS

Mild and strong stress-induced defecation requires BLA activity

Previous studies have demonstrated that stress activates BLA neurons under both mild and intense conditions, such as in the open field test (OFT) and acute restraint stress (ARS).[14, 15] Additionally, stress induces defecation in rodents, a behavior distinct from normal bowel movements, reflecting emotional responses.[16] This stress-induced defecation has long been considered as one of the indicators of negative emotional states.[17, 18] However, the relationship between stress-induced BLA activity and defecation remains unclear.

To investigate this relationship, we measured BLA neuronal activity and defecation behavior in the same mice under three conditions: home cage (minimal stress), OFTs (mild stress), and ARS (intense stress) (Fig. 1a). OFTs were conducted in arenas of varying sizes (40 cm, 35 cm, and 10 cm) to assess the impact of spatial size as a potential variable. Similar to ARS, OFTs induced stress-induced defecation and c-Fos (a marker of neuronal activity) expression,[14] both of which increased with stress levels (Fig. 1b, c; supplementary information, Fig. S1), showing a strong positive correlation (Fig. 1d).

Surprisingly, subtle differences in stress intensity between the 40 cm and 35 cm OFTs led to a distinct increase in BLA c-Fos expression (Fig. 1b), which is also reflected as decreased time spent in the box's center and extended total

running distance in the 35 cm OFT (Fig. 1e, f), which are two conventional

indicators. This phenomenon may be attributed to the smaller arena size (35

cm), which likely induced greater stress in mice by limiting exploratory

behavior.[19] Stress-induced defecation successfully differentiated these mild

variations in stress levels (Fig. 1c), akin to the above two conventional

indicators. These findings supported the perspective that 35 cm OFT induced

more stress than 40 cm OFT, and defecation can be a comparably effective

measure of stress in mild conditions. Furthermore, defecation locations in the

40 cm OFT (Fig. 1g) exhibited a comparable preference for the peripheral zone,

consistent with the typical peripheral zone preference observed in mice (Fig.

1h),[14] further suggesting that stress-induced defecation is a structured and non-

random behavior linked to BLA activity.

To further verify this phenomenon, we employed a chemogenetic approach

to selectively inhibit excitatory neurons in the BLA by expressing hM4Di

receptors (Fig. 1i). This manipulation reduced ARS-induced c-Fos expression

in the BLA (Fig. 1j) and significantly suppressed stress-induced defecation in

both OFT and ARS (Fig. 1k). Except for defecation bots, we assessed

defecation rate (proportion of mice defecating) and latency (time to first pellet)

in the OFT. Chemogenetic inhibition reduced the defecation rate and prolonged

the defecation latency (Fig. 1l), while no changes were observed in time spent

in the box's center and total running distance in the OFT (Fig. 1m, n). Sham

controls showed no effects (Fig. 1l, o).

In summary, BLA activation is essential for stress-induced defecation.

Defecation bouts, rate, and latency serve as quantifiable, sensitive behavioral

measures of stress, reflecting both stress intensity and BLA activation. We

define this combination as the "negative emotion index," a set of non-invasive

behavioral indicators of stress.

Systemic NVC dysfunction via SMC-specific GluN2D loss exaggerates

negative emotion under mild stress

Our previous research identifies a specialized structure within the

neurovascular unit (NVU), termed the Neuro-smooth Muscle Junction (NsMJ),

where neuronal axon terminals directly contact smooth muscle cells (SMCs)

and regulate rCBF via neurotransmitter release (Fig. 2a).[20] We demonstrate

that SMC-specific knockout of *Grin1*, the core NMDA receptor subunit,

significantly reduces functional hyperemia in the barrel cortex (S1BF) during

whisker stimulation.[20] However, the non-core NMDA receptor subunits

contributing to receptor formation in SMCs remain unidentified.[21]

To address this, we first examined the expression of all NMDA receptor

subunits in brain SMCs. We found that only *Grin2d* (encoding GluN2D) and

Grin1 (encoding GluN1) were detectable (Fig. 2b), a result further confirmed by

RNA_{scope} (Fig. 2c; Supplementary information, Fig. S2a, b).[22] We then

generated SMC-specific *Grin2d* conditional knockout (*Grin2d-cKO*) mice to

assess the role of GluN2D in mediating NVC in both the superficial S1BF and

deep BLA (Fig. 2d-k). In *Grin2d-cKO* mice, GluN2D expression was selectively

reduced in SMCs, without affecting other brain cell types (Fig. 2d;

Supplementary information, Fig. S2c-e).

Consistent with our prior findings that *Grin1* deletion impairs NVC, loss of

Grin2d also disrupted NVC in S1BF during whisker stimulation (Fig. 2e-g). To

further explore the impact of *Grin2d* loss on rCBF dynamics in the BLA, we

developed a novel paradigm to detect BLA-local NVC in freely moving mice.

This approach allowed simultaneous monitoring of BLA rCBV, neuronal Ca^{2+}

dynamics, and defecation behavior during the OFT (Fig. 2h).

Although regional cerebral blood volume (rCBV) measurements using

rhodamine-labeled plasma are well-established in superficial brain regions,

their application in deep regions like the BLA required validation.[23] 3D

reconstruction revealed abundant arteriole in the BLA, notably the

corticoamygdaloid artery (coamg), which originates from the internal carotid

artery (Supplementary information, Fig. S3, Video S1), providing structural

support for BLA rCBV detection in dilating and contracting arterioles. We

confirmed that fiber photometry reliably detected rhodamine fluorescence in the

BLA, with a linear signal increase across concentrations from 2.5 to 20 mg/ml

(Supplementary information, Fig. S4a, b). For long-term monitoring, we

selected a 5 mg/ml concentration, as its signal remained stable for about 900

seconds post-injection (Supplementary information, Fig. S4c). All subsequent

rCBV recordings were performed under this optimized condition.

In control mice, NVC in the BLA exhibited robust functional hyperemia

coupled with neuronal activity during OFT exposure (Fig. 2i-k). However,

Grin2d-cKO mice displayed significantly reduced functional hyperemia, despite

elevated neuronal Ca^{2+} signals and c-Fos expression, indicating a decoupling

of rCBV from neuronal activity in the BLA (Fig. 2i-k, Supplementary information,

Fig. S5). Given the correlation between BLA activity and defecation (Fig. 1d),

the heightened BLA neuronal Ca^{2+} and c-Fos might suggest exaggerated

negative emotional responses in *Grin2d-cKO* mice. Indeed, these mice

exhibited significantly increased defecation bouts during the OFT, especially

within the first 5 minutes, reaching 12-fold higher than controls (Fig. 2l).

Additionally, the defecation rate increased, and latency was shortened (Fig. 2m;

Supplementary information, Table S1).

Notably, *Grin2d-cKO* mice also showed a reduction in rearing bouts (Fig.

2n), a classic indicator of heightened negative emotional states.[19] Both

behavioral changes (defecation, rearing) suggested amplified stress responses

in *Grin2d-cKO* mice under mild stress. Importantly, basal rCBV in the BLA,

locomotion, area preference, routine food/water consumption, defecation,

urination, and intestinal health remained unchanged in *Grin2d-cKO* mice, ruling

out confounding factors' impact (Supplementary information, Fig. S6, S7a-n).

These basal physiological parameters further confirmed that *Grin2d-cKO* mice

were in good general health.

To further probe emotional flexibility in *Grin2d-cKO* mice, we used OFT

arenas of varying sizes shown in Fig. 1a to assess *Grin2d-cKO* mice's response

to increasing spatial stress intensities (Fig. 2m, o; Supplementary information,

Table S1). In control mice, the negative emotion index scaled with stress

intensity: as arena size decreased, the negative emotion index shifted (Figs. 1c,

2m, o; Supplementary information, Table S1). In contrast, *Grin2d-cKO* mice

exhibited a rigid, exaggerated emotional response: Their defecation bouts

peaked immediately upon entering the standard 40 cm OFT and remained

unchanged across smaller arena sizes (Fig. 2o). Similarly, defecation rate and

latency failed to reflect the spatial stress gradient (Fig. 2m; Supplementary

information, Table S1). These findings indicated impaired emotional flexibility,

or an emotion rigidity-like state, in NVC-deficient mice.[3, 24]

To exclude non-specific effects, we performed a series of control assays.

We verified that SMC contractile proteins (α-SMA and smoothelin) remained

unaffected (Supplementary information, Fig. S8) in *Grin2d-cKO* mice. Moreover,

GluN2D expression was absent in peripheral vascular SMCs of control mice

(Supplementary information, Fig. S9), ruling out systemic side effects of *Grin2d-*

cKO. Tamoxifen administration alone had no impact on NVC (Supplementary

information, Fig. S10).

In conclusion, GluN2D expression in SMCs was essential for maintaining

functional NVC in both BLA and S1BF. Loss of GluN2D in cerebral SMCs

impaired NVC, decoupled it from neuronal activity, and triggered exaggerated

negative emotional responses under mild stress. These results underscored

the importance of normal NVC for maintaining balanced emotional responses.

Convergent effects of distinct systemic NVC dysfunction models on
214

negative emotional reactivity to mild stress
215

Relying on a single genetic model of NVC dysfunction can lead to biased

conclusions. To address this, we examined a panel of models with selective

disruptions in distinct cell types within the NVU (Fig. 2a), targeting both

established and previously uncharacterized pathways that mediate NVC, and

assessed their impact on stress-induced negative emotional responses.

We first generated double cKO of *Grin1* and *Grin2d* in SMCs, which did not

exacerbate NVC dysfunction beyond that observed with deletion of either of the

genes, functionally confirming that GluN1 and GluN2D likely constitute the

NMDA receptor complex in SMCs for NVC regulation (Fig. 3a-c). Furthermore,

we generated mice with *Grin2d* deletion across all mural cells, including arterial

and venous SMCs and pericytes. Compared to *Grin2d* deletion restricted to

arteriolar SMCs (*SMA-CreER* driven), pan-mural cell deletion (*Pdgfrβ-CreER*

driven) did not further impair functional hyperemia in the S1BF during whisker

stimulation (Fig. 3b, c). Consistent with these results, negative emotion indices

were not further altered in double cKO and pan-mural cell-specific *Grin2d-cKO*

mice (Fig. 3d, e; Supplementary information, Table S1). To rule out systemic

confounds, we confirmed that *Grin2d-cKO* mice did not alter blood-brain barrier

(BBB) permeability, which reported to cause NVC-deficiency (Supplementary

information, Fig. S7o, p).[25]

Next, we disrupted NVC by deleting *Ptgs2* (encoding COX-2) selectively in

neurons, as pharmacological inhibition and *Ptgs2* knockout both impair NVC.[20,]

[26-28] Consistently, neuron-specific *Ptgs2-cKO mice* exhibited significantly

reduced functional hyperemia in the S1BF and shifted negative emotion indices,

closely mirroring the effects of SMC-specific *Grin2d-cKO* (Fig. 3d–h;

Supplementary information, Table S1). In contrast, although COX-2 mRNA and

protein were detected in SMCs (Supplementary information, Fig. S11), SMC-

specific *Ptgs2-cKO* did not affect NVC or emotional responses, highlighting cell-

specific mechanisms.

Then, we deleted *Acta2* in SMCs, encoding α-SMA, a key contractile

protein in SMCs.[29] Surprisingly, only partial reduction of α-SMA was achieved

(It may be due to the fact that N-terminal acetylation of actin is irreversible,

thereby reducing its degradation efficiency in cells[30-33]), contrasting with the

near-complete knockout observed in *Grin2d-cKO* and *Ptgs2-cKO* models (Figs.

2d, 3f, and 3i; Supplementary information, Fig. S1c). However, this partial loss

was sufficient to induce NVC impairments similar to those observed in other

models with near-complete genetic ablation (Fig. 3g, h). Consistently, *Acta2-*

cKO mice showed exaggerated negative emotional responses, mirroring the

phenotypes observed in both *Grin2d-cKO* and *Ptgs2-cKO* (Neuron) models (Fig.

3d, e, g, h).

Lastly, we also explored the impact of a high-fat diet (HFD) treatment,

known to induce NVC dysfunction and emotional dysregulation (Fig. 3j-n).[25, 34]

In this independent, clinically relevant model,[35] 4 weeks of HFD feeding

increased stress-induced defecation bouts, rate, and shorten defecation

latency despite reduced food intake and the absence of BBB disruption (Fig.

3d-e; Supplementary information, Fig. S12).[36]

The consistent emergence of heightened emotional reactivity across six

distinct models of NVC dysfunction—spanning both whole-brain and whole-

body manipulations—supports the idea that intact NVC is a necessity and

actively modulates emotional states under mild stress.

BLA-specific optogenetic NVC suppression amplifies negative emotional

responses to mild stress

The whole-scale NVC dysfunction models discussed above, which require

prolonged induction periods, lack relative spatial and temporal precision in BLA

NVC impairment. These NVC dysfunction models limit the ability to specifically

assess the necessity of intact NVC in BLA for maintaining emotional

homeostasis. To overcome this limitation, we employed optogenetic technology,

which allows precise, temporally controlled manipulations, to acutely suppress

NVC in the BLA by constricting arterioles.[37]

We first generated SMC-specific ChR2-expressing mice (*SMA-*
CreER:Ai32) and used a fiber-optic setup similar to that in Fig. 2h to locally

activate ChR2-expressing SMCs in the BLA while simultaneously monitoring

rCBV dynamics (Fig. 4a, b). After optimizing optogenetic protocols in the home

cage with minimal stress, we identified a protocol that reliably and reversibly

decreased BLA-local rCBV by approximately 3%, with rCBV rapidly returning

to baseline after the cessation of stimulation (Fig. 4c, d). This approach enabled

precise, reversible control of deep brain arteriole contraction in freely behaving

mice.

Optogenetic constriction of arterioles in the deep brain attenuated the

stress-induced increase in rCBV specifically in the BLA during OFT (Fig. 4e-g).

Fourier transform analysis of rCBV dynamics confirmed that OFT stress elicited

functional hyperemia in the BLA, which was effectively suppressed by vascular

optogenetics (Supplementary information, Fig. S13). This acute manipulation

provided a model of localized NVC dysfunction in the BLA, presumably sparing

NVC in other brain regions.

Notably, this local NVC disruption led to a significant shift in the negative

emotion index, notably with an over tenfold increase in the defecation latency

(Fig. 4h, i), consistent with the phenotype observed in whole-brain/global NVC

manipulation models (Fig. 2, 3). Behavioural analysis also revealed that

vascular optogenetics did not affect locomotion anxiety-like states

(Supplementary information, Fig. S14a, b). Together, our acute BLA-specific

optogenetic model and the chronic whole-brain NVC dysfunction models

converge to demonstrate that intact NVC in the BLA was necessary for the

proper modulation of negative emotional responses.

In addition to blocking stress-induced rCBF increases above baseline that

couple with activated BLA during OFT stress, we asked whether transient

below-baseline BLA rCBF dynamics—still far from ischemic—could also trigger

emotional shifts. In other words, would decoupling basal blood supply from

neuronal activity in the BLA lead to an emotional response, even without

external stimuli?

To investigate this, we conducted the same experimental paradigm with a

3% rCBV decrease, but this time in the home cage context (Fig. 4j). ChR2$^+$ mice

exhibited sustained rCBV reduction during optogenetic stimulation period (Fig.

4j, k). Surprisingly, while home cage conditions typically produced only routine

bowel movements (0.23 fecal pellets/15 min), optogenetic activation induced a

marked increase in defecation frequency and bouts (2 fecal pellets/15 min; Fig.

4l), representing a nearly tenfold elevation and reaching about half the level

seen during ARS. Due to limited visibility caused by corncob bedding in the

home cage, we quantified total fecal pellet output at the end of testing but were

unable to measure defecation latency.

Notably, neuronal activity in the BLA was elevated by approximately tenfold

(Fig. 4m, n) following the optogenetic rCBV reduction both in the home cage

and OFT that did not involve ischemia (hypoxia) in the home cage (Supplementary information, Fig. S14c, d). This heightened BLA activity may explain the increased defecation bouts (Fig. 4l). These results demonstrated that targeted below-baseline rCBV in the BLA alone is sufficient to evoke heightened negative emotional states. The below-baseline rCBV manipulation in the home cage recapitulated the NVC dysfunction seen in *Grin2d-cKO* mice during OFT stress in two respects: BLA hyperactivity and increased defecation (Fig. 2 and 3). This suggested that the mice behave as if experiencing stress, despite being in a low-stress home cage environment.

In summary, the convergence of these two lines of loss-of-function evidence, in both OFT and home cage contexts, established that NVC in the BLA is both necessary (in the OFT, Fig. 4e-i) and sufficient (in the home cage, Fig. 4j-l) for the expression of negative emotional states. These findings suggested a direct causal link between BLA NVC and emotional regulation.

Whole-brain and BLA-local disruptions of the SMC GluN2D partner caldasmon heighten negative emotion under mild stress

To explore the molecular mechanisms by which NVC modulates negative emotions, we focused on GluN2D and its downstream signaling in SMCs. Using co-immunoprecipitation and mass spectrometry, we identified proteins interacting with GluN2D in vascular SMCs. KEGG pathway analysis revealed

that caldesmon, an actin-binding protein critical for vascular contraction, was enriched in pathways related to *Vascular Smooth Muscle Contraction* (Fig. 5a).[38] Western blotting confirmed the presence of caldesmon in the protein complex immunoprecipitated by anti-GluN2D antibody (Fig. 5b).

Caldesmon regulates SMC contractility, and its disruption could impair arteriole dilation, thus affecting NVC.[39] Both mRNA and protein analysis indicated that *Cald*/caldesmon is expressed exclusively by perivascular mural cells,[40] ensuring that siRNA-mediated knockdown (KD) minimally impacts the neuronal circuit directly (Fig. 5c-e). This approach could not be applied to *Grin2d*, as it is also expressed in neurons. Whole-brain *Cald* KD, via CSF delivery of siRNA, attenuated NVC in S1BF during whisker stimulation (Fig. 5f-k), confirming *Cald* KD as an effective strategy for disrupting NVC.

BLA-targeted *Cald* KD did not alter caldesmon expression in other regions, such as the nearby VMH (Fig. 5g-i), and functional hemodynamics data confirmed the spatial specificity of this approach (Fig. 5j, k). Behaviorally, global and BLA-targeted *Cald* KD led to BLA neuronal activation and exaggerated negative emotional responses (Fig. 5l-o). This phenotype closely resembled that of *Grin2d-cKO* mice in two ways: BLA hyperactivation (c-Fos expression) and increased defecation behaviors (Fig. 2i-m, 5l-o). Importantly, *Cald* KD in the BLA of *Grin2d-cKO* mice did not further affect the negative emotion index (Fig. 5n, o), suggesting that caldesmon functions downstream of GluN2D in SMCs.

Finally, S1BF-restricted *Cald* KD did not affect BLA caldesmon expression

and the negative emotion index (Supplementary information, Fig. S15). These

findings underscored the region-specific role of BLA NVC in emotional

modulation and highlight that proper NVC function in the BLA is particularly

critical for modulating negative emotions. Together, our results identify

caldesmon as a key effector linking GluN2D to SMC contractility, with BLA NVC

impairment was sufficient to promote emotional distress during the OFT.

BLA-targeted pharmacological rescue of NVC normalized negative hyper-

responsiveness to mild stress

To rescue BLA NVC dysfunction in genetic models, we administered HET0016,

a cytochrome P450 inhibitor that blocks 20-HETE, a vasoconstrictor.[41, 42] We

first confirmed its NVC-enhancing effect through the whisker stimulation assay

after global i.p. injection (Fig. 6a). Dose-dependent assays showed that 0.1 and

1 mg/kg doses significantly reduced stress-induced defecation bouts in ARS

(Supplementary information, Fig. S16a). We adopted 1 mg/kg as the standard

dose and confirmed that Het0016 enhanced S1BF hyperemia upon the whisker

stimulation (Supplementary information, Fig. S16b, c).

In *Grin2d-cKO* mice, both systemic (1 mg/kg, i.p.) and local BLA (10 μM)

HET0016 administration normalized excessive negative emotion indices and

suppressed rearing bouts (Fig. 6a-f). These results demonstrated that localized

pharmacological NVC enhancement can reverse heightened negative emotional responses in whole-brain NVC dysfunction models, underscoring BLA NVC's role in emotional modulation and therapeutic potential. By contrast, HET0016 did not rescue phenotypes in *Acta2-cKO* mice, which directly affects the contract machine of SMCs (Fig. 6g-i), suggesting that SMC contractility is crucial for the observed effects. Moreover, this result indicated that HET0016 likely did not act directly on BLA neurons.

HET0016 was also tested in HFD mice, where local BLA injection partially restored negative emotion indices (Supplementary information, Fig. S17), although the effect was weaker than in *Grin2d-cKO* mice, likely due to broader brain and vascular alterations. Together, these findings indicated that the HET0016 rescue strategy is effective, but requires responsive arterioles.

A topology-based estimation framework of BLA activity for negative emotional reactivity to mild and strong stressors

To better classify exaggerated emotional responses caused by NVC dysfunction, we developed an estimation framework based on the link between BLA neuronal activation and stress intensity (Fig. 7). We used seven contexts with increasing stress levels, ranging from minimal stress (home cage) to intense stress (predator-like threat), simulated by the traditional looming assay[43] and a new virtual reality eagle diving task (VR_{eagle}) (Fig. 7a).

Contextual mapping revealed a spectrum of BLA activity levels, reflected

in c-Fos activation patterns that integrated anatomical distribution and intensity

(Fig. 7a). By comparing these patterns across contexts, we created a similarity

matrix (Supplementary information, Table S2, See Material and methods).

Unsupervised clustering algorithms grouped the seven contexts into four stress

levels (0-3), matching expectations: level 0 (home cage), level 1 (OFT), level 2

(ARS), and level 3 (Looming, VR VR_{eagle}) (Fig. 7a).

Notably, a custom-designed complex OFT—using high-contrast white

flooring to heighten stress—was placed between OFT and predator-threat

paradigms at level 2 with ARS (Fig. 7a; Supplementary information, Fig.

S18a).[44] Looming and VR_{eagle} were both classified as level 3, with VR_{eagle}

slightly higher (Fig. 7a; Supplementary information, Fig. S18b, c). This

framework proved reliable for estimating stress levels across different contexts.

Next, we examined emotional over-responsiveness in NVC dysfunction

models (Fig. 7b). NVC dysfunction disrupted c-Fos activation patterns, which

were crucial for the estimation framework in normal mice. In the *Grin2d-cKO*

mice, the characteristic gradient of BLA c-Fos activity was lost across the seven

contexts (Fig. 7b). Only the home-cage context was comparable between

genotypes (Fig. 7c), aligning with previous findings of similar basal

characteristics (Supplementary information, Fig. S6a).

This framework revealed that *Cald* KD and *Grin2d-cKO* mice were the most

similar groups (Fig. 7b, d, e), consistent with caldesmon's role alongside GluN2D (Fig. 5). Despite impaired NVC, both groups exhibited hyper-activation of BLA neurons. Additionally, the framework ranked "cKO_OFT_40" at Level_2, despite these mice experiencing Level_1 stress, indicating hyper-reactivity, which matched observed behaviors (Fig. 2l, m, Fig. 3d, e). Furthermore, arteriole optogenetics caused a two-level increase in stress (from Level_0 to Level_2) (Fig. 7f), indicating that NVC-deficient mice responded excessively to mild stressors.

Unexpectedly, *Grin2d-cKO* mice exhibited diminished responses to the most intense stressors, with their stress level dropping from Level 3 to Level 2 in response to looming and VR$_{eagle}$ paradigms (Fig. 7g, h) (Supplementary information, Fig. S19a). Moreover, while defecation served as a robust index in level 1-2, it failed to distinguish between NVC-impaired and control mice under strong stress (Supplementary information, Fig. S19b, c). This paradoxical hypo-responsiveness under strong stress contrasted with the hyper-reactivity observed under mild stress, highlighting context-dependent, bipolar manifestations of NVC dysfunction. These findings warrant further physiological and behavioral investigation.

NVC dysfunction induces hypo-reactivity: accelerated recovery from strong, virtual predator stress

Innate fear responses to natural predators (e.g., eagles and cats) follow a five-step process: (1) predator detection, (2) threat evaluation, (3) selection of defensive strategy, (4) execution of defense (e.g., fight or flee), and (5) termination of defense and recovery.[45] The fifth step is particularly important: premature termination increases vulnerability, while delayed termination leads to unnecessary energy loss.[45]

Compared to conventional stress paradigms, innate fear assays like Looming and VR_{eagle} are more complex and multidimensional. Due to the limitation of the defecation behavior, to quantify negative emotional responses, we introduced refined behavioral indices—locomotion speed, freezing duration, and pupil diameter—to provide a comprehensive view of emotional dynamics.

We first assessed behavior in the Looming assay, simulating predator approach. When mice reached a speed of 13–15 cm/s, a rapidly expanding black disk was triggered (Supplementary information, Fig. S18b). During baseline (no stimulus), NVC-impaired and control mice showed no difference in locomotion performance (Fig. 8a blue zone; Supplementary information, Fig. S20a), suggesting intact motor function. Upon stimulus, both groups exhibited freezing behavior, reducing speed to 0–2 cm/s (Fig. 8a yellow zone), indicating normal initiation of fear responses (Step 1-4).

Focusing on step 5—termination of defense and recovery—velocity heatmaps showed earlier recovery in NVC-impaired mice (Fig. 8b).

Quantitatively, they exhibited significantly faster locomotion speed within 21 seconds post-stimulus (Fig. 8c). Additionally, NVC-impaired mice increased rearing bouts after stimulation (Fig. 8d), supporting accelerated recovery. These findings suggested that NVC impairment did not affect the initiation of fear responses but hastens emotional recovery, likely due to lower BLA activation (Fig. 7b; Supplementary information, Fig. S19a).

Next, we upgraded the Looming assay to the VR$_{eagle}$ assay to evaluate each stage of fear responses in greater detail. Pupil diameter, a sensitive index, was added to assess the emotional impact of NVC impairment under more ecologically relevant conditions.[46] Mice were head-fixed and allowed to move freely on a float ball (Supplementary information, Fig. S18c), with real-time tracking of locomotion speed and pupil diameter (Supplementary information, Video S2). During the baseline, no significant differences were observed between control and NVC-impaired mice (Supplementary information, Fig. S20b, c).

The assay had two stages: (1) eagle call (sound only), and (2) rapid visual dive of an eagle image. In stage (1), both groups exhibited slight freezing, but no significant pupil changes (Fig. 8e blue zone), indicating mild threat perception and intact early-stage processing. In stage (2), both groups showed robust fear responses—freezing and pupil dilation (Fig. 8e yellow zone; Supplementary information, Fig. S20d), confirming normal sensory processing and behavioral execution (steps 1–4).

However, after stimulus termination, we observed significantly faster

recovery in NVC-impaired mice. We defined the "defensive offset phase" as the

interval between stimulus offset and the point where pupil size line and

locomotion speed line in the diagram traces crossed (Fig. 8e brown vertical line

to blue/ magenta vertical lines). This phase was markedly shorter in NVC-

impaired mice (Fig. 8f). Across both Looming and VR$_{eagle}$ paradigms, the time

to baseline recovery (Locomotion speed change for the Looming assay, and

pupil size change for the VR$_{eagle}$ assay) was ~12 s in controls, but reduced by

~59% in NVC-impaired mice (Fig. 8g). These results demonstrated a consistent,

cross-paradigm reduction in emotional recovery time, potentially linked to lower

BLA activation levels.

To assess whether the attenuated negative emotion phenotype is predator-

specific, we developed a Cat$_{Video}$ assay presenting a different predatory threat

(Supplementary information, Fig. S18d). Head-fixed mice ran on a treadmill

while watching a cat stalking and pouncing on prey (Supplementary information,

Video S3); retreat behavior and tail rattling were recorded as defensive

readouts.

Consistent with previous paradigms, control and NVC-impaired mice did

not differ in defecation bouts (Supplementary information, Fig. S19b, c). Control

mice exhibited strong defensive behaviors—retreat and tail rattling—indicating

threat perception (Fig. 8h-l). In contrast, most NVC-impaired mice showed little

or no defensive response, such as significantly reduced retreat behavior and

tail rattling (Fig. 8h, j-l).

These results confirmed that the diminished innate fear response caused by NVC impairment was not predator-specific but generalized across various aversive contexts. The Cat$_{Video}$ assay, alongside the Looming and VR$_{eagle}$ data, validated that NVC impairment consistently lowers stress levels in strong stress contexts, leading to blunted defense behavior. This highlighted that intact NVC was crucial for calibrating the appropriate emotional intensity and duration during threat exposure, while its disruption elevated the response threshold and shortens defensive states.

Evaluation of NVC-dysfunction mice's emotional responsiveness to strong, real social aggressor

To extend our findings from virtual paradigms to real-world threats, we designed an aggressive social stress assay to assess how NVC-dysfunction mice respond to actual social stressors. This approach allowed us to evaluate whether NVC-impaired mice exhibit mismatched responses when confronted with real-life stress.

C57BL/6 mice, with either intact or impaired NVC, were co-housed with pre-screened aggressive CD1 mice in a new cage divided into small and large compartments by a metal lid (Fig. 9a). The mice were free to move between zones during a 10-minute trial, and social interactions—particularly fighting—

were recorded. As in previous assays, defecation bouts did not differentiate

NVC status (Supplementary information, Fig. S19b, c). NVC-normal mice

engaged in approximately 20 seconds of fighting (~4% of the session time) (Fig.

9b), which is less than what is typically observed in classic social defeat models

where C57BL/6 mice are introduced into the home cage of resident CD1

aggressors.[47] Based on the estimation framework, these mice were classified

at stress Level_1 (Fig. 9c).

In contrast, NVC-impaired mice were classified as stress Level_0—

equivalent to baseline conditions—even while facing aggressive social stress

(Fig. 9d). This indicated emotional hypo-responsiveness. Behavioral analysis

confirmed this observation: although both groups avoided sharing the small

space with the CD1 aggressor, NVC-impaired mice showed significantly

reduced avoidance (Fig. 9e). While 98% of control mice fled during fights, only

37% of NVC-impaired mice actively escaped (Fig. 9f), suggesting an impaired

defensive drive. Additionally, NVC-impaired mice experienced fewer attacks

(1.9 vs. 4.6 fights per mouse on average) (Fig. 9g). However, both groups

exhibited similar levels of sniffing toward the CD1 mice (Fig. 9h), ruling out

sensory deficits.

Together, these findings reinforce the crucial role of NVC in generating

context-appropriate negative emotional responses. Disruption of NVC resulted

in misaligned stress reactivity, weakened defense mechanisms, and blunted

emotional responses—highlighting a mechanistic link between cerebrovascular

553 dysfunction and affective disorders.

Genetic NVC enhancement prevents strong CRS-induced NVC dysfunction and confers resistance to the resulting emotional distress

To further explore the intrinsic genetic regulators that could enhance NVC and

mitigate stress-induced emotional dysregulation, we focused on identifying

anion channels that might promote NVC. As we had previously shown, NVC

impairment was induced by targeting cation channels (NMDA receptors) in

SMCs (Fig. 2a). We hypothesized that inhibiting anion channels could enhance

NVC and potentially counteract stress-induced negative emotional responses.

Through transcriptomic database screening, we identified *Glrb* (encoding

GlyRβ), a ligand-gated chloride channel, as being highly expressed in venous

SMCs (Supplementary information, Fig. S21).[40] RNA$_{scope}$ and immunostaining

further confirmed that GlyRβ was predominantly localized in venous SMCs (Fig.

10a–d). To assess its function, we generated *Pdgfrβ-CreER:Glrb$^{f/f}$* mice to

delete *Glrb* specifically in pan-mural cells. GlyRβ expression was significantly

reduced in these mice (Fig. 10c, e–f). Additionally, we identified that GlyRβ

colocalized with gephyrin *in vitro* (Fig. 10g), a scaffolding protein commonly

found in inhibitory postsynaptic sites, suggesting that GlyRβ may cluster on the

post-junctional membrane of glycinergic NsMJ. This hypothesis was supported

by confocal laser scanning microscopy (Fig. 10h) and electron microscopy (Fig.

10i), which showed frequent direct contact of neuronal axons onto venular

SMCs.

In functional assays, whisker stimulation in *Glrb-cKO* mice led to enhanced

functional hyperemia (Fig. 10j, k), indicating potentiated NVC. However, this

effect was absent when *Glrb* was deleted specifically in arteriolar SMCs

(Supplementary information, Fig. S22a, b), highlighting the crucial role of non-

arteriolar mural cells. Behaviorally, NVC-enhanced mice exhibited similar

emotional indices to controls under mild stress (OFT) but showed significantly

fewer defecation bouts under ARS, suggesting enhanced emotional resilience

(Fig. 10m). This effect was absent in SMC-specific *Glrb-cKO* mice

(Supplementary information, Fig. S22c).

Given that NVC enhancement conferred resilience to stress, we tested

whether genetically restoring NVC could reverse the effects of *Grin2d-cKO*. We

generated a double cKO model targeting both *Grin2d* and *Glrb* in mural cells

(*Pdgfrβ-CreER:Grin2d$^{f/f}$:Glrb$^{f/f}$*) to restore the balance between excitatory and

inhibitory channels (Fig. 10l). Results confirmed that compared to *Grin2d-cKO*

mice, the dual knockouts preserved whisker-stimulation-induced hyperemia

(Fig. 10j, k) and normalized emotional indices in the OFT (Fig. 10m, n;

Supplementary information, Table S1), indicating that NVC restoration can

reverse stress hypersensitivity.

To rule out off-target effects of *Glrb* deletion, we used the NVC inhibitor

pinacidil, a Kir6.1 activator.[48, 49] Intraperitoneal injection of pinacidil reduced

functional hyperemia (Supplementary information, Fig. S23a–c) but also

decreased basal cerebral perfusion (Supplementary information, Fig. S23d),

suggesting systemic side effects. Global administration of pinacidil abolished

ARS-induced defecation in all mice (Supplementary information, Fig. S23e),

likely due to effects on SMCs in the cardiac and gastrointestinal systems.[50] To

isolate NVC-specific effects, we locally injected pinacidil into the BLA

(Supplementary information, Fig. S23a). This intervention abolished the

resilience conferred by NVC enhancement, returning defecation bouts to

baseline levels (Supplementary information, Fig. S23f), confirming that the

resilience originated from local BLA NVC enhancement rather than systemic

actions.

Finally, we examined whether enhanced NVC could provide long-term

resilience to chronic stress, using a chronic restraint stress (CRS) model (Fig.

10o).[51, 52] After CRS exposure, control mice exhibited anxiety-like behaviors

(reduced open-arm time in the EPM), constipation, and attenuated functional

hyperemia (Fig. 10j, k, p, q; Supplementary information, Fig. S24). These

effects were absent in NVC-enhanced mice, indicating resistance to CRS (Fig.

10j, k, p, q). In the forced swimming test (FST), NVC-enhanced mice exhibited

reduced immobility regardless of CRS history (Fig. 10r), consistent with ARS-

induced defecation bouts remaining unchanged post-CRS (Supplementary

information Fig. S25). These data further confirmed their enhanced emotional

resilience.

Locomotion and anxiety-like assessments excluded locomotor or basal

anxiety confounds (Supplementary information, Fig. S26a, b). However, NVC-

enhanced mice exhibited weight loss during CRS (Supplementary information,

Fig. S26c), suggesting incomplete protection. Since CRS is known to induce

BBB disruption and emotional dysfunction, we also evaluated BBB integrity.[53-

55] Notably, 28 days of CRS did not induce IgG leakage, suggesting intact BBB

integrity (Supplementary information, Fig. S27). These results suggested that

enhancing NVC confers protection against both acute and chronic stress.

DISCUSSION

In this study, we demonstrate that bidirectional control of NVC at both systemic

and region-specific levels establishes its necessity and sufficiency for

modulating negative affect. Thus, NVC exerts causal control over negative-

emotion programming, thereby conceptually changing our understanding of

NVC. Far from being a passive read-out of neuronal activity, BLA-local NVC

operates as an indispensable neurovascular modulator of the BLA network.

Disrupting this modulator in the BLA—but not in the primary somatosensory

cortex—selectively alters negative emotional outputs, underscoring a region-

specific modulation. These findings redefine NVC as an active component of

affective circuitry rather than a mere hemodynamic by-product and prompt a re-

evaluation of its broader roles throughout the brain.

This study introduces a fundamentally independent therapeutic avenue by

targeting the mural cell-expressed *Grin2d*, *Cald*, and *Glrb* genes. In particular,

Glrb-cKO mice has been shown to confer resilience to chronic aversive stress,

underscoring the need to develop optimized mural cell–specific delivery

systems. Because optogenetically reducing baseline rCBV below in the BLA is

sufficient to induce negative emotion (Fig. 4j-n), future pharmacological or

neuromodulatory strategies that stabilize or increase this baseline blood flow

may offer viable treatments for affective disorders. Collectively, our findings

open a new path for intervening in psychopathological progression by

enhancing NVC, thus complementing currently existing therapeutic approaches.

Our findings demonstrate that baseline hemodynamic states alone can

modulate emotion (Fig. 4j-n). This implies hemodynamic or vascular factors as

potential initiators of negative affect, aligned with recent causal evidence linking

cardiac rate to affective status.[56] This finding also suggest that resting-state

fMRI measures of amygdala connectivity reflect not only synaptic interactions

but also baseline vascular dynamics, underscoring the need to integrate

hemodynamic factors into translational models of affective circuitry.[57]

However, a key limitation of this study is that the mechanism by which SMC

contraction induced by optogenetics triggers BLA neuronal c-Fos upregulation

and negative affect remains unclear. Previous studies have reported that

increased hemodynamic flux into penetrating arterioles in the cortical cortex can suppress action potential firing in surrounding neurons, hinting at a potential role for vascular-neuronal coupling (VNC) in mediating these effects.[58-60] This raises the possibility that vascular changes may directly influence neuronal activity through mechanical or metabolic signaling pathways, a hypothesis warranting further investigation to dissect the causal chain linking vascular dynamics to affective states.

We developed a refined negative emotion index (bouts, latency, and rate) as a non-invasive and reliable behavioral readout of emotional responses across multiple contexts. Unlike the time-in-center parameter–which only applies to the OFT–this defecation index can be readily applied to diverse behavioral tests. Previous studies have identified a direct monosynaptic circuit between the amygdala and Barrington's nucleus (Bar), a brainstem center that may be implicated in defecation control.[61, 62] Based on this, we hypothesized that optogenetic stimulation of SMCs in the BLA, through yet unknown mechanisms, activates BLA neurons, which then signal to Bar directly or indirectly (Supplementary information, Fig. S28).

Importantly, our single integrated platform accomplished all five of the following in the same freely behaving mouse: bidirectional optical/pharmacological control of NVC, simultaneous recording of neuronal activity, concurrent measurement of hemodynamic signals, precise targeting of deep-brain regions (e.g., BLA, Bar), and free behavior monitoring. By

integrating these capabilities into one system, we usher NVC research into a

new era–enabling causal, brain region-specific dissections of neurovascular

"calculations" underlying affective behavior.

Traditional neuronal computation studies have primarily focused on

decoding neuronal electrical activity at the synaptic level, emphasizing how

neuronal networks transmit and integrate electrical signals to regulate

perception, decision-making, and behavior.[63] We propose that the

completeness of neuronal computation could extend beyond synaptic activity

and integrate "neurovascular modulatory contribution" to reveal how vascular

and hemodynamic changes modulate neuronal circuits in a flexible and fine-

tuning manner. The concept of neurovascular computing not only expands our

understanding of "computation" in neuroscience but may also offer new

biological insights for future artificial intelligence architectures that aim to be

able to flexibly and finely tune affective responses.[64]

Negative emotion linked to innate fear is a key survival mechanism. For

example, in high-stress situations such as combat or natural disasters, the

ability to regulate emotions profoundly influences decision-making and survival

outcomes.[45, 65] Our findings show that accelerated recovery from defensive

states in the NVC-dysfunction model may increase vulnerability to subsequent

threats (Fig.8),[66] offering insight into the evolutionary conservation of NVC

across species, as appropriate regulation of innat responses can be a matter

of life or death in the animal kingdom and, at times, in human society as well.[67]

MATERIALS AND METHODS

Animal and animal care

All mouse procedures complied with the Institutional Animal Care and Use Committee (IACUC) guidelines of the School of Life Sciences, Westlake University. All mice were housed in the Westlake University Laboratory Animal Resource Centre with standard animal facilities, which provided a 12-hour light/dark cycle at 25 °C. No more than six mice were fed in each cage, and standard chow/water was provided to the mice *ad libitum*. All the genetic information of mice is in Supplementary information, Table S3.

Home cage monitor assay

The blood pressure of mice was measured by Kent Noninvasive blood pressure measurement system for animals (Kent/Biopac, USA) before the home cage monitor assay. Each home cage only contains four mice. In each cycle (three days), 60 g of food, 340 g of water (including 100 g bottle weight), and 100 g of padding were provided. Weight of mice, food/water consumption, fecal particles, and urination amount (counted by increased padding weight) were measured by calculating the mean data. Three cycles of monitoring were performed for each home cage. The intestine system inspection was performed after three cycles of monitor. After the assay, mice were fed alone in a clean new cage for

90 min and then used for brain extraction to detect c-Fos expression.

Open field test

The setup of open field test (OFT) referred to published papers.[14, 19] Each

mouse was placed in the central area of a 40*40 cm, 35*35 cm, or 10*10 cm

box for the OFT. The camera recording was 15 min. For the chemogenetic

manipulation assay, the C21 intraperitoneal injection was 45 min before the

OFT. After the assay, mice were fed alone in a clean new cage for 90 min and

then used for brain extraction to detect c-Fos expression. For the BLA NVC

recording, optic fibers were implanted at BLA (AP: -1.6 mm; ML: +/-3.45 mm;

DV: -4.0 mm) with the injection of hSyn-Gcamp-8f virus (*SMA-creER:Ai32* mice

do not need virus injection. Detailed information about virus mentioned later)

two weeks before the assay. Before the formal assay, three days of adaptation

were performed. On the day of the assay, orbital injection of Rhodamine (Sigma,

Cat# R9379-1G, USA) 100 µl 5 mg/ml 15 minutes before the recording. 405 nm

light was used for bassline control, 473 nm light recording for the Gcamp-8f

signal (For *SMA-creER:Ai32* and *SMA-creER:Ai47* mice, using 473 nm light for

optogenetic stimulation), and 580 nm light recording for the Rhodamine

signal. Recording sustained 15 minutes (For NVC recoding)/45 minutes (For

optogenetic stimulation plus rCBV recording). Untouched-rearing bouts

measurement in the 1st 5 min. Defecation bouts measurement keep 15 min.

Tube stress assay

A 50ml tube was used to restrain the mice to perform acute restraint stress (ARS) and chronic restraint stress (CRS).[54, 68] The tube cover and wall will be drilled to ensure normal breathing. For ARS, the mice were restrained in the tube for 15 min, and then the fecal production was measured. For the ARS plus chemogenetic manipulation assay, the C21 intraperitoneal injection was 45 min before the tube stress assay. After the assay, mice were fed alone in a clean new cage for 90 min and then used for brain extraction to detect c-Fos expression. For CRS, the mice were restrained in the tube for 180 min/day (random time for restraint), constant for 28 days. Fecal production and Bristol Score was measured at the end of each seven days within 60 minutes in a empty home cage box.

***In vivo* virus injection**

rAAV2/4-Glyt2-EGFP-WPRE-hGHpolyA (5E+12VG/ml) was used to label glycinergic neurons. Two weeks before the assay, bilateral medullary reticular formation injection (500 nl on each side, AP: -3.5 mm; ML: +/-0.84 mm; DV: -2.8 mm)[69] was administered. AAV2/5-mCaMKIIa-hM4D(Gi)-EGFP-ER2-WPRE-pA(2.30E+12VG/ml) and AAV2/5-mCaMKIIa-EGFP-ER2-WPRE-pA(2.30E+12VG/ml) were used for the chemogenetical manipulation. Two

weeks before the assay, bilateral amygdala injection (150 nl on each side, AP:

-1.6 mm; ML: +/-3.45 mm; DV: -4.1 mm) was administered. AAV2/9-hSyn-

jGCaMP8f-WPRE-pA was used for the neuronal calcium measurement. Two

weeks before the assay, bilateral amygdala injection (150 nl on each side, AP:

-1.6 mm; ML: +/-3.45 mm; DV: -4.1 mm) was administered.

Immunofluorescence staining

For Immunofluorescence (IF) staining, after anesthesia with 1% pentobarbital

sodium, the mice were perfused with 1X PBS (5 min) and 4% PFA in 1X PBS

(5 min) through the left ventricle to empty blood and fix tissue. Afterwards, tissue

was extracted and fixed by 4% PFA in 1X PBS for 6 hours at 4 °C. For samples

like the brain and peripheral organs, samples were sliced into 50 μm (Leica,

Histo-VT1200S, DEU). Then, treated samples with 1% Triton X-100 (Diamond,

Cat# A110694-0500, CHN) and 0.25% NDS (Jackson Resources, Cat# 017-

000-121, USA) in 1X PBS for 60 min at room temperature. After that, samples

were rewashed three times with 1X PBS.

In the staining process, the primary antibody diluted with 0.1% Triton X-

100 in 1X PBS was used to incubate samples overnight at 4 °C. After the

incubation, samples were washed with 1X PBS 10 min (3 replicates). The

corresponding secondary fluorescence-conjugated antibody diluted with 0.1%

Triton X-100 in 1X PBS was used to incubate samples for 120 min at room

temperature. After the incubation, samples were washed with 1X PBS for 10

min (3 replicates). After that, 1000X diluted DAPI (Beyotime, Cat# C1002, CHN)

was used for nucleus staining. Finally, samples were covered by a glass slice

with the mounting medium (SouthernBiotech, 0100-01, USA). Zeiss LSM800

(Zeiss, LSM800, DEU) microscopy was used for photography. A list of

antibodies is in Supplementary Table S4.

RT-PCR

For cultured cells, after obtaining pure sorted EGFP$^+$ cells from the cortex, cells

were cultured in 6 cm dishes (Corning, Cat# 430166, USA) with SMCM. After

cells reached around 90% density, Trizol (Sangon Biotec, Cat# B511311-0100,

CHN) was used for the RNA extraction. For brain tissue, after mice were

anesthetized with ice, the cortex or cerebellum of the mice was directly taken

out, and TRIzol was used for RNA extraction. After extraction, all RNA samples

were reverse transcribed to cDNA (50 ng/μl) using HiScript III 1st Strand cDNA

Synthesis Kit (Vazyme, Cat# R312-02, CHN). For the Real-Time Polymerase

Chain Reaction (RT-PCR), each reaction contained 10 μl of Green Taq Mix

(Vazyme, Cat# P131-AA, CHN), 2 μl of cDNA, 2 μl of primer pairs (10 μM), and

6 μl of RNAase-free water. Agarose gel electrophoresis used 2 % agarose gel

(BIOWESTE, Cat# 111860, ESP) and ran at 220 V for 30 min. RT-PCR gel

images were obtained from a Tanon imager (Tanon, 1600 Geil imager, CHN). A

list of primer pairs is in Supplementary Table S5.

RNA$_{scope}$ in situ hybridisation

Horizontal sections (15 µm) of tamoxifen administrated *SMA-creER:Ai14* (for

Grin2d), *Pdgfrβ-creER:Ai14* (for *Glrb*) and C57/B6 *wild-type* (for *Grin2d* and

Grin1 probe validation) mice brains were obtained by CM1950 (Lecia, Histo-

CM1950, DEU). All RNA scope *in situ* hybridisation productions were obtained

from ACDbio, including RNA$_{scope}$ Multiplex Fluorescent Reagent Kit v2 (ACDbio,

Cat# 323100, USA), *Grin2d* probe (ACDbio, Cat# 425951-C3, USA), *Grin1*

probe (ACDbio, Cat# 431611, USA), and *Glrb* probe (ACDbio, Cat# 1225261-

C1, USA). All operations were carried out following the reagent instructions of

ACDbio company and the published paper.[20, 70]

siRNA cisternal magna/BLA local injection

Cisterna magna injection protocol referred to Cannula Implantation into the

cisterna magna of Rodents (https://doi.org/10.3791/57378) and paper

published by our laboratory.[20] 1 nM siRNA was mixed with FITC (10 mg/ml,

Sigma, FD40S-1G, USA) for injection (20 µl in total). The scalp was opened 10

min after the injection, and the microscope was used to detect the FITC signal

(Zeiss, Axio Zoom v16, DEU) in perivascular spaces. Only successfully injected

mice remain. Whisker stimulation was performed 48 hours after siRNA injection,

followed by 40 cm OFT. As for BLA local siRNA injection, 150nl of 1nM siRNA

was injected in two sides of BLA (AP: -1.55 mm; ML: +/-3.40 mm; DV: -4.00

mm), and 40 cm OFT was performed 48 hours after siRNA injection. Brain slice

collection was performed 90 minutes after OFT for IF staining. Sequence of

siRNA can be found in the Supplementary information, Table S5.

Cell culture

SMA-creER:Ai47 Post-born 0-day (P0) mice were anesthetized with ice.

Surgical tweezers were used to quickly extract the cortex and then transfer it

into the 0.25% Trypsin (Thermo Scientific, Cat# 25200072, USA) for 15 min at

37 °C. Next, FBS (Gibico, Cat# 10099, USA) was added to block digestion.

Then, cells were filted by a 40 μm filter (Falcon, Cat# 352340, CHN). Finally,

cells were centrifuged 1,600*g at 4 °C for 5 min and resuspended by SMCM

(ScienCell, Cat# 1101, USA) in T25 flasks (Corning, Cat# 430639, USA). When

cells reached approximately 80% full of the bottom surface of T25 flasks,

tamoxifen was administrated to cells. Tamoxifen was dissolved in DMSO

(Solarbio, Cat# D8371, CHN), and the final concentration of tamoxifen was 5

μM. After a four-day culture, EGFP in cells were expressed. Cells were sorted

by MA-900 (Sony, FCCF-MBR, JPN) to obtain EGFP+ cells. Sorted cells were

cultured in SMCM for further assay. The purity of cells sorted by the same

method was verified in a paper published by our laboratory.[71] For data in Fig.

10f, samples were directly sorted from Pdgfrβ$^+$ cells in brain lysis by using Anti-

Pdgfrβ antibody (without cell culture). Then, Pdgfrβ$^+$ cells were quickly

resuspended by Trizol for RNA extraction. A list of antibodies is in Table S44. In

Fig. 10g, Human Umbilical Vein Smooth Muscle Cells (HU-vSMC, Sciencell,

Cat# 8020, USA) were used for cell culture.

Tamoxifen and drug administration for mice

Tamoxifen (MCE, Cat# HY-13757A, USA) was dissolved in corn oil (Aladdin,

Cat# C116023-500ml, CHN) and orally administered to mice (10 mg/ml, 200 μl

per mouse per day). For *Acta2* cKO, tamoxifen was given from P21 to P24.

Otherwise, tamoxifen was orally administered to mice from P60 to P63.

For drug delivery, intraperitoneal injection was used for global

administration. The total volume is 200 μl. For local administration into the BLA

(AP: -1.6 mm; ML: +/-3.45 mm; DV: -4.0 mm), a cannula drug delivery system

(RWD, Cat# 800-00245-00, CHN) was employed. The concentrations used for

local delivery were 10 μM for Het0016 and 50 μM for pinacidil. The total delivery

volume was 1 μl (at a rate of 250 nl/min), and behavioral assays were

conducted 5 minutes after Het0016 administration or 25 minutes after pinacidil

administration.

Western-blot

For tissue sample collection, samples were ground with RIPA (Thermo Scientific, Cat# 89901, USA) in the tube on ice for 30 min. For cells, after they reached around 90% density, the RIPA was used for protein extraction on ice for 30 min. Protease Inhibitor Cocktail (CWBIO, Cat# CW22008, CHN) was added into RIPA to prevent protease. After that, samples were centrifuged at 12,000*g at 4 °C for 5 min, and the supernatant was collected. Concentrations of proteins were measured by Micro BCA Protein Assay kit (CWBIO, Cat# CW2011S, CHN), and all samples were boiled with SDS-PAGE loading buffer (CWBIO, Cat# CW0027S, CHN) for 10 min at 100 °C. 8% SDS-polyacrylamide electrophoresis gels were used for gel running. After 90 min, gels running at 120 V (80 V for the first 30 min), gels were transferred to 0.45 μm PVDF membranes (Millipore, Cat# IPVH00010, USA) at 400 mA for 120 min. After transfer, PVDF membranes were blocked by 5% non-fat milk in 1X TBST buffer for 60 min at room temperature with a mild horizontal shake.

After the block, the primary antibody diluted with 5% non-fat milk in 1X TBST buffer was used to incubate the membrane overnight at 4 °C with a mild horizontal shake. After the incubation, the membrane was washed by 1X TBST 10min (3 replicates). The corresponding secondary HRP-conjugated antibody diluted with 5% non-fat milk in 1X TBST buffer was used to incubate the membrane for 45 minutes at room temperature with a mild horizontal shake. After the incubation, the membrane was washed with 1X TBST for 10 min (3

replicates). ECL kits (Abbkine, Cat# BMU102-CN, CHN) and gel image system

(GE, GEL Imaging System AI680RGB v.2.0, USA) were used for exposure

photography. A list of antibodies is in Supplementary information, Table S4.

Whisker stimulation

Mice were anaesthesia with 1% pentobarbital sodium and open the scalp, then

mice stable fixated on the holder. The electro-motor-driven brush stimulated the

whisker longitudinally for 1 minute at a frequency of 12 Hz. The principle of laser

speckle contrast imaging has been described in the previous paper, and the

operation referred to in the published paper.[20, 23, 72] Laser Speckle Image

System (RWD, RFSLI ZW/RFSLI III, CHN) was used to record the CBF change

in the ROI of the opposite barrel cortex of mice. The recording parameters were

512*512 resolution, 10 FPS, and 2-minute recording.

Home cage optogenetic stimulation and recording

Optogenetic stimulation referred published papers.[37, 73, 74] *SMA-creER:Ai32*

and *SMA-creER:Ai47* mice were used for optogenetic stimulation. Tamoxifen

was administered to mice 18 days before tha assay. Then, optic fibers were

implanted at BLA (AP: -1.6 mm; ML: +/-3.45 mm; DV: -4.0 mm) two weeks

before the assay. Before the formal assay, three days of adaptation were

performed. On the day of the assay, orbital injection of Rhodamine 100 µl 5

mg/ml 15 minutes before the recording. 405 nm light was used for bassline

control, 580 nm light recording for Rhodamine signal for 45 min, and 473 nm

light for optogenetic stimulation was performed at the 2nd 15 min of the 45 min

recording. The parameters of the stimulation protocol included the stimulation

period, interval period, recording period, total frequency, and laser power. The

optimal protocol shown in Fig. 4c is 85 ms stimulation, 5 ms interval, 5 ms

recording, 10 Hz frequency, 0.939 mW laser power. During the 45-minute

recording, mice always stayed in their home cage, but all fecal particles were

removed before the assay. After 45 min recording, fecal particles were recorded.

Tissue transparency

Mouse brain tissues were washed three times with 1×PBS at room temperature,

each lasting two hours, and were processed on a shaker at 40 rpm.

Subsequently, a commercial tissue clearing kit (Nuohai, Cat# NH-CR-210701,

CHN) was used to perform the clearing procedure according to the

manufacturer's standard protocol. The cleared tissues were subjected to three-

dimensional imaging using a light-sheet fluorescence microscope (Nuohai,

Cat# LS18, CHN) with a laser wavelength of 561 nm. After imaging, the

acquired data were preliminarily processed using LS18 image stitching

software to generate complete three-dimensional image datasets. The datasets

were then imported into ImageJ software for downsampling, with a

downsampling factor of 0.25 in the x, y, and z directions. The downsampled

datasets were converted into coronal section stacks using the reslice function

in ImageJ. The converted datasets were further imported into the BIRDS plugin

and registered against the whole-brain atlas (Allen Mouse Brain Common

Coordinate Framework version 3, CCFv3). After registration, Imaris Surface

modules corresponding to different brain regions were created based on the

registration results. Using the Surface modules of each brain region, the

acquired three-dimensional data were segmented, and the Surface modules of

vasculature for each brain region were generated based on the fluorescent

signals. Finally, vascular volume data were obtained by analyzing the volumes

of arterial vessels and vasculature in each brain region.

Immunoprecipitation assay

tdTomato$^+$ sorted aSMCs from P0 *SMA-creER:Ai14* mice were used for

lentivirus infection and further immunoprecipitation (IP) assay. Sorted cells were

cultured in 6 cm dishes with SMCM. After cells reach around 90% density, add

lentivirus (Lenti-EF1-EGFP-CMV-grin2d-3XFlag-WPRE) into the culture and

keep for one week to over-express GluN2D. MA-900 sorted out EGFP and

tdTomato double-positive cells, which were lentivirus-infected aSMCs. These

cells were then cultured in 10 cm dishes. After cells reached around 90%

density, cells were collected by RIPA reagent. The Protein G Magnetic beads

kit (Epizyme, Cat# YJ002, CHN) was used for the immunoprecipitation assay.

GluN2D and IgG control antibodies were mixed with Protein G Magnetic beads

to pull down GluN2D and its related protein. Pull-down products were verified

by Western-blot and analyzed through mass spectrometry. A list of antibodies

is in Supplementary information, Table S5.

High-fat diet challenge assay

Set up of High-fat diet (HFD) challenge referred to published papers.[25, 34, 36]

Normal fat die (NFD) mice used as control. Mice were given high-fat food

(SYSE, Cat# PD6001, CHN) for four weeks. The weight of mice and food

consumption were measured before the assay and at the end of each week.

The home cage fecal production was measured using the same method

mentioned in the *home cage monitor assay* part. 40 cm OFT assay and whisker

stimulation assay were performed after four weeks of HFD challenge.

BBB leakage inspection (evens blue and IgG)

Assay setup referred to published papers.[36, 75] 2% evens blue (Merck,

Cat#E2129, USA) dissolved in saline was used for orbital injection (4 ul/g). After

45 min, the mouse was anesthetized, and the brain was extracted for

appearance inspection for BBB leakage using a microscope (Zeiss, Axio Zoom

v16, DEU). The brain was weighted and then homogenized using 1 ml of 50%

trichloroacetic acid (Merck, Cat#190780, USA). Then, samples were

centrifuged at 10,000*g at 4 °C for 20 min, supernatants were collected, and

absorbance detection at 620 nm was performed. For IgG inspection methods,

see the *Immunofluorescence staining* part above.

Colorectal inspection

The ENDOQ Colonoscopy system took representative images of colorectal

tissue. A professional physician performed scoring of the endoscopic images in

a double-blind manner to ensure unbiased assessment.

Hematoxylin and eosin staining

For hematoxylin and eosin (HE) staining, after anesthesia with 1% pentobarbital

sodium, the mice were perfused with 1X PBS (5 min) and 4% PFA in 1X PBS

(5 min) through the left ventricle to empty blood and fix tissue. Afterward, tissue

was extracted and fixed by 4% PFA in 1X PBS for 6 hours at 4 °C. After a

paraffin block, tissues were sliced into 8 μm sections and examined following

HE staining.

Topology-based estimation framework

The topology-based estimation framework is based on the Structural Similarity

Index (SSIM) and the cluster analysis of SSIM (See *Statistical analysis software*

and code part for details). In brief, translating BLA's c-Fos positive cells location

and density with different contexts into SSIM. Then, the cluster analysis was

used to classify the SSIM of different contexts to build a library of topology-

based estimation framework. To predict a subject's "real" stress level, the SSIM

of this subject will be clustered together with the topology-based estimation

framework library to locate the subject's stress level.

1004

Complex-Open field test

The complex-open field test (complex OFT) setup referred to a published

paper.[44, 76] Each mouse was placed in the central area of the left part of the

complex OFT box. One side of the complex OFT box has a white bottom (the

mouse is put on this side) and a black bottom on the other. The camera

recording was 20 min. After the assay, mice were fed alone in a clean new cage

for 90 minutes and then used for brain extraction to detect c-Fos expression.

1012

1013 **Looming assay**

1014 The assay was carried out in a 35 cm*35 cm OFT box. The settings of the assay

1015 were based on the published paper.[43] Modifications include: Modifications

1016 include:

- mice will be put in the OFT box to adapt for 10 minutes before starting the assay;

- we only use a single camera at the top of the box to record;

- we stimulate the mice when they move and are not close to the box wall rather than continuously stimulating them.

Each mouse performs one trial of stimulation. After the assay, mice were fed alone in a clean new cage for 90 min and then used for brain extraction to detect c-Fos expression.

VR$_{eagle}$ assay

Assay set up referred published paper.[43] The mouse head was fixed with a fixator two weeks before the assay. The mice received daily adaptive training on the VR system one week before the assay. On the day of the assay, the mice were fixed on the VR system and given 5 min of adaptation. Unless stimulated, the screen in front of the mouse will always play the regular landscape video. The stimulation of the eagle lasts for three seconds. The eagle dives down from far to near and plays the voice of the eagle at the beginning of the stimulation. The criterion for the stimulation was that the mice moved for the first time after 5 min of adaptation. The assay ended 2 min after the stimulation. The pupil changes and mice's locomotion speed were recorded simultaneously. After the assay, mice were fed alone in a clean new cage for 90 min and then used for

1038 brain extraction to detect c-Fos expression.

1039

Cat_{video} stimulation assay

The assay was carried out in a 40*40 cm OFT box. The mouse head was fixed

with a fixator and trained to fit the treadmill two weeks before the assay. On the

day of the assay, the mice were anesthetized with isoflurane and fixed on the

treadmill. After the mouse wakes up, play the video of cat hunting on the iPad

Air (Apple. Cat# 3rd generation, USA) in front of the mouse. A camera directly

above was used to record the assay.

1047

Aggressive social stress assay

The setup of the home cage social assay referred to published papers.[47, 76-79]

Selected aggressive CD1 mice never exposed to *Grin2d$^{f/f}$* or *SMA-*

creER:Grin2d$^{f/f}$ mice were used for the assay. Before the assay, simultaneously

put a single *Grin2d$^{f/f}$* or *SMA-creER:Grin2d$^{f/f}$* mouse and a CD1 mouse into a

cage. The cage contains new padding and a metal lid on the top. One side of

the metal lid is used to place food and water bottles, so the height is lower than

the other. A camera was used to record two mice's behaviours for 10 min. After

the assay, the mice were separated again. *Grin2d$^{f/f}$* or *SMA-creER:Grin2d$^{f/f}$*

mice were fed alone in a clean new cage for 90 min and then used for brain

extraction to detect c-Fos expression.

1059

Serial block-face scanning electron microscopy image

The specimen was obtained in the same batch as the previously published

paper from our laboratory. The details in the *SBF-SEM imaging* part of the

methods in that paper. In brief, the ascending venule was identified using X-ray

microscopy (Zeiss, XRM, Xradia 520 Versa, DEU) at ×4 and 80 kV

magnification. The specimen adhered to an aluminium metal pin using

conductive epoxy (Chemtronics, CW2400, USA). Then, it was further precisely

trimmed into a block of 600*600*700 μm^3 using a glass knife and a right-angled

diamond knife (DiATOME, Trim 90, CHE) after coating with gold-palladium

(Leica, EM ACE200, DEU) and a final trim using a 3View ultramicrotome (Gatan,

USA). Image acquisition was performed using a BSE detector on a Carl Zeiss

Gemini 300 scanning electron microscope at 2 kV voltage and approximately

1.8*10−3 mBar in high-vacuum mode with focus charge compensation. The

specimens were sectioned at a 50–75 nm thickness and a cutting speed of 0.1

mm/s.

1075

1076 **Elevated plus-maze test**

1077 The setup of elevated plus-maze (EPM) referred to published papers.[70, 80] The

1078 EPM comprises two open arms (65 cm * 5 cm) and two close arms (65 cm * 5

1079 cm). Each mouse was placed in the centre, facing one of the two open arms,

1080 and allowed to explore for 5 min. Time spent in open arms was recorded.

1081

1082 **Force Swimming Test**

1083 The Force Swimming Test (FST) referred to a published paper.[81] In brief, mice

1084 were put into the glass water tank with 25 °C clear water. Measuring the

1085 swimming time of mice in 5 minutes and then putting the mouse back in the

1086 home cage.

1087

Statistical analysis software and code

All the image processing and analysis of fluorescence intensity used Zeiss Zen

Lite (v3.4) and ImageJ (v1.53u). All whisker stimulation original data were

measured by RWD LSIS-self software (v5.0). All the original behavioral data for

OFT/complex OFT/EPM were recorded using Noldus. Data recording and

analysis of Looming and VR$_{eagle}$ assay was performed using self-compiled

codes provided by Bayonne company. All statistical data were processed by

Numbers (v12.2). All figures were made by Prism (v9.4.1) and Adobe Illustrator

2024 (v28.3). CBF peak mean value, heat maps, mass spectrometry data

inclusion criteria, cross-correlation, and prediction index (based on the

Structural Similarity Index, SSIM) were analyzed by self-compiled MATLAB

codes. Besides, the KEGG pathways, and cluster analysis of SSIM were using

self-compiled R code. All available codes can be found at

https://github.com/JialabEleven/Functional_hyperemia_regulates_emotion.

1102

REFERENCES

1 Roelofs K, Bramson B, Toni I. A neurocognitive theory of flexible emotion

control: The role of the lateral frontal pole in emotion regulation. *Ann N Y Acad*

Sci 2023; **1525**:28-40.

2 Bortolon C, Chen S, Bonanno GA. Components of emotion regulation

flexibility and psychosis: The association between psychosis-proneness and

context sensitivity. *Br J Clin Psychol* 2023; **62**:82-95.

3 D'Agostino A, Covanti S, Rossi Monti M, Starcevic V. Reconsidering Emotion

Dysregulation. *Psychiatr Q* 2017; **88**:807-825.

4 Easdale-Cheele T, Parlatini V, Cortese S, Bellato A. A Narrative Review of the

Efficacy of Interventions for Emotional Dysregulation, and Underlying Bio-

Psycho-Social Factors. *Brain Sci* 2024; **14**.

5 De Prisco M, Oliva V, Fico G *et al.* Emotion dysregulation in bipolar disorder

compared to other mental illnesses: a systematic review and meta-analysis.

Psychol Med 2023; **53**:7484-7503.

6 Geng H, Xu P, Aleman A, Qin S, Luo YJ. Dynamic Organization of Large-scale

Functional Brain Networks Supports Interactions Between Emotion and

Executive Control. *Neurosci Bull* 2024; **40**:981-991.

7 Szaflarski JP, Allendorfer JB, Heyse H, Mendoza L, Szaflarski BA, Cohen N.

Functional MRI of facial emotion processing in left temporal lobe epilepsy.

Epilepsy Behav 2014; **32**:92-99.

Etkin A, Wager TD. Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 2007; **164**:1476-1488.

Mathias EJ, Kenny A, Plank MJ, David T. Integrated models of neurovascular coupling and BOLD signals: Responses for varying neural activations. *Neuroimage* 2018; **174**:69-86.

Iadecola C. The Neurovascular Unit Coming of Age: A Journey through Neurovascular Coupling in Health and Disease. *Neuron* 2017; **96**:17-42.

Pico-Perez M, Radua J, Steward T, Menchon JM, Soriano-Mas C. Emotion regulation in mood and anxiety disorders: A meta-analysis of fMRI cognitive reappraisal studies. *Prog Neuropsychopharmacol Biol Psychiatry* 2017; **79**:96-104.

Shen J, Wang D, Wang X *et al.* Neurovascular Coupling in the Dentate Gyrus Regulates Adult Hippocampal Neurogenesis. *Neuron* 2019; **103**:878-890 e873.

Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception II: Implications for major psychiatric disorders. *Biol Psychiatry* 2003; **54**:515-528.

Hale MW, Hay-Schmidt A, Mikkelsen JD, Poulsen B, Shekhar A, Lowry CA. Exposure to an open-field arena increases c-Fos expression in a distributed anxiety-related system projecting to the basolateral amygdaloid complex. *Neuroscience* 2008; **155**:659-672.

15 Kovacs LA, Schiessl JA, Nafz AE, Csernus V, Gaszner B. Both Basal and

Acute Restraint Stress-Induced c-Fos Expression Is Influenced by Age in the

Extended Amygdala and Brainstem Stress Centers in Male Rats. *Front Aging*

Neurosci 2018; **10**:248.

16 Kakol-Palm D, Brusberg M, Sand E *et al.* Role of tachykinin NK(1) and NK(2)

receptors in colonic sensitivity and stress-induced defecation in gerbils. *Eur J*

Pharmacol 2008; **582**:123-131.

17 Taguchi R, Shikata K, Furuya Y *et al.* Selective corticotropin-releasing factor

1 receptor antagonist E2508 reduces restraint stress-induced defecation and

visceral pain in rat models. *Psychoneuroendocrinology* 2017; **75**:110-115.

18 Monarca RI, Silva RFB, Gabriel SI, Cerveira AM, von Merten S. The

Presence of a Shelter in an Open Field Test Has Differential Effects on the

Behavior and Stress Response of Two Mouse Species. *J Exp Zool A Ecol Integr*

Physiol 2025; **343**:480-492.

19 Sturman O, Germain PL, Bohacek J. Exploratory rearing: a context- and

stress-sensitive behavior recorded in the open-field test. *Stress* 2018; **21**:443-

452.

20 Zhang D, Ruan J, Peng S *et al.* Synaptic-like transmission between neural

axons and arteriolar smooth muscle cells drives cerebral neurovascular

coupling. *Nat Neurosci* 2024; **27**:232-248.

21 Laurie DJ, Bartke I, Schoepfer R, Naujoks K, Seeburg PH. Regional,

developmental and interspecies expression of the four NMDAR2 subunits,

examined using monoclonal antibodies. *Brain Res Mol Brain Res* 1997; **51**:23-32.

Monyer H, Burnashev N, Laurie DJ, Sakmann B, Seeburg PH. Developmental and regional expression in the rat brain and functional properties of four NMDA receptors. *Neuron* 1994; **12**:529-540.

Li J, Zhang Y, Zhang D *et al.* Ca(2+) oscillation in vascular smooth muscle cells control myogenic spontaneous vasomotion and counteract post-ischemic no-reflow. *Commun Biol* 2024; **7**:332.

Thompson RA. Methods and measures in developmental emotions research: some assembly required. *J Exp Child Psychol* 2011; **110**:275-285.

Huber G, Ogrodnik M, Wenzel J *et al.* Telmisartan prevents high-fat diet-induced neurovascular impairments and reduces anxiety-like behavior. *J Cereb Blood Flow Metab* 2021; **41**:2356-2369.

Lecrux C, Toussay X, Kocharyan A *et al.* Pyramidal neurons are "neurogenic hubs" in the neurovascular coupling response to whisker stimulation. *J Neurosci* 2011; **31**:9836-9847.

Lacroix A, Toussay X, Anenberg E *et al.* COX-2-Derived Prostaglandin E2 Produced by Pyramidal Neurons Contributes to Neurovascular Coupling in the Rodent Cerebral Cortex. *J Neurosci* 2015; **35**:11791-11810.

Niwa K, Araki E, Morham SG, Ross ME, Iadecola C. Cyclooxygenase-2 contributes to functional hyperemia in whisker-barrel cortex. *J Neurosci* 2000; **20**:763-770.

29 Hill RA, Tong L, Yuan P, Murikinati S, Gupta S, Grutzendler J. Regional Blood

Flow in the Normal and Ischemic Brain Is Controlled by Arteriolar Smooth

Muscle Cell Contractility and Not by Capillary Pericytes. *Neuron* 2015; **87**:95-

110.

30 Heissler SM, Chinthalapudi K. Structural and functional mechanisms of actin

isoforms. *FEBS J* 2025; **292**:468-482.

31 Mayer A, Siegel NR, Schwartz AL, Ciechanover A. Degradation of proteins

with acetylated amino termini by the ubiquitin system. *Science* 1989; **244**:1480-

1483.

32 Ree R, Varland S, Arnesen T. Spotlight on protein N-terminal acetylation.

Exp Mol Med 2018; **50**:1-13.

33 Kashina AS. Regulation of actin isoforms in cellular and developmental

processes. *Semin Cell Dev Biol* 2020; **102**:113-121.

34 Li W, Prakash R, Chawla D *et al.* Early effects of high-fat diet on

neurovascular function and focal ischemic brain injury. *Am J Physiol Regul*

Integr Comp Physiol 2013; **304**:R1001-1008.

35 Yang S, Webb AJS. Reduced neurovascular coupling is associated with

increased cardiovascular risk without established cerebrovascular disease: A

cross-sectional analysis in UK biobank. *J Cereb Blood Flow Metab*

2024:271678x241302172.

36 Chang HC, Tai YT, Cherng YG *et al.* Resveratrol attenuates high-fat diet-

induced disruption of the blood-brain barrier and protects brain neurons from

apoptotic insults. *J Agric Food Chem* 2014; **62**:3466-3475.

Hauglund NL, Andersen M, Tokarska K *et al.* Norepinephrine-mediated slow vasomotion drives glymphatic clearance during sleep. *Cell* 2025; **188**:606-622 e617.

Foster DB, Shen LH, Kelly J, Thibault P, Van Eyk JE, Mak AS. Phosphorylation of caldesmon by p21-activated kinase. Implications for the Ca(2+) sensitivity of smooth muscle contraction. *J Biol Chem* 2000; **275**:1959-1965.

Yao YB, Xiao CF, Lu JG, Wang C. Caldesmon: Biochemical and Clinical Implications in Cancer. *Front Cell Dev Biol* 2021; **9**:634759.

Vanlandewijck M, He L, Mae MA *et al.* A molecular atlas of cell types and zonation in the brain vasculature. *Nature* 2018; **554**:475-480.

Liu X, Li C, Falck JR, Roman RJ, Harder DR, Koehler RC. Interaction of nitric oxide, 20-HETE, and EETs during functional hyperemia in whisker barrel cortex. *Am J Physiol Heart Circ Physiol* 2008; **295**:H619-631.

Renic M, Klaus JA, Omura T *et al.* Effect of 20-HETE inhibition on infarct volume and cerebral blood flow after transient middle cerebral artery occlusion. *J Cereb Blood Flow Metab* 2009; **29**:629-639.

Shang C, Chen Z, Liu A *et al.* Divergent midbrain circuits orchestrate escape and freezing responses to looming stimuli in mice. *Nat Commun* 2018; **9**:1232.

Shoji H, Miyakawa T. Effects of test experience, closed-arm wall color, and illumination level on behavior and plasma corticosterone response in an

elevated plus maze in male C57BL/6J mice: a challenge against conventional interpretation of the test. *Mol Brain* 2021; **14**:34.

Silva BA, Gross CT, Graff J. The neural circuits of innate fear: detection, integration, action, and memorization. *Learn Mem* 2016; **23**:544-555.

Wang L, Chen IZ, Lin D. Collateral pathways from the ventromedial hypothalamus mediate defensive behaviors. *Neuron* 2015; **85**:1344-1358.

Bigi S, Maestripieri D, Aloe L, Alleva E. NGF decreases isolation-induced aggressive behavior, while increasing adrenal volume, in adult male mice. *Physiol Behav* 1992; **51**:337-343.

Zhou H, Neudecker V, Perez-Zoghbi JF, Brambrink AM, Yang G. Age-dependent cerebral vasodilation induced by volatile anesthetics is mediated by NG2(+) vascular mural cells. *Commun Biol* 2024; **7**:1519.

Bowen RM, York NW, Padawer-Curry J, Bauer AQ, Lee JM, Nichols CG. Control of neurovascular coupling by ATP-sensitive potassium channels. *J Cereb Blood Flow Metab* 2025; **45**:1130-1143.

York NW, Parker H, Xie Z *et al.* Kir6.1- and SUR2-dependent KATP overactivity disrupts intestinal motility in murine models of Cantu syndrome. *JCI Insight* 2020; **5**.

Lee S, Kang BM, Shin MK *et al.* Chronic Stress Decreases Cerebrovascular Responses During Rat Hindlimb Electrical Stimulation. *Front Neurosci* 2015; **9**:462.

Chiba S, Numakawa T, Ninomiya M, Richards MC, Wakabayashi C, Kunugi

H. Chronic restraint stress causes anxiety- and depression-like behaviors, downregulates glucocorticoid receptor expression, and attenuates glutamate release induced by brain-derived neurotrophic factor in the prefrontal cortex. *Prog Neuropsychopharmacol Biol Psychiatry* 2012; **39**:112-119.

Dion-Albert L, Dudek KA, Russo SJ, Campbell M, Menard C. Neurovascular adaptations modulating cognition, mood, and stress responses. *Trends Neurosci* 2023; **46**:276-292.

Chen M, Ma S, Liu H *et al.* Brain region-specific action of ketamine as a rapid antidepressant. *Science* 2024; **385**:eado7010.

Lee S, Kang BM, Kim JH *et al.* Real-time in vivo two-photon imaging study reveals decreased cerebro-vascular volume and increased blood-brain barrier permeability in chronically stressed mice. *Sci Rep* 2018; **8**:13064.

Hsueh B, Chen R, Jo Y *et al.* Cardiogenic control of affective behavioural state. *Nature* 2023; **615**:292-299.

Roy AK, Shehzad Z, Margulies DS *et al.* Functional connectivity of the human amygdala using resting state fMRI. *Neuroimage* 2009; **45**:614-626.

Kim KJ, Ramiro Diaz J, Iddings JA, Filosa JA. Vasculo-Neuronal Coupling: Retrograde Vascular Communication to Brain Neurons. *J Neurosci* 2016; **36**:12624-12639.

Jammal Salameh L, Bitzenhofer SH, Hanganu-Opatz IL, Dutschmann M, Egger V. Blood pressure pulsations modulate central neuronal activity via mechanosensitive ion channels. *Science* 2024; **383**:eadk8511.

60 Moore CI, Cao R. The hemo-neural hypothesis: on the role of blood flow in

information processing. *J Neurophysiol* 2008; **99**:2035-2047.

61 Hou XH, Hyun M, Taranda J *et al.* Central Control Circuit for Context-

Dependent Micturition. *Cell* 2016; **167**:73-86.e12.

62 Tateno F, Sakakibara R, Kishi M *et al.* Brainstem Stroke and Increased Anal

Tone. *Low Urin Tract Symptoms* 2012; **4**:161-163.

63 Hassabis D, Kumaran D, Summerfield C, Botvinick M. Neuroscience-

Inspired Artificial Intelligence. *Neuron* 2017; **95**:245-258.

64 Park S, Whang M. Empathy in Human-Robot Interaction: Designing for

Social Robots. *Int J Environ Res Public Health* 2022; **19**.

65 Apfelbach R, Blanchard CD, Blanchard RJ, Hayes RA, McGregor IS. The

effects of predator odors in mammalian prey species: a review of field and

laboratory studies. *Neurosci Biobehav Rev* 2005; **29**:1123-1144.

66 Bateman AW, Vos M, Anholt BR. When to defend: antipredator defenses and

the predation sequence. *Am Nat* 2014; **183**:847-855.

67 Liu Y, Wu Y, Yang Q. The impact of stress on the risk decision-making

process. *Psychophysiology* 2024; **61**:e14595.

68 Seo JS, Park JY, Choi J *et al.* NADPH oxidase mediates depressive behavior

induced by chronic stress in mice. *J Neurosci* 2012; **32**:9690-9699.

69 Khalil V, Faress I, Mermet-Joret N, Kerwin P, Yonehara K, Nabavi S.

Subcortico-amygdala pathway processes innate and learned threats. *Elife* 2023;

12.

70 Yu XD, Zhu Y, Sun QX *et al.* Distinct serotonergic pathways to the amygdala

underlie separate behavioral features of anxiety. *Nat Neurosci* 2022; **25**:1651-

1663.

71 Li X, Zhou L, Zhang X *et al.* Proteins secreted by brain arteriolar smooth

muscle cells are instructive for neural development. *Mol Brain* 2022; **15**:97.

72 Jin Y, Shi P, Wang Y *et al.* Precise control of embolic stroke with magnetized

red blood cells in mice. *Commun Biol* 2022; **5**:136.

73 Chao TH, Zhang WT, Hsu LM, Cerri DH, Wang TW, Shih YI. Computing

hemodynamic response functions from concurrent spectral fiber-photometry

and fMRI data. *Neurophotonics* 2022; **9**:032205.

74 Oyarzabal EA, Hsu LM, Das M *et al.* Chemogenetic stimulation of tonic locus

coeruleus activity strengthens the default mode network. *Sci Adv* 2022;

8:eabm9898.

75 Hou W, Yao J, Liu J *et al.* USP14 inhibition promotes recovery by protecting

BBB integrity and attenuating neuroinflammation in MCAO mice. *CNS Neurosci*

Ther 2023; **29**:3612-3623.

76 Arakawa H, Arakawa K, Blanchard DC, Blanchard RJ. Scent marking

behavior in male C57BL/6J mice: sexual and developmental determination.

Behav Brain Res 2007; **182**:73-79.

77 Arakawa H, Arakawa K, Blanchard DC, Blanchard RJ. A new test paradigm

for social recognition evidenced by urinary scent marking behavior in C57BL/6J

mice. *Behav Brain Res* 2008; **190**:97-104.

78 Borelli KG, Blanchard DC, Javier LK, Defensor EB, Brandao ML, Blanchard

RJ. Neural correlates of scent marking behavior in C57BL/6J mice: detection

and recognition of a social stimulus. *Neuroscience* 2009; **162**:914-923.

79 McQuaid RJ, Audet MC, Anisman H. Environmental enrichment in male CD-

1 mice promotes aggressive behaviors and elevated corticosterone and brain

norepinephrine activity in response to a mild stressor. *Stress* 2012; **15**:354-360.

80 Liu J, Li D, Huang J *et al.* Glutamatergic Neurons in the Amygdala Are

Involved in Paclitaxel-Induced Pain and Anxiety. *Front Psychiatry* 2022;

13:869544.

81 Warden MR, Selimbeyoglu A, Mirzabekov JJ *et al.* A prefrontal cortex-

brainstem neuronal projection that controls response to behavioural challenge.

Nature 2012; **492**:428-432.

1333

ACKNOWLEDGMENTS

We thank Prof. Bo Li for editing the manuscript and providing suggestions. We

thank Prof. Yanmei Tao and Dr. Xu Hu for sharing *Nestin-cre* and *Emx1-cre*

mice. We thank Yi Zhu for sharing psychological knowledge to facilitate our

understanding of data. We thank Yuxiao Niu for the KEGG map drawing. We

thank Westlake University Laboratory Animal Resource Centre for guiding

behaviour assays. We thank the Microscopy Core Facility of Westlake

University for imaging guidance. We thank the Mass Spectrometry &

Metabolomics Core Facility (MSMCF) of Westlake University for mass data

collection. We thank the Flow Cytometry Core Facility (FCCF) for cell sorting assistance. This work was supported by Westlake Laboratory of Life Sciences and Biomedicine (Project No. 202309002), Westlake University startup funding, the Westlake Education Foundation, Zhejiang, Key R&D Program of Zhejiang (2024SSYS0031), and the National Natural Science Foundation of China (project No. 32170961).

AUTHOR CONTRIBUTIONS

J.-M.J. conceived the project, designed the experiments, and supervised this study. J.Y.R. designed the key experiments, did primary experiments, obtained and organized the majority of original data, and wrote the manuscript draft. X.H.Q. participated in behavioural assay design and operations, siRNA cisterna magna injection, gut health inspection assay, BBB integrity inspection, caldesmon expression inspection, and cluster analysis of similarity index. Y.Y.Z. wrote the main MATLAB code for prediction framework. H.Q.X. participated in the Looming assay and VR$_{eagle}$ assay, as well as in blood pressure measurement. D.D.Z. provided primer pairs of *Grin1*, raw data of electron microscope, and *SMA-creER:Grin1$^{f/f}$* 40 cm OFT data. W.T.W. contributed to the writing of MATLAB code with Y.Y.Z.. B.R.Z. participated in the genotyping of mice and cell sorting. D.P.L. helped the mouse brain section.

CONFLICT OF INTERESTS

The authors declare no competing interests.

 FIGURE LEGENDS

 Fig.1 The negative emotion index. a Schematic diagrams of the home cage,

 OFT, and ARS. **b-c** BLA neuronal activation levels (**b**) and defecation bouts (**c**)

 under paradigms shown in **a**. Each dot represents one mouse. Student's t-test

 (**b**) and Mann-Whitney test (**c**) were used. **** indicates P<0.0001. **d** Linear

 correlation analysis between **b** and **c**. The original data were from **b** and part of

 c. **e** Time spent by mice in the center area of the OFT's box. Each dot

 represents one mouse. Student's t-test. **f** Total running distance by mice in the

 OFT. Each dot represents one mouse. Student's t-test. **g** Heatmap of mice's

 zone preference and defecation locations in the 40 cm OFT. **h** Statistical graph

 of **g**, data obtained from sixteen mice. **i** Schematic diagram of BLA

 chemogenetic inhibition during ARS. **j** Statistical graph of **i**. Each dot represents

 one mouse. Student's t-test. **k** Mice defecation bouts during BLA chemogenetic

 inhibition. Each dot represents one mouse. Mann-Whitney test. **l** Defecation

 rate and latency in the OFT. Number of mice is indicated in the figure. Gehan-

 Breslow-Wilcoxon test. **m** Time spent by mice in the center area of the OFT's

 box. Each dot represents one mouse. Student's t-test. **n** Total running distance

 by mice in the OFT. Each dot represents one mouse. Student's t-test. **o**

 Defecation bouts in the OFT following injection of the control virus. Each dot

 represents one mouse. Mann-Whitney test. Data are presented as mean ± SEM.

 See also Supplementary information Fig. S1.

Fig.2 GluN2D in SMCs regulates NVC and modulates negative emotions.

a Schematic diagram of the NsMJ within the NVU. **b** Schematic (top) and statistical graph (bottom) of RT-PCR. Each dot represents one mouse. **c** Schematic of RNA$_{scope}$. The inserted histogram shows the fluorescence intensity of the region indicated by the white arrows. The yellow dashed line indicates the boundary of the arteriole SMCs. **d** Schematic (left) and statistical graph (right) of MCA IF staining. Each dot represents one MCA. Six mice per group. Student's t-test. **** indicates P<0.0001. **e** Schematic diagram of the mouse whisker stimulation experiment. **f** Functional hyperemia in the S1BF during the whisker stimulation. **g** Statistical graph of the functional hyperemia plateau phase in **f**. Each dot represents one mouse. Student's t-test. **** indicates P<0.0001. **h** Schematic of the OFT combined with real-time neuronal activity, and rCBV monitoring in the BLA along with the negative emotion index monitor of freely moving mice. **i-j** rCBV (**i**) and Ca^{2+} (**j**) change in the BLA during the OFT. 0–900 seconds represents the duration of the OFT. **k** Statistical graphs of **i** and **j**. Each dot represents one mouse. Student's t-test. **l** Defecation bouts in every five minutes in a 40 cm OFT. Each dot represents one mouse. Mann-Whitney test. **m** Defecation rate and latency in the OFT. The number of mice is indicated in the figure. Gehan-Breslow-Wilcoxon test. **n** 1st 5 min rearing bouts in a 40 cm OFT. Each dot represents one mouse. Mann-Whitney test. **o** Defecation bouts in mice during OFTs of varying sizes and ARS. Each dot represents one mouse. Mann-Whitney test. **** indicates P<0.0001. Data are

presented as mean±SEM. Data from **o** reuse the data of *Grin2d^{f/f}* mice from **Fig.**

1c for comparison. **l** and **o** use same raw data of defecation bouts of *Grin2d^{f/f}*

and *SMA-creER:Grin2d^{f/f}* mice. Data are presented as mean±SEM. See also

Supplementary information Figs. S2-10.

Fig.3 Multiple NVC dysfunction models lead to excessive negative

emotions. a Schematic diagram of the double cKO mice mating strategy. **b**

Functional hyperemia in the S1BF during the whisker stimulation. **c** Statistical

graph of the functional hyperemia plateau phase in **b**. Each dot represents one

mouse. Student's t-test. **** indicates P<0.0001. **d** Defecation bouts in the OFT

and ARS in mice. Each dot represents one mouse (the number in parentheses

also indicates the number of mice). Mann-Whitney test. **e** Defecation rate and

latency in the OFT. The number of mice is indicated in the figure. Gehan-

Breslow-Wilcoxon test. **f** Schematic diagram of IF staining in the S1BF (left).

The orange dashed line indicates the location of the magnified area in the upper

right corner of the image. The right panel is a magnified view of the left

schematic. NeuN⁻ cells refer to non-neuronal cells expressing COX-2 within the

white dashed line in the left magnified image. Each dot represents one mouse.

Student's t-test. **g** Functional hyperemia in the S1BF during the whisker

stimulation. **h** Statistical graph of the functional hyperemia plateau phase in **g**.

Each dot represents one mouse. Student's t-test. **i** Schematic (top) and

statistical graph (bottom) of α-SMA. Each dot represents one mouse. Student's

t-test. **j** Schematic diagram of the HFD-induced model. **k** Body weight changes

of mice during HFD modelling. Each group sixteen mice. Student's t-test. ****

indicates P<0.0001. **l** mice fat and lean changes during HFD modelling. Each

dot represents one mouse. Student's t-test. **** indicates P<0.0001. **m**

Functional hyperemia in the S1BF during the whisker stimulation. **n** Statistical

graph of the functional hyperemia plateau phase in **m**. Each dot represents one

mouse. Student's t-test. Data from **b-e** reuse the data of *Grin2d^{f/f}* and *SMA-*

creER:Grin2d^{f/f} mice from **Fig. 2f, g, l, m** for comparison. NFD: normal fat diet.

Data are presented as mean±SEM. See also Supplementary information Figs.

S7, 11 ,12.

1444

Fig.4 Optogenetic stimulation of BLA SMCs. a Schematic diagram of

optogenetic stimulation in a freely moving mouse within the home cage. **b**

Schematic diagram showing the implanted position of the fiber ceramic ferrule.

c Schematic diagram illustrating the rCBV inhibition in the BLA under different

optogenetic protocols. **d** Statistical graph of **c**. Each point represents the

average change data from one mouse during the stimulation period. Student's

t-test. **e** Schematic diagram of optogenetic stimulation in a freely moving mouse

undergoing the OFT. **f** Schematic diagram illustrating the rCBV inhibition in the

BLA during the OFT. **g** Statistical graph of **f**. Each point represents the average

change data from one mouse during the stimulation period. Student's t-test. **h**

Defecation bouts in freely moving mice during optogenetic stimulation in the

OFT. Each point represents one mouse. Mann-Whitney test. **i** Defecation rate

and latency in the OFT. The dashed line represents the median defecation

latency. Twelve Control mice, and ten ChR2-overexpressing mice. Gehan-

Breslow-Wilcoxon test. **j** Experimental design (top) of optogenetic stimulation

(15 minutes) in a freely moving mouse within the home cage, and

corresponding changes in BLA rCBF (bottom). **k** Statistical graph of **j**. Each

point represents the average change from one mouse during optogenetic

stimulation period. Student's t-test. **l** Defecation bouts of mice. Each point

represents one mouse. Mann-Whitney test. **m** Schematic diagram of IF staining

in the BLA following optogenetic stimulation in the home cage and OFT. Each

point represents one mouse. Student's t-test. **n** Statistical graph of **m**. Each

point represents one mouse. Student's t-test. Data are expressed as mean ±

SEM. See also Supplementary information Figs. S13, 14.

1469

Fig.5 Caldesmon in SMCs regulates NVC and negative emotions. a KEGG

analysis of proteomic data from GluN2D IP products in SMCs. **b** IP products

detection using GluN2D and caldesmon antibodies. **c** Schematic illustrating

whole brain and BLA-specific caldesmon expression manipulation. **d**

Schematic of caldesmon IF staining. **e** Statistical graph of **d**. Each dot

represents one mouse. Student's t-test. **** indicates P<0.0001. **f**

Representative image (upper) and quantitative analysis (lower) of *Cald*

expression in the S1BF following whole-brain *Cald siRNA* injection. Each dot

represents one mouse. Student's t-test. **g-h** Schematic of IF staining for

caldesmon in the BLA (**g**) and ventromedial hypothalamus (VMH) (**h**) following

BLA local *Cald* siRNA injection. **i** Statistical graph of **f-h**. Each dot represents

one mouse. Student's t-test. **j** Functional hyperemia in the S1BF during the

whisker stimulation. **k** Statistical graph of the functional hyperemia plateau

phase in **j**. Each dot represents one mouse. Student's t-test. **l** Schematic of IF

staining after OFT in mice injected with *Cald* siRNA into the BLA. **m** Statistical

graph of **l**. Each dot represents one mouse. Student's t-test. **n** Defecation bouts

in the OFT and ARS. Each dot represents one mouse. Mann–Whitney test. ****

indicates $P<0.0001$. **o** Defecation rate and latency in the OFT. Number of mice

is indicated in the figure. Gehan–Breslow–Wilcoxon test. Data from **n** and **o**

reuse *Grin2d^{f/f}* and *SMA-creER:Grin2d^{f/f}* mice from **Fig. 2m, o** as controls. Data

are presented as mean ± SEM. See also Supplementary information Figs. S15.

1491

Fig 6. Pharmacological enhanced NVC rescues consequences of NVC

impairment. a Schematic diagram of drug intraperitoneal injection and BLA

local injection. **b-e** Effects of Het0016 administered via intraperitoneal injection

(**b, c**) and BLA local injection (**d, e**) on the negative emotion index in the OFT.

Each dot represents one mouse. **b, d** use Mann-Whitney test. **c, e** use Gehan-

Breslow-Wilcoxon test. **** indicates $P<0.0001$. **f** Effects of Het0016 on 1st 5

min rearing bouts. Each dot represents one mouse. Mann-Whitney test. **g**

Defecation bouts of *Pdgfrβ-creER:Acta2^{f/f}* mice after BLA local Het0016

injection in the OFT. Mann-Whitney test. **k** Defecation rate and defecation

latency of *Pdgfrβ-creER:Acta2$^{f/f}$* mice in the OFT after BLA local Het0016

injection. The number of mice is indicated in the figure. Gehan-Breslow-

Wilcoxon test. **i** Effects of Het0016 on 1st 5 min rearing bouts. Each dot

represents one mouse. Mann-Whitney test. Data are presented as mean ± SEM.

See also supplementary information, Figs. S16, 17.

Fig 7. Topology-Based estimation framework. a-b SSIM-based unbiased

classification of stress levels for seven standard contexts in control (**a**) and NVC

deficiency mice (**b**). **c-h** The predicted position of every single treatment in the

unbiased classification system. Black font represents *Grin2d$^{f/f}$* mice. Pink font

represents *SMAcreER:Grin2d$^{f/f}$* mice. Red font represents *Cald*-KD mice (**e**)

and ChR2$^+$ mice (**f**). Pink arrow presents the location of *SMAcreER:Grin2d$^{f/f}$*

mice, *Cald*-KD mice, and ChR2$^+$ mice with unbiased classification in the

Topology-Based estimation framework. See also Supplementary information

Figs. S18, S19.

Fig. 8 NVC deficiency prompted an early relaxation of defense upon

predator-like threat. a-d Looming assay. Temporal profile of locomotion speed

dynamics of the two genotypic mice before (blue) and during (yellow)

stimulation in **a**. Heatmap that profiled the locomotion speed dynamics for 30

seconds in **b**. The yellow box marks the 3-sec looming duration. Cumulative

time used for relaxation from a freezing status following looming stimulation in

c. Rearing bouts of each group in **d**. Fifteen mice for each group. **e-g** VR_{eagle}

stimulation. Temporal changes of the normalized pupil size and locomotion

speed during and after VR_{eagle} stimulation in **e**. Eagle call only present during

the first 1.5s (blue) and eagle started to dive at 1.5 second (yellow).

Representative images of VR_{eagle} video and mouse eye with pupil clear

visualized. The statistical data of the defensive offset phase for **e** in **f**. The

statistical data of pupil size full-recovery latency and looming locomotion full-

recovery latency in **g**. For VR_{eagle}, seven mice for each group; for Looming,

fifteen mice for each group. **h-l** Cat_{video} assay. The locomotion pattern during

the Cat_{video} assay in **h**. Black bar marks retreat behavior; magenta lines indicate

tail rattling. Two vertical dash lines highlight the time points when the cat

appeared and pounced. Mouse numbering is on the left. Quantification of

retreat behaviors of $Grin2d^{f/f}$ mice before and after cat appearance in **i**. Retreat

rate of two groups upon in Cat_{video} stimulation in **j**. Retreat rate referred to the

fraction of mice that exhibited at least one retreat behavior of total mice.

Cumulative retreat time curve in **k**. Quantification of tail rattling times in **l**. Twelve

mice for each group. Student's t-test. Data are presented as mean ± SEM. See

also supplementary information, Fig. S20.

Fig 9. NVC-deficient mice show hypo-responses to aggressive social stress. a Diagram of a home cage from a side view. **b** Total fighting time. each dot indicates one mouse. **c-d** Predicted positions of the two genotypic mice experience aggressive social stress. **e** Total time of staying with CD1 mouse in the small space. **f-g** Flee rate of *Grin2d^{f/f}* and *SMA-creER:Grin2d^{f/f}* mice (counted as per mouse) after fighting in **f**, and the normalized flee rate (counted as per fight event) after each fighting in **g**. Numbers in the graph indicates mouse number (**f**) and events (**g**). **h** Sniffing duration. Eight mice for *Grin2d^{f/f}* and ten mice for *SMA-creER:Grin2d^{f/f}*. Student's t-test. In **c-d**, black font represents *Grin2d^{f/f}* mice in **Fig. 7**. Blue (**c**) and pink (**d**) font represent *Grin2d^{f/f}* mice (**c**) and *SMAcreER:Grin2d^{f/f}* mice (**d**). Blue (**c**) and pink (**d**) arrow present the location of *Grin2d^{f/f}* mice (**c**) and *SMAcreER:Grin2d^{f/f}* mice (**d**) with unbiased classification in the Topology-Based estimation framework. Data are presented as mean ± SEM.

Fig. 10 Enhanced NVC in transgenic mouse model prevents strong CRS-induced NVC dysfunction and resistance emotional distress. a Schematic of RNA_{scope} of *Glrb* mRNA expression in venous SMCs. **b** Statistical graph of **a**. Each dot represents a vessel. Four mice per group. Mann-Whitney test. **** indicates $P < 0.0001$. **c** GlyRβ expression in venous SMCs. **d** Statistical graph of **c** (comparison between arterioles and venules in control mice). Each dot represents a vessel. Four mice per group. Mann-Whitney test. **e** Statistical

graph of **c** (comparison between venules of control and cKO mice). Each dot

represents a vessel. Four mice per group. Student's t-test. **f** Schematic (top)

and quantification (bottom) of *Glrb* expression in Pdgfrβ⁺ cells from the cortex.

Each dot represents one mouse. Student's t-test. **g** Co-localization of GlyRβ

with Gephyrin in HU-vSMC. The upper right panel shows the co-localization

rate. **h** Co-localization of GlyRβ protein with glycinergic axon terminals. The

orange dashed box indicates the zoom-in area in the upper right. Lower right is

histogram displays the fluorescence intensity across the region indicated by the

white arrow above. **i** Sequential electron microscopy sections of a ascending

venule in mouse cerebral cortex. White arrows indicate structural contacts

between axonal terminal and venous SMCs. **j** Functional hyperemia in the

S1BF during the whisker stimulation. **k** Statistical graph of the functional

hyperemia plateau phase in **j**. Each dot represents one mouse. Student's t-test.

l NVC-neutrality mice mating strategy. **m** Defecation bouts during the OFT and

ARS. Each dot represents one mouse. Mann-Whitney test. **** indicates

P<0.0001. **n** Defecation rate and defecation latency of mice in the OFT. The

number of mice is indicated in the figure. Gehan-Breslow-Wilcoxon test. **o**

Schematic diagram of the CRS model. **p** Fecal appearance of mice following

CRS. BS: Bristol score. **q** Bristol score and defecation bouts during CRS

modelling. Fiftheen mice per group. Mann-Whitney test. **** indicates P<0.0001.

r Immobility time of mice in the FST. Each dot represents one mouse. Student's

t-test. **** indicates P<0.0001. Data from **j, k, m, n** reuse the data of *Pdgfrβ-*

1586   *creER:Grin2d^{f/f}* mice from **Fig. 3b-e** for comparison. Data are presented as

1587   mean ± SEM. See also supplementary information, Figs. S21-27.

1588

SUPPLEMENTARY INFOEMATION

Supplementary figure legends

Fig. S1 Schematic illustration of BLA IF staining in conditions shown in Fig. 1a, b.

Fig. S2 Supplementary data for *Grin2d-cKO*. a Schematic of RNA$_{scope}$ of *Grin1* and *Grin2d* expression in the cerebellum. **b** Statistical graph of **a**. Each dot represents one mouse. **c-d** Schematic (top) and statistical graph (bottom) of GluN2D experiments in cerebral SMCs of mouse brain parenchyma (**c**) and whole cortex lysis (**d**). Each point represents one mouse. Student's t-test. **e** Schematic illustration (top) and quantitative analysis (bottom) of GluN2D. Each dot represents one mouse. Student's t-test. **** indicates P<0.0001. Data are presented as mean ± SEM.

Fig. S3 Vascular topology in the BLA of the mouse. a The schematic ventral view illustration of vascular topology is adapted from Figure 14.1 on page 460 of book *The Mouse Nervous System.* Translucent gray area indicates BLA. **b** The coronal view of the BLA. Green, yellow, and red lines indicate arterial distribution in the BAa, and BAv, and LA part of the BLA, respectively.

Fig. S4 Validation for detecting plasma rhodamine fluorescence intensity.

a Real-time fluorescence intensity measurement following retro-orbital

intravenous injection of various concentrations of rhodamine. **b** Linear

correlation statistical graph of **a**. Each point represents the average

fluorescence intensity from 120 to 150 seconds post-injection for a specific

rhodamine concentration. **c** Real-time fluorescence intensity measurements

following retro-orbital intravenous injection of 5 mg/mL rhodamine. The

experimental period ranged from 900 to 3600 seconds post-injection.

1617

1618 **Fig. S5 Neuronal activation in the BLA of *Grin2d-cKO* mice. a** Schematic

1619 illustration of IF staining in the BLA after the OFT. **b** Statistical graph of **a**. Each

1620 dot represents one mouse. Control mice data in **b** reuse data from **Fig. 1b** for

1621 comparison. Student's t-test. Data are presented as mean ± SEM.

1622

Fig. S6 Supplementary data on baseline rCBF and OFT in *Grin2d-cKO*

mice. a rCBF in the BLA of mice before entering the OFT box. Each dot

represents one mouse. Student's t-test. **b** Total running distance by mice during

the OFT. Each dot represents data from one mouse. Student's t-test. **c** Time

spent by mice in the central area of the OFT box. Each dot represents data from

one mouse. Student's t-test. Data are presented as mean ± SEM.

1629

Fig. S7 Verification of mouse metabolic status, gut health, and BBB integrity. a-g Food consumption, water intake, number of fecal pellets, daily fecal weight, weight per pellet, urine output, body weight changes, and blood pressure measured in the home cage. Each data point in figures **a-f** represents the average value for four mice within a cage. Each data point in figures **g-h** represents an individual mouse. Mann-Whitney test was applied for nonparametric data analysis, while Student's t-test was used otherwise. **i** Schematic illustration of mouse intestinal length, appearance, and HE staining under microscopy. **j-n** Statistical graphs of **i**. Shown are the body weight (**j**), intestinal length (**k**), intestinal length to body weight ratio (**l**), and total defecation count over 24 hours for 8-week-old experimental mice (**m**). Each data point represents one mouse. Student's t-test. **o** Appearance examination of the brain following retro-orbital intravenous injection of Evans Blue. **p** Statistical graph of **o**. Each data point represents one mouse. Student's t-test. Data are presented as mean ± SEM.

Fig. S8 SMC's marker protein validation in *Grin2d-cKO* mice. a Schematic illustration (top) and quantitative analysis (bottom) of α-SMA. Each dot represents one mouse. Student's t-test. **b** Statistical graph of Smoothelin IF staining from **Fig. 2d**. Each dot represents one MCA. Four mice per group. Student's t-test. Data are presented as mean ± SEM.

1651

Fig. S9 Validation of GluN2D expression levels in peripheral organs. a Low-magnification schematic diagrams of tissue sections from the heart, liver, spleen, lung, kidney, small intestine, and bladder of *SMA-creER:Ai14* mice (top). The yellow dashed box indicates the area enlarged (bottom). White arrows indicate the regions of interest for fluorescence intensity analysis in **b**. **b** Fluorescence intensity distribution in the regions indicated by the white arrows in **a**. **c** Statistical analysis of GluN2D IF staining intensity in GluN2D-positive cells and SMCs from tissue sections of **a**. Each dot represents one mouse. Student's t-test. **** indicates $P<0.0001$. Data are presented as mean ± SEM.

1661

Fig. S10 The effect of tamoxifen on NVC. a Functional hyperemia in the S1BF during the whisker stimulation. **b** Statistical graph of the functional hyperemia plateau phase in **a**. Each data point represents one mouse. Student's t-test. Data are presented as mean ± SEM.

1666

Fig. S11 IF staining of penetrating arteries in the mouse cerebral cortex. CD31 is the marker of endothelial cells.

1669

Fig. S12 Supplementary figure for HFD modelling. a Statistical graph of daily

1671 food consumption per cage. Each point represents the average of five mice per

1672 cage. Student's t-test. **b** Appearance examination of the brain following retro-

1673 orbital intravenous injection of Evans Blue (left) and statistical graph (right) of

1674 left. Each point represents one mouse. Student's t-test. Data are presented as

1675 mean ± SEM.

1676

1677 **Fig. S13 BLA blood spectral energy analysis. a** Representative image of

1678 mouse BLA blood spectral energy. **b** Statistical graph of **a**. Each data point

1679 represents one mouse. Student's t-test. Data are presented as mean ± SEM.

1680

Fig. S14 Optogenetic stimulation supplementary data. a Total running

distance by mice in the OFT. Each dot represents data from one mouse.

Student's t-test. **b** Time spent by mice in the center area of the OFT's box. **c**

Schematic diagram of IF staining for the BLA after optogenetic stimulation in

the home cage. Stroke area Hif-1α from stroke model mice served as a positive

control. **d** Statistical graph of **c**. Each data point represents one mouse.

Student's t-test. Data are presented as mean ± SEM.

1688

1689 **Fig. S15 S1BF *Cald* KD does not impact emotional responses. a** Schematic

1690 illustration showing the injection site of *Cald* siRNA in the S1BF and its spatial

1691 relationship with the BLA (orange dashed line). **b-c** Schematic diagram of IF

staining in the S1BF (**b**) BLA (**c**) after *Cald* siRNA injection into the S1BF. **d** Statistical analysis of **b-c**. Each dot represents one mouse. Student's t-test. **e** The defecation bouts in the OFT. Each dot represents one mouse. Mann-Whitney test. **f** Defecation rate and latency in the OFT. The number of mice is indicated in the figure. Gehan-Breslow-Wilcoxon test. Data are presented as mean ± SEM.

Fig. S16 Het0016 concentration test. a Effects of different concentrations of Het0016 on the defecation bouts in ARS. Each dot represents one mouse. Mann-Whitney test. **b** Functional hyperemia in the S1BF during the whisker stimulation. **c** Statistical graph of the functional hyperemia plateau phase in **b**. Each dot represents one mouse. Paired Student's t-test. Data are presented as mean±SEM.

Fig. S17 The effects of Het0016 in HFD-induced obese mice. a The defecation bouts in the OFT in HFD-induced obese mice. Each dot represents one mouse. Mann-Whitney test. **b** Defecation rate and latency in the OFT in HFD-induced obese mice. The number of mice is indicated in the figure. Gehan-Breslow-Wilcoxon test. Data are presented as mean±SEM.

Fig. S18 Schematic diagrams of behavioural apparatus and setups. a

1713 Diagram of the complex OFT chamber. **b** Diagram of the Looming assay setup.

1714 **c** Diagram of the VR$_{eagle}$ assay. **d** Diagram of the Cat$_{video}$ assay.

1715

Fig. S19 Neuronal activity changes and defecation bouts of mice in

different contexts. a Schematic diagram of neuronal activity changes and c-

Fos activation topography in the BLA of mice across different behavioral

paradigms. Each dot represents one mouse. Student's t-test. Data are

presented as mean ± SEM. **b** Normalized defecation bouts per 15 minutes in

different contexts. Each dot represents one mouse. Mann-Whitney test. Social:

aggressive social stress assay. **c** P_value of **b**. Schematic topological diagram

from **a** reuse the diagram from **Fig.7a, b**. Data are presented as mean ± SEM.

1724

Fig. S20 Supplementary data from the Looming and VR$_{eagle}$ assays. a

Statistical analysis of mice's locomotion speed before stimulation in the

Looming assay. Each dot represents one mouse. Student t-test. **b** Statistical

analysis of mice's locomotion speed before stimulation in the VR$_{eagle}$ assay.

Each dot represents one mouse. Student t-test. **c** Statistical analysis of mice's

pupil size before stimulation in the VR$_{eagle}$. Each dot represents one mouse.

Student t-test. **d** Statistical analysis of mice's pupil size at the last moment of

stimulation in the VR$_{eagle}$. Each dot represents one mouse. Student t-test. Data

are presented as mean ± SEM.

1734

1735 **Fig. S21 The expression level of *Glrb* in the transcriptome database.** vSMC

1736 refers to venous SMCs, aaSMC refers to arteriolar SMCs, and aSMC refers to

1737 arterial SMCs. Data are presented as mean ± SEM.

1738

Fig. S22 Functional hyperemia and behavioral supplementary data in

SMC-specific *Glrb-cKO* mice. a Functional hyperemia in the S1BF during the

whisker stimulation. **b** Statistical graph of the functional hyperemia plateau

phase in **a**. Each data point represents one mouse. Student's t-test. **c**

Defecation bouts during ARS. Each data point represents one mouse. Mann-

Whitney test. Data are presented as mean ± SEM.

1745

Fig. S23 Pinacidil inhibits NVC enhancement. a Schematic illustration of

intraperitoneal and BLA local injection of pinacidil. **b** Functional hyperemia in

the S1BF during the whisker stimulation. **c** Statistical graph of the functional

hyperemia plateau phase in **b**. Each dot represents one mouse. Paired

Student's t-test. **d** Absolute rCBF in the S1BF of mice before the whisker

stimulation. Each dot represents one mouse. Paired Student's t-test. **e-f**

Defecation bouts of mice during ARS after intraperitoneal injection (**e**) and BLA

local injection (**f**) of pinacidil. Each dot represents one mouse. Mann–Whitney

test. Data are presented as mean ± SEM.

1755

Fig. S24 Statistics of the time spent in the open arms of the EMP (left) and fecal pellet quality (right) in mice after CRS modelling. Each dot represents one mouse. Student's t-test was used for the left panel statistics, and Mann-Whitney test was used for the right panel statistics. BS: Bristol score. Data are presented as mean ± SEM.

1761

Fig. S25 Defecation bouts of mice in ARS. Each dot represents one mouse. Mann-Whitney test. **** indicates $P < 0.0001$. *$Glrb^{f/f}$* and *$Pdgfr\beta$-creER:$Glrb^{f/f}$* data are reused from **Fig. 10m** as controls. Data are presented as mean ± SEM.

1765

Fig. S26 Locomotion, anxiety-like, and weight of NVC-enhancement mice. a Total running distance by mice in the OFT. Each dot represents one mouse. Student's t-test. **b** Time spent by mice in the center area of the OFT's box. Each dot represents one mouse. Student's t-test. **c** Body weight changes of mice during the CRS modelling. Fifteen mice per group. Student's t-test. Data are presented as mean ± SEM.

1772

Fig. S27 BBB integrity following the CRS modelling. a Schematic illustration of IF staining in the S1BF and the BLA. **b** Statistical graph of **a**. Each dot

1775 represents one mouse. Student's t-test. Data are presented as mean ± SEM.

1776

1777 **Fig. S28 Schematic illustration of the BLA-Bar axis.** Dotted arrows and grey

1778 font represent inferred neuronal circuits or functions.

1779

1780 **Supplementary tables**

1781 **Supplementary Table S1. P-value of Defecation rate-latency curve**

1782 Sub table 1 for **Fig. 2m**

P_value	1	2	3	4	5	6
1	1	0.0031	0.0019	****	****	****
2		1	0.1934	0.3581	0.1384	0.1189
3			1	0.8685	0.3193	0.3458
4				1	0.1585	0.1083
5					1	0.9278
6						1

1. OFT_40 *Grin2d$^{f/f}$*

2. OFT_35 *Grin2d$^{f/f}$*

3. OFT_10 *Grin2d$^{f/f}$*

4. OFT_40 *SMA-creER:Grin2d$^{f/f}$*

5. OFT_40 *SMA-creER:Grin2d$^{f/f}$*

6. OFT_40 *SMA-creER:Grin2d^{f/f}*

**** indicates P<0.0001

1790

1791   Sub table 2 for **Fig. 3e**

P_value	1	2	3	4	5	6	7	8	9	10	11	12
1	1	****	0.0002	0.0823	0.0005	0.8480	****	0.4924	0.4906	****	0.5680	****
2		1	0.7333	0.0092	0.7209	0.0002	0.0786	0.0010	0.0043	0.4279	0.0010	0.2679
3			1	0.0047	0.7747	0.0001	0.2580	0.0014	0.0028	0.7549	0.0010	0.4218
4				1	0.0130	0.1038	****	0.2713	0.2961	0.0009	0.0548	0.0005
5					1	0.0004	0.2596	0.0042	0.0078	0.9480	0.0022	0.5377
6						1	****	0.5712	0.5970	****	0.4793	****
7							1	****	****	0.2935	0.0003	0.8118
8								1	0.9720	0.0001	0.2299	0.0001
9									1	0.0008	0.2695	0.0005
10										1	0.0003	0.5583
11											1	0.0005
12												1

1. *Grin2d^{f/f}*

2. *SMA-creER:Grin2d^{f/f}*

3. *Pdgfrβ-creER:Grin2d^{f/f}*

4. *Grin1^{f/f}*

5. *SMA-creER:Grin1^{f/f}*

6. *Ptgs2^{f/f}*

7. *Emx1-cre:Ptgs2^{f/f}*

8. *SMA-creER:Ptgs2^{f/f}*

9. *Acta2^{f/f}*

10. *Pdgfrβ-creER:Acta2^{f/f}*

11. NFD

12. HFD

**** indicates P<0.0001

1805

1806 Sub table 3 for **Fig. 5p**

P_value	1	2	3	4	5	6	7
1	1	****	0.9518	****	****	0.0008	****
2		1	0.0001	0.6713	0.4083	0.7646	0.4077
3			1	****	****	0.0009	****
4				1	0.5745	0.8881	0.9999
5					1	0.99990	0.6926
6						1	0.6353
7							1

1. *Grin2d^{f/f}*

2. *SMA-creER:Grin2d^{f/f}*

3. Scramble

4. *Cald* KD (Brain)

5. *Cald* KD (BLA)

6. Scramble+ *SMA-creER:Grin2d$^{f/f}$*

7. *Cald* KD (BLA) + *SMA-creER:Grin2d$^{f/f}$*

**** indicates P<0.0001

1815

1816　　Sub table 4 for **Fig. 6c**

P_value	1	2	3	4
1	1	0.1650	****	0.6470
2		1	****	0.2934
3			1	****
4				1

1817　　1. *Grin2d$^{f/f}$* +Saline

1818　　2. *Grin2d$^{f/f}$* +Het0016

1819　　3. *SMA-creER:Grin2d$^{f/f}$*+Saline

1820　　4. *SMA-creER:Grin2d$^{f/f}$*+ Het0016

1821　　**** indicates P<0.0001

1822

1823　　Sub table 5 for **Fig. 6e**

P_value	1	2	3	4
1	1	0.0679	0.0003	0.2123
2		1	****	0.3173
3			1	****
4				1

1824   5. *Grin2d^{f/f}* +Saline

1825   6. *Grin2d^{f/f}* +Het0016

1826   7. *SMA-creER:Grin2d^{f/f}*+Saline

1827   8. *SMA-creER:Grin2d^{f/f}*+ Het0016

1828   **** indicates P<0.0001

1829

1830

1831   Sub table 6 for **Fig. 10n**

P_value	1	2	3	4
1	1	0.0012	0.6285	0.4703
2		1	0.0035	0.0244
3			1	0.2637
4				1

1832   1. Floxp

1833   2. *Pdgfrß-creER:Grin2d^{f/f}*

1834   3. *Pdgfrß-creER:Glrb^{f/f}*

1835   4. *Pdgfrβ-creER:Grin2d^{f/f}:Glrb^{f/f}*

1836   **** indicates $P<0.0001$

1837

1838 **Supplementary Table S2. SSIM for Fig.7**

SSIM	HC	OFT_40	complex OFT	ARS	Cat$_{video}$	Looming	VR$_{eagle}$
HC	1	0.0359	0	0	0.0019	0.0015	0.0205
OFT_40		1	0.0038	0.0011	0.2150	0.1550	0.79800
complex OFT			1	0.2190	0.1050	0.0461	0.0002
ARS				1	0.0183	0.0084	0
Cat$_{video}$					1	0.9440	0.0778
Looming						1	0.0542
VR$_{eagle}$							1

1839

1840 **Supplementary Table S3. Mice information**

Abbreviation	Full name	Source
Ai32	B6;129S-Gt(ROSA)26Sortm32(CAG-COP4*H134R/EYFP)Hze/J	Hailan Hu, Zhejiang University

Nestin-cre	B6.Cg-Tg(Nes-cre)1Kln/J	Yanmei Tao and Xu Hu, Hangzhou Normal University/ Southeast University
Emx1-cre	B6.129S2-Emx1tm1(cre)Krj/J	Yanmei Tao and Xu Hu, Hangzhou Normal University/ Southeast University
SMA-creER	Tg(Acta2-cre/ERT2)12Pcn	Bo Zhou, ShanghaiInstitutes for Biological Sciences, Chinese Academy of Sciences
Pdgfrβ-creER	Tg(Pdgfrb-cre/ERT2)6096Rha	Xiang Yu, Peking University
Ai47	Gt(ROSA)26Sortm47(CAG-EGFP*)Hze	Zilong Qiu, Chinese Academy of Sciences Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences.

Ai14	B6.Cg-Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J	Woo-Ping Ge, Chinese Institute of Brain Research (CIBR)
Grin2d	B6/JGpt-Grin2dem1Cflox/Gpt	GemPharmatech
Grin1	Grin1tm2Stl	Yangang Sun, Chinese Academy of Sciences Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences
Ptgs2	C57BL/6JCya-Ptgs2em1flox/Cya	Cyagen
Acta2	C57BL/6JGpt-Acta2em1Cflox/Gpt	GemPharmatech
Glrb	C57BL/6JGpt-Glrbem1Cflox/Gpt	GemPharmatech

1841

1842 **Supplementary Table S4. Antibodies information**

Name	Host	Cat#	Dilution	Company

Anti-NMDAR2D Antibody	Mouse	MAB5578	1:400 (IF) 1:2,500 (WB) 1:1,000 (IP)	Merck
Anti-Smoothelin antibody	Rabbit	ab219652	1:400 (IF)	abcam
α-Smooth Muscle Actin (ACTA2) Rabbit mAb	Rabbit	A17910	1:2,500 (WB)	ABclonal
ACE2 Recombinant Rabbit Monoclonal Antibody (SN0754)	Rabbit	MA5-32307	1:400 (IF)	Invitrogen
Gephyrin Polyclonal Antibody	Rabbit	PA5-29036	1:400 (IF)	Invitrogen
BD Pharmingen™ Purified Rat Anti-Mouse CD31	Rat	557355	1:200 (IF)	BD
Anti-NeuN purified Antibody	Guinea Pig	ABN90P	1:800 (IF)	Merck
COX2/PTGS2 Rabbit pAb	Rabbit	A1253	1:400 (IF)	ABclonal
c-Fos antibody	Rabbit	226 003	1:1,000 (IF)	SYSY
Recombinant Alexa Fluor® 488 Anti-Caldesmon/CDM antibody	Rabbit	ab208116	1:400 (IF)	abcam

			1:2,500 (WB)	
iFluor™ 488 Conjugated Goat anti-mouse IgG polyclonal Antibody	Goat	HA1125	1:400 (IF)	HUABIO
Mouse Aminopeptidase N/CD13 Antibody	Goat	AF2335	1:200 (IF)	R&D
CD140b (PDGFRB) Monoclonal Antibody (APB5), APC, eBioscience™	Rat	17-1402-82	1:50 (FCM)	Invitrogen
Mice IgG	Mouse	A7028	1:1,000 (IP)	Beyotime
Donkey anti-Mouse IgG (H+L) ReadyProbes™ Secondary Antibody, Alexa Fluor™ 488	Donkey	R37114	1:1,000 (IF)	Invitrogen
Donkey anti-Rabbit IgG (H+L) ReadyProbes™ Secondary Antibody, Alexa Fluor™ 488	Donkey	R37118	1:1,000 (IF)	Invitrogen
Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 546	Donkey	A10040	1:1,000 (IF)	Invitrogen

Donkey anti-Rat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488	Donkey	A21208	1:1,000 (IF)	Invitrogen
Donkey anti-Rat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 555	Donkey	A78945	1:1,000 (IF)	Invitrogen
Donkey anti-Rat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 647	Donkey	A48272	1:1,000 (IF)	Invitrogen
Alexa Fluor® 647 AffiniPure™ Donkey Anti-Guinea Pig IgG (H+L)	Donkey	706-605-148	1:1,000 (IF)	Jackson ImmunoResearch
β-Tubulin Mouse mAb	Mouse	AC021	1:5,000 (WB)	ABclonal
Goat Anti-Mouse IgG, HRP Conjugated	Goat	CW0102	1:10,000 (WB)	CWBIO
Goat Anti-Rabbit IgG, HRP Conjugated	Goat	CW0103	1:10,000 (WB)	CWBIO
DDDDK-Tag Rabbit mAb	Rabbit	AE063	1:400 (IF)	ABclonal

| Hif-1α | | | Rabbit | ER1802-41 | 1:100 (IF) | HUABIO |

1843

Supplementary Table S5. Primer pairs information

Name		Sequence
Grin1	F	AAGAACCTGCAGCAGTACCA
	R	GCAGCAGGACTCATCAGTGT
Grin2a	F	TCAGGGACTTTGGAAAACGCT
	R	AAGTCATGCAGGGTTGGTTG
Grin2b	F	ACTAACTATCAATGAAGAACGGT
	R	CCAGACCCCAGAGTAACCA
Grin2c	F	TGTGGGCCTTCTTCGCTGTCATCT
	R	TGCCATTAGGTACCGTGCCAAAAC
Grin2d	F	TGGAGGAGTACGACTGGACAT
	R	CGCACTGACACTACGGAGC
Grin3a	F	CTGCACACCAGTCAGAGGTT
	R	TTCGTTGGTTGTCGTGACTC

Grin3b	F	TGGGGGTCTACCACTTGTCA
	R	TGGATCTTCTGGCTCGTGTG
Glrb	F	TTCAAAGGCATTCCTGTTGATGT
	R	TGAGCCTCTGAAGTCACTAGG
Cald	F	GCCGTTCAAGTGCTTCACTC
	R	TGTCAATCTTGGAGACTACTGCT
18s	F	CGGACACGGACAGGATTGACA
	R	CCAGACAAATCGCTCCACCAACTA
Cald (siRNA)	F	GUUUGAUCCGACCAUAACAGA
	R	UGUUAUGGUCGGAUCAAACUC

1845

1846 **Supplementary Video**

1847 **Supplementary Video S1. Topological distribution of arterioles in the BLA.**

1848 **Supplementary Video S2. VR$_{eagle}$ stimulation assay.**

1849 **Supplementary Video S3. Cat$_{video}$ stimulation assay.**

1850

Figure 1

a

Stress level
Low — High

HC OFT ARS

HC: Home cage
OFT: Open field test
ARS: Acute restraint stress

b

Routine | Stress-induced
c-Fos positive cells in BLA per mm²
****
****
0.0004
0.0017
HC 40 35 10 ARS
OFT

c

Routine | Stress-induced
Defecation bouts
****
0.0404
0.0404
0.0359
HC 40 35 10 ARS
OFT

d

R=0.8254
P<0.0001
c-Fos positive cells in BLA per mm²
Stress-induced defecation bouts

e

Time in the center (s)
0.0062
OFT_40 OFT_35

f

Running distance (cm)
0.0056
OFT_40 OFT_35

g

Short — Long ● Defecation

h

Rate (%)
Center / Peripheral
88.2% 78.5%
11.8% 21.5%
Defecation Time

i

EGFP-hM4Di
c-Fos
DAPI
Vehicle C21
800µm 200µm

j

c-Fos positive cells in BLA per mm²
0.0003
Vehicle C21

k

Defecation bouts
OFT ARS
0.0187 0.0126
V C V C
V: Vehicle
C: C21

l

— hM4Di-EGFP_Vehicle (N=10)
— hM4Di-EGFP_C21 (N=9)
—○— EGFP_Vehicle (N=7)
—○— EGFP_C21 (N=7)

Defecation rate (%)
ns
0.0208
Defecation latency (s)

m

hM4Di-EGFP
Time in the center (s)
ns
Vehicle C21

n

hM4Di-EGFP
Running distance (cm)
ns
Vehicle C21

o

EGFP
Defecation bouts
ns
Vehicle C21

a Neurovascular interface NsMJ / Neuron / Cation/anion Receptors / Bouton / Glu/Gly / SMC / Lumen / Neurovascular unit (NVU)

b 1: P0 cerebellum / 2: Parenchymal aSMCs sorted from adult brains / Ladder / Grin1 Grin2a Grin2b Grin2c Grin2d Grin3a Grin3b / NMDAR subunits / 18s / mRNA expression level (%) / P0 cerebellum / Parenchymal aSMCs / Grin1 Grin2a Grin2b Grin2c Grin2d Grin3a Grin3b

c RNAscope_SMA-creER:Ai14 / tdTomato / Grin2d / DAPI / Intensity (a.u) / Distance (μm) / 10 μm

d Grin2d^{f/f} / SMA-creER:Grin2d^{f/f} / Smoothelin / GluN2D / DAPI / 10 μm / GluN2D expression level in MCA / ****

e

f Flox control / SMA-creER:Grin1^{f/f} / SMA-creER:Grin2d^{f/f} / ΔCBF (%) / Whisker stimulation / Seconds

g ΔCBF (%) / **** / **** / ns

h Gcamp8f (Neuron) / Rhodamine (rCBV) / OFT free-moving mouse BLA recording

i CBV_Grin2d^{f/f} / CBV_SMA-creER:Grin2d^{f/f} / ΔCBV% / Seconds

j Ca^{2+}_Grin2d^{f/f} / Ca^{2+}_SMA-creER:Grin2d^{f/f} / ΔCa^{2+}% / Seconds

k Grin2d^{f/f} / SMA-creER:Grin2d^{f/f} / 1st 5-min AUC (a.u. ×10^4) / CBV / Ca^{2+} / 0.0170 / 0.0380

l Grin2d^{f/f} / SMA-creER:Grin2d^{f/f} / 11.81 folds / 0.0025 / 3.50 folds / 0.0075 / 3.00 folds / 0.0389 / Defecation bouts / 0-5min 5-10min 10-15min

m OFT_40 35 10 / Grin2d^{f/f}(N=16/19/14) / SMA-creER:Grin2d^{f/f}(N=16/19/14) / Defecation rate (%) / Defecation latency (s) / 0.0001

n Grin2d^{f/f} / SMA-creER:Grin2d^{f/f} / **** / Rearing bouts

o Grin2d^{f/f} / SMA-creER:Grin2d^{f/f} / ns / 0.0311 / 0.0404 / ns / ns / 0.0359 / Defecation bouts / HC OFT(40) OFT(35) OFT(10) ARS / **** / ns ns ns

a KEGG Pathway Enrichment_Top10

b

c Cisterna magna or local injection

Brain-specific/BLA local siRNA injection

d Caldesmon DAPI

e

f

g BLA injection

h VMH(BLA injectopn)

i

j

k

l

m

n

o

a *Pdgfrβ-creER:Ai14*
Glrb mRNA / tdTomato | Glrb mRNA

b Number of signal dots per cell
aSMCs
vSMCs
****

c *Pdgfrβ-creER:Ai14*
GlyRβ / tdTomato | GlyRβ
Pdgfrβ-creER:Ai14:Glrb^{f/f}
GlyRβ / tdTomato | GlyRβ

d GLRβ cluster number per 10 μm
Arterioles
Venules
0.0007

e GlyRRβ relative expression level
0.0045
● Glrb^{f/f} ● Pdgfrβ-creER:Glrb^{f/f}

f Glrb relative expression level
Glrb / 18s
Adult cortex
0.0001
● Adult cerebral cortex
● Glrb^{f/f} ● Pdgfrβ-creER:Glrb^{f/f}

g Gephyrin / GlyRβ / ACE2 / DAPI

h GlyT2-EGFP / GlyRβ / CD13 / DAPI
Intensity (a.u) vs Distance (μm)
CD13 / GlyRβ / GlyT2_EGFP

i 1, 2, 3, 4 — SMC, EC, Lumen

j △CBF (%) vs Seconds — Whisker stimulation
● Glrb^{f/f}
● Pdgfrβ-creER:Glrb^{f/f}
● Pdgfrβ-creER:Grin2d^{f/f}
● Pdgfrβ-creER:Grin2d^{f/f}:Glrb^{f/f}
● CRS_Glrb^{f/f}
● CRS_Pdgfrβ-creER:Glrb^{f/f}

k △CBF (%)
ns
**** 0.0172
0.0051
0.0156
ns

l *Pdgfrβ-creER:Grin2d^{f/f}* (Deficiency) X *Pdgfrβ-creER:Glrb^{f/f}* (Enhancement)
→ *Pdgfrβ-creER:Grin2d^{f/f}:Glrb^{f/f}* NVC-neutrality

m Defecation bouts — OFT | ARS
ns
**** 0.0364
ns
****
● Floxp (N=24)
● Pdgfrβ-creER:Grin2d^{f/f}(N=17)
● Pdgfrβ-creER:Glrb^{f/f}(N=15)
● Pdgfrβ-creER:Grin2d^{f/f}:Glrb^{f/f}(N=13)

n Defecation rate (%) vs Defecation latency (s)
0.0299
0.0013
ns

o Chronic restraint stress (CRS)model
Random restraint 2-3 hours/day X 28 days

p BS 3 / CRS - / Glrb^{f/f}
BS 2 / CRS + / Glrb^{f/f}
BS 3 / CRS + / Pdgfrβ-creER:Glrb^{f/f}

q Bristol Score / Defecation bouts (HC-like) vs Day
ns
ns
Bristol score: ****, 0.0001, ns
Defecation bouts: ****, ****, ns
● Naive_Glrb^{f/f}
● CRS_Glrb^{f/f}
● CRS_Pdgfrβ-creER:Glrb^{f/f}

r Immobility time in 300s (s)
● Naive_Glrb^{f/f} ● Naive_Pdgfrβ-creER:Glrb^{f/f}
● CRS_Glrb^{f/f} ● CRS_Pdgfrβ-creER:Glrb^{f/f}
0.0002
0.0289
ns
****