Macrophage epigenetic memories of early life injury drive neonatal nociceptive priming.

Adam J. Dourson[1], Adewale O. Fadaka[1], Anna M. Warshak[1], Aditi Paranjpe[2], Benjamin Weinhaus[3], Luis F. Queme[1], Megan C. Hofmann[1], Heather M. Evans[4], Omer A. Donmez[5], Carmy Forney[5], Matthew T. Weirauch[5,6,7], Leah C. Kottyan[5,6,8], Daniel Lucas[3,8], George S. Deepe, Jr.[9], Michael P. Jankowski[1,8,10]*

[1] Department of Anesthesia, Division of Pain Management, Cincinnati Children's Hospital Medical Center, [2] Biomedical Informatics, Cincinnati Children's Hospital Medical Center, [3] Division of Experimental Hematology and Cancer Biology, Cincinnati Children's Medical Center, [4] Division of Infectious Diseases, University of Cincinnati, [5] Center for Autoimmune Genomics and Etiology, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, United States of America. [6] Division of Allergy and Immunology, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, United States of America. [7] Divisions of Biomedical Informatics and Developmental Biology, Cincinnati Children's Hospital Medical Center. [8] Department of Pediatrics, University of Cincinnati, College of Medicine, Cincinnati, OH 45229. [9] Department of Medicine, Division of Infectious Diseases, University of Cincinnati College of Medicine, [10] Pediatric Pain Research Center, Cincinnati Children's Hospital Medical Center

*Corresponding author:

Michael P. Jankowski
3333 Burnet Avenue, MLC6016
Cincinnati, Ohio 45229-3026
Telephone: (513) 803-7966
E-mail: michael.jankowski@cchmc.org

Conflict of interest:

The authors declare no competing financial interests.

Abstract

The developing peripheral nervous and immune systems are functionally distinct from adults.

These systems are vulnerable to early life injury, which influences outcomes related to

nociception following subsequent injury later in life (i.e., "neonatal nociceptive priming"). The

underpinnings of this phenomenon are largely unknown, although previous work indicates that

macrophages are epigenetically trained by inflammation and injury. We found that macrophages

are both necessary and partially sufficient to drive neonatal nociceptive priming possibly due to

a long-lasting epigenetic remodeling. The p75 neurotrophic factor receptor (NTR) was an

important effector in regulating neonatal nociceptive priming through modulation of the

inflammatory profile of rodent and human macrophages. This "pain memory" was long lasting in

females and could be transferred to a naïve host to alter sex-specific pain-related behaviors.

This study reveals a novel mechanism by which acute, neonatal post-surgical pain drives a

peripheral immune-related predisposition to persistent pain following a subsequent injury.

Introduction

Peripheral injury activates several biological systems to evoke an appropriate motor and affective response to sensory stimuli. To generate and maintain nociceptive signals, there is remarkable crosstalk between cellular systems to create distinct molecular patterns that depend on the given aversive event[1,2]. Importantly, these systems undergo significant developmental changes especially in early life[3,4], a period in which nociceptors and immune cells display unique properties compared to their adult counterparts[5-10].

Critical periods are constrained time points of high plasticity in which typical development is persistently altered by external input. Clinical studies indicate that early life surgery or time spent in the neonatal intensive care unit (NICU) alters the development of the somatosensory system[11,12]. In these patients, injury in adulthood has a higher risk of complications including increased pain management medication use and longer hospital stays. Rodent drug studies revealed that the peripheral afferent input is necessary for this phenomenon termed "neonatal nociceptive priming"[13]. However, other peripheral and developmental factors that influence this experience are unknown.

Immune cells, particularly macrophages, modulate nociceptive responses. For example, patient recovery from surgery correlates with their pre-operative inflammatory profile[14-20]. In rodents, the depletion or activation/transfer of macrophages results in differential pain-like behaviors. These studies indicate a key role for neuroimmune signaling in the nociceptive response[5,20-23]. Evidence also suggests that macrophages "learn" in response to multiple exposures of the same or similar adverse agents, likely through epigenetic remodeling[24-26]. While these studies

indicate a cellular "memory" of these innate cells, it is unknown if there is a similar effect after

injury, or whether this "memory" may persistently affect behavior.

Aversive stimuli within the first week of life alters sensory neuron development by modifying key

signaling pathways, such as neurotrophic factors[6,7,27]. Similar effects may be present in immune

cells after an injury, however, much less is understood[28,29]. p75NTR (also known as nerve

growth factor receptor (NGFR)) modulates broad neurotrophic factor signaling pathways[30],

transcription, and the function of neurons[31] and immune cells[28]. Neurotrophins, therefore, have

critical effects on the development of the nervous and immune systems and could play a dual

role in both neonatal inflammation and pain.

Pediatric post-surgical pain is a major clinical health problem that is complicated by underlying

differences in patients, diagnostic uncertainty, and developmental concerns[32-35]. There is often a

choice between patients suffering the negative effects of pain (intensity or unpleasantness and

developmental consequences) or the side effects of current treatments (NSAIDs, steroids,

anticonvulsants, opioids, etc.[36,37]). These and other studies clearly demonstrate that improved

therapeutics designed for early life pain are necessary for precision care[38]. Here, we aimed to

investigate how early life injury alters the peripheral neuroimmune system to "prime" animals to

re-injury later in life. We determined the necessity and sufficiency of macrophages for neonatal

nociceptive priming, and we evaluated the epigenetic changes that occur in macrophages after

neonatal injury, which may underlie this phenomenon. We hypothesized that injury-induced

epigenetic changes to chromatin accessibility in developing macrophages contributes to

neonatal nociceptive priming.

Results

Macrophages are necessary for neonatal nociceptive priming.

Previous data indicate that macrophages are important for acute nociception in adults and

neonates[5,21,39]. To establish the role of neonatal macrophages in the induction of neonatal

nociceptive priming, we used macrophage fas-induced apoptosis (MaFIA) animals to ablate

macrophages early in life and tested animals before and after both neonatal and adolescent

incision (Fig. 1A). In this animal, *Csf1r*-driven expression of the FK506 binding protein 1A

induces cell apoptosis when dimerized, which is temporally controlled by the injection of a

designer chemical and inducer of dimerization, AP20187 (AP). Importantly, when injected,

Csf1r+ cells undergo apoptosis, but within days of stopping the injection, new monocytes

derived from the bone marrow begin to replenish the lost cells[40]. We found that systemic

injection of AP drastically reduced splenic macrophages but not microglia in the spinal cord

(Supp. Fig. 1A-B), as has been similarly noted in adults[41,42]. This knockout strategy also

prevented macrophage infiltration into injured muscle one day after a neonatal incision (Fig. 1B-

C). Both vehicle and AP treated groups displayed significant paw guarding one day after a

neonatal incision, with no statistical difference between groups (Fig. 1D, *p<0.001 vs. baseline

(BL), p=0.071 n.inc+AP vs. n.inc+vehicle). However, while vehicle treated animals displayed a

reduction in mechanical withdrawal thresholds to muscle squeezing one day after incision

(*p=0.014 vs. BL), AP treated animals were hyposensitive compared to baseline and controls

(Fig. 1E, *p=0.007 vs. BL, ^p=0.028 vs. n.inc+vehicle).

We then allowed different animals to age to P35 when sensory neurons and immunological

functions are more developed[6,7,43,44]. At this time point, we first performed evaluations to

determine any long-lasting effects of early life incision. Prior to a second incision at P35, we

evaluated muscle integrity and LysM+ cell presence (Supp. Fig. 1C). Evans Blue Dye (EBD),

when administered peripherally, is unable to penetrate into myofiber bundles unless they are

damaged or undergoing repair[45]. Two days after adolescent injury, we observed that about half

of the myofibers were positive for EBD. However, we found significantly less EBD staining within

myofibers in animals that received an early life injury or early life sham (Supp. Fig. 1D-E,

*$p<0.001$ vs. two days post incision). In addition, there were very few LysM+ cells still present in

the muscle of monocyte/macrophage reporter animals (LysM;tdTom) at P35 after neonatal

sham (n.sham) or incision surgery (n.inc) (Supp. Fig. 1F, $p=0.648$ vs. n.sham).

To determine the effect of early life injury in the presence or absence of macrophages on re-

injury pain-like behaviors, we evaluated MaFIA animals treated with AP or vehicle early in life

that received either a P7 sham or incision surgery. After aging and prior to a second incision, we

observed no differences between any groups for paw guarding or muscle squeezing. One day

after the P35 incision, we confirmed that even in neonatal AP treated animals, muscle *Csf1r+*,

GFP expressing macrophages are present in the injured muscle (see Fig. 1B, C). At this same

time point, we found expected acute pain-like behaviors from all groups. However, n.sham

animals from both treatments returned to baseline levels of paw guarding after day three (Fig.

1F, *$p<0.05$ vs. BL for all groups) and muscle withdrawal thresholds after day seven (Fig. 1G,

*$p<0.001$ vs. BL for all groups). We detected no effect of neonatal AP alone on nociceptive

development in n.sham+AP animals (n.sham+vehicle vs. n.sham+AP). Unlike n.sham animals,

n.inc+vehicle animals displayed long-lasting pain-like behaviors after a second incision through

day 14 for guarding (Fig. 1F, *$p<0.001$ vs. BL, ^$p<0.05$ vs. n.sham+vehicle, #$p<0.001$ vs.

n.inc.+AP) and day 21 for muscle squeezing (Fig. 1G, *$p<0.001$ vs. BL, ^$p<0.005$ vs.

n.sham+vehicle, #$p<0.005$ vs. n.inc+AP). Early life ablation of macrophages rescued this

persistent pain-like phenotype with n.inc+AP animals displaying guarding and mechanical

hypersensitivity only through day seven (Fig. 1F-G, guarding *$p<0.05$ vs. BL; squeezing

*$p<0.001$ vs. BL). Animals of this group were not significantly different at any time point

compared to controls.

To evaluate if the ablation of *Csf1r*+ macrophages early in life alters the sensitization of sensory

neurons following dual incision, we performed intracellular recordings of individual sensory

neurons in an *ex vivo* intact muscle/tibial-nerve/DRG/SC preparation (Fig. 2A). We evaluated

multiple peripheral response properties seven days after the second incision in vehicle and AP

treated mice where we saw a robust block of both spontaneous guarding and evoked muscle

squeezing behaviors (see Figs. 1F-G). We found no difference in the conduction velocity (CV) of

thinly myelinated group III afferents in animals treated with AP versus vehicle seven days after

the second incision (Fig. 2B, p=0.987). However, unmyelinated group IV afferents showed a

significantly slower CV in AP treated animals versus vehicle treated animals (Fig. 2C,

*p=0.048). After mechanical stimulation of receptive fields, we observed no change in average

mechanical force necessary to evoke an action potential (Fig. 2D, p=0.430).

Overall, we observed no difference in the firing dynamics on all neurons collectively (Supp. Fig.

2), but upon the analysis of subtypes, we found that chemosensitive and mechanically sensitive

fibers (without cold responsiveness[46]), had an overall greater mechanical firing rate that was

reduced following incubation with noxious metabolites (Fig. 2E, *p=0.031). Vehicle controls did

not show a pre/post-metabolite difference. We further uncovered that control nociceptors

displayed a direct increase in firing rate as we applied greater forces to the receptive field as

has been previously reported[47]. However, in animals with an early in life restricted knockout of

macrophages, we found no association between firing rate and force applied (Fig. 2F, *p=0.019,

main effect within n.inc+vehicle). We found no differences in thermal or chemical firing rate or

instantaneous frequencies between groups (Supp. Fig. 2). The proportion of responders in AP

treated animals resulted in a significantly higher number of responders to mechanical stimuli

(42% increased to 61%) with a corresponding, non-significant, reduction (43% decreased to

27%) in the number of non-responder "silent" neurons (Fig. 2G, *p=0.049, #p=0.090) compared

to vehicle controls. We observed no change in the proportion of polymodal neurons (Supp. Fig.

2). Together, these data indicate that the temporary loss of macrophages during neonatal injury

changes inherent sensory neuron dynamics and decreases the ability of nociceptors to change

their responsiveness over increasing mechanical forces.

Adoptive transfer of "primed" macrophages drives increased mechanical
hypersensitivity.

We next wanted to determine if macrophages isolated from neonatal incised animals were

sufficient to evoke a "primed" response in older mice. To do this we performed adoptive transfer

(AT) experiments in which macrophages from one animal (donor) were isolated and transferred

to a sex and age-matched naïve host (Fig. 3A). The *Csf1r* driver in MaFIA animals also

promotes GFP expression so we used this strain to isolate and sort peritoneal GFP+

macrophages (Fig. 3B). We transferred the isolated cells into to the right hind paw of naïve

hosts at P35. We first ensured that the AT was accepted by dissecting the site of injection as

well as other immune-related tissues. We found GFP+ cells in WT hosts three weeks after the

AT specifically at the muscle injection site and not in the nearby bone marrow, lymph node, nor

spleen (Fig. 3C). After confirming the transfer, we found that AT of unprimed macrophages by

itself (n.sham+AT alone) into an uninjured paw had no effect on paw guarding, but injection of

primed cells (n.inc.+AT alone) by itself induced low-level paw guarding that lasted for one day

(Fig. 3D, *p=0.044 vs. BL), and slightly reduced muscle squeezing thresholds, with no

difference between groups (Fig. 3E, *p<0.05 Day 1, Day 3 vs. BL). To determine how exposure

to injury might evoke a differential behavioral response, we also paired the AT with a P35

incision in different animals. We found that primed cells (n.inc.+AT with P35 inc), but not

unprimed cells (n.sham+AT with P35 inc), when transferred into an injury environment

increased mechanical hypersensitivity compared to controls measured by muscle squeezing

(Fig. 3E, ^p<0.01 Day 1, Day 3 vs. n.sham+AT alone). This effect was not observed in paw

guarding where donor neonatal injury did not impact host behavioral outcomes, indicating a

modality-directed phenotype. These data indicate that an early life injury drives changes to

macrophages which are sufficient to drive some of the enhanced behavioral phenotypes due to

injury in a host environment.

Early life incision alters the peripheral macrophage epigenetic landscape.

To determine what molecular changes might be occurring in these immune cells because of

early life incision, we collected peritoneal macrophages from three different conditions of

reporter animals (LysM;tdTom): naïve animals with macrophages isolated at P7 (naïve isolated

at P7), naïve macrophages isolated at P35, and P7 incised animals with macrophages isolated

at P35 (P7 inc, isolated at P35 (primed)) (Fig. 4A). To test if early life injury altered the

epigenetic landscape of macrophages, we performed an Assay for Transposase-Accessible

Chromatin sequencing (ATAC-seq) to measure chromatin accessibility. Quality control

confirmed our isolation and predicted that the sequenced cells were macrophages (Supp. Fig.

3). Principle component analysis (PCA) revealed distinct clustering of the three groups,

indicating an overall effect of early life incision (Fig. 4B). We repeated these same conditions to

evaluate gene expression using RNA-seq and found somewhat similar clustering in our bulk

RNA-seq PCA (Fig. 4C). We evaluated the top 100 differentially accessible chromatin sites (Fig.

4D) and top 100 differentially expressed genes (DEG) (Fig. 4E) from age-matched

macrophages in both the P35 naïve and primed groups. We obtained 10 candidate genes that

were both differentially accessible and expressed in the same direction because of early life

incision (Fig. 4F). From this sequencing data, we found that a promising candidate, NGFR (aka.

p75NTR), was differentially regulated in macrophages due to neonatal injury (Fig. 4G-H).

p75NTR has long been of interest for nervous system development and nociception, and it has

roles in macrophage differentiation and function[28,31]. We therefore decided to test whether

macrophage p75NTR could modulate neonatal nociceptive priming phenotypes.

To do this, we crossed the inducible LysMCreERT2 animal with a p75NTR floxed animal[48]. All

littermates were first incised at P7 to induce priming. Beginning at P33 and continuing until P37,

animals were injected with tamoxifen to eliminate p75NTR from LysM+ cells. At P35, all animals

received a second incision to evoke the primed macrophage and behavioral responses.

Behavior was evaluated prior to and for three weeks following the P35 incision (Fig. 5A). To

verify our injection protocol, we obtained samples following the final injection of tamoxifen at

P37. We found an overall decrease of p75NTR mRNA from total peritoneal isolated cells in

knockout animals compared to controls (-43±51% LysMCreERT2;p75fl/fl vs. controls). A P35

incision resulted in a number of F4/80+ macrophages within the injured tissue, of which 31%

were positive for p75NTR in control genotypes (Fig. 5B). The knockout of p75NTR from LysM+

cells resulted in a significant reduction in co-stained (F4/80+ and p75+) macrophages (Fig. 5C,

*p<0.05 vs. controls). Behaviorally, we observed that the knockout of p75NTR specifically in

macrophages did not prevent long-lasting paw guarding (Fig. 5D), but significantly blunted the

severity of the mechanical sensitivity one day following injury compared to controls (Fig. 5E;

^p=0.01 vs. controls). We further determined that hypersensitivity of the contralateral paw

(Supp. Fig. 3, contralateral data from Fig. 1) was also inhibited by p75NTR knockout in LysM+

cells (Supp. Fig. 3). These data indicate that p75NTR has a partial but critical role in the

macrophage-driven cellular memory that drives nociceptive priming.

Knockout of p75NTR in macrophages alters neuroimmune communication in mouse and
in human iPSCs.

We then wanted to test if macrophage p75NTR modulated nociceptive priming through its role

in inflammation in a system that could maintain a long-term immune memory. To do this, we

began evaluating bone marrow derived macrophages (BMDMs) and how they respond to

neurotrophins. First, we isolated bone marrow from naïve animals at P35, differentiated them

into macrophages[49], and measured cytokine release in response to different stimuli using

cytokine arrays. We tested the effect of knocking down p75NTR using an siRNA strategy

(example arrays Fig. 6A; siRNA transfection Supp. Fig. 4). We tested the effect of stimulation of

two neutrophins, NGF and BDNF, which both bind to p75NTR in addition to TrkA and TrkB,

respectively. We compared these to negative (vehicle) and positive (lipopolysaccharide and

interferon gamma (LPS+IFNγ)) controls (Fig. 6B; full data Supp. Fig. 4). In murine cells, the

knockdown of p75NTR in macrophages alone (sip75+Vehicle (Veh)) affected both pro- and anti-

inflammatory cytokines. If stimulated with LPS+IFNγ, the response was predominately pro-

inflammatory. NGF did not appear to alter many proteins alone in control cultures (Supp. Fig. 4),

however, NGF+p75NTR knockdown caused a significant down regulation of pro-inflammatory

cytokine release and increased anti-inflammatory cytokine production (*$p<0.05$ vs. siCON+Veh,

^$p<0.05$ vs. sip75+LPS+IFNγ, #$p=0.05$ vs. sip75+Veh, ##$p>0.05$ vs siP75+Veh; Fig. 6C). Fewer

changes were observed when BDNF or a combination of NGF and BDNF was used (Supp. Fig.

4). Together with our previous data, this may indicate that the epigenetic regulation of p75NTR

in macrophages that are exposed to early life injury could promote pro-inflammatory/pro-

nociceptive factor release and the targeting of this receptor can reverse this profile.

To test if these effects are conserved in human cells to influence sensory function, we then

performed similar experiments with NGF and p75NTR knockdown in human iPSC-derived

macrophages (Fig. 6D). We extracted the media of vehicle or NGF stimulated iPSC-derived

macrophages that were treated with either control or p75 targeting siRNAs and applied it directly

to human iPSC-derived sensory neurons (Fig. 6E) loaded with the calcium indicator

Rhodamine-2 (Rhod-2; Fig. 6F). After 24 hour incubation, we discovered no difference between

control groups that received conditioned media from macrophages with NGF or vehicle, but we

did find a decreased calcium response to ATP in iPSC derived sensory neurons that were

previously incubated with conditioned media from human macrophages treated with sip75+NGF

(Fig. 6G-H). No differences between groups were found after treatment with capsaicin (Fig. 6I).

Analyzed cells were confirmed to be viable and healthy by responding to high KCl stimuli (Fig.

6G inserts). Together with our previous data, this suggests that neuroimmune signaling in

murine and human cells is impacted by p75 expression.

Transplant of neonatal "primed" bone marrow alters female incision-related behaviors.

Trained immunity describes the concept of a long-lasting cellular memory that allows the

immune system to display an enhanced response to subsequent insults. To determine the

persistence of neonatal priming *in vivo*, we repeated priming experiments from Figure 1.

However, in these experiments, we waited 20 weeks following neonatal incision, when animals

were P147, to perform the repeated HP incision followed by three weeks of behavioral testing.

We found a persistent effect of neonatal priming in a sex specific manner. While the effect of

neonatal incision was lost in the aged males (Fig. 7A, B), re-injured females displayed long-

lasting spontaneous paw guarding (Fig. 7C ^p=0.02 n.inc+P147 inc female vs. n.sham+P147

female at day 7) and mechanical hypersensitivity (Fig. 7D ^p<0.01 n.inc+p147 inc female vs.

n.sham+P147 female at days 7 and 21) compared to neonatal females that received a sham

surgery.

We next hypothesized that underlying changes to bone marrow cells caused by an early life

injury could be a mechanism of this retained behavioral and molecular "memory." Therefore, we

isolated mouse BMDMs at P35 from neonatal incised or sham controls. We found no difference

in the total number of bone marrow cells isolated due to an incision (Supp. Fig. 4). However,

when we stimulated these cultured macrophages with LPS+IFNγ as an artificial second "injury"

(Fig. 7E) we observed that early life injury "primed" these cells as they responded with a greater

pro-inflammatory profile compared to n.sham controls (Fig. 7F, *p<0.05 vs. n.sham+vehicle,

^p<0.05 vs. n.sham+LPS+IFNγ (stim'd)). These results indicate that incision induces a long-

lasting (at least 28 days) reprogramming of hematopoietic progenitors to generate macrophages

that are primed to respond more potently to inflammation.

Finally, to test if hematopoietic stem cells (HSCs) are a mechanism through which this long-term

memory is maintained and functionally relevant to nociception, particularly in females, we

performed a bone marrow transplant (BMT) experiment. To do this, we performed a neonatal

sham or incision, allowed the animals to age until P35 (where we detected an altered BMDM

expression pattern), and then performed a BMT into naïve adolescent mice (Fig. 7G). To regain

full immune functional activity following a BMT, animals were allowed to recover for 16 weeks

(P147) when almost all hematopoietic cells are derived from the donated hematopoietic stem

cells[50,51]. At this time point animals were subjected to a hind paw incision and were monitored

for another three weeks. We successfully transferred the bone marrow evidenced by the vast

majority of CD45.2+ donor cells and a minority of CD45.1+ host cells (Fig. 7H). Prior to incision

later in life, we detected an expected, but non-significant, increase in p75NTR transcript from

bulk isolated peritoneal cells (128 ± 28%, One way ANOVA n=4-5/group). In this assay we could

not isolate macrophages specifically so other immune cells such as B and T cells are

contributing to these values and likely diluting the effect. Interestingly, we found that if we

analyze female and male peritoneal cells separately, regardless of condition, females had 480 ±

25% more p75NTR than males (p=0.016, ANOVA on Ranks, Dunn's, n=4-5/sex). When we

evaluated all animal behavior independent of sex following later in life incision, we uncovered

little evidence of enhanced pain-like behaviors in either group with quick recovery in both paw

guarding and muscle squeezing (Supp. Fig. 5). Both groups displayed guarding behavior for

only one day (p<0.001 vs. BL) and variable mechanical hypersensitivity for 1-7 days (p<0.05 vs.

BL; Supp. Fig. 5). However, when we separated animals by sex, although the length of injury-

induced behaviors was not different between groups and there was no acceleration or delayed

onset sensitivity in males (Fig. 7I-J), we found a delayed onset of n.sham+BMT female-specific

mechanical sensitivity. However, the n.inc+BMT (primed) females displayed a leftward shift from

controls (Fig. 7K-L). One day after injury, female n.sham+BMT controls were not mechanically

sensitive but female n.inc+BMT primed-cell recipient females had significantly lower thresholds

than controls (^p=0.013 vs. controls). Together, these data describe an effect of early life

incision on HSCs that can influence evoked pain-like behaviors for over 100 days in females.

Therefore, the predisposition to chronic pain due to neonatal priming is enhanced in a sex-

specific manner partially due to an HSC-related immune memory of early life injury.

Discussion

In this study, we uncovered a mechanism in which the peripheral immune system can form and

maintain a "pain memory" following a neonatal hind paw incision to modulate prolonged

responses to repeated injury (graphical abstract). Macrophages were found to be both

necessary (Figs. 1-2) and partially sufficient (Fig. 3) to drive neonatal nociceptive priming

responses at the behavioral and afferent levels. This single, local insult dramatically altered the

systemic macrophage epigenetic/transcriptomic landscape (Fig. 4) and function in part through

the p75NTR receptor in mouse and human cells (Figs. 5-6). The molecular effects of early life

injury were also evident in BMDMs, and following a BMT, which induced a sex-specific

alteration in pain like behavior that was functionally relevant long-term in hosts (Fig. 7). This

study is the first to show a macrophage cellular memory formed in neonates that modulates pain

related outcomes. Together, these data help us to understand how the distinct cellular systems

are changed by an aversive event in early development to influence nociceptive processing

across the lifespan.

Previous work has evaluated the vulnerable periods that exist in the somatosensory and

immune systems[5,38,52-57] as well as genetic risk factors in children following surgery[32,58]. After

neonatal injury, the inhibitory circuits and microglial activity of the spinal cord dorsal horn

(SCDH) are dysregulated leading to hyperactivity of spinal output[59-64]. Moriarty et al found that a

peripheral nerve block was sufficient to prevent neonatal nociceptive priming, while systemic

morphine injection was not[13]. We previously demonstrated that peripheral growth hormone,

which is a vital player in neonatal nociceptive development [65], can be used to prevent neonatal

nociceptive priming [5]. This highlights the importance of peripheral input to induce nociceptive

priming and our work extends these findings to capture a neuroimmune mechanism.

Following an injury there are both local and global responses in the immune and nervous

systems[38]. Macrophages are one of the first immune cells at the injury site to promote

inflammation and subsequent wound healing. This is accomplished through damage associated

molecular patterns (DAMPs) which can be stimulated by, and also control, cytokine and growth

factor responses. Both DAMPs and the effector proteins released may also be pro-nociceptive

such as high-mobility group box 1 (HMGB1), IL1β and NGF[66-71]. ATP is another DAMP that is

present in our *ex vivo* preparation metabolite mixture which we found to alter the mechanical

firing rate of chemosensitive neurons in animals that lacked macrophages during an early life

incision (Fig. 2[67]). It may be that desensitization to a high metabolic/noxious tissue environment

was a contributor to the rescuing of priming by early life macrophage ablation (Fig. 1). The

conduction velocity slowing of group IV sensory neurons because of macrophage ablation is an

interesting result. There are reports of activity dependent slowing of cutaneous unmyelinated

fibers[72], but this is unlikely to be the driving factor here because of the lack of spontaneous

activity in these cells. There may be differences in the size of the population of neurons that

could explain this phenomenon and could correlate with the differences in proportions of each

type of fiber response detected (Fig. 2). AT experiments (Fig. 3) showed that macrophages are

only partially sufficient to induce priming. This could suggest that other cells, including other

infiltrating or resident immune cells, "nociceptive" Schwann cells[73], or even sensory neurons

themselves, may contribute neonatal nociceptive priming. However, these concepts would need

further investigation.

Neonatal macrophages are unique compared to the adult, with a distinct epigenetic landscape

(Fig. 4) and differential responses to stimuli[14,52,9,10,24,74]. The plasticity of the macrophage

epigenome is critical to differentiate monocytes to macrophages, and acute stimulation of these

cells induces epigenetic modifications[75-78]. Changes in macrophages at the epigenetic level is

one way in which these cells become either "trained" or "tolerant" to subsequent stimuli[25,26 57].

Recent work by Cobo et al reveals that infant nociceptive reflexes to noxious and non-noxious

stimuli are altered by current or previous infection[79]. It is possible that an early life aversive

event such as a surgical incision (or an infection) may generate molecular patterns that induce

epigenetic priming in macrophages as was observed in the current study. Strategies to interrupt

neuroimmune signaling may prove beneficial for these adverse injury-related effects[80-82]. It will

be necessary to determine the factors that epigenetically train macrophages for future drug

targeting.

While the priming initiation mechanism is yet unknown, we determined that p75NTR is one

critical endpoint of the epigenetic modifications. In early development, neurotrophin signaling is

necessary for sensory neuron specialization into distinct populations[83-86], which can be

modulated by p75NTR expression[31]. Macrophage p75NTR has been less studied, but HSCs

express high levels of neutrophins, and we and others found that the inflammatory response

from macrophages is directly affected by a loss of p75NTR (Fig. 6[29,87,88]). Developed immune

cells regularly contain low levels of neurotropic factor receptors, but during inflammation they

are robustly, and transiently, increased[5,28]. NGF in human derived macrophages modulates

p75NTR expression and drives its co-localization with TrkA, which we found to have persistent

upregulation in macrophages due to early life incision (data availability at NCBI: GSE224209)[88].

Further, Khodorova et al demonstrated that intraplantar NGF injections evoke expected pain-like

behaviors, but that a general inhibition of the p75NTR receptor rescued the phenotype[89].

Combined with our data, this indicates that immune cell p75NTR increases their pro-

inflammatory signature and that the ablation of this receptor can modulate pro-inflammatory

signaling, blunt the ability of macrophages to communicate to sensory neurons, and partially

prevent prolonged pain-like behaviors (Figs. 5-6).

Circulating immune cells that infiltrate injury sites are constantly replenished through the

hematopoietic stem cell pool in the bone marrow. These are highly plastic cells that are

differentially altered by injury type[90], infection[91] and development[28,92]. Our data support the

notion that peripheral injury can alter these stem cells long-term, and changes in these cells are

part of a system to maintain immune-related neonatal nociceptive priming (Figs. 4-7), however

more work will be necessary to fully understand these findings. The mechanism in which HSCs

are notified of noxious stimuli is currently unknown, although there are potential avenues to be

tested including direct immune or neuronal signaling[93], or indirect signals in the blood or

cerebral spinal fluid[94]. Regardless, the results we find are reminiscent of trained immunity where

an initial event is epigenetically retained in innate immune cells to enhance responses to

subsequent activation[26]. In the context of immunity, the outcome is increased inflammation,

while here the outcome was enhanced nociception. Evaluation of the functional effects of injury

on HSCs by BMT indicated a female-specific hastening of mechanical hypersensitivity.

Interestingly, we detected no effect in spontaneous paw guarding between early life incised and

control animals indicating a modality specific effect (Fig. 7). It may be that "primed"

macrophages have direct signaling roles that involve mechanical sensation as we saw similar,

mechanically directed, effects with AT (Fig. 3) and p75NTR knockout (Fig. 5) behaviors. An

understanding of how stem cells may be changed by peripheral injury may have important

clinical considerations for a number of conditions that impact childhood development.

There are multiple technical points to consider. The depletion of macrophages in MaFIA animals

alters animal activity and, in some studies, withdrawal thresholds[39,95]. In our neonatal behavior,

we detected no changes in baseline withdrawal reflexes nor guarding scores due to the

macrophage knockout (Fig.1C-D). However, we did observe that when we directly manipulated

macrophage presence, there was an impact on withdrawal reflexes, but not spontaneous

behaviors (Figs. 1D, 3E, and our previous work[5]). Since withdrawal thresholds are one of the

most utilized behavioral readouts, it may be prudent to consider multiple behavioral assays

when performing experiments depleting macrophages[5,21,39,95,96]. This confound is not a major

concern for adolescent/adult behaviors, however, because the animal's immune system had

recovered before the second injury (Fig. 1). Similar concerns are also avoided when performing

adoptive transfer experiments because we tested latent changes in primed macrophages, a

major strength of this study.

We considered sex as a critical factor throughout our study. In our RNA-seq, we noticed some

indications of differences between the sexes, especially in the early life incised condition. We do

not have the power to critically evaluate these against one another, however, there are multiple

lines of evidence that suggest the immune-modulated injury response differs between males

and females[e.g.97]. In our evaluations in aged animals, with or without a BMT, we observed

female-specific impact of the early life injury (Fig. 7). Transfer of HSCs alone was not sufficient

to replicate the precise timing of hypersensitivity as if the host had been injured as a neonate,

but primed HSC cells did produce a leftward shift of mechanical sensitivity, again only in

females. Recently dos Santos et al found a delayed onset hypersensitivity in aged animals

following an incision injury as well as blunted hyperalgesic priming in aged animals. This was

connected with changes in inflammation in older rodents compared to younger animals[98]. We

found a similar delay following a BMT in female incision at P147, but not in non-BMT males or

females. Nevertheless, results reveal that the relatively minor neonatal hind paw incision injury

causes "priming" of the bone marrow and resulted in a female specific long-lasting phenotype.

Indeed, when evaluating the levels of p75NTR in our BMT animals, we found that isolated

female peritoneal cells had significantly more p75NTR than males. These data, along with

substantial literature on the heightened prevalence of chronic pain in females[99], warrant

additional studies that examine sex differences in injury-induced changes in immune cells over

time to help better understand these phenomena.

Clinical studies have shown that neonatal injuries have the potential to cause a "primed"

somatosensory system, and lead to long-term alterations in nociceptive processing[11,12,100,101].

The potential therapeutic ability of p75NTR inhibitors has been recently reviewed by Malik et al

indicating a range of functional neurological disorders that are affected by p75NTR, while

Norman and McDermott review potential therapeutic strategies in blocking p75NTR signaling for

chronic pain[102,103]. New technologies that allow for the specific targeting of macrophage p75NTR

could provide the specificity to prevent the priming effect on this cell type, while avoiding off-

target effects of the nervous system[104]. Here, we discovered a change in the epigenetic

landscape of macrophages due to early life incision that modulated neonatal nociceptive

priming. Future studies designed to analyze the immune "pain memory" may lead to important

therapeutic interventions for pediatric post-surgical pain.

Author contributions:

A.J.D. and M.P.J. conceived and designed the study and wrote the manuscript. A.J.D.

completed most of the experiments and collected most of the data. A.J.D. performed the

analyses with input from M.P.J. A.F. completed the experiments at P147 and the human culture

experiments. A.M.W. completed various experiments including some IHC and qPCR. A.P.

completed bioinformatics data processing. L.F.Q. assisted in ex vivo recordings and

interpretations of data. G. D. and H.E. facilitated experiments utilizing bone marrow isolation,

differentiation and stimulation of macrophages. G.D. helped with the conceptualization of the

study. D. L. and B.W. assisted with the BMT experiment planning and execution. O.D., C.F.,

M.T.W., and L.C.K. helped perform ATAC-seq and RNA-seq processing. M.C.H. managed the

colony and performed genotyping assays.

Acknowledgements:

We would like to recognize others who have provided support for this project including: The

Research Flow Cytometry Core in the Division of Rheumatology and the Confocal Imaging Core

at CCHMC; Dr. Yoon at the Ohio State University who provided to us the p75NTRfl/fl animals;

The Center for Autoimmune Genomics and Etiology (CAGE) at CCHMC and; The

Comprehensive Mouse and Cancer Core for their assistance in the completion of the bone

marrow transplants and care for the animals. This work was supported by grants from the NIH

to M.P.J. (R01NS105715 and R01NS113965), A.J.D. (F31NS122494), D.L. (R01 HL160614),

L.C. K. (P30 AR070549), and a CCHMC ARC award (#53632) to L.C.K. and M.T.W.. D.L. is a

scholar of the Leukemia and Lymphoma Society.

 Figure Legends

 Figure 1: Early life ablation of macrophage alters responses to re-injury later in life.

 A. Schematic of the experimental design used in MaFIA mice. **B-C.** The treatment of MaFIA

 animals with AP early in life was sufficient to ablate macrophages at the injury site one day after

 the neonatal incision compared to vehicle treated controls. Macrophages are present in the hind

 paw of a P35 injured animal even after neonatal AP treatment (***$p<0.001$ vs. n.inc+vehicle and

 n.inc+AP+P35 re-inc.; one way ANOVA,Tukey's; n=3 per group). Scale bar, 10μM. **D.** Paw

 guarding one day after a neonatal incision with or without macrophage ablation indicated an

 effect of day (F=103.418) but no difference between groups (*$p<0.001$ vs. BL for AP and vehicle

 treated animals; $p=0.071$ AP treated vs. vehicle treated). **E.** Muscle squeezing in the same

 animals showed an interaction for withdrawal thresholds (F=16.647). Vehicle treated animals

 had a significant reduction from baseline (BL) (*$p=0.014$ vs. BL) while incision+AP animals were

 hyposensitive and significantly different from controls (*$p=0.007$ vs. BL; groups were ^$p=0.028$

 vs. controls; two-way RM ANOVA; n=10-11/group). **F.** Animals that received an early life

 ablation of macrophages (n.inc+AP) did not have altered baseline scores at adolescence, but

 the response to a second injury revealed a significant effect of dual incision (guarding F=5.635,

 muscle squeezing F=5.303). Groups that received a sham as neonates (n.sham+vehicle,

 n.sham+AP) guarded through day 3 (*$p<0.01$ vs. BL) while vehicle treated dually incised

 animals guarded through day 14 (*$p<0.001$ vs. BL). Animals from the neonatal incision

 (n.inc+AP) group returned to BL guarding scores after seven days (*$p<0.05$ vs. BL). At days

 three, seven, and 14, the n.inc+vehicle group was significantly different compared to all other

 groups (^$p<0.05$ n.inc+vehicle vs. controls, #$p<0.001$ vs. n.inc+AP). **G.** In the same group of

 animals, a similar pattern was observed for muscle squeezing withdrawal thresholds. N.sham

 operated groups were hypersensitive through day 7 (*$p<0.005$ vs. BL). The n.inc+vehicle group

 showed reduced thresholds through day 14 (*$p<0.001$ vs. BL), but n.inc+AP treated animals

were hypersensitive only through day 3 (*p<0.001 vs. BL). At days three, seven and 14, the

n.inc+vehicle group had significantly lower thresholds compared to all other groups (^p<0.005

vs controls, #p<0.005 vs. n.inc+AP). Colored horizontal lines indicate the duration of

significance compared to BL for each group. Data shown as mean +/- SEM.

Figure 2. Macrophage depletion in neonates alters sensory neuron sensitization

following repeated incision.

A. Representative image of the intact electrophysiological preparation containing the injured

hind paw muscle, tibial nerve, DRGs and SC. The search electrode (SE) and recording

electrode (RE) are marked. **B.** There is no difference in CVs between groups in group III muscle

afferents (p=0.987 n.inc+AP vs. n.inc+vehicle, n=34 and 39 respectively, one way ANOVA,

Tukey's) but (**C**) there is a significant reduction of group IV conduction velocities (H=3.896) in

the neonatal treated AP group compared to vehicle (*p=0.048 n.inc+AP vs. n.inc+vehicle, n=16

and 11 respectively, ANOVA on Ranks, Dunn's). **D.** There are no differences in overall

mechanical thresholds between groups (p=0.430 n.inc+AP vs. n.inc+vehicle, n=31 and 22

respectively, ANOVA on Ranks, Dunn's). **E.** There is no change in mechanical firing rate pre- to

post- metabolite treatment in vehicle treatment group, unlike that seen in the AP treatment

group (H=4.934, n.inc+vehicle: p=0.223, n.inc+AP: *p=0.031, ANOVA on Ranks, Dunn's). **F.**

There is a significant increase in firing rate over mechanical forces applied in the vehicle

treatment (H=5.760) but not in the AP treatment group (n.inc+vehicle: *p=0.019, One way

ANOVA on Ranks, Dunn's. n.inc+AP: p=0.149, One way ANOVA on Ranks, Dunn's, "X"

indicates no cell responded to the applied force). **G.** There is an increase in the number of

mechanically sensitive cells in AP treated animals compared to vehicle treated animals

(*p=0.049 n.inc+AP vs. n.inc+vehicle, Chi-Square on mechanical responders), with a

corresponding non-significant decrease in cells that responded to no stimulus ("silent",

#p=0.090 n.inc+AP vs. n.inc+vehicle, Chi-Square on silent neurons). Data shown as mean +/-

SEM.

Figure 3: Adoptive transfer of macrophages from neonatal incised animals increases

pain-like behaviors.

A. Outline of the experiment design for AT experiments. **B.** Example of the sorting of live GFP+

peritoneal macrophages from MaFIA mice. **C.** Example images demonstrating the presence of

GFP+ donated cells (arrows) in the hindpaw of a host 21 days after injection, but not in the bone

marrow (ipsilateral tibia), lymph node (potineal), or spleen. Scale bars, 25μM; BM, 150μM. **D.**

For guarding behaviors, there was an effect (F=3.132) of macrophage exposure to injury. Both

primed (n.inc+AT with P35 inc) and unprimed (n.sham+AT with P35 inc) cells injected into an

incision site induced prolonged guarding (*p<0.05 vs. BL). In the absence of injury at P35,

primed (n.inc+AT alone) cells induced one day of guarding (*p=0.044 vs. BL) while unprimed

cells (n.sham+AT alone) did not induce detectable behavior. There was a significant difference

between n.inc+AT with P35 inc and n.sham+AT alone at days 1, 3 and 7 (^p<0.001 vs.

n.sham+AT alone) and between n.sham+AT with P35 inc vs. n.sham+AT alone at days 1-14

(#p<0.05 vs. n.sham). **E.** The same animals were then tested for muscle withdrawal thresholds

by muscle squeezing which resulted in a significant interaction (F=4.318). All groups had a

significant reduction in withdrawal thresholds that varied over time depending on condition

(*p<0.05 vs. BL). N.inc+AT with P35 incision was significantly different than n.sham+AT alone at

days 1 and 3 (^p<0.01 vs. n.sham+AT alone). N.sham+AT with P35 incision vs. n.sham+AT

alone treatment showed an effect at day 14 (#p=0.015 vs. n.sham+AT alone). Two-way RM

ANOVA, Tukey's post hoc, n=8-11/group. Colored horizontal lines indicate the duration of

significance compared to BL for each group. Data shown as mean +/- SEM.

Figure 4. Early life incision alters the epigenetic and mRNA landscape of LysM+ cells.

A. Example sorting of peritoneal macrophages from P7 naïve, P35 naïve, and P7 incised

animals isolated at P35. **B.** Principle component analysis from ATAC-seq datasets

demonstrating distinct clustering of groups. **C.** Principle component analysis form the RNA-seq

dataset shows similar distinct clustering of groups. **D.** The top 100 differentially accessible

chromatin regions in macrophages in P35 naïve cells compared to P35 primed cells (age

matched). **E.** The top 100 differentially expressed genes (DEGs) for the same conditions. **F.** A

table of factors that had differentially accessible regions and DEGs. NGFR (p75NTR) is

indicated. **G.** ATAC-seq indicating higher reads of NGFR (*p=0.001 vs. controls, one-way

ANOVA, Tukey's, n=3-4/group), and (**H**) transcripts per million (TPM) data for RNA-seq

(*p=0.008 vs. controls). Data shown as mean +/- SEM.

Figure 5. Macrophage specific knockout of p75NTR after neonatal incision impacts
neonatal nociceptive priming.

A. Schematic of knocking out p75NTR from macrophages following neonatal incision. **B.**

Representative images of p75NTR labeling (green) co-stained with F4/80 (macrophages, white)

in the injured hind paw. Yellow arrows indicate double positive staining. Scale Bar, 25μM. **C.**

There is a significant reduction (F=4.012) in the percentage of double positive cells in

LysM;p75fl/fl animals compared to other genotypes (*p<0.05 vs. controls (p75+/fl and p75fl/fl)

and LysM;p75+/fl, one way ANOVA, Tukey's, n=3-5/group). **D.** We found no effect of tamoxifen

prior to the second injury (No Tam vs. BL time points) on paw guarding. Following the second

incision there was robust paw guarding (F=26.258), with a faster recovery in the

LysM;p75NTRfl/fl mice (*p<0.05 vs. BL, #p=0.05 vs BL p75fl/fl, Two-way RM ANOVA, Tukey's).

E. No effect of tamoxifen on mechanical withdrawal thresholds was found prior to injury (No

Tam vs. BL time points), however there was an effect following a second incision (F=40.516).

Both groups displayed mechanical sensitivity on days one, three and seven (*p<0.01 vs. BL).

The p75fl/fl genotype continued to display significant sensitivities at day 14 while LysM;p75fl/fl

animals did not (p75fl/fl: *p=0.002 vs. BL, LysM;p75fl/fl: p=0.063 vs. BL). At one day following

second incision, LysM;p75fl/fl animals were less hypersensitive than controls (^p=0.01 vs.

controls). Two-way RM ANOVA, Tukey's, n=7/group.). Colored horizontal lines indicate the

duration of significance compared to BL for each group. Data shown as mean +/- SEM.

**Figure 6. p75NTR expression in macrophages alters the inflammatory signature of
macrophages and neuronal activity.**

A. Examples of the protein arrays that were run on the lysates of BMDMs that received a

knockdown of p75NTR and were stimulated with indicated factors. **B.** Heatmap indicating the

protein expression detected across each stimulus, n=4/group. **C.** Examples of significantly

regulated proteins detected from the protein array (*p<0.05 vs. siCON+Veh, ^p<0.05 vs.

sip75+LPS, #p=0.05 vs. sip75+Veh, ##p>0.05 vs sip75+Veh; one-way ANOVA, Tukey's or

ANOVA on Ranks, Dunn's. n=4/group). **D-F.** Representative images of iPSC-derived

macrophages, sensory neurons and Rhod-2 responses to ATP. Arrows indicate increased

responses. Arrowheads indicated new responses. **G.** Example calcium transients from iPSC

derived sensory neurons in response to ATP, capsaicin or KCl stimulation. The arrow indicates

when the ATP or capsaicin was added, and the brackets indicate the part of the trace that was

analyzed to calculate changes in fluorescence intensity in direct response to a stimulus. Scale

Bar, 20μM. **H.** Exposure of iPSC derived sensory neurons to media from macrophages with loss

of p75NTR treated with NGF caused decreased responsiveness of neurons to ATP (F=4.257)

(*p<0.04 vs. all other conditions, one way ANOVA with Tukey's post hoc, n=15 cells per group).

I. No changes in calcium responses to capsaicin (F=2.702) were observed (one way ANOVA).

Data shown as mean +/- SEM or percent change from controls with variance.

Figure 7. Bone marrow transplant from neonatal incised animals transfers a "pain

memory" to naïve animals.

A, C. There is a main effect of group (F=6.96) and day (F=27.20) for guarding with post hoc

analysis revealing that n.inc+P147 inc females guard through day 14 (*p<0.05 vs. BL) and

significantly more than n.sham +P147 females at day 7 (^p=0.02). n.sham+P147 females guard

through day 3 (*p<0.001 vs. BL) while both male groups only guard at day one (*p<0.05 vs. BL).

Two-way RM ANOVA, Tukey's. **B, D.** In the same animals for muscle squeezing there is an

interaction effect (F=3.50). n.inc+P147 inc female have significantly reduced thresholds from

baseline for three weeks (*p<0.05 vs. BL) and are significantly more hypersensitive than

n.sham+P147 inc females at days seven and 21 (^p<01 vs. n.sham+P147 inc female).

N.sham+P147 inc females have reduced thresholds at days one, three and 14, while males

from both groups are hypersensitive at day one only (*p<0.05 vs. BL). Two-way RM ANOVA,

Tukey's, n=11-13/injury condition, 4-7/sex in condition. **E.** A schematic indicating timeline for

neonatal nociceptive priming experiments separating injury by 140 days. **F.** Analyses of the

transcripts from BMDMs after stimulation with LPS+IFNγ (*p<0.05 vs. n.sham no stimulation

controls, ^p<0.05 vs. n.sham with stimulation (stim'd), n=3-8/group, Two-way ANOVA, Tukey's).

G. Schematic of the bone marrow transplant experiments. **H.** Representative cell sorting of

living cells, 45.1+ host remaining cells and 45.2+ transferred cells. **I, K.** After recovery from the

BMT, there is a main effect of day (F=22.90). Animals display no guarding behaviors prior to

incision. Following an incision both n.sham and n.inc groups from both sexes guarded for one

day only (p<0.01 vs. BL, Two-way RM ANOVA, Tukey's). **J, L.** In the muscle squeezing assay,

n.inc+P35 BMT+P147 inc. females are hypersensitive to baseline at days one and three

(*p<0.05 vs. BL) and have significantly lower thresholds than controls at day one (^p=0.013 vs.

n.sham+P35 BMT+P147 inc female). n.sham+P35 BMT+P147 inc females do not begin to

display sensitivity until day seven (*p=0.001 vs. BL). N.sham+P35 BMT+P147 inc males have

reduced thresholds from BL at days one and three (*p<0.05 vs. BL). Two-way RM ANOVA,

Tukey's, n=10-11/injury condition, 4-6/sex in condition. Colored horizontal lines indicate the

duration of significance compared to BL for each group. Data shown as mean +/- SEM.

Methods

Animals. Animals of both sexes were used for all studies. Neonatal animals were considered at

postnatal day 7 (P7). Adolescent animals begun experimentation at P35-P42. Adult animals

were analyzed at P147. Sex was analyzed as a critical biological factor for each experiment,

and differences are noted in figure legends. In an effort to reduce stress, neonatal animals were

kept with the dam when not actively analyzed behaviorally, and experiments were kept as short

as possible (~1 hour, never more than 1.5 hours). In all behavioral experiments, mice were

acclimated to the room and were performed by the same experimenter under the same

conditions including time of day, location, lighting and temperature. Animals were used +/- 1 day

from the indicated age. All animals were kept in an environment-controlled facility at Cincinnati

Children's Hospital Medical Center with free access to food and water while on a 14/10-hour

light/dark cycle.

C57/Bl6 (CD45.2$^+$) mice were bred in-house or purchased from Jax (Stock#: 000664).

B6.SJL/BoyJ (CD45.1$^+$) mice were bred in the CCHMC transplant core. Macrophage fas-

induced apoptosis (MaFIA) animals were purchased from Jax (Stock#: 005070) and bred in-

house. Myeloid/Macrophage reporter animals were generated by crossing the LysM-Cre

positive animal (see Jax Stock#: 004781) with the tdTomato (tdTom) reporter mouse (B6.Cg-

Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J) purchased from The Jackson Laboratory (Stock#: 007914).

Myeloid/Macrophage specific p75NTR knockout animals were generated by crossing the

tamoxifen inducible LysM-CreERT2 animal (Jax 031674) to the p75NTRfl/fl animal that was

kindly donated to us by Dr. Sung Yoon at the Ohio State University. All procedures were

approved by the CCHMC Institutional Animal Care and Use Committee in compliance with

AALAC approved practices.

Behavioral Measures. Neonatal animals were transferred from their home cage to opaque

chambers fitted with a translucent lid. They were acclimated for 10 minutes prior to any

assessments. Adolescent animals were transferred from their home cage to raised chambers

fitted with black side walls, translucent viewing front, back and lids, and a grid mesh bottom.

These animals were acclimated for 30 minutes prior to any assessments. Behavioral data were

gathered at neonatal baseline and one day (1d) following early life injury. Further data was

gathered at adolescent baseline and 1d, 3d, 7d, 14d and 21d following adolescent injury.

Behavioral assessments were performed as described before[5] and below, and included a

measure of spontaneous pain-like behavior as well as evoked muscle squeezing pain-like

behavior.

Spontaneous pain-like behavior was measured as described previously by qualitatively

assessing paw weight preference on a scale of zero to two, where zero is no guarding after

injury, 1 indicates a shift in weight off of the injured paw, and 2 indicates a complete removal of

the injured paw from the platform[5]. Assessments were made for a duration of 1 minute every 5

minutes for 30 minutes (neonates) or 1 hour (adolescents).

Muscle-directed withdrawal scores were measured using a digital pressure device (IITC Life

Science Inc. Woodland Hills, CA, USA) with a dulled probe attachment ~2 mm wide at the tip. In

this assay, the animal was lightly scruffed and was held upright. One at a time, the hind paws

were inserted between the two arms of the device. The top arm has a flat platform to hold the

dorsal hindpaw in place, while the bottom has the dulled probe. Slowly, the medial plantar paw

was pressed by the probe while the animal and force were observed. When the animal gave a

robust withdrawal response, the force in grams was recorded. This was repeated for a total of

three trials for each animal, separated by at least 5 minutes between trials for recovery. The

maximum squeezing force implemented to a neonate was 150 grams and for an adolescent was

350 grams.

Injections. To induce cell apoptosis in MaFIA animals, the designed drug AP20187 (AP; B/B

Homodimerizer purchased from Takara #635058) was first prepared according to

manufacturer's directions in 100% ethanol. Prior to each experiment, fresh vehicle was

prepared according to manufacturer's directions for 10 mg/kg which contained 10% PEG-400,

86% Tween and 4% drug/vehicle. All animals were injected with a final volume of 30 μL by

intraperitoneal (IP) injections. The drug or vehicle was added to the mixture immediately before

injecting the animal. AP20187 has no known impact on wildtype (WT) animals and previous

reports demonstrate that neonates injected at P1, P2, P3 and P7 show robust knockout of Csf1r

expressing cells in tissue[41,105]. Tamoxifen (1000 μg at 10 mg/mL) was injected in all animals

within a litter of potential LysM-CreERT2;p75NTRfl/fl genotype. The injection protocol began at

P33 and was completed i.p. daily for 5 days. Animals injected with Evan's Blue Dye (EBD,

Sigma E2129-10) were injected as adolescence 16-20 hours prior to dissection with 200 μL of

1% EBD in sterile 0.9% saline.

Surgical Hind Paw Incisions. Animals were anesthetized with 2-3% isoflurane. Once

sufficiently anesthetized, animals were placed on their backs and their right hind paw was taped

to a sterile cloth. The dorsal skin was disinfected using Chlorhaxiderm Scrub and 70% isopropyl

alcohol. A longitudinal incision was made through the hairy skin lateral to the main saphenous

innervation territory. The incision was continued in between the bones through to the body of

the flexor digitorum brevis muscles. The muscles were then manipulated using blunt dissection

techniques with #5 (neonate) and #3 (adolescent) forceps. The plantar skin was left untouched.

The skin was sutured using 7-0 or 6-0 sutures for neonates and adolescents, respectively.

Sham surgeries were performed exactly the same, except animals received only a suture

through intact hairy hindpaw skin with no incision. Dual incision regimes were performed at both

P7 and P35-P42, or P147 to the same hindpaw muscle.

Dissections. For peripheral sites, the animal was heavily anesthetized with ketamine and

xylazine (100 and 16 mg/kg, respectively) and the tissue was removed. This included hind paw

muscle, tibia, potineal lymph node and spleen. For neural tissue, the animal was first cardiac

perfused with ice-cold saline prior to dissection where ipsilateral DRGs were removed.

Realtime PCR and Protein Arrays. RNA was isolated from BMDM cultures at the indicated

time points using RNeasy Mini Kits (Qiagen Stock#: 74104). All isolation were performed exactly

according to the manufacturer's instructions. 500ng of RNA was reverse transcribed to cDNA

and realtime PCR was performed using SYBR Green Master Mix on a StepOne realtime PCR

system (Applied Biosystems). Quantitative PCR was analyzed by the ΔΔcycle threshold (CT)

method with normalization to GAPDH. Differences in expression and standard error are

determined from the normalized ΔΔCt. This is used to calculate fold change between conditions

and values are then converted to a percent change where 2-fold = 100% change[106]. Cytokines

from BMDMs were analyzed using proteome profilers (R&D Systems ARY006) loaded with 700

μLs of media. One membrane was removed due to a lack of positive control expression (siCON

stim LPS+IFNy). Data was first normalized to the average positive control before percent

change from experimental control was calculated.

ATAC-Seq and RNA-seq. Peritoneal macrophages were isolated as indicated below from P7

naïve, P35 naïve, and P7 incised isolated at P35 animals. Samples were pooled across two

animals of the same age and sex. Samples for ATAC- and RNA-Seq were processed by the

sequencing core at CCHMC. Data were analyzed by the Bioinformatics Collaborative Services

at CCHMC. Briefly, shared peaks between samples were obtained by merging peaks at 50%

overlap with BEDtools v2.27.0 between replicates (peaks in all biological replicates) and

treatment (merge at 50% all replicate peaks). These were then converted to Gene Transfer

Format and peaks were counted with feature counts v1.6.2 (Rsubread package). Raw counts

were normalized as transcripts per million and peaks were annotated by the nearest or

overlapping gene and genomic features using an in-house script defined by: Promoter = 1kb

upstream and downstream from peak; upstream = between 21 and 1kb from peak; priority =

promoter>upstream>exon>intron. The R package DESeq2 v1.26.0 was used for differential

analysis, with significance set to log2 fold change of 0.58 and FDR less than or equal to 0.05.

RNA-seq was prepared similarly, where FASTQ formats were aligned to mm9 by STAR v2.6.le

and stripped of duplicates by Sambamba v0.6.8. Read counts and differential expression were

defined as before. Plots were generated by ggplot2 or PlotsReasy from CCHMC.

781
782

Immunohistochemistry. Freshly dissected tissue was frozen in OCT medium on dry ice or

snap frozen in liquid N_2 for skeletal muscle and was then stored at -80ºC. On a cryostat, tissue

was sectioned at 20 µm and was melted onto a slide. Tissue was fixed in 4% paraformaldehyde,

washed and blocked prior to primary antibody (WGA 1:500 (Invitrogen W32466); Dystrophin

1:250 (Abcam ab15277); p75NTR 1:1000 (Cell Signaling 82385); F4/80 1:500 (Abcam ab6640))

incubation overnight. The next day, the tissue was washed and stained with secondary

antibodies before cover slipping and stained with DAPI (Fisher Scientific 17985-50.) For tissue

that was genetically marked with a fluorescent dye, the tissue was post-fixed, washed, and

mounted with media containing DAPI. Analysis was completed on Nikon's NIS Elements where

two-three nonconsecutive sections across two-three slides were averaged to create a score for

each animal. Quantification of GFP+ macrophages in Fig. 1 were performed using Image J

following similar procedures.

Macrophage Isolation. *Peritoneal:* Animals were rapidly killed by cervical dislocation and

cleaned with 70% ethanol. The peritoneum was injected with 1 mL (neonate) or 5 mLs

(adolescent) of 3% fetal bovine serum (FBS) until bloated. The midsection was gently

massaged to dislodge any cells, and then peritoneal fluid was extracted. Extracted peritoneal

serum was processed for RNA extraction immediately, or sorted by the Flow Cytometry core at

CCHMC by the LysMtdTom or MaFIA (GFP+) expression depending on the experiment and

animal used. Cells were sorted in 2% FBS and into 50% FBS or PBS. Dead cells were marked

with 7AAD and were removed from selection. Gating strategies were placed by the flow core

and examples are presented.

Bone marrow derived macrophages (BMDM): Animals were prepared as above. The right leg

was then detached from the body and placed into a dish containing HBSS. The femur and tibia

were isolated, and the bone marrow was visualized by making horizontal cuts on either side of

the knee. The bone marrow was flushed out of the bone with HBSS and a 22-gauge needle.

HBSS with cells was collected, spun at 4ºC and 1500 RPM for 5 minutes. The supernatant was

discarded, and RPMI buffer (Catalog #11875093) supplemented with 10% fetal bovine serum

(35-010-CV Lot 35010171) and gentamycin was added to resuspend the cells. GM-CSF (60

ng/mL) was added, and cells were incubated for 5 days at 37ºC and 5% CO_2. Additional media

and GM-CSF was added halfway through the incubation. After maturation, cells were treated

with siRNAs (Origene 10 ng/uL) and stimulants (NGF, BDNF 50 ng/uL or LPS+IFNγ 10 ng/uL

each) as indicated in the figure legends. Knockdown efficiency tested across three duplexes,

achieved 90% by PCR. Cells were stimulated for 24 hours before washing, trypsinizing, and/or

scraping of the plates. Media was saved for protein array analyses. Cells were counted using a

hemocytometer and then resuspended in buffer for RNA isolation using the RNeasy Mini Kit.

Adoptive Transfer. Peritoneal macrophages from MaFIA animals were isolated as described

above. ~80k cells were sorted into PBS. These were pelleted and reconstituted in 10 µLs of

sterile PBS. Naïve hosts were anesthetized and the right hindpaw was injected with all of the

cells directly into either intact muscle, or the wound of a hindpaw incision before suturing.

Animals were monitored following the transplant for adverse effects and behavioral responses.

Bone Marrow Transplant. Tibia and femur bone marrow was collected using sterile tools from

adolescent animals. Under a sterile hood, bones were flushed with PBS and treated with red

blood cell lysis buffer until a transparent pellet was formed after centrifugation at 1100 RMP for

4 minutes at 4ºC. After reconstitution, cells were passed through a 70 µm filter and were

counted on a hemocytometer. One million cells were transferred to a naïve age and sex

matched host that was previously exposed to radiation (administered in split doses, the first at

700 rads/700 cGy followed 3 hours later with a second dose of 4.75 Gy (475 rads/475 cGy)) to

prepare for the transplant. Animals were monitored following the transplant in their immune

compromised state. We successfully completed 28 out of 32 BMTs. The animals were allowed

to recover for 16 weeks prior to further experimentation, to ensure a functional immune

response after transplant[50,51].

Human induced pluripotent stem cell preparations and calcium imaging. Human induced

pluripotent stem cells (iPSC) derived CD14+ monocytes (ATCC, Manassas, VA, Cat No.

DYS0100) were plated in macrophage differentiation medium (RPMI-1640 Medium /10% Fetal

Bovine Serum (FBS), supplemented with 100 ng/ml human M-CSF, and 1% Pen/Strep) in 24

well plate at a density of 10^5 cells/ml according to manufacturer's directions. Monocytes were

differentiated into macrophages within seven days. Half medium change and siRNA treatment

were done at day three and day five, respectively. Briefly, 10 nM control (siCON), or siRNAs

against p75NTR (sip75; Origene SR321107) were incubated with macrophages at day five in

macrophage differentiation medium. At day six, macrophages treated with siRNAs were

stimulated with either 50 ng/ml NGF or vehicle in macrophage differentiation medium for 24

hours. At day seven, the media was extracted and used for stimulation of the sensory neurons.

Human iPSC-Derived Sensory Neuron Progenitors (AXOL; ax0555) were cultured on

SureBond-XF-coated 96 well plate with Neural Plating Medium (AXOL, product codes: ax0053

and ax0033) for 24 hours as recommended by the manufacturer. Briefly, full media change was

achieved by replacing the neural plating medium with sensory neuron maintenance medium

supplemented with growth factors (25 ng/mL Glial-Derived Neurotrophic Factor (ax139855), 25

ng/mL Nerve Growth Factor (ax139789), 10 ng/mL Brain-Derived Neurotrophic Factor

(ax139800), and 10 ng/mL Neurotrophin-3 (ax139811) and sensory maturation maximizer

(ax0058) at a seeding density of 50000/cm^2. Cells were treated with 2.5 µg/ml mitomycin C at

day three after which the media was replaced with sensory neuron maintenance medium

supplemented with growth factors and sensory maturation maximizer. Half media change was

done at every three days for 21 days. All iPSC cultures and incubations were carried out in 5%

CO_2/ 37°C.

For imaging, cells were pretreated with 5 µg/ml Rhodamine (in DMSO, ThermoFisher R1244)

for 30-45 minutes and further dosed with equal volume of siRNA treated and stimulated

macrophage media. Neuronal activities were visualized using wide-field Nikon Ti2 inverted

SpectraX with 10X LWD Phase high quality objective. To analyze changes in fluorescence,

regions of interest (ROI) were randomly drawn around 15 cells across triplicate conditions.

Calcium transients were measured for after stimulation with ATP (200µM) followed by capsaicin

(1µM). KCl (300µM) was used as a positive control for cell viability. ΔF/FO were calculated

where ΔF/FO = (Fmax – FO) / FO where Fmax: maximum intensity during stimulation and FO:

the average intensity of immediately prior to stimulation.

***Ex vivo* Preparation.** This preparation has recently been described in detail[95]. In this study,

pups were treated with either AP or vehicle as described above and were subjected to a P7

incision followed by a P35-P42 incision. Briefly, 7d after the second incision we dissected the

animal HP muscle/tibial nerve/DRG/SC in continuity in ice cold oxygenated (95%O2/5% CO2)

artificial cerebral spinal fluid (aCSF; 127.0 mM NaCl, 1.9mM KC l, 1.2 mM KH2PO4, 1.3 mM

MgSO4, 2.4 mM CaCl2, 26.0 mM NaHCO3, and 10.0 mM D-glucose: O2aCSF). The

preparation was transferred to a new dish where the muscle and DRGs were isolated in

separate circulating baths of O2aCSF. The temperature was slowly raised to 32°C. In L3 or L4

an electrode was filled with 5% neurobiotin and 1M potassium acetate for sharp electrode single

unit recordings using Quartz electrodes (impedance>150MΩ). A suction electrode was placed

on the side of the tibial nerve to send search stimuli. An impaled cell was identified by the

presence of an action potential (AP) recording from the search stimulus. A receptive field on the

muscle was then located using a concentric bipolar electrode. If found, the cell was counted in

our analyses and mechanical, thermal and chemical stimuli were applied in that order. First von

Frey fiber thresholds were identified using fibers from 0.07-10g of force applied for ~1-2

seconds. If the cell did not respond to the maximum, physical push was applied to determine if

the cell would respond to any mechanical force. Next ice-cold saline (~2°C) was applied to the

circulating bath over the receptive field, followed by hot ~53°C saline. After washout, a low

concentration mixture of lactic acid, protons and ATP (applied just before stimulation) were

oxygenated and flooded the muscle chamber through an in-line heater for 2 minutes (15 mM

lactate, 1μM ATP, pH 7.0). This was repeated with a high concentration of the same metabolites

(50 mM lactate, 5 μM ATP, pH 6.6). Mechanical and thermal responses were then repeated.

Activity was recorded by the Spike2 program (Cambridge Electronic Design) and was analyzed

offline. At the end of the preparation the nerve length from suction electrode to DRG was

measured to calculate conduction velocity (CV), where a Group IV afferent was defined as

being (≤1.2 m/s) or Group III afferents (1.2–14 m/s). Mechanical thresholds were set at the

minimum force required to evoke an action potential, firing rate (FR) was calculated as the

maximum number events that occurred over a 200 ms bin, and instantaneous frequency (IF)

was calculated to determine the maximum response to a stimulus. Distribution analyses were

completed to determine if there was a shift in the number of responders in our conditions.

Statistical Analyses. Data was analyzed using SigmaPlot software (v14.5). Critical

significance value was set to $\alpha < 0.05$. All data were first checked for normality by Shapiro-Wilk

to determine parametric or nonparametric tests. Individual tests used are described in the figure

legends. Behavioral data of the same animal over time and between groups were analyzed

using a two-way repeated measures (RM) ANOVA. In situations in which different animals were

compared across multiple time points and groups, such as PCR data, a two-way ANOVA was

used. Data at a single time points across multiple groups including much of the

electrophysiological data were analyzed by a one-way ANOVA. Proportions data were analyzed

using a Chi-squared or Fisher's exact test as marked in the figure legends. Graphs were made

using GraphPad Prism (v9). In all experiments in which there is a potential for bias including

behavior, electrophysiology, cell culture and IHC the investigator was blinded to the conditions.

For behavior, animals were assigned to groups by a random number generator. Biological and

technical replicates were completed for each experiment and are appropriately marked in the

figure legends.

920
921
922
923

References.

Udit, S., Blake, K. & Chiu, I. M. Somatosensory and autonomic neuronal regulation of the immune response. *Nature Reviews Neuroscience* **23**, 157-171 (2022). https://doi.org/10.1038/s41583-021-00555-4

Tamari, M., Ver Heul, A. M. & Kim, B. S. Immunosensation: Neuroimmune Cross Talk in the Skin. *Annu Rev Immunol* (2021). https://doi.org/10.1146/annurev-immunol-101719-113805

Karshikoff, B., Tadros, M. A., Mackey, S. & Zouikr, I. Neuroimmune modulation of pain across the developmental spectrum. *Current opinion in behavioral sciences* **28**, 85-92 (2019). https://doi.org/10.1016/j.cobeha.2019.01.010

Spencer, S. J., Galic, M. A. & Pittman, Q. J. Neonatal programming of innate immune function. *Am J Physiol Endocrinol Metab* **300**, E11-18 (2011). https://doi.org/10.1152/ajpendo.00516.2010

Dourson, A. J. *et al.* Early Life Nociception is Influenced by Peripheral Growth Hormone Signaling. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 4410-4427 (2021). https://doi.org/10.1523/jneurosci.3081-20.2021

Fitzgerald, M. The development of nociceptive circuits. *Nature Reviews Neuroscience* **6**, 507-520 (2005). https://doi.org/10.1038/nrn1701

Jankowski, M. P. *et al.* Age-dependent sensitization of cutaneous nociceptors during developmental inflammation. *Molecular pain* **10**, 34 (2014). https://doi.org/10.1186/1744-8069-10-34

Jones, L. *et al.* Widespread nociceptive maps in the human neonatal somatosensory cortex. *eLife* **11**, e71655 (2022). https://doi.org/10.7554/eLife.71655

Kumar, S. K. & Bhat, B. V. Distinct mechanisms of the newborn innate immunity. *Immunology letters* **173**, 42-54 (2016). https://doi.org/10.1016/j.imlet.2016.03.009

Winterberg, T. *et al.* Distinct phenotypic features of neonatal murine macrophages. *European journal of immunology* **45**, 214-224 (2015). https://doi.org/10.1002/eji.201444468

Walker, S. M. *et al.* Somatosensory function and pain in extremely preterm young adults from the UK EPICure cohort: sex-dependent differences and impact of neonatal surgery. *British journal of anaesthesia* **121**, 623-635 (2018). https://doi.org/10.1016/j.bja.2018.03.035

Hermann, C., Hohmeister, J., Demirakca, S., Zohsel, K. & Flor, H. Long-term alteration of pain sensitivity in school-aged children with early pain experiences. *Pain* **125**, 278-285 (2006). https://doi.org/10.1016/j.pain.2006.08.026

Moriarty, O., Harrington, L., Beggs, S. & Walker, S. M. Opioid analgesia and the somatosensory memory of neonatal surgical injury in the adult rat. *British journal of anaesthesia* **121**, 314-324 (2018). https://doi.org/10.1016/j.bja.2017.11.111

Fragiadakis, G. K. *et al.* Patient-specific Immune States before Surgery Are Strong Correlates of Surgical Recovery. *Anesthesiology* **123**, 1241-1255 (2015). https://doi.org/10.1097/aln.0000000000000887

15 Laudanski, K. *et al.* Acquired immunological imbalance after surgery with
cardiopulmonary bypass due to epigenetic over-activation of PU.1/M-CSF. *J Transl Med*
16, 143 (2018).
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5970449/pdf/12967_2018_Article_1518.
pdf>.

16 Koennecke, L. A., Zito, M. A., Proescholdt, M. G., van Rooijen, N. & Heyes, M. P.
Depletion of systemic macrophages by liposome-encapsulated clodronate attenuates
increases in brain quinolinic acid during CNS-localized and systemic immune activation.
Journal of neurochemistry **73**, 770-779 (1999).

17 Shutov, L. P. *et al.* The Complement System Component C5a Produces Thermal
Hyperalgesia via Macrophage-to-Nociceptor Signaling That Requires NGF and TRPV1.
The Journal of neuroscience : the official journal of the Society for Neuroscience **36**,
5055-5070 (2016). https://doi.org/10.1523/jneurosci.3249-15.2016

18 Wang, Y. R., Mao, X. F., Wu, H. Y. & Wang, Y. X. Liposome-encapsulated clodronate
specifically depletes spinal microglia and reduces initial neuropathic pain. *Biochemical*
and biophysical research communications **499**, 499-505 (2018).
https://doi.org/10.1016/j.bbrc.2018.03.177

19 Mert, T. *et al.* Macrophage depletion delays progression of neuropathic pain in diabetic
animals. *Naunyn-Schmiedeberg's archives of pharmacology* **379**, 445-452 (2009).
https://doi.org/10.1007/s00210-008-0387-3

20 Shepherd, A. J. *et al.* Angiotensin II Triggers Peripheral Macrophage-to-Sensory Neuron
Redox Crosstalk to Elicit Pain. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience **38**, 7032-7057 (2018). https://doi.org/10.1523/jneurosci.3542-
17.2018

21 Shepherd, A. J. *et al.* Macrophage angiotensin II type 2 receptor triggers neuropathic
pain. *Proceedings of the National Academy of Sciences of the United States of America*
115, E8057-E8066 (2018). https://doi.org/10.1073/pnas.1721815115

22 Pannell, M. *et al.* Adoptive transfer of M2 macrophages reduces neuropathic pain via
opioid peptides. *Journal of neuroinflammation* **13**, 262 (2016).
https://doi.org/10.1186/s12974-016-0735-z

23 Jupelli, M. *et al.* Chlamydia pneumoniae infection in mice induces chronic lung
inflammation, iBALT formation, and fibrosis. *PloS one* **8**, e77447 (2013).
https://doi.org/10.1371/journal.pone.0077447

24 Bermick, J. R. *et al.* Neonatal monocytes exhibit a unique histone modification
landscape. *Clinical epigenetics* **8**, 99 (2016). https://doi.org/10.1186/s13148-016-0265-7

25 Norouzitallab, P., Baruah, K., Biswas, P., Vanrompay, D. & Bossier, P. Probing the
phenomenon of trained immunity in invertebrates during a transgenerational study, using
brine shrimp Artemia as a model system. *Sci Rep* **6**, 21166 (2016).
https://doi.org/10.1038/srep21166

26 Netea, M. G. *et al.* Trained immunity: A program of innate immune memory in health and
disease. *Science (New York, N.Y.)* **352**, aaf1098 (2016).
https://doi.org/10.1126/science.aaf1098

27   Fitzgerald, M. Developmental biology of inflammatory pain. *British journal of anaesthesia*
75, 177-185 (1995). https://doi.org/10.1093/bja/75.2.177

28   Bracci-Laudiero, L. & De Stefano, M. E. in *Neurotoxin Modeling of Brain Disorders—Life-*
long Outcomes in Behavioral Teratology   (eds Richard M. Kostrzewa & Trevor Archer)
125-152 (Springer International Publishing, 2016).

29   Kobayashi, H. & Mizisin, A. P. Nerve growth factor and neurotrophin-3 promote
chemotaxis of mouse macrophages in vitro. *Neuroscience letters* **305**, 157-160 (2001).
https://doi.org/10.1016/s0304-3940(01)01854-7

30   Sajanti, A. *et al.* A comprehensive p75 neurotrophin receptor gene network and pathway
analyses identifying new target genes. *Scientific Reports* **10**, 14984 (2020).
https://doi.org/10.1038/s41598-020-72061-z

31   Chen, Z. *et al.* p75 Is Required for the Establishment of Postnatal Sensory Neuron
Diversity by Potentiating Ret Signaling. *Cell Rep* **21**, 707-720 (2017).
https://doi.org/10.1016/j.celrep.2017.09.037

32   Dourson, A. J. *et al.* Genetic and epigenetic mechanisms influencing acute to chronic
postsurgical pain transitions in pediatrics: Preclinical to clinical evidence. *Canadian*
Journal of Pain (2022). https://doi.org/10.1080/24740527.2021.2021799

33   Fortier, M. A., MacLaren, J. E., Martin, S. R., Perret-Karimi, D. & Kain, Z. N. Pediatric
pain after ambulatory surgery: where's the medication? *Pediatrics* **124**, e588-595 (2009).
https://doi.org/10.1542/peds.2008-3529

34   Rabbitts, J. A., Fisher, E., Rosenbloom, B. N. & Palermo, T. M. Prevalence and
Predictors of Chronic Postsurgical Pain in Children: A Systematic Review and Meta-
Analysis. *J Pain* **18**, 605-614 (2017). https://doi.org/10.1016/j.jpain.2017.03.007

35   Neville, A. *et al.* Diagnostic uncertainty in pediatric chronic pain: nature, prevalence, and
consequences. *Pain reports* **5**, e871 (2020).
https://doi.org/10.1097/pr9.0000000000000871

36   Angst, M. S. *et al.* Aversive and reinforcing opioid effects: a pharmacogenomic twin
study. *Anesthesiology* **117**, 22-37 (2012).
https://doi.org/10.1097/ALN.0b013e31825a2a4e

37   Martin, L., Jimenez, N. & Lynn, A. A review of perioperative anesthesia and analgesia for
infants: updates and trends to watch [version 1; referees: 2 approved]. *F1000Research* **6**
(2017). https://doi.org/10.12688/f1000research.10272.1

38   Dourson, A. J. & Jankowski, M. P. Developmental impact of peripheral injury on
neuroimmune signaling. *Brain Behav Immun* **113**, 156-165 (2023).
https://doi.org/10.1016/j.bbi.2023.07.002

39   Yu, X. *et al.* Dorsal root ganglion macrophages contribute to both the initiation and
persistence of neuropathic pain. *Nat Commun* **11**, 264 (2020).
https://doi.org/10.1038/s41467-019-13839-2

40   Shepherd, A. J. *et al.* Macrophage angiotensin II type 2 receptor triggers neuropathic
pain. *Proc Natl Acad Sci U S A* **115**, E8057-e8066 (2018).
https://doi.org/10.1073/pnas.1721815115

Kalymbetova, T. V. *et al.* Resident alveolar macrophages are master regulators of arrested alveolarization in experimental bronchopulmonary dysplasia. *The Journal of pathology* **245**, 153-159 (2018). https://doi.org/10.1002/path.5076

Burnett, S. H. *et al.* Conditional macrophage ablation in transgenic mice expressing a Fas-based suicide gene. *Journal of leukocyte biology* **75**, 612-623 (2004). https://doi.org/10.1189/jlb.0903442

Geissmann, F. *et al.* Development of monocytes, macrophages, and dendritic cells. *Science (New York, N.Y.)* **327**, 656-661 (2010). https://doi.org/10.1126/science.1178331

Lissner, M. M. *et al.* Age-Related Gene Expression Differences in Monocytes from Human Neonates, Young Adults, and Older Adults. *PloS one* **10**, e0132061 (2015). https://doi.org/10.1371/journal.pone.0132061

Millay, D. P. *et al.* Genetic and pharmacologic inhibition of mitochondrial-dependent necrosis attenuates muscular dystrophy. *Nature medicine* **14**, 442-447 (2008). https://doi.org/10.1038/nm1736

Jankowski, M. P., Rau, K. K., Ekmann, K. M., Anderson, C. E. & Koerber, H. R. Comprehensive phenotyping of group III and IV muscle afferents in mouse. *Journal of neurophysiology* **109**, 2374-2381 (2013). https://doi.org/10.1152/jn.01067.2012

Koerber, H. R. *et al.* Cutaneous C-polymodal fibers lacking TRPV1 are sensitized to heat following inflammation, but fail to drive heat hyperalgesia in the absence of TPV1 containing C-heat fibers. *Molecular pain* **6**, 58 (2010). https://doi.org/10.1186/1744-8069-6-58

Follis, R. M. *et al.* Metabolic Control of Sensory Neuron Survival by the p75 Neurotrophin Receptor in Schwann Cells. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **41**, 8710-8724 (2021). https://doi.org/10.1523/jneurosci.3243-20.2021

Subramanian Vignesh, K., Landero Figueroa, J. A., Porollo, A., Caruso, J. A. & Deepe, G. S., Jr. Granulocyte macrophage-colony stimulating factor induced Zn sequestration enhances macrophage superoxide and limits intracellular pathogen survival. *Immunity* **39**, 697-710 (2013). https://doi.org/10.1016/j.immuni.2013.09.006

May, M., Slaughter, A. & Lucas, D. Dynamic regulation of hematopoietic stem cells by bone marrow niches. *Curr Stem Cell Rep* **4**, 201-208 (2018). https://doi.org/10.1007/s40778-018-0132-x

Ojielo, C. I. *et al.* Defective phagocytosis and clearance of Pseudomonas aeruginosa in the lung following bone marrow transplantation. *Journal of immunology (Baltimore, Md. : 1950)* **171**, 4416-4424 (2003). https://doi.org/10.4049/jimmunol.171.8.4416

Spencer, S. J., Martin, S., Mouihate, A. & Pittman, Q. J. Early-life immune challenge: defining a critical window for effects on adult responses to immune challenge. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* **31**, 1910-1918 (2006). https://doi.org/10.1038/sj.npp.1301004

Reynolds, M. L. & Fitzgerald, M. Long-term sensory hyperinnervation following neonatal skin wounds. *The Journal of comparative neurology* **358**, 487-498 (1995). https://doi.org/10.1002/cne.903580403

54   Walker, S. M., Meredith-Middleton, J., Cooke-Yarborough, C. & Fitzgerald, M. Neonatal
inflammation and primary afferent terminal plasticity in the rat dorsal horn. *Pain* **105**,
185-195 (2003).

55   Walker, S. M. Long-term effects of neonatal pain. *Seminars in fetal & neonatal medicine*
24 (2019). https://doi.org/10.1016/j.siny.2019.04.005

56   Walker, S. M. Early life pain—effects in the adult. *Current Opinion in Physiology* **11**, 16-
24 (2019). https://doi.org/https://doi.org/10.1016/j.cophys.2019.04.011

57   Zhong, X. S. *et al.* Neonatal Colonic Inflammation Epigenetically Aggravates Epithelial
Inflammatory Responses to Injury in Adult Life. *Cellular and molecular gastroenterology*
and hepatology **6**, 65-78 (2018). https://doi.org/10.1016/j.jcmgh.2018.02.014

58   Chidambaran, V. *et al.* Enrichment of Genomic Pathways Based on Differential DNA
Methylation Associated With Chronic Postsurgical Pain and Anxiety in Children: A
Prospective, Pilot Study. *J Pain* **20**, 771-785 (2019).
https://doi.org/10.1016/j.jpain.2018.12.008

59   Brewer, C. L., Li, J., O'Conor, K., Serafin, E. K. & Baccei, M. L. Neonatal Injury Evokes
Persistent Deficits in Dynorphin Inhibitory Circuits within the Adult Mouse Superficial
Dorsal Horn. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience **40**, 3882-3895 (2020). https://doi.org/10.1523/jneurosci.0029-20.2020

60   Moriarty, O. *et al.* Priming of Adult Incision Response by Early-Life Injury: Neonatal
Microglial Inhibition Has Persistent But Sexually Dimorphic Effects in Adult Rats. *The*
Journal of Neuroscience **39**, 3081 (2019). https://doi.org/10.1523/JNEUROSCI.1786-
18.2019

61   van den Hoogen, N. J. *et al.* Repeated touch and needle-prick stimulation in the
neonatal period increases the baseline mechanical sensitivity and postinjury
hypersensitivity of adult spinal sensory neurons. *Pain* **159**, 1166-1175 (2018).
https://doi.org/10.1097/j.pain.0000000000001201

62   Schwaller, F., Beggs, S. & Walker, S. M. Targeting p38 Mitogen-activated Protein
Kinase to Reduce the Impact of Neonatal Microglial Priming on Incision-induced
Hyperalgesia in the Adult Rat. *Anesthesiology* **122**, 1377-1390 (2015).
https://doi.org/10.1097/aln.0000000000000659

63   Beggs, S., Currie, G., Salter, M. W., Fitzgerald, M. & Walker, S. M. Priming of adult pain
responses by neonatal pain experience: maintenance by central neuroimmune activity.
Brain **135**, 404-417 (2012). https://doi.org/10.1093/brain/awr288

64   Xu, Y., Moulding, D., Jin, W. & Beggs, S. Microglial phagocytosis mediates long-term
restructuring of spinal GABAergic circuits following early life injury. *Brain, Behavior, and*
Immunity **111**, 127-137 (2023). https://doi.org/https://doi.org/10.1016/j.bbi.2023.04.001

65   Ford, Z. K. *et al.* Systemic growth hormone deficiency causes mechanical and thermal
hypersensitivity during early postnatal development. *IBRO Reports* **6**, 111-121 (2019).
https://doi.org/https://doi.org/10.1016/j.ibror.2019.02.001

66   Kato, J. & Svensson, C. I. in *Progress in Molecular Biology and Translational Science*
Vol. 131 (eds Theodore J. Price & Gregory Dussor) 251-279 (Academic Press, 2015).

67   Comish, P. B., Carlson, D., Kang, R. & Tang, D. Damage-Associated Molecular Patterns
and the Systemic Immune Consequences of Severe Thermal Injury. *The Journal of*
Immunology **205**, 1189 (2020). https://doi.org/10.4049/jimmunol.2000439

68   McMahon, S. B., Cafferty, W. B. & Marchand, F. Immune and glial cell factors as pain
mediators and modulators. *Experimental neurology* **192**, 444-462 (2005).
https://doi.org/10.1016/j.expneurol.2004.11.001

69   Basbaum, A. I., Bautista, D. M., Scherrer, G. & Julius, D. Cellular and molecular
mechanisms of pain. *Cell* **139**, 267-284 (2009). https://doi.org/10.1016/j.cell.2009.09.028

70   Ren, K. & Dubner, R. Interactions between the immune and nervous systems in pain.
Nature medicine **16**, 1267-1276 (2010). https://doi.org/10.1038/nm.2234

71   Philippou, A., Maridaki, M., Theos, A. & Koutsilieris, M. Cytokines in muscle damage.
Advances in clinical chemistry **58**, 49-87 (2012). https://doi.org/10.1016/b978-0-12-
394383-5.00010-2

72   Campero, M., Bostock, H., Baumann, T. K. & Ochoa, J. L. Activity-dependent slowing
properties of an unmyelinated low threshold mechanoreceptor in human hairy skin.
Neuroscience letters **493**, 92-96 (2011). https://doi.org/10.1016/j.neulet.2011.02.012

73   Abdo, H. *et al.* Specialized cutaneous Schwann cells initiate pain sensation. *Science*
(New York, N.Y.) **365**, 695-699 (2019). https://doi.org/doi:10.1126/science.aax6452

74   McKelvey, R., Berta, T., Old, E., Ji, R.-R. & Fitzgerald, M. Neuropathic Pain Is
Constitutively Suppressed in Early Life by Anti-Inflammatory Neuroimmune Regulation.
The Journal of Neuroscience **35**, 457-466 (2015). https://doi.org/10.1523/jneurosci.2315-
14.2015

75   Wallner, S. *et al.* Epigenetic dynamics of monocyte-to-macrophage differentiation.
Epigenetics & chromatin **9**, 33 (2016). https://doi.org/10.1186/s13072-016-0079-z

76   Foster, S. L., Hargreaves, D. C. & Medzhitov, R. Gene-specific control of inflammation
by TLR-induced chromatin modifications. *Nature* **447**, 972-978 (2007).
https://doi.org/10.1038/nature05836

77   Lavin, Y. *et al.* Tissue-resident macrophage enhancer landscapes are shaped by the
local microenvironment. *Cell* **159**, 1312-1326 (2014).
https://doi.org/10.1016/j.cell.2014.11.018

78   Fanucchi, S., Domínguez-Andrés, J., Joosten, L. A. B., Netea, M. G. & Mhlanga, M. M.
The Intersection of Epigenetics and Metabolism in Trained Immunity. *Immunity* **54**, 32-43
(2021). https://doi.org/https://doi.org/10.1016/j.immuni.2020.10.011

79   Cobo, M. M. *et al.* Early life inflammation is associated with spinal cord excitability and
nociceptive sensitivity in human infants. *Nature communications* **13**, 3943 (2022).
https://doi.org/10.1038/s41467-022-31505-y

80   Pinho-Ribeiro, F. A. *et al.* Blocking Neuronal Signaling to Immune Cells Treats
Streptococcal Invasive Infection. *Cell* **173**, 1083-1097.e1022 (2018).
https://doi.org/10.1016/j.cell.2018.04.006

81   Perner, C. *et al.* Substance P Release by Sensory Neurons Triggers Dendritic Cell
Migration and Initiates the Type-2 Immune Response to Allergens. *Immunity* **53**, 1063-
1077.e1067 (2020). https://doi.org/https://doi.org/10.1016/j.immuni.2020.10.001

Filtjens, J. *et al.* Nociceptive sensory neurons promote CD8 T cell responses to HSV-1 infection. *Nat Commun* **12**, 2936 (2021). https://doi.org/10.1038/s41467-021-22841-6

Mendell, L. M. Neurotrophins and sensory neurons: role in development, maintenance and injury. A thematic summary. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **351**, 463-467 (1996). https://doi.org/10.1098/rstb.1996.0043

Lewin, G. R., Ritter, A. M. & Mendell, L. M. On the role of nerve growth factor in the development of myelinated nociceptors. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **12**, 1896-1905 (1992).

Sharma, N. *et al.* The emergence of transcriptional identity in somatosensory neurons. *Nature* **577**, 392-398 (2020). https://doi.org/10.1038/s41586-019-1900-1

Luo, W. *et al.* A hierarchical NGF signaling cascade controls Ret-dependent and Ret-independent events during development of nonpeptidergic DRG neurons. *Neuron* **54**, 739-754 (2007). https://doi.org/10.1016/j.neuron.2007.04.027

Susaki, Y. *et al.* Functional properties of murine macrophages promoted by nerve growth factor. *Blood* **88**, 4630-4637 (1996).

Williams, K. S., Killebrew, D. A., Clary, G. P., Seawell, J. A. & Meeker, R. B. Differential regulation of macrophage phenotype by mature and pro-nerve growth factor. *Journal of neuroimmunology* **285**, 76-93 (2015). https://doi.org/10.1016/j.jneuroim.2015.05.016

Khodorova, A., Nicol, G. D. & Strichartz, G. The p75NTR signaling cascade mediates mechanical hyperalgesia induced by nerve growth factor injected into the rat hind paw. *Neuroscience* **254**, 312-323 (2013). https://doi.org/10.1016/j.neuroscience.2013.09.046

Leitão, L. *et al.* Bone marrow cell response after injury and during early stage of regeneration is independent of the tissue-of-injury in 2 injury models. *The FASEB Journal* **33**, 857-872 (2019). https://doi.org/https://doi.org/10.1096/fj.201800610RR

Glatman Zaretsky, A., Engiles, J. B. & Hunter, C. A. Infection-induced changes in hematopoiesis. *Journal of immunology (Baltimore, Md. : 1950)* **192**, 27-33 (2014). https://doi.org/10.4049/jimmunol.1302061

Leitão, L. *et al.* The alliance between nerve fibers and stem cell populations in bone marrow: life partners in sickness and health. *The FASEB Journal* **33**, 8697-8710 (2019). https://doi.org/https://doi.org/10.1096/fj.201900454R

Cai, R. *et al.* Panoptic imaging of transparent mice reveals whole-body neuronal projections and skull–meninges connections. *Nature Neuroscience* **22**, 317-327 (2019). https://doi.org/10.1038/s41593-018-0301-3

Mazzitelli, J. A. *et al.* Cerebrospinal fluid regulates skull bone marrow niches via direct access through dural channels. *Nature Neuroscience* **25**, 555-560 (2022). https://doi.org/10.1038/s41593-022-01029-1

Queme, L. F. *et al.* Disruption of Hyaluronic Acid in Skeletal Muscle Induces Decreased Voluntary Activity via Chemosensitive Muscle Afferent Sensitization in Male Mice. *eNeuro* **9** (2022). https://doi.org/10.1523/eneuro.0522-21.2022

96   Walker, S. M., Tochiki, K. K. & Fitzgerald, M. Hindpaw incision in early life increases the
hyperalgesic response to repeat surgical injury: critical period and dependence on initial
afferent activity. *Pain* **147**, 99-106 (2009). https://doi.org/10.1016/j.pain.2009.08.017

97   Sorge, R. E. *et al.* Different immune cells mediate mechanical pain hypersensitivity in
male and female mice. *Nat Neurosci* **18**, 1081-1083 (2015).
https://doi.org/10.1038/nn.4053

98   dos Santos, N. L. *et al.* Age and Sex Drive Differential Behavioral and Neuroimmune
Phenotypes During Postoperative Pain. *Neurobiology of Aging* (2022).
https://doi.org/https://doi.org/10.1016/j.neurobiolaging.2022.09.008

99   Mogil, J. S. Qualitative sex differences in pain processing: emerging evidence of a
biased literature. *Nature Reviews Neuroscience* **21**, 353-365 (2020).
https://doi.org/10.1038/s41583-020-0310-6

100   Peters, J. W. *et al.* Does neonatal surgery lead to increased pain sensitivity in later
childhood? *Pain* **114**, 444-454 (2005). https://doi.org/10.1016/j.pain.2005.01.014

101   Williams, M. D. & Lascelles, B. D. X. Early Neonatal Pain-A Review of Clinical and
Experimental Implications on Painful Conditions Later in Life. *Front Pediatr* **8**, 30 (2020).
https://doi.org/10.3389/fped.2020.00030

102   Malik, S. C. *et al.* In vivo functions of p75(NTR): challenges and opportunities for an
emerging therapeutic target. *Trends Pharmacol Sci* **42**, 772-788 (2021).
https://doi.org/10.1016/j.tips.2021.06.006

103   Norman, B. H. & McDermott, J. S. Targeting the Nerve Growth Factor (NGF) Pathway in
Drug Discovery. Potential Applications to New Therapies for Chronic Pain. *J Med Chem*
60, 66-88 (2017). https://doi.org/10.1021/acs.jmedchem.6b00964

104   Zhao, Z., Ukidve, A., Kim, J. & Mitragotri, S. Targeting Strategies for Tissue-Specific
Drug Delivery. *Cell* **181**, 151-167 (2020).
https://doi.org/https://doi.org/10.1016/j.cell.2020.02.001

105   Burnett, S. H. *et al.* Development of peritoneal adhesions in macrophage depleted mice.
The Journal of surgical research **131**, 296-301 (2006).
https://doi.org/10.1016/j.jss.2005.08.026

106   Queme, L. F., Weyler, A. A., Cohen, E. R., Hudgins, R. C. & Jankowski, M. P. A dual
role for peripheral GDNF signaling in nociception and cardiovascular reflexes in the
mouse. *Proc Natl Acad Sci U S A* **117**, 698-707 (2020).
https://doi.org/10.1073/pnas.1910905116

1247

Figure 1

B.

MΦ
Dystr.

n.inc+Vehicle@P8   n.inc+AP@P8   n.inc+AP+P35 re-inc.@P36

C.

Mean macrophage numbers per section

***

n.inc+Veh   n.inc+AP   n.inc+AP+P35 re-inc

D. Paw Guarding

Guarding Score

*   *

Baseline   Day 1

E. Mechanical Thresholds

Withdrawal Thresholds (grams)

*   *^

Baseline   Day 1

F. Paw Guarding

Guarding Score

*

^#   ^#   ^#

Baseline Day 1 Day 3 Day 7 Day 14 Day 21

G. Mechanical Thresholds

Withdrawal Thresholds (grams)

*^#   ^#   ^#

*

Baseline Day 1 Day 3 Day 7 Day 14 Day 21

—●— n.inc + vehicle   —■— n.inc + AP   —▲— n.sham + vehicle   —▼— n.sham + AP

Figure 2

Figure 3

Figure 4

Figure 5

A.

P7:
-n.sham
-n.inc

Time

NGFR/p75

Groups:
-p75 fl/fl (no LysMcreRT2)
-LysM;p75 fl/fl

P35:
-inc
-Behavior

P33-37
-Tamoxifen

B.

F4/80
p75

C.

p75+ Macrophages

Percent f4/80+ and p75+

Controls   LysM;   LysM;
            p75+/fl p75fl/fl

D. **Paw Guarding**

Guarding Score

No Tam BL Day 1 Day 3 Day 7 Day 14 Day 21

*   #

E. **Mechanical Thresholds**

Withdrawal Thresholds (grams)

No Tam BL Day 1 Day 3 Day 7 Day 14 Day 21

^
*

p75fl/fl   LysM;p75fl/fl

Figure 6

A.

sip75+Vehicle   sip75+LPS   sip75+NGF

Pos. Con.   TNFα   IL1ra   TNFα   IL1ra   TNFα   IL1ra

B.

BCA1
C5a
GCSF
GMCSF (in bath)
I309
Eotaxin
ICAM1
IFNγ
IL1α
IL1β
IL1ra
IL2
IL3
IL4
IL5
IL6
IL7
IL10
IL13
IL12
IL16
IL17
IL23
IL27
IP10
ITAC
CXCL1
MCSF
MCP1
MCP5
MIG
MIP1α
MIP1β
MIP2
RANTES
SDF1
TARC
TIMP1
TNFα
TREM1
NEG

siCON Vehicle   sip75 Vehicle   sip75 LPS+IFNγ   sip75 NGF

-1   0   1

C.

% Change from siCON+Vehicle			
Protein	sip75+Veh	sip75+LPS	sip75+NGF
TNFα	572 ± 469%	8392 ± 5043%*	24 ± 54%^
IL1β	253 ± 153%	322 ± 154%	34 ± 39%#
IL1α	-68 ± 14%	24 ± 88%	-78 ± 16%
IL6	1036 ± 920%	19944 ± 9191%*	31 ± 44%
IL1ra	-31 ± 36%	4104 ± 3619%	193 ± 189%
IFNγ	76 ± 10%*	177 ± 39%*	14 ± 16%
IL4	140 ± 133%	35 ± 14%	98 ± 33%*
IL10	91 ± 30%*	113 ± 38%	57 ± 33%##

D. hMacrophage

E. hNeuron

F. ATP Responses

Pre-stim   Post-stim

siCON+NGF

sip75+NGF

G.

siCON+NGF (ATP Response)   KCl Added

ATP Added

siCON+NGF (CAP Response)   KCl Added

CAP Added

sip75+NGF (ATP Response)   KCl Added

ATP Added

sip75+NGF (CAP Response)   KCl Added

CAP Added

H. ATP Responses

dF/F

siCON+Veh   sip75+Veh   siCON+NGF   sip75+NGF

I. Capsaicin Responses

dF/F

siCON+Veh   sip75+Veh   siCON+NGF   sip75+NGF

Figure 7

Graphical Abstract