1    **Title: Selection on many loci drove the origin and spread of a key innovation**
2

3    **Authors:** Sean Stankowski[1,2]*, Zuzanna B. Zagrodzka[1], Martin D. Garlovsky[3], Arka Pal[2], Daria
4    Shipilina[2,4], Diego Garcia Castillo[2], Alan Le Moan[5,6], Erica Leder[6], James Reeve[6], Kerstin
5    Johannesson[6], Anja M. Westram[2,7], Roger K. Butlin[1,6].
6
**Affiliations:**
[1]1Ecology and Evolutionary Biology, School of Biosciences, University of Sheffield,
Sheffield, UK
[2] Institute of Science and Technology Austria (ISTA), Klosterneuburg, Austria
[3]Department of Applied Zoology, Faculty of Biology, Technische Universität Dresden, 01069
Dresden, Germany
[4]Department of Ecology and Genetics, Program of Evolutionary Biology, Uppsala University,
Uppsala, Sweden
[5]CNRS & Sorbonne Université, Station Biologique de Roscoff, Roscoff, France
[6] Department of Marine Sciences, Tjärnö, University of Gothenburg, Strömstad, Sweden
[7] Faculty of Biosciences and Aquaculture, Nord University, N-8049 Bodø, Norway
*Corresponding author: sean.stankowski@ista.ac.at

**Abstract: Key innovations are fundamental to biological diversification, but their genetic**
**architecture is poorly understood. A recent transition from egg-laying to live-bearing in**
***Littorina* snails provides the opportunity to study the architecture of a young innovation.**
**Samples do not cluster by reproductive mode in a genome-wide phylogeny, but local**
**genealogical analysis revealed numerous genomic regions where all live-bearers carry the**
**same core haplotype. Associated regions show evidence for live-bearer-specific positive**
**selection, and are enriched for genes that are differentially expressed between egg-laying**
**and live-bearing reproductive systems. Ages of selective sweeps suggest live-bearing alleles**
**accumulated gradually, involving selection at different times in the past. Our results**
**suggest that innovation can have a polygenic basis, and that novel functions can evolve**
**gradually, rather than in a single step.**

Main text: Evolution is a gradual process, but occasionally results in sudden changes in
form and function that allow organisms to exploit new ecological opportunities (*1, 2*). These
game-changing traits—including flight, vision, and the bearing of live offspring—are known as
'key innovations'. Key innovations are all around us, and have catalyzed the diversification of
many groups (*1*). Despite their significance, we know surprisingly little about the origins and
genetic architecture of most innovations (*1*). This is because most originated deep in the past,
making it difficult to disentangle causal loci from the countless genetic changes that accumulated
up to the present.
A recent transition in female reproductive mode offers a rare opportunity to study the
genetic architecture of a young innovation (*3*). We focus on a clade of intertidal gastropods
(Genus *Littorina*), where the ancestral state is to lay a large egg-mass but one species gives birth
to live young (Fig. 1A, fig. S1) (*4, 5*). Egg-layers have a gland that embeds egg-capsules into a
protective jelly. In the live-bearer, *L. saxatilis*, this structure has evolved into a brood pouch
where embryos develop inside the mother. Live-bearing is a recent innovation in the littorinidae
and is thought to allow snails to reproduce in areas where eggs are exposed to harsh conditions
(*5*). This is reflected in the much broader ecological and geographic distribution of *L. saxatilis*

compared with the two egg-laying sister species, *L. arcana and L. compressa* (*5*) (Fig. 1B and
1C). Egg-laying and live-bearing species have adapted in parallel to contrasting environments (*5*,
*6*), partly decoupling reproductive mode from other axes of phenotypic divergence (Fig. 1B).
There is also evidence for occasional hybridization between egg-layers and live-bearers (*7, 8*).
These features provide an opportunity to identify and study the genetic changes underlying the
live-bearing innovation.

**Figure 1. Variation in reproductive mode in *Littorina*.** (A) Anatomical differences between modes (B) Egg-
layers reproduce during a limited breeding season, while live-bearers release offspring year-round. The two
egg-layers share their habitats with ecotypes of the live-bearer, *L. saxatilis*. (C) Approximate distributions of the
modes, highlighting the broader distribution of live birth. (D) Existing hypothesis for the origin of live birth. (E)
ML phylogenetic tree based on whole-genome sequences (108 individuals and 18.5 million variable sites).
Bootstrap support for key nodes is shown.
59

***Live-bearing snails do not form a monophyletic group***
We used whole-genome sequences from 108 individuals to test the existing hypothesis of a
single origin of live-bearing (Fig. 1D ) (*4*). Surprisingly, live-bearers formed two separate clades
in a phylogenetic tree: one containing all *L. saxatilis* from Spain (hereafter 'Spanish *saxatilis*'),
and another including all other *L. saxatilis* ('Northern *saxatilis*') that was sister to egg-laying *L.*
*arcana* (Fig. 1E). The discordance between evolutionary relationships and reproductive mode
(also seen in PCAs, fig. S6) has several possible explanations, including two genetically
independent transitions between egg-laying and-live birth. However, given the close
relationships of these taxa, a single origin could have been followed by the sharing of causal

alleles between lineages via gene exchange and selection (*9*). For example, live-bearing could
first have evolved in Spain, whereupon causal alleles spread to the north, introgressing into the
genetic background of the resident egg-laying lineage. In this case, we would expect genealogies
for loci causing live birth to be strongly discordant from the genome-wide tree, with samples
grouping by reproductive mode (*9*).
74
***Topology weighting reveals rampant genealogical discordance and loci associated with***
***reproductive mode***
With this expectation in mind, we used topology weighting (Fig. 2A) to identify genomic regions
associated with reproductive mode. For each genomic window, topology weighting calculates
the relative weight of the three possible subtrees (Fig. 2C, fig. S8): (*i*) the background topology,
Tb, observed in our genome-wide analysis, (ii) the reproduction topology, Tr, where samples
cluster by reproductive mode, and (iii) the control topology, Tc, which is of no specific interest
except that it provides a control for distinguishing incomplete lineage sorting (ILS) from other
processes that cause genealogical discordance. We used non-overlapping 100-SNP windows
(mean size 5.8 kb, fig. S7), and calculated topology weights (10) for each window by sampling
10,000 subtrees (Fig. 2A). We took the novel approach of analyzing the joint distribution of
topology weights in a ternary plot (Fig. 2A) and used simulations to understand how different
factors shape the ternary distribution of weights (Fig. 2B; Supplementary text, figs. S9—S19;
tables S3 & S4).

**Figure 2. Topology weighting reveals genomic regions associated with reproductive mode.** (A) For each
genomic window, we inferred a full tree including all haplotypes, and then sampled and classified 10k 'subtrees'
by randomly picking one haplotype per group. Topology weights are the proportions of each topology among all
subtrees. Windows were then plotted in a ternary plot based on their topology weights. (B) Simulated

distributions of weights. A greater opportunity for lineage sorting (i - iii) biases the distribution toward the
topology that matches the demographic history. Incomplete lineage sorting yields genealogies that are a better
fit to one of the discordant trees, but the distribution is always symmetrical between the left and right half
triangles. Additional factors, including gene flow, create a bias toward one of the discordant genealogies
(panels iv - vi). (C) Possible topologies and the empirical distribution of weights for the 154,971, 100 SNP
windows. Hexagonal bins are colored according to window count. (D) Counts of windows in the left and right
half triangles, with the asymmetry quantified using $D_{LR}$. Further division into sub-triangles reveals left-right
asymmetry throughout the distribution. Asterisks indicate significant asymmetry between corresponding left-
and right-sided sub-triangles. (E) Distributions of weights > 0.7.

We expected the empirical distribution of weights to be biased toward Tb, because lineage sorting results in concordance between the demographic history and underlying gene trees (*10*) (Fig. 2B, Supplementary text). However, the observed bias was only slight (Tb = 0.380, Tc = 0.310, Tr = 0.308), with just 62 of ~155,000 genomic regions perfectly fitting Tb (*i.e.*, Tb = 1) (Fig. 2C). Instead, the bulk of the distribution fell close to the center of the triangle, revealing extensive ILS due to rapid diversification relative to the effective population size (*10*, *11*). Thus, although well-supported statistically, the genome-wide tree is a very poor predictor of evolutionary relationships at any given genomic region.

We found substantial left-right asymmetry in the distribution of weights (Fig. 2D). Such a bias is not expected to arise from ILS, because there is an equal chance that a given gene tree will more closely resemble either alternative topology (Fig. 2B, supplementary text) (*10*). We detected asymmetry using a new statistic, $D_{LR}$ (Fig. 2D, fig. S19). A genome-wide test, performed by calculating $D_{LR}$ between the two halves of the triangle, revealed a 3.4% excess of windows shifted toward the control topology ($D_{LR}$ = 0.034, permutation test p = 1e-5). $D_{LR}$ calculated between analogous left- and right-side sub-triangles, revealed that this asymmetry was driven by an excess of trees with a small bias toward Tc (Fig. 2D, table S5). Further exploration showed that this bias is due to several previously identified chromosomal inversions, where one arrangement is more common in Spanish *L. saxatilis* and *L. arcana*, and the other is more common in *L. compressa* and Northern *L. saxatilis* ($D_{LR}$ for regions outside inversions = -0.007, p = 0.074) (figs. S21—S24, table S6, Supplementary text).

Much stronger asymmetry was observed between the extreme left and right sub-triangles, corresponding to windows that strongly fit one of the alternative topologies (Fig. 2D). However, the asymmetry was in the opposite direction to the genome-wide pattern, with a large excess of windows strongly biased toward the reproduction tree compared with the control tree (Tr > 0.7 = 1151 windows vs. 461 for Tc; $D_{LR}$ = -0.43, p = 1e-5). A total of 88 windows perfectly fit the reproduction topology (*i.e.*, Tr = 1), compared with 0 windows that perfectly fit the control topology ($D_{LR}$ = 1.00, p = 1e-5; Fig. 2E, table S5).

### *Evidence for live-bearer specific positive selection*

Although neutral gene flow can generate strong asymmetry under some circumstances, we are unable to explain the observed Tr bias without invoking natural selection (supplementary text). We found strong additional evidence for live-bearer-specific positive selection in these regions. First, window-based estimates of nucleotide diversity ($\pi$) in live-bearers decreased substantially with increasing Tr weight (Fig. 3A). We found no such relationship in egg-layers. Eighty-four of the 88 (95%) perfectly associated regions showed reduced $\pi$ in live-bearers (mean $\pi_{\text{live-bearer}}$ = 0.0029 vs $\pi_{\text{Egg-layer}}$ = 0.0065; paired Wilcoxon test, p = 1.313e-15, Fig. 3A, fig. S25), consistent with selection having purged diversity from live-bearing haplotypes (*12*). Although this pattern could in principle result from a live-bearer-specific demographic bottleneck, we can rule this out

because live-bearers and egg-layers have similar levels of genome-wide diversity (mean π live-bearer = 0.0065 vs. π egg-layer = 0.0062; fig. S26). Further, relationships between π and the other weights (Ts and Tc) were weak, and similar for both groups, confirming that reduced π in live-bearers is specific to Tr rather than being a general feature of windows with extreme weights (fig. S27). The site-frequency spectra (SFS) and sample-size-corrected estimates of private alleles for perfectly associated regions provide further evidence for selection (Fig 3B & C; figs. S28—S30; tables S9 and S10): the live-bearer SFS was strongly skewed toward rare variants (Tajima's D = -1.89, 95% CIs -1.77 – -2.01; fig. S29), the majority of which (80%) were private to the group. Both results are expected during the phase when diversity is recovered by mutation after a selective sweep (*13*). In contrast, the SFS for egg-layers was much closer to the neutral expectation (Tajima's D = -0.24, 95% CIs -0.037 – -0.437), with polymorphic sites being 2.14 times more abundant in egg-layers after accounting for the difference in sample size.

**Figure 3. Evidence for positive selection on haplotypes associated with live birth.** (A) Relationship between π and Tr for both reproductive modes. Triangles on right: genome-wide π. Violin plots: distributions of π for windows where Tr = 1. Letters show mean values of π for egg-layers and live-bearers. (B) Folded SFS for each mode in perfectly associated regions, projected at the same sample size for comparison. (C) Estimates of

Tajima's D with 95% CIs for perfectly associated regions. (D) Unrooted trees for example windows where Tr = 1. (E) Variation across two example contigs that contain a window where Tr = 1 (span of the orange box). The tree associated with each region is shown. Top panel: $F_{ST}$ between egg-layers and live-bearers in 3kb sliding windows (30 bp step). TrARG shows the results of topology weighting applied to marginal trees obtained from inferred ancestral recombination graphs (ARGs). Purple arrows show fixed differences between modes. Middle panel: π and $d_{xy}$ in sliding windows. Bottom panel: traces of time to the most recent common ancestor (TMRCA) obtained from ARGs. Bold lines: median estimates; Envelopes: 95% CIs. The red box shows the inferred length of the core haplotype block associated with live birth.

We characterized footprints of selection within contigs to more accurately estimate the number and size of candidate regions (Fig. 3E). The 88 perfectly associated windows mapped to 50 contigs in our genome assembly (mean 1.7 ± sd 1.5 windows per contig; table S8). Associated regions were narrow, mostly spanning less than 20 kb (mean 12 kb ± sd 14.4 kb). Sliding-window analysis of each contig generally revealed clear peaks of allele frequency differentiation ($F_{ST}$) and sequence divergence ($d_{xy}$) between the groups, as well as valleys of nucleotide diversity (π) in live-bearers (Fig. 3E; fig. S33). We also inferred ancestral recombination graphs (ARGs) for selected contigs to refine candidate regions (Fig. 3E). Unlike the trees for windows of arbitrary size and position, each marginal tree in an ARG corresponds to an inferred non-recombining segment of the genome (*14*). Thus, by applying topology weighting to the sequence of marginal trees, we were able to identify more precisely the segment of genome retained by all live bearing samples following the sweep. In both cases, the core live-bearing haplotype spanned less than 2 kb. Live-bearers showed much shallower coalescence in these regions than egg-layers, as expected following a selective sweep (Fig. 3E).

***Mode-associated regions are widespread and enriched for genes that are differentially expressed between reproductive systems***

The assignment of contigs to a genetic map revealed that reproductive-mode-associated windows are widespread across the genome, rather than co-localizing to one or a few genomic regions (Fig. 4a; table S11). As expected for a polygenic trait, the number of mode-associated windows on each LG was strongly predicted by LG size (Tr > 0.7, r = 0.79, p < 0.0001; Tr > 0.9, r = 0.71, p < 0.005). Associated windows were also widespread within linkage groups, in some cases with strong associations near opposite ends of the same LG (Fig. 4B).

Candidate regions also showed strong enrichment of genes that are differentially expressed between live-bearing and egg-laying reproductive tissues. To identify differentially expressed genes (DEGs), we collected reproductively mature female *L. arcana* and Northern *L. saxatilis* from a single location to control for environmental effects, and compared transcriptomes from whole reproductive systems (brood pouch vs jelly gland) and a non-reproductive control tissue (foot). We identified 1,598 DEGs, the majority of which showed differential expression between the reproductive tissues (1,297) (Fig. 4C, fig. S36). Of these, 66.1% (858) showed higher expression in the brood pouch of live bearers (Fig. 4D). To test for the enrichment of DEGs in regions associated with reproductive mode, we binned each DEG according to the Tr score of its associated genomic region (Fig. 4E). We found that the proportion of reproductive mode DEGs strongly increased with increasing Tr weight (Spearman's rho = 0.903, p = 9e-04; table S12)

Gene ontology analysis and functional annotation suggest that the transition to live-birth involved genes with diverse functions. Separate GO analyses conducted on a sequence-based gene set (574 genes in regions where Tr > 0.7) and expression-based gene set (1,450 reproductive mode DEGs) yielded 37 enriched gene ontology terms, including transmembrane

transport, calcium ion binding, and ion channel activity (Fig. S37). We examined the putative
functions of the 27 genes found in both sets in more detail (table S13). These included genes
putatively associated with antibacterial activity (lectin L6-like protein; higher expression in
brood pouch), the synthesis of mucin-type oligosaccharides (GALNT10-like; higher expression
in brood pouch), the formation of structural tissue (IFB-like and CMP-like, both higher
expression in brood pouch), and two secretary genes involved in egg-mass production in another
marine snail (both with lower expression in brood pouch).
212

**Figure 4. Candidate regions are widespread across the genome and enriched for genes that are
differentially expressed between reproductive systems.** (A) The number of high Tr windows (Tr > 0.7)
assigned to each of the 17 *L. saxatilis* LGs. The circles show the expected number given the total assigned of
windows to each LG. Asterisks indicate when the observed number is unlikely to be recovered by chance (p <
0.05). (B) Distribution of high Tr windows across LGs. Vertical blue lines indicate map positions that are
enriched for high Tr windows. (C) Number of genes that showed differential expression (DE) and the number of
DE genes in each expression class. (D) Clustering of reproductive tissue libraries based on patterns of
expression. (E) The proportion of genes in each DE class after binning each gene according to the Tr weight.

*Conclusions*

Our analyses show that live-bearing, a key innovation, is associated with selection on many loci, as in the only comparable analysis in *Zootoca* lizards (*15*). Although our genome-wide analysis hinted at two independent origins of live-bearing, the alleles associated with this trait have a single origin and have spread across space and genetic background. We found evidence that selection has acted on differences in gene expression, driving the origin of the live-bearing brood pouch. Other loci associated loci may underpin physiological changes that contribute to the difference in mode, such as differences in embryo retention-time (*16*), or may underlie other adaptations the became beneficial as live-bearing spread. Approximate estimates of the timing of selective sweeps at live-bearing loci, based on the accumulation of private mutations ($T = \pi_w/2\mu$), span a broad range from ~20 k to 200 k generations before present, with a median time of 70 k generations BP (roughly 35k years BP, assuming 2 generations per year) (fig. S38). Thus, our results suggest that alleles associated with live-bearing accumulated gradually over the last 100 k years. These findings are relevant to the long-standing debate about the genetic basis of evolutionary novelty. Because key innovations are not visible to selection before they arise, models of saltational evolution invoke large-effect macromutations to explain their sudden appearance (*17*). However, our results show that innovation can have a polygenic basis, and suggest that novel functions can arise as the end product of sustained quantitative evolution, rather than in a single evolutionary step.

**Acknowledgments:** Juan Galindo, Mauricio Montaño-Rendón, Natalia Mikhailova, April Blakeslee, Einar Arnason and Petri Kemppainen provided samples. Richard Turney, Graciela Sotelo, Jenny Larsson, Thomas Broquet and Stéphane Loisel helped with sample collection and processing. We thank Sci Ani for providing and allowing us to modify the cartoons shown in Fig. 1. Mark Dunning helped develop bioinformatic pipelines. Rui Faria, Hernan Morales, Vitor Sousa, Joost Raeymaekers, Aaron Comeault and Nick Barton provided advice and discussion.

**Funding:**
National Environment Research Council grant NE/P001610/1 (RKB)
European Research Council grant ERC-2015-AdG693030-BARRIERS (RKB)

**Author contributions:**
Conceptualization: SS, AMW, KJ, RKB
Analysis: SS, ZBZ, MG, AP, DS, DGC, EL, JR, ALM
Writing – original draft: SS, RKB
Writing – review & editing: SS, ZBZ, MG, AP, DS, DGC, TB, ALM, EL, JR, KJ, AMW, RKB

**Competing interests:**
Authors declare that they have no competing interests.

**Data and materials availability:**
Sequence data are available on the short-read archive (SRA). All other data and analysis scripts are available on Github at https://github.com/seanstankowski/Littorina_reproductive_mode.

**List of supplementary materials:**
Materials and Methods

**References and notes**

1.    A. Wagner, *The Origins of Evolutionary Innovations: A Theory of Transformative Change*
*in Living Systems* (OUP Oxford, 2011).
2.    A. H. Miller, J. T. Stroud, J. B. Losos, The ecology and evolution of key innovations.
*Trends Ecol. Evol.* (2022), doi:10.1016/j.tree.2022.09.005.
3.    C. M. Whittington, J. U. Van Dyke, S. Q. T. Liang, S. V. Edwards, R. Shine, M. B.
Thompson, C. E. Grueber, Understanding the evolution of viviparity using intraspecific
variation in reproductive mode and transitional forms of pregnancy. *Biol. Rev. Camb.*
*Philos. Soc.* **97**, 1179–1192 (2022).
4.    D. G. Reid, P. Dyal, S. T. Williams, A global molecular phylogeny of 147 periwinkle
species (Gastropoda, Littorininae). *Zool. Scr.* **41**, 125–136 (2012).
5.    D. G. Reid, Systematics and evolution of Littorina. The Ray Society, London, 463 p (1996).
6.    K. Johannesson, Evolution in Littorina: ecology matters. *J. Sea Res.* **49**, 107–117 (2003).
7.    S. Stankowski, A. M. Westram, Z. B. Zagrodzka, I. Eyres, T. Broquet, K. Johannesson, R.
K. Butlin, The evolution of strong reproductive isolation between sympatric intertidal
snails. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **375**, 20190545 (2020).
8.    T. Warwick, A. J. Knight, R. D. Ward, Hybridisation in the Littorina saxatilis species
complex (Prosobranchia : Mollusca). *Hydrobiologia*. **193**, 109–116 (1990).
9.    M. W. Hahn, L. Nakhleh, Irrational exuberance for resolved species trees. *Evolution*. **70**, 7–
17 (2016).
10.   W. P. Maddison, Gene Trees in Species Trees. *Syst. Biol.* **46**, 523–536 (1997).
11.   R. R. Hudson, Gene genealogies and the coalescent process. *Oxford surveys in evolutionary*
*biology*. **7**, 44 (1990).
12.   J. M. Maynard Smith, J. Haigh, The hitch-hiking effect of a favourable gene. *Genet. Res.*
**23**, 23–35 (1974).
13.   J. M. Braverman, R. R. Hudson, N. L. Kaplan, C. H. Langley, W. Stephan, The hitchhiking
effect on the site frequency spectrum of DNA polymorphisms. *Genetics*. **140**, 783–796
(1995).
14.   D. Shipilina, A. Pal, S. Stankowski, Y. F. Chan, N. H. Barton, On the origin and structure of
haplotype blocks. *Mol. Ecol.* (2022), doi:10.1111/mec.16793.
15.   H. Recknagel, M. Carruthers, A. A. Yurchenko, M. Nokhbatolfoghahai, N. A. Kamenos, M.
M. Bain, K. R. Elmer, The functional genetic architecture of egg-laying and live-bearing
reproduction in common lizards. *Nat Ecol Evol*. **5**, 1546–1556 (2021).
16.   R. Shine, Reptilian Reproductive Modes: The Oviparity-Viviparity Continuum.
*Herpetologica*. **39**, 1–8 (1983).
17.   G. Theißen, Saltational evolution: hopeful monsters are here to stay. *Theory Biosci.* **128**,
43–51 (2009).
***The following refrences are cited in the supplementary materials***
18.   K. Johannesson, Z. Zagrodzka, R. Faria, A. Marie Westram, R. K. Butlin, Is embryo
abortion a post-zygotic barrier to gene flow between Littorina ecotypes? *J. Evol. Biol.* **33**,
342–351 (2020).

19. R. K. Butlin, M. Saura, G. Charrier, B. Jackson, C. André, A. Caballero, J. A. Coyne, J.
Galindo, J. W. Grahame, J. Hollander, P. Kemppainen, M. Martínez-Fernández, M. Panova,
H. Quesada, K. Johannesson, E. Rolán-Alvarez, Parallel evolution of local adaptation and
reproductive isolation in the face of gene flow. *Evolution*. **68**, 935–949 (2014).
20. H. E. Morales, R. Faria, K. Johannesson, T. Larsson, M. Panova, A. M. Westram, R. K.
Butlin, Genomic architecture of parallel ecological divergence: Beyond a single
environmental contrast. *Sci Adv*. **5**, eaav9963 (2019).
21. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics*. **30**, 2114–2120 (2014).
22. A. M. Westram, M. Rafajlović, P. Chaube, R. Faria, T. Larsson, M. Panova, M. Ravinet, A.
Blomberg, B. Mehlig, K. Johannesson, R. Butlin, Clines on the seashore: The genomic
architecture underlying rapid divergence in the face of gene flow. *Evol Lett*. **2**, 297–309
(2018).
23. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
*arXiv [q-bio.GN]* (2013), (available at http://arxiv.org/abs/1303.3997).
24. G. Tischler, *biobambam2: Tools for early stage alignment file processing* (Github;
https://github.com/gt1/biobambam2).
25. A. McKenna, M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K.
Garimella, D. Altshuler, S. Gabriel, M. Daly, M. A. DePristo, The Genome Analysis
Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
*Genome Res*. **20**, 1297–1303 (2010).
26. P. Danecek, J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham,
T. Keane, S. A. McCarthy, R. M. Davies, H. Li, Twelve years of SAMtools and BCFtools.
*Gigascience*. **10** (2021), doi:10.1093/gigascience/giab008.
27. A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*. **30**, 1312–1313 (2014).
28. E. M. Ortiz, *vcf2phylip v2.0: convert a VCF matrix into several matrix formats for
phylogenetic analysis* (2019; https://zenodo.org/record/2540861).
29. A. Rambaut, FigTree. Tree Figure Drawing Tool. *http://tree.bio.ed.ac.uk/software/figtree/*
(2009) (available at https://ci.nii.ac.jp/naid/10029324212/).
30. C. C. Chang, C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell, J. J. Lee, Second-
generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience*. **4**, 7
(2015).
31. S. H. Martin, S. M. Van Belleghem, Exploring Evolutionary Relationships Across the
Genome Using Topology Weighting. *Genetics*. **206**, 429–438 (2017).
32. B. L. Browning, S. R. Browning, Improving the accuracy and efficiency of identity-by-
descent detection in population data. *Genetics*. **194**, 459–471 (2013).
33. S. Guindon, J.-F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk, O. Gascuel, New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321 (2010).
34. G. Yu, D. K. Smith, H. Zhu, Y. Guan, T. T.-Y. Lam, Ggtree : An r package for visualization
and annotation of phylogenetic trees with their covariates and other associated data.
*Methods Ecol. Evol.* **8**, 28–36 (2017).
35. S. H. Martin, K. K. Dasmahapatra, N. J. Nadeau, C. Salazar, J. R. Walters, F. Simpson, M.
Blaxter, A. Manica, J. Mallet, C. D. Jiggins, Genome-wide evidence for speciation with
gene flow in Heliconius butterflies. *Genome Res.* **23**, 1817–1828 (2013).

36. F. Baumdicker, G. Bisschop, D. Goldstein, G. Gower, A. P. Ragsdale, G. Tsambos, S. Zhu, B. Eldon, E. C. Ellerman, J. G. Galloway, A. L. Gladstein, G. Gorjanc, B. Guo, B. Jeffery, W. W. Kretzschumar, K. Lohse, M. Matschiner, D. Nelson, N. S. Pope, C. D. Quinto-Cortés, M. F. Rodrigues, K. Saunack, T. Sellinger, K. Thornton, H. van Kemenade, A. W. Wohns, Y. Wong, S. Gravel, A. D. Kern, J. Koskela, P. L. Ralph, J. Kelleher, Efficient ancestry and mutation simulation with msprime 1.0. *Genetics*. **220** (2022), doi:10.1093/genetics/iyab229.

37. J. Kelleher, K. Lohse, Coalescent Simulation with msprime. *Methods Mol. Biol.* **2090**, 191–230 (2020).

38. G. Gower, A. P. Ragsdale, G. Bisschop, R. N. Gutenkunst, M. Hartfield, E. Noskova, S. Schiffels, T. J. Struck, J. Kelleher, K. R. Thornton, Demes: a standard format for demographic models. *Genetics*. **222** (2022), doi:10.1093/genetics/iyac131.

39. J. Kelleher, K. R. Thornton, J. Ashander, P. L. Ralph, Efficient pedigree recording for fast population genetics simulation. *PLoS Comput. Biol.* **14**, e1006581 (2018).

40. S. H. Martin, J. W. Davey, C. D. Jiggins, Evaluating the Use of ABBA–BABA Statistics to Locate Introgressed Loci. *Mol. Biol. Evol.* **32**, 244–257 (2015).

41. R. E. Green, J. Krause, S. E. Ptak, A. W. Briggs, M. T. Ronan, J. F. Simons, L. Du, M. Egholm, J. M. Rothberg, M. Paunovic, S. Pääbo, Analysis of one million base pairs of Neanderthal DNA. *Nature*. **444**, 330–336 (2006).

42. J. Kelleher, Y. Wong, A. W. Wohns, C. Fadil, P. K. Albers, G. McVean, Inferring whole-genome histories in large population datasets. *Nat. Genet.* **51**, 1330–1338 (2019).

43. L. Speidel, M. Forest, S. Shi, S. R. Myers, A method for genome-wide genealogy estimation for thousands of samples. *Nat. Genet.* **51**, 1321–1329 (2019).

44. M. D. Rasmussen, M. J. Hubisz, I. Gronau, A. Siepel, Genome-wide inference of ancestral recombination graphs. *PLoS Genet.* **10**, e1004342 (2014).

45. N. E. Hamilton, M. Ferry, ggtern: Ternary diagrams using ggplot2. *J. Stat. Softw.* **87**, 1–17 (2018).

46. R. Faria, P. Chaube, H. E. Morales, T. Larsson, A. R. Lemmon, E. M. Lemmon, M. Rafajlović, M. Panova, M. Ravinet, K. Johannesson, A. M. Westram, R. K. Butlin, Multiple chromosomal rearrangements in a hybrid zone between Littorina saxatilis ecotypes. *Mol. Ecol.* **28**, 1375–1393 (2019).

47. A. M. Westram, R. Faria, K. Johannesson, R. Butlin, Using replicate hybrid zones to understand the genomic basis of adaptive divergence. *Mol. Ecol.* **30**, 3797–3814 (2021).

48. T. Jombart, adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*. **24**, 1403–1405 (2008).

49. P. Kemppainen, C. G. Knight, D. K. Sarma, T. Hlaing, A. Prakash, Y. N. Maung Maung, P. Somboon, J. Mahanta, C. Walton, Linkage disequilibrium network analysis (LDna) gives a global view of chromosomal inversions, local adaptation and geographic structure. *Mol. Ecol. Resour.* **15**, 1031–1045 (2015).

50. F. Tajima, Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*. **123**, 585–595 (1989).

51. P. Danecek, A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E. Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, R. Durbin, 1000 Genomes Project Analysis Group, The variant call format and VCFtools. *Bioinformatics*. **27**, 2156–2158 (2011).

52. S. Andrews, Others, FastQC: a quality control tool for high throughput sequence data
(2010).
53. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019).
54. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R.
Durbin, 1000 Genome Project Data Processing Subgroup, The Sequence Alignment/Map
format and SAMtools. *Bioinformatics*. **25**, 2078–2079 (2009).
55. M. Pertea, G. M. Pertea, C. M. Antonescu, T.-C. Chang, J. T. Mendell, S. L. Salzberg,
StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat.*
*Biotechnol.* **33**, 290–295 (2015).
56. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics*. **26**, 139–140
(2010).
57. S. X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and
plants. *Bioinformatics*. **36**, 2628–2629 (2020).
58. G. Pertea, M. Pertea, GFF Utilities: GffRead and GffCompare. *F1000Res*. **9** (2020),
doi:10.12688/f1000research.23297.2.
59. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics*. **26**, 841–842 (2010).
60. H. Tavares, A. Whibley, D. L. Field, D. Bradley, M. Couchman, L. Copsey, J. Elleouet, M.
Burrus, C. Andalo, M. Li, Q. Li, Y. Xue, A. B. Rebocho, N. H. Barton, E. Coen, Selection
and gene flow shape genomic islands that control floral guides. *Proc. Natl. Acad. Sci. U. S.*
*A*. **115**, 11006–11011 (2018).
61. M. Panova, A. M. H. Blakeslee, A. W. Miller, T. Mäkinen, G. M. Ruiz, K. Johannesson, C.
André, Glacial history of the North Atlantic marine snail, Littorina saxatilis, inferred from
distribution of mitochondrial DNA lineages. *PLoS One*. **6**, e17511 (2011).
62. K. Janson, Genetic and environmental effects on the growth rate of Littorina saxatilis. *Mar.*
*Biol.* **69**, 73–78 (1982).