Intracellular replication of *Pseudomonas aeruginosa* in epithelial cells requires suppression

of the caspase-4 inflammasome

Abby R Kroken[1,2]*, Keith A Klein[1], Patrick S Mitchell[3], Vincent Nieto[2], Eric J Jedel[2], David J

Evans[2,4], Suzanne M J Fleiszig[2, 5]

1. Department of Microbiology and Immunology, Loyola University Chicago, Maywood, IL

USA

2. Herbert Wertheim School of Optometry & Vision Science, University of California,

Berkeley, CA USA

3. Department of Microbiology, University of Washington, Seattle, WA, USA

4. College of Pharmacy, Touro University California, Vallejo, CA USA

5. Graduate Groups in Vision Sciences, Microbiology, and Infectious Diseases & Immunity,

University of California, Berkeley, CA USA

* Corresponding author. Department of Microbiology and Immunology, Loyola University

Chicago, Maywood, IL, USA. Email: akroken@luc.edu

Abstract

Pathogenesis of *Pseudomonas aeruginosa* infections can include bacterial survival inside epithelial cells. Previously, we showed this involves multiple roles played by the type three-secretion system (T3SS), and specifically the effector ExoS. This includes ExoS-dependent inhibition of a lytic host cell response that subsequently enables intracellular replication. Here, we studied the underlying cell death response to intracellular *P. aeruginosa*, comparing wild-type to T3SS mutants varying in capacity to induce cell death and that localize to different intracellular compartments. Results showed that corneal epithelial cell death induced by intracellular *P. aeruginosa* lacking the T3SS, which remains in vacuoles, correlated with activation of NF-κB as measured by p65 relocalization and TNFα transcription and secretion. Deletion of caspase-4 through CRISPR-Cas9 mutagenesis delayed cell death caused by these intracellular T3SS mutants. Caspase-4 deletion also countered more rapid cell death caused by T3SS effector-null mutants still expressing the TSSS apparatus that traffic to the host cell cytoplasm, and in doing so rescued intracellular replication normally dependent on ExoS. While HeLa cells lacked a lytic death response to T3SS mutants, it was found to be enabled by interferon gamma treatment. Together, these results show that epithelial cells can activate the noncanonical inflammasome pathway to limit proliferation of intracellular *P. aeruginosa*, not fully dependent on bacterially-driven vacuole escape. Since ExoS inhibits the lytic response, the data implicate targeting of caspase-4, an intracellular pattern recognition receptor, as another contributor to the role of ExoS in the intracellular lifestyle of *P. aeruginosa*.

Importance

Pseudomonas aeruginosa can exhibit an intracellular lifestyle within epithelial cells in vivo and in

vitro. The type three secretion system (T3SS) effector ExoS contributes via multiple mechanisms,

including extending the life of invaded host cells. Here, we aimed to understand the underlying

cell death inhibited by ExoS when *P. aeruginosa* is intracellular. Results showed that intracellular

P. aeruginosa lacking T3SS effectors could elicit rapid cell lysis via the non-canonical

inflammasome pathway. Caspase-4 contributed to cell lysis even when the intracellular bacteria

lacked the entire T33S and were consequently unable to escape vacuoles, representing a naturally

occurring subpopulation during wildtype infection. Together, the data show the caspase-4

inflammasome as an epithelial cell defense against intracellular *P. aeruginosa*, and implicate its

targeting as another mechanism by which ExoS preserves the host cell replicative niche.

Introduction

Pseudomonas aeruginosa is an opportunistic pathogen able to cause life- and vision-threatening infections. While often referred to as an extracellular pathogen, *P. aeruginosa* can adopt an intracellular lifestyle in various epithelial cell types, including corneal (1), bronchial (2), HeLa cells (3) and in vivo (4). The ability of *P. aeruginosa* to replicate inside a host cell depends on the type three secretion system (T3SS), with multiple roles played by the T3SS effector ExoS (5-7) encoded by only invasive strains (8). ExoS is a bifunctional protein with both RhoGAP and ADP ribosyltransferase (ADPr) activity. Broad substrate specificity of its ADPr domain (9, 10) enables ExoS to impact multiple host processes (11, 12). Demonstrated effects of ExoS include inactivation of host cell proliferative/survival signaling, disassembling the host cell cytoskeleton, freezing of intracellular membrane trafficking, and disruption of ROS generation (9, 13-16). While the molecular components of host cells targeted by ExoS to enable *P. aeruginosa* to replicate intracellularly have not yet been elucidated, its ADPr activity is required, and mechanisms shown include inhibition of autophagy (17) and evasion of lysosomes (18). Recently, we reported another role for the ADPr activity of ExoS in intracellular persistence: preservation of host cell viability (19), which was further explored here.

Eukaryotic host cells can sense and respond to intracellular pathogens (20, 21). Responses can include initiation of regulated cell death pathways (22), which may subsequently be suppressed by the bacteria to preserve their intracellular niche (23). One type of host response is pyroptosis, a form of inflammatory lytic cell death (20). Multiple pyroptotic pathways have been identified, each sensing different pathogen-associated molecular patterns (PAMPs) (24-27) or pathogen-

associated aberrant activities (28, 29). The sensor converges on the activation of a cysteine-aspartic

protease: caspase-1 for canonical inflammasome pathways (30, 31), and caspase-4/5 for later-

discovered pathways designated noncanonical (32-34). Inflammatory caspases cleave and activate

gasdermin D (35, 36), which assembles into pores in host plasma membranes (37). This leads to

release of cytokines IL-18 and IL-1β (38), and total lysis for many cell types (39).

P. aeruginosa effectors ExoS and ExoT can reduce IL-1β secretion in macrophages (40-42),

suggesting an ability to interfere with inflammasome activation. Responses of macrophages and

epithelial cells can differ. While inflammasomes are well-studied in macrophages and other

myeloid cells, the repertoire of inflammasomes in epithelial cells is usually limited, and their

functions less-well studied (43). One inflammasome pathway studied in some corneal diseases is

NLRP3 (44, 45), which can detect numerous stimuli (46) including ionic flux from pore formation

(47), lysosomal damaging agents (48), and bacterial RNA (49). However, not all relevant studies

verified which corneal cell type expressed NLRP3, or explicitly determined if NLRP3 was

activated, versus other inflammasome pathways that could also yield mature IL-1β. Recently, both

caspase-4, which detects cytoplasmic LPS (50), and NLRP1, which detects pathogenic enzymatic

activities and dsRNA (28, 51), were shown to be expressed and functional in the human corneal

epithelium (52, 53). The Protein Atlas RNAseq data set for the corneal epithelial cell line hTCEpi

(54) provides evidence for expression of caspase-4, caspase-5 (an additional LPS sensor (33, 55)),

NLRP1, and NLRC4 (although its required sensor protein NAIP (56) was not detected) (57). While

NLRP3, Pyrin, and AIM2 inflammasomes were not detected in these cells, this does not preclude

upregulation upon specific stimuli or in vivo (57).

Having shown that the T3SS effector ExoS inhibits rapid cell lysis induced by intracellular *P.*

aeruginosa in corneal epithelial cells (19), we investigated the mechanisms underlying host cell

death elicited and modulated by intracellular *P. aeruginosa*. In doing so, we also leveraged the

finding that *P. aeruginosa* mutants lacking the entire T3SS (Δ*exsA* mutants) remain toxic to

corneal epithelial cells (58), contrasting with HeLa and CHO cells (59, 60). Results showed that

caspase-4 is required for rapid corneal epithelial cell death in response to *P. aeruginosa* invasion,

and in this way limits accumulation of an intracellular population of bacteria. While caspase-4-

dependent death occurred even if bacteria lacked the ability to leave vacuoles/enter the cytoplasm,

the response was faster when they could. Moreover, the death response was enabled in otherwise

unresponsive HeLa cells after stimulation with IFN-γ, which has been shown to limit cytoplasmic

sub-populations of *Salmonella* in a manner dependent on caspase-4 and GBP proteins (61). Since

ExoS inhibits the cell death response to intracellular *P. aeruginosa*, these results implicate ExoS

targeting a caspase-4-dependent response as another contributor to its well-established role in

intracellular survival by *P. aeruginosa*.

Results

***P. aeruginosa* lacking the T3SS kill corneal epithelial cells but not HeLa cells.**

Generally, it is thought that *P. aeruginosa* mutants missing the T3SS (i.e., Δ*exsA* mutants) do not

kill epithelial cells, as shown for HeLa and CHO cells (59, 60). However, our published work has

shown that Δ*exsA* mutants are able to kill corneal epithelial cells, with contributions made by the

intracellular population (58). Recognizing that differences between cell types could assist in

deciphering mechanisms, we compared corneal epithelial cells (hTCEpi) (54) to HeLa cells using

the same methods. Impact of wild type PAO1, ΔexsA mutants (lacking the entire T3SS), and ΔexoSTY mutants (lacking all known T3SS effectors) were studied. After a 3-hour invasion period, extracellular bacteria were eliminated using the non-cell permeable antibiotic amikacin (3). Cell death rates were measured with a FIJI macro that counts propidium iodide-positive nuclei (permeabilized cells) and Hoechst-labeled nuclei (total cells) and reports a ratio over a 20-hour post-infection period (19). The results confirmed that the two cell types were differentially susceptible. Death rates for corneal epithelial cells were statistically similar between wild type and ΔexsA mutants (**Figure 1A-B**), whereas HeLa cells infected with ΔexsA mutants showed very little cell death even 20-h post infection (**Figure 1C-D**). In both cell types, the ΔexoSTY mutant yielded the most rapid cell death as noted previously (19). Thus, corneal epithelial cells have an intrinsic response to T3SS-null bacteria that is absent in HeLa cells, which may underly cell death occurring in response to T3SS-positive bacteria.

Dynamics of cell death in bacterially-occupied cells indicates roles for both T3SS machinery and T3SS effectors.

Visual inspection of infected cells suggested that not all cells were occupied by bacteria, and that some non-occupied cells also died. Recognizing that this could skew results in bulk analyses of cell death rates, we performed additional experiments to analyze only bacterially-occupied cells. To enable this, we employed methods that allowed simultaneous detection of viable intracellular bacteria and host cell (19). Wild type (PAO1) was compared to mutants lacking the entire T3SS (ΔexsA mutants), and ΔexoSTY mutants lacking only T3SS effectors, using quantitative time lapse imaging over 20 h. Since ΔexsA mutants do not kill HeLa cells, this analysis was only done using corneal epithelial cells.

A method of detecting intracellular bacteria is needed, as amikacin-killed bacteria adhere to the

dishes and cells, making intracellular bacteria impossible to distinguish if using constitutive

fluorophore expression. To detect Δ*exsA* mutants inside cells, we previously used an arabinose-

inducible green fluorescent protein (GFP) expression vector pBAD-GFP (58). This method

involves induction of GFP expression using arabinose induction only after bacteria have invaded

cells and extracellular populations are killed (using amikacin). To keep experimental methods

consistent across samples, we explored the feasibility of using the induction method to also study

wild type and the Δ*exoSTY* mutants. In prior studies, these strains were visualized using a reporter

for the T3SS (3), because T3SS effectors are needed for intracellular replication (5, 19).

Unfortunately, fewer bacterial cells exhibited cytoplasmic spread using the arabinose induction

method compared to the T3SS reporter method (**Supplemental Figure 1A**). Moreover, while the

host cell death rate was similar for wild type PAO1 using both methods, it trended lower for

Δ*exoSTY* mutants expressing arabinose-induced GFP (**Supplemental Figure 1B)**. This suggested

GFP induction impacts the T3SS or some other factor relevant to cytoplasmic entry or spread: a

critical step for intracellular infection by wild type and Δ*exoSTY* mutants, but not for Δ*exsA*

mutants (which remain in vacuoles (58)). Indeed, T3SS effector secretion was reduced *in vitro*

using EGTA stimulation in the presence of arabinose when bacteria were also transformed with

pBAD-GFP (**Supplemental Figure 1C**). Thus, the T3SS expression plasmid pJNE05 was used to

study intracellular wild type and Δ*exoSTY* mutants, reserving arabinose induction of GFP for the

Δ*exsA* mutant. A limitation of this approach is that a sub-population of intracellular wild type

bacteria can remain T3SS-negative (58) and would be undetected in this analysis. However, the

advantage of this approach is that T3SS-dependent outcomes for intracellular bacteria can be

spatially isolated and examined independently, and then compared to homogenous populations of

Δ*exsA* mutants missing the T3SS apparatus.

Representative images from a time-lapse experiment using Δ*exsA* mutants are shown in **Figure**

2A. As expected, Δ*exsA* mutants localized to vacuoles inside cells (58). Most bacteria-occupied

cells died before the end of the 20 h assay, as shown by propidium iodide labeling of the nucleus.

However, **Figure 2B** shows an invaded cell still alive at 20 hours (no propidium iodide labeling),

representing a minority of cells at this time point. Videos of these time lapse experiments are

available in **Supplemental Movie 1**, which includes cells visualized in **Figures 2B and 2C** at 20

hours post infection. As shown previously, both wild type PAO1 and Δ*exoSTY* mutant infected

cells (shown in **Figure 2D)** replicated in the host cell cytoplasm after bacterial escape from

vacuoles (19). In both cases, most invaded cells died by the end of the assay, the Δ*exoSTY* mutant

doing so even more rapidly, expected due to lack of ExoS which normally counters cell death (19).

Hours following host cell death, propidium iodide of intracellular bacterial bodies eventually

replaced GFP signal (**Figure 2C and Supplemental Movie 1)**.

A computational approach was used to measure when populations of invaded cells died (19), and

a super violin plot visualization script developed by Kenny and Schoen was used to show

independent replicates as stripes single plots (62). This analysis examined 2,710 total invaded cells

over 6 replicate experiments. Results confirmed that intracellular Δ*exsA* mutants killed their

occupied cells at a broad range of time points, and the mean value of invaded cell survival times

was unexpectedly similar to wild type PAO1 (**Figure 3E**). The analysis also confirmed that the

intracellular Δ*exoSTY* mutant infections caused cell lysis at a more rapid rate, significantly

different from both wild type PAO1 and Δ*exsA* mutants (19). Thus, while cell death can be driven

by vacuolar contained bacteria, it was more rapid if bacteria could access the cytoplasm while

lacking effectors.

T3SS-null mutant bacteria engage NF-κB signaling in corneal epithelial cells.

To explore molecular mechanisms involved in *P. aeruginosa*-driven cell death, a real time PCR

array with genes involved in regulated cell death pathways was used to study candidate host

responses. Results showed that TNFα and BCL2A1 were upregulated in excess of fifty-fold in

both cell lines: (**Figure 3**). In the corneal cells, wild type and both mutants upregulated these genes,

while in HeLa cells, only the PAO1Δ*exoSTY* mutant had that impact.

Since TNFα and BCL2A1 can be regulated by NF-κB, activation of NF-κB was measured by

observing translocation of p65 to the nucleus (**Figure 4A**). In corneal epithelial cells, infection

with each tested strain caused p65 translocation, correlating with positive target gene transcription.

In HeLa cells, only infection with Δ*exoSTY* mutants led to p65 translocation (**Figure 4B**). The

result with wild type PAO1 infection was difficult to discern due to cell rounding and low

cytoplasmic area; however, p65 appeared excluded from the nuclear region of HeLa cells (**Figure**

4B, inset). Quantification of p65-positive nuclei was accomplished with an ImageJ macro (see

methods) (**Figure 4C and D**). This finding correlated with the transcriptional data, and is

suggestive of a mechanism in which the T3SS effectors limit p65 translocation in HeLa cells, but

not corneal epithelial cells. Since TNFα is associated with both apoptotic and necrotic cell death,

TNFα secretion from infected cells was measured by ELISA. The results showed that while wild

type PAO1 stimulated TNFα secretion from corneal epithelial cells, higher levels were detected

after infection with the Δ*exoSTY* mutant, and the Δ*exsA* mutant causing an intermediate response

(**Figure 4E**). TNFα was unable to be detected from HeLa cells infected with any strain.

Having found TNFα secretion levels correlated with cell death rates, we next asked if there was a

direct causative relationship that might also explain why some cells without visible intracellular

bacteria could also be killed. Thus, media from naïve cells or Δ*exsA* mutant-infected cells was

replaced with filter-sterilized media from Δ*exoSTY* mutant-infected cells (high TNFα). The

outcome showed that this did not enhance host cell death rates (**Figure 4F**), suggesting cell death

is not driven by a secreted factor such as TNFα, and requires bacteria present.

Caspase-4 is involved in cell death response to intracellular *P. aeruginosa*.

The morphology of invaded cell death included membrane integrity loss, intact nucleus, and

transient blebbing, all suggestive of pyroptosis (**Figure 2**) (22). Since caspase-4 can restrict

numerous other Gram-negative bacteria (61, 63), and the pathogen *Shigella* inhibits the caspase-4

inflammasome by multiple mechanisms to survive (64-66), and corneal epithelial cells were shown

to express caspase-4 (53), we tested the hypothesis that caspase-4 was involved in driving cell

death in response to intracellular *P. aeruginosa*. We generated *CASP4* knockout corneal epithelial

cells, and loss of caspase-4 protein was confirmed by Western blot (**Supplemental Figure 2**).

Control cell lines with non-targeting guide RNA were also generated using an identical selection

strategy and grown up as monoclonal lines; each exhibited death rates consistent with wild type,

non-transduced hTCEpi cells (**Supplemental Figure 3**).

Invasion by wild type PAO1 yielded similar cell death timing in both *CASP4* knockout cells as

wild type corneal epithelial cells (**Figure 5A, B, C**). In contrast, Δ*exoSTY* mutant-invaded cells

experienced significantly increased survival times (**Figure 5A, B, D**), which allowed substantial

accumulation of intracellular bacteria. A smaller but significant increase of invaded cell survival

times were also observed with PAO1Δ*exsA* mutant-occupied *CASP4* knockout cells (**Figure 5E**).

Bacterial replication was measured using summed GFP area contained in live cell masked region

generated by the FIJI macro (19). Results indicate increased bacterial replication in *CASP4*

knockout cells, with the most substantial increase observed in cells invaded by Δ*exoSTY* mutants

(**Figure 5F-H**). These data implicated the non-canonical inflammasome as a dominant response

to cytoplasmic invasion of T3SS-positive *P. aeruginosa* independently of the T3SS effectors,

and also show involvement in responding to vacuole-occupying bacteria. Unchanged cell death

timing for wild-type PAO1 in *CASP4* knockout cells aligns with our previously published data

showing that the T3SS effector exotoxins, specifically the ADP ribosyltransferase activity of

ExoS (19), counters cell death. Of note, while *CASP4* knockout reduced corneal epithelial cell

death associated with intracellular Δ*exsA* mutants, many of the cells still died by the end of the

20-hour imaging time frame (**Figure 1 and 5E**). This implies additional mechanisms for

recognizing vacuolar/T3SS-negative *P. aeruginosa* in addition to caspase-4 which are also

absent from HeLa cells.

Prior studies implicate Exotoxin S in delaying host cell lysis in response to invasion (19), thus

we examined the Δ*exoS* mutant in both wt corneal epithelial cells and *CASP4* knockout cells to

evaluate whether there is a role for ExoT and ExoY in suppressing caspase-4-mediated lysis or

supporting cytoplasmic bacterial grown (**Supplemental Figure 4**). Comparing to the

aforementioned study which tested strains encoding a single effector (e.g., ΔexoST or Δ*exoSY*)

(19), the ΔexoS mutant showed a similar phenotype to Δ*exoSY* mutant, which expresses only

ExoT: bacteria were unable to replicate substantially within in wt corneal epithelial cells

(**Supplemental Figure 4**). Bacterial mutant replication was rescued in *CASP4* knockout cells.

However, the timing of Δ*exoS*-invaded wt cell survival was irregular across the timeframe

observed (**Supplemental Figure 4C**), which may relate to ExoT's ability to block invasion (67,

68), or interfere with other inflammasomes (41). However, ExoT and ExoY were unable to

contribute to intracellular replication in cells with caspase-4.

IFN-γ-stimulation enables HeLa cells to respond to intracellular *P. aeruginosa*. HeLa cells

support cytoplasmic hyper-replication of *P. aeruginosa* before significant impact on host cell

viability occurs (3, 19). The bacterium *Salmonella* accomplishes a similar feat (69). Santos and

colleagues demonstrated that IFN-γ stimulation causes HeLa cells to lyse in response to

Salmonella entry into the cytosol, which was associated with upregulation of GBP proteins and

stimulation of the caspase-4 inflammasome. (61). We examined the impact of IFN-γ stimulation

on HeLa cells containing intracellular *P. aeruginosa* (**Figure 6**). The kinetics of wild type PAO1-

induced host cell death remained unchanged. However, the rate of cell death increased for both

PAO1Δ*exoSTY* and for PAO1Δ*exsA* infections (**Figure 1 and 6**) (59). IFN-γ-treated HeLa cells

occupied by Δ*exsA* mutants died over varied timepoints during the 20 hours of infection similar to

results for corneal epithelial cells. Aligning with the quick timing of cell lysis, PAO1Δ*exoSTY*

mutants were no longer able to replicate effectively in the cytoplasm of IFN-γ-treated HeLa cells.

Thus, IFN-γ-treated HeLa cells resembled corneal epithelial cells in their response to intracellular

P. aeruginosa.

Discussion

Invasive (ExoS-expressing) strains of *P. aeruginosa* can thrive inside a wide range of host cells

(2, 3, 5). After invading an epithelial cell, ExoS plays multiple roles in supporting intracellular

survival and replication, including inhibition of host cell lysis (19). This study aimed to

understand the underlying host response inhibited by ExoS. Knowing *P. aeruginosa* diversifies

intracellularly into vacuolar and cytoplasmic populations, bacteria in both locations trigger cell

death, their locations are determined by T3SS expression state, and that host cell lysis can be

suppressed by ExoS, we used T3SS mutants to interrogate mechanisms. Results showed that cell

death in response to cytoplasmic *P. aeruginosa* lacking T3SS effectors involved caspase-4; its

deletion inhibiting rapid pyroptotic lysis. Deletion of caspase-4 also partially alleviated cell death

caused by bacteria lacking the T3SS that were restricted to vacuoles (Δ*exsA* mutants). We further

found that while HeLa cells lack this response to intracellular *P. aeruginosa*, it could be induced

by IFN-γ, a stimulator of numerous genes including factors that assist in activation of the

caspase-4 inflammasome (61). This suggests one difference between corneal epithelial cells and

HeLa cells may be among interferon stimulated genes (such as GBP1-4), or the ability of corneal

epithelial cells to promptly upregulate inflammatory genes in response to *P. aeruginosa* (**Figure**

3). Since ExoS inhibits cell death in response to cytoplasmic bacteria (19), these findings

implicate ExoS-mediated interference with the non-canonical inflammasome pathway. ExoS is

only capable of prolonging cell survival to the same level as Δ*exsA* mutant-invaded cell death,

suggesting its role is restricted to the host response to cytoplasmic bacteria.

Since *P. aeruginosa* is often assumed an extracellular pathogen, previous studies of host

epithelial cell death have been from the perspective of extracellular bacteria. In that regard, ExoS

alone has cytotoxic activities. HeLa cells can undergo apoptotic cell death based on both

caspase-3 (59) and caspase-8 activity (70), and activation of apoptosis through JNK1 and

cytochrome c release (71). This was confirmed using a mammalian expression system for

introducing ExoS in the absence of bacteria, which showed it was sufficient to induce HeLa cell

apoptosis even in the absence of bacterial interaction (72). Placing our findings into the context

of these earlier HeLa cell studies, we note differences in our experimental systems: while they

explored mechanisms for cell death caused by ExoS, we studied cell death triggered by live

intracellular bacteria lacking ExoS. Were intracellular bacteria present in these assays, HeLa

cells would not have responded to them without stimulation. Our experiments also provide an

analysis of only invaded cells excluding extracellular bacteria to focus on impact of intracellular

bacteria, and used corneal cells which naturally respond to bacterial pathogen-associated

molecular patterns. We then showed that IFN-γ-stimulated HeLa cells respond in the same

manner: ExoS inhibits rather than drives a host cell death response. In sum, while introducing

ExoS in isolation is a useful surrogate for *in vitro* experimentation to address specific questions,

other factors modify the impact of ExoS in other situations. Despite differences in experimental

models, it remains important to reconcile outcomes. The innate ability of ExoS to both drive and

inhibit cell death, involving different types of regulated cell death, and correlating with presence

or absence of internalized *P. aeruginosa,* is interesting and worth exploring. Indeed, it could aid

in determining mechanisms by which ExoS blocks pyroptosis, given crosstalk between

pyroptosis and apoptosis (73).

The specific mechanism by which the ADPr activity of ExoS interferes with caspase-4-mediated

cell lysis remains to be identified. Recently, *Shigella flexneri* was shown to block the non-

canonical inflammasome by multiple mechanisms: caspase-4 inactivation from ADP-

riboxanation by the effector OspC3 (64, 65), ubiquitylating gasdermin D (74), and GBP1 (75) to

target each for degradation. ExoS is unique among bacterial ADP ribosyltransferases for having

numerous host substrates, though currently none are known to interact with the caspase-4/11

pathway (76). Since there are many possibilities to investigate, comprehensive identification of

uncharacterized ExoS substrates and their potential involvement will require a separate study.

Considering that *CASP4* knockout does not fully prevent cell death triggered by vacuole-bound

T3SS mutants (**Figure 6**), it would also be worth exploring which additional cell death pathways

are involved.

In summary, the results of this study demonstrate a role for caspase-4 in limiting intracellular

colonization of *P. aeruginosa* in ocular surface cells, a phenomenon triggered by the presence of

intracellular bacteria but inhibited by the T3SS effector ExoS. Importantly, the data support the

notion that *P. aeruginosa* can function as both an intracellular and extracellular pathogen,

suggestive of a T3SS effector function devoted to inhibition of an *intracellular* pattern

recognition receptor, consistent with its unusual capacity to adapt to its environment, survive

hardship, and exist ubiquitously. Ultimately, development of effective strategies to prevent or

treat the devastating infections caused by *P. aeruginosa* will require a thorough appreciation of

these complexities.

Methods

Bacterial strains and cell lines. *P. aeruginosa* strain PAO1 and isogenic mutants of *exsA*, or

ExoS/ExoT/ExoY, or ExoS were used for all infection experiments (77). Plasmids used for

visualization were pJNE05 (78) and pBAD-GFP (58). Plasmid selection was achieved with 100

µg/ml gentamicin. Corneal epithelial cells (hTCEpi) (54) were maintained in KGM-2 media

(Lonza) lacking gentamicin. HeLa cells were maintained in phenol red-free DMEM (Gibco) with

10% FBS.

CRISPR-Cas9 knockout cell line. Lentiviral particles for transduction were generated from

293T cells transfected with psPAX2 (Didier Trono, Addgene plasmid # 12260) and pMD.2

(Didier Trono, Addgene plasmid # 12259), combined with either Cas9 expression vector plenti-

Cas9-blast (Feng Zhang, Addgene plasmid # 52962) (79) or a guide RNA vector plenti-LIC-Puro

which was adapted for ligation-independent cloning (gifted by Moritz Gaidt) (80). Guide RNA

sequences targeting caspase-4 (5'- CCACAGAAAAAGCCACTTA - 3') or a non-targeting

control sequence (5'- GACGGAGGCTAAGCGTCGCA - 3') were cloned into plenti-LIC-Puro

using ligation independent cloning. After six hours, media on 293T cells was replaced with

KGM-2. Media was collected at 48 hours and passed through a 0.45 µm filter and added directly

to hTCEpi cells, which were then centrifuged at 1200g for 90 minutes at 37°C. Antibiotic

selection with blasticidin (50 µg/ml) or puromycin (10 µg/ml) was performed after two days. For

generation of monoclonal lines, single cells were seeded in 96-well plates. Isolated colonies were

identified and grown up over approximately 12 days before transfer into 6-well plates to avoid

differentiation induced by crowding. *CASP4* knockout candidates were screened by Western

blot: lysates from a 6-well plate were collected in 150 µl RIPA buffer, frozen and thawed, and

then clarified by centrifugation for 20 minutes at 4ºC. Samples were run on 10% stain free gels

(Bio-Rad) and total protein visualized as a loading control prior to transfer to 0.2 µm PVDF

membrane. Membrane was blocked with Bio-Rad EveryBlot block and probed with anti-caspase-

4 antibody (Santa Cruz, sc-56056, 1:1000). Blot was washed using TBST, and antibody detected

with goat-anti-rabbit HRP (Bio-Rad, 1706515, 1:5000).

Infection experiments. One day preceding experiments, hTCEpi cells were plated on No. 1.5

glass-bottom 24 well plates (MatTek) in KGM-2 at 75% confluence with 1.15mM calcium to

induce differentiation (54). HeLa cells were maintained in DMEM with 10% FBS and plated at

50% confluence on optical plastic 8-well chambered coverslips (ibidi). Where indicated, 50

ng/ml IFN-γ (Peprotech) was added at 16 hours prior to infections of HeLa cells and maintained

throughout the infection and imaging.

Bacterial suspensions were made in PBS from 16-hour lawns grown on TSA media at 37°C. The

absorbance at 540 nm was measured, and MOI of 10 was calculated using OD540 of 1 equal to 4

x 10^8 CFU per mL. Bacteria were directly to cell culture media and allowed to infect for 3 hours.

For live imaging, Hoechst (0.8 µg/ml) was added to the cells prior to infections and allowed to

label the cells during the 3-hour infection period. Media was replaced with media containing the

antibiotic amikacin (200 µg/ml) and propidium iodide (0.8 µg/ml). Where needed, a 1:10

dilution of 10% arabinose in media was added at 3.75 hours post infection.

RT-PCR arrays. Cells in 10cm dishes were infected with PAO1, PAO1Δ*exoSTY*, or

PAO1Δ*exsA* for 3 hours at an MOI of 10, and media was replaced for 1 hour with 200 µg/ml

amikacin. At 4 hours, cells were washed twice in PBS and collected in 1ml TriReagent (Sigma).

RNA was purified using Zymo Direct-zol RNA MiniPrep. cDNA synthesis was performed with

the RT2 first strand kit (Qiagen) using 0.5µg RNA. RT PCR was performed on a Roche

LightCycler 96 using the The RT² Profiler™ PCR Array Human Cell Death PathwayFinder

(PAHS-212Z, Qiagen)w, and analyzed in Qiagen's Data Center Analysis Portal.

(https://geneglobe.qiagen.com/us/analyze).

Fixed immunostaining. Cells were seeded on No. 1.5 glass coverslips and infected as described

above. At four hours post infection, cells were washed twice in PBS and fixed in 4%

paraformaldehyde in PBS for 10 m. Cells were washed twice in PBS, and neutralized with 150

mM glycine in PBS for 10 m. Cells were washed twice in PBS, and permeabilized/blocked (5%

FBS, 2.5% cold fish skin gelatin, 0.1% TritonX-100, 0.05% Tween-20 in PBS) for 1 h. Antibody

in solution (2.5% FBS, 1.25% cold fish skin gelatin, 0.1% TritonX-100, 0.05% Tween-20 in

PBS) was incubated overnight at 4°C. Cells were washed 4 times (5 minutes), and secondary

antibodies were added in antibody solution for 1 h. Cells were washed once, labeled with DAPI

for 5 minutes, and washed twice (5 minutes). Coverslips were mounted using ProLong Diamond.

Microscopy. Images for Figs 1 and 4A, B were captured on a Nikon Ti-E inverted microscope

equipped with a Lumencor Spectra X LED Light Engine illumination source. All other images

were captured on a Ti2-E inverted microscope with X-Cite XYLIS XT720S Broad Spectrum

LED Illumination System. Both systems used Nikon perfect focus, an Okolab stage-top

incubation chamber, a DS-Qi2 CMOS camera, and CFI Plan Apochromat Lambda D 40X air NA

0.95 objective. Time lapse fields were selected between hours 3 and 4 without observing

fluorescence channels. Eight fields per condition were imaged hourly from 4 to 20 hours post

infection.

Image Analysis. Time-lapse images were computationally analyzed using two custom macros

written for the FIJI package of Image J aspreviously described (19). The code for measuring

rates of cell death for the whole population, and set of macros for tracking the invasion state of

cells is available in a GitHub repository: https://github.com/Llamero/Nuclei_analysis-macro.

Subsequent Python scripts for exporting the analysis TIF file and processing are available in the

following GitHub repository: https://github.com/abbykroken/cell_survival_with_bacteria. An

Image J macro for measuring a ratio of p65 stain intensity within the nucleus and the periplasmic

region has been made available in the following GitHub repository:

https://github.com/abbykroken/Nucleus-to-cell-ratio.

In vitro T3SS effector secretion. Bacteria were grown for 5 hours in 5ml TSB supplemented

with 100mM MSG and 1% glycerol at 37ºC with shaking at 200 rpm. EGTA (2mM) was added

to induce T3 secretion. OD 540 nm readings were taken prior to supernatant protein

concentration and used to normalize volumes. Bacteria were centrifuged at 12000g to clarify 1ml

of supernatant. Proteins were concentrated using TCA precipitation: 250 µl of 100% cold TCA

was added to 1 ml supernatant for 30 minutes and centrifuged at 14000g for 5 minutes. The

pellet was washed with 1 ml cold acetone twice and suspended in 4x laemmli buffer normalized

to starting OD (max volume 50 µl). Proteins were visualized using 10% stain-free gels (Bio-

Rad).

Statistics. Statistical analyses were performed and data presented using Graph Pad Prism 9. Super

violin plots were prepared using scripts published by Kenny and Schoen (62) and output images

aligned on axes generated in Graph Pad Prism 9. Data were shown as a mean ± standard deviation

(SD) of 3-6 independent experiments unless otherwise indicated. Comparison of two groups was

performed by Student's t-test, three or more groups by One-way ANOVA with Tukey's post-hoc

analysis. Comparison between two groups for total cell death rates overtime was performed by

multiple column t-tests for each timepoint, and the two samples that were compared are specified

in the figure legends. In each instance, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.005$ and **** $P < 0.001$.

Acknowledgements

Thanks to Arne Rietsch (Case Western Reserve University, OH) for providing *P. aeruginosa*

strains and mutants, Timothy L Yahr (University of Iowa, IA) for plasmid pJNE05, and Danielle

M Robertson (University of Texas Southwestern, TX) for hTCEpi cells. Thanks to Benjamin E

Smith (University of California, Berkeley, CA) for continued support of FIJI image analysis

macros. We also thank Russell E Vance (University of California, Berkeley, CA) for helpful

suggestions during the course of this project.

ARK, PSM, VN, DJE, and SMJF designed the experiments; ARK, KAK, PSM, and EJJ performed

the experiments; ARK, KAK, PSM, VN, DJE, SMJF analyzed and interpreted the data; ARK,

DJE, and SMJF wrote the manuscript; ARK, DJE, and SMJF supervised the study.

This work was supported by the National Institutes of Health R01EY011221 (SMJF),

R01EY034239 (ARK), F32 EY025969 (ARK), DP2 AI154432 (PSM), P30 DK089507 (PSM),

F32 EY029152 (VN), and the Mallinckrodt Foundation (PSM). The funding agencies had no role

in the study design, data collection and interpretation, or decision to submit the work for

publication.

Figure 1. Host cell death rates from *P. aeruginosa* mutant infections. **(A)** Corneal epithelial

cells (hTCEpi) were infected with indicated strain: wt PAO1, PAO1Δ*exoSTY*, or PAO1Δ*exsA* for

3 hours at an MOI equal to 10. Hoechst (blue) was added to label nuclei. Non-associated bacteria

were removed, and media with amikacin and propidium iodide (red) was added to kill extracellular

bacteria. Cells were imaged hourly from 4 to 20 hours post infection. Select fields from indicated

times are shown. Scale bar = 50 μm. **(B)** The percent of cells positive for propidium iodide at each

timepoint was determined by using a custom FIJI macro to segment all nuclei from the Hoechst

channel, and all nuclei of dead cells in the propidium iodide channel. Multiple column t-tests were

performed comparing wt PAO1 and PAO1Δ*exsA*, and SD error bars are displayed from three

replicates. **(C)** Experiment was performed identically as in panel A, but using HeLa cells. Scale

bar = 50 μm. **(D)** Analysis was performed as in panel B, using the data obtained in panel C.

Figure 2. Intracellular Δ*exsA* mutants kill invaded cells at a similar rate to wt PAO1-invaded

cells. (A) Bacteria were transformed with an arabinose-inducible GFP plasmid. Corneal epithelial

cells (hTCEpi) were infected at an MOI of 10 and extracellular bacteria eliminated at 3 hours post

infection using amikacin. A final concentration of 1% arabinose was used to induce GFP in

surviving intracellular bacteria beginning at 3.5 hours post infection. Time lapse imaging was

conducted hourly from 4 to 20 hours, and host cell nuclei detected with Hoechst and propidium

iodide to determine time of death. Scale bar = 50 μm. Insets 1-2 show the boxed cell at 6 or 11

hours post infection, respectively. **(B)** An example of a live invaded cell from 20 hours post

infection. **(C)** Example of a dead invaded cell from 20 hours post infection, where the PI channel

has been saturated such that PI-positive bacteria can be visualized. Scale bar = 50 μm. **(D)** The

T3SS-GFP reporter pJNE05 was used to visualize wt PAO1 and PAO1Δ*exoSTY* infections, as

described previously (19), to compare to PAO1Δ*exsA* infections. Scale bar = 50 µm. Time lapses

from panels A through D are available as **Supplemental Movie 2**. **(E)** A computational analysis

approach was used to segregate only invaded cells, and determine their survival time in hours.

PAO1 condition includes 881 cells over six replicates; PAO1Δ*exoSTY*, 967 cells; PAO1Δ*exsA*,

862 cells. The survival times of all invaded cells were combined a single super violin plot. Mean

survival times with SD error bars are displayed, and significance determined by One-way

ANOVA. *** $P < 0.005$.

Figure 3. Transcriptional responses of infected host cells. (A) Corneal epithelial cells (hTCEpi)

were infected with wt PAO1, PAO1Δ*exoSTY*, or PAO1Δ*exsA* for 3 hours at an MOI equal to 10.

Non-associated bacteria were removed, and media with amikacin was added for 1 additional hour.

At four hours, RNA was purified from infected cells, and RT PCR analysis performed using a

commercial array for genes associated with specific cell death pathways. **(B)** Experiment

performed identically to panel A, except using HeLa cells.

Figure 4. NF-κB signaling and the role of secreted factors in cell death. (A) Corneal epithelial

cells (hTCEpi) were infected with wt PAO1, PAO1Δ*exoSTY*, or PAO1Δ*exsA* for 3 hours at an

MOI equal to 10. Then media was replaced with amikacin-containing media. Cells were fixed at

4 hours post infection and p65 localization was determined using immunofluorescent staining.

Nuclei were labeled with DAPI. Scale bar = 50 µm. **(B)** Experiment performed identically to panel

A, using HeLa cells. **(C)** Quantification of p65-positive nuclei from corneal epithelial cells, and

(**D**) HeLa cells. PAO1-infected HeLa cells were not able to be accurately quantified (indicated

using "ND") by automated analysis due to the small and irregular cytoplasmic area. (**E**) Corneal

epithelial cells were infected as described in panel A. Supernatant of infected cells was collected

at 4 hours post infection and TNFα measured by ELISA. **(D)** Supernatants of corneal epithelial

cells infected with PAO1Δ*exoSTY*, PAO1Δ*exsA*, or uninfected were collected at 4 hours post

infection and sterilized with 0.22 µm filter, and used as the replacement media on uninfected cells

or PAO1Δ*exsA* infected cells at 3 hours post infection, combined with amikacin and propidium

iodide. Hoechst was used to label all nuclei. The cell death rates were measured by time lapse

imaging, error bars display SD from three replicates.

Figure 5. Caspase-4 limits intracellular replication and persistence of T3SS mutant bacteria.

(A) Corneal epithelial cells (hTCEpi) or (B) cells knocked out for caspase-4 were infected with wt

PAO1, PAO1Δ*exoSTY* (each visualized by T3SS-GFP reporter) or PAO1Δ*exsA* (pBAD-GFP

induced) at an MOI of 10. Nuclei were labeled with Hoechst and propidium iodide. Extracellular

bacteria were eliminated with amikacin at 3 hours post infection, and imaged hourly from 4 to 20

hours post infection. Images from 4-9 hours post infection are shown. Scale bar = 50 µm. Full-

field time lapses are available as **Supplemental Movie 3**. (**B-D**) (C-D) Cells from indicated

infections were analyzed with a computational approach to segregate only invaded cells and

measure survival times in hours. Six replicates were combined into a single super violin plot. Mean

survival times with SD error bars are displayed, and significance determined by Student's t-test.

(F-H) GFP area was summed within the boundaries of live cells using the same data set as panels

C-E, and normalized to PAO1 (wt) at 4 hours post infection.

Figure 6. IFN-γ-stimulated HeLa cells are not permissive for PAO1Δ*exoSTY* intracellular

replication nor PAO1Δ*exsA* intracellular persistence. (A) HeLa cells were treated with 50

ng/ml IFN-γ for 16 hours prior to infection with wt PAO1, PAO1Δ*exoSTY* (each using the T3SS-

GFP reporter) or PAO1Δ*exsA* (pBAD-GFP induced) at an MOI of 10. Nuclei were labeled with

Hoechst and propidium iodide. Extracellular bacteria were eliminated with amikacin at 3 hours

post infection, and imaged hourly from 4 to 20 hours post infection. Select time frames are shown

at either 10 or 15 hours post infection. Scale bar = 50 µm. Full-field time lapses are available as

Supplemental Movie 4. **(B)** The rate of cell death was determined using host cell nuclear stains.

All panels are from the same experiment with only indicated bacterial strain shown. Error bars

show SD from three replicates. Multiple column t-tests were performed comparing IFN-γ-treated

cells to untreated.

Supplemental Figure 1. GFP expression from plasmid pBAD-GFP suppresses the T3SS. (A)

wt PAO1 was transformed with the T3SS reporter plasmid pJNE05, or the GFP inducible plasmid

pBAD-GFP and imaged hourly in time lapse. Example fields are shown in which bacteria

expressing GFP under the T3SS reporter exhibit intracellular spread and replication, whereas those

with inducible GFP show only limited intracellular replication. (B) The cell death rates of wt PAO1

or PAO1Δ*exoSTY* expressing GFP by either plasmid were measured, showing a trend of reduced

cell killing rates for PAO1Δ*exoSTY* when GFP expression is induced. (C) wt PAO1 with indicated

plasmid was grown for 5 hours in LB media with EGTA to induce the T3SS, and arabinose to

induce GFP. Supernatant proteins were concentrated by TCA precipitation, and the bands

constituting ExoS and ExoT are indicated.

Supplemental Figure 2. Confirmation of caspase-4 deletion. Lysates from wt hTCEpi cells or

caspase-4 candidate knockout cells were collected in RIPA buffer and clarified. Total protein was

visualized using Bio-Rad stain free gels prior to transfer. Blot was probed with anti-caspase-4

antibody.

Supplemental Figure 3. Control monoclonal cells for CRISPR-Cas9 mutagenesis. A control

cell line was generated using non-targeting guide RNA, and monoclonal line generated. Cells were

infected with wt PAO1, PAO1Δ*exoSTY* (each using the T3SS-GFP reporter) or PAO1Δ*exsA*

(pBAD-GFP induced) at an MOI of 10. Nuclei were labeled with Hoechst and Propidium iodide.

Extracellular bacteria were eliminated with amikacin at 3 hours post infection, and imaged hourly

from 4 to 20 hours post infection. Images were analyzed with a computational approach to

segregate invaded cells and measure survival times in hours. Three replicates were combined into

a single super violin plot. Mean survival times with SD error bars are displayed.

Supplemental Figure 4. Host cell survival time and intracellular replication of single exotoxin

deletion of ExoS. (A) Corneal epithelial cells or cells lacking caspase-4 were infected with wt

PAO1, PAO1Δ*exoSTY,* or PAO1Δ*exoS* (each visualized by T3SS-GFP reporter) at an MOI of 10.

Nuclei were labeled with Hoechst and propidium iodide. Extracellular bacteria were eliminated

with amikacin at 3 hours post infection, and imaged hourly from 4 to 20 hours post infection.

Images from 4-10 hours are shown. Scale bar = 50 µm. (B) GFP area was summed within the

boundaries of live cells and normalized to PAO1 (wt) at 4 hours post infection. (C) Cells from

indicated infections were analyzed with a computational approach to segregate only invaded cells

and measure survival times in hours. Mean survival times with SD error bars are displayed, and

significance determined by Student's t-test.

Supplemental Movie 1. Corneal epithelial cells invaded by *P. aeruginosa* and mutants.
Corneal epithelial cells (hTCEpi) were infected with wt PAO1, PAO1Δ*exoSTY* (each expressing
GFP by T3SS reporter), or PAO1Δ*exsA* (inducible GFP) for 3 hours at an MOI equal to 10. Then
media was replaced with amikacin-containing media. Arabinose was added to PAO1Δ*exsA*
condition at 3.5 hours post infection. Imaging was conducted hourly from 4 to 20 hours.

**Supplemental Movie 2. *CASP4* knockout corneal epithelial cells invaded by *P. aeruginosa*
and mutants.** Corneal epithelial cells (hTCEpi) or *CASP4* knockout cells were infected with wt
PAO1, PAO1Δ*exoSTY* (each expressing GFP by T3SS reporter), or PAO1Δ*exsA* (inducible GFP)
for 3 hours at an MOI equal to 10. Then media was replaced with amikacin-containing media.
Arabinose was added to PAO1Δ*exsA* condition at 3.5 hours post infection. Imaging was conducted
hourly from 4 to 20 hours.

Supplemental Movie 3. IFN-γ-stimulated HeLa cells invaded by *P. aeruginosa* and mutants.
HeLa cells were treated with 50 ng/ml IFN-γ for 16 hours prior to infection with wt PAO1,
PAO1Δ*exoSTY* (each using the T3SS-GFP reporter) or PAO1Δ*exsA* (pBAD-GFP induced) at an
MOI of 10. Nuclei were labeled with Hoechst and Propidium iodide. Extracellular bacteria were
eliminated with amikacin at 3 hours post infection, and imaged hourly from 4 to 20 hours post
infection. Arabinose was added to PAO1Δ*exsA* condition at 3.5 hours post infection. Imaging was
conducted hourly from 4 to 20 hours. IFN-γ was maintained during infection and imaging.

602 **References**
603

1. Angus AA, Lee AA, Augustin DK, Lee EJ, Evans DJ, Fleiszig SM. 2008. Pseudomonas
aeruginosa induces membrane blebs in epithelial cells, which are utilized as a niche for
intracellular replication and motility. Infect Immun 76:1992-2001.
2. Jolly AL, Takawira D, Oke OO, Whiteside SA, Chang SW, Wen ER, Quach K, Evans DJ,
Fleiszig SM. 2015. Pseudomonas aeruginosa-induced bleb-niche formation in epithelial cells is
independent of actinomyosin contraction and enhanced by loss of cystic fibrosis transmembrane-
conductance regulator osmoregulatory function. mBio 6:e02533.
3. Kroken AR, Chen CK, Evans DJ, Yahr TL, Fleiszig SMJ. 2018. The Impact of ExoS on
Pseudomonas aeruginosa Internalization by Epithelial Cells Is Independent of fleQ and Correlates
with Bistability of Type Three Secretion System Gene Expression. mBio 9.
4. Fleiszig SM, Zaidi TS, Fletcher EL, Preston MJ, Pier GB. 1994. Pseudomonas aeruginosa
invades corneal epithelial cells during experimental infection. Infect Immun 62:3485-93.
5. Angus AA, Evans DJ, Barbieri JT, Fleiszig SM. 2010. The ADP-ribosylation domain of
Pseudomonas aeruginosa ExoS is required for membrane bleb niche formation and bacterial
survival within epithelial cells. Infect Immun 78:4500-10.
6. Sana TG, Baumann C, Merdes A, Soscia C, Rattei T, Hachani A, Jones C, Bennett KL, Filloux A,
Superti-Furga G, Voulhoux R, Bleves S. 2015. Internalization of Pseudomonas aeruginosa Strain
PAO1 into Epithelial Cells Is Promoted by Interaction of a T6SS Effector with the Microtubule
Network. mBio 6:e00712.
7. Garai P, Berry L, Moussouni M, Bleves S, Blanc-Potard AB. 2019. Killing from the inside:
Intracellular role of T3SS in the fate of Pseudomonas aeruginosa within macrophages revealed by
mgtC and oprF mutants. PLoS Pathog 15:e1007812.
8. Fleiszig SM, Zaidi TS, Preston MJ, Grout M, Evans DJ, Pier GB. 1996. Relationship between
cytotoxicity and corneal epithelial cell invasion by clinical isolates of Pseudomonas aeruginosa.
Infect Immun 64:2288-94.
9. Coburn J, Dillon ST, Iglewski BH, Gill DM. 1989. Exoenzyme S of Pseudomonas aeruginosa
ADP-ribosylates the intermediate filament protein vimentin. Infect Immun 57:996-8.
10. Riese MJ, Goehring UM, Ehrmantraut ME, Moss J, Barbieri JT, Aktories K, Schmidt G. 2002.
Auto-ADP-ribosylation of Pseudomonas aeruginosa ExoS. J Biol Chem 277:12082-8.
11. Goehring UM, Schmidt G, Pederson KJ, Aktories K, Barbieri JT. 1999. The N-terminal domain
of Pseudomonas aeruginosa exoenzyme S is a GTPase-activating protein for Rho GTPases. J Biol
Chem 274:36369-72.
12. Iglewski BH, Sadoff J, Bjorn MJ, Maxwell ES. 1978. Pseudomonas aeruginosa exoenzyme S: an
adenosine diphosphate ribosyltransferase distinct from toxin A. Proc Natl Acad Sci U S A
75:3211-5.
13. McGuffie EM, Frank DW, Vincent TS, Olson JC. 1998. Modification of Ras in eukaryotic cells
by Pseudomonas aeruginosa exoenzyme S. Infect Immun 66:2607-13.
14. Simon NC, Barbieri JT. 2014. Exoenzyme S ADP-ribosylates Rab5 effector sites to uncouple
intracellular trafficking. Infect Immun 82:21-8.
15. Maresso AW, Deng Q, Pereckas MS, Wakim BT, Barbieri JT. 2007. Pseudomonas aeruginosa
ExoS ADP-ribosyltransferase inhibits ERM phosphorylation. Cell Microbiol 9:97-105.
16. Deng Q, Barbieri JT. 2008. Modulation of host cell endocytosis by the type III cytotoxin,
Pseudomonas ExoS. Traffic 9:1948-57.
17. Rao L, De La Rosa I, Xu Y, Sha Y, Bhattacharya A, Holtzman MJ, Gilbert BE, Eissa NT. 2021.
Pseudomonas aeruginosa survives in epithelia by ExoS-mediated inhibition of autophagy and
mTOR. EMBO Rep 22:e50613.
18. Heimer SR, Evans DJ, Stern ME, Barbieri JT, Yahr T, Fleiszig SM. 2013. Pseudomonas
aeruginosa utilizes the type III secreted toxin ExoS to avoid acidified compartments within
epithelial cells. PLoS One 8:e73111.

19. Kroken AR, Gajenthra Kumar N, Yahr TL, Smith BE, Nieto V, Horneman H, Evans DJ, Fleiszig
SMJ. 2022. Exotoxin S secreted by internalized Pseudomonas aeruginosa delays lytic host cell
death. PLoS Pathog 18:e1010306.
20. Jorgensen I, Miao EA. 2015. Pyroptotic cell death defends against intracellular pathogens.
Immunol Rev 265:130-42.
21. Ulland TK, Ferguson PJ, Sutterwala FS. 2015. Evasion of inflammasome activation by microbial
pathogens. J Clin Invest 125:469-77.
22. Galluzzi L, Vitale I, Aaronson SA, Abrams JM, Adam D, Agostinis P, Alnemri ES, Altucci L,
Amelio I, Andrews DW, Annicchiarico-Petruzzelli M, Antonov AV, Arama E, Baehrecke EH,
Barlev NA, Bazan NG, Bernassola F, Bertrand MJM, Bianchi K, Blagosklonny MV, Blomgren
K, Borner C, Boya P, Brenner C, Campanella M, Candi E, Carmona-Gutierrez D, Cecconi F,
Chan FK, Chandel NS, Cheng EH, Chipuk JE, Cidlowski JA, Ciechanover A, Cohen GM,
Conrad M, Cubillos-Ruiz JR, Czabotar PE, D'Angiolella V, Dawson TM, Dawson VL, De
Laurenzi V, De Maria R, Debatin KM, DeBerardinis RJ, Deshmukh M, Di Daniele N, Di Virgilio
F, Dixit VM, Dixon SJ, et al. 2018. Molecular mechanisms of cell death: recommendations of the
Nomenclature Committee on Cell Death 2018. Cell Death Differ 25:486-541.
23. Cunha LD, Zamboni DS. 2013. Subversion of inflammasome activation and pyroptosis by
pathogenic bacteria. Front Cell Infect Microbiol 3:76.
24. Rayamajhi M, Zak DE, Chavarria-Smith J, Vance RE, Miao EA. 2013. Cutting edge: Mouse
NAIP1 detects the type III secretion system needle protein. J Immunol 191:3986-9.
25. Unterholzner L, Keating SE, Baran M, Horan KA, Jensen SB, Sharma S, Sirois CM, Jin T, Latz
E, Xiao TS, Fitzgerald KA, Paludan SR, Bowie AG. 2010. IFI16 is an innate immune sensor for
intracellular DNA. Nat Immunol 11:997-1004.
26. Shi J, Zhao Y, Wang Y, Gao W, Ding J, Li P, Hu L, Shao F. 2014. Inflammatory caspases are
innate immune receptors for intracellular LPS. Nature 514:187-92.
27. Martinon F, Agostini L, Meylan E, Tschopp J. 2004. Identification of bacterial muramyl
dipeptide as activator of the NALP3/cryopyrin inflammasome. Curr Biol 14:1929-34.
28. Sandstrom A, Mitchell PS, Goers L, Mu EW, Lesser CF, Vance RE. 2019. Functional
degradation: A mechanism of NLRP1 inflammasome activation by diverse pathogen enzymes.
Science 364.
29. Xu H, Yang J, Gao W, Li L, Li P, Zhang L, Gong YN, Peng X, Xi JJ, Chen S, Wang F, Shao F.
2014. Innate immune sensing of bacterial modifications of Rho GTPases by the Pyrin
inflammasome. Nature 513:237-41.
30. Poyet JL, Srinivasula SM, Tnani M, Razmara M, Fernandes-Alnemri T, Alnemri ES. 2001.
Identification of Ipaf, a human caspase-1-activating protein related to Apaf-1. J Biol Chem
276:28309-13.
31. Martinon F, Burns K, Tschopp J. 2002. The inflammasome: a molecular platform triggering
activation of inflammatory caspases and processing of proIL-beta. Molecular Cell 10:417-426.
32. Kayagaki N, Warming S, Lamkanfi M, Vande Walle L, Louie S, Dong J, Newton K, Qu Y, Liu J,
Heldens S, Zhang J, Lee WP, Roose-Girma M, Dixit VM. 2011. Non-canonical inflammasome
activation targets caspase-11. Nature 479:117-21.
33. Vigano E, Diamond CE, Spreafico R, Balachander A, Sobota RM, Mortellaro A. 2015. Human
caspase-4 and caspase-5 regulate the one-step non-canonical inflammasome activation in
monocytes. Nat Commun 6:8761.
34. Downs KP, Nguyen H, Dorfleutner A, Stehlik C. 2020. An overview of the non-canonical
inflammasome. Mol Aspects Med 76:100924.
35. He WT, Wan H, Hu L, Chen P, Wang X, Huang Z, Yang ZH, Zhong CQ, Han J. 2015.
Gasdermin D is an executor of pyroptosis and required for interleukin-1beta secretion. Cell Res
25:1285-98.
36. Kayagaki N, Stowe IB, Lee BL, O'Rourke K, Anderson K, Warming S, Cuellar T, Haley B,
Roose-Girma M, Phung QT, Liu PS, Lill JR, Li H, Wu J, Kummerfeld S, Zhang J, Lee WP,

704   Snipas SJ, Salvesen GS, Morris LX, Fitzgerald L, Zhang Y, Bertram EM, Goodnow CC, Dixit
705   VM. 2015. Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. Nature
706   526:666-71.

707 37.   Ding J, Wang K, Liu W, She Y, Sun Q, Shi J, Sun H, Wang DC, Shao F. 2016. Pore-forming
708   activity and structural autoinhibition of the gasdermin family. Nature 535:111-6.

709 38.   Karmakar M, Minns M, Greenberg EN, Diaz-Aponte J, Pestonjamasp K, Johnson JL, Rathkey
710   JK, Abbott DW, Wang K, Shao F, Catz SD, Dubyak GR, Pearlman E. 2020. N-GSDMD
711   trafficking to neutrophil organelles facilitates IL-1beta release independently of plasma
712   membrane pores and pyroptosis. Nat Commun 11:2212.

713 39.   DiPeso L, Ji DX, Vance RE, Price JV. 2017. Cell death and cell lysis are separable events during
714   pyroptosis. Cell Death Discov 3:17070.

715 40.   Galle M, Schotte P, Haegman M, Wullaert A, Yang HJ, Jin S, Beyaert R. 2008. The
716   Pseudomonas aeruginosa Type III secretion system plays a dual role in the regulation of caspase-
717   1 mediated IL-1beta maturation. J Cell Mol Med 12:1767-76.

718 41.   Mohamed MF, Gupta K, Goldufsky JW, Roy R, Callaghan LT, Wetzel DM, Kuzel TM, Reiser J,
719   Shafikhani SH. 2022. CrkII/Abl phosphorylation cascade is critical for NLRC4 inflammasome
720   activity and is blocked by Pseudomonas aeruginosa ExoT. Nat Commun 13:1295.

721 42.   Ince D, Sutterwala FS, Yahr TL. 2015. Secretion of Flagellar Proteins by the Pseudomonas
722   aeruginosa Type III Secretion-Injectisome System. J Bacteriol 197:2003-11.

723 43.   Churchill MJ, Mitchell PS, Rauch I. 2022. Epithelial Pyroptosis in Host Defense. J Mol Biol
724   434:167278.

725 44.   Lian H, Fang X, Li Q, Liu S, Wei Q, Hua X, Li W, Liao C, Yuan X. 2022. NLRP3
726   Inflammasome-Mediated Pyroptosis Pathway Contributes to the Pathogenesis of Candida
727   albicans Keratitis. Front Med (Lausanne) 9:845129.

728 45.   Zheng Q, Tan Q, Ren Y, Reinach PS, Li L, Ge C, Qu J, Chen W. 2018. Hyperosmotic Stress-
729   Induced TRPM2 Channel Activation Stimulates NLRP3 Inflammasome Activity in Primary
730   Human Corneal Epithelial Cells. Invest Ophthalmol Vis Sci 59:3259-3268.

731 46.   Kelley N, Jeltema D, Duan Y, He Y. 2019. The NLRP3 Inflammasome: An Overview of
732   Mechanisms of Activation and Regulation. Int J Mol Sci 20.

733 47.   Mariathasan S, Weiss DS, Newton K, McBride J, O'Rourke K, Roose-Girma M, Lee WP,
734   Weinrauch Y, Monack DM, Dixit VM. 2006. Cryopyrin activates the inflammasome in response
735   to toxins and ATP. Nature 440:228-32.

736 48.   Hornung V, Bauernfeind F, Halle A, Samstad EO, Kono H, Rock KL, Fitzgerald KA, Latz E.
737   2008. Silica crystals and aluminum salts activate the NALP3 inflammasome through phagosomal
738   destabilization. Nature Immunology 9:847-856.

739 49.   Kanneganti TD, Ozoren N, Body-Malapel M, Amer A, Park JH, Franchi L, Whitfield J, Barchet
740   W, Colonna M, Vandenabeele P, Bertin J, Coyle A, Grant EP, Akira S, Nunez G. 2006. Bacterial
741   RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3. Nature
742   440:233-6.

743 50.   Kayagaki N, Wong MT, Stowe IB, Ramani SR, Gonzalez LC, Akashi-Takamura S, Miyake K,
744   Zhang J, Lee WP, Muszynski A, Forsberg LS, Carlson RW, Dixit VM. 2013. Noncanonical
745   inflammasome activation by intracellular LPS independent of TLR4. Science 341:1246-9.

746 51.   Bauernfried S, Scherr MJ, Pichlmair A, Duderstadt KE, Hornung V. 2021. Human NLRP1 is a
747   sensor for double-stranded RNA. Science 371.

748 52.   Griswold AR, Huang HC, Bachovchin DA. 2022. The NLRP1 Inflammasome Induces Pyroptosis
749   in Human Corneal Epithelial Cells. Invest Ophthalmol Vis Sci 63:2.

750 53.   Xu S, Liu X, Liu X, Shi Y, Jin X, Zhang N, Li X, Zhang H. 2021. Wedelolactone ameliorates
751   Pseudomonas aeruginosa-induced inflammation and corneal injury by suppressing caspase-
752   4/5/11/GSDMD-mediated non-canonical pyroptosis. Exp Eye Res 211:108750.

54. Robertson DM, Li L, Fisher S, Pearce VP, Shay JW, Wright WE, Cavanagh HD, Jester JV. 2005.
Characterization of growth and differentiation in a telomerase-immortalized human corneal
epithelial cell line. Invest Ophthalmol Vis Sci 46:470-8.

55. Bitto NJ, Baker PJ, Dowling JK, Wray-McCann G, De Paoli A, Tran LS, Leung PL, Stacey KJ,
Mansell A, Masters SL, Ferrero RL. 2018. Membrane vesicles from Pseudomonas aeruginosa
activate the noncanonical inflammasome through caspase-5 in human monocytes. Immunol Cell
Biol 96:1120-1130.

56. Reyes Ruiz VM, Ramirez J, Naseer N, Palacio NM, Siddarthan IJ, Yan BM, Boyer MA,
Pensinger DA, Sauer JD, Shin S. 2017. Broad detection of bacterial type III secretion system and
flagellin proteins by the human NAIP/NLRC4 inflammasome. Proc Natl Acad Sci U S A
114:13242-13247.

57. Thul PJ, Akesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, Alm T, Asplund A,
Bjork L, Breckels LM, Backstrom A, Danielsson F, Fagerberg L, Fall J, Gatto L, Gnann C, Hober
S, Hjelmare M, Johansson F, Lee S, Lindskog C, Mulder J, Mulvey CM, Nilsson P, Oksvold P,
Rockberg J, Schutten R, Schwenk JM, Sivertsson A, Sjostedt E, Skogs M, Stadler C, Sullivan
DP, Tegel H, Winsnes C, Zhang C, Zwahlen M, Mardinoglu A, Ponten F, von Feilitzen K, Lilley
KS, Uhlen M, Lundberg E. 2017. A subcellular map of the human proteome. Science 356.

58. Kumar NG, Nieto V, Kroken AR, Jedel E, Grosser MR, Hallsten ME, Mettrucio MME, Yahr TL,
Evans DJ, Fleiszig SMJ. 2022. Pseudomonas aeruginosa Can Diversify after Host Cell Invasion
to Establish Multiple Intracellular Niches. mBio 13:e0274222.

59. Kaufman MR, Jia J, Zeng L, Ha U, Chow M, Jin S. 2000. Pseudomonas aeruginosa mediated
apoptosis requires the ADP-ribosylating activity of exoS. Microbiology (Reading) 146 (Pt
10):2531-2541.

60. Vallis AJ, Finck-Barbancon V, Yahr TL, Frank DW. 1999. Biological effects of Pseudomonas
aeruginosa type III-secreted proteins on CHO cells. Infect Immun 67:2040-4.

61. Santos JC, Boucher D, Schneider LK, Demarco B, Dilucca M, Shkarina K, Heilig R, Chen KW,
Lim RYH, Broz P. 2020. Human GBP1 binds LPS to initiate assembly of a caspase-4 activating
platform on cytosolic bacteria. Nat Commun 11:3276.

62. Kenny M, Schoen I. 2021. Violin SuperPlots: visualizing replicate heterogeneity in large data
sets. Mol Biol Cell 32:1333-1334.

63. Zamyatina A, Heine H. 2020. Lipopolysaccharide Recognition in the Crossroads of TLR4 and
Caspase-4/11 Mediated Inflammatory Pathways. Front Immunol 11:585146.

64. Kobayashi T, Ogawa M, Sanada T, Mimuro H, Kim M, Ashida H, Akakura R, Yoshida M,
Kawalec M, Reichhart JM, Mizushima T, Sasakawa C. 2013. The Shigella OspC3 effector
inhibits caspase-4, antagonizes inflammatory cell death, and promotes epithelial infection. Cell
Host Microbe 13:570-583.

65. Li Z, Liu W, Fu J, Cheng S, Xu Y, Wang Z, Liu X, Shi X, Liu Y, Qi X, Liu X, Ding J, Shao F.
2021. Shigella evades pyroptosis by arginine ADP-riboxanation of caspase-11. Nature
doi:10.1038/s41586-021-04020-1.

66. Goers L, Kim K, Canning PJ, Stedman TC, Mou X, Ernst NH, Coers J, Lesser CF. 2022. *Shigella*
IpaH9.8 limits GBP1-dependent LPS release from intracytosolic bacteria to suppress caspase-4
activation. bioRxiv doi:10.1101/2022.11.14.516511:2022.11.14.516511.

67. Cowell BA, Chen DY, Frank DW, Vallis AJ, Fleiszig SM. 2000. ExoT of cytotoxic Pseudomonas
aeruginosa prevents uptake by corneal epithelial cells. Infect Immun 68:403-6.

68. Garrity-Ryan L, Kazmierczak B, Kowal R, Comolli J, Hauser A, Engel JN. 2000. The arginine
finger domain of ExoT contributes to actin cytoskeleton disruption and inhibition of
internalization of Pseudomonas aeruginosa by epithelial cells and macrophages. Infect Immun
68:7100-13.

69. Knodler LA, Nair V, Steele-Mortimer O. 2014. Quantitative assessment of cytosolic Salmonella
in epithelial cells. PLoS One 9:e84681.

70.   Alaoui-El-Azher M, Jia J, Lian W, Jin S. 2006. ExoS of Pseudomonas aeruginosa induces
apoptosis through a Fas receptor/caspase 8-independent pathway in HeLa cells. Cell Microbiol
8:326-38.
71.   Jia J, Alaoui-El-Azher M, Chow M, Chambers TC, Baker H, Jin S. 2003. c-Jun NH2-terminal
kinase-mediated signaling is essential for Pseudomonas aeruginosa ExoS-induced apoptosis.
Infect Immun 71:3361-70.
72.   Jia J, Wang Y, Zhou L, Jin S. 2006. Expression of Pseudomonas aeruginosa toxin ExoS
effectively induces apoptosis in host cells. Infect Immun 74:6557-70.
73.   Taabazuing CY, Okondo MC, Bachovchin DA. 2017. Pyroptosis and Apoptosis Pathways
Engage in Bidirectional Crosstalk in Monocytes and Macrophages. Cell Chem Biol 24:507-514
e4.
74.   Luchetti G, Roncaioli JL, Chavez RA, Schubert AF, Kofoed EM, Reja R, Cheung TK, Liang Y,
Webster JD, Lehoux I, Skippington E, Reeder J, Haley B, Tan MW, Rose CM, Newton K,
Kayagaki N, Vance RE, Dixit VM. 2021. Shigella ubiquitin ligase IpaH7.8 targets gasdermin D
for degradation to prevent pyroptosis and enable infection. Cell Host Microbe
doi:10.1016/j.chom.2021.08.010.
75.   Li P, Jiang W, Yu Q, Liu W, Zhou P, Li J, Xu J, Xu B, Wang F, Shao F. 2017. Ubiquitination and
degradation of GBPs by a Shigella effector to suppress host defence. Nature 551:378-383.
76.   Barbieri JT, Sun J. 2004. Pseudomonas aeruginosa ExoS and ExoT. Rev Physiol Biochem
Pharmacol 152:79-92.
77.   Sun Y, Karmakar M, Taylor PR, Rietsch A, Pearlman E. 2012. ExoS and ExoT ADP
ribosyltransferase activities mediate Pseudomonas aeruginosa keratitis by promoting neutrophil
apoptosis and bacterial survival. J Immunol 188:1884-95.
78.   Urbanowski ML, Brutinel ED, Yahr TL. 2007. Translocation of ExsE into Chinese hamster ovary
cells is required for transcriptional induction of the Pseudomonas aeruginosa type III secretion
system. Infect Immun 75:4432-9.
79.   Sanjana NE, Shalem O, Zhang F. 2014. Improved vectors and genome-wide libraries for CRISPR
screening. Nat Methods 11:783-784.
80.   Schmidt T, Schmid-Burgk JL, Hornung V. 2015. Synthesis of an arrayed sgRNA library targeting
the human genome. Sci Rep 5:14987.

Figure 1

Figure 1. Host cell death rates from *P. aeruginosa* mutant infections. **(A)** Corneal epithelial cells (hTCEpi) were infected with indicated strain: wt PAO1, PAO1Δ*exoSTY*, or PAO1Δ*exsA* for 3 hours at an MOI equal to 10. Hoechst (blue) was added to label nuclei. Non-associated bacteria were removed, and media with amikacin and propidium iodide (red) was added to kill extracellular bacteria. Cells were imaged hourly from 4 to 20 hours post infection. Select fields from indicated times are shown. Scale bar = 50 μm. **(B)** The percent of cells positive for propidium iodide at each timepoint was determined by using a custom FIJI macro to segment all nuclei from the Hoechst channel, and all nuclei of dead cells in the propidium iodide channel. Multiple column t-tests were performed comparing wt PAO1 and PAO1Δ*exsA*, and SD error bars are displayed from three replicates. **(C)** Experiment was performed identically as in panel A, but using HeLa cells. Scale bar = 50 μm. **(D)** Analysis was performed as in panel B, using the data obtained in panel C.

Figure 2

A PAO1Δ*exsA*, pBAD-GFP

Hoechst / PI / GFP / DIC

4 hpi 5 hpi 6 hpi 7 hpi

8 hpi 9 hpi 10 hpi 11 hpi

inset 1 inset 2 **B** **C**

Hoechst / PI / GFP

6 hpi 11 hpi 20 hpi 20 hpi

D PAO1, T3SS reporter

Hoechst / PI / GFP / DIC

4 hpi 5 hpi 6 hpi 7 hpi

PAO1Δ*exoSTY*, T3SS reporter

4 hpi 5 hpi 6 hpi 7 hpi

E

ns

*** ***

Survival time of invaded cells (hours)

PAO1 Δ*STY* Δ*exsA*

Figure 2. Intracellular Δ*exsA* mutants kill invaded cells at a similar rate to wt PAO1-invaded cells. (A) Bacteria were transformed with an arabinose-inducible GFP plasmid. Corneal epithelial cells (hTCEpi) were infected at an MOI of 10 and extracellular bacteria eliminated at 3 hours post infection using amikacin. A final concentration of 1% arabinose was used to induce GFP in surviving intracellular bacteria beginning at 3.5 hours post infection. Time lapse imaging was conducted hourly from 4 to 20 hours, and host cell nuclei detected with Hoechst and propidium iodide to determine time of death. Scale bar = 50 µm. Insets 1-2 show the boxed cell at 6 or 11 hours post infection, respectively. **(B)** An example of a live invaded cell from 20 hours post infection. **(C)** Example of a dead invaded cell from 20 hours post infection, where the PI channel has been saturated such that PI-positive bacteria can be visualized. Scale bar = 50 µm. **(D)** The T3SS-GFP reporter pJNE05 was used to visualize wt PAO1 and PAO1Δ*exoSTY* infections, as described previously (19), to compare to PAO1Δ*exsA* infections. Scale bar = 50 µm. Time lapses from panels A through D are available as **Supplemental Movie 2**. **(E)** A computational analysis approach was used to segregate only invaded cells, and determine their survival time in hours. PAO1 condition includes 881 cells over six replicates; PAO1Δ*exoSTY*, 967 cells; PAO1Δ*exsA*, 862 cells. The survival times of all invaded cells were combined a single super violin plot. Mean survival times with SD error bars are displayed, and significance determined by One-way ANOVA. *** $P < 0.005$.

Figure 3

Figure 3. Transcriptional responses of infected host cells. (A) Corneal epithelial cells (hTCEpi) were infected with wt PAO1, PAO1Δ*exoSTY*, or PAO1Δ*exsA* for 3 hours at an MOI equal to 10. Non-associated bacteria were removed, and media with amikacin was added for 1 additional hour. At four hours, RNA was purified from infected cells, and RT PCR analysis performed using a commercial array for genes associated with specific cell death pathways. **(B)** Experiment performed identically to panel A, except using HeLa cells.

Figure 4.

Figure 4. NF-κB signaling and the role of secreted factors in cell death. (A) Corneal epithelial cells (hTCEpi) were infected with wt PAO1, PAO1Δ*exoSTY*, or PAO1Δ*exsA* for 3 hours at an MOI equal to 10. Then media was replaced with amikacin-containing media. Cells were fixed at 4 hours post infection and p65 localization was determined using immunofluorescent staining. Nuclei were labeled with DAPI. Scale bar = 50 μm. **(B)** Experiment performed identically to panel A, using HeLa cells. **(C)** Quantification of p65-positive nuclei from corneal epithelial cells, and **(D)** HeLa cells. PAO1-infected HeLa cells were not able to be accurately quantified (indicated using "ND") by automated analysis due to the small and irregular cytoplasmic area. **(E)** Corneal epithelial cells were infected as described in panel A. Supernatant of infected cells was collected at 4 hours post infection and TNFα measured by ELISA. **(D)** Supernatants of corneal epithelial cells infected with PAO1Δ*exoSTY*, PAO1Δ*exsA*, or uninfected were collected at 4 hours post infection and sterilized with 0.22 μm filter, and used as the replacement media on uninfected cells or PAO1Δ*exsA* infected cells at 3 hours post infection, combined with amikacin and propidium iodide. Hoechst was used to label all nuclei. The cell death rates were measured by time lapse imaging, error bars display SD from three replicates.

Figure 5

Figure 5. Caspase-4 limits intracellular replication and persistence of T3SS mutant bacteria. (A) Corneal epithelial cells (hTCEpi) or (B) cells knocked out for caspase-4 were infected with wt PAO1, PAO1Δ*exoSTY* (each visualized by T3SS-GFP reporter) or PAO1Δ*exsA* (pBAD-GFP induced) at an MOI of 10. Nuclei were labeled with Hoechst and propidium iodide. Extracellular bacteria were eliminated with amikacin at 3 hours post infection, and imaged hourly from 4 to 20 hours post infection. Images from 4-9 hours post infection are shown. Scale bar = 50 μm. Full-field time lapses are available as **Supplemental Movie 3**. (**B-D**) (C-D) Cells from indicated infections were analyzed with a computational approach to segregate only invaded cells and measure survival times in hours. Six replicates were combined into a single super violin plot. Mean survival times with SD error bars are displayed, and significance determined by Student's t-test. (F-H) GFP area was summed within the boundaries of live cells using the same data set as panels C-E, and normalized to PAO1 (wt) at 4 hours post infection.

Figure 6

Figure 6. IFN-γ-stimulated HeLa cells are not permissive for PAO1Δ*exoSTY* intracellular replication nor PAO1Δ*exsA* intracellular persistence. (A) HeLa cells were treated with 50 ng/ml IFN-γ for 16 hours prior to infection with wt PAO1, PAO1Δ*exoSTY* (each using the T3SS-GFP reporter) or PAO1Δ*exsA* (pBAD-GFP induced) at an MOI of 10. Nuclei were labeled with Hoechst and propidium iodide. Extracellular bacteria were eliminated with amikacin at 3 hours post infection, and imaged hourly from 4 to 20 hours post infection. Select time frames are shown at either 10 or 15 hours post infection. Scale bar = 50 μm. Full-field time lapses are available as **Supplemental Movie 4**. **(B)** The rate of cell death was determined using host cell nuclear stains. All panels are from the same experiment with only indicated bacterial strain shown. Error bars show SD from three replicates. Multiple column t-tests were performed comparing IFN-γ-treated cells to untreated.

Supplemental Figure 1

Supplemental Figure 1. GFP expression from plasmid pBAD-GFP suppresses the T3SS. (A) wt PAO1 was transformed with the T3SS reporter plasmid pJNE05, or the GFP inducible plasmid pBAD-GFP and imaged hourly in time lapse. Example fields are shown in which bacteria expressing GFP under the T3SS reporter exhibit intracellular spread and replication, whereas those with inducible GFP show only limited intracellular replication. (B) The cell death rates of wt PAO1 or PAO1Δ*exoSTY* expressing GFP by either plasmid were measured, showing a trend of reduced cell killing rates for PAO1Δ*exoSTY* when GFP expression is induced. (C) wt PAO1 with indicated plasmid was grown for 5 hours in LB media with EGTA to induce the T3SS, and arabinose to induce GFP. Supernatant proteins were concentrated by TCA precipitation, and the bands constituting ExoS and ExoT are indicated.

Supplemental Figure 2

Supplemental Figure 2. Confirmation of caspase-4 deletion. Lysates from wt hTCEpi cells or caspase-4 candidate knockout cells were collected in RIPA buffer and clarified. Total protein was visualized using Bio-Rad stain free gels prior to transfer. Blot was probed with anti-caspase-4 antibody.

Supplemental Figure 3. Control monoclonal cells for CRISPR-Cas9 mutagenesis. A control cell line was generated using non-targeting guide RNA, and monoclonal line generated. Cells were infected with wt PAO1, PAO1Δ*exoSTY* (each using the T3SS-GFP reporter) or PAO1Δ*exsA* (pBAD-GFP induced) at an MOI of 10. Nuclei were labeled with Hoechst and Propidium iodide. Extracellular bacteria were eliminated with amikacin at 3 hours post infection, and imaged hourly from 4 to 20 hours post infection. Images were analyzed with a computational approach to segregate invaded cells and measure survival times in hours. Three replicates were combined into a single super violin plot. Mean survival times with SD error bars are displayed.

Supplemental Figure 4

Supplemental Figure 4. Host cell survival time and intracellular replication of single exotoxin deletion of ExoS. (A) Corneal epithelial cells or cells lacking caspase-4 were infected with wt PAO1, PAO1Δ*exoSTY*, or PAO1Δ*exoS* (each visualized by T3SS-GFP reporter) at an MOI of 10. Nuclei were labeled with Hoechst and propidium iodide. Extracellular bacteria were eliminated with amikacin at 3 hours post infection, and imaged hourly from 4 to 20 hours post infection. Images from 4-10 hours are shown. Scale bar = 50 μm. (B) GFP area was summed within the boundaries of live cells and normalized to PAO1 (wt) at 4 hours post infection. (C) Cells from indicated infections were analyzed with a computational approach to segregate only invaded cells and measure survival times in hours. Mean survival times with SD error bars are displayed, and significance determined by Student's t-test.