Repeated *Plasmodium vivax* blood stage infection provides sterile protection against

homologous challenge in non-human primates

[1,2,3]Nicanor Obaldía III*, [3,4]Joao Luiz Da Silva Filho, [1]Marlon Núñez, [5]Katherine A. Glass,

[5]Tate Oulton, [3,4]Fiona Achcar, [6]Grennady Wirjanata, [2]Manoj Duraisingh, [7]Philip Felgner,

[5]Kevin K.A. Tetteh, [6]Zbynek Bozdech, [3]Thomas D. Otto, [2,3,4]Matthias Marti*

[1]Departamento de Investigaciones en Parasitologia, Instituto Conmemorativo Gorgas de

Estudios de la Salud, Panamá City, Republic of Panamá.

[2]Department of Immunology and Infectious Diseases, Harvard TH Chan School of Public

Health, Harvard University, Boston, MA, United States.

[3]Wellcome Centre for Integrative Parasitology, School of Infection & Immunity, College of

Medical, Veterinary and Life Sciences. University of Glasgow, Glasgow, UK.

[4]Institute of Parasitology, VetSuisse and Medical Faculty, University of Zurich, Zurich,

Switzerland

[5]Department of Immunology and Infection, London School of Hygiene and Tropical Medicine,

London, UK

[6]School of Biological Sciences, Nanyang Technological University, Singapore.

[7]Institute for Immunology, University of California, Irvine, CA, USA.

Address correspondence to:

* nobaldia@gorgas.gob.pa (NO); Matthias.Marti@uzh.ch (MM)

Abstract

The malaria parasite *Plasmodium vivax* remains a major global public health challenge, causing major morbidity across tropical and subtropical regions. Several candidate vaccines are in preclinical and clinical trials, however no vaccine against *P. vivax* malaria is approved for use in humans. Here we assessed whether *P. vivax* strain-transcendent immunity can be achieved by repeated infection in *Aotus* monkeys. For this purpose, we repeatedly infected six animals with blood stages of the *P. vivax* Salvador 1 (SAL-1) strain until sterile immune, and then challenged with the AMRU-1 strain. Sterile immunity was achieved in 4/4 (100%) *Aotus* monkeys after two homologous infections with the SAL-1 strain, while partial protection against a heterologous AMRU-1 challenge was achieved in 3/3 (100%) monkeys. IgG levels based on *P. vivax* lysate ELISA and protein microarray increased with repeated infections and correlated with the level of homologous protection. Analysis of parasite transcriptional profiles across inoculation levels provided no evidence of major antigenic switching upon homologous or heterologous challenge. In contrast, we observed significant transcriptional differences in the *P. vivax* core gene repertoire between SAL-1 and AMRU-1. Together with the strain-specific genetic diversity between SAL-1 and AMRU-1 these data suggest that the protection upon heterologous challenge is due to strain differences (at genome and transcriptome level) rather than immune evasion by antigenic switching. Our study provides the first benchmark for achieving sterile immunity in the *Aotus*/*P. vivax* non-human primate model, and a template for establishing correlates of immunity to test the efficacy of candidate blood stage *P. vivax* malaria vaccines.

Author summary

*Plasmodium vi*vax is the most widespread human malaria parasite. Elimination efforts are complicated due to the peculiar biology of *P. vivax* including dormant liver forms, cryptic reservoirs in bone marrow and spleen and a large asymptomatic infectious reservoir in affected populations. Currently there is no vaccine against malaria caused by *P. viv*ax. Here we induce sterile immunity by repeated *P. vivax* infection with the SAL-1 strain in non-human primates. In contrast, heterologous challenge with the AMRU-1 strain only provided partial protection. Antibody levels against a crude antigen and a protein microarray correlated with the level of homologous protection. Parasite transcriptional profiles across inoculation levels failed to show major antigenic switching across SAL-1 infections or upon heterologous challenge, instead suggesting other mechanisms of immune evasion. Our study provides the first benchmark for achieving sterile immunity in the *Aotus*/*P. vivax* non-human primate model, and a template for establishing correlates of immunity to test the efficacy of candidate blood stage *P. vivax* malaria vaccines.

Introduction

Malaria is caused by parasites of the genus *Plasmodium* that are transmitted to humans by the bite of the female anopheles mosquito. Currently, approximately 241 million cases and 0.6 million deaths from malaria occur worldwide, an increase of 12% from the previous year [1]. Most deaths are due to infection with *Plasmodium falciparum*, the most pathogenic of the species, especially in children under the age of five living in sub-Saharan Africa [1-3].

After the elimination of *P. falciparum*, *Plasmodium vivax* is expected to remain a major cause of morbidity and mortality outside of Africa, especially in Central and South America, Asia, and the Pacific Islands [4-6]. This is due in part to its peculiar biology, including silent parasite liver forms known as hypnozoites that can cause relapses and major

parasite reservoirs in bone marrow and spleen that may act as an unobserved pathogenic

biomass and source for recrudescence [7-13]. Complete removal of the parasite from the

human reservoir is therefore challenging [5, 14], underscoring the need for innovative

therapeutic strategies including the development of an effective vaccine [15, 16].

Vivax malaria impacts the health of individuals of all ages causing repeated febrile

episodes and severe anemia [15, 17], clinical severity including hemolytic coagulation

disorders, jaundice, nephropathology, rhabdomyolysis, porphyria, cerebral malaria, splenic

rupture [4], and Acute Respiratory Distress Syndrome (ARDS)[18-20]. Fatal *P. vivax* cases

are reported from all endemic regions across the globe [3, 15, 21]. Compounding the

epidemiology of the disease, *P vivax* malaria transmission is intermittent and acquired

immunity is short and strain-specific [15]. Even in low transmission regions, it is common to

find individuals with asymptomatic parasitemia suggestive of natural premunition – a

phenomenon resulting from a delicate host-parasite equilibrium in individuals with acquired

immunity [15, 22]. Epidemiological studies have demonstrated that repeated exposure

increases clinical immunity and decreases parasite density and frequency of clinical episodes.

For example, individuals subjected to malariotherapy with *P. vivax* for treatment of

neurosyphilis rapidly developed immunity after repeated blood stage infections [8, 23-25],

and such repeated infection provided strain transcending protection [26]. The development of

a vaccine against malaria with at least 75% protective efficacy is one of the two main

objectives identified in the roadmap adopted by the global vaccine action plan until 2030

[27]. Such an effective *P. vivax* vaccine should provide long-term and strain-transcending

immunity. Current *P. vivax* vaccine studies are focused at inducing a stronger antibody

response in combination with an already robust T-cell response [28]. To date, there is no

vaccine against *P. vivax* approved for use in humans [15]. Immunization by repeated

infection with live blood stage parasites has been used in humans [29, 30] and non-human

primates against a *P. falciparum* heterologous challenge [31, 32]. The same approach also

produced heterologous cross-protection against *Plasmodium chabaudi* infection in mice [33].

However, such an approach has not been applied to *P. vivax* to the best of our knowledge.

To assess whether strain-transcendent immunity can be achieved by repeated blood

stage infection in *P. vivax*, we used the *Aotus* non-human primate model. The aims of our

study were to determine i) how many repeated homologous infections are required for control

of parasitemia and development of sterile immunity, and ii) whether strain-transcending

immunity could be achieved. Our study provides important insights into protective immune

responses and evasion mechanisms and contributes to the design of a whole organism

erythrocytic vaccine for *P. vivax*.

Results

***P. vivax* blood stage infection induces sterile immunity to homologous challenge**

To evaluate the level of protection against repeated *P. vivax* blood stage infection, six *Aotus*

monkeys (MN30014, MN30034, MN32028, MN32047, MN25029, MN29012) were infected

intravenously with 50,000 parasites of the *P. vivax* SAL-1 strain and monitored until peak

parasitemia (**Figure 1A and S1**). SAL-1 strain was originally isolated from a patient in El

Salvador in the late 1960s and adapted to *Aotus* monkeys by W.E. Collins [34]. During the

first infection, all six animals were positive by day 6 post inoculation (PI) and parasitemia

increased steadily to more than $100 \times 10^3/\mu L$ infected red blood cells (iRBCs)/µL (mean ± sd

$= 100,198 \pm 43,661/\mu L$) until days 13-14 PI, when the animals were treated with a curative

course of chloroquine (CQ) (**Figure 1B**). Sixty-five days PI, one animal (MN29012) was

removed from the study due to malaria unrelated causes (**Figure 1A and S1**).

Eighty-five days PI the remaining five animals (MN30014, MN30034, MN32028,

MN32047, MN25029) and the donor from the first inoculation (MN29041) were infected

with SAL-1 using the same inoculum size of 50,000 parasites i.v. This time, by day 91 (day 6

PI of inoculation level II, D6 PI II), all animals were positive by blood smear but parasitemia

remained low with a mean peak of 2,332/µL between days 94-95 (D9-10 PI II) (**Figure 1B).**

A similar pattern was observed when total parasite load was measured by qPCR (18s rRNA

levels) and parasite biomass by ELISA (pLDH levels) after this second inoculation (**Figure**

S2A, B). Two animals (MN32047 and MN30034) self-cured on day 98 (D13 PI II) and 102

(D17 PI II) respectively, and a third animal (MN30014) became negative for two days

between days 98-99 (D13-14 PI II) but recrudesced on day 100 (D15 PI II) and was treated

with CQ on day 105 (D20 PI II) while still positive at the level of <10 parasites /µL.

Meanwhile, MN29041 that had controlled its parasitemia until day 98 (D13 PI II), became

negative on day 99 (D14 PI II), but recrudesced the next day, reaching a parasitemia level of

11,500/µL on day 105 (D20 PI II) when it was treated with CQ [35]. All animals received CQ

treatment on day 105 (D20 PI II). Two animals (MN30034 and MN29041) were excluded

following CQ treatment - MN30034 on day 169 (D114 PI II) due to severe anemia (Hct% =

20) and kindney failure, and MN29041 for malaria unrelated causes on day 143 (D58 PI II).

On day 166 the remaining 4 original animals (MN30014, MN32028, MN32047 and

MN25029) plus a malaria naïve infection control (MN32029), were re-inoculated a third time

with SAL-1 and followed up as described above (**Figure 1B**). This time, all animals except

for the control (MN32029) that had a peak parasitemia of 95,550/µL on day 179 (D13 PI III)

remained negative and did not require CQ treatment. Of note, MN32028 had to be removed

from the experiment on day 254 (D88 PI III) due to anemia and kidney failure (**Figure 1A**).

Altogether, these experiments demonstrate that repeated homologous *P. vivax*

infection confers full protection (or sterile immunity) against a homologous challenge.

Partial protection to heterologous challenge after repeated homologous infections

To determine the difference in protection between homologous and heterologous infections,

we challenged on experimental day 276 the three remaining monkeys that went through three

SAL-1 inoculations (MN30014, MN32047 and MN25029) plus a new malaria naïve infection

control (MN31029) and the donor of the second SAL-1 infection (MN27050) with the CQ

resistant AMRU-1 strain (**Figure 1A, B**). The AMRU-I strain was originally isolated from a

patient in Papua New Guinea in 1989 [36].

This time all animals became positive. First, the 2 controls (MN27050 and MN31029)

were positive on day 283 (D7 PI IV) with peak parasitemia of 131.5 x 10^3/µL and 180 x

10^3/µL on day 290 (D14 PI IV), respectively when they were treated with MQ. Meanwhile,

MN25029 became positive four days later on day 287 (D11 PI IV) with a lower (10-fold)

peak parasitemia of 11.4 x 10^3/µL on day 290 (D14 PI IV), clearing on day 296 (D20 PI IV)

and treated with CQ on day 297 (D21 PI IV). Similarly, MN30014 became positive on day

295 (D19 PI IV) with a 100-fold lower peak parasitemia of 1,700/µL on day 297 (D21 PI IV)

compared to the peak parasitemia of the controls. The animal was treated with MQ on day

304 (D28 PI IV) for moderate anemia (Hct% = 27.4) and thrombocytopenia (PLT = 54 x

10^3/µL), while still positive at 1,510 parasites /µL. In contrast, MN32047 was positive only

once on day 292 (D16 PI IV) with less than 10 parasites /µL and was treated with MQ for

severe anemia (Hct% = 16) on day 297 (D21 PI IV).

Altogether, this experiment revealed partial protection in 3/3 (100%) of the monkeys

to a heterologous *P. vivax* challenge in sterile homologous immune animals. Partial

protection was characterized by a delay of 4-12 days in patency and reduced parasitemia

compared to the controls and a delay of 5-13 days in patency compared to the first

homologous SAL-1 challenge. To further investigate the difference between repeated

homologous and heterologous infections, we used survival analysis to assess the probability

of the test subjects not requiring treatment at each inoculation level (**Figure 1C**). Median

time to treatment was established at 14, 20 and none for homologous inoculation levels I-III,

respectively, and 21 days for the heterologous challenge. Further analysis of various

parasitemia-related parameters including mean days patent, mean day of peak, mean peak

parasitemia and the Total Area Under the parasitemia Curve (AUC) (**Figure S2**), indicated

that the level of protection against the heterologous challenge in inoculation level IV was

similar to protection after one homologous challenge (i.e., inoculation level II). Indeed, the

mean days of patency was shorter in infection level IV (unpaired t-test = 3.060; df = 6; $p =$

0.0222) while the mean day to peak parasitemia was longer compared to level II (unpaired t-

test = 3.032; df = 6; $p = 0.0230$). No significant difference was found in peak parasitemia

(unpaired t-test = 2.191; df = 6; $p = 0.0709$) and AUC (unpaired t-test = 2.409; df = 6; $p =$

0.0526) between level II and IV (**Figure S2**).

Severe anemia upon *P. vivax* heterologous challenge in sterile homologous immune

Aotus

Next, we investigated the longitudinal dynamics of hematological parameters and selected

blood chemistry during the repeated *P. vivax* infections (**Figure 2 and Table S1**). During the

first inoculation we observed a temporary but significant reduction of both hematocrit and

platelet counts that coincided with peak parasitemia, as has been previously observed in

Aotus [37] and humans experimentally infected with *P. vivax* [38] (**Figure 2A, B -**

inoculation level I). During the second homologous infection, and with partial immunity

ensuing, all the animals had hematological values within the normal range at peak

parasitemia on day 20 PI, when they were treated with CQ for three days (**Figure 2A, B -**

inoculation level II). Of note, MN25029 developed mild anaemia (Hct% = 34.7) and severe

thrombocytopenia (39 x 10^3/μL). During the third homologous infection, none of the animals

became parasitemic and their hematocrit and platelet counts remained stable (note MN25029

again developed a moderate thrombocytopenia (90 x 10^3/µL)) on day 14 PI (**Figure 2A, B –**

inoculation level III). In contrast, the heterologous *P. vivax* AMRU-1 strain challenge

triggered anemia and thrombocytopenia in all the animals (**Figure 2A, 2B - inoculation level**

IV). For instance, mild to moderate anemia developed in two animals (MN30014 and

MN32047) by day 7 PI, even though both animals had undetectable or subpatent parasitemia.

Later, on day 28 PI MN30014 developed moderate anemia and severe thrombocytopenia with

a parasitemia of 1510/µL and needed treatment with MQ to end the experiment. Similarly,

MN32047 developed severe anemia (Hct% = 19.3) on day 18 PI while still negative by light

microscopy and needed treatment with MQ on day 21 PI to end the experiment. In contrast,

MN25029 developed severe thrombocytopenia (24 x 10^3/ µL) at peak parasitemia

(11,430/µL) on day 14 PI, even though, its Hct% remained within normal limits (Hct% = 45),

but later developed moderate anemia (Hct% = 26.2) on day 18 PI when it was still positive at

<10 µL and was treated with MQ on day 21 to end the experiment.

Taken together, these data support previous studies observing the development of severe

anemia (hematocrit < 50% of baseline) and thrombocytopenia (< 50 x 10^3 x µL) in *P. vivax-*

infected *Aotus* monkeys around days 12–15 PI [37]. Indeed, 2/3 (67%) of the remaining

original animals (MN30014, MN32047) and a control (inoculated once) (MN27050) showed

a Reticulocyte Production Index (RPI) below 1.0 suggestive of bone marrow

dyserythropoiesis [39] before inoculation level IV (**Figure 2D**), while only 1/3 (33%) of the

original animals (MN25029) was over an RPI of 1.0 with a Hct% of 45.

Antibody levels increase with repeated infections

In a next series of experiments, we analyzed the development of antibodies against a crude *P.*

vivax lysate across repeated infections (**Figure 3A, Table S2**). After the first inoculation with

P. vivax SAL-1 total antibody (Ab) levels reached a mean of 3.1 Log10 arbitrary ELISA units

(day 28 PI), decreasing slightly to 2.9 Log10 ELISA units by day 84 PI. After the second

homologous inoculation (day 84 PI) Ab levels peaked to 4 Log10 ELISA units on day 114 PI,

decreasing slightly again to 3.5 Log10 ELISA units by day 165 PI (**Figure 3B**). After the

third homologous inoculation (day 165 PI) when all the animals were sterile protected against

challenge (**Figure 3B**), Ab levels remained over 4.0 Log10 ELISA units until day 275 PI

when the animals were challenged with the heterologous *P. vivax* AMRU-1 strain. This time

a booster response was observed with Ab levels increasing to 4.3 Log10 ELISA units by day

304 PI (**Figure 3B**). Interestingly, Ab levels appear to be negatively correlated with

parasitemia (**Figure 3C**). In summary, the dynamics of mean parasitemia and ELISA titers

during inoculation levels I-IV suggest that an ELISA titer of 3-4 arbitrary Log10 units would

fully protect against challenge with a homologous but only partially protect against a

heterologous strain of *P. vivax* (**Figure 3D**). These correlates of protection provide a

benchmark for efficacy testing of *P. vivax* blood stage candidate vaccines in the *Aotus* model.

Quantification of antigen responses using a *P. vivax* protein microarray

Antibody responses during repeated infection (log2 (antigen reactivity/no DNA control

reactivity)) show that 66 out of 244 *P. vivax* antigens in the protein microarray demonstrated

reactivity above 0 in 10% of all samples analyzed (**Figure 4A**). When we compared the

antibody levels for these 66 antigens for all time points in inoculation III (the final

homologous challenge) vs inoculation IV (the heterologous challenge) for the three monkeys

that completed the entire experiment, there were no differentially reactive antigens (paired t-

test with FDR correction). It is possible that there are differentially reactive antigens that

were not identified in this study due to the limited number of antigens tested and/or the small

sample size.

Within inoculation levels I-III, the number of reactive antigens (antigen breadth) was

significantly increased at days 14, 21, and/or 28 when compared with the pre-inoculation

antigen breadth (**Figure 4B**, $p < 0.05$ Wilcoxon matched pairs test). The trend for increased

antigen breadth over time appears similar but not as drastic for the heterologous infection

with the *P. vivax* AMRU-1 in inoculation IV. When we calculated the area under the curve of

antigen breadth for each inoculation level for the three monkeys which completed all four

inoculations, inoculation III and IV were both significantly higher than inoculation I (and

were not different from each other) (**Figure 4C**, $p < 0.05$ repeated measures ANOVA with

paired sample post hoc t-tests). These data show that repeated infections of the homologous

strain *P. vivax* SAL-1 (inoculation levels I-III) increase the breath of the immune response as

the number of infections increased, and that the breadth remained high during heterologous

challenge with *P. vivax* AMRU-1. We also found that the ELISA titer for the crude lysate

correlated well with antigen breadth, however correlations were only significant at

inoculation level II; days 99 (Pearson R = 0.98, significant at $p < 0.005$) and 114 (Pearson R

= 0.86, trend at $p = 0.062$) (**Figure 4D and S3**).

Longitudinal follow up during repeated infection revealed major immunogenic

antigens (Ags) by protein microarray. Indeed, seven targets have significantly higher

antibody responses at inoculation level III compared to inoculation level I (**Figure S4**, **Table**

S3), including the Early Transcribed Membrane Protein (ETRAMP) [PVX_090230],

Parasitophorous vacuolar protein 1 (PV1) [PVX_092070], Merozoite Surface Protein 1

(MSP-1) [099980, fragments 2 & 3], and three Plasmodium Exported Proteins

[PVX_121930, PVX_083560 & PVX_121935]. These data indicate that the targets may

function as surrogates or correlates of sterile immunity in *Aotus*.

The maintenance of antigen breadth after heterologous challenge (inoculation IV)

highlights the presence of homologous or cross-reactive antigens between the two isolates.

However, amino acid sequences for all 7 targets (6 genes) were identical, except for a region

of 14 amino acids in one of the Plasmodium exported proteins (PVX_083560). This suggests

that the partial protection observed following heterologous challenge may not be due to these

proteins, or that transcription of some of these targets may be variable between the isolates.

Genetic diversity rather than immune evasion determines level of strain transcendent

protection

Our data so far suggest that protection from the homologous challenge is antibody mediated,

however the limited resolution of the ELISA and protein array data cannot explain the lower

protection after the heterologous challenge. As an alternative approach we investigated

possible immune evasion mechanisms on genomic and transcriptional level.

For this purpose, we performed whole genome sequencing of both SAL-1 and AMRU-1

strains to improve the strain-transcendent coverage of the existing *P. vivax* microarray

platform [40]. Selective WGA enabled targeted amplification of the AT-rich subtelomeres of

AMRU-1 and SAL-1 strains used in this study (**Figure 5A**). Assembly and annotation

generated continuous subtelomere sequences for SAL-1 and AMRU-1. The number of

contigs in the original SAL-1 dropped from 2748 to 113, highlighting the continuity of the

PacBio assembly (**Figure 5B**). After the annotation with Companion [41] the improved

assembly increased the number of *pir* genes for SAL-1 from 124 to 425, demonstrating that

long reads better represent the number of variable gene families in subtelomeric regions.

Comparison of the *pir* gene repertoire across strains revealed 593 and 425 *pir* genes in

AMRU-1 and SAL-1, respectively, compared to over 1000 in the PvP01 reference strain

(**Figure 5C**). This difference in number may be because the reference strain came straight

from patient infection while SAL-1 and AMRU-1 may have adapted during repeated

passages through monkeys. Finally, the proportion of *pir* subtypes remains constant across

strains as previously reported [42].

With this information in hand, we complemented the existing microarray probe set that

was generated for the *P. vivax* core genome [40] with probes for the SAL-1 and AMRU-I

subtelomeric genes. Next, we investigated whether the virulent phenotype upon heterologous

AMRU-1 infection was a result of immune evasion. Differential gene expression (DGE)

analysis and principal component analysis (PCA) of the expressed genes revealed greater

differences in both core and *pir* genes when comparing AMRU-1 parasites from heterologous

challenges (after 3 inoculations with SAL-1) with SAL-1 parasites during the homologous

challenges (**Figures 6A – left panel, 6B**). We also compared DGE of AMRU-1 parasites

between animals previously infected with three SAL-1 inoculations with i) animals

previously infected with only one SAL-1 inoculation and ii) with the malaria naïve infection

control. Interestingly, only a small number of changes in core and *pir* genes was observed

across these comparisons (**Figure 6A – right panel**). The clear overall similarity of sample

distribution in the PCA plots based on DGE of core (**Figure 6B - left panel**) or *pir* genes

(**Figure 6B – right panel**) suggests that the repeated SAL-1 infections do not induce

extensive *pir* gene switching either in SAL-1 or AMRU-1 parasites. Rather, there appear to

be significant strain-specific differences in both core and *pir* expression between SAL-1 and

AMRU-1. Further analysis using a *pir* gene network revealed no apparent changes in *pir* gene

expression across SAL-1 challenges or upon AMRU-1 challenge (**Figure 6C**).

Altogether, the transcriptional analysis does not indicate that AMRU-1 parasites

actively evade the antibody mediated protection induced by SAL-1 homologous challenges

by antigenic switching. Thus, the lower protection observed after the heterologous challenge

is likely due to major genetic and hence antibody epitope variation between these two

geographically separated strains [43].

Discussion

Previous trials of *P. falciparum* and *P. vivax* vaccine candidates have demonstrated the utility

of the *Aotus* model in supporting vaccine development [44-46]. Various asexual stage

vaccine candidate antigens have been subjected to testing in *Aotus [45, 47-55]*, but only a few

have shown some level of efficacy in human clinical trials [15, 56]. Development of highly

effective strain-transcendent immunity against malaria is a universal goal of vaccine

developers [57]. Recently, whole organism blood stage malaria vaccines have gained

prominence as an alternative to subunit vaccines [58, 59]. One major advantage of

vaccination using whole blood stage parasites is the multiplicity of immunogenic antigens,

including those conserved across strains that may be able to induce strain transcendent

immunity [60, 61].

To assess whether strain-transcendent immunity can be achieved by repeated blood stage

infection with *P. vivax*, and to investigate the correlates of protection during repeated

infection, we infected six *Aotus* monkeys with the *P. vivax* SAL-1 strain until sterile

protected and then challenged with the AMRU-1. We demonstrate that repeated whole blood

stage infection with a homologous *P. vivax* strain (i.e., same strain) induces sterile immunity

in *Aotus* monkeys after only two infections. In contrast, *Aotus* monkeys infected with *P.*

falciparum needed between three to four [62] and six to seven [31] repeated infections to

achieve sterile immunity. This is consistent with previous observations made during

malariotherapy in patients with neurosyphilis demonstrating that immunity to *P. falciparum*

is acquired more slowly than to *P. vivax* or *P. malariae* [26]. Interestingly, *Saimiri sciureus*

boliviensis monkeys immunized with irradiated sporozoites of *P. vivax* SAL-1 and challenged

four to nine times with homologous viable sporozoites over a period of almost four years

showed sterile protection [63]. However, all animals remained susceptible when challenged

345 with SAL-1 blood stage parasites, suggesting that humoral immunity is a correlate of

346 protection against repeated blood stage infections.

347  Furthermore, our study demonstrates that the sterile immunity achieved after repeated

348 infection with a homologous strain was only partially protective after a heterologous

349 challenge (i.e., different strain). Similar observations have been reported for *P. falciparum* in

350 *Aotus* [31]. In both cases, heterologous challenges resulted in severe anemia and

351 thrombocytopenia irrespective of parasitemia level. Such hematological manifestations in

352 semi-immune *Aotus* monkeys with low or subpatent *P. falciparum* parasitemia have been

353 attributed in the past to clearance of non-infected RBCs mediated by autoantibodies [64-67],

354 sequestration of infected RBCs, bone marrow suppression [64, 65] and immune-mediated

355 thrombocytopenia [37, 68-70]. The pernicious severe anemia without thrombocytopenia

356 observed in the monkey MN32047 during subpatent parasitemia may have been the result of

357 immune complex disease, or of dyserythropoiesis due to bone marrow infection, as

358 previously described in humans and *Aotus* monkeys infected with *P. falciparum*, *P. malariae*

359 *and P. brasilianum* [68, 71-77].

360  A *P. vivax* IVTT protein microarray revealed antibody responses against major

361 immunogenic blood stage antigens (Ags) in this study. Immune reactivity to individual

362 antigens and antibody breath in sera from these animals increased with each inoculation level

363 and were statistically significantly different between inoculation levels I and III, when the

364 animals achieved sterile immunity to a homologous SAL-1 challenge. Among the most

365 significant asexual blood stages antigens detected by the protein microarray were ETRAMP

366 (PVX_090230) located in chromosome 5, and two MSP1 fragments: PVX_099980_s4 and

367 PVX_099980_s2 located on chromosome 7, the latter, a leading vaccine candidate that has

368 been identified as a major determinant of strain-specific protective immunity [78].

In this study, animals with sterile immunity to a *P. vivax* SAL-1 homologous

challenge were partially protected against a heterologous AMRU-1 strain. This difference in

protection may have been the result of highly polymorphic antigens associated with essential

parasite phenotypes such as red cell invasion or due to immune evasion, as it has been

suggested in recent genomic studies of *P. vivax* parasites from distinct geographic origin such

as SAL-1 and AMRU-1 [43, 79]. Finding conserved and cryptic (not exposed to the immune

system) epitopes involved in essential phenotypes that could be targeted by strain

transcending neutralizing antibodies represents a possible way forward [80].

In contrast to *P. falciparum* that utilizes the variant PfEMP1 antigens to induce

cytoadherence and avoid splenic clearance of blood stage parasites, limited vascular

sequestration occurs in most other *Plasmodium* species investigated so far. At least in *P.*

vivax, this process may be mediated by *P. vivax* orthologs of the *Plasmodium* interspersed

repeat (PIR) variant antigens [42]. In our study, gene expression analysis along multiple

infections allowed correlating *pir* gene expression with the immune response across

infections to illuminate parasite immune evasion mechanisms during heterologous challenge.

Interestingly, only minor changes in *pir* gene variant expression were observed across all the

different inoculation levels, whether homologous or heterologous. Further analysis using a

pir gene network confirmed no apparent changes in *pir* gene expression in AMRU-1

parasites, regardless of the nature of the previous infections. Together, the transcriptional

analysis does not indicate that *P. vivax* actively evades the antibody-mediated protection

through antigenic switching. These findings are in accordance with previous studies that have

shown no significant difference when comparing the sera of single *versus* repeated infection

in patients for VIR antigens and question their role in immune evasion [81, 82]. The partial

protection observed in the heterologous AMRU-1 challenges may therefore be due to major

genetic differences and hence antibody epitope variation between the two strains [43].

394  In conclusion, our study contributes to our understanding of the pathogenesis of *P. vivax*-

395 induced anemia, *P. vivax* asexual blood stage antigen discovery and correlates of protection,

396 as well as possible immune evasion mechanisms. Finally, we establish a benchmark for *P.*

397 *vivax* protective immunity in the *Aotus* monkey model, providing an important criterion for

398 vaccine development [31].

Materials and methods

Ethics statement

The experimental protocol entitled "Induction of sterile protection by blood stage repeated

infections in *Aotus* monkeys against subsequent challenge with homologous and heterologous

Plasmodium vivax strains" was approved and registered at the ICGES Institutional Animal

Care and Use Committee (CIUCAL) under accession number CIUCAL-01/2016. The

experiment was conducted in accordance with the Animal Welfare Act and the Guide for the

Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources,

National Research Council [83], and the laws and regulations of the Republic of Panama.

Animals and parasites

Twelve laboratory bred (lab-bred) adult male and female "spleen-intact" *Aotus l. lemurinus*

lemurinus Panamanian owl monkeys of karyotypes VIII and IX were used in the study [84].

The animals were cared and maintained as described elsewhere [85]. Isolates of *P. vivax*

SAL-1 originally adapted to splenectomized *Aotus* monkeys by W.C. Collins in 1972 [34]

and further adapted to spleen intact *A. l. lemurinus* [37, 45], and of *P. vivax* AMRU-1 from

Papua New Guinea (PNG) originally adapted to splenectomized *Aotus* by R.D. Cooper in

1994 [86, 87], and further adapted to spleen intact *A. l. lemurinus* by Obaldia N.III. in 1997

[88] were used.

Briefly, each frozen stabilate of SAL-1 and AMRU-1 was thawed, washed three times

with incomplete RPMI medium, and resuspended in 1 ml of RPMI medium. This suspension

was used for intravenous (i.v.) inoculation into the saphenous vein of a donor animal using a

25-gauge butterfly needle catheter attached to a 3-ml syringe. When the level of parasitemia

reached a peak around days 12-15 post-inoculation (PI), a dilution of blood was made in

RPMI to get a total inoculum of 50,000 parasites/ml. All animals received 1 ml of the

inoculum through the saphenous vein.

In total 12 spleen intact lab-bred animals were used in this experiment. Six monkeys

(three male and three females; MN30014, MN30034, MN32028, MN32047, MN25029,

MN29012) were repeatedly infected with *P. vivax* SAL-1 (Homologous challenge) for three

times (Levels I-III) and another six animals served as either donors or were assigned as

infection or naïve controls. Donors and controls were reassigned back into subsequent

inoculation levels as depicted in **Figures 1A** and **S1**. Three SAL-1 homologous sterile

immune monkeys from the original six, plus one infected once with SAL-1 and one malaria

naive control, were re-challenged (Level IV) with the heterologous CQ resistant and *Aotus*

adapted *P. vivax* AMRU-1 strain [37]. The animals were treated with CQ at 15 mg/kg for

three consecutive days during inoculation level I-III and a drug wash out period of 70 days

was kept between inoculation levels I and II and 65 days between levels II and III. No CQ

treatment was instituted in inoculation level III. To treat the *P. vivax* AMRU-1 CQ resistant

strain, inoculated animals on inoculation level IV and at the end of the experiment were

treated with MQ at 25 mg/kg orally once.

General procedures

Five days after infection, the animals were monitored for any signs of clinical disease and

bled 5 µL from a prick made with a lancet in the marginal ear vein to measure daily

parasitemia. Parasitemia was determined using thick blood smear stained with Giemsa as

described in the Earle and Perez (1932) technique [89]. Blood samples were also collected at

regular intervals from the femoral vein to assess humoral immune responses against *P. vivax*

blood stage proteins, for complete blood count (CBC) and blood chemistry (liver and renal

panel), for collection of parasite DNA on FTA® Elute cards (Whatman, Florham Park, NJ.

USA) and for RNA in TRizol® solution (Invitrogen, Carlsbad, CA, USA) for molecular

biology studies. The animals were treated with Mefloquine (MQ) at 25 mg/kg orally by

gastric intubation to end the experiment.

Criteria for parasitemia

For this study patency was defined as the first of three consecutive positive days after

inoculation. Clearance was defined as the first of three consecutive negative days.

Recrudescence was defined as the first of three consecutive positive days after a period of

clearance. Positivity of $<10/\mu L$ for less than three days was considered evidence of subpatent

infection.

Criteria for anemia and thrombocytopenia

For this study we classified anemia based on the hematocrit % as mild (Hct% = 31-36),

moderate (Hct% = 25-30), or severe (Hct% < 25). Thrombocytopenia was considered mild if

platelet counts were between 149-100 x $10^3/\mu L$, moderate if between 99 and 50 x $10^3/\mu L$ or

severe if < 50 x $10^3/\mu L$.

Drug treatment

CQ was administered orally for three consecutive days at 10 mg/Kg daily at peak

parasitemia. Rescue treatment with MQ was triggered if the hematocrit reached 50% of

baseline or hemoglobin was < 8 gm/dL, platelets were < 50 x 10^3/µL or the animals remained

positive by LM after day 28 PI [37].

Serology

Serum ELISA. P. vivax SAL-1 antigen was prepared from *Aotus* iRBCs purified by

Percoll™(GE Healthcare Bio-Sciences AB, Uppsala, Sweden) cushion (47%) centrifugation

as described [90] and adsorbed at 5 µg/mL concentration diluted in PBS pH 7.4 to a 96 well

plate at 4-8 degrees Celsius overnight. The plates were blocked with 5% skimmed milk in

PBS-0.05% Tween for 2 hours. Serum samples were added to the plate at a dilution of 1/100

in dilution buffer and incubated for one hour, washed further 5 times with PBS pH 7.4 and

incubated for one hour with Goat anti-monkey (Rhesus macaque) (Abcam cat # a112767),

diluted 1:2000 in PBS pH 7.4. 100 uL per well of the OPD substrate solution (P9029-50G,

Sigma-Aldrich, St. Louis, MI, USA) was added to the plate and incubated for 30 minutes

away from light and the reaction was stopped with 50 µL of sulfuric acid 3N. To detect the

antigen–antibody reactivity, the plates were then read immediately at 492 nm in a ELx808

Plate reader (BioTek®, Winooski, VT, USA).

pLDH ELISA. To measure *P. vivax* lactate dehydrogenase levels (PvLDH) in the monkey

plasma samples, ELISA was performed using a matching pair of capture and detection

antibodies (Mybiosource, San Diego, CA). Briefly, 96-well microtiter plate was coated with

mouse monoclonal anti-pLDH (clone #M77288) at a concentration of 2µg/mL in PBS (pH

7.4) and incubated overnight at 4 °C. The plate was washed and incubated with blocking

buffer (PBS-BSA 1% - reagent diluent) at room temperature for 2hrs. After washing, samples

were diluted 1:2, added to the plate and incubated for 2hrs. Next, plates were washed and

HRP-conjugated anti-pLDH detection antibody (clone #M12299), diluted 1:1000 in blocking

buffer, was incubated for 1hr at room temperature Plates were washed and incubated for 15

min with substrate solution (OPD), the reaction was stopped adding sulphuric acid 2.5M.

Optical density was determined at 450 nm. Cut-off of positivity was defined by correcting

absorbance values generated in the plasma samples from blank values (plate controls).

Schizont extracts were used to perform standard curves and lower absorbance values were in

the range of O.D = 0.01–0.02. All positive monkey samples gave O.D. values equal to or

higher than 0.05.

Protein microarray and hybridization. The construction of the protein microarray was

conducted using methods as described elsewhere [91]. Briefly, coding sequences were PCR

amplified from *P. vivax* SAL-1 genomic DNA and cloned into the PXT7 plasmid using

homologous recombinant as complete or overlapping fragments, the resulting plasmids

(n=244) were expressed in an *Escherichia.coli* based *in vitro* transcription/translation (IVTT)

reactions, and the completed reactions printed onto nitrocellulose-coated microarray slides

(Grace Bio-Labs, USA). Serum samples mixed with 1/100 blocking buffer (ArrayIt Corp,

USA) supplemented with *E. coli* lysate (Genscript, USA). The diluted serum samples were

incubated with the protein arrays overnight at 4^0C, followed by incubation with a goat anti-

human IgG Texas Red secondary antibody (Southern Biotech, USA). The arrays were

scanned using a Genepix 4300A scanner (Molecular Devices, UK) at 5µm resolution and a

wavelength of 594nm [92].

Protein microarray data processing and analysis. Raw median fluorescent intensity was local

background corrected using the normexp function (offset = 50, method = "mle", limma R

package). All data was log transformed (base 2) and normalized as a ratio of the signal for

each spot to the mean of the no DNA control spot within each sample. The number of

antigens that have reactivity above 0 in at least 10% of samples was calculated and included

in the heatmap (generated in Microsoft excel). Seropositive antigens for each sample were

defined as those with reactivity above the mean of the sample specific No DNA control spots

+ 3SD. These seropositive antigens were totalled for each sample to determine the antigen

breadth (number of reactive antigens). The antigen breadth AUC was calculated using the

trapezoid rule after limiting the data to only the same number of time points for all

inoculations. Pearson's correlations were performed for available ELISA titers and antigen

breadth. All statistics and plots were done using R unless otherwise specified.

PacBio Whole Genome Sequencing and analysis

P. vivax AMRU-1 and SAL-1 were amplified with selective whole genome amplification

(sWGA), using primers specific for the subtelomeres that are enriched for low GC content

[93]. Amplified DNA was used for PacBio sequencing using a commercial protocol

(Genscript). We obtained 373,772 subreads with an N50 (type of median length of the reads)

of 13,119 bp for SAL-1 and 325,996 subreads with an N50 of 12,035 bp for AMRU-1. The

reads were mapped with BWA MEM for quality control (**Figure 5A**) and then assembled

with canu [94] (parameter: genomeSize=32m ErrorRate=0.10 gnuplotTested=true useGrid=0

-pacbio-raw, version January 2018). The assemblies generated 113 and 103 contigs with an

N50 of 50k and 41k and the largest contig be 195kb and 140kb for SAL-1 and AMRU-1,

respectively. For annotation, the assemblies were loaded into Companion [41], using PvP01

as reference strain (June 2018, Augustus cut-off set to 0.4). The genome and its annotation

can be found at http://cellatlas.mvls.gla.ac.uk/Assemblies/.

For the Gephi analysis, we extracted all the genes annotated as *pir* from the two

Companion runs, merged them with the *pir* genes of PvP01 and performed an all-against-all

BLASTp (-F F, Evalue 1e-6). The results were parsed into the open source software Gephi to

produce **Figure 5C**. For graphical representation, a force atlas algorithm was run and then the

global identity cut-off was set to 32% and the Fruchterman Reingold algorithm was run.

Gene expression microarray and gene expression analysis

pir gene probe development: Sequences from sWGA, representing mostly the AT-rich subtelomeres and excluding the mitochondrial genes, were used as input for probe design using OligoRankPick [95](oligo size=60, %GC=40). The oligos that were overlapping with core genome oligos from the existing *P. vivax* microarray [40] were removed (12 for SAL-1 and 6 for AMRU-1). The final list of new oligos contains 929 SAL-1 probes and 701 AMRU-1 probes, amongst which 8 match two SAL-1 genes and 5 ma tch two AMRU-1 genes (Full list as **Table S5**).

RNA preparation and microarray hybridization. Cell pellets from the blood samples collected at different time points during SAL-1 or AMRU-1 inoculations were stored in trizol. RNA was extracted and processed to be run in a customized microarray assay detecting both core and subtelomoeric genes. The previously described microarray hybridization protocol was used for this study, with several modifications [96]. In brief, 100 ng of cDNA was used for subsequent 10 rounds of amplification to generate aminoallyl-coupled cDNA for the hybridizations as described [96]. 17µl (~ 5 µg) of each Cy-5-labelled (GE Healthcare) cDNA of the sample and an equal amount of Cy-3-labelled (GE Healthcare) cDNA of the reference pool were then hybridized together on customized microarray chip using commercially available hybridization platform (Agilent) for 20 h at 70 °C with rotation at 10 rpm. Microarrays were washed and immediately scanned using Power Scanner (Tecan) at 10 µm resolution and with automated photomultiplier tubes gain adjustments to balance the signal intensities between both channels. The reference pool used for microarray was a mixture of 3D7 parasite strain RNA collected every 6 h during 48 h of the full IDC.

Microarray analysis. To quantify microarray data signals, intensities were first corrected using an adaptive background correction using the method "normexp" and offset 50 using the Limma package in R[97]. Next, we performed within-array loess normalization followed by

quantile-normalization between samples/arrays. Each gene expression was estimated as the

average of log2 ratios (Cy5/Cy3) of representative probes, thus intensities or log-ratios could

be comparable across arrays. Finally, probes with signal showing median foreground

intensity less than 2-fold of the median background intensity at either Cy5 (sample RNA) or

Cy3 (reference pool RNA) channel were assigned missing values. Fold changes and standard

errors of relative gene expression were estimated by fitting a linear model for each gene,

followed by empirical Bayes smoothing to the standard errors. Next, the average log 2-

expression level for each gene across all the arrays was calculated using the topTable

function of the limma package. In parallel we adjusted p-values for multiple testing using the

Benjamini and Hochberg's method to control the false discovery rate. The lists of

differentially expressed genes (DEGs) for each of the comparisons were extracted by defining

a cut-off of adjusted p-values < 0.001 and fold change > 1. The log fold change and adjusted

p-values were graphed in volcano plots, using the EnhancedVolcano package in R. From the

lists of DEGs, we matched and highlighted those related to 3 categories: (i) surface proteins

related to parasite invasion, determined as syntenic orthologs with the *P. falciparum*

exportome (refs); (ii) proteins whose expression is spleen-dependent (ref); and (iii) *P. vivax*

IVTT antigens inducing high antibody responses as determined by the protein array.

Acknowledgements

588 Funding for this study was provided in part by core funds from the Gorgas Memorial Institute

589 of Health Studies and the Sistema Nacional de Investigación de Panamá, SENACYT-SNI

590 awarded to N.O., Panamá, Panamá. This work was also supported by Wolfson Merit Royal

591 Society Award (to M.M.) and Wellcome Trust Center Award 104111 (to F.A., J.L.S.F.,

592 T.D.O. and M.M.).

Competing interests

The authors declare that they have no financial or non-financial competing interests.

Author Contributions

Conceptualization: Nicanor Obaldia, Matthias Marti.

Data curation: Nicanor Obaldia, Joao Luiz Da Silva Filho, Kevin K.A. Tetteh, Thomas D. Otto, Matthias Marti.

Formal analysis: Nicanor Obaldia, Joao Luiz Da Silva Filho, Katherine A. Glass, Tate Oulton, Kevin K.A. Tetteh, Thomas D. Otto, Matthias Marti.

Funding acquisition: Nicanor Obaldia, Matthias Marti.

Investigation: Nicanor Obaldia, Marlon Nuñez, Joao Luiz Da Silva Filho, Katherine A. Glass, Tate Oulton, Fiona Achcar, Grennady Wirjanata, Manoj Duraisingh, Philip Felgner, Kevin K.A. Tetteh, Zbynek Bozdech, Thomas D. Otto, Matthias Marti.

Methodology: Nicanor Obaldia, Matthias Marti.

Project administration: Nicanor Obaldia.

Resources: Nicanor Obaldia, Philip Felgner, Kevin K.A. Tetteh, Zbynek Bozdech, Thomas D. Otto, Matthias Marti.

Supervision: Nicanor Obaldia.

Validation: Nicanor Obaldia.

Visualization: Nicanor Obaldia, Joao Luiz Da Silva Filho, Kevin K.A. Tetteh, Thomas D. Otto, Matthias Marti.

Writing – original draft: Nicanor Obaldia, Matthias Marti.

Writing – review & editing: Nicanor Obaldia, Joao Luiz Da Silva Filho, Matthias Marti.

References

References

1.   WHO. World Malaria Report 2021. 2021.

2.   Schuldt NJ, Amalfitano A. Malaria vaccines: focus on adenovirus based vectors.

Vaccine. 2012;30(35):5191-8. Epub 2012/06/12. doi: 10.1016/j.vaccine.2012.05.048.

PubMed PMID: 22683663.

3.   WHO. World Malaria Report 2019. 2019.

4.   Bourgard C, Albrecht L, Kayano A, Sunnerhagen P, Costa FTM. Plasmodium vivax

Biology: Insights Provided by Genomics, Transcriptomics and Proteomics. Front Cell Infect

Microbiol. 2018;8:34. Epub 20180208. doi: 10.3389/fcimb.2018.00034. PubMed PMID:

29473024; PubMed Central PMCID: PMCPMC5809496.

5.   Price RN, Commons RJ, Battle KE, Thriemer K, Mendis K. Plasmodium vivax in the

Era of the Shrinking P. falciparum Map. Trends Parasitol. 2020;36(6):560-70.

6.   Battle KE, Baird JK. The global burden of Plasmodium vivax malaria is obscure and

insidious. PLoS Med. 2021;18(10):e1003799. Epub 2021/10/08. doi:

10.1371/journal.pmed.1003799. PubMed PMID: 34618814; PubMed Central PMCID:

PMCPMC8496786.

7.   Fernandez-Becerra C, Aparici-Herraiz I, Del Portillo HA. Cryptic erythrocytic

infections in Plasmodium vivax, another challenge to its elimination. Parasitol Int.

2021:102527. Epub 2021/12/14. doi: 10.1016/j.parint.2021.102527. PubMed PMID:

34896615.

8.   Adams JH, Mueller I. The Biology of Plasmodium vivax. In: Wirth DF, Alonso P,

editors. Malaria: Biology in the Era of Eradication. Cold Spring Harbor, NY: Cold Spring

Harbor Laboratory Press; 2017. p. 43-54.

9.   Bantuchai S, Imad H, Nguitragool W. Plasmodium vivax gametocytes and

transmission. Parasitol Int. 2021:102497. Epub 2021/11/09. doi:

10.1016/j.parint.2021.102497. PubMed PMID: 34748969.

10.   Baro B, Deroost K, Raiol T, Brito M, Almeida AC, de Menezes-Neto A, et al.

Plasmodium vivax gametocytes in the bone marrow of an acute malaria patient and changes

in the erythroid miRNA profile. PLoS Negl Trop Dis. 2017;11(4):e0005365. doi:

10.1371/journal.pntd.0005365. PubMed PMID: 28384192; PubMed Central PMCID:

PMC5383020.

11.   Obaldia N, 3rd, Meibalan E, Sa JM, Ma S, Clark MA, Mejia P, et al. Bone Marrow Is

a Major Parasite Reservoir in Plasmodium vivax Infection. MBio. 2018;9(3). doi:

10.1128/mBio.00625-18. PubMed PMID: 29739900; PubMed Central PMCID:

PMCPMC5941073.

12.   Markus MB. New Evidence for Hypnozoite-Independent Plasmodium vivax Malarial

Recurrences. Trends Parasitol. 2018;34(12):1015-6. doi: 10.1016/j.pt.2018.08.010. PubMed

PMID: 30213708.

13.   Baird JK. African Plasmodium vivax malaria improbably rare or benign. Trends

Parasitol. 2022. Epub 2022/06/07. doi: 10.1016/j.pt.2022.05.006. PubMed PMID: 35667992.

14.   Baird JK. Resistance to therapies for infection by Plasmodium vivax. Clin Microbiol

Rev. 2009;22(3):508-34. Epub 2009/07/15. doi: 10.1128/CMR.00008-09. PubMed PMID:

19597012; PubMed Central PMCID: PMCPMC2708388.

15.   De SL, Ntumngia FB, Nicholas J, Adams JH. Progress towards the development of a

P. vivax vaccine. Expert Rev Vaccines. 2021. Epub 2021/01/23. doi:

10.1080/14760584.2021.1880898. PubMed PMID: 33481638.

16.   White M, Chitnis CE. Potential role of vaccines in elimination of Plasmodium vivax.

Parasitol Int. 2022;90:102592. Epub 2022/05/01. doi: 10.1016/j.parint.2022.102592. PubMed

PMID: 35489701.

17.   White NJ. What causes malaria anemia? Blood. 2022;139(15):2268-9. Epub

2022/04/15. doi: 10.1182/blood.2021015055. PubMed PMID: 35420692.

18.   Val F, Avalos S, Gomes AA, Zerpa JEA, Fontecha G, Siqueira AM, et al. Are

respiratory complications of Plasmodium vivax malaria an underestimated problem? Malar J.

2017;16(1):495. Epub 2017/12/24. doi: 10.1186/s12936-017-2143-y. PubMed PMID:

29273053; PubMed Central PMCID: PMCPMC5741897.

19.   Zain Ul A, Qadeer A, Akhtar A, Rasheed A. Severe acute respiratory distress

syndrome secondary to Plasmodium vivax malaria. J Pak Med Assoc. 2016;66(3):351-3.

Epub 2016/03/13. PubMed PMID: 26968294.

20.   Sharma R, Suneja A, Yadav A, Guleria K. Plasmodium vivax Induced Acute

Respiratory Distress Syndrome - A Diagnostic and Therapeutic Dilemma in Preeclampsia. J

Clin Diagn Res. 2017;11(4):QD03-QD4. Epub 2017/06/03. doi:

10.7860/JCDR/2017/25529.9710. PubMed PMID: 28571216; PubMed Central PMCID:

PMCPMC5449862.

21.   Baird JK. Severe and fatal vivax malaria challenges 'benign tertian malaria' dogma.

Ann Trop Paediatr. 2009;29(4):251-2. Epub 2009/11/28. doi:

10.1179/027249309X12547917868808. PubMed PMID: 19941746.

22.   Mohan K, Stevenson MN. Acquired Immunity to Asexual Blood Stages. In: I.W. S,

editor. Malaria: Parasite Biology, Pathogenesis and Protection. Washington D.C.: ASM

Press; 1998. p. 467-93.

23.   Snounou G, Perignon JL. Malariotherapy--insanity at the service of malariology. Adv

Parasitol. 2013;81:223-55. Epub 2013/02/07. doi: 10.1016/B978-0-12-407826-0.00006-0.

PubMed PMID: 23384625.

24.   Mueller I, Galinski MR, Baird JK, Carlton JM, Kochar DK, Alonso PL, et al. Key

gaps in the knowledge of Plasmodium vivax, a neglected human malaria parasite. Lancet

Infect Dis. 2009;9(9):555-66. Epub 2009/08/22. doi: S1473-3099(09)70177-X [pii]

10.1016/S1473-3099(09)70177-X. PubMed PMID: 19695492.

25.   Jeffery GM. Epidemiological significance of repeated infections with homologous

and heterologous strains and species of Plasmodium. Bull World Health Organ.

1966;35(6):873-82. Epub 1966/01/01. PubMed PMID: 5298036; PubMed Central PMCID:

PMCPMC2476277.

26.   Doolan DL, Dobano C, Baird JK. Acquired immunity to malaria. Clin Microbiol Rev.

2009;22(1):13-36, Table of Contents. Epub 2009/01/13. doi: 10.1128/CMR.00025-08.

PubMed PMID: 19136431; PubMed Central PMCID: PMCPMC2620631.

27.   Moorthy VS, Newman RD, Okwo-Bele JM. Malaria vaccine technology roadmap.

Lancet. 2013;382(9906):1700-1. Epub 2013/11/19. doi: 10.1016/S0140-6736(13)62238-2.

PubMed PMID: 24239252.

28.   Hodgson SH, Choudhary P, Elias SC, Milne KH, Rampling TW, Biswas S, et al.

Combining Viral Vectored and Protein-in-adjuvant Vaccines Against the Blood-stage

Malaria Antigen AMA1: Report on a Phase 1a Clinical Trial. Mol Ther. 2014;22(12):2142-

54. doi: 10.1038/mt.2014.157. PubMed PMID: 25156127; PubMed Central PMCID:

PMC4250079.

29.   Wykes M, Good MF. A case for whole-parasite malaria vaccines. Int J Parasitol.

2007;37(7):705-12. Epub 2007/04/06. doi: 10.1016/j.ijpara.2007.02.007. PubMed PMID:

17408673.

30.   Pombo DJ, Lawrence G, Hirunpetcharat C, Rzepczyk C, Bryden M, Cloonan N, et al.

Immunity to malaria after administration of ultra-low doses of red cells infected with

Plasmodium falciparum. Lancet. 2002;360(9333):610-7. Epub 2002/09/21. doi:

10.1016/S0140-6736(02)09784-2. PubMed PMID: 12241933.

31.   Jones TR, Obaldia N, 3rd, Gramzinski RA, Hoffman SL. Repeated infection of Aotus

monkeys with Plasmodium falciparum induces protection against subsequent challenge with

homologous and heterologous strains of parasite. Am J Trop Med Hyg. 2000;62(6):675-80.

PubMed PMID: 11304053.

32.   Siddiqui WA, Taylor DW, Kan SC, Kramer K, Richmond-Crum S. Partial protection

of Plasmodium falciparum-vaccinated Aotus trivirgatus against a challenge of a heterologous

strain. Am J Trop Med Hyg. 1978;27(6):1277-8. Epub 1978/11/01. doi:

10.4269/ajtmh.1978.27.1277. PubMed PMID: 103450.

33.   Elliott SR, Kuns RD, Good MF. Heterologous immunity in the absence of variant-

specific antibodies after exposure to subpatent infection with blood-stage malaria. Infect

Immun. 2005;73(4):2478-85. Epub 2005/03/24. doi: 10.1128/IAI.73.4.2478-2485.2005.

PubMed PMID: 15784594; PubMed Central PMCID: PMCPMC1087398.

34.   Collins WE, Contacos PG, Krotoski WA, Howard WA. Transmission of four Central

American strains of Plasmodium vivax from monkey to man. J Parasitol. 1972;58(2):332-5.

Epub 1972/04/01. PubMed PMID: 4623380.

35.   Gomes PS, Bhardwaj J, Rivera-Correa J, Freire-De-Lima CG, Morrot A. Immune

Escape Strategies of Malaria Parasites. Front Microbiol. 2016;7:1617. Epub 2016/11/02. doi:

10.3389/fmicb.2016.01617. PubMed PMID: 27799922; PubMed Central PMCID:

PMCPMC5066453.

36. Rieckmann KH, Davis DR, Hutton DC. Plasmodium vivax resistance to chloroquine?

Lancet. 1989;2(8673):1183-4. Epub 1989/11/18. doi: 10.1016/s0140-6736(89)91792-3.

PubMed PMID: 2572903.

37. Obaldia N, 3rd. Clinico-pathological observations on the pathogenesis of severe

thrombocytopenia and anemia induced by Plasmodium vivax infections during antimalarial

drug efficacy trials in Aotus monkeys. The American journal of tropical medicine and

hygiene. 2007;77(1):3-13. Epub 2007/07/11. PubMed PMID: 17620623.

38. Woolley SD, Marquart L, Woodford J, Chalon S, Moehrle JJ, McCarthy JS, et al.

Haematological response in experimental human Plasmodium falciparum and Plasmodium

vivax malaria. Malar J. 2021;20(1):470. Epub 2021/12/22. doi: 10.1186/s12936-021-04003-7.

PubMed PMID: 34930260; PubMed Central PMCID: PMCPMC8685492.

39. Wickramasinghe SN, Looareesuwan S, Nagachinta B, White NJ. Dyserythropoiesis

and ineffective erythropoiesis in Plasmodium vivax malaria. Br J Haematol. 1989;72(1):91-9.

PubMed PMID: 2660903.

40. Bozdech Z, Mok S, Hu G, Imwong M, Jaidee A, Russell B, et al. The transcriptome

of Plasmodium vivax reveals divergence and diversity of transcriptional regulation in malaria

parasites. Proc Natl Acad Sci U S A. 2008;105(42):16290-5. Epub 20081013. doi:

10.1073/pnas.0807404105. PubMed PMID: 18852452; PubMed Central PMCID:

PMCPMC2571024.

41. Steinbiss S, Silva-Franco F, Brunk B, Foth B, Hertz-Fowler C, Berriman M, et al.

Companion: a web server for annotation and analysis of parasite genomes. Nucleic Acids

Res. 2016;44(W1):W29-34. Epub 20160421. doi: 10.1093/nar/gkw292. PubMed PMID:

27105845; PubMed Central PMCID: PMCPMC4987884.

42. Auburn S, Bohme U, Steinbiss S, Trimarsanto H, Hostetler J, Sanders M, et al. A new

Plasmodium vivax reference sequence with improved assembly of the subtelomeres reveals

an abundance of pir genes. Wellcome Open Res. 2016;1:4. Epub 20161115. doi:

10.12688/wellcomeopenres.9876.1. PubMed PMID: 28008421; PubMed Central PMCID:

PMCPMC5172418.

43.   Buyon LE, Santamaria AM, Early AM, Quijada M, Barahona I, Lasso J, et al.

Population genomics of Plasmodium vivax in Panama to assess the risk of case importation

on malaria elimination. PLoS Negl Trop Dis. 2020;14(12):e0008962. Epub 2020/12/15. doi:

10.1371/journal.pntd.0008962. PubMed PMID: 33315861; PubMed Central PMCID:

PMCPMC7769613 was unable to confirm their authorship contributions. On their behalf, the

corresponding author has reported their contributions to the best of their knowledge.

44.   Collins WE. The Owl Monkey as a Model for Malaria. In: Baer JFW, R.E.; Kakoma,

I., editor. Aotus The Owl Monkey. San Diego, CA: Academic Press Inc.; 1994. p. 217-44.

45.   Obaldia N, 3rd, Stockelman MG, Otero W, Cockrill JA, Ganeshan H, Abot EN, et al.

A Plasmodium vivax Plasmid DNA- and Adenovirus-Vectored Malaria Vaccine Encoding

Blood-Stage Antigens AMA1 and MSP142 in a Prime/Boost Heterologous Immunization

Regimen Partially Protects Aotus Monkeys against Blood-Stage Challenge. Clin Vaccine

Immunol. 2017;24(4). doi: 10.1128/CVI.00539-16. PubMed PMID: 28179404; PubMed

Central PMCID: PMCPMC5382831.

46.   Douglas AD, Baldeviano GC, Lucas CM, Lugo-Roman LA, Crosnier C, Bartholdson

SJ, et al. A PfRH5-based vaccine is efficacious against heterologous strain blood-stage

Plasmodium falciparum infection in aotus monkeys. Cell Host Microbe. 2015;17(1):130-9.

doi: 10.1016/j.chom.2014.11.017. PubMed PMID: 25590760; PubMed Central PMCID:

PMCPMC4297294.

47.   Gramzinski RA, Millan CL, Obaldia N, Hoffman SL, Davis HL. Immune response to

a hepatitis B DNA vaccine in Aotus monkeys: a comparison of vaccine formulation, route,

and method of administration. Mol Med. 1998;4(2):109-18. Epub 1998/04/16. PubMed

PMID: 9508788; PubMed Central PMCID: PMC2230303.

48.   Herrera MA, Rosero F, Herrera S, Caspers P, Rotmann D, Sinigaglia F, et al.

Protection against malaria in Aotus monkeys immunized with a recombinant blood-stage

antigen fused to a universal T-cell epitope: correlation of serum gamma interferon levels with

protection. Infect Immun. 1992;60(1):154-8. PubMed PMID: 1370271; PubMed Central

PMCID: PMCPMC257516.

49.   Herrera S, Herrera MA, Certa U, Corredor A, Guerrero R. Efficiency of human

Plasmodium falciparum malaria vaccine candidates in Aotus lemurinus monkeys. Mem Inst

Oswaldo Cruz. 1992;87 Suppl 3:423-8. PubMed PMID: 1343722.

50.   Herrera S, Herrera MA, Corredor A, Rosero F, Clavijo C, Guerrero R. Failure of a

synthetic vaccine to protect Aotus lemurinus against asexual blood stages of Plasmodium

falciparum. Am J Trop Med Hyg. 1992;47(5):682-90. PubMed PMID: 1449209.

51.   Inselburg J, Bzik D, Li W, Green K, Kansopon J, Hahm B, et al. Protective immunity

induced in Aotus monkeys by recombinant SERA proteins of Plasmodium falciparum. Infect

Immun. 1991;59:1247-50.

52.   Jones TR, Narum DL, Gozalo AS, Aguiar J, Fuhrmann SR, Liang H, et al. Protection

of Aotus monkeys by Plasmodium falciparum EBA-175 region II DNA prime-protein boost

immunization regimen. J Infect Dis. 2001;183(2):303-12. doi: 10.1086/317933. PubMed

PMID: 11110648.

53.   Jones TR, Obaldia N, 3rd, Gramzinski RA, Charoenvit Y, Kolodny N, Kitov S, et al.

Synthetic oligodeoxynucleotides containing CpG motifs enhance immunogenicity of a

peptide malaria vaccine in Aotus monkeys. Vaccine. 1999;17(23-24):3065-71. Epub

1999/08/26. PubMed PMID: 10462241.

54.   Siddiqui WA. An effective immunization of experimental monkeys against a human

malaria parasite, Plasmodium falciparum. Science. 1977;197(4301):388-9. Epub 1977/07/22.

doi: 10.1126/science.406671. PubMed PMID: 406671.

55.   Singh S, Miura K, Zhou H, Muratova O, Keegan B, Miles A, et al. Immunity to

recombinant plasmodium falciparum merozoite surface protein 1 (MSP1): protection in

Aotus nancymai monkeys strongly correlates with anti-MSP1 antibody titer and in vitro

parasite-inhibitory activity. Infection and immunity. 2006;74(8):4573-80. Epub 2006/07/25.

doi: 10.1128/IAI.01679-05. PubMed PMID: 16861644; PubMed Central PMCID:

PMC1539572.

56.   Beeson JG, Kurtovic L, Dobano C, Opi DH, Chan JA, Feng G, et al. Challenges and

strategies for developing efficacious and long-lasting malaria vaccines. Sci Transl Med.

2019;11(474). Epub 2019/01/11. doi: 10.1126/scitranslmed.aau1458. PubMed PMID:

30626712.

57.   Elias SC, Collins KA, Halstead FD, Choudhary P, Bliss CM, Ewer KJ, et al.

Assessment of immune interference, antagonism, and diversion following human

immunization with biallelic blood-stage malaria viral-vectored vaccines and controlled

malaria infection. J Immunol. 2013;190(3):1135-47. Epub 20130104. doi:

10.4049/jimmunol.1201455. PubMed PMID: 23293353; PubMed Central PMCID:

PMCPMC3672846.

58.   Stanisic DI, Good MF. Whole organism blood stage vaccines against malaria.

Vaccine. 2015;33(52):7469-75. Epub 2015/10/03. doi: 10.1016/j.vaccine.2015.09.057.

PubMed PMID: 26428451.

59.   Good MF, Stanisic DI. Whole parasite vaccines for the asexual blood stages of

Plasmodium. Immunol Rev. 2020;293(1):270-82. Epub 2019/11/12. doi: 10.1111/imr.12819.

PubMed PMID: 31709558.

60.   Raja AI, Cai Y, Reiman JM, Groves P, Chakravarty S, McPhun V, et al. Chemically

Attenuated Blood-Stage Plasmodium yoelii Parasites Induce Long-Lived and Strain-

Transcending Protection. Infect Immun. 2016;84(8):2274-88. Epub 2016/06/02. doi:

10.1128/IAI.00157-16. PubMed PMID: 27245410; PubMed Central PMCID:

PMCPMC4962623.

61.   Stanisic DI, Ho MF, Nevagi R, Cooper E, Walton M, Islam MT, et al. Development

and Evaluation of a Cryopreserved Whole-Parasite Vaccine in a Rodent Model of Blood-

Stage Malaria. mBio. 2021;12(5):e0265721. Epub 2021/10/20. doi: 10.1128/mBio.02657-21.

PubMed PMID: 34663097; PubMed Central PMCID: PMCPMC8524336.

62.   Udeinya IJ, Miller LH, McGregor IA, Jensen JB. Plasmodium falciparum strain-

specific antibody blocks binding of infected erythrocytes to amelanotic melanoma cells.

Nature. 1983;303(5916):429-31. doi: 10.1038/303429a0. PubMed PMID: 6343885.

63.   Collins WE, Skinner JC, Millet P, Broderson JR, Filipski VK, Morris CL, et al.

Reinforcement of immunity in Saimiri monkeys following immunization with irradiated

sporozoites of Plasmodium vivax. Am J Trop Med Hyg. 1992;46(3):327-34. doi:

10.4269/ajtmh.1992.46.327. PubMed PMID: 1558272.

64.   Egan AF, Fabucci ME, Saul A, Kaslow DC, Miller LH. Aotus New World monkeys:

model for studying malaria-induced anemia. Blood. 2002;99(10):3863-6. Epub 2002/05/03.

doi: 10.1182/blood.v99.10.3863. PubMed PMID: 11986251.

65.   Jones TR, Stroncek DF, Gozalo AS, Obaldia N, 3rd, Andersen EM, Lucas C, et al.

Anemia in parasite- and recombinant protein-immunized aotus monkeys infected with

Plasmodium falciparum. Am J Trop Med Hyg. 2002;66(6):672-9. Epub 2002/09/13. doi:

10.4269/ajtmh.2002.66.672. PubMed PMID: 12224573.

66.   Mourao LC, Cardoso-Oliveira GP, Braga EM. Autoantibodies and Malaria: Where

We Stand? Insights Into Pathogenesis and Protection. Front Cell Infect Microbiol.

2020;10:262. Epub 2020/07/01. doi: 10.3389/fcimb.2020.00262. PubMed PMID: 32596165;

PubMed Central PMCID: PMCPMC7300196.

67.   Mourao LC, Medeiros CMP, Cardoso-Oliveira GP, Roma P, Aboobacar J, Rodrigues

BCM, et al. Effects of IgG and IgM autoantibodies on non-infected erythrocytes is related to

ABO blood group in Plasmodium vivax malaria and is associated with anemia. Microbes

Infect. 2020;22(8):379-83. Epub 2020/02/26. doi: 10.1016/j.micinf.2020.02.003. PubMed

PMID: 32097712.

68.   Weller RE, Collins WE, Buschbom RL, Malaga CA, Ragan HA. Impaired renal

function in owl monkeys (Aotus nancymai) infected with Plasmodium falciparum. Mem Inst

Oswaldo Cruz. 1992;87 Suppl 3:435-42. Epub 1992/01/01. doi: 10.1590/s0074-

02761992000700073. PubMed PMID: 1343724.

69.   Nagatake T, Broderson JR, Tegoshi T, Collins WE, Aikawa M. Renal pathology in

owl monkeys vaccinated with Plasmodium falciparum asexual blood-stage synthetic peptide

antigens. Am J Trop Med Hyg. 1992;47(5):614-20. Epub 1992/11/01. doi:

10.4269/ajtmh.1992.47.614. PubMed PMID: 1449202.

70.   Iseki M, Broderson JR, Pirl KG, Igarashi I, Collins WE, Aikawa M. Renal pathology

in owl monkeys in Plasmodium falciparum vaccine trials. Am J Trop Med Hyg.

1990;43(2):130-8. Epub 1990/08/01. doi: 10.4269/ajtmh.1990.43.130. PubMed PMID:

2202223.

71.   Houba V. Immunopathology of nephropathies associated with malaria. Bull World

Health Organ. 1975;52(2):199-207. Epub 1975/01/01. PubMed PMID: 1083308; PubMed

Central PMCID: PMCPMC2366359.

72.   Chalifoux LV, Bronson RT, Sehgal P, Blake BJ, King NW. Nephritis and hemolytic

anemia in owl monkeys (Aotus trivirgatus). Vet Pathol. 1981;18(Suppl 6):23-37. Epub

1981/04/01. doi: 10.1177/0300985881018s0603. PubMed PMID: 7344245.

73.   Sato Y, Yanagita M. Renal anemia: from incurable to curable. Am J Physiol Renal

Physiol. 2013;305(9):F1239-48. Epub 2013/07/26. doi: 10.1152/ajprenal.00233.2013.

PubMed PMID: 23884144.

74.   Weller RE. Infectious and Noninfectious Diseases of Owl Monkeys. In: Baer JFW,

R.E.; Kakoma, I., editor. Aotus the Owl Monkey. San Diego, CA, USA.: Academic Press,

INC.; 1994. p. 178-211.

75.   Voller A, Davies DR, Hutt MS. Quartan malarial infections in Aotus trivirgatus with

special reference to renal pathology. Br J Exp Pathol. 1973;54(5):457-68. Epub 1973/10/01.

PubMed PMID: 4202259; PubMed Central PMCID: PMCPMC2072562.

76.   Hutt MS, Davies DR, Voller A. Malarial infections in Aotus trivirgatus with special

reference to renal pathology. II. P. falciparum and mixed malaria infections. Br J Exp Pathol.

1975;56(5):429-38. Epub 1975/10/01. PubMed PMID: 813757; PubMed Central PMCID:

PMCPMC2072788.

77.   Silva-Filho JL, Dos-Santos JC, Judice C, Beraldi D, Venugopal K, Lima D, et al.

Total parasite biomass but not peripheral parasitaemia is associated with endothelial and

haematological perturbations in Plasmodium vivax patients. Elife. 2021;10. Epub 20210929.

doi: 10.7554/eLife.71351. PubMed PMID: 34585667; PubMed Central PMCID:

PMCPMC8536259.

78.   Fonseca JA, Cabrera-Mora M, Singh B, Oliveira-Ferreira J, da Costa Lima-Junior J,

Calvo-Calle JM, et al. A chimeric protein-based malaria vaccine candidate induces robust T

cell responses against Plasmodium vivax MSP119. Sci Rep. 2016;6:34527. Epub 20161006.

doi: 10.1038/srep34527. PubMed PMID: 27708348; PubMed Central PMCID:

PMCPMC5052570.

79.   Ford A, Kepple D, Abagero BR, Connors J, Pearson R, Auburn S, et al. Whole

genome sequencing of Plasmodium vivax isolates reveals frequent sequence and structural

polymorphisms in erythrocyte binding genes. PLoS Negl Trop Dis. 2020;14(10):e0008234.

Epub 20201012. doi: 10.1371/journal.pntd.0008234. PubMed PMID: 33044985; PubMed

Central PMCID: PMCPMC7581005.

80.   Kar S, Sinha A. Plasmodium vivax Duffy Binding Protein-Based Vaccine: a Distant

Dream. Front Cell Infect Microbiol. 2022;12:916702. Epub 20220713. doi:

10.3389/fcimb.2022.916702. PubMed PMID: 35909975; PubMed Central PMCID:

PMCPMC9325973.

81.   Jemmely NY, Niang M, Preiser PR. Small variant surface antigens and Plasmodium

evasion of immunity. Future Microbiol. 2010;5(4):663-82. doi: 10.2217/fmb.10.21. PubMed

PMID: 20353305.

82.   Fernandez-Becerra C, Bernabeu M, Castellanos A, Correa BR, Obadia T, Ramirez M,

et al. Plasmodium vivax spleen-dependent genes encode antigens associated with

cytoadhesion and clinical protection. Proc Natl Acad Sci U S A. 2020;117(23):13056-65.

Epub 20200521. doi: 10.1073/pnas.1920596117. PubMed PMID: 32439708; PubMed Central

PMCID: PMCPMC7293605.

83.   NRC. Guide for the Care and Use of Laboratory Animals. (US) NAP, editor.

Washington DC: National Research Council (US) Institute for Laboratory Animal Research;

1996.

84.   Ma NS, Rossan RN, Kelley ST, Harper JS, Bedard MT, Jones TC. Banding patterns

of the chromosomes of two new karyotypes of the owl monkey, Aotus, captured in Panama.

Journal of medical primatology. 1978;7(3):146-55. Epub 1978/01/01. PubMed PMID:

101668.

85.   Obaldia N. Long-term effect of a simple nest-box on the reproductive efficiency and

other life traits of an Aotus lemurinus lemurinus monkey colony: an animal model for malaria

research. J Med Primatol. 2011. Epub June 18, 2011.

86.   Cooper RD. Studies of a chloroquine-resistant strain of Plasmodium vivax from

Papua New Guinea in Aotus and Anopheles farauti s.l. J Parasitol. 1994;80(5):789-95. Epub

1994/10/01. PubMed PMID: 7931914.

87.   Sullivan JS, Morris CL, Richardson BB, Galland GG, Jennings VM, Kendall J, et al.

Adaptation of the AMRU-1 strain of Plasmodium vivax to Aotus and Saimiri monkeys and to

four species of anopheline mosquitoes. J Parasitol. 1999;85(4):672-7. Epub 1999/08/26.

PubMed PMID: 10461947.

88.   Obaldia N, Rossan R, Cooper R, Kyle D, Nuzum E, Rieckmann K, et al. WR 238605,

chloroquine, and their combinations as blood schizonticides against a chloroquine-resistant

strain of Plasmodium vivax in Aotus monkeys. Am J Trop Med Hyg. 1997;56:508-10.

89.   Earle WC, Perez M. Enumeration of parasites in the blood of malarial patients. J Lab

Clin Med. 1932:1124-30.

90.   Andrysiak PM, Collins WE, Campbell GH. Concentration of Plasmodium ovale- and

Plasmodium vivax-infected erythrocytes from nonhuman primate blood using Percoll

gradients. Am J Trop Med Hyg. 1986;35(2):251-4. Epub 1986/03/01. PubMed PMID:

3006527.

91.   Zhou AE, Berry AA, Bailey JA, Pike A, Dara A, Agrawal S, et al. Antibodies to

Peptides in Semiconserved Domains of RIFINs and STEVORs Correlate with Malaria

Exposure. mSphere. 2019;4(2). Epub 20190320. doi: 10.1128/mSphere.00097-19. PubMed

PMID: 30894432; PubMed Central PMCID: PMCPMC6429043.

92.   van den Hoogen LL, Walk J, Oulton T, Reuling IJ, Reiling L, Beeson JG, et al.

Antibody Responses to Antigenic Targets of Recent Exposure Are Associated With Low-

Density Parasitemia in Controlled Human Plasmodium falciparum Infections. Front

Microbiol. 2018;9:3300. Epub 20190116. doi: 10.3389/fmicb.2018.03300. PubMed PMID:

30700984; PubMed Central PMCID: PMCPMC6343524.

93.   Oyola SO, Ariani CV, Hamilton WL, Kekre M, Amenga-Etego LN, Ghansah A, et al.

Whole genome sequencing of Plasmodium falciparum from dried blood spots using selective

whole genome amplification. Malar J. 2016;15(1):597. Epub 20161220. doi: 10.1186/s12936-

016-1641-7. PubMed PMID: 27998271; PubMed Central PMCID: PMCPMC5175302.

94.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu:

scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation.

Genome Res. 2017;27(5):722-36. Epub 20170315. doi: 10.1101/gr.215087.116. PubMed

PMID: 28298431; PubMed Central PMCID: PMCPMC5411767.

95.   Hu G, Llinas M, Li J, Preiser PR, Bozdech Z. Selection of long oligonucleotides for

gene expression microarrays using weighted rank-sum strategy. BMC Bioinformatics.

2007;8:350. Epub 20070919. doi: 10.1186/1471-2105-8-350. PubMed PMID: 17880708;

PubMed Central PMCID: PMCPMC2099447.

96.   Bozdech Z, Mok S, Gupta AP. DNA microarray-based genome-wide analyses of

Plasmodium parasites. Methods Mol Biol. 2013;923:189-211. doi: 10.1007/978-1-62703-

026-7_13. PubMed PMID: 22990779.

97.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers

differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids

Res. 2015;43(7):e47. Epub 20150120. doi: 10.1093/nar/gkv007. PubMed PMID: 25605792;

PubMed Central PMCID: PMCPMC4402510.

Figures and legends

Fig. 1. Experimental timeline, parasite dynamics and survival analysis.

A. Experimental timeline of infection and challenge. *: died of malaria unrelated causes. **:

anemia and renal failure. **B**. Peripheral parasitemia across the experiment. Panels I-III show

individual parasitemia of *Aotus* monkeys repeatedly infected with *P. vivax* SAL-1

(inoculations I to III). Panel IV shows *Aotus* challenged with *P. vivax* AMRU-1 (inoculation

987 IV). Inoculated control animals were treated at peak parasitemia. **C**. Probability of no

988 treatment of *Aotus* repeatedly infected with the homologous *P. vivax* SAL-1 and heterologous

989 *P. vivax* AMRU-1 strains at each inoculation level. *p* values for survival curve comparison

990 were obtained using the Log-rank (Mantel-Cox) test. Survival curves for homologous

991 infection 1 shown in blue; homologous infection 2 shown in red; homologous infection 3

992 shown in green. *P. vivax* AMRU-1 heterologous infection 4 shown in black. CBC: red blood

993 cell count. CQ: chloroquine, at 15 mg/kg oral for 3 days. MQ: mefloquine, at 25 mg/kg oral

994 once. C: malaria naïve control. C1: control, once inoculated with *P. vivax*. PI: post

995 inoculation.

Fig. 2. Hematological and parasite parameters. Panels **A-C** show hematocrit levels (Hct%)

(**A**), platelet counts (**B**) and combined data from A, B and mean parasitemia (**C**) across

inoculation levels I to IV. Panel **D** shows the percentage of reticulocytes and the Reticulocyte

Production Index (RPI) at infection level IV. RPI = Reticulocyte Absolute Count/

Reticulocyte Maturation Correction. Reticulocyte Absolute Count = Hct% / 45 x

Reticulocyte %. T = CQ: chloroquine, at 15 mg/kg oral for 3 days; MQ at 25 mg/kg once for

rescue treatment of *P. vivax* AMRU-1 infections in panel C.

1005

Fig. 3. ELISA titers of *Aotus* repeatedly infected with *P. vivax* blood stages. A. Crude

antigen checkerboard titration. *P. vivax* SAL-1 antigen was prepared from *Aotus* iRBCs

purified by Percoll cushion (47%) centrifugation and adsorbed to the plate wells diluted in

PBS pH 7.4 at a concentration of 5 µg / mL. Secondary antibodies (peroxidase conjugated

Goat anti-monkey, Rhesus macaque) was diluted 1:2000 in PBS pH 7.4., and optical density

(OD) read using a 492 nm filter. **B.** Mean ELISA* titers of *Aotus* immunized by repeated

infection with the homologous SAL-1 and challenged with the heterologous AMRU-1 strains

of *P. vivax*. I-IV indicates inoculation level, each with inoculum of 50 x 10^3 iRBCs. Level I-

III infection with homologous SAL-1. Level IV indicates infection with heterologous

AMRU-1. **C.** Pearson correlation analysis of mean ELISA titers at inoculation levels I (n =

6), II (n = 5) and III (n = 4) showed a high negative correlation vs mean (*r* – 0.98) and

cumulative (AUC) parasitemia (r = -0.98), and a moderate positive correlation vs peak

parasitemia (r = 0.63). **D.** Combined plot of mean parasitemia and ELISA titers with *Aotus*

repeatedly infected with the homologous SAL-1 (Infection I-III) and challenged with the

heterologous AMRU-1 (Infection IV).

1021

1022

Fig. 4. **Protein microarray**. **A**. Shown are antibody responses (log2(antigen reactivity / no

DNA control reactivity)) to 66 out of 244 *P. vivax* IVTT antigens with reactivity above 0 in

10% of all samples. Thus, zero represents equal or lower reactivity than the mean of the no

DNA control spots. Antigens are ordered from highest to lowest overall mean. Samples are

ordered top to bottom by inoculation level, day, and then by monkey. **B**. Antigen breadth

(number of *P. vivax* reactive antigens) by post-infection day at each inoculation level (I-IV).

Antigens were considered reactive if the reactivity was higher than the mean + 3SD of the no

DNA control spots for that sample. * indicates a significantly higher antigen breadth at that

day than at baseline (day -1) within each inoculation ($p<0.05$, Wilcoxon matched pairs test,

one-sided). **C.** Area under the curve (AUC) of the antigen breadth at each inoculation level

for the three monkeys that completed the experiment. **D.** Pearson correlation of ELISA titer at

each day post-infection versus antigen breadth. *p* values shown are from t-tests with the null

hypothesis that the correlation coefficient equals 0.

1036

1037

Fig. 5. PacBio Whole Genome Sequencing (WGA) of *Plasmodium vivax* SAL-1 and

AMRU-1. A. Artemis screenshot. Shown is one arm of *P. vivax* PvP01 chromosome 14, with

PacBio reads mapped (SAL-1 in blue and AMRU-1 in red). Most of the coverage occurs in

subtelomeric regions, demonstrating the specificity of the sWGA. **B**. Circos plot of one

representative SAL-1 contig that contains mostly *pir* genes. The contig maps to chromosome

1 of *P. vivax* reference PvP01. Gray lines show syntenic matches of *pir* genes between the

two strains. **C.** Gephi plot showing *pir* genes from AMRU-1 (red), SAL-1 (blue) and PvP01

reference (gray). Genes are connected if they share at least 32% global identity.

1046

1047

1048 **Fig. 6**. **Parasite gene expression comparisons across infection regimes. A.** Differential

1049 gene expression (DGE) across core genes. Volcano plots show DGE between infection

regimes. *Left*: DGE of core genes between SAL-1 inoculation II vs SAL-1 inoculation I

(black) and between AMRU-1 inoculation IV vs the averaged expression of SAL-1 during

the homologous challenges (blue). *Right*: DGE of core genes across AMRU-I infection

regimes. Yellow dots represent DGE between AMRU-1 parasites from *Aotus* monkeys

previously infected with three SAL-1 inoculations (AMRU3Sal) vs AMRU-1 parasites from

naïve *Aotus* monkeys (AMRUNaive). Black dots represent DGE between AMRU3Sal vs

AMRU-1 parasites from *Aotus* monkeys previously infected with only one SAL-1 inoculation

(AMRU1Sal). Blue dots represent DGE between AMRU1Sal vs AMRUNaive. Each dot

represents one annotated *P. vivax* core gene and is displayed according to the fold-change in

expression (x-axis, in log2) and statistical significance (y-axis, in negative logarithm to the

base 10 of the *p*-value). **B.** Principal Component Analysis (PCA) of the parasite core gene

(left panel) and *pir* gene (right panel) expression profiles from each biological replicate,

coloured according to the corresponding group: SAL-1 parasites at day 14 PI of the first

inoculation (gray); SAL-1 parasites at day 14 PI of the second inoculation (black); AMRU-1

parasites at day 14 PI from *Aotus* monkeys previously infected with three SAL-1 inoculations

(light blue dots); gene expression of AMRU-1 parasites at day 28 PI from *Aotus* monkeys

previously infected with three SAL-1 inoculations (blue dots); gene expression of AMRU-1

parasites at day 1 PI from naïve *Aotus* monkeys (orange dots); gene expression of AMRU-1

parasites at day 14 PI from naïve *Aotus* monkeys (red dots). **C.** *pir* gene network analysis

comparing *P. vivax pir* gene expression in SAL-1 *vs* AMRU-1 infections in *Aotus* monkeys.

Same network as in **Figure 5C**, except that larger circles indicate *pir* gene expression level.

Left panel: *pir* expression in SAL-1 parasites at day 14 PI of the first inoculation across

individual monkeys. Right panel: comparison of *pir* expression in monkey MN32047

between SAL-1 parasites at day 14 PI of the first inoculation AMRU-1 parasites at day 14 PI

of the fourth inoculation.

1075 **Supplementary figures and legends**

1076

1077 **Fig. S1. Experimental scheme**. Diagram depicting repeated infection of *Aotus* monkeys with

1078 the homologous *P. vivax* SAL-1 and challenge with heterologous AMRU-1 strain.

1079 Inoculation level, inoculation day, donor monkey, monkey number, and number of animals

1080 remaining from the original group of six inoculated are shown.

1081

Figure S2. Parasite load and biomass across animals. A Parasite load. Panels I-III show

parasite load (qPCR 18sRNA in copies x µL) across inoculation levels I-III in individual

monkeys infected with *P. vivax* SAL-1. Panel IV shows inoculation level IV, i.e., individual

monkeys infected with *P. vivax* AMRU-1. **B**. Parasite biomass. Panels I-III show parasite

biomass (pLDH ng/mL) across inoculation levels I-III in individual monkeys infected with *P.*

vivax SAL-1. Panel IV shows inoculation level IV, i.e., individual monkeys infected with *P.*

vivax AMRU-1. **C.** Parasitemia parameters across inoculation levels I-IV (Mean ± SD)**.** Left:

Days patent. Mid left: Day of Peak. Mid right: Peak parasitemia. Right: Area under the curve

(AUC). *P* value; unpaired t-test with equal standard deviation.

1094

Fig. S3. Protein microarray and ELISA titer. Pearson correlation of ELISA titer at each

day post-infection vs antigen breadth. *p* values shown are from t-tests with the null

hypothesis that the correlation coefficient equals 0.

1098

1099

Fig. S4. Protein microarray individual responses. Reactivity of *Aotus* sera repeatedly infected with *P. vivax* blood stages against selected immunogenic targets. PVX_090230 = ETRAMP; PVX_099980_s3 & PVX_099980_s4 = MSP1; PVX_092070 = PV1. All data is log2(antigen reactivity / no DNA control reactivity). Notably, the dynamic of antibody acquisition varies across individual antigens and monkeys.

Supplementary tables

Table S1. Summary hematological and parasitemia values of *Aotus* repeatedly infected with *P. vivax* SAL-1 (inoculation levels I-III) and challenged with the heterologous *P. vivax* strain AMRU-1 (inoculation level IV).

Table S2. Mean ELISA titers of *Aotus* repeatedly infected with *P. vivax* SAL-1 (inoculation levels I-III) and challenged with the heterologous *P. vivax* strain AMRU-1 (inoculation level IV).

Table S3. *P. vivax* immunogenic targets with significantly higher antibody levels at inoculation level III vs inoculation level I.

Table S4: Normalised microarray data across monkeys and time points.

Table S5. Microarray probe set for SAL-1 and AMRU-1.