Conflicts are represented in a cognitive space to reconcile domain-general and domain-specific cognitive control

Guochun Yang,[1,2,3,4] Haiyan Wu,[5] Qi Li,[6] Xun Liu,[1,2*] Zhongzheng Fu,[7,8] Jiefeng Jiang[3,4]

[1] CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing 100101, China.

[2] Department of Psychology, University of Chinese Academy of Sciences, Beijing 100101, China.

[3] Department of Psychological and Brain Sciences, University of Iowa, Iowa City, IA 52242, USA.

[4] Cognitive Control Collaborative, University of Iowa, Iowa City, IA 52242, USA.

[5] Centre for Cognitive and Brain Sciences and Department of Psychology, University of Macau, Taipa, Macau 999078, China.

[6] Beijing Key Laboratory of Learning and Cognition, School of Psychology, Capital Normal University, Beijing 100048, China.

[7] Department of Neurosurgery, Cedars-Sinai Medical Center, Los Angeles, CA 90048, USA.

[8] Division of Humanities and Social Sciences, California Institute of Technology, Pasadena, CA 91125, USA.

*Correspondence: liux@psych.ac.cn

Abstract
Cognitive control resolves conflict between task-relevant and -irrelevant information
to enable goal-directed behavior. As conflict can arise from different sources (e.g.,
sensory input, internal representations), how a finite set of cognitive control processes
can effectively address huge array of conflict remains a major challenge. We
hypothesize that different conflict can be parameterized and represented as distinct
points in a (low-dimensional) cognitive space, which can then be resolved by a
limited set of cognitive control processes working along the dimensions. To test this
hypothesis, we designed a task with five types of conflict that could be conceptually
parameterized along one dimension. Over two experiments, both human performance
and fMRI activity patterns in the right dorsolateral prefrontal (dlPFC) support that
different types of conflict are organized in a cognitive space. The findings suggest that
cognitive space can be a dimension reduction tool to effectively organize neural
representations of conflict for cognitive control.
36
37 Keywords: cognitive control, cognitive space, domain-general, domain-specific,
38 conflict
39

Introduction

Cognitive control enables humans to behave purposefully by modulating neural
processing to resolve conflict between task-relevant and task-irrelevant information.
For example, when naming the color of the word "BLUE" printed in red ink, we are
likely to be distracted by the word meaning, because reading a word is highly
automatic in daily life. To keep our attention on the color, we need to mobilize the
cognitive control processes to resolve the conflict between the color and word by
boosting/suppressing the processing of color/word meaning. As task-relevant and
task-irrelevant information can come from different sources, the sources of conflict
and how they should be resolved can vary greatly[1]. For example, conflict may occur
between items of sensory information, such as between a red light and a police officer
signaling cars to pass. Alternatively, conflict may occur between sensory and motor
information, such as when a voice on the left asks you to turn right. The large variety
of conflict sources implies that there may be unlimited number of conflicts. A key
unsolved question in cognitive control is how our brain efficiently resolves a nearly
infinite number of different types of conflict.
A first step to addressing this question is to examine the commonalities and/or
dissociations across different types of conflict that can be categorized into different
domains. Examples of the domains of conflict include experimental paradigm[2,3],
sensory modality[4,5], or conflict type regarding the dimensional overlap of conflict
processes[6,7].
Two solutions to resolving a wide range of conflict types are proposed. They
differ based on whether the same cognitive control mechanisms are applied across
domains. On the one hand, the *domain-general* cognitive control theories posit that

the frontoparietal cortex adaptively encodes task information and can thus flexibly
implement control strategies for different types of conflict. This is supported by the
generalizable control adjustment (i.e., encountering a conflict trial from one type can
facilitate conflict resolution of another type)[2,8] and similar neural patterns[9,10] across
distinct conflict tasks. A broader domain-general view holds that the frontoparietal
brain regions/networks are widely involved in multiple control demands well beyond
the conflict domain[11,12], which explains the remarkable flexibility in human behaviors.
However, since domain-general processes are by definition likely shared by different
tasks, when we need to handle multiple task demands at the same time, the efficiency
of both tasks would be impaired due to resource competition or interference[13].
Therefore, the domain-general processes is evolutionarily less advantageous for
humans to deal with the diverse situations requiring high efficiency[14]. On the other
hand, the *domain-specific* theories argue that different types of conflict are handled by
distinct cognitive control processes (e.g., where and how information processing
should be modulated)[15,16]. However, according to the domain-specific view, the
potentially unlimited conflict situations require a large variety of preexisting control
processes, which is biologically implausible[17].
To reconcile the two theories, researchers recently proposed that cognitive control
might be a mixture of domain-general and domain-specific processes. For instance,
Freitas et al.[18] found that trial-by-trial adjustment of control can generalize across two
conflict domains to different degrees, leading to domain-general (strong
generalization) or domain-specific (weak or no generalization) conclusions depending
on the task settings of the consecutive conflict. Similarly, different brain networks
may show domain-generality (i.e., representing multiple conflicts) or domain-
specificity (i.e., representing individual conflicts separately)[7,19]. Even within the same
brain area (e.g., medial frontal cortex), Fu et al.[20] found that the neural population
activity can be factorized into orthogonal dimensions encoding both domain-general
and domain-specific conflict information, which can be selectively read out by
downstream brain regions. While the mixture view provides an explanation for the
contradictory findings[21], it suffers the same criticism as domain-specific cognitive
control theories, as it still requires unlimited cognitive control processes to fully cover
all possible conflicts.
A key to reconciling domain-general and domain-specific cognitive control is to
organize the nearly infinite possible types of conflict using a system with limited,
dissociable dimensions. A construct with a similar function is the *cognitive space*[22],
which extends the idea of cognitive map[23] to the representation of abstract
information. Critically, the cognitive space view holds that the representations of
different abstract information are organized continuously and the locations of
representations in the cognitive space are determined by the similarity among the
represented information[22].
In the human brain, it has been shown that abstract[23,24] and social[25] information
can be represented in a cognitive space. For example, social hierarchies with two
independent scores (e.g., popularity and competence) can be represented in a 2D
cognitive space (one dimension for each score), such that each social item can be

located by its score in the two dimensions[25]. In the field of cognitive control, recent
studies have begun to conceptualize different control states within a cognitive space[26].
For example, Fu et al.[20] mapped different conflict conditions to locations in a
low/high dimensional cognitive space to demonstrate the domain-general/domain-
specific problems; Grahek et al.[27] used a cognitive space model of cognitive control
settings to explain behavioral changes in the speed-accuracy tradeoff. However, the
cognitive spaces proposed in these studies were only applicable to a limited number
of control states involved in their designs. Therefore, it remains unclear whether there
is a cognitive space that can explain an unlimited number of control states, similar to
that of the spatial location[22] and non-spatial knowledge[23]. A challenge to answering
this question lies in how to construct control states with continuous levels of
similarity. Our recent work[28] showed that it is possible to manipulate continuous
conflict similarity by using a mixture of two independent conflict types with varying
ratios, which can be used to further examine the behavioral and neural evidence for
the cognitive space view. It is also unclear how the cognitive space of cognitive
control is encoded in the brain, although that of spatial locations and non-spatial
abstract knowledge has been relatively well investigated in the medial temporal lobe,
medial prefrontal and orbitofrontal system[22,23]. Recent research has suggested that the
abstract task structure could be encoded and implemented by the frontoparietal
network[29,30], but whether a similar neural system encodes the cognitive space of
cognitive control remains untested.
We hypothesize that different types of conflict are represented as points in a
cognitive space. The dimensions in the cognitive space of conflict can be the
aforementioned *domains*, in which domain-specific cognitive control processes are
defined. For a specific type of conflict, its location in the cognitive space can be
parameterized using a limited number of coordinates, which reflect how much control
is needed for each of the domain-specific cognitive control processes. The cognitive
space can also represent different types of conflict with low dimensionality[26,31].
Different domains can be represented conjunctively in a single cognitive space to
achieve domain-general cognitive control, as conflict from different sources can be
resolved using the same set of cognitive control processes. We further hypothesize
that the cognitive space representing different types of conflict may be located in the
frontoparietal network due to its essential roles in conflict resolution[20,32] and abstract
task representation[30].
In this study, we adjusted the paradigm from our previous study[28] by including
transitions of trials from five different conflict types, which enabled us to test if these
conflict types are organized in a cognitive space (Fig. 1A). Specifically, on each trial,
an arrow, pointing either upwards or downwards, was presented on one of the 10
possible locations on the screen. Participants were required to respond to the pointing
direction of the arrow (up or down) by pressing either the left or right key.
Importantly, conflict from two sources can occur in this task. On one hand, the
vertical location of the arrow can be incongruent with the direction (e.g., an up-
pointing arrow on the lower half of the screen), resulting spatial Stroop conflict[6,33].
On the other hand, the horizontal location of the arrow can be incongruent with the

response key (e.g., an arrow requiring left response presented on the right side of the
screen), thus causing Simon conflict[33,34]. As the arrow location rotates from the
horizontal axis to the vertical axis, spatial Stroop conflict increases, and Simon
conflict decreases. Therefore, the 10 possible locations of the arrow give rise to five
conflict types with unique blend of spatial Stroop and Simon conflict[28]. As the
increase in spatial Stroop conflict is perfectly correlated with the decrease in Simon
conflict, we can use a 1D cognitive space to represent all five conflict types.

Fig. 1. Experimental design. (A) The left panel shows the orthogonal stimulus-response mappings of
the two participant groups. In each group the stimuli were only displayed at two quadrants of the
circular locations. One group were asked to respond with the left button to the upward arrow and with
the right button to the downward arrow presented in the to-left and bottom-right quadrants, and the
other group vice versa. The right panel shows the time course of one example trial. The stimuli were
displayed for 600 ms, preceded and followed by fixation crosses that lasted for 1400 ms in total. (B)
Examples of the five types of conflict, each containing congruent and incongruent conditions. The
arrows were presented at locations along five orientations with isometric polar angles, in which the
vertical location introduces the spatial Stroop conflict, and the horizontal location introduces the Simon
conflict. Dashed lines are shown only to indicate the location of arrows and were not shown in the
experiments. (C) The definition of the angular difference between two conflict types and the conflict
similarity. The angle θ is determined by the acute angle between two lines that cross the stimuli and the
central fixation. Therefore, stimuli of the same conflict type form the smallest angle of 0, and stimuli

between Conflict 1 and Conflict 5 form the largest angle of 90°, and others are in between. Conflict
similarity is defined by the cosine value of θ.

One way to parameterize (i.e., defining a coordinate system) the cognitive space
is to encode each conflict type by the angle of the axis connecting its two possible
stimulus locations (Fig. 1B). Within this cognitive space, the similarity between two
conflict types can be quantified as the cosine value of their angular difference (Fig.
1C). If the conflict types are organized as a cognitive space in the brain, the similarity
between conflict types in the cognitive space should be reflected in both the behavior
and similarity in the neural representations of conflict types. Our data from two
experiments using this experimental design support both predictions: using behavioral
data, we found that the influence of congruency (i.e., whether the task-relevant and
task-irrelevant information indicate the same response) from the previous trial to the
next trial increases with the conflict similarity between the two trials. Using fMRI
data, we found that more similar conflict showed higher multivariate pattern similarity
in the right dorsolateral prefrontal cortex (dlPFC).

Results

Conflict type similarity modulated behavioral congruency sequence effect (CSE)

Experiment 1.
We conducted a behavioral experiment (n = 33, 18 females) to examine how CSEs
across different conflict types are influenced by their similarity. First, we validated the
experimental design by testing the congruency effects. All five conflict types showed
robust congruency effects such that the incongruent trials were slower and less
accurate than the congruent trials (Note S1; Fig. S1 A/B). To test the influence of
similarity between conflict types on behavior, we examined the CSE in consecutive
trials. Specifically, the CSE was quantified as the interaction between previous and
current trial congruency and can reflect how (in)congruency on the previous trial
influences cognitive control on the current trial[35,36]. It has been shown that the CSE
diminishes if the two consecutive trials have different conflict types[37-39]. Similarly, we
tested whether the size of CSE increases as a function of conflict similarity between
consecutive trials. To this end, we organized trials based on a 5 (previous trial conflict
type) × 5 (current trial conflict type) × 2 (previous trial congruency) × 2 (current trial
congruency) factorial design, with the first two and the last two factors capturing
between-trial conflict similarity and the CSE, respectively. The cells in the 5 × 5
matrix were mapped to different similarity levels according to the angular difference
between the two conflict types (Fig. 1C). As shown in Fig. 2, the CSE, measured in
both reaction time (RT) and error rate (ER), scaled with conflict similarity.
To test the modulation of conflict similarity on the CSE, we constructed a linear
mixed effect model to predict RT/ER in each cell of the factorial design using a
predictor encoding the interaction between the CSE and conflict similarity (see
Methods). The results showed a significant effect of conflict similarity (RT: $\beta = 0.10$

± 0.01, $t(1978) = 15.82$, $p < .001$, $\eta_p^2 = .120$; ER: $\beta = 0.15 \pm 0.02$, $t(1978) = 7.84$, $p <$
$.001$, $\eta_p^2 = .085$, Fig. S2B/E). In other words, the CSE increased with the conflict
similarity between two consecutive trials. The conflict similarity modulation effect
remained significant after regressing out the influence of physical proximity between
the stimuli of consecutive trials (Note S2). As a control analysis, we also compared
this approach to a two-stage analysis that first calculated the CSE for each previous $\times$
current trial conflict type condition and then tested the modulation of conflict
similarity on the CSEs[28]. The two-stage analysis also showed a strong effect of
conflict similarity (RT: $\beta = 0.58 \pm 0.04$, $t(493) = 14.74$, $p < .001$, $\eta_p^2 = .383$; ER: $\beta =$
0.36 ± 0.05, $t(493) = 7.01$, $p < .001$, $\eta_p^2 = .321$, Fig. S2A/D). Importantly, individual
modulation effects of conflict similarity were positively correlated between the two
approaches (RT: $r = 0.48$; ER: $r = 0.86$, both $ps < 0.003$, one-tailed), indicating the
consistency of the estimated conflict similarity effects across the two approaches.

Fig. 2. The conflict similarity modulation on the behavioral CSE in Experiment 1. (A) RT and (B)
ER are plotted as a function of congruency types on trial n−1 and trial n. Each column shows one
similarity level, as indicated by the defined angular difference between two conflict types. Error bars
are standard errors. C = congruent; I = incongruent; RT = reaction time; ER = error rate.

Experiment 2.
Behavioral results. We next conducted an fMRI experiment using a shorter version of
the same task with a different sample (n = 35, 17 females) to seek neural evidence of
how different conflict types are organized. Using behavioral data, we first validated
the experimental design by testing congruency effects in each of the five conflict

types (Note S1; Fig. S1 C/D). We then tested the modulation of conflict similarity on
the behavioral CSE using the linear mixed effect model as in Experiment 1 (except
the two-stage method). Results showed a significant effect of conflict similarity
modulation (RT: $\beta = 0.24 \pm 0.04$, $t(1148) = 6.36$, $p < .001$, $\eta_p^2 = .096$; ER: $\beta = 0.33 \pm$
0.06, $t(1206) = 5.81$, $p < .001$, $\eta_p^2 = .124$, Fig. S2C/F), thus replicating the results of
Experimental 1 and setting the stage for fMRI analysis. As in Experiment 1, the
conflict similarity modulation effect remained significant after regressing out the
influence of physical proximity between the stimuli of consecutive trials (Note S2).

Brain activations modulated by conflict type dissimilarity with univariate analyses

Fig. 3. The congruency effect and parametric modulation effect detected by uni-voxel analyses.
Results displayed are thresholded with voxel-wise one-tailed $p < .005$ and cluster-size > 20 voxels. The
congruency effect denotes the higher activation in incongruent than congruent condition. The positive
parametric modulation effect (I_pm – C_pm) denotes the higher activation when the conflict type
contained a higher ratio of Simon conflict component (bottom left panel). The negative parametric
modulation effect [converted to positive with – (I_pm – C_pm)] denotes the higher activation when the
conflict type contained a higher ratio of spatial Stroop conflict component (bottom right panel). I =
incongruent; C = congruent; pm = parametric modulator.

In the fMRI analysis, we first replicated the classic congruency effect by searching for
brain regions showing higher univariate activation in incongruent than congruent
conditions (GLM1, see Methods). Consistent with the literature[20,40], this effect was
observed in the pre-supplementary motor area (pre-SMA) and anterior cingulate
cortex (ACC) areas (Fig. 3, Table S1). We then tested the encoding of conflict type as

a cognitive space by identifying brain regions with activation levels parametrically
covarying with the coordinates (i.e., axial angle relative to the horizontal axis) in the
hypothesized cognitive space. As shown in Fig. 1B, change in the angle corresponds
to change in spatial Stroop and Simon conflicts in opposite directions. Accordingly, in
the left middle frontal gyrus (MFG), fMRI activation scaled with the increase in
spatial Stroop conflict, whereas the right inferior parietal sulcus (IPS) and the right
dorsomedial prefrontal cortex (dmPFC) displayed positive correlation between fMRI
activation and Simon conflict (Fig. 3, Fig. S3, Table S1).
To further test if the univariate results explain the conflict similarity modulation
of the behavioral CSE (slope in Fig. S2C), we conducted brain-behavioral correlation
analyses for regions identified above. Regions with higher spatial Stroop/Simon
modulation effects were expected to trigger higher behavioral conflict similarity
modulation effect on the CSE. However, none of the three regions (i.e., left MFG,
right IPS and right dmPFC, Fig. 3) were positively correlated with the behavioral
performance, all $p_{FDR} > .762$, one-tailed. In addition, since the conflict type difference
covaries with the orientation of the arrow location at the individual level (e.g., the
stimulus in a higher level of Simon conflict is always closer to the horizontal axis, see
Fig. S4), the univariate modulation effects may not reflect purely conflict type
difference. To further tease these factors apart, we used multivariate analyses.

Multivariate patterns of the right dlPFC encodes the conflict similarity

283
Fig. 4. The conflict type effect. (A) Brain regions surviving the FDR-correction ($p_{FDR} < 0.05$ and $p <$
0.001) across the 360 regions (criterion 1). Labeled regions are those meeting the criterion 2. (B) The
regions showing stronger encoding of conflict type in the incongruent than congruent conditions

(criterion 2). ** $p_{FDR} < .01$, *** $p_{FDR} < .001$. (C) The brain-behavior correlation of the right 8C

(criterion 3). (D) Illustration of the different encoding strength of conflict type similarity in incongruent

versus congruent conditions of right 8C. l = left; r = right.

The hypothesis that the brain encodes conflict types in a cognitive space predicts that

similar conflict types will have similar neural representations. To test this prediction,

we computed the representational similarity matrix (RSM) that encoded correlations

of blood-oxygen-level dependent (BOLD) signal patterns between each pair of

conflict type (conflict 1, 2, 3, 4 and 5, as shown in Fig. 1B) × congruency (congruent,

incongruent) × arrow direction (up, down) × run × subject combinations for each of

the 360 cortical regions from the Multi-Modal Parcellation (MMP) cortical atlas[41,42].

The RSM was then submitted to a linear mixed-effect model as the dependent

variable to test whether the representational similarity in each region was modulated

by various experimental variables (e.g., conflict type, spatial orientation, stimulus,

response, etc., see Methods). The linear mixed-effect model was used to de-correlate

conflict type and spatial orientation leveraging the between-subject manipulation of

stimulus locations (Fig. S4).

To validate this method, we applied this analysis to test the effects of

response/stimulus features and found that representational similarity of the BOLD

signal significantly covaried with whether two response/spatial location/arrow

directions were the same most strongly in bilateral motor/visual/somatosensory areas,

respectively (Fig. S5). We then identified the cortical regions encoding conflict type

as a cognitive space by testing whether their RSMs can be explained by the similarity

between conflict types. Specifically, we applied three independent criteria: (1) the

cortical regions should exhibit a statistically significant positive conflict similarity

effect on the RSM; (2) the conflict similarity effect should be stronger in incongruent

than congruent trials to reflect flexible adjustment of cognitive control demand when

conflict is present; and (3) the conflict similarity effect should be positively correlated

with the behavioral conflict similarity modulation effect on the CSE (see *Behavioral*

Results of Experiment 2). The first criterion revealed several cortical regions encoding

the conflict similarity, including the 8C area (a subregion of dlPFC[42]), a47r, TPOJ3

and V3CD in the right hemisphere, and the 6r, 7Am, 24dd, VMV1, VMV2, 7Pl, 23c

and 25 areas in the left hemisphere (p_{FDR}s < 0.05, with raw ps < 0.001, one-tailed, Fig.

4A). We next tested whether these regions were related to cognitive control by

comparing the strength of conflict similarity effect between incongruent and

congruent conditions (criterion 2). Results revealed that the left lateral area 7P (7P1),

left ventromedial visual area 1 (VMV1), left dorsal area 24d (24dd), right Brodmann

area 8C (8C), and right V3CD met this criterion, p_{FDR}s < .01, one-tailed (Table 1, Fig.

4B), suggesting that the representation of conflict type was strengthened when

conflict was present (e.g., Fig. 4D). The inter-subject brain-behavioral correlation

analysis (criterion 3) showed that the strength of conflict similarity effect on RSM

scaled with the modulation of conflict similarity on the CSE (slope in Fig. S2C) in

right 8C ($r = 0.43$, $p_{FDR} = .027$, one-tailed, Fig. 4C) but not in the other regions (all

$p_{FDR} > .632$, one-tailed). In addition, we did not find evidence supporting the encoding

of congruency in the right 8C area (see Note S5), suggesting that the right 8C area
specifically represents conflict similarity. In sum, we found converging evidence
supporting that the right dlPFC (8C area) encoded conflict similarity, which further
supports the hypothesis that conflict types are represented in a cognitive space.

Multivariate patterns of visual and oculomotor areas encode stimulus orientation

Fig. 5. The axial orientation effect. (A) Brain regions surviving the FDR-correction ($p_{FDR} < 0.05$ and
$p < 0.001$) across the 360 regions (criterion 1). Labeled regions are those meeting the criterion 2. (B)
The regions showing stronger encoding of orientation in the incongruent than congruent conditions
(criterion 2). * $p_{FDR} < .05$, ** $p_{FDR} < .01$, *** $p_{FDR} < .001$.

To tease apart the representation of conflict type from that of perceptual information,
we tested the modulation of the spatial orientations of stimulus locations on RSM
using the aforementioned RSA. We also applied three independent criteria: (1) the
cortical regions should exhibit a statistically significant orientation effect on the RSM;
(2) the conflict similarity effect should be stronger in incongruent than congruent
trials; and (3) the orientation effect should not interact with the CSE, since the
orientation effect was dissociated from the conflict similarity effect and was not
expected to influence cognitive control. We observed increasing fMRI
representational similarity between trials with more similar orientations of stimulus
location in the occipital cortex, such as right V1, bilateral V2 and V3, right V4, left
area temporoparietooccipital junction 3 (TPOJ3) and right PHT areas (FDR corrected
ps < 0.05 and raw ps < 0.001). We also found the same effect in several oculomotor
related regions, including the left frontal eye field (FEF), anterior 6m (6ma), area
intraparietal 2 (IP2), right parietal area F (PF) and bilateral 5m, as well as other
regions (Fig. 5A). Then we tested if any of these brain regions were related to the
conflict representation by comparing their encoding strength between incongruent and
congruent conditions. Results showed that the right V1, bilateral V2, left FEF, left
IP2, right hippocampus (H) and right PF encoded stronger orientation effect in the
incongruent than the congruent condition, p_{FDR}s < .05, one-tailed (Table1, Fig. 5B).
We then tested if any of these regions was related to the behavioral performance, and
results showed that none of them positively correlated with the behavioral conflict
similarity modulation effect, all $p_{FDR} > .675$, one-tailed. Thus all regions are
consistent with the criterion 3. Like the right 8C area, none of the reported areas
directly encoded congruency (see Note S5). Taken together, we found that the visual

and oculomotor regions encoded orientations of stimulus location in a continuous
manner and that the encoding strength was stronger when conflict was present.
To explore the relation between conflict type and orientation representations, we
conducted representational connectivity (i.e., the similarity between two RSMs of two
regions)[43] analyses and found that among the orientation effect regions, the right V1
and bilateral V2 showed significant representational connectivity with the right 8C
compared to the controlled regions (including those encoding orientation effect but
not showing larger encoding strength in incongruent than congruent conditions, as
well as three other regions encoding none of our defined effects in the main RSA, see
Methods). Compared with the largest connectivity strength in the controlled regions
(i.e., the left V3, β = 0.1447 ± 0.0069), we found higher connectivity in the left V2, β
= 0.1645 ± 0.0060, $t(34)$ = 4.86, right V1, β = 0.1628 ± 0.0065, $t(34)$ = 4.54, and
right V2, β = 0.1678 ± 0.0074, $t(34)$ = 5.65, all p_{FDR} < .001, one-tailed (Fig. S6).
379
380 **Table 1. Summary statistics of regions showing larger encoding strength in**
381   **incongruent than congruent conditions for the conflict type and**
382   **orientation effects.**

Region name	$t(34)$	β(SD)	Cohen's d	p_{FDR}
Conflict type effect				
left 7P1	3.13	0.0049 ± 0.0016	0.53	.011
left VMV1	3.96	0.0077 ± 0.0019	0.67	.002
left 24	7.82	0.0094 ± 0.0012	1.32	< .001
right 8C	3.15	0.0073 ± 0.0023	0.53	.011
right V3CD	2.86	0.0057 ± 0.0020	0.48	.017
Orientation effect				
left V2	3.20	0.0107 ± 0.0033	0.54	.007
left FEF	2.97	0.0066 ± 0.0022	0.50	.010
left IP2	5.73	0.0129 ± 0.0022	0.97	.001
right V1	2.70	0.0060 ± 0.0022	0.46	.014
right V2	3.26	0.0083 ± 0.0025	0.55	.007
right H	2.79	0.0037 ± 0.0013	0.47	.014
right PF	5.31	0.0097 ± 0.0018	0.90	< .001

Discussion

Understanding how different types of conflict are resolved is essential to answer how
cognitive control achieves adaptive behavior. However, the dichotomy between
domain-general and/or domain-specific processes presents a dilemma[15,21].
Reconciliation of the two views also suffers from the inability to fully address how
infinite conflict can be resolved by a limited set of cognitive control processes. In this
study, we hypothesized that this issue can be addressed if conflict is organized as a
cognitive space. Leveraging the well-known dissociation between the spatial Stroop
and Simon conflict[44-46], we designed five conflict types that are systematically

different from each other. The cognitive space hypothesis predicted that the
representational proximity/distance between two conflict types scales with their
similarities/dissimilarities, which was tested at both behavioral and neural levels.
Behaviorally, we found that the CSEs were linearly modulated by conflict similarity
between consecutive trials, replicating and extending our previous study[28]. BOLD
activity patterns in the right dlPFC further showed that the representational similarity
between conflict types was modulated by their conflict similarity, and that strength of
the modulation was positively associated with the modulation of conflict similarity on
the behavioral CSE. We also observed that activity in three brain regions (right IPS,
right dlPFC and left MFG) was parametrically modulated by the conflict type
difference, though they did not directly explain the behavioral results. Additionally,
we found that the visual regions encoded the spatial orientation of the stimulus
location, which might provide the essential concrete information to determine the
conflict type. Together, these results support the hypothesis that the conflicts are
organized in a cognitive space that enables a limited set of cognitive control processes
to resolve infinite possible types of conflict.

Fig. 6. Illustration of the hypothesized dimensionalities of different representations. The shade of
the red color indicates the degree of dimensionality (i.e., how many dimensions are needed to represent
different states). The dimensionality of domain-general representation is extremely low, with all
representations compressed to one dot. The dimensionality of domain-specific representation is
extremely high, with each control state encoded in a unique and orthogonal dimension. The
dimensionality of the organized representation is modest, enabling distant states to be separated but
also allowing close states to share representations. The solid arrows show the axes of different
dimensions. The dashed arrows indicate how the representational dimensionality can be reduced by
projecting the independent dimensions to a common dimension.

Conventionally, the domain-general view of control suggests a common
representation for different types of conflict (Fig. 6, left), while the domain-specific
view suggests dissociated representations for different types (Fig. 6, right). Previous
research on this topic often adopts a binary manipulation of conflict[21] (i.e., each
domain only has one conflict type) and thus is not suitable to test the cognitive space

hypothesis. Here, we parametrically manipulated the similarity of conflict in different
conflict types and demonstrated that the two theories can be reconciled as a cognitive
space[22] (Fig. 6, middle). Specifically, the cognitive space provides a solution to use a
single cognitive space organization to encode different types of conflict that are close
(domain-general) or distant (domain-specific) to each other. It also shows the
potential for how unlimited conflict types can be coded using limited resources (i.e.,
as different points in a low-dimensional cognitive space). Moreover, geometry can
also emerge in the cognitive space[20], which will allow for decomposition of a conflict
type (e.g., how much conflict in each of the dimensions in the cognitive space) so that
it can be mapped into the limited set of cognitive control processes. Such geometry
enables fast learning of cognitive control settings from similar conflict types by
providing a measure of similarity (e.g., as distance in space).
If the dimensionality of the cognitive space of conflict is extremely high, the
cognitive space solution would suffer the same criticism as the domain-specificity
theory. We argue that the dimensionality is manageable for the human brain, as task
information unrelated to differentiating conflicts can be removed. For example, the
Simon conflict can be represented in a space consisting of spatial location, stimulus
information and responses. Thus, the dimensionality of the cognitive space of conflict
should not exceed the number of represented features. The dimensionality can be
further reduced, as humans selectively represent a small number of features when
learning task representations (e.g., spatial information is reduced to the horizontal
dimension from the 3D space we live in)[47]. The reduced dimensionality does not only
require less effort to represent the conflict, but also facilitates generalization of
cognitive control settings among different conflict types[26].
Although our finding of cognitive space in the right dlPFC differs from other
cognitive space studies[24,25,48] that highlighted the orbitofrontal and hippocampus
regions, it is consistent with the cognitive control literature. The prefrontal cortex has
long been believed to be a key region of cognitive control representation[49-51] and is
widely engaged in multiple task demands[12,52]. However, it is not until recently that the
multivariate representation in this region has been examined. For instance, Vaidya et
al.[29] reported that frontal regions presented latent states that are organized
hierarchically. Freund et al.[32] showed that dlPFC encoded the target and congruency
in a typical color-word Stroop task. Taken together, we suggest that the right dlPFC
might flexibly encode a variety of cognitive spaces to meet the dynamic task
demands. In addition, we found no such representation in the left dlPFC (Note S6),
indicating a possible lateralization. Previous studies showed that the left dlPFC was
related to the expectancy-related attentional set up-regulation, while the right dlPFC
was related to the online adjustment of control[53,54], which is consistent with our
findings. Moreover, the right PFC also represents a composition of single rules[55],
which may explain how the spatial Stroop and Simon types can be jointly encoded in
a single space.
We found that participants with stronger conflict representation as cognitive
space in right dlPFC have also adjusted their conflict control to a greater extent based
on the conflict similarity (Fig 4C). The finding suggests that the cognitive space

organization of conflict guides cognitive control to adjust behavior. Previous studies
have shown that participants may adopt different strategies to represent a task, with
the model-based strategies benefitting goal-related behaviors more than the model-
free strategies[56]. Similarly, we propose that the cognitive space could serve as a
mental model to assist fast learning and efficient organization of cognitive control
settings. With the organization of a cognitive space, a new conflict can be quickly
assigned a location in the cognitive space, which will facilitate the development of
cognitive control settings for this conflict by interpolating nearby conflicts and/or
projecting the location to axes representing different cognitive control processes. On
the other hand, without a cognitive space, there would be no measure of similarity
between conflict on different trials, hence limiting the ability of fast learning of
cognitive control setting from similar trials.
The cognitive space in the right dlPFC appears to be an abstraction of concrete
information from the visual regions. We found that the right V1 and bilateral V2
encoded the spatial orientation of the target location (Fig. 5) and showed strong
representational connectivity with the right dlPFC (Fig. S6), suggesting that there
might be information exchange between these regions. We speculate that the
representation of spatial orientation may have provided the essential perceptual
information to determine the conflict type (Fig. 1) and thus served as the critical input
for the cognitive space. The conflict type representation further incorporates the
stimulus-response mapping rules to the spatial orientation representation, so that
vertically symmetric orientations can be recognized as the same conflict type (Fig.
S4). In other words, the representation of conflict type involves the compression of
perceptual information[57], which is consistent with the idea of a low-dimensional
representation of cognitive control[26,31]. The compression and abstraction processes
might be why the frontoparietal regions are the top of hierarchy of information
processing[58] and why the frontoparietal regions are widely engaged in multiple task
demands[59].
With conventional univariate analyses, we observed that the overall congruency
effect was located at the medial frontal regions (i.e., pre-SMA and ACC), which is
consistent with previous studies[20,40]. Beyond that, we also found regions that can be
parametrically modulated by conflict type difference, including right IPS, right dlPFC
(modulated by Simon difference) and left MFG (modulated by spatial Stroop
difference). The lateralization of these regions is consistent with a previous finding[19],
which highlighted the difference of Stroop and Simon types with brain activities at
different hemispheres. The scaling of brain activities based on conflict difference is
potentially important to the representational organization of different types of conflict.
However, we didn't observe their brain-behavioral relevance. One possible reason is
that the conflict (dis)similarity is a combination of (dis)similarity of spatial Stroop and
Simon conflicts, but each univariate region only reflects difference along a single
conflict domain. Also likely, the representational geometry is more of a multivariate
problem than what univariate activities can capture[60]. Future studies may adopt
approaches such as repetition suppression induced fMRI adaptation[26] to test the role
of univariate activities in task representations.

One limitation of this study needs to be noted. To parametrically manipulate the
conflict similarity levels, we adopted the spatial Stroop-Simon paradigm that enables
parametrical combinations of spatial Stroop and Simon conflicts. However, since this
paradigm is a two-alternative forced choice design, the behavioral CSE is not a pure
measure of adjusted control but could be partly confounded by bottom-up factors such
as feature integration[61]. Future studies may replicate our findings with a multiple-
choice design with confound-free trial sequences[62].
In sum, we showed that the cognitive control can be organized in an abstract
cognitive space that is represented in the right dlPFC and guides cognitive control to
adjust goal-directed behavior. The cognitive space hypothesis reconciles the long-
standing debate between the domain-general and domain-specific views of cognitive
control and provides a parsimonious and more broadly applicable framework for
understanding how our brains efficiently and flexibly represents multiple task
settings.
527
528

Materials and Methods

Subjects

In Experiment 1, we enrolled thirty-three college students (19-28 years old, average of
21.5 ± 2.3 years old; 19 males). In Experiment 2, thirty-six college students were
recruited, and one subject was excluded due to not following task instructions. The
final sample of Experiment 2 consisted of thirty-five participants (19-29 years old,
average of 22.3 ± 2.5 years old; 17 males). The sample sizes were determined based
on our previous study[28]. All participants reported no history of psychiatric or
neurological disorders and were right-handed, with normal or corrected-to-normal
vision. The experiments were approved by the Institutional Review Board of the
Institute of Psychology, Chinese Academy of Science. Informed consent was obtained
from all subjects.
541

Method Details

Experiment 1

Experimental Design.   We adopted a modified spatial Stroop-Simon task[28] (Fig. 1).
The task was programmed with the E-prime 2.0 (Psychological Software Tools, Inc.).
The stimulus was an upward or downward black arrow (visual angle of ~ 1°)
displayed on a 17-inch LCD monitor with a viewing distance of ~60 cm. The arrow
appeared inside a grey square at one of ten locations with the same distance from the
center of the screen, including two horizontal (left and right), two vertical (top and
bottom), and six corner (orientations of 22.5°, 45° and 67.5°) locations. The distance
from the arrow to the screen center was approximately 3°. To dissociate orientation of

stimulus locations and conflict types (see below), participants were randomly
assigned to two sets of stimulus locations (one included top-right and bottom-left
quadrants, and the other included top-left and bottom-right quadrants).
Each trial started with a fixation cross displayed in the center for 100−300 ms,
followed by the arrow for 600 ms and another fixation cross for 1100−1300 ms (the
total trial length was fixed at 2000 ms). Participants were instructed to respond to the
pointing direction of the arrow by pressing a left or right button and to ignore its
location. The mapping between the arrow orientations and the response buttons was
counterbalanced across participants. The task design introduced two possible sources
of conflict: on one hand, the direction of the arrow is either congruent or incongruent
with the vertical location of the arrow, thus introducing a spatial Stroop conflict[33,63],
which contains the dimensional overlap between task-relevant stimulus and task-
irrelevant stimulus[1]; on the other hand, the response (left or right button) is either
congruent or incongruent with the horizontal location of the arrow, thus introducing a
Simon conflict[33,34], which contains the dimensional overlap between task-irrelevant
stimulus and response[1]. Therefore, the five polar orientations of the stimulus location
(from 0 to 90°) defined five unique combinations of spatial Stroop and Simon
conflicts, with more similar orientations having more similar composition of conflict.
More generally, the spatial orientation of the arrow location relative to the center of
the screen forms a cognitive space of different blending of spatial Stroop and Simon
conflict.
The formal task consisted of 30 runs of 101 trials each, divided into three sessions
of ten runs each. The participants completed one session each time and all three
sessions within one week. Before each session, the participants performed training
blocks of 20 trials repeatedly until the accuracy reached 90% in the most recent block.
The trial sequences of the formal task were pseudo-randomly generated to ensure that
each of the task conditions and their transitions occurred with equal number of trials.

Experiment 2

Experimental Design. The apparatus, stimuli and procedure were identical to
Experiment 1 except for the changes below. The stimuli were back projected onto a
screen (with viewing angle being ~3.9° between the arrow and the center of the
screen) behind the subject and viewed via a surface mirror mounted onto the head
coil. Due to the time constraints of fMRI scanning, the trial numbers decreased to a
total of 340, divided into two runs with 170 trials each. To obtain a better
hemodynamic model fitting, we generated two pseudo-random sequences optimized
with a genetic algorithm[64] conducted by the NeuroDesign package[65] (see Note S3 for
more detail). In addition, we added 6 seconds of fixation before each run to allow the
stabilization of the hemodynamic signal, and 20 seconds after each run to allow the
signal to drop to the baseline.
Before scanning, participants performed two practice sessions. The first one
contained 10 trials of center-displayed arrow and the second one contained 32 trials
using the same design as the main task. They repeated both sessions until their
performance accuracy for each session reached 90%, after which the scanning began.

 fMRI Image acquisition and preprocessing

Functional imaging was performed on a 3T GE scanner (Discovery MR750) using
echo-planar imaging (EPI) sensitive to BOLD contrast [in-plane resolution of 3.5 $\times$
3.5 mm^2, 64 $\times$ 64 matrix, 37 slices with a thickness of 3.5 mm and no interslice skip,
repetition time (TR) of 2000 ms, echo-time (TE) of 30 ms, and a flip angle of 90°]. In
addition, a sagittal T1-weighted anatomical image was acquired as a structural
reference scan, with a total of 256 slices at a thickness of 1.0 mm with no gap and an
in-plane resolution of 1.0 $\times$ 1.0 mm^2.
Before preprocessing, the first three volumes of the functional images were
removed due to the instability of the signal at the beginning of the scan. The
anatomical and functional data were preprocessed with the fMRIprep 20.2.0[66]
(RRID:SCR_016216), which is based on Nipype 1.5.1[67] (RRID:SCR_002502).
Specifically, BOLD runs were slice-time corrected using 3dTshift from AFNI
20160207[68] (RRID:SCR_005927). The BOLD time-series were resampled to the
MNI152NLin2009cAsym space without smoothing. For a more detailed description
of preprocessing, see Note S4. After preprocessing, we resampled the functional data
to a spatial resolution of 3 $\times$ 3 $\times$ 3 mm^3. All analyses were conducted in volumetric
space, and surface maps are produced with Connectome Workbench
(https://www.humanconnectome.org/software/connectome-workbench) for display
purpose only.

Quantification and Statistical Analysis

Behavioral analysis

Experiment 1. RT and ER were the two dependent variables analyzed. As for RTs,
we excluded the first trial of each block (0.9%, for CSE analysis only), error trials
(3.8%), trials with RTs beyond three *SD*s or shorter than 200 ms (1.3%) and post-
error trials (3.4%). For the ER analysis, the first trial of each block and trials after an
error were excluded. To exclude the possible influence of response repetition, we
centered the RT and ER data within the response repetition and response alternation
conditions separately by replacing condition-specific mean with the global mean for
each subject.
To examine the modulation of conflict similarity on the CSE, we organized trials
based on a 5 (previous trial conflict type) $\times$ 5 (current trial conflict type) $\times$ 2 (previous
trial congruency) $\times$ 2 (current trial congruency) factorial design. As conflict similarity
is commutive between conflict types, we expected the previous by current trial
conflict type factorial design to be a symmetrical (e.g., a conflict 1-conflict 2
sequence in theory has the same conflict similarity modulation effect as a conflict 2-
conflict 1 sequence), resulting a total of 15 conditions left for the first two factors of
the design (i.e., previous $\times$ current trial conflict type). For each previous $\times$ current
trial conflict type condition, the conflict similarity between the two trials can be
quantified as the cosine of their angular difference. In the current design, there were
five possible angular difference levels (0, 22.5°, 42.5°, 67.5° and 90°, see Fig. 1C).

We further coded the previous by current trial congruency conditions (hereafter
abbreviated as CSE conditions) as CC, CI, IC and II, with the first and second letter
encoding the congruency (C) or incongruency (I) on the previous and current trial,
respectively. As the CSE is operationalized as the interaction between previous and
current trial congruency, it can be rewritten as a contrast of (CI – CC) – (II – IC). In
other words, the load of CSE on CI, CC, II and IC conditions is 1, –1, –1 and 1,
respectively. To estimate the modulation of conflict similarity on the CSE, we built a
regressor by calculating the Kronecker product of the conflict similarity scores of the
15 previous × current trial conflict similarity conditions and the CSE loadings of
previous × current trial congruency conditions. This regressor was regressed against
RT and ER data separately, which were normalized across participants and CSE
conditions. The regression was performed using a linear mixed-effect model, with the
intercept and the slope of the regressor for the modulation of conflict similarity on the
CSE as random effects (across both participants and the four CSE conditions). As a
control analysis, we built a similar two-stage model[28]. In the first stage, the CSE [i.e.,
(CI – CC) – (II – IC)] for each of the previous × current trial conflict similarity
condition was computed. In the second stage, CSE was used as the dependent variable
and was predicted using conflict similarity across the 15 previous × current trial
conflict type conditions. The regression was also performed using a linear mixed
effect model with the intercept and the slope of the regressor for the modulation of
conflict similarity on the CSE as random effects (across participants).
Experiment 2. Behavioral data was analyzed using the same linear mixed effect model
as Experiment 1, with all the CC, CI, IC and II trials as the dependent variable. In
addition, to test if fMRI activity patterns may explain the behavioral representations
differently in congruent and incongruent conditions, we conducted the same analysis
to measure behavioral modulation of conflict similarity on the CSE using congruent
(CC and IC) and incongruent (CI and II) trials separately.

Estimation of fMRI activity with univariate general linear model (GLM)

To estimate voxel-wise fMRI activity for each of the experimental conditions, the
preprocessed fMRI data of each run were analyzed with the GLM. We conducted
three GLMs for different purposes. GLM1 aimed to validate the design of our study
by replicating the engagement of frontoparietal activities in conflict processing
documented in previous studies[7,19], and to explore the cognitive space related regions
that were parametrically modulated by the conflict type. Preprocessed functional
images were smoothed using a 6-mm FWHM Gaussian kernel. We included
incongruent and congruent conditions as main regressors and appended a parametric
modulator for each condition. The modulation parameters for Conf 1, Conf 2, Conf 3,
Conf 4, and Conf 5 trials were −2, −1, 0, 1 and 2, respectively. In addition, we also
added event-related nuisance regressors, including error/missed trials, outlier trials
(slower than three SDs of the mean or faster than 200 ms) and trials within two TRs
of significant head motion (i.e., outlier TRs, defined as standard DVARS > 1.5 or FD
> 0.9 mm from previous TR)[41]. On average there were 1.2 outlier TRs for each run.
These regressors were convolved with a canonical hemodynamic response function

(HRF) in SPM 12 (http://www.fil.ion.ucl.ac.uk/spm). We further added volume-level
nuisance regressors, including the six head motion parameters, the global signal, the
white matter signal, the cerebrospinal fluid signal, and outlier TRs. Low-frequency
signal drifts were filtered using a cutoff period of 128 s. The two runs were regarded
as different sessions and incorporated into a single GLM to get more power. This
yielded two beta maps (i.e., a main effect map and a parametric modulation map) for
the incongruent and congruent conditions, respectively and for each subject. At the
group level, paired t-tests were conducted between incongruent and congruent
conditions, one for the main effect and the other for the parametric modulation effect.
Since the spatial Stroop and Simon conflict change in the opposite direction to each
other, a positive modulation effect would reflect a higher brain activation when there
is more Simon conflict, and a negative modulation effect would reflect a higher brain
activation for more spatial Stroop conflict. To avoid confusion, we converted the
modulation effect of spatial Stroop to positive by using a contrast of $[- (I\_pm -$
$C\_pm)]$ throughout the results presentation. Results were thresholded by 3dclust
function in AFNI [69] with voxel-wise $p < .005$ and cluster-size > 20 voxels, which was
supposed to produce a desirable balance between Type I and II error rates[70]. To
visualize the parametric modulation effects, we conducted a similar GLM (GLM2),
except we used incongruent and congruent conditions from each conflict type as
separate regressors with no parametric modulation. Then we extracted beta
coefficients for each regressor and each participant with regions observed in GLM1 as
regions of interest, and finally got the incongruent−congruent contrasts for each
conflict type at the individual level. We reported the results in Fig. 3, Table S1, and
Fig. S3. Visualization of the uni-voxel results was made by the MRIcron
(https://www.mccauslandcenter.sc.edu/mricro/mricron/).
The GLM3 aimed to prepare for the representational similarity analysis (see
below). There were several differences compared to GLM1. The unsmoothed
functional images after preprocessing were used. This model included 20 event-
related regressors, one for each of the 5 (conflict type) × 2 (congruency condition) × 2
(arrow direction) conditions. The event-related nuisance regressors were similar to
GLM1, but with additional regressors of response repetition and post-error trials to
account for the nuisance inter-trial effects. To fully expand the variance, we
conducted one GLM analysis for each run. After this procedure, a voxel-wise fMRI
activation map was obtained per condition, run and subject.

Representational similarity analysis (RSA)

To measure the neural representation of conflict similarity, we adopted the RSA.
RSAs were conducted on each of the 360 cortical regions of a volumetric version of
the MMP cortical atlas[42]. To de-correlate the factors of conflict type and orientation of
stimulus location, we leveraged the between-subject manipulation of stimulus
locations and conducted RSA in a cross-subject fashion (Fig. S4)[60,71]. The beta
estimates from GLM3 were noise-normalized by dividing the original beta
coefficients by the square root of the covariance matrix of the error terms[72]. For each
cortical region, we calculated the Pearson's correlations between fMRI activity

patterns for each run and each subject, yielding a 1400 (20 conditions × 2 runs × 35
participants) × 1400 RSM. The correlations were calculated in a cross-voxel manner
using the fMRI activation maps obtained from GLM3 described in the previous
section. Similar to the behavioral analyses, we assumed the conflict similarity
between two trials is commutive and hence collapsed the RSM along the diagonal and
converted the lower triangle into a vector, which was then z-transformed and
submitted to a linear mixed effect model as the dependent variable. The linear mixed
effect model also included regressors of conflict similarity and orientation similarity.
Importantly, conflict similarity was based on how Simon and spatial Stroop conflict
are combined and hence was calculated by first rotating all subject's stimulus location
to the top-right and bottom-left quadrants, whereas orientation was calculated using
original stimulus locations. As a result, the regressors representing conflict similarity
and orientation similarity were de-correlated. Similarity between two conditions was
measured as the cosine value of the angular difference. Other regressors included a
target similarity regressor (i.e., whether the arrow directions were identical), a
response similarity regressor (i.e., whether the correct responses were identical); a
spatial Stroop distractor regressor (i.e., vertical distance between two stimulus
locations); a Simon distractor regressor (i.e., horizontal distance between two stimulus
locations). Additionally, we also included three regressors denoting the similarity of
Run (i.e., whether two conditions are within the same run), Subject (i.e., whether two
conditions are within the same subject), and Group (i.e., whether two conditions are
within the same subject group, according to the stimulus-response mapping). We also
added two regressors including ROI-mean fMRI activations for each condition of the
pair to remove the possible uni-voxel influence on the RSM. A last term was the
intercept. The intercept and slopes of the regressors were set as random effects at the
subject level. Individual effects for each regressor were also extracted from the model
for statistical inference and brain-behavioral correlation analyses. In brain-behavioral
analyses, only the RT was used as behavioral measure to be consistent with the fMRI
results, where the error trials were regressed out.
The statistical significance of these beta estimates was determined with one-
sample t-tests (one-tailed). Multiple comparison correction was applied with false
discovery rate (FDR) approach[73] across all cortical regions ($p_{FDR} < 0.05$), together
with a threshold of 0.001 for each region. To test if the representation strengths are
different between congruent and incongruent conditions, we also conducted the RDM
analyses using only congruent and incongruent trials separately. Individual effects
were extracted from each model and tested using a paired t-test. To visualize the
difference, we plotted the effect-related patterns (the predictor multiplied by the slope,
plus the residual) as a function of the similarity levels (Fig. 4D).

Representational connectivity analysis

To explore the possible relevance between the conflict type and the orientation
effects, we conducted representational connectivity[43] between regions showing
evidence encoding conflict similarity and orientation similarity. Similar to the RSA
mentioned above, the z-transformed RSM vector of each region were extracted and

submitted to a mixed linear model, with the RSM of the conflict type region (i.e., the
right 8C) as the dependent variable, and the RSM of one of the orientation regions
(e.g., bilateral V2) as the predictor. Intercept and the slope of the regressor were set as
random effects at the subject level, and individual coefficients of the slope were
extracted for further statistical analysis. The mixed effect model was conducted for
each pair of regions, respectively. Considering there might be strong intrinsic
correlations across the RSMs induced by the nuisance factors, such as the within-
subject similarity, we added two sets of regions as control. First, we selected regions
without showing any effects of interest (i.e., $q_{FDR} > 0.05$ for all the conflict type,
orientation, congruency, target, response, spatial Stroop distractor and Simon
distractor effects). Second, we selected regions of orientation effect meeting the first
but not the second criterion, to account for the potential correlation between regions
of the two partly orthogonal regressors (Fig. S6). Existence of representational
connectivity was defined by a higher connectivity slope than any of the control
regions with paired-t tests.
780

Acknowledgement

782 We thank Eliot Hazeltine for valuable comments on a previous version of this
783 manuscript. The work was supported by the National Natural Science Foundation of
784 China and the German Research Foundation (NSFC 62061136001/DFG TRR-169) to
785 X.L. and China Postdoctoral Science Foundation (2019M650884) to G.Y.
786

Reference

1. Kornblum, S., Hasbroucq, T., and Osman, A. (1990). Dimensional overlap:
cognitive basis for stimulus-response compatibility--a model and taxonomy.
Psychol. Rev. *97*, 253-270. 10.1037/0033-295x.97.2.253.
2. Freitas, A.L., Bahar, M., Yang, S., and Banai, R. (2007). Contextual adjustments
in cognitive control across tasks. Psychol. Sci. *18*, 1040-1043. 10.1111/j.1467-
9280.2007.02022.x.
3. Magen, H., and Cohen, A. (2007). Modularity beyond perception: evidence from
single task interference paradigms. Cogn. Psychol. *55*, 1-36.
10.1016/j.cogpsych.2006.09.003.
4. Yang, G., Nan, W., Zheng, Y., Wu, H., Li, Q., and Liu, X. (2017). Distinct
cognitive control mechanisms as revealed by modality-specific conflict
adaptation effects. J. Exp. Psychol. Hum. Percept. Perform. *43*, 807-818.
10.1037/xhp0000351.
5. Hazeltine, E., Lightman, E., Schwarb, H., and Schumacher, E.H. (2011). The
boundaries of sequential modulations: evidence for set-level control. J. Exp.
Psychol. Hum. Percept. Perform. *37*, 1898-1914. 10.1037/a0024662.
6. Liu, X., Banich, M.T., Jacobson, B.L., and Tanabe, J.L. (2004). Common and
distinct neural substrates of attentional control in an integrated Simon and spatial

Stroop task as assessed by event-related fMRI. NeuroImage *22*, 1097-1106.
10.1016/j.neuroimage.2004.02.033.

7. Jiang, J., and Egner, T. (2014). Using neural pattern classifiers to quantify the
modularity of conflict-control mechanisms in the human brain. Cereb Cortex *24*,
1793-1805. 10.1093/cercor/bht029.

8. Kan, I.P., Teubner-Rhodes, S., Drummey, A.B., Nutile, L., Krupa, L., and
Novick, J.M. (2013). To adapt or not to adapt: the question of domain-general
cognitive control. Cognition *129*, 637-651. 10.1016/j.cognition.2013.09.001.

9. Peterson, B.S., Kane, M.J., Alexander, G.M., Lacadie, C., Skudlarski, P., Leung,
H.C., May, J., and Gore, J.C. (2002). An event-related functional MRI study
comparing interference effects in the Simon and Stroop tasks. Brain Res. Cogn.
Brain Res. *13*, 427-440. 10.1016/s0926-6410(02)00054-x.

10. Wu, T., Spagna, A., Chen, C., Schulz, K.P., Hof, P.R., and Fan, J. (2020).
Supramodal Mechanisms of the Cognitive Control Network in Uncertainty
Processing. Cereb. Cortex *30*, 6336-6349. 10.1093/cercor/bhaa189 %J Cerebral
Cortex.

11. Assem, M., Glasser, M.F., Van Essen, D.C., and Duncan, J. (2020). A Domain-
General Cognitive Core Defined in Multimodally Parcellated Human Cortex.
Cereb. Cortex *30*, 4361-4380. 10.1093/cercor/bhaa023.

12. Cole, M.W., Reynolds, J.R., Power, J.D., Repovs, G., Anticevic, A., and Braver,
T.S. (2013). Multi-task connectivity reveals flexible hubs for adaptive task
control. Nat. Neurosci. *16*, 1348-1355. 10.1038/nn.3470.

13. Musslick, S., and Cohen, J.D. (2021). Rationalizing constraints on the capacity
for cognitive control. Trends Cogn Sci *25*, 757-775. 10.1016/j.tics.2021.06.001.

14. Cosmides, L., and Tooby, J. (1994). Origins of domain specificity: The evolution
of functional organization. In Mapping the mind: Domain specificity in cognition
and culture, L.A. Hirschfeld, and S.A. Gelman, eds. (Cambridge University
Press).

15. Egner, T. (2008). Multiple conflict-driven control mechanisms in the human
brain. Trends Cogn. Sci. *12*, 374-380. 10.1016/j.tics.2008.07.001.

16. Kim, C., Chung, C., and Kim, J. (2012). Conflict adjustment through domain-
specific multiple cognitive control mechanisms. Brain Res. *1444*, 55-64.
10.1016/j.brainres.2012.01.023.

17. Abrahamse, E., Braem, S., Notebaert, W., and Verguts, T. (2016). Grounding
cognitive control in associative learning. Psychol. Bull. *142*, 693-728.
10.1037/bul0000047.

18. Freitas, A.L., and Clark, S.L. (2015). Generality and specificity in cognitive
control: conflict adaptation within and across selective-attention tasks but not
across selective-attention and Simon tasks. Psychol. Res. *79*, 143-162.
10.1007/s00426-014-0540-1.

19. Li, Q., Yang, G., Li, Z., Qi, Y., Cole, M.W., and Liu, X. (2017). Conflict
detection and resolution rely on a combination of common and distinct cognitive
control networks. Neurosci. Biobehav. Rev. *83*, 123-131.
10.1016/j.neubiorev.2017.09.032.

20. Fu, Z., Beam, D., Chung, J.M., Reed, C.M., Mamelak, A.N., Adolphs, R., and
Rutishauser, U. (2022). The geometry of domain-general performance monitoring
in the human medial frontal cortex. Science *376*, eabm9922.
10.1126/science.abm9922.
21. Braem, S., Abrahamse, E.L., Duthoo, W., and Notebaert, W. (2014). What
determines the specificity of conflict adaptation? A review, critical analysis, and
proposed synthesis. Front. Psychol. *5*, 1134. 10.3389/fpsyg.2014.01134.
22. Bellmund, J.L.S., Gardenfors, P., Moser, E.I., and Doeller, C.F. (2018).
Navigating cognition: Spatial codes for human thinking. Science *362*.
10.1126/science.aat6766.
23. Behrens, T.E.J., Muller, T.H., Whittington, J.C.R., Mark, S., Baram, A.B.,
Stachenfeld, K.L., and Kurth-Nelson, Z. (2018). What Is a Cognitive Map?
Organizing Knowledge for Flexible Behavior. Neuron *100*, 490-509.
10.1016/j.neuron.2018.10.002.
24. Schuck, N.W., Cai, M.B., Wilson, R.C., and Niv, Y. (2016). Human Orbitofrontal
Cortex Represents a Cognitive Map of State Space. Neuron *91*, 1402-1412.
10.1016/j.neuron.2016.08.019.
25. Park, S.A., Miller, D.S., Nili, H., Ranganath, C., and Boorman, E.D. (2020). Map
Making: Constructing, Combining, and Inferring on Abstract Cognitive Maps.
Neuron *107*, 1226-1238 e1228. 10.1016/j.neuron.2020.06.030.
26. Badre, D., Bhandari, A., Keglovits, H., and Kikumoto, A. (2021). The
dimensionality of neural representations for control. Curr Opin Behav Sci *38*, 20-
28. 10.1016/j.cobeha.2020.07.002.
27. Grahek, I., Leng, X., Fahey, M.P., Yee, D., and Shenhav, A. (2022). Empirical
and Computational Evidence for Reconfiguration Costs During Within-Task
Adjustments in Cognitive Control. In 44.
28. Yang, G., Xu, H., Li, Z., Nan, W., Wu, H., Li, Q., and Liu, X. (2021). The
congruency sequence effect is modulated by the similarity of conflicts. J. Exp.
Psychol. Learn. Mem. Cogn. *47*, 1705-1719. 10.1037/xlm0001054.
29. Vaidya, A.R., Jones, H.M., Castillo, J., and Badre, D. (2021). Neural
representation of abstract task structure during generalization. Elife *10*, 1-26.
10.7554/eLife.63226.
30. Vaidya, A.R., and Badre, D. (2022). Abstract task representations for inference
and control. Trends Cogn Sci *26*, 484-498. 10.1016/j.tics.2022.03.009.
31. MacDowell, C.J., Tafazoli, S., and Buschman, T.J. (2022). A Goldilocks theory
of cognitive control: Balancing precision and efficiency with low-dimensional
control states. Curr Opin Neurobiol *76*, 102606. 10.1016/j.conb.2022.102606.
32. Freund, M.C., Bugg, J.M., and Braver, T.S. (2021). A Representational Similarity
Analysis of Cognitive Control during Color-Word Stroop. J. Neurosci. *41*, 7388-
7402. 10.1523/JNEUROSCI.2956-20.2021.
33. Lu, C.H., and Proctor, R.W. (1995). The influence of irrelevant location
information on performance: A review of the Simon and spatial Stroop effects.
Psychon Bull Rev *2*, 174-207. 10.3758/BF03210959.

34. Simon, J.R., and Small, A.M., Jr. (1969). Processing auditory information:
interference from an irrelevant cue. J. Appl. Psychol. *53*, 433-435.
10.1037/h0028034.

35. Egner, T. (2007). Congruency sequence effects and cognitive control. Cogn.
Affect. Behav. Neurosci. *7*, 380-390. 10.3758/cabn.7.4.380.

36. Schmidt, J.R., and Weissman, D.H. (2014). Congruency sequence effects without
feature integration or contingency learning confounds. PLoS One *9*, e102337.
10.1371/journal.pone.0102337.

37. Torres-Quesada, M., Funes, M.J., and Lupianez, J. (2013). Dissociating
proportion congruent and conflict adaptation effects in a Simon-Stroop
procedure. Acta Psychol. (Amst.) *142*, 203-210. 10.1016/j.actpsy.2012.11.015.

38. Akcay, C., and Hazeltine, E. (2011). Domain-specific conflict adaptation without
feature repetitions. Psychon Bull Rev *18*, 505-511. 10.3758/s13423-011-0084-y.

39. Egner, T., Delano, M., and Hirsch, J. (2007). Separate conflict-specific cognitive
control mechanisms in the human brain. NeuroImage *35*, 940-948.
10.1016/j.neuroimage.2006.11.061.

40. Botvinick, M.M., Cohen, J.D., and Carter, C.S. (2004). Conflict monitoring and
anterior cingulate cortex: an update. Trends Cogn. Sci. *8*, 539-546.
10.1016/j.tics.2004.10.003.

41. Jiang, J., Wang, S.F., Guo, W., Fernandez, C., and Wagner, A.D. (2020).
Prefrontal reinstatement of contextual task demand is predicted by separable
hippocampal patterns. Nat Commun *11*, 2053. 10.1038/s41467-020-15928-z.

42. Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub,
E., Ugurbil, K., Andersson, J., Beckmann, C.F., Jenkinson, M., et al. (2016). A
multi-modal parcellation of human cerebral cortex. Nature *536*, 171-178.
10.1038/nature18933.

43. Kriegeskorte, N., Mur, M., and Bandettini, P. (2008). Representational similarity
analysis - connecting the branches of systems neuroscience. Front Syst Neurosci
2, 4. 10.3389/neuro.06.004.2008.

44. Li, Q., Nan, W., Wang, K., and Liu, X. (2014). Independent processing of
stimulus-stimulus and stimulus-response conflicts. PLoS One *9*, e89249.
10.1371/journal.pone.0089249.

45. Wang, K., Li, Q., Zheng, Y., Wang, H., and Liu, X. (2014). Temporal and
spectral profiles of stimulus-stimulus and stimulus-response conflict processing.
NeuroImage *89*, 280-288. 10.1016/j.neuroimage.2013.11.045.

46. Liu, X., Park, Y., Gu, X., and Fan, J. (2010). Dimensional overlap accounts for
independence and integration of stimulus-response compatibility effects. Atten.
Percept. Psychophys. *72*, 1710-1720. 10.3758/APP.72.6.1710.

47. Niv, Y. (2019). Learning task-state representations. Nat Neurosci *22*, 1544-1553.
10.1038/s41593-019-0470-8.

48. Constantinescu, A.O., O'Reilly, J.X., and Behrens, T.E.J. (2016). Organizing
conceptual knowledge in humans with a gridlike code. Science *352*, 1464-1468.
10.1126/science.aaf0941.

49. Miller, E.K., and Cohen, J.D. (2001). An integrative theory of prefrontal cortex
function. Annu. Rev. Neurosci. *24*, 167-202. 10.1146/annurev.neuro.24.1.167.

50. Milner, B. (1963). Effects of Different Brain Lesions on Card Sorting - Role of
Frontal Lobes. Arch Neurol-Chicago *9*, 90-&. DOI
10.1001/archneur.1963.00460070100010.

51. Mansouri, F.A., Buckley, M.J., and Tanaka, K. (2007). Mnemonic function of the
dorsolateral prefrontal cortex in conflict-induced behavioral adjustment. Science
318, 987-990. 10.1126/science.1146384.

52. Duncan, J. (2010). The multiple-demand (MD) system of the primate brain:
mental programs for intelligent behaviour. Trends Cogn. Sci. *14*, 172-179.
10.1016/j.tics.2010.01.004.

53. Vanderhasselt, M.A., De Raedt, R., and Baeken, C. (2009). Dorsolateral
prefrontal cortex and Stroop performance: tackling the lateralization. Psychon
Bull Rev *16*, 609-612. 10.3758/PBR.16.3.609.

54. Friehs, M.A., Klaus, J., Singh, T., Frings, C., and Hartwigsen, G. (2020).
Perturbation of the right prefrontal cortex disrupts interference control.
NeuroImage *222*, 117279. 10.1016/j.neuroimage.2020.117279.

55. Reverberi, C., Gorgen, K., and Haynes, J.D. (2012). Compositionality of rule
representations in human prefrontal cortex. Cereb Cortex *22*, 1237-1246.
10.1093/cercor/bhr200.

56. Rmus, M., Ritz, H., Hunter, L.E., Bornstein, A.M., and Shenhav, A. (2022).
Humans can navigate complex graph structures acquired during latent learning.
Cognition *225*, 105103. 10.1016/j.cognition.2022.105103.

57. Flesch, T., Juechems, K., Dumbalska, T., Saxe, A., and Summerfield, C. (2022).
Orthogonal representations for robust context-dependent task performance in
brains and neural networks. Neuron. 10.1016/j.neuron.2022.01.005.

58. Gilbert, C.D., and Li, W. (2013). Top-down influences on visual processing. Nat
Rev Neurosci *14*, 350-363. 10.1038/nrn3476.

59. Duncan, J. (2013). The structure of cognition: attentional episodes in mind and
brain. Neuron *80*, 35-50. 10.1016/j.neuron.2013.09.015.

60. Freund, M.C., Etzel, J.A., and Braver, T.S. (2021). Neural Coding of Cognitive
Control: The Representational Similarity Analysis Approach. Trends Cogn. Sci.
25, 622-638. 10.1016/j.tics.2021.03.011.

61. Hommel, B., Proctor, R.W., and Vu, K.P. (2004). A feature-integration account
of sequential effects in the Simon task. Psychol. Res. *68*, 1-17. 10.1007/s00426-
003-0132-y.

62. Braem, S., Bugg, J.M., Schmidt, J.R., Crump, M.J.C., Weissman, D.H.,
Notebaert, W., and Egner, T. (2019). Measuring Adaptive Control in Conflict
Tasks. Trends Cogn. Sci. *23*, 769-783. 10.1016/j.tics.2019.07.002.

63. MacLeod, C.M. (1991). Half a century of research on the Stroop effect: an
integrative review. Psychol. Bull. *109*, 163-203.

64. Wager, T.D., and Nichols, T.E. (2003). Optimization of experimental design in
fMRI: a general framework using a genetic algorithm. NeuroImage *18*, 293-309.
10.1016/S1053-8119(02)00046-0.

65. Durnez, J., Blair, R., and Poldrack, R.A. (2018). Neurodesign: Optimal
Experimental Designs for Task fMRI. BioRxiv, 119594. 10.1101/119594.
66. Esteban, O., Markiewicz, C.J., Blair, R.W., Moodie, C.A., Isik, A.I., Erramuzpe,
A., Kent, J.D., Goncalves, M., DuPre, E., Snyder, M., et al. (2019). fMRIPrep: a
robust preprocessing pipeline for functional MRI. Nat. Methods 16, 111-116.
10.1038/s41592-018-0235-4.
67. Gorgolewski, K., Burns, C.D., Madison, C., Clark, D., Halchenko, Y.O.,
Waskom, M.L., and Ghosh, S.S. (2011). Nipype: a flexible, lightweight and
extensible neuroimaging data processing framework in python. Front.
Neuroinform. 5, 13. 10.3389/fninf.2011.00013.
68. Jenkinson, M., Bannister, P., Brady, M., and Smith, S. (2002). Improved
Optimization for the Robust and Accurate Linear Registration and Motion
Correction of Brain Images. NeuroImage 17, 825-841. 10.1006/nimg.2002.1132.
69. Cox, R.W., and Hyde, J.S. (1997). Software tools for analysis and visualization of
fMRI data. NMR Biomed. 10, 171-178. 10.1002/(sici)1099-
1492(199706/08)10:4/5<171::Aid-nbm453>3.0.Co;2-l.
70. Lieberman, M.D., and Cunningham, W.A. (2009). Type I and Type II error
concerns in fMRI research: re-balancing the scale. Soc. Cogn. Affect. Neurosci.
4, 423-428. 10.1093/scan/nsp052.
71. van Baar, J.M., Chang, L.J., and Sanfey, A.G. (2019). The computational and
neural substrates of moral strategies in social decision-making. Nat. Commun. 10,
1483. 10.1038/s41467-019-09161-6.
72. Nili, H., Wingfield, C., Walther, A., Su, L., Marslen-Wilson, W., and
Kriegeskorte, N. (2014). A toolbox for representational similarity analysis. PLoS
Comput. Biol. 10, e1003553. 10.1371/journal.pcbi.1003553.
73. Genovese, C.R., Lazar, N.A., and Nichols, T. (2002). Thresholding of statistical
maps in functional neuroimaging using the false discovery rate. NeuroImage 15,
870-878. 10.1006/nimg.2001.1037.

Supplementary Notes

Note S1. Behavioral congruency effects

To test the congruency effects for the five conflict types, we conducted 5 (conflict type) $\times$ 2
(congruency) repeated-measure ANOVAs with RT and ER from both experiments. The results
are displayed in Supplementary Fig. 1.
6
Experiment 1.
For the RT, we observed a significant main effect of Congruency, $F(1, 32) = 407.70$, $p < .001$,
$\eta_p^2 = .93$, a significant main effect of Conflict Type, $F(4, 128) = 6.32$, $p < .001$, $\eta_p^2 = .16$, and an
interaction between Conflict Type and Congruency, $F(4, 128) = 27.86$, $p < .001$, $\eta_p^2 = .47$.
Simple effect analyses showed that participants responded more slowly in incongruent
conditions than in congruent conditions for all conflict types, $p_{FDR}s < .001$. Additionally, the
congruency effect of the Type 2, 3 and 4 were significantly larger than that of the Type 1, and
the congruency effect of the Type 2 and 3 were significantly larger than that of the Type 5, $p_{FDR}s$
$< .05$.
Similar results were found with the ER. We observed a significant main effect of
Congruency, $F(1, 32) = 56.83$, $p < .001$, $\eta_p^2 = .64$, a significant main effect of Conflict Type, $F(4,$
$128) = 6.29$, $p < .001$, $\eta_p^2 = .16$, and an interaction between Conflict Type and Congruency, $F(4,$
$128) = 13.23$, $p < .001$, $\eta_p^2 = .29$. Simple effect analyses showed that participants were more
error-prone in incongruent conditions than in congruent conditions for all conflict types, $p_{FDR}s$
$< .001$. The congruency effect of the Type 2, 3 and 4 were significantly larger than that of the
Type 1, and the congruency effect of the Type 3 and 4 were significantly larger than that of the
Type 5, $p_{FDR}s < .05$.
24
Experiment 2.
For the RT, We observed a significant main effect of Congruency, $F(1, 34) = 149.71$, $p < .001$,
$\eta_p^2 = .81$, a significant main effect of Conflict Type, $F(4, 136) = 10.11$, $p < .001$, $\eta_p^2 = .23$, and an
interaction between Conflict Type and Congruency, $F(4, 136) = 7.63$, $p < .001$, $\eta_p^2 = .18$. Simple
effect analyses showed that participants responded more slowly in incongruent conditions than in
congruent conditions for all conflict types, $p_{FDR}s < .001$. The congruency effect of the Type 4
condition was larger than that of Type 1, and Type 3 and Type 4 were significantly larger than
that of the Type 5, $p_{FDR}s < .05$.
For the ER, we only observed a significant main effect of Congruency, $F(1, 34) = 29.80$, p
$< .001$, $\eta_p^2 = .47$. All the types showed a larger error rate in incongruent than congruent
conditions ($p_{FDR}s < .001$), except that the Type 1 only showed a marginal significance (p_{FDR}
$= .062$).
In sum, we observed strong behavioral congruency effects in both experiments. The findings
indicate that these conflict conditions indeed engaged cognitive control[1].

Note S2. *Modulation of conflict similarity on behavioral CSEs cannot be explained by the*
physical proximity

In our design, the conflict similarity might be confounded by the physical proximity between
stimulus (i.e., the arrow) of two consecutive trials. That is, when arrows of the two trials appear
at the same quadrant, a higher conflict similarity also indicates a higher physical proximity (Fig.
1A). Although the opposite is true if arrows of the two trials appear at different quadrants, it is
possible the behavioral effects can be biased by the within quadrant trials. To examine if the
physical distance has confounded the conflict similarity modulation effect, we conducted an
additional analysis.
We defined the physical angular difference across two trials as the difference of their polar
angles relative to the origin. Therefore, the physical angular difference could vary from 0 to
180°. For each CSE conditions (i.e., CC, CI, IC and II), we grouped the trials based on their
physical angular distances, and then averaged trials with the same previous by current conflict
type transition but different orders (e.g., Conf 2−Conf 3 and Conf 3−Conf 2) within each subject.
The data were submitted to a mixed-effect model with the conflict similarity, physical proximity
(i.e., the opposite of the physical angular difference) as fixed-effect predictors, and subject and
CSE condition as random effects. Results showed significant conflict similarity modulation
effects in both Experiment 1 (RT: $\beta = 0.09 \pm 0.01$, $t(7812) = 13.74$, $p < .001$, $\eta_p^2 = .025$; ER: $\beta =$
0.09 ± 0.01, $t(7812) = 7.66$, $p < .001$, $\eta_p^2 = .018$) and Experiment 2 (RT: $\beta = 0.21 \pm 0.02$, $t(3956)$
$= 9.88$, $p < .001$, $\eta_p^2 = .043$; ER: $\beta = 0.20 \pm 0.03$, $t(4201) = 6.11$, $p < .001$, $\eta_p^2 = .038$). Thus, the
observed modulation of conflict similarity on behavioral CSEs cannot be explained by physical
proximity.

Note S3. The fMRI sequence generation approach

Two sequences of 170 trials each were generated independently with the
NeuroDesign package [2]. Each sequence was initialized as 10 consecutive sub-blocks
of each condition (incongruent and congruent) for each conflict type (Conf 1, Conf 2,
Conf 3, Conf 4 and Conf 5). The contrasts of interest were the main effect of
congruency (i.e., [1 −1 1 −1 1 −1 1 −1 1 −1]) and the parametric effect (i.e., [−2 −2 −1
−1 0 0 1 1 2 2]). The order was optimized after 5000 cycles of crossover, mutation,
immigration, fitness, and natural selection. The final number of trials for different
conflict types varied from 64 to 73.

Note S4. fMRI data preprocessing

Results included in this manuscript come from preprocessing performed using
fMRIPrep 20.2.0 (RRID:SCR_016216)[3], which is based on Nipype 1.5.1
(RRID:SCR_002502)[4].

Anatomical data preprocessing. The T1-weighted (T1w) image was corrected for
intensity non-uniformity (INU) with N4BiasFieldCorrection[5], distributed with ANTs
2.3.3 (RRID:SCR_004757)[6], and used as T1w-reference throughout the workflow.
The T1w-reference was then skull-stripped with a Nipype implementation of the
antsBrainExtraction.sh workflow (from ANTs), using OASIS30ANTs as target
template. Brain tissue segmentation of cerebrospinal fluid (CSF), white-matter (WM)
and gray-matter (GM) was performed on the brain-extracted T1w using fast (FSL
5.0.9, RRID:SCR_002823)[7]. Volume-based spatial normalization to one standard
space (MNI152NLin2009cAsym) was performed through nonlinear registration with
antsRegistration (ANTs 2.3.3), using brain-extracted versions of both T1w reference
and the T1w template. The following template was selected for spatial normalization:
ICBM 152 Nonlinear Asymmetrical template version 2009c [RRID:SCR_008796;
TemplateFlow ID: MNI152NLin2009cAsym][8].

Functional data preprocessing. For each of the 5 BOLD runs found per subject
(across all tasks and sessions), the following preprocessing was performed. First, a
reference volume and its skull-stripped version were generated using a custom
methodology of fMRIPrep. Susceptibility distortion correction (SDC) was omitted.
The BOLD reference was then co-registered to the T1w reference using flirt (FSL
5.0.9)[9] with the boundary-based registration[10] cost-function. Co-registration was
configured with nine degrees of freedom to account for distortions remaining in the
BOLD reference. Head-motion parameters with respect to the BOLD reference
(transformation matrices, and six corresponding rotation and translation parameters)
are estimated before any spatiotemporal filtering using mcflirt (FSL 5.0.9)[11]. BOLD
runs were slice-time corrected using 3dTshift from AFNI 20160207
(RRID:SCR_005927)[11]. The BOLD time-series (including slice-timing correction
when applied) were resampled onto their original, native space by applying the
transforms to correct for head-motion. These resampled BOLD time-series will be
referred to as preprocessed BOLD in original space, or just preprocessed BOLD. The
BOLD time-series were resampled into standard space, generating a preprocessed
BOLD run in MNI152NLin2009cAsym space. First, a reference volume and its skull-
stripped version were generated using a custom methodology of fMRIPrep. Several
confounding time-series were calculated based on the preprocessed BOLD: framewise
displacement (FD), DVARS and three region-wise global signals. FD was computed
using two formulations following Power (absolute sum of relative motions)[11] and
Jenkinson (relative root mean square displacement between affines, Jenkinson et al.[9]).
FD and DVARS are calculated for each functional run, both using their
implementations in Nipype (following the definitions by Power et al.[11]). The three

global signals are extracted within the CSF, the WM, and the whole-brain masks.
Additionally, a set of physiological regressors were extracted to allow for component-
based noise correction (CompCor)[12]. Principal components are estimated after high-
pass filtering the preprocessed BOLD time-series (using a discrete cosine filter with
128s cut-off) for the two CompCor variants: temporal (tCompCor) and anatomical
(aCompCor). tCompCor components are then calculated from the top 2% variable
voxels within the brain mask. For aCompCor, three probabilistic masks (CSF, WM
and combined CSF+WM) are generated in anatomical space. The implementation
differs from that of Behzadi et al. in that instead of eroding the masks by 2 pixels on
BOLD space, the aCompCor masks are subtracted a mask of pixels that likely contain
a volume fraction of GM. This mask is obtained by thresholding the corresponding
partial volume map at 0.05, and it ensures components are not extracted from voxels
containing a minimal fraction of GM. Finally, these masks are resampled into BOLD
space and binarized by thresholding at 0.99 (as in the original implementation).
Components are also calculated separately within the WM and CSF masks. For each
CompCor decomposition, the k components with the largest singular values are
retained, such that the retained components time series are sufficient to explain 50
percent of variance across the nuisance mask (CSF, WM, combined, or temporal).
The remaining components are dropped from consideration. The head-motion
estimates calculated in the correction step were also placed within the corresponding
confounds file. The confound time series derived from head motion estimates and
global signals were expanded with the inclusion of temporal derivatives and quadratic
terms for each[12]. Frames that exceeded a threshold of 0.5 mm FD or 1.5 standardised
DVARS were annotated as motion outliers. All resamplings can be performed with a
single interpolation step by composing all the pertinent transformations (i.e. head-
motion transform matrices, susceptibility distortion correction when available, and co-
registrations to anatomical and output spaces). Gridded (volumetric) resamplings were
performed using antsApplyTransforms (ANTs), configured with Lanczos
interpolation to minimize the smoothing effects of other kernels[13]. Non-gridded
(surface) resamplings were performed using mri_vol2surf (FreeSurfer).
Many internal operations of fMRIPrep use Nilearn 0.6.2 (RRID:SCR_001362)[14],
mostly within the functional processing workflow. For more details of the pipeline,
see the section corresponding to workflows in fMRIPrep's documentation.

Note S5. *The multivariate representations of conflict type and orientation are*
different from the congruency effect.

An explanation to the stronger encoding of conflict type in incongruent than
congruent condition (Fig. 3B/D) in right 8C area may be the encoding of congruency.
To test this possibility, we first tested the univariate congruency effect (incongruent
minus congruent) using the parametric modulating GLM1 that was used to estimate
fMRI activation levels of conflict type × congruency conditions. We observed no
univariate congruency effect in the right 8C region, $t(34) = -0.03$, $p = .513$, one-
tailed. We further tested the possibility that the congruency effect may be manifested
in behavioral relevance. To this end, we extracted the congruency effect (incongruent
minus congruent) on encoding strength of conflict similarity for each subject from the
mixed-effect model based on the cross-subject RSA (see the *Representational*
similarity analysis of Methods in the main text) and correlated it with the behavioral
congruency effect, averaged across the five conflict types (i.e., the main effect
reported in the Note S1). No significant correlation was observed ($r = 0.14$, $p = .380$,
one-tailed). Taken together, these results suggested that the neural encoding strength
of conflict type does not reflect the level of cognitive control engagement, but the
dynamic adjustment of cognitive control instead.
Similarly, we tested whether those regions with stronger encoding of orientation
in incongruent than congruent condition (i.e., bilateral V2, left FEF, left IP2, right V1,
right H and right PF) reflects the congruency effect. We observed no uni-voxel
congruency effect in any of these regions, all $p_{FDR} > .998$, one-tailed. In addition, the
orientation effect was not correlated to the behavioral congruency in any of the
regions, all $p_{FDR} > .608$, one-tailed. Together with our finding that there was no
correlation between the strength of orientation encoding and the conflict similarity
modulation on behavioral CSEs in any of these regions (see the *Multivariate patterns*
of visual and oculomotor areas encode stimulus orientation of Results in the main
text), these results indicate that the encoding of orientation effect did not reflect the
encoding of congruency or conflict type. Instead, we speculate that the encoding of
orientations provides perceptual information to determine the conflict type.

Note S6. The lateralization of conflict type representation

We observed the right 8C but not the left 8C represented the conflict type similarity.
A further test is to show if there is a lateralization. We tested several regions of the
left dlPFC, including the i6-8, 8Av, 8C, p9-46v, 46, 9-46d, a9-46v[15]. We found that
none of these regions show the representation of conflict type, all $p_{FDR} > .99$. These
results indicate that the conflict type is specifically represented in the right dlPFC.

Supplementary Figures

Fig. S1. The congruency effects of Experiment 1 (A and B) and Experiment 2 (C and
D). Error bars denote the standard errors of mean. Conf 1 to 5 denotes the five conflict
types. Small insets on top of panel A denote an example of stimuli positions for each
conflict type. RT = reaction time; ER = error rate.

Fig. S2. The conflict similarity modulation on performance of Experiment 1 (A, B, D
and E) and Experiment 2 (C and F), respectively. A and D are scatter plots of CSE
[i.e., (CI−CC) − (II−IC)] for RT and ER as a function of the cosine similarity,
respectively. In B, C, E and F, the cosine similarity and RT / ER are normalized
across conflict similarity levels within each of the four CSE conditions (i.e., CC, II,
CI and IC). Conflict similarity for CC and II conditions are reversed (multiplied by
−1), such that for all the four CSE conditions, higher conflict similarity is expected to
be associated with worse performance (see *Behavioral analysis* in Methods). Each dot
represents a subject. The thin colored lines in B, C, E and F are the fitted lines for
each of the four CSE conditions, and the thick black lines are the fitted lines
collapsing across all CSE conditions. For panel C and F some similarity levels are
missing because of the limited trial numbers in the experimental design in Experiment
2. CSE = congruency sequence effect; RT = reaction time; ER = error rate; CI =
congruent (trial n−1)-incongruent (trial n); IC = incongruent-congruent; CC =
congruent-congruent; II = incongruent-incongruent.

204
Fig. S3. Neural congruency effect (I−C) by GLM2 [see the *Estimation of fMRI*
activity with univariate general linear model (GLM) of Methods in the main text],
plotted as a function of conflict type in different cortical ROIs. The ROIs were
selected because they show a statistically significant congruency effects or parametric
modulation effects when analyzed using the univariate GLM1.The pre-SMA and
ACC showed overall congruency effects regardless of the conflict type (upper panel);
the right IPS and right dmPFC were positively modulated by the conflict type and the
left MFG was negatively modulated by the conflict type (lower panel). Conf 1 to 5
denotes the five conflict types, from the spatial Stroop to the Simon. Pre-SMA = pre-
supplementary motor area; ACC = anterior cingulate cortex; IPS = inferior parietal
sulcus; dmPFC = dorsomedial prefrontal cortex; MFG = middle frontal gyrus.

$$RSM_{brain} = \beta_0 + \beta_1 RSM_{ConflictType} + \beta_2 RSM_{Orientation} + \beta_3 RSM_{Congruency} + \beta_4 RSM_{Target} + \beta_5 RSM_{Response} +$$
$$\beta_6 RSM_{StroopDistractor} + \beta_7 RSM_{SimonDistractor} + \beta_8 RSM_{Run} + \beta_9 RSM_{Subject} + \beta_{10} RSM_{Group} +$$
$$\beta_{11} RSM_{UnivoxelRow} + \beta_{12} RSM_{UnivoxelColumn} + \varepsilon$$

Fig. S4. The cross-subject RSA model and the rationale. The RSM is calculated as the
Pearson's correlation between each pair of conditions and the 35 subjects. For 17
subjects, the stimuli were displayed on the top-left and bottom-right quadrants, and
they were asked to respond with left hand to the upward arrow and right hand to the
downward arrow. For the other 18 subjects, the stimuli were displayed on the top-
right and bottom-left quadrants, and they were asked to respond with left hand to the
downward arrow and right hand to the upward arrow. Within each subject, the
conflict type and orientation regressors were perfectly covaried. For instance, the
same conflict type will always be on the same orientation. To de-correlate conflict
type and orientation effects, we conducted the RSA across subjects from different
groups. For example, the dashed ellipses highlight the conditions that are orthogonal
to each other on the orientation representation, response, and Simon distractor, when
their conflict type, target and spatial Stroop distractor are the same. The dashed boxes
show the possible target locations for different conditions. RSM = representational
similarity matrix.

Fig. S5. The cortical regions showing different effects in the main RSA. (A) The target effect reflects the above chance encoding of upward and downward arrow directions, and is most strongly encoded in the visual, memory and semantic regions, possibly because producing a goal-direct response to the stimulus require processing in all these regions. (B) the response effect reflects the above chance encoding of left and right responses, and is most strongly encoded in motor regions. (C) the spatial Stroop distractor effect reflects the above chance encoding of vertical location of the stimulus, and is most strongly encoded in left visual regions. (D) the Simon distractor effect reflects the above chance encoding of horizontal locations of the stimulus, and is most strongly encoded at the right visual regions, among others. All p-values are FDR-corrected ($p_{FDR} < 0.05$ and raw $p < 0.001$) across the 360 cortical ROIs. Brighter colors denote stronger effects as indicated by the opposite of log-transformed p values.

Fig. S6. The representational connectivity between the right 8C area and the cortical regions showing significant encoding of orientation. The black bars represent regions showing both the overall orientation effect and higher encoding of orientation in incongruent than congruent conditions; the grey bars are regions showing only the overall orientation effect but not higher encoding of orientation in incongruent than congruent conditions; and the white bars are regions not showing any of the effects of interest (i.e., $q_{FDR} > 0.05$ for all the conflict type, orientation, congruency, target, response, spatial Stroop distractor and Simon distractor effects). The grey and white bars show controlled regions. Error bars are the standard error of the mean. The dashed line indicates the 95% confidence interval of the highest connectivity of controlled regions (i.e., left V3). l = left, r = right.

258 **Supplementary Tables**

259 *Table S1.* Brain activations for the uni-voxel parametric analysis in GLM1 (voxel-
260 wise one-tailed $p < .005$, cluster > 20)

Region	L/R	MNI coordinate (mm)			Volume (No. of voxels)	MaxZ	BA
		x	y	z			
incongruent > congruent							
Pre-supplementary motor area	R	12	12	73	60	4.07	6
Anterior cingulate cortex	L	−3	26	34	24	3.17	24
Positive parametric modulator (linear Simon effect)							
Inferior parietal sulcus	R	52	−64	33	68	3.06	39
Dorsomedial prefrontal cortex	R	15	57	42	43	2.96	9
Negative parametric modulator (linear spatial Stroop effect)							
Middle frontal gyrus	L	−27	45	25	29	3. 06	46

262 Notes. L = left; R = right; BA = Brodmann area.

Supplementary References

1. Freund MC, Etzel JA, Braver TS. Neural Coding of Cognitive Control: The Representational Similarity Analysis Approach. *Trends Cogn Sci* **25**, 622-638 (2021).

2. Durnez J, Blair R, Poldrack RA. Neurodesign: Optimal Experimental Designs for Task fMRI. *BioRxiv*, 119594 (2018).

3. Esteban O*, et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat Methods* **16**, 111-116 (2019).

4. Gorgolewski K*, et al.* Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in python. *Front Neuroinform* **5**, 13 (2011).

5. Tustison NJ*, et al.* N4ITK: improved N3 bias correction. *IEEE Trans Med Imaging* **29**, 1310-1320 (2010).

6. Avants BB, Epstein CL, Grossman M, Gee JC. Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Med Image Anal* **12**, 26-41 (2008).

7. Zhang Y, Brady M, Smith S. Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Trans Med Imaging* **20**, 45-57 (2001).

8. Fonov VS, Evans AC, McKinstry RC, Almli CR, Collins DL. Unbiased nonlinear average age-appropriate brain templates from birth to adulthood. *NeuroImage* **47**, (2009).

9. Jenkinson M, Smith S. A global optimisation method for robust affine registration of brain images. *Med Image Anal* **5**, 143-156 (2001).

10. Greve DN, Fischl B. Accurate and robust brain image alignment using boundary-based registration. *NeuroImage* **48**, 63-72 (2009).

11. Jenkinson M, Bannister P, Brady M, Smith S. Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage* **17**, 825-841 (2002).

12. Behzadi Y, Restom K, Liau J, Liu TT. A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage* **37**, 90-101 (2007).

13. Lanczos C. Evaluation of Noisy Data. *Journal of the Society for Industrial and Applied Mathematics Series B Numerical Analysis* **1**, 76-85 (1964).

14. Abraham A*, et al.* Machine learning for neuroimaging with scikit-learn. *Front Neuroinform* **8**, 14 (2014).

15. Freund MC, Bugg JM, Braver TS. A Representational Similarity Analysis of Cognitive Control during Color-Word Stroop. *J Neurosci* **41**, 7388-7402 (2021).