**Utilizing high resolution ribosome profiling for the global investigation of gene**

**expression in *Chlamydomonas reinhardtii***

Vincent Leon Gotsmann[1], Michael Kien Yin Ting[2], Nadin Haase[3], Sophia Rudorf[3], Reimo

Zoschke[2], and Felix Willmund[1]*

Author's institution(s)/affiliation(s):

[1] Molecular Genetics of Eukaryotes, RPTU Kaiserslautern-Landau, Paul-Ehrlich-Str. 23,

67663 Kaiserslautern, Germany.

[2] Max Planck Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam-

Golm, Germany.

[3] Institute of Cell Biology and Biophysics, Leibniz University Hanover, Herrenhäuser-Str.

2, 30419 Hanover, Germany.

15
16

**\*Corresponding author:** Felix Willmund (willmund@bio.uni-kl.de)

**Keywords:** ribosome profiling, next generation sequencing, ribosomes, protein synthesis,

translation, transcript accumulation

**Short title:** Cell-wide ribosome profiling in *Chlamydomonas*

**Abstract**

Ribosome profiling (Ribo-seq) is a powerful method for the deep analysis of translation

mechanisms and regulatory circuits during gene expression in plant cells. Here, we

established an optimized and high resolution Ribo-seq protocol for the unicellular model

alga *Chlamydomonas reinhardtii* (Chlamydomonas). Comparing different nuclease

treatments for the extraction and sequencing of ribosome-protected fragments (RPFs)

and parallel RNA-seq, provided deep insight into translational dynamics and post-

transcriptional control of gene expression, thoroughly covering more than 10,000 different

transcripts. Our high quality Ribo-seq protocol captures the 3-nucleotide movement of

elongating ribosomes along nuclear and chloroplast transcripts. Detailed analysis of the

ribosomal offsets on transcripts uncovers presumable transition states during

translocation of elongating ribosomes within the 5´- and 3´- sections of transcripts and

features of eukaryotic translation termination. Furthermore, drastic differences between

cytosolic and chloroplast offset distributions underline the different nature of the

chloroplast translation mechanism. Analyses of RPF distribution along specific transcripts

revealed characteristic patterns of translation elongation exemplified for the major light

harvesting complex proteins, LHCs. Moreover, our Ribo-seq data can be utilized to survey

coding sequence annotations and the expression preference of alternatively spliced

transcripts in Chlamydomonas. These features will be easily accessible for the research

community since we attached our ribosome profiling data to the most recent

Chlamydomonas reference genome.

**Introduction**

Translation is accomplished via ribosomes, highly conserved macromolecular

ribonucleoprotein machines that decode genetic information into linear polypeptide chains

resulting in an almost unlimited diversity of proteins. Protein synthesis is the origin for the

quantitative and qualitative determination of a proteome´s diversity, which ultimately

shapes the biochemical character of a cell. It is now clear that translation output does not

strictly follow the changes of transcript abundance, but instead underlies strict regulatory

control (Pechmann et al., 2013; Payne, 2015; Slobodin and Dikstein, 2020). The

importance of this regulation is illustrated by the fact that bacteria invest about 50% of

their available energy for protein biogenesis and that more than 10% of all yeast proteins

may be somehow involved in this process (Russell and Cook, 1995; Costanzo et al.,

2000). Translational regulation is observed throughout an organism's life span and

includes central regulatory circuits that control the cell cycle, tissue differentiation and

development, acclimation to environmental changes and the integration of external

signals (reviewed in Pechmann et al., 2013; Zoschke and Bock, 2018; Teixeira and

Lehmann, 2019; Wang and Amoyel, 2022). Thus, understanding the regulatory principles

and mechanisms of translation is a central task of modern biology. With the tremendous

advances in next generation sequencing technologies, it is now possible to systematically

assay the status of cellular translation by a method called ribosome profiling (Ribo-seq)

(Ingolia et al., 2009). The technology is based on the observation that ribosomes occupy

precise sections of a translated mRNA (Wolin and Walter, 1988). These mRNA sections

(termed ribosome footprints or Ribosome Protected Fragments, RPFs) are extracted by

nucleolytic removal of un-occupied mRNA in cell lysates or isolated ribosomes. RPFs are

then identified through next generation sequencing and mapping to a reference

transcriptome or genome. Thus, Ribo-seq benefits from the same dynamic range as RNA-

seq, and provides a rich and precise positional information of translating ribosomes from

the physiological state when cells were harvested (Ingolia et al., 2009; Ingolia, 2014; Brar

70 and Weissman, 2015). Parallel sequencing of all expressed transcripts via RNA-seq

71 allows the comparison of RNA accumulation with translation and thereby to determine the

72 fraction of translated transcripts, which is often referred to as "translation efficiency".

73 Assuming that each RPF represents a translating ribosome and thus a synthesized

74 nascent polypeptide, mRNAs with high average RPF coverages can be considered as

75 highly translated transcripts. In addition, local over-representation of specific RPFs within

76 a specific protein-coding sequence (CDS) are commonly interpreted as regions of slow or

77 even halted translation (Ingolia et al., 2009; Ingolia, 2014; Brar and Weissman, 2015).

78 Compared to other techniques used in systems biology, ribosome profiling goes far

79 beyond the targeted assays of radio-labelling of nascent polypeptides or polysome-

80 loading for specific mRNAs, and regularly provides much deeper dataset information than

81 proteomic analyses via mass-spectrometry (Ingolia, 2014; Brar and Weissman, 2015).

82 Yet, ribosome profiling faces the challenges to distinguish active translation events from

83 non-translating (paused or stalled) ribosomes. However, high-quality Ribo-seq datasets,

84 with populations of precise read-length distribution allow the determination of active

85 translation events if RPFs are aligned relative to the ribosomal Peptidyl-site (P-site). This

86 reveals a 3-nucleotide periodic movement of translating ribosome along a translated CDS

87 by over-accumulation of reads containing the first nucleotide of a decoded codon in the

88 P-site. Thus, it is possible to distinguish ribosome pausing/stalling from translation and

89 therefore even enables *ab initio* annotation of uncharacterized protein-coding sequences

90 (Ingolia et al., 2009; Calviello et al., 2016; Hsu et al., 2016).

91   Ribo-seq has been applied to different plant species with variable gene coverage

92 and quality, including *Chlamydomonas reinhardtii* (Chlamydomonas hereafter) (Chung et

93 al., 2015), *Arabidopsis thaliana*, (Liu et al., 2013; Juntawong et al., 2014; Merchante et

al., 2015; Hsu et al., 2016; Lukoszek et al., 2016; Chen et al., 2022), maize (Lei et al.,

2015; Chotewutmontri and Barkan, 2018), and other crop plants (Wu et al., 2019; Yang et

al., 2020; Yang et al., 2021; Chiu et al., 2022). Studying translational regulation in plants

is of specific interest, due to the interplay of three semi-autonomous genomes (nuclear,

chloroplast and mitochondrial), for which many aspects are not fully understood to date.

Chloroplast gene expression seems particularly dependent on co-translational regulation,

which facilitates the fast adjustment of the photosynthesis machinery to environmental

changes and the stoichiometric assembly of chloroplast multi-subunit complexes which

often contain nuclear and chloroplast encoded subunits (Eberhard et al., 2002; Zoschke

and Bock, 2018; Fujita et al., 2019). Recently, ribosome profiling revealed the specific

adjustments made to chloroplast gene expression during acclimation to different light and

temperature regimes (Chotewutmontri and Barkan, 2018; Schuster et al., 2019;

Chotewutmontri and Barkan, 2020; Gao et al., 2022; Trösch et al., 2022). We have

previously applied a high-resolution microarray approach for the fast and cost-efficient

analysis of chloroplast translation in Chlamydomonas, which allowed for the directly

comparison of transcript accumulation and translation output between the alga and land

plants (Trösch et al., 2018; Trösch et al., 2022). We now established Ribo-seq for the

deep analyses of the three Chlamydomonas genomes. Ribo-seq had been described for

Chlamydomonas before (Chung et al., 2015). However, we aimed to optimize the method

to obtain a deeper coverage of genes, with super-resolution tri-nucleotide periodicity for

both nucleus- and chloroplast-encoded genes and to provide a publicly accessible dataset

aligned to the most recent Chlamydomonas genome (Craig et al., 2022).

**Results and Discussion**

*Establishing super resolution ribosome profiling for Chlamydomonas reinhardtii*

Ribosome profiling was performed from logarithmically grown Chlamydomonas cultures

that were kept under mixotrophic conditions and moderate light (see Materials and

Methods). Our previous experience with targeted chloroplast ribosome profiling (Trösch

et al., 2018; Trösch et al., 2022) showed that addition of the elongation inhibitors

chloramphenicol (CAP) and cycloheximide (CHX) (stalling 70S and 80S ribosomes,

respectively) is crucial to prevent ribosome run-off during harvesting (see Materials and

Methods). However, by short incubation of the inhibitors just during harvest, we reduced

the exposure time to the minimum. For RPF generation by nucleolytic digest of RNA that

is not protected by a ribosome, we applied RNase I at room temperature (23 °C) or 4°C,

consistent with most Ribo-seq studies on other organisms (Ingolia et al., 2012). Our

previous chloroplast ribosome profiling approaches (using MNase) revealed that the best

digestion (i.e. polysome to monosome dissociation) was achieved when ribosomes were

purified prior nuclease treatment (Trösch et al., 2018). We thus compared RNase I

digestion on pre-purified ribosomes both at 4°C (condition i.) and 23°C (condition ii.) with

samples containing whole cell lysates (condition iii.) (Figure 1A). RNase I was applied at

a concentration of 1 unit (U) per micro gram (µg) RNA, which is similar to published Ribo-

seq approaches performed in yeast and Arabidopsis (Chartron et al., 2016; Hsu et al.,

2016; Döring et al., 2017). However, the RNase I concentration was far below the >9 U

per µg RNA that was previously applied for Chlamydomonas (Chung et al., 2015). By

using the lowest possible concentrations of RNase I, experimental costs can be reduced.

Moreover, over digestion of ribosomes may cause an over-proportional content of

contaminating rRNA fragments, which then dominate sequencing libraries. In the previous

Chlamydomonas Ribo-seq study by Chung et al. (2015), contaminating rRNA fragments

accounted for >90% of all sequenced reads. Our pilot runs had contaminating rRNA levels

between 80 and 90%, mainly deriving from the cytosolic 5.8S rRNA and the chloroplast

23S rRNA. To further reduce the amount of contaminating rRNA species, we generated

anti-sense biotinylated oligos that allow for the specific depletion of these species. For

digest condition iii., we additionally tested RNase H treatment, which cleaves hybrids of

the anti-sense oligos and the target rRNA contaminants. Other than the standard

procedure, we performed rRNA depletion prior to dual linker ligation and were thus able

to provide a higher input of RPFs in the sample, relative to contaminating rRNAs, which

then allows for lowering the number of PCR amplification steps during final sequencing

library generation. The final PCR amplification step of the sequencing library preparation

is known to be a major source of bias by over-representing certain reads (Aird et al., 2011).

Read duplicates were defined as reads that share the same sequence and the same

unique molecular identifier tags (UMI tags), four random bases located between a RPF

and the sequencing adapters at both of its termini (Fu et al., 2018). Resulting sequencing

libraries were thus corrected by deduplicating the final data sets after the genome

mapping step.

Parallel to Ribo-seq, transcript accumulation was determined by sequencing whole

cell RNA samples on the same platform. Ideally, RNA samples are harvested just prior to

ribosome profiling experiments from the same culture. While typical transcriptomic studies

enrich mRNA pools by polyadenylation-enrichment, we here sequenced fragmented small

RNAs that are derived from the full set of transcripts including plastid transcripts, which

frequently have no or short polyadenylation signal sequences and are thus poorly

enriched by standard protocols (Komine et al., 2000; Gallaher et al., 2018).

Ribosome profiling and transcript samples were sequenced with a depth of 40 M

and 20 M reads, respectively, which on average resulted in 25.7% of reads mapping to

annotated CDSs of the most recent Chlamydomonas reference genome v6.1 (Craig et al.,

2022). On average, 64% mapped to putative rRNA loci and 4.2% mapped to annotated

untranslated regions upstream or downstream of CDSs. Despite the application of

different rRNA depletion methods, the rRNA contamination was very similar in all samples.

Of the sequenced reads that mapped annotated CDS a minor fraction (<1%) matched to

more than 10 positions and were removed from further analyses. Of the reads mapping

to CDSs, 96.24% mapped to transcripts of the nuclear genome, 3.63% mapped to

chloroplast-encoded transcripts, and 0.12% to mitochondrial transcripts.

Transcript accumulation was determined by averaging reads per CDS, relative to

the length of each CDS and the total read numbers of the dataset (Reads Per Kilobase

per Million, RPKM), a parameter that is widely used for transcriptome analyses and

accounts for a higher likelihood of measuring RNA fragments of long versus short

sequences (Conesa et al., 2016). This is of particular importance for the sequencing of

small and fragmented RNAs, as commonly accompanied with Ribo-seq. For Ribo-seq,

averaging RPFs across CDS has been previously used to correct for local variations of

RPF distribution to determine the level of translation output per transcript (Ingolia et al.,

2011; Ingolia, 2014). However, the short RPF molecules represent absolute counts of

ribosome-protected RNA segments in a sample. Since the number of ribosomes bound to

a transcript should majorly be determined by the transcripts initiation and translation rate

instead of their length, RPKM normalization may distort the data, especially by over-

representing read counts for transcripts considerably shorter than 1000 nt and under-

representing very long transcripts (Ingolia et al., 2009). Thus, translation output was

determined by normalizing the RPF counts per CDS relative to total read depth of the

sequencing dataset (Count Per Million, CPM), placing the expression values into a

ribosome-centric context that represents the proportion of ribosomes per transcripts

relative to the total ribosome pool. This is consistent with recent Ribo-seq studies (Hsu et

al., 2016). *Pearson* correlation coefficients between CPM values and transcript length is

near to zero in all samples and proves the absence of any linear monotonic relationship

between these variables. Thus, the effects of transcript length on the read count, if

existent, are much weaker than other influences and therefore negligible (Supplemental

Figure S1A).

Based on CPM and RPKM values, respectively, the three nuclease digestion

conditions (i. to iii.) showed high reproducibility between the Ribo-seq and RNA-seq

experiments, respectively (r = 0.98 and r = 0.99, Figure 1B). This agrees with previous

observations that Ribo-seq tolerates variations in RNase digest (Gerashchenko and

Gladyshev, 2017).

We next aimed to determine the depth of the Ribo-seq approach. Low coverage

Ribo-seq frequently results in ribosome profiles covering only the transcripts with highest

ribosome occupancy (i.e. translation output), whereas moderate or lowly translated CDS

may have partial RPF coverage along the respective sequence (here termed gene body

coverage, Figure 1C, left panel). For Ribo-seq with a good depth, many CDS should show

decent gene body coverage (Figure 1C, right panel). The depth was determined for all

Chlamydomonas transcripts of the v6 genome (32670 transcripts considering the products

of gene copies). For digest condition iii., more than 10,000 transcripts had at least 50% of

their CDSs sequence covered with two or more RPFs, while 6437 transcripts had over

90% of their CDS covered with at least one RPF (Figure 1D). 1508 transcripts are covered

over >90% with an average of more than 10 RPFs/nt, hence representing the group of

transcripts with the most complete gene body coverage combined with a high basal

coverage (see also Figure 4B). Categorizing the transcripts in this way allows for the

assessment of the general quality of a data set, as well as the selection of transcripts of

similar quality for specific analyses without having to rely on expression values alone. The

gene body coverage was highest for digest condition iii. (Figure 1D and Supplemental

Figure S1B and S1C), and comparably good for the coverage of RNA fragments for the

RNA-seq experiment (Figure 1E). Of note, even transcripts with low CPM correlated well

between experimental conditions, demonstrating good reproducibility that is comparable

to the pioneer experiments performed with yeast (Figure 1B) (Ingolia et al., 2009).

The read length distribution of the RPFs mapping to annotated CDSs of the nuclear

genome displayed high comparability to published Ribo-seq approaches with other

organisms. RNase I treatment typically results in RPFs with a length between 28 and 30

nt. Here, digest of purified ribosomes at 4°C resulted in RPFs with a predominant length

of 31 nt. Digest condition ii. caused a broader distribution of RPFs between 28 and 30 nt,

while digest in cell lysate resulted in a sharp peak of RPFs with a length of 30 nt (Figure

2A). Such a sharp peak at 30 nt is typically found in Ribo-seq from mammalian cells when

harvested in the presence of CHX (Wu et al., 2019; Sharma et al., 2021). The RPF size

distribution of the 8 mitochondrial transcripts needs to be considered with caution due to

the low coverage but appears bimodal with peaks at 22 nt and 33 nt under all conditions.

Similar size distributions were observed from previously described mitochondrial Ribo-

seq and could stem from the bulkiness of the membrane associated ribosomes (Rooijers

et al., 2013). For chloroplast transcripts, the RPF size distribution is less defined and may

be the result of diverse translational states, as previously reported for maize and

Arabidopsis (Chotewutmontri and Barkan, 2016; Gawronski et al., 2018; Fujita et al.,

2019). The sharpest peak for chloroplast RPFs is seen under condition ii., albeit the

predominant RPFs of 28 nt is found under all treatment conditions (Figure 2A). While

consistent with the observations in other organisms, our RPF size distribution varies from

the previous Ribo-seq data with Chlamydomonas, which reported a general shift towards

a smaller RPF sizes, maybe as consequence from their harsh RNase I treatment (Chung

et al., 2015).

243           A hallmark of high-resolution Ribo-seq data is the triplet-wise 3-nt periodicity

of RPFs that represents slowdown of ribosome movement as ribosomes decode a

respective codon (Ingolia, 2016). To calculate periodicity, it is necessary to determine the

5'-offset for every RPF length species, which is the number of bases located 5' to the P-

site (the translational state of polypeptide bond formation) of a respective RPF. This is

usually done by considering all RPFs that fully enclose an annotated start codon and

determining the frequency of all occurring 5'-offsets for every RPF length separately. The

5'-offset, most frequently occurring for a specific RPF length, is then declared to be the

true 5'-offset of this species. This strategy is based on the fact that RPF occupancy of

transcripts usually inclines dramatically at the borders between 5'-untranslated regions

and CDS, with the start codon often having an exceptionally high coverage, which is due

to the slow process of translation initiation. The 5'-offsets determined in this way are used

to calculate the exact P-site position and reading frame of every RPF. By this, 3-nt

periodicity can be nicely visualized in meta-plots combining all genes while 3-nt plots of

individual transcripts can be noisy, depending on translation levels. The exact 5'-offset

varies in an organism- and protocol-specific manner and is usually 12 to 13 nt.

Alternatively, the P-site position has also been determined by calculating the 3'-offset of

the ribosomal Aminoacyl-site (A-site), applying the same strategy to RPFs mapping to

annotated stop codons.

With the majority of cytosolic RPFs with one read length (Figure 2A), all three

RNase I digest conditions yielded a clear frame preference of "frame 0" for nucleus-

encoded transcripts, meaning that the base triplet found at the P-site of a RPF is for the

for the majority of RPFs in-frame with the annotated CDS. Again, lysate digest (iii.)

resulted in the best frame preference values (>68% of RPFs in frame 0, 19.4% in frame 1

and 12.5% in frame 2), suggesting that this condition yields Ribo-seq data of the best

quality (Figure 2B). Due to the differences between 80S and 70S ribosomes, 3-nt

periodicity needs to be determined separately for chloroplast translation. For chloroplast

transcripts, the frame preference was less obvious, albeit it was clearest for digest

condition iii. (Figure 2C). Considering the overall quality of sample iii. (the similar degree

of rRNA contamination between all three samples and the high correlation with the other

samples on CPM level), it can be concluded that RNase H mediated rRNA depletion has

no adverse effects on the Ribo-seq sample quality and seems to be similarly effective.

To get a better insight into the RPF offsets and periodicity, different lengths of the

RPFs were plotted against their offset frequencies upon P-site and A-site alignment for

the dataset of condition iii. (Figure 2D). The triplet-wise movements of the 5´ends with an

offset of 12 nt during initiation is clearly visible for cytosolic footprints and most prominent

for the 30 nt RPFs of digestion conditions iii., consistent with the observation in other

eukaryotic organisms (Lauria et al., 2018). Likewise, triplet-wise movements of the 3´-

ends are also detectable and are in frame with the stop codon. Here, a 9 nt long 3´-offset

seems the predominant form prior to termination with the stop codon in the A-site of the

ribosome. However, due to the 12 nt 5'-offset and 3 nt occupied by the P- and A-sites

respectively, a 3'-offset of 12 nt was expected. This discrepancy could be a result of CHX

treatment, which locks ribosomes in a pre-translocation conformation after peptide bond

formation (Wu et al., 2019) so that RPFs on non-stop codons may exceed the natural

termination-caused accumulation on stop-codons. Moreover, a slow termination process

might arrest ribosomes on the codon just upstream of the stop, prolonging ribosome dwell

time at this site. At the same time, ribosomes that already entered termination at the time

of sampling are likely to complete the process and dissociate from the transcript. CHX

does not inhibit termination, which in turn would dramatically lower the number of FPs

mapped to stop codons with their A-site. Interestingly, RPFs with a length of 21 nt are

detectable upon initiation (asterisks, Figure 2D). These fragments have been previously

characterized as footprints of ribosomes in an open ratchet conformation during

translocation, even accumulating as abundant species without CHX treatment in yeast

(Lareau et al., 2014). During elongation, ribosomes undergo massive conformational

changes by rotating the large subunit relative to the small subunit (Frank and Agrawal,

2000; Zhang et al., 2009). The fact that the RPFs produced from the open-ratchet

conformation have the same 5'-offsets as the main RPF species despite being

considerably shorter, suggests that the RNase exclusively removes nucleotides from the

3´-end of RPFs in this conformation (and could indicate that the center of the rotation is

located towards the 5'-direction, seen relative to the RPF). Cycloheximide was shown to

prevent binding of the next aminoacylated tRNA in the A-site during peptide bond

formation (Schneider-Poetsch et al., 2010) and thus stabilizes ribosomes in a specific

stage during the elongation cycle, in which RPFs of 28-30 nt are the prevalent form.

Interestingly, human 80S ribosomes are not exclusively arrested in one conformation

upon CHX treatment (Sharma et al., 2021), comparable to the situation seen here for

Chlamydomonas 80S ribosomes (Figure 2D, upper panel). These 20-22 nt RPFs might

derive from a post-translocation ribosome species (Lareau et al., 2014). The presence of

both species demonstrates active movement of ribosomes along the CDS. The absence

of the "open-ratchet" 21 nt footprints starting from two peptide-bond formations prior to

termination, may support the idea from above that ribosomes near stop codons are

paused, which can prevent the collision between ribosomes (Figure 2D upper panel,

dashed box). Interestingly, the frame preference for frame 0 of the stop-minus-2 codon is

the highest in the whole metagene periodicity plot (79.6% Figure 2B), which could denote

a short stalling at this codon before translocation. However, for the next codons the frame

0 preference drops considerably to 55.8%, especially in favor of frame 1 (36.2%) and then

to 33.1% at the stop codon, clearly indicating that translation dynamics change towards

the termination step. Moreover, these characteristics are accompanied by a striking

continuous decline of absolute numbers of reads mapping to these codons on the meta-

gene level (for digest iii: stop-2: 19890, stop-1: 6786, stop: 522).

The prokaryotic-type chloroplast RPFs 5'-offsets seem to be much more diverse

than their cytosolic counterparts, apparent by their multiple clusters on the heat map

(Figure 2D, bottom panel). These might resemble populations of initiation complexes in

different states or in complex with transcript specific RNA processing factors and *trans*-

acting factors that interact with ribosomes (Westrich et al., 2021). Without clear criteria

how to distinguish and interpret these clusters, 5'-offset definition is problematic, since

multiple 5'-offset frequencies of comparable amplitude exist for many RPF lengths.

However, the chloroplast 3'-offsets seem less scattered and reveals a main RPF species

of 28 nt length with 11 nt 3'-offset. This is in accordance with the RPF length distribution

of all chloroplast RPFs, where the 28 nt species also represents the main species,

although the distribution is much broader than for the cytosolic RPFs. Similar to the

cytosolic offset heat map, a triplet-wise movement of ribosomes can also be observed for

the terminating chloroplast ribosomes in the 3'-offset heatmap for 29 nt RPFs, albeit with

much lower clarity. Altogether, the observations made for chloroplast offsets and RPF

length distribution might indicate a certain heterogeneity of the chloroplast ribosome pool,

which is an interpretation that would be in line with the large number of proteins that

interact with chloroplast ribosome found in our previous study (Westrich et al., 2021) and

the large number of RNA-binding proteins known or suspected to regulate the translation

of specific chloroplasts transcripts (Nickelsen et al., 2014). This is remarkable considering

the very small number of transcripts encoded on the chloroplast genome (72). Together,

this might suggest that translation in the chloroplast of Chlamydomonas is heavily

influenced by transcript-specific features rather than following a universal scheme

(Zoschke and Bock, 2018).

*Differences in translation output between cytosolic and organellar transcripts*

We next compared the translation levels of the organellar versus nuclear transcripts.

Chlamydomonas cells contain on average 83 plastid genomes within the large cup-

shaped chloroplast occupying about 40% of the cellular volume. The mitochondrial

genomes exist in 130 copies, also greatly outnumbering the one haploid nuclear genome

(Gallaher et al., 2018). Despite the rather low contribution of organellar RPFs relative to

the total RPF pool, both chloroplast and mitochondrial transcripts had higher CPM values

when compared to the average CPM values of nuclear genes (Figure 3A). A similar trend

is seen for transcript levels (Figure 3B), suggesting that organellar genes are expressed

at considerably higher levels compared to most nuclear genes (Forsythe et al., 2022).

With the high reproducibility of Ribo-seq and RNA-seq data, we chose to average

the datasets for estimating the levels of transcript accumulation and translation output. A

direct comparison between translation output and transcript accumulation is frequently

referred to as translation efficiency (TE) and normalizes RPF values over transcript

accumulation. Contrasting translation over transcript data of the Chlamydomonas genes

in a scatter plot and highlighting their subcellular localization showed that nuclear

transcripts encoding chloroplast-localized proteins have higher expression values when

compared to proteins of cytosolic or Endoplasmic Reticulum localization (Figure 3C),

again supporting the notion that many chloroplast proteins are expressed at high levels.

Interestingly, most nucleus-encoded transcripts showed similar levels between transcript

accumulation and translation (distributed along the diagonal, Figure 3C). In contrast,

chloroplast- and mitochondria-encoded transcripts showed higher transcript accumulation

levels, when compared to translation, which agrees with the published data of profound

potential translational regulation within the chloroplast (Sun and Zerges, 2015; Trösch et

al., 2018; Zoschke and Bock, 2018). Remarkably, translational regulation seems also

predominant for the few mitochondrial genes.

Column plots of the individual chloroplast transcript and translation values show

that even the weakest expressed known chloroplast transcripts have CPM values ~30,

which is within the median range of nucleus-encoded transcripts (Figure 4A versus 3A).

Not surprisingly, transcripts that encode the core photosystem I subunits PsaA and PsaB,

the photosystem II core subunits PsbA/D1,PsbB/CP47, PsbC/CP43, PsbD/D2, the

Rubisco large subunit, RbcL, and the translation elongation factor TufA show the highest

translation levels, consistent with our previous chloroplast ribosome profiling experiments

and the examination of protein accumulation levels (Schroda et al., 2015; Trösch et al.,

2018). We further detected transcripts of the uncharacterized chloroplast Open Reading

Frames (ORF) *orf202* (positioned within the transposon Wendy I), *orf854* (positioned

within the transposon Wendy II), *orf528*, and *i-creI* (Gallaher et al., 2018), however,

translation output was low or even undetectable (for *i-crel*), suggesting that these

transcripts are barely translated in Chlamydomonas cultures, at least under standard

conditions (Figure 4A). The direct TE ratios of the chloroplast transcripts again show the

imbalance between chloroplast transcript accumulation and translation output, suggesting

that the long-living chloroplast transcripts present a buffering capacity for the fast and

efficient adjustment of translation levels during environmental changes, as for example

seen during light and temperature acclimation and most prominent for PsbA/D1 synthesis

(Chotewutmontri and Barkan, 2018; Schuster et al., 2019; Trösch et al., 2022).

We next determined translation output levels of nucleus-encoded transcripts, only

considering the longest transcript variant if multiple splice forms existed. Of the transcripts

with the highest translation output, 54 protein products have a known or predicted

chloroplast localization and only 34 are present in the nucleus or cytosol, again showing

the high accumulation levels of chloroplast proteins (Supplemental Figure S2A). Amongst

those, transcripts of the light harvesting complex proteins (LHCs) have the highest

translation output, consistent with their high protein accumulation levels in plant cells

(Dall'Osto et al., 2015). Generally, *LHC* translation showed comparable CPM translation

output values, whereas transcript accumulation was rather variable. For the transcripts of

the high-light induced photoprotective LHCSR3.1 and LHCSR3.3 proteins, we could not

determine translation profiles (Supplemental Figure S2B), suggesting that these proteins

are not synthesized under the investigated growth conditions (Bonente et al., 2012).

Besides photosynthesis related transcripts, the list with top-most translation further

contained transcripts encoding proteins such as the Isocitrate lyase (ICL1, CPM 4656,

rank 5), a key enzyme of the glyoxylate cycle (Plancke et al., 2014), the thiazole

biosynthesis enzyme (THI4, CPM 3230, rank 13) (Moulin et al., 2013) or the Methionine

Adenosyl-transferase (METM1, CPM 2856, rank 26), confirming their central role in

Chlamydomonas cells. Interestingly, the $B_{12}$-independent methionine synthase enzyme,

METE1, seems to be higher expressed, when compared to the $B_{12}$-dependent METH1

isoform (CMP 1655 versus 990) (Bunbury et al., 2020). METE1 is less common than

METH1 within the algal clade and makes these species independent form external $B_{12}$

supply (Bertrand et al., 2012).

*Detection of open reading frames by ribosome profiling facilitate Chlamydomonas*

*genome annotation*

Local elongation rates are known to vary considerably during translation along a transcript,

depending on tRNA availability, mRNA secondary structure, the amino acid sequence of

the resulting polypeptide and co-translational folding (Gloge et al., 2014; Kramer et al.,

2019). With the robust gene body coverage over Chlamydomonas transcripts (Figure 1D),

we aimed to learn more from the dynamics of translation elongation, derived from the RPF

distribution over individual coding sequences (ribosome profiles). Ribosome profiles were

highly comparable between the three RNase I treatment conditions i.-iii., as indicated by

median *Pearson* correlations between data sets ii. and iii. of ~0.62 for nucleus-encoded

transcripts and >0.85 for chloroplast-encoded transcripts. Although RNA-seq read

distributions over individual CDS were even more comparable (median of *r*-value 0.87 for

nucleus-encoded and 0.99 for chloroplast-encoded transcripts) among the RNA-seq data

sets, they showed no correlation (median *r*-value close to zero) to the respective ribosome

profiles (Figure 5A and Supplemental Figure S3A). Certainly, methodological-driven

differences between Ribo-seq and RNA-seq cannot be fully excluded, but it can be

assumed that the ribosome profiles image transcript-specific translation dynamics and are

likely not the result of sequencing biases.

The correlation coefficients for specific transcripts comparing two experiments

(both *Pearson* and *Spearman*) stably range between 0.7 and 1 for the majority of ribosome

profiles that have an average coverage >100 reads per nucleotide, suggesting that low

correlation coefficients are the consequence of sequencing depth limitations and thus

limited observability of lowly translated transcripts rather than the true absence of a

characteristic translation dynamic of these transcripts (Supplemental Figure S4).

Importantly, more than 10,000 transcripts have *r*-values >0.7 (Figure 5B), again showing

the high reproducibility between the RNase I digest conditions for transcripts with

moderate and high RPF coverage (Supplemental Figure S4). In fact, the *r*-value

distribution shows that these direct comparisons of read coverage per transcript might be

well suited for selecting good cut-off ranges for a deeper analyses of ribosome profiles. In

many previous studies, simply minimum CPM values (e.g. 1) were taken as cut-off.

Ribosome profiles of the highly translated *LHC* transcripts showed some

remarkable differences between the individual *LHC* species. Efficiently controlling the

expression of *LHC* genes is essential for regulating the antenna size of the photosynthesis

machinery in order to capture photons under low irradiance conditions (increased

expression) or avoid over-excitation and thus photodamage under high irradiance

conditions (reduced expression) (Dall'Osto et al., 2015). Of the transcripts encoding the

21 different LHC members serving Photosystem I or II (LHCA and LHCB, respectively),

20 were found amongst the top 100 highest translated transcripts (not considering splice

variants). Only the transcript encoding LHCB7, a rather newly discovered protein

containing the unusual number of 4 transmembrane domains (Klimmek et al., 2006) had

low CPM values of ~16 (Supplemental Figure S2B). The variations in *LHC* transcript levels

result in considerably different TE values (Supplemental Figure S2B), suggesting that

translational regulation might be achieved by different strategies for expression of the *LHC*

species. Indeed, the ribosome profiles are considerably different between transcripts

encoding the different LHCs. For example, *LHCBM6* showed an exceptionally strong peak

of RPFs covering the first 7 codons, which exceeds the average RPF density over the

remaining CDS by more than 4-fold (Figure 5C). While "initiation peaks" are commonly

accumulating over the first few codons of a CDS, their amplitudes are usually within the

dynamic range of the remaining CDS. Such high peaks, in contrast, could point to a

regulative mechanism stalling ribosomes during or shortly after initiation. Indeed,

*LHCBM6* was shown to be translationally repressed under moderate light conditions by a

co-translational regulator termed NAB1 (Nucleic Acid Binding protein). NAB1 itself is

tuned by a regulatory circuit sensing carbon dioxide supply levels in Chlamydomonas, for

precisely adjusting the antenna composition of photosystem II (Mussgnug et al., 2005;

Blifernez-Klassen et al., 2021). Similarly, *LHCBM4* is recognized by NAB1, also

presenting a strong RPF peak over the initiation codon, while LHCBM1 (not a target of

NAB1) has no over-proportional initiation peak (Figure 5C), despite sharing a 75% amino

acid sequence identity with LHCBM6. Translational regulation during *LHC* expression is

not unique to Chlamydomonas, as *Lhcb4.2*, *Lhcb4.3* and *Lhcb6* are also subject to post-

transcriptional regulation in Arabidopsis (Floris et al., 2013). We searched all transcripts

that had a gene body coverage of at least 70% (8757 nucleus-encoded, 66 chloroplast-

encoded, 6 mitochondria-encoded) for the occurrence of comparable initiation peaks

(average coverage of the first seven codons at least four times higher than the average

coverage of the remaining CDS). Around 38% of selected nucleus-encoded transcripts

had comparable RPF accumulation peaks near the initiation codon, possibly hinting to

more co-translational regulation in the cytosol than previously anticipated. In contrast, only

10% (6 transcripts) of the chloroplast-encoded genes showed such peaks (Figure 5D).

However, it cannot be excluded that these peaks present artifacts of CHX treatment of the

Chlamydomonas cultures during harvest. Upon establishment of the Ribo-seq protocol,

several reports demonstrated that pre-incubation of yeast cultures with CHX may create

a bias by inducing increased RPF accumulation at initiation codons, which could be the

consequence of stalled elongation but unaffected initiation through CHX treatment (Ingolia

et al., 2009; Gerashchenko and Gladyshev, 2014; Sharma et al., 2021). However, a recent

direct comparison of CHX-treated and untreated conditions demonstrated that CHX

causes no initiation-peak bias in human cells, and that even the strong accumulation in

yeast might derive from slightly altered protocols between laboratories (Sharma et al.,

2021). For Chlamydomonas, CHX biases should result in general initiation peaks, but

these peaks are only found in a fraction of transcripts, independent of their translation

activity (e.g. *LHCs*). Thus, varying accumulation of RPFs around the initiation codon may

be an interesting indicator for altered initiation regulation between transcripts in the

cytosol.

Lastly, we aimed to exploit our Ribo-seq data for surveying correct CDS

annotations and transcript splice variants within the recently released Chlamydomonas

genome, version 6.1 (Craig et al., 2022). In previous genome versions, several genes

were falsely annotated, ignoring upstream start codons that were possibly the genuine

initiation site of translation (Cross, 2015; Craig et al., 2022). In the new version, the first

in-frame ATG codon was generally considered as the initiation codon, in accordance with

the scanning model of initiating ribosomes (Hinnebusch, 2011). Systematic comparison

of the reference start site with the onset of RPF coverage, confirmed that start codon

assignment has been clearly improved between genome version 5.6 and 6.1. However,

we detected gene models in which the first ATG appears not to be the genuine initiation

site. For example, transcripts encoding LHCA4 (Cre10.g452050) do not initiate from the

first ATG (Figure 6A) and were correctly annotated in the previous genome versions.

Another example is PRPL3 (Cre10.g417700) (Figure 6B). In addition, we determined

some gene models in which RPF accumulation did not correlate with the cognate ATG

start codon, possibly hinting to a translation start from non-cognate start sites as

exemplified by the *PDI2* CDS (Cre01.g033550, Figure 6C). In this case, the ribosome

profile starts with a prominent peak upstream of the annotated start codon located within

the annotated 5'-UTR while the annotated start codon is poorly covered compared with

the remaining CDS. Here, initiation from the non-cognate TTG could be an explanation

(Cao and Slavoff, 2020), which is located in-frame with the annotated coding sequence at

two positions within the area covered by the peak. Ribo-seq was already used before to

mine for non-cognate start sites in other organisms (Brar and Weissman, 2015).

Alternatively, these RPFs could point to the presence of upstream Open Reading Frames

for *PDI2*. *PDI2* represents a good example for detecting alternative splicing. Transcript 1

of *PDI2*, possesses a RPF gap between exon 3 and 4 (379 nt downstream of the

transcription start site), which is spliced out in transcript versions 2-4 (Figure 6C). In

addition, a sudden drop of RPFs is seen at the end of exon 3 in transcript 2 and 4. Based

on distribution of RPFs and the transition of ribosome profiles, transcript 5 has the most

consistent profile and might hence be the prevalently expressed variant in

Chlamydomonas cells, grown under standard laboratory conditions. In future, intelligent

modelling and peak deconvolution algorithms could be potentially applied to Ribo-seq

data to estimate the stoichiometric ratio of different splice variants' translation output.

*Concluding remarks*

Taken together, ribosome profiling is a fascinating tool for advancing our understanding

of post-transcriptional regulation of gene expression. Our Ribo-seq approach has the

resolution and depth to deeply study translation of the nuclear and chloroplast transcripts

in Chlamydomonas cells. Furthermore, Chlamydomonas is a well-suited organism for

understanding system level changes of gene expression throughout the cell cycle or

following environmental changes. Ribosome profiles can be of specific help, for a first

understanding of the many transcripts, which were not experimentally investigated before.

To make our data easily accessible in the research community, we have now linked our

Ribo-seq data to the recent Chlamydomonas genome CC-4532 v6.1 (https://phytozome-

next.jgi.doe.gov), thereby allowing individuals to conveniently browse the ribosome

profiles of their genes of interest. With the new genome release, the number of

alternatively spliced transcripts have been significantly expanded. Visualization of

ribosome profiles may help to uncover the dominant transcript variants, which are

translated under standard laboratory conditions. Certainly, additional experimental

approaches are required to understand unexpected features such as translation initiation

from non-cognate start sites, RPFs along upstream open reading frames, mechanism of

translational regulation, and spatial organization of protein synthesis.

**Materials and Methods**

*Cell growth and harvest*

*Chlamydomonas reinhardtii* CC-1690 cells were grown mixotrophically in Tris-Acetate

Phosphate (TAP) medium (Kropat et al., 2011) to mid-log phase (4 to $5 \times 10^6$ cells per mL)

under constant light of 80 mmol $m^{-2}$ $s^{-1}$ (MASTER LEDtube HF 1,200 mm UO 16W830 T8

and 16W840 T8, Philips) and 25°C on a rotary shaker. Immediately before harvest, 100

µg/mL chloramphenicol (CAP) and cycloheximide (CHX) was added and cells were rapidly

chilled by pouring over -80°C cold silicon ice cubes until 4°C culture temperature was

reached. Subsequently, cells were pelleted by 2 min centrifugation at 4,000 g and 4°C,

washed in ice-cold TKM+ buffer (50 mM acetate buffered Tris pH 8.0, 150 mM KCl, 10

mM $MgCl_2$, 100 µg/mL CAP, 100 µg/mL CHX) and pelleted again using the same

centrifugation conditions. Afterwards, the pellet was resuspended in freezing buffer (TKM+

supplemented with 100 mM phenyl methyl sulfonyl fluoride (PMSF) and 16% of 25x Roche

protease inhibitor cocktail) by slow pipetting in 1/2000 of the initial culture volume and

flash frozen by dripping the cell suspension into liquid nitrogen. Frozen cells were stored

at -80°C until further use.

*Ribosome profiling and RNA sampling*

Frozen cells were ground using a bead mill with nitrogen-cooled steel containers and

beads for two times 2 min at 27 Hz frequency and cooling in liquid nitrogen between

rounds. The cell powder was mixed with equal volumes of 2x concentrated lysis buffer (2x

concentrated TKM+ buffer supplemented with 2 mM dithiotreitol (DTT), 2% Triton X-100

and 20% sucrose) and incubated for 5 min at 4°C before pelleting cell debris at 8,000 g,

4°C for 10 min. For nucleolytic digest conditions i. and ii., lysate was layered on a 2.5 mL

64% sucrose cushion prepared in TKM+ buffer supplemented with 1 mM DTT and

ultracentrifuged for 3 h at 60,000 rpm and 4°C in a Ti70 rotor. The resulting polysome

pellet was briefly rinsed with 500 µL ice-cold diethyl pyro carbonate (DEPC)-treated water

and resuspended overnight in 100 µL ice-cold ribosome buffer (2x concentrated TKM+

buffer diluted with 25x concentrated Roche protease inhibitor cocktail to a total amount of

16%, 1 mM PMSF and 0.1% Triton X-100). Remaining insoluble debris were removed by

centrifuged for 1 min at 1,500 g before addition of 1 U of RNase I (Ambion) per µg RNA

and 0.134 U TURBO DNase (Invitrogen). Nucleolytic digest was performed for 1 h at 4°C

(condition i.) or 23°C (condition ii.) and stopped by addition of 0.4 U per applied unit of

RNase I of SUPERase•In RNase Inhibitor (Invitrogen). For condition iii., 1 U of Ambion

RNase I (Invitrogen) per µg RNA and 0.134 µL per µL lysate of TURBO DNase (Invitrogen)

was directly added to the lysate and incubated for 1 h at 4°C. The digest was stopped by

addition of 0.4 U of SUPERase•In RNase Inhibitor (Invitrogen) per used unit of RNase I.

The lysate was centrifuged again for 10 min at 8,000 g and 4°C to remove any cell debris

that may have precipitated during the nuclease digest. For all conditions, monosomes

were pelleted through a 750 µL 30% sucrose cushion prepared in TKM+ buffer

supplemented with 1 mM DTT for 30 min in a S150AT rotor at 72,000 rpm and 4°C. The

resulting ribosome pellets were resuspended in 100 µL of ice-cold ribosome buffer, then

supplemented with 15 mM ethylene diamine tetra-acetic acid (EDTA), pH 8.0 and instantly

mixed with 750 µL Invitrogen TRIzol reagent (Invitrogen) and incubated at room

temperature for 5 min. The solution was mixed with 150 µL chloroform and incubated for

2 min at room temperature before 15 min centrifugation at 20,000 g and 4°C. The

supernatant was mixed with 3 µL GlycoBlue co-precipitant (Invitrogen), 1:10 volume of 3

M sodium acetate, pH 5.5, 1 volume isopropanol and incubated overnight at -20°C. The

precipitated RNA was pelleted by 1 h centrifugation at 20,000 g and 4°C, briefly washed

with ice-cold 80% ethanol and centrifuged again for 30 min at the same parameters. The

resulting RNA pellet was briefly dried and dissolved in 11 µL of nuclease-free water.

For RNA-seq samples, 10 mL of culture were harvested by 2 min centrifugation at

4,000 g at room temperature (before harvesting the culture for footprint isolation), shock-

freezing in liquid nitrogen and storage at -80°C. Samples were lysed in prewarmed (50°C)

lysis buffer (50 mM Tris-HCl pH 8.0, 200 mM NaCl, 20 mM EDTA pH 8.0 and 2% SDS)

and incubated for 2 min at 50°C. Immediately after incubation, 500 µL of TRIzol

(Invitrogen) reagent were added and the mixture was incubated for 5 min at room

temperature. The suspension was mixed with 200 µL chloroform and incubated for

another 5 min at room temperature before 3 min centrifugation at 12,000 g and room

temperature. The nucleic acid containing phase was transferred to a fresh tube and mixed

with 1.5 volumes of ethanol. RNA was purified by the Macherey Nagel NucleoSpin RNA

Plant kit.

*Ribosome footprint extraction and rRNA depletion*

50 µg RNA from the ribosomal extractions were diluted to a volume of 20 µL with nuclease-

free water, mixed with 2x formamide RNA loading buffer (90% deionized formamide, 20

mM Tris-HCl, pH 7.5, 0.5 M EDTA, 0.04% bromophenol blue in ethanol) and denatured

for 90 s at 80°C. The samples were instantly put on ice and then loaded onto 12% urea-

TBE-gels (90 mM Tris, 9 mM boric acid, 2 mM EDTA, 12% 19:1 40% acrylamide-

bisacrylamide mix, 8 M urea). Gels were run at 200 V for approximately 1 h and incubated

with SYBR Gold nucleic acid stain (Invitrogen), diluted 1:10,000 in TBE buffer (90 mM

Tris, 9 mM boric acid, 2 mM EDTA). Ribosome footprints were excised from the gels in a

size range from 20 – 40 nt according to a RNA size marker. Gel pieces were crushed and

incubated overnight with 400 µL footprint elution buffer (300 mM sodium acetate, pH 5.5,

0.25% sodium dodecyl sulfate and 1 mM EDTA, pH 8.0) at room temperature. The eluate

was collected, mixed with 3 µL GlycoBlue co-precipitant, 1 volume of isopropanol and

incubated overnight at -20°C. The precipitated RNA was pelleted by 1 h centrifugation at

20,000 g and 4°C, briefly washed with ice-cold 70% ethanol and centrifuged again for 30

min at the same parameters. The resulting RNA pellet was briefly dried and dissolved in

20 µL of nuclease-free water. For ribosome footprint samples, abundant rRNA fragments

were depleted by incubating 100 - 200 ng of ribosome footprints with 1.19 µL of 100 µM

biotinylated rRNA-oligo depletion mix (consisting of the oligonucleotides:

AATATGCGTTCAAAGATTCGATGATTCACG,

TAGCTCTAGAATTACTACGGTTATCCGAGTA,

TACCCGACGCTGAGGCAGACATGCTCTTGG,

GATTCGTGAAGTTATCATGATTCACCGCA,

ACGGGATGAATCTCAGTGGATCGTAGCA,

CGATCTAGCCGTCTTAGAGCTAGAAGCAGG) with 4 µL formamide, 1 µL 20x

concentrated hybridization buffer (300 mM sodium citrate, 3 M NaCl), 2 µL 0.5 M EDTA,

pH 8.0 and in a total volume of 20 µL. The mixture was hybridized in a thermocycler by

heating up to 80°C for 5 min and subsequent cooling in 5°C steps for 2 min each until

35°C was reached in the device. Afterwards, 60 µL of hybridization buffer, supplemented

with 20% formamide was added to the tube and the sample was depleted twice for 15 min

using 15 µL of Dynabeads MyOne Streptavidin T1 magentic beads (Invitrogen) freshly

washed and prepared according to the manufacturers manual. The supernatant was

mixed with 3 µL GlycoBlue co-precipitant, 2.5 volumes of ethanol and incubated overnight

at -20°C. Footprints were pelleted by 1 h centrifugation at 20,000 g and 4°C, briefly

washed with ice-cold 70% ethanol and centrifuged again for 10 min at the same

parameters. The resulting pellet was briefly dried and dissolved in 41 µL of nuclease-free

water. Excess oligos were removed by 30 min DNase digest at 37°C upon addition of 5

µL 10x TURBO DNase buffer, 2 µL Invitrogen TURBO DNase and 2 µL SUPERase•In

RNase Inhibitor (Invitrogen). After digest, footprints were mixed with 150 µL of 100 mM

NaCl and purified again by phenol-chloroform precipitation and final precipitation

overnight at -20°C using 2.5 volumes of ethanol and 2 µL GlycoBlue co-precipitant.

Footprints were pelleted by 1 h centrifugation at 20,000 g and 4°C, briefly washed with

ice-cold 70% ethanol and centrifuged again for 10 min at the same parameters. The

resulting pellet was briefly dried and dissolved in 36.5 µL of nuclease-free water.

For sequencing library preparation, ribosome footprints were denatured at 65°C for 5 min

in a thermocycler and immediately transferred to ice. The sample was then mixed with 5

µL 10 x T4-polynucleotide kinase buffer A, 2.5 µL T4-polynucleotide kinase (Thermo

Scientific), 2 µL SUPERase•In RNase Inhibitor and incubated for 10 min at 37°C.

Subsequently, 5 µL of 10 mM ATP was added and the sample was incubated for additional

30 min at 37°C. After phosphorylation, the footprints were mixed with 150 µL 100 mM

NaCl and purified again by phenol-chloroform precipitation and final precipitation

overnight at -20°C using 2.5 volumes of ethanol and 2 µL GlycoBlue co-precipitant.

Footprints were pelleted by 1 h centrifugation at 20,000 g and 4°C, briefly washed with

ice-cold 70% ethanol and centrifuged again for 10 min at the same parameters. The

resulting pellet was briefly dried and dissolved in 11.5 µL of nuclease-free water. The

whole sample was used as input for the NEXTFLEX Small RNA-seq Kit v3 (Perkin Elmer)

to generate libraries according to the kit's manual. For RNA-seq, the purified RNA was

later used as input for the Zymo-Seq RiboFree Total RNA library kit for library preparation

according to the manufacturer's manual.

*Processing of raw sequencing data*

Demultiplexed FASTQ files were derived from an Illumina NextSeq 550 system aiming for

40 M reads per footprint sample and 20 M reads per RNA sample at a read length of 75

nt. Samples were processed through a custom-made pipeline in form of a Linux shell

script calling several bioinformatic tools and custom Python scripts. First operation of the

pipeline was removing 3' adapter sequences using cutadapt for RNA-seq and Ribo-seq,

respectively    (Martin,    2011)    with    parameters    [--minimum-length=9,    -

a="TGGAATTCTCGGGTGCCAAGG" ]. After removal of 3' adapter sequences, the 3' and

5' unique molecular identifier tags (UMI tags) were removed from each read's sequence

and shifted to the rear of the header row by UMI tools (Smith et al., 2017) with parameters

[extract --bc-pattern=NNNN] for 5' UMI tags and [extract --bc-pattern=NNNN –3prime] for

3' UMI tags. Remaining reads were filtered for sequence length using cutadapt again with

parameters [--minimum length=20 --maximum-length=39] to remove any read that is

unlikely to represent a real ribosome footprint. After size filtering, a custom-made Python

script corrected the UMI-tags that were previously shifted to the read headers in the form

[header line_NNNN_NNNN] to [header line_NNNNNNNN], allowing UMI tools in a later

step to deduplicate the data set based on both UMI tags. In a first mapping step, reads

were aligned to a set of *C. reinhardtii* non-coding RNA FASTA sequences provided by

ENSEMBL plants together with the genome version 5.5 to remove contaminating RNA

species using STAR with parameters [--runMode alignReads --outSAMtype BAM

SortedByCoordinate --outFilterMultimapNmax 20 --outFilterMismatchNmax 3 --

outReadsUnmapped Fastx --alignIntronMax 7000 --twopassMode Basic]. Reads that did

not align to the ncRNA set were copied to a new FASTQ file that was mapped against the

*C. reinhardtii* genome version 6.1 using STAR (Dobin et al., 2013) with parameters [--

runMode             alignReads             –outSAMtype             BAM

SortedByCoordinate --outFilterMismatchNoverLmax 0.1 --outReadsUnmapped Fastx --

alignIntronMax 8000 –outSAMmultNmax 1 –outMultimapperOrder Random]. After

genome mapping, the resulting BAM file was deduplicated using UMI-tools' dedup

function and afterwards indexed using samtools' index function (Li et al., 2009).

*Data processing*

All analyses on the NGS data sets including calculations of gene body coverage,

periodicity, P-site offsets, organellar read length distributions, ribosome profiles and gene

expression metrics were performed using a custom-made object-oriented Python module

that makes extensive use of the packages pysam/HTSlib (https://github.com/pysam-

developers/pysam), pandas (McKinney, 2010), NumPy (Harris et al., 2020), SciPy

(Virtanen et al., 2020), matplotlib (Hunter, 2007) and seaborn (Waskom, 2021) among

multiple others. For CPM and RPKM calculations in Ribo-seq data sets, only reads

mapping to the annotated CDS of a transcript were considered. Although we are

convinced that RPKM is not a suitable measure to quantify gene expression on the

translatome level, we also calculated RPKM values for the interested reader based on the

length of the annotated ORF of a transcript only (Supplemental Dataset). For any

calculations on RNA-seq data sets, reads mapping to the whole transcript were

considered and also the full length of the mature transcript was used for RPKM

calculations. Translation efficiency values were simply calculated by dividing an CDSs

CPM value from a footprint data set by the respective transcripts RPKM value from the

affiliated RNA-seq data set.

For the 5'-P-site / 3'-A-site offset calculations, all reads fully enclosing either an

annotated start codon (5'-P-site offset) or an annotated stop codon (for 3'-A-site offset)

were taken into consideration separately for the nuclear and the chloroplast genome.

These reads were sorted according to the length of their alignment to the genome, which

was considered as read length to avoid miscalculations due to clipped or masked

sequences. For every read length species, the position of the reads 5' end or 3' end

relative to the position of the first base of the start codon (for 5'-P-site offsets) or the last

base of the stop codon (for 3'-A-site offsets) was determined as the offset. Afterwards, the

most frequently occurring offset for a read length species was determined to be the true

offset of this species and was used in the P-site calculation. To display the results of the

offset calculations, the count matrix of both 5'-P-site offsets and 3'-A-site offsets were

normalized separately to the total number of counts each and plotted as heatmap.

For calculation of the P-sites first position of reads mapping to nucleus-encoded

CDSs, all reads mapping to a CDS were filtered from the data set and for every read, the

5'-most mapping position of the read's alignment + the 5'-P-site offset determined for the

reads alignment length was determined to be the first P-site position in case of transcripts

encoded on the + strand. For reads encoded on the minus strand, the first P-site position

was determined to be the 3'-most mapping position of the RPF alignment minus the

determined 5'-P-site offset. In all cases, splicing was considered and whenever a spliced

read was found, the calculation was adjusted according to the number of genomic

positions spanned by the splice junction of the read. For reads mapping to chloroplast-

encoded transcripts on the + strand, the first P-site position was determined to be the 3'-

most mapping position – (3'-A-site offset + 5). For reads mapping to chloroplast-encoded

transcripts on the - strand, the first P-site position was calculated to be the 5'-most

mapping position + (3'-A-site offset + 5). The metagene P-site profiles of nucleus-encoded

transcripts were calculated considering only the 2,000 most expressed transcripts of a

dataset, according to their CPM values. If multiple splice variants of the same gene

occurred in the list, only the longest variant (annotated as transcript 1 in the genome) was

considered for the calculation to avoid analyzing the same reads multiple times. The

profiles were constructed by calculating the first position of the P-site for all reads mapping

to the first 30 and the last 30 codons of considered transcript CDSs. For a ribosome

footprint to be in-frame with the annotated CDS, the first position of the P-site must be

located on the first base of a codon. According to this logic, the calculated P-site positions

were categorized into "in-frame", "+1 shifted" and "+2 shifted" and summed up for all

codons of considered transcript CDSs. To calculate the frame preference per codon, the

counts in each category in a codon were divided by the sum of all categories for every

codon separately. For calculation of the chloroplast metagene P-site profile, all protein-

coding chloroplast transcripts were considered.

The extension of gene body coverage versus a respective cut-off was calculated

as the portion of an CDS that was covered by the number of reads equal to or greater

than the respective cut-off. This procedure was applied to all annotated CDSs using

increasing cut-offs from 1 to 10 reads to monitor the stepwise decrease of gene-body

coverage. For every applied cut-off, the CDSs were categorized as being covered over

>10% of their length to >90% of their length in 10% steps and CDSs falling into every

category were counted and displayed in a heat map to estimate both completeness and

strength of CDS coverage in the data set.

To calculate the read length distribution, all reads mapping to the respective

genome were filtered and the length of their alignment to the genome was taken as read

length. After counting the number of all possible read lengths for each genome separately,

these numbers were normalized by the total number of reads mapping to the respective

genome. For read biotype distribution, the reads were further categorized if they map to

an annotated coding sequence, 5'- or 3'-UTR or intergenic region. To handle reads

overlapping two regions of different category, for example 5'-UTR and coding sequence

as seen regularly in initiation peaks, the read was assigned the category that had the

larger share on the read's alignment. In rare cases, when a read mapped to different

categories in two different splice variants of the same transcript, the read was assigned to

both categories.

**Supplemental data**

**Supplemental Figure S1:** Features of ribosome protected fragment coverage.

**Supplemental Figure S2:** Transcript accumulation, translation output and translation

efficiency of nuclear genes.

**Supplemental Figure S3:** Correlation between nuclease digestion conditions, transcript

accumulation and translation.

**Supplemental Figure S4:** Correlation analysis of RPF count between nuclease digest

conditions.

**Supplemental Figure S5:** Detection of abnormal initiation peaks in the three genomes of

*Chlamydomonas reinhardtii.*

**Supplemental Dataset.** List of gene body coverage, RPKM and CPM of Ribo-seq and

RNA-seq experiments.

**Acknowledgements**

We thank Fabian Ries, Claudia Herkt and Peter Emelin for critical discussion of the data

and the manuscript. Martin Simon, Jaro Schmitt and Günter Kramer for help with next

generation sequencing. We further thank Jeremy Phillips and David Goodstein from the

Joint Genome Institute for making our data available for the public on the Phytozome

website.

## Funding

This work was supported by the Deutsche Forschungsgemeinschaft grant 437345987 (WI

3477/3-1, ZO 302/5-1, RU 2266/2-1 to F.W., R.Z. and S.R., respectively) and the

Forschungsschwerpunkt BioComp to F.W.

## Author contribution:

V.L.G. performed ribosome profiling experiments, bioinformatic data analyses and wrote

parts of the manuscript. M.K.Y.T. and R.Z. helped with setting up ribosome profiling, N.H.

and S.R. conducted correlation analyses. F.W. designed experiments, analyzed data, and

wrote the manuscript.

## Data availability

All gene accession codes are given in the Supplemental Dataset and are in accordance

with the latest Chlamydomonas genome v6.1 annotation (https://phytozome-

next.jgi.doe.gov) or the GenBank/EMBL data libraries. The full Ribo-seq and RNA-seq

data are accessible via the Dataplant consortium.

## Competing interests

The authors declare no competing interests.

## References

**Aird D, Ross MG, Chen WS, Danielsson M, Fennell T, Russ C, Jaffe DB, Nusbaum C, Gnirke A** (2011) Analyzing and minimizing PCR amplification bias in Illumina sequencing libraries. Genome Biol **12:** R18

**Bertrand EM, Allen AE, Dupont CL, Norden-Krichmar TM, Bai J, Valas RE, Saito MA** (2012) Influence of cobalamin scarcity on diatom molecular physiology and identification of a cobalamin acquisition protein. Proc Natl Acad Sci U S A **109:** E1762-1771

**Blifernez-Klassen O, Berger H, Mittmann BGK, Klassen V, Schelletter L, Buchholz T, Baier T, Soleimani M, Wobbe L, Kruse O** (2021) A gene regulatory network

for antenna size control in carbon dioxide-deprived *Chlamydomonas reinhardtii*
cells. Plant Cell **33:** 1303-1318

**Bonente G, Pippa S, Castellano S, Bassi R, Ballottari M** (2012) Acclimation of
*Chlamydomonas reinhardtii* to different growth irradiances. J Biol Chem **287:** 5833-
5847

**Brar GA, Weissman JS** (2015) Ribosome profiling reveals the what, when, where and
how of protein synthesis. Nat Rev Mol Cell Biol **16:** 651-664

**Bunbury F, Helliwell KE, Mehrshahi P, Davey MP, Salmon DL, Holzer A, Smirnoff N,**
**Smith AG** (2020) Responses of a Newly Evolved Auxotroph of Chlamydomonas
to B(12) Deprivation. Plant Physiol **183:** 167-178

**Calviello L, Mukherjee N, Wyler E, Zauber H, Hirsekorn A, Selbach M, Landthaler M,**
**Obermayer B, Ohler U** (2016) Detecting actively translated open reading frames
in ribosome profiling data. Nat Methods **13:** 165-170

**Cao X, Slavoff SA** (2020) Non-AUG start codons: Expanding and regulating the small
and alternative ORFeome. Exp Cell Res **391:** 111973

**Chartron JW, Hunt KC, Frydman J** (2016) Cotranslational signal-independent SRP
preloading during membrane targeting. Nature **536:** 224-228

**Chen H, Alonso JM, Stepanova AN** (2022) A Ribo-Seq Method to Study Genome-Wide
Translational Regulation in Plants. Methods Mol Biol **2494:** 61-98

**Chiu CW, Li YR, Lin CY, Yeh HH, Liu MJ** (2022) Translation initiation landscape profiling
reveals hidden open-reading frames required for the pathogenesis of tomato yellow
leaf curl Thailand virus. Plant Cell **34:** 1804-1821

**Chotewutmontri P, Barkan A** (2016) Dynamics of chloroplast translation during
chloroplast differentiation in maize. PLoS Genet **12:** e1006106

**Chotewutmontri P, Barkan A** (2018) Multilevel effects of light on ribosome dynamics in
chloroplasts program genome-wide and *psbA*-specific changes in translation.
PLoS Genet **14:** e1007555

**Chotewutmontri P, Barkan A** (2020) Light-induced *psbA* translation in plants is triggered
by photosystem II damage via an assembly-linked autoregulatory circuit. Proc Natl
Acad Sci U S A **117:** 21775-21784

**Chung BY, Hardcastle TJ, Jones JD, Irigoyen N, Firth AE, Baulcombe DC, Brierley I**
(2015) The use of duplex-specific nuclease in ribosome profiling and a user-friendly
software package for Ribo-seq data analysis. RNA **21:** 1731-1745

**Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A,**
**Szczesniak MW, Gaffney DJ, Elo LL, Zhang X, Mortazavi A** (2016) A survey of
best practices for RNA-seq data analysis. Genome Biol **17:** 13

**Costanzo MC, Hogan JD, Cusick ME, Davis BP, Fancher AM, Hodges PE, Kondu P,**
**Lengieza C, Lew-Smith JE, Lingner C, Roberg-Perez KJ, Tillberg M, Brooks**
**JE, Garrels JI** (2000) The yeast proteome database (YPD) and *Caenorhabditis*
*elegans* proteome database (WormPD): comprehensive resources for the
organization and comparison of model organism protein information. Nucleic Acids
Res **28:** 73-76

**Craig RJ, Gallaher SD, Shu S, Salome P, Jenkins JW, Blaby-Haas CE, Purvine SO,**
**O'Donnell S, Barry K, Grimwood J, Strenkert D, Kropat J, Daum C, Yoshinaga**
**Y, Goodstein DM, Vallon O, Schmutz J, Merchant SS** (2022) The
Chlamydomonas Genome Project, version 6: reference assemblies for mating type
*plus* and *minus* strains reveal extensive structural mutation in the laboratory. Plant
Cell

**Cross FR** (2015) Tying Down Loose Ends in the *Chlamydomonas* Genome: Functional
Significance of Abundant Upstream Open Reading Frames. G3 (Bethesda) **6:** 435-
446
**Dall'Osto L, Bressan M, Bassi R** (2015) Biogenesis of light harvesting proteins. Biochim
Biophys Acta **1847:** 861-871
**Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson**
**M, Gingeras TR** (2013) STAR: ultrafast universal RNA-seq aligner. Bioinformatics
**29:** 15-21
**Döring K, Ahmed N, Riemer T, Suresh HG, Vainshtein Y, Habich M, Riemer J, Mayer**
**MP, O'Brien EP, Kramer G, Bukau B** (2017) Profiling Ssb-Nascent Chain
Interactions Reveals Principles of Hsp70-Assisted Folding. Cell **170:** 298-311 e220
**Eberhard S, Drapier D, Wollman FA** (2002) Searching limiting steps in the expression
of chloroplast-encoded proteins: relations between gene copy number,
transcription, transcript abundance and translation rate in the chloroplast of
*Chlamydomonas reinhardtii*. The Plant journal : for cell and molecular biology **31:**
149-160
**Floris M, Bassi R, Robaglia C, Alboresi A, Lanet E** (2013) Post-transcriptional control
of light-harvesting genes expression under light stress. Plant Mol Biol **82:** 147-154
**Forsythe ES, Grover CE, Miller ER, Conover JL, Arick MA, 2nd, Chavarro MCF, Leal-**
**Bertioli SCM, Peterson DG, Sharbrough J, Wendel JF, Sloan DB** (2022)
Organellar transcripts dominate the cellular mRNA pool across plants of varying
ploidy levels. Proc Natl Acad Sci U S A **119:** e2204187119
**Frank J, Agrawal RK** (2000) A ratchet-like inter-subunit reorganization of the ribosome
during translocation. Nature **406:** 318-322
**Fu Y, Wu PH, Beane T, Zamore PD, Weng Z** (2018) Elimination of PCR duplicates in
RNA-seq and small RNA-seq using unique molecular identifiers. BMC Genomics
**19:** 531
**Fujita T, Kurihara Y, Iwasaki S** (2019) The Plant Translatome Surveyed by Ribosome
Profiling. Plant Cell Physiol **60:** 1917-1926
**Gallaher SD, Fitz-Gibbon ST, Strenkert D, Purvine SO, Pellegrini M, Merchant SS**
(2018) High-throughput sequencing of the chloroplast and mitochondrion of
*Chlamydomonas reinhardtii* to generate improved de novo assemblies, analyze
expression patterns and transcript speciation, and evaluate diversity among
laboratory strains and wild isolates. Plant J **93:** 545-565
**Gao Y, Thiele W, Saleh O, Scossa F, Arabi F, Zhang H, Sampathkumar A, Kuhn K,**
**Fernie A, Bock R, Schottler MA, Zoschke R** (2022) Chloroplast translational
regulation uncovers nonessential photosynthesis genes as key players in plant cold
acclimation. Plant Cell **34:** 2056-2079
**Gawronski P, Jensen PE, Karpinski S, Leister D, Scharff LB** (2018) Plastid ribosome
pausing is induced by multiple features and is linked to protein complex assembly.
Plant Physiol
**Gerashchenko MV, Gladyshev VN** (2014) Translation inhibitors cause abnormalities in
ribosome profiling experiments. Nucleic Acids Res **42:** e134
**Gerashchenko MV, Gladyshev VN** (2017) Ribonuclease selection for ribosome profiling.
Nucleic Acids Res **45:** e6
**Gloge F, Becker AH, Kramer G, Bukau B** (2014) Co-translational mechanisms of protein
maturation. Curr Opin Struct Biol **24:** 24-33

Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser E, Taylor J, Berg S, Smith NJ, Kern R, Picus M, Hoyer S, van Kerkwijk MH, Brett M, Haldane A, Del Rio JF, Wiebe M, Peterson P, Gerard-Marchant P, Sheppard K, Reddy T, Weckesser W, Abbasi H, Gohlke C, Oliphant TE (2020) Array programming with NumPy. Nature **585:** 357-362

Hinnebusch AG (2011) Molecular mechanism of scanning and start codon selection in eukaryotes. Microbiol Mol Biol Rev **75:** 434-467, first page of table of contents

Hsu PY, Calviello L, Wu HL, Li FW, Rothfels CJ, Ohler U, Benfey PN (2016) Super-resolution ribosome profiling reveals unannotated translation events in Arabidopsis. Proc Natl Acad Sci U S A

Hunter J (2007) Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering **9:** 90-95

Ingolia NT (2014) Ribosome profiling: new views of translation, from single codons to genome scale. Nat Rev Genet **15:** 205-213

Ingolia NT (2016) Ribosome Footprint Profiling of Translation throughout the Genome. Cell **165:** 22-33

Ingolia NT, Brar GA, Rouskin S, McGeachy AM, Weissman JS (2012) The ribosome profiling strategy for monitoring translation in vivo by deep sequencing of ribosome-protected mRNA fragments. Nat Protoc **7:** 1534-1550

Ingolia NT, Ghaemmaghami S, Newman JR, Weissman JS (2009) Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. Science **324:** 218-223

Ingolia NT, Lareau LF, Weissman JS (2011) Ribosome profiling of mouse embryonic stem cells reveals the complexity and dynamics of mammalian proteomes. Cell **147:** 789-802

Juntawong P, Girke T, Bazin J, Bailey-Serres J (2014) Translational dynamics revealed by genome-wide profiling of ribosome footprints in *Arabidopsis*. Proc Natl Acad Sci U S A **111:** E203-212

Klimmek F, Sjodin A, Noutsos C, Leister D, Jansson S (2006) Abundantly and rarely expressed *Lhc* protein genes exhibit distinct regulation patterns in plants. Plant Physiol **140:** 793-804

Komine Y, Kwong L, Anguera MC, Schuster G, Stern DB (2000) Polyadenylation of three classes of chloroplast RNA in *Chlamydomonas reinhadtii*. RNA **6:** 598-607

Kramer G, Shiber A, Bukau B (2019) Mechanisms of Cotranslational Maturation of Newly Synthesized Proteins. Annu Rev Biochem **88:** 337-364

Kropat J, Hong-Hermesdorf A, Casero D, Ent P, Castruita M, Pellegrini M, Merchant SS, Malasarn D (2011) A revised mineral nutrient supplement increases biomass and growth rate in *Chlamydomonas reinhardtii*. Plant J **66:** 770-780

Lareau LF, Hite DH, Hogan GJ, Brown PO (2014) Distinct stages of the translation elongation cycle revealed by sequencing ribosome-protected mRNA fragments. Elife **3:** e01257

Lauria F, Tebaldi T, Bernabo P, Groen EJN, Gillingwater TH, Viero G (2018) riboWaltz: Optimization of ribosome P-site positioning in ribosome profiling data. PLoS Comput Biol **14:** e1006169

Lei L, Shi J, Chen J, Zhang M, Sun S, Xie S, Li X, Zeng B, Peng L, Hauck A, Zhao H, Song W, Fan Z, Lai J (2015) Ribosome profiling reveals dynamic translational landscape in maize seedlings under drought stress. Plant J **84:** 1206-1218

**Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,**
**Durbin R, Genome Project Data Processing S** (2009) The Sequence
Alignment/Map format and SAMtools. Bioinformatics **25:** 2078-2079
**Liu MJ, Wu SH, Wu JF, Lin WD, Wu YC, Tsai TY, Tsai HL, Wu SH** (2013) Translational
landscape of photomorphogenic *Arabidopsis*. Plant Cell **25:** 3699-3710
**Lukoszek R, Feist P, Ignatova Z** (2016) Insights into the adaptive response of
*Arabidopsis thaliana* to prolonged thermal stress by ribosomal profiling and RNA-
Seq. BMC Plant Biol **16:** 221
**Martin M** (2011) Cutadapt removes adapter sequences from high-throughput sequencing
reads. 2011 **17:** 3
**McKinney W** (2010) Data Structures for Statistical Computing in Python. *In* Proceedings
of the 9th Python in Science Conference, pp 56-61
**Merchante C, Brumos J, Yun J, Hu Q, Spencer KR, Enriquez P, Binder BM, Heber S,**
**Stepanova AN, Alonso JM** (2015) Gene-specific translation regulation mediated
by the hormone-signaling molecule EIN2. Cell **163:** 684-697
**Moulin M, Nguyen GT, Scaife MA, Smith AG, Fitzpatrick TB** (2013) Analysis of
*Chlamydomonas* thiamin metabolism in vivo reveals riboswitch plasticity. Proc Natl
Acad Sci U S A **110:** 14622-14627
**Mussgnug JH, Wobbe L, Elles I, Claus C, Hamilton M, Fink A, Kahmann U,**
**Kapazoglou A, Mullineaux CW, Hippler M, Nickelsen J, Nixon PJ, Kruse O**
(2005) NAB1 is an RNA binding protein involved in the light-regulated differential
expression of the light-harvesting antenna of *Chlamydomonas reinhardtii*. Plant
Cell **17:** 3409-3421
**Nickelsen J, Bohne A-V, Westhoff P** (2014) Chloroplast gene expression - translation.
49-78
**Payne SH** (2015) The utility of protein and mRNA correlation. Trends Biochem Sci **40:** 1-
3
**Pechmann S, Willmund F, Frydman J** (2013) The ribosome as a hub for protein quality
control. Mol Cell **49:** 411-421
**Plancke C, Vigeolas H, Hohner R, Roberty S, Emonds-Alt B, Larosa V, Willamme R,**
**Duby F, Onga Dhali D, Thonart P, Hiligsmann S, Franck F, Eppe G, Cardol P,**
**Hippler M, Remacle C** (2014) Lack of isocitrate lyase in *Chlamydomonas* leads to
changes in carbon metabolism and in the response to oxidative stress under
mixotrophic growth. Plant J **77:** 404-417
**Rooijers K, Loayza-Puch F, Nijtmans LG, Agami R** (2013) Ribosome profiling reveals
features of normal and disease-associated mitochondrial translation. Nat Commun
**4:** 2886
**Russell JB, Cook GM** (1995) Energetics of bacterial growth: balance of anabolic and
catabolic reactions. Microbiol Rev **59:** 48-62
**Schneider-Poetsch T, Ju J, Eyler DE, Dang Y, Bhat S, Merrick WC, Green R, Shen**
**B, Liu JO** (2010) Inhibition of eukaryotic translation elongation by cycloheximide
and lactimidomycin. Nat Chem Biol **6:** 209-217
**Schroda M, Hemme D, Mühlhaus T** (2015) The *Chlamydomonas* heat stress response.
Plant J **82:** 466-480
**Schuster M, Gao Y, Schöttler MA, Bock R, Zoschke R** (2019) Limited Responsiveness
of Chloroplast Gene Expression during Acclimation to High Light in Tobacco. Plant
Physiol

**Sharma P, Wu J, Nilges BS, Leidel SA** (2021) Humans and other commonly used model organisms are resistant to cycloheximide-mediated biases in ribosome profiling experiments. Nat Commun **12:** 5094

**Slobodin B, Dikstein R** (2020) So close, no matter how far: multiple paths connecting transcription to mRNA translation in eukaryotes. EMBO Rep **21:** e50799

**Smith T, Heger A, Sudbery I** (2017) UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. Genome Res **27:** 491-499

**Sun Y, Zerges W** (2015) Translational regulation in chloroplasts for development and homeostasis. Biochim Biophys Acta **1847:** 809-820

**Teixeira FK, Lehmann R** (2019) Translational Control during Developmental Transitions. Cold Spring Harb Perspect Biol **11**

**Trösch R, Barahimipour R, Gao Y, Badillo-Corona JA, Gotsmann VL, Zimmer D, Mühlhaus T, Zoschke R, Willmund F** (2018) Commonalities and differences of chloroplast translation in a green alga and land plants. Nat Plants **4:** 564-575

**Trösch R, Ries F, Westrich LD, Gao Y, Herkt C, Hoppstädter J, Heck-Roth J, Mustas M, Scheuring D, Choquet Y, Räschle M, Zoschke R, Willmund F** (2022) Fast and global reorganization of the chloroplast protein biogenesis network during heat acclimation. Plant Cell **34:** 1075-1099

**Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat I, Feng Y, Moore EW, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I, Quintero EA, Harris CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P, SciPy C** (2020) SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods **17:** 261-272

**Wang R, Amoyel M** (2022) mRNA Translation Is Dynamically Regulated to Instruct Stem Cell Fate. Front Mol Biosci **9:** 863885

**Waskom M** (2021) Seaborn: statistical data visualization. Journal of Open Source Software **6:** 3021

**Westrich LD, Gotsmann VL, Herkt C, Ries F, Kazek T, Trosch R, Armbruster L, Mühlenbeck JS, Ramundo S, Nickelsen J, Finkemeier I, Wirtz M, Storchova Z, Räschle M, Willmund F** (2021) The versatile interactome of chloroplast ribosomes revealed by affinity purification mass spectrometry. Nucleic Acids Res **49:** 400-415

**Wolin SL, Walter P** (1988) Ribosome pausing and stacking during translation of a eukaryotic mRNA. EMBO J **7:** 3559-3569

**Wu CC, Zinshteyn B, Wehner KA, Green R** (2019) High-Resolution Ribosome Profiling Defines Discrete Ribosome Elongation States and Translational Regulation during Cellular Stress. Mol Cell **73:** 959-970 e955

**Wu HL, Song G, Walley JW, Hsu PY** (2019) The Tomato Translational Landscape Revealed by Transcriptome Assembly and Ribosome Profiling. Plant Physiol **181:** 367-380

**Yang X, Cui J, Song B, Yu Y, Mo B, Liu L** (2020) Construction of High-Quality Rice Ribosome Footprint Library. Front Plant Sci **11:** 572237

**Yang X, Song B, Cui J, Wang L, Wang S, Luo L, Gao L, Mo B, Yu Y, Liu L** (2021) Comparative ribosome profiling reveals distinct translational landscapes of salt-sensitive and -tolerant rice. BMC Genomics **22:** 612

**Zhang W, Dunkle JA, Cate JH** (2009) Structures of the ribosome in intermediate states of ratcheting. Science **325:** 1014-1017

**Zoschke R, Bock R** (2018) Chloroplast Translation: Structural and Functional
Organization, Operational Control and Regulation. Plant Cell **30:** 745-770
1049

  **Figures and legends**

1051

**Figure 1: Ribo-seq provides a broad coverage of expressed genes**
(A) Illustration of the experimental setup, testing nucleolytic conditions. Polysome purification and subsequent RNase I digest for 1h at 4°C (i.) or 1h at 23°C (ii.) was performed from Chlamydomonas samples, respectively. Alternatively, cell lysate was directly supplemented with RNase I (iii.). (B) Scatter plots representing the reproducibility of Ribo-seq (left panels) and RNA-seq experiments (right panels). Shown are the R-values and CPM/RPKM values. (C) Schematic representation of the concept of gene body coverage (left panel: incomplete coverage that strongly declines with an increasing read cut-off; right panel: good coverage that is robust against increasing read cut-offs). (D) and (E) Cumulative heatmaps visualizing the number of genes (given in the respective panel) covered to a certain minimal extent (rows) by a certain minimal number of reads (columns) for ribosome footprints (C) and transcripts (D). Ribo-seq data shown are derived from the RNase I digest in lysate (condition iii.). The respective data for the other treatments are shown in Supplemental Figure 1.

**Figure 2: Tri-nucleotide periodicity of cellular translation**
(A) Length distribution of RPFs for each organelle from the three different nuclease treatments tested (i. and ii.: digest of purified ribosomes at 4°C and 23 °C, respectively, iii.: digest in lysate at 4°C). (B) and (C) metagene relative frame preference of nuclear and chloroplast-encoded transcripts per codon for all three nuclease treatments. Codons 1-30 represent the first 30 codons of each transcript while codons 31-60 represent the last 30 codons of each transcript. Frame 0 corresponds to AUG in the ribosomal P-site. (D) Heatmaps representing normalized 5'-P-site offset and 3'-A-site offset counts for nucleus- and chloroplast-mapped RPFs that fully enclose a start or stop codon. Columns represent 5'/3'-RPF end positions relative to the start or stop codon while rows represent the RPF lengths. The shades of red are proportional to the number of RPF ends mapping onto a tile. Arrows indicate the triplet-wise movement of ribosomes in early or late elongation state. Asterisks indicate the tiles harboring the most 5'- or 3'-ends of the 21 and 30 nt RPF-species. Dashed boxes indicate the sudden absence of 21 nt RPFs for the last two codons before translation terminates.

1083

1084

**Figure 3: Chloroplast-encoded proteins are highly expressed**
(A) Violin plots of the RPF data, displaying the CPM (count per million reads) distribution for genes of each plant cell genome, following the three different RNase I treatments (i. and ii.: digest of purified ribosomes at 4°C and 23°C, respectively, iii.: digest in lysate at 4°C). Values with a CPM cut-off <2 (log$_2$) are excluded. (B) Violin plots, displaying distribution of RPKM (reads per kilobase of exon per million reads mapped) for RNA values with a cut-off >2 (log$_2$) for each plant cell genome in replicate 3 exemplary. (C) Scatterplots of the averaged Ribo-seq (CPM) and RNA-seq (RPKM) data for different subcellular-localized proteins. Coloring indicates subcellular localization and origin of the encoded proteins.

1095

**Figure 4: Translation output of chloroplast- and nucleus-encoded transcripts**
(A) Mean read intensity/TE ratio (per respective coding sequence) of chloroplast-encoded transcripts on the level of RPF, RNA and translation efficiency. The transcripts are grouped by function of the encoded proteins (Cyt$b_6f$= cytochrome $b_6f$ complex, PS I = photosystem I, PS II = photosystem II, polym. = RNA polymerase, chl. synth. = chlorophyll

synthesis, misc. = miscellaneous). (B) Values of the 70 most highly expressed nucleus-
encoded genes on the footprint level. Only the longest splice variants were considered.
Respective translation efficiency for these transcripts is given below. For comparison,
transcripts with the top 70 highest TE values are given in Supplementary Figure S2C.
Error bars indicate standard deviation of the three experiments (treatment i. to iii. for
ribosome footprint, and biological replicate 1-3 for RNA-seq), and are shown unidirectional
to improve readability.
1109

1110

1111

**Figure 5: Putative translational regulation of individual transcripts**
(A) Boxplots representing the distribution of Pearson's r correlation coefficients comparing the RNA gene body coverage and ribosome profiles of all chloroplast- and nucleus-encoded transcripts between replicates (R1 and R2 for RNA) and between two types of RNase I treatments (ii and iii for ribosome footprints). (B) Average *Pearson* and *Spearman* correlation coefficients depending on the number n of considered transcripts. Transcripts were ordered by descending RPF count per transcript, n most highly covered transcripts were selected, correlation coefficients were computed for each transcript in this subset, and the average was plotted. For correlation coefficients of individual transcripts see Supplemental Figure S4A. (C) Ribosome profiles of LHCBM1 exemplary for a profile with even footprint distribution and LHCBM6 as an example for a profile with an extreme initiation peak. Red parts of the profile represent coverage within the 5'-UTRs, green parts represent 3'-UTRs and alternating blue shades represent exons within the CDS. Black arrows on the x-axis indicate start and stop codons. (D) Stacked column chart representing the fraction of ribosome profiles containing extreme initiation peaks (defined as transcripts with an average coverage in first 7 codons exceeding by at least 4 times the average coverage of the whole CDS). Nucleus-, chloroplast- and mitochondria-encoded transcripts are plotted separately containing only transcripts with a gene body coverage of >70% in the footprint dataset (transcripts with extreme initiation peak are 3346 out of 8757 for nuclear, 7 out of 66 for chloroplast and 0 out of 7 for mitochondria).

1132

1133

**Figure 6: Utilizing ribosome profiles for coding sequence annotation**
Exemplary ribosome profiles for determining initiation sites and alternatively spliced
transcripts. Red parts of the profiles represent coverage within the 5'-UTRs, green parts
represent 3'-UTRs and alternating blue shades represent exons within the CDS. Black
arrows on the x-axis indicate start and stop codons. (A) and (B) ribosome profiles of
LHCA4 and RPL3 as an example of wrong start codon prediction, respectively. The red
arrow indicates the annotated start codon while the black arrows indicate true start and
stop codons. (C) Ribosome profiles of PDI2 (Cre01.g033550) transcript variants
demonstrating how ribosome profiles can be utilized to identify putative non-cognate start
codon initiation and estimate the dominance of transcript variants.

1145 **Supplemental Figures and Legends:**

1146

1147

1148

1149

**Supplemental Figure S1: Features of ribosome protected fragment coverage**
Supporting Figure 1. (A) scatter plots representing CPM values of transcripts compared
to their length for all three Ribo-seq data sets. R represents the Pearson correlation
coefficient between both variables for each data set separately. (B) and (C) heatmaps
representing transcript numbers categorized by their extent of gene body coverage in
dependence of specific read count cut-offs for Ribo-seq data sets i. and ii.
1156

1157

1158

1159

**Supplemental Figure S2: Transcript accumulation, translation output and translation efficiency of nuclear genes**

Supporting Figure 4. (A) Subcellular distribution of the top 100 expressed nucleus-encoded transcripts on the translation level. If multiple transcript variants of the same gene occurred in the list, only the longest occurring variant was considered. (B) Translation output, transcript abundance and translation efficiency of light harvesting protein encoding transcripts across all three Ribo-seq data sets, considering only the longest transcript variant, respectively. Mean values averaging all three Ribo-seq data are plotted, error bars denote standard deviations. (C) Translation efficiency and translation output of the 80 most efficiently translated nucleus-encoded transcripts. Mean values averaging all three Ribo-seq data are plotted, error bars denote standard deviations. If multiple transcript variants of the same gene occurred in the list, only the longest occurring variant was considered.

1173

1174

1175

1176

1177

**Supplemental Figure S3: Correlation between nuclease digestion conditions,**
**transcript accumulation and translation**
Supporting Figure 5. Boxplots representing the distribution of Pearson correlation
coefficients between ribosome profiles and RNA-seq read distribution of the same
transcripts each in different data sets. Boxes represent the 0.25 to 0.5 and the 0.5 to 0.75
quartiles, lines represent the median values and whiskers represent the remaining data.
Outliers are defined as values being smaller or greater than the respective median ±1.5
times the interquartile range and are depicted as points R1/2/3 indicate different RNA-seq
data sets.

1187

1188

**Supplemental Figure S4: Correlation analysis of RPF count between nuclease digest conditions**

Supporting Figure 5. (A) Relation between correlation coefficient comparing two experimental conditions and average read counts. For each transcript, counts were summed over all positions and then divided by the total number of nucleotides in the transcript. The correlation is high unless the average read counts are low. The distribution of *r*-values might be well suited for selecting good cut-off ranges for a deeper analyses of ribosome profiles. (B) Examples of ribosome profiles with medium and high read counts for all three experimental conditions. Scatter plot of read counts for each nucleotide for comparison of two experimental conditions. Perfectly identical profiles would correspond to a straight line in the scatter plot.

1200

 **Supplemental Figure S5: Detection of abnormal initiation peaks in the three**
 **genomes of *Chlamydomonas reinhardtii***
 Supporting Figure 5. (A) Violin plots representing the distribution of ratios between
 average per nucleotide coverage within the first seven codons and the average per
 nucleotide coverage of the remaining CDS for each organelle's transcripts separately.
 Thick lines within the violins represent the interquartile range while thin lines represent the
 whiskers with a cut-off of 1.5 times the interquartile range. (B) Scatter plot with lines
 representing the increase of the percentile borders of the distributions shown in (A). In
 both cases, only transcripts with a gene-body coverage of at least 70% were considered
 (8757 nucleus-encoded, 66 chloroplast-encoded, 6 mitochondrial-encoded) to minimize
 the risk of misinterpretations due to low or absent coverage in either of both regions.
 Calculation of a 90[th]-percentile border for mitochondrial transcripts was not possible due
 to the low number of transcripts in consideration.