Utilizing high resolution ribosome profiling for the global investigation of gene

expression in *Chlamydomonas*

Vincent Leon Gotsmann[1], Michael Kien Yin Ting[2], Nadin Haase[3], Sophia Rudorf[3], Reimo

Zoschke[2], and Felix Willmund[1]*

Author's institution(s)/affiliation(s):

[1] Molecular Genetics of Eukaryotes, RPTU Kaiserslautern-Landau, Paul-Ehrlich-Str. 23,

67663 Kaiserslautern, Germany.

[2] Max Planck Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam-

Golm, Germany.

[3] Institute of Cell Biology and Biophysics, Leibniz University Hanover, Herrenhäuser-Str.

2, 30419 Hanover, Germany.

15
16

*Corresponding author: Felix Willmund (willmund@staff.uni-marburg.de)

Keywords: ribosome profiling, next generation sequencing, ribosomes, protein synthesis,

translation, transcript accumulation

Short title: Cell-wide ribosome profiling in *Chlamydomonas*

Abstract

Ribosome profiling (Ribo-seq) is a powerful method for the deep analysis of translation

mechanisms and regulatory circuits during gene expression. Here, we established an

optimized and high resolution Ribo-seq protocol for the unicellular model alga

Chlamydomonas reinhardtii (Chlamydomonas). Comparing different nuclease treatments

for the extraction and sequencing of ribosome-protected fragments (RPFs) and parallel

RNA-seq, provided deep insight into translational dynamics and post-transcriptional

control of gene expression, thoroughly covering more than 10,000 different transcripts.

Our high quality Ribo-seq protocol captures the 3-nucleotide movement of elongating

ribosomes along nuclear and chloroplast transcripts. Detailed analysis of the ribosomal

offsets on transcripts uncovers presumable transition states during translocation of

elongating ribosomes within the 5´- and 3´- sections of transcripts and features of

eukaryotic translation termination. These offsets reveal drastic differences between the

nature of cytosolic and chloroplast translation mechanisms. Chloroplast translation is

further characterized by heterogenous RPF size distribution. We found that local

accumulation of small RPFs correlates with local slowdown of *psbA* translation, possibly

revealing an uncharacterized regulator step during PsbA/D1 synthesis. Further analyses

of RPF distribution along specific cytosolic transcripts revealed characteristic patterns of

translation elongation exemplified for the major light harvesting complex proteins, LHCs.

Moreover, our Ribo-seq data can be utilized to survey coding sequence annotations and

the expression preference of alternatively spliced transcripts in Chlamydomonas. We

made these features easily accessible for the research community by attaching our Ribo-

seq data to the most recent Chlamydomonas reference genome.

Introduction

Translation is accomplished via ribosomes, highly conserved macromolecular ribonucleoprotein machines that decode genetic information into linear polypeptide chains resulting in an almost unlimited diversity of proteins. Protein synthesis is the origin for the quantitative and qualitative determination of a proteome´s diversity, which ultimately shapes the biochemical character of a cell. It is now clear that translation output does not strictly follow the changes of transcript abundance, but instead underlies strict regulatory control (Pechmann et al., 2013; Payne, 2015; Slobodin and Dikstein, 2020). The importance of this regulation is illustrated by the fact that bacteria invest about 50% of their available energy for protein biogenesis and that more than 10% of all yeast proteins may be somehow involved in this process (Russell and Cook, 1995; Costanzo et al., 2000). Translational regulation is observed throughout an organism's life span and includes central regulatory circuits that control the cell cycle, tissue differentiation and development, acclimation to environmental changes and the integration of external signals (reviewed in Pechmann et al., 2013; Zoschke and Bock, 2018; Teixeira and Lehmann, 2019; Wang and Amoyel, 2022). Thus, understanding the regulatory principles and mechanisms of translation is a central task of modern biology. With the tremendous advances in next generation sequencing technologies, it is now possible to systematically assay the status of cellular translation by a method called ribosome profiling (Ribo-seq) (Ingolia et al., 2009). The technology is based on the observation that ribosomes occupy precise sections of a translated mRNA (Wolin and Walter, 1988). These mRNA sections (termed ribosome footprints or Ribosome Protected Fragments, RPFs) are extracted by nucleolytic removal of un-occupied mRNA in cell lysates or isolated ribosomes. RPFs are then identified through next generation sequencing and mapping to a reference

transcriptome or genome. Thus, Ribo-seq benefits from the same dynamic range as RNA-

seq, and provides a rich and precise positional information of translating ribosomes from

the physiological state when cells were harvested (Ingolia et al., 2009; Ingolia, 2014; Brar

and Weissman, 2015). Parallel sequencing of all expressed transcripts via RNA-seq

allows the comparison of RNA accumulation with translation and thereby to determine the

fraction of translated transcripts, which is often referred to as "translation efficiency".

Assuming that each RPF represents a translating ribosome and thus a synthesized

nascent polypeptide, mRNAs with high average RPF coverages can be considered as

highly translated transcripts. In addition, local over-representation of specific RPFs within

a specific protein-coding sequence (CDS) are commonly interpreted as regions of slow or

even halted translation (Ingolia et al., 2009; Ingolia, 2014; Brar and Weissman, 2015).

Compared to other techniques, ribosome profiling goes far beyond the targeted assays of

radio-labelling of nascent polypeptides or polysome-loading for specific mRNAs, and

regularly provides much deeper dataset information than proteomic analyses via mass-

spectrometry (Ingolia, 2014; Brar and Weissman, 2015). Yet, ribosome profiling faces the

challenges to distinguish active translation events from non-translating (paused or stalled)

ribosomes. However, high-quality Ribo-seq datasets, with populations of precise read-

length distribution allow the determination of active translation events if RPFs are aligned

relative to the ribosomal Peptidyl-site (P-site). This reveals a 3-nucleotide periodic

movement of translating ribosome along a translated CDS by over-accumulation of reads

containing the first nucleotide of a decoded codon in the P-site. Thus, it is possible to

distinguish ribosome pausing/stalling from translation and therefore even enables *ab initio*

annotation of uncharacterized protein-coding sequences (Ingolia et al., 2009; Calviello et

al., 2016; Hsu et al., 2016).

Ribo-seq has been applied to different plant species with variable gene coverage

and quality, including *Chlamydomonas reinhardtii* (Chlamydomonas hereafter) (Chung et

al., 2015), *Arabidopsis thaliana*, (Liu et al., 2013; Juntawong et al., 2014; Merchante et

al., 2015; Hsu et al., 2016; Lukoszek et al., 2016; Chen et al., 2022), maize (Lei et al.,

2015; Chotewutmontri and Barkan, 2018), and other crop plants (Wu et al., 2019; Yang et

al., 2020; Yang et al., 2021; Chiu et al., 2022). Studying translational regulation in plants

is of specific interest, due to the interplay of three semi-autonomous genomes (nuclear,

chloroplast and mitochondrial), for which many aspects are not fully understood to date.

Chloroplast gene expression seems particularly dependent on co-translational regulation,

which facilitates the fast adjustment of the photosynthesis machinery to environmental

changes and the stoichiometric assembly of chloroplast multi-subunit complexes which

often contain nuclear and chloroplast encoded subunits (Eberhard et al., 2002; Zoschke

and Bock, 2018; Fujita et al., 2019). Recently, ribosome profiling revealed the specific

adjustments made to chloroplast gene expression during acclimation to different light and

temperature regimes (Chotewutmontri and Barkan, 2018; Schuster et al., 2019;

Chotewutmontri and Barkan, 2020; Gao et al., 2022; Trösch et al., 2022). We have

previously applied a high-resolution microarray approach for the fast and cost-efficient

analysis of chloroplast translation in Chlamydomonas, which allowed for the direct

comparison of transcript accumulation and translation output between the alga and land

plants (Trösch et al., 2018; Trösch et al., 2022). We now established Ribo-seq for the

deep analyses of the three Chlamydomonas genomes. Ribo-seq had been described for

Chlamydomonas before, however, with limited depth and analysis (Chung et al., 2015).

However, we aimed to optimize the method to obtain a deeper coverage of genes, with

super-resolution tri-nucleotide periodicity for both nucleus- and chloroplast-encoded

genes and to provide a publicly accessible dataset aligned to the most recent

Chlamydomonas genome (Craig et al., 2022).

Results and Discussion

Establishing super resolution ribosome profiling for Chlamydomonas reinhardtii

Ribosome profiling was performed from logarithmically grown Chlamydomonas cultures

that were kept under mixotrophic conditions and moderate light (see Materials and

Methods). Our previous experience with targeted chloroplast ribosome profiling (Trösch

et al., 2018; Trösch et al., 2022) showed that addition of the elongation inhibitors

chloramphenicol (CAP) and cycloheximide (CHX) (stalling 70S and 80S ribosomes,

respectively) is crucial to prevent ribosome run-off during harvesting (see Materials and

Methods). However, by short incubation of the inhibitors just during harvest, we reduced

the exposure time to the minimum. For nucleolytic digest of RNA that is not protected by

a ribosome, we applied RNase I, consistent with most Ribo-seq studies on other

organisms (Ingolia et al., 2012). Our previous chloroplast ribosome profiling approaches

(using MNase) revealed that the best digestion (i.e. polysome to monosome dissociation)

was achieved when ribosomes were purified prior nuclease treatment (Trösch et al.,

2018). We thus compared RNase I digestion on pre-purified ribosomes both at 4°C

(condition i.) and 23°C (condition ii.) with samples containing whole cell lysates (condition

iii.) (Figure 1A). RNase I was applied at a concentration of 1 unit (U) per micro gram (µg)

RNA, which is similar to published Ribo-seq approaches performed in yeast and

Arabidopsis (Chartron et al., 2016; Hsu et al., 2016; Döring et al., 2017). However, the

RNase I concentration was far below the >9 U per µg RNA that was previously applied for

Chlamydomonas (Chung et al., 2015). By using the lowest possible concentrations of

RNase I, experimental costs can be reduced. Moreover, over digestion of ribosomes may

cause an over-proportional content of contaminating rRNA fragments, which then

dominate sequencing libraries. In the previous Chlamydomonas Ribo-seq study by Chung

et al. (2015), contaminating rRNA fragments accounted for >90% of all sequenced reads.

Our pilot runs had contaminating rRNA levels between 80 and 90%, mainly deriving from

the cytosolic 5.8S rRNA and the chloroplast 23S rRNA. To further reduce the amount of

contaminating rRNA species, we generated anti-sense biotinylated oligos that allow for

the specific depletion of these species. For digest condition iii., we additionally tested

RNase H treatment, which cleaves hybrids of the anti-sense oligos and the target rRNA

contaminants. Other than the standard procedure, we performed rRNA depletion prior to

dual linker ligation and were thus able to provide a higher input of RPFs in the sample,

relative to contaminating rRNAs, which then allows for lowering the number of PCR

amplification steps during final sequencing library generation. Importantly, all these steps

were similar to a Ribo-seq protocol that we established for Arabidopsis and tobacco in

parallel (Ting et al., 2023). The final PCR amplification step of the sequencing library

preparation is known to be a major source of bias by over-representing certain reads (Aird

et al., 2011). Read duplicates were defined as reads that share the same sequence and

the same unique molecular identifier tags (UMI tags), four random bases located between

a RPF and the sequencing adapters at both of its termini (Fu et al., 2018). Resulting

sequencing libraries were thus corrected by deduplicating the final data sets after the

genome mapping step.

Parallel to Ribo-seq, transcript accumulation was determined by sequencing whole

cell RNA samples on the same platform. Ideally, RNA samples are harvested just prior to

ribosome profiling experiments from the same culture. While typical transcriptomic studies

enrich mRNA pools by polyadenylation-enrichment, we here sequenced fragmented small

RNAs that are derived from the full set of transcripts including plastid transcripts, which

frequently have no or short polyadenylation signal sequences and are thus poorly

enriched by standard protocols (Komine et al., 2000; Gallaher et al., 2018).

Ribosome profiling and transcript samples were sequenced with a depth of 40 M

and 20 M reads, respectively, which on average resulted in 25.7% of reads mapping to

annotated CDSs of the most recent Chlamydomonas reference genome v6.1 (Craig et al.,

2022). On average, 64% mapped to putative rRNA loci and 4.2% mapped to annotated

untranslated regions upstream or downstream of CDSs. Despite the application of

different rRNA depletion methods, the rRNA contamination was very similar in all samples.

Of the sequenced reads that mapped annotated CDS a minor fraction (<1%) matched to

more than 10 positions and were removed from further analyses. Of the reads mapping

to CDSs, 96.24% mapped to transcripts of the nuclear genome, 3.63% mapped to

chloroplast-encoded transcripts, and 0.12% to mitochondrial transcripts.

Transcript accumulation was determined by averaging reads per CDS, relative to

the length of each CDS and the total read numbers of the dataset (Reads Per Kilobase

per Million, RPKM), a parameter that is widely used for transcriptome analyses and

accounts for a higher likelihood of measuring RNA fragments of long versus short

sequences (Conesa et al., 2016). This is of particular importance for the sequencing of

small and fragmented RNAs, as commonly accompanied with Ribo-seq. Averaging RPFs

across the respective CDS corrects for local variations of RPF distribution and allows to

get a proxy for the level of translation output per transcript (Ingolia et al., 2011; Ingolia,

2014). For better comparability and consistent with previous Ribo-seq studies, RPKM was

also determined for Ribo-seq data.

Based on RPKM values, the three nuclease digestion conditions (i. to iii.) showed

high reproducibility between the Ribo-seq and RNA-seq experiments, respectively (r =

0.99, Figure 1B). This agrees with previous observations that Ribo-seq tolerates variations

in RNase digest (Gerashchenko and Gladyshev, 2017).

We next aimed to determine the depth of the Ribo-seq approach. Low coverage

Ribo-seq frequently results in ribosome profiles covering only the transcripts with highest

ribosome occupancy (i.e. translation output), whereas moderate or lowly translated CDS

may have partial RPF coverage along the respective sequence (here termed gene body

coverage, Figure 1C, left panel). For Ribo-seq with a good depth, many CDS should show

decent gene body coverage (Figure 1C, right panel). The depth was determined for all

Chlamydomonas transcripts of the v6 genome (32670 transcripts considering the products

of gene copies). For digest condition iii., more than 10,000 transcripts had at least 50% of

their CDSs sequence covered with two or more RPFs, while 6437 transcripts had over

90% of their CDS covered with at least one RPF (Figure 1D). 1508 transcripts are covered

over >90% with an average of more than 10 RPFs/nt, hence representing the group of

transcripts with the most complete gene body coverage combined with a high basal

coverage (see also Figure 4B). Categorizing the transcripts in this way allows for the

assessment of the general quality of a data set, as well as the selection of transcripts of

similar quality for specific analyses without having to rely on expression values alone. The

gene body coverage was highest for digest condition iii. (Figure 1D and Supplemental

Figure S1), and comparably good for the coverage of RNA fragments for the RNA-seq

experiment (Figure 1E).

The read length distribution of the RPFs mapping to annotated CDSs of the nuclear

genome displayed high comparability to published Ribo-seq approaches with other

organisms. RNase I treatment typically results in RPFs with a length between 28 and 30

nt. Here, digest of purified ribosomes at 4°C resulted in RPFs with a predominant length

of 31 nt. Digest condition ii. caused a broader distribution of RPFs between 28 and 30 nt,

while digest in cell lysate resulted in a sharp peak of RPFs with a length of 30 nt (Figure

2A). Such a sharp peak at 30 nt is typically found in Ribo-seq from mammalian cells when

harvested in the presence of CHX (Wu et al., 2019; Sharma et al., 2021). The sharp peak

is even more prominent if only those RPFs are considered that map to the CDS

(Supplemental Figure S2A). This is most evident for the size distribution mapping to the

CDS of the 8 mitochondrial transcripts. Here, RPF accumulations peaks at a size of 33 nt

(Supplemental Figure S2A). The larger size of mitochondrial RPFs was described before

and could stem from the bulkiness of the membrane associated ribosomes (Rooijers et

al., 2013). Whereas, cytosolic and mitochondrial ribosomes produced sharp RPF peaks,

the RPF size distribution of chloroplast transcripts is less defined and may be the result of

diverse translational states, as previously reported for maize and Arabidopsis

(Chotewutmontri and Barkan, 2016; Gawronski et al., 2018; Fujita et al., 2019) (see

below). The sharpest peak for chloroplast RPFs is seen under condition ii., albeit the

predominant RPFs of 28 nt is found under all treatment conditions (Figure 2A). While

consistent with the observations in other organisms, our RPF size distribution varies from

the previous Ribo-seq data with Chlamydomonas, which reported a general shift towards

smaller RPF sizes, maybe as consequence from their harsh RNase I treatment (Chung et

al., 2015).

233   A hallmark of high-resolution Ribo-seq data is the triplet-wise 3-nt periodicity

of RPFs that represents slowdown of ribosome movement as ribosomes decode a

respective codon (Ingolia, 2016). To calculate periodicity, it is necessary to determine the

5'-offset for every RPF length species, which is the number of bases located 5' to the P-

site (the translational state of polypeptide bond formation) of a respective RPF. This is

usually done by considering all RPFs that fully enclose an annotated start codon and

determining the frequency of all occurring 5'-offsets for every RPF length separately. The

5'-offset, most frequently occurring for a specific RPF length, is then declared to be the

true 5'-offset of this species. This strategy is based on the fact that RPF occupancy of

transcripts usually inclines dramatically at the borders between 5'-untranslated regions

and CDS, with the start codon often having an exceptionally high coverage, which is due

to the slow process of translation initiation (Supplemental Figure S2B). The 5'-offsets

determined in this way are used to calculate the exact P-site position and reading frame

of every RPF. By this, 3-nt periodicity can be nicely visualized in meta-plots combining all

genes while 3-nt plots of individual transcripts can be noisy, depending on translation

levels. The exact 5'-offset varies in an organism- and protocol-specific manner and is

usually 12 to 13 nt. Alternatively, the P-site position has also been determined by

calculating the 3'-offset of the ribosomal Aminoacyl-site (A-site), applying the same

strategy to RPFs mapping to annotated stop codons.

With the majority of cytosolic RPFs with one read length (Figure 2A), all three

RNase I digest conditions yielded a clear frame preference of "frame 0" for nucleus-

encoded transcripts, meaning that the base triplet found at the P-site of a RPF is for the

for the majority of RPFs in-frame with the annotated CDS. Again, lysate digest (iii.)

resulted in the best frame preference values (>68% of RPFs in frame 0, 19.4% in frame 1

and 12.5% in frame 2), suggesting that this condition yields Ribo-seq data of the best

quality (Figure 2B and Supplemental Figure S2B). Due to the differences between 80S

and 70S ribosomes, 3-nt periodicity needs to be determined separately for chloroplast

translation. For chloroplast transcripts, the frame preference was less obvious, albeit it

was clearest for digest condition iii. (Figure 2C and Supplemental Figure S2B).

Considering the overall quality of sample iii. (the similar degree of rRNA contamination

between all three samples and the high correlation with the other samples on RPKM level),

it can be concluded that RNase H mediated rRNA depletion has no adverse effects on the

Ribo-seq sample quality and seems to be similarly effective.

To get a better insight into the RPF offsets and periodicity, different lengths of the

RPFs were plotted against their offset frequencies upon P-site and A-site alignment for

the dataset of condition iii. (Figure 2D). The triplet-wise movements of the 5´ends with an

offset of 12 nt during initiation is clearly visible for cytosolic RPFs and most prominent for

the 30 nt RPFs of digestion conditions iii., consistent with the observation in other

eukaryotic organisms (Lauria et al., 2018). Likewise, triplet-wise movements of the 3´-

ends are also detectable and are in frame with the stop codon. Here, a 9 nt long 3´-offset

seems the predominant form prior to termination with the stop codon in the A-site of the

ribosome. However, due to the 12 nt 5'-offset and 3 nt occupied by the P- and A-sites

respectively, a 3'-offset of 12 nt was expected. This discrepancy could be a result of CHX

treatment, which locks ribosomes in a pre-translocation conformation after peptide bond

formation (Wu et al., 2019) so that RPFs on non-stop codons may exceed the natural

termination-caused accumulation on stop-codons. Moreover, a slow termination process

might arrest ribosomes on the codon just upstream of the stop, prolonging ribosome dwell

time at this site. At the same time, ribosomes that already entered termination during

sampling are likely to complete the process and dissociate from the transcript. CHX does

not inhibit termination, which in turn would dramatically lower the number of FPs mapped

to stop codons with their A-site. Interestingly, RPFs with a length of 21 nt are detectable

upon initiative (asterisks, Figure 2D). These fragments have been previously

characterized as RPFs of ribosomes in an open ratchet conformation during translocation,

even accumulating as abundant species without CHX treatment in yeast (Lareau et al.,

2014). During elongation, ribosomes undergo massive conformational changes by

rotating the large subunit relative to the small subunit (Frank and Agrawal, 2000; Zhang

et al., 2009). The fact that the RPFs produced from the open-ratchet conformation have

the same 5'-offsets as the main RPF species despite being considerably shorter, suggests

that the RNase exclusively removes nucleotides from the 3´-end of RPFs in this

conformation (and could indicate that the center of the rotation is located towards the 5'-

direction, viewed relative to the RPF). Cycloheximide was shown to prevent binding of the

next aminoacylated tRNA in the A-site during peptide bond formation (Schneider-Poetsch

et al., 2010) and thus stabilizes ribosomes in a specific stage during the elongation cycle,

in which RPFs of 28-30 nt are the prevalent form. Interestingly, human 80S ribosomes are

not exclusively arrested in one conformation upon CHX treatment (Sharma et al., 2021),

comparable to the situation seen here for Chlamydomonas 80S ribosomes (Figure 2D,

upper panel). These 20-22 nt RPFs might derive from a post-translocation ribosome

species (Lareau et al., 2014). The presence of both species demonstrates active

movement of ribosomes along the CDS. The absence of the "open-ratchet" 21 nt RPFs

starting from two peptide-bond formations prior to termination, may support the idea from

above that ribosomes near stop codons are paused, which can prevent the collision

between ribosomes (Figure 2D upper panel, dashed box). Interestingly, the frame

preference for frame 0 of the stop-minus-2 codon is the highest in the whole metagene

periodicity plot (79.6% Figure 2B), which could denote a short stalling at this codon before

translocation. However, for the next codons the frame 0 preference drops considerably to

55.8%, especially in favor of frame 1 (36.2%) and then to 33.1% at the stop codon, clearly

indicating that translation dynamics change towards the termination step. Moreover, these

characteristics are accompanied by a continuous decline of absolute numbers of reads

mapping to these codons on the meta-gene level (for digest iii: stop-2: 19890, stop-1:

6786, stop: 522).

The prokaryotic-type chloroplast RPFs 5'-offsets seem to be much more diverse

than their cytosolic counterparts, apparent by their multiple clusters on the heat map

(Figure 2D, bottom panel). These might resemble populations of initiation complexes in

different states or in complex with transcript specific RNA processing factors and *trans-*

acting factors that interact with ribosomes (Westrich et al., 2021). Without clear criteria

how to distinguish and interpret these clusters, 5'-offset definition is problematic, since

multiple 5'-offset frequencies of comparable amplitude exist for many RPF lengths.

However, the chloroplast 3'-offsets seem less scattered and reveals a main RPF species

of 28 nt length with 11 nt 3'-offset. This is in accordance with the RPF length distribution

of all chloroplast RPFs, where the 28 nt species also represents the main species,

although the distribution is much broader than for the cytosolic RPFs. Similar to the

cytosolic offset heat map, a triplet-wise movement of ribosomes can also be observed for

the terminating chloroplast ribosomes in the 3'-offset heatmap for 29 nt RPFs, albeit with

much lower clarity. Altogether, the observations made for chloroplast offsets and RPF

length distribution might indicate a certain heterogeneity of the chloroplast ribosome pool,

which is an interpretation that would be in line with the large number of proteins that

interact with chloroplast ribosome found in our previous study (Westrich et al., 2021) and

the large number of RNA-binding proteins known or suspected to regulate the translation

of specific chloroplasts transcripts (Nickelsen et al., 2014). This is remarkable considering

the very small number of transcripts encoded on the chloroplast genome (72). Together,

this might suggest that translation in the chloroplast of Chlamydomonas is heavily

influenced by transcript-specific features rather than following a universal scheme

(Zoschke and Bock, 2018).

Differences in translation output between cytosolic and organellar transcripts

We next compared the translation levels of the organellar versus nuclear transcripts.

Chlamydomonas cells contain on average 83 plastid genomes within the large cup-

shaped chloroplast occupying about 40% of the cellular volume. The mitochondrial

genomes exist in 130 copies, also greatly outnumbering the one haploid nuclear genome

(Gallaher et al., 2018). Despite the rather low contribution of organellar RPFs relative to

the total RPF pool, both chloroplast and mitochondrial transcripts had higher RPKM values

when compared to the average RPKM values of nuclear genes (Figure 3A). A similar trend

is seen for transcript levels (Figure 3B), suggesting that organellar genes are expressed

at considerably higher levels compared to most nuclear genes (Forsythe et al., 2022).

346   With the high reproducibility of Ribo-seq and RNA-seq data, we chose to average

the datasets for estimating the levels of transcript accumulation and translation output. A

direct comparison between translation output and transcript accumulation is frequently

referred to as translation efficiency (TE) and normalizes RPF values over transcript

accumulation. Contrasting translation over transcript data of the Chlamydomonas genes

in a scatter plot and highlighting their subcellular localization showed that nuclear

transcripts encoding chloroplast-localized proteins have higher expression values when

compared to proteins of cytosolic or Endoplasmic Reticulum localization (Figure 3C),

again supporting the notion that many chloroplast proteins are expressed at high levels.

Interestingly, most nucleus-encoded transcripts showed similar levels between transcript

accumulation and translation (distributed along the diagonal, Figure 3C). In contrast,

chloroplast- and mitochondria-encoded transcripts showed higher transcript accumulation

levels, when compared to translation, which agrees with the published data of profound

potential translational regulation within the chloroplast (Sun and Zerges, 2015; Trösch et

al., 2018; Zoschke and Bock, 2018). Remarkably, translational regulation seems also

predominant for the few mitochondrial genes.

Column plots of the individual chloroplast transcript and translation values show

that even the weakest expressed known chloroplast transcripts have RPKM values ~10,

which is within the median range of nucleus-encoded transcripts (Figure 4A versus 3A).

Not surprisingly, transcripts that encode the core photosystem I subunits PsaA and PsaB,

the photosystem II core subunits PsbA/D1,PsbB/CP47, PsbC/CP43, PsbD/D2, the

Rubisco large subunit, RbcL, the CF_0 ATPase subunit AtpH, and the translation elongation

factor TufA show the highest translation levels, consistent with our previous chloroplast

ribosome profiling experiments and the examination of protein accumulation levels

(Schroda et al., 2015; Trösch et al., 2018). We further detected transcripts of the

uncharacterized chloroplast Open Reading Frames (ORF) *orf202* (positioned within the

transposon Wendy I), *orf854* (positioned within the transposon Wendy II), *orf528*, and *i-*

crel (Gallaher et al., 2018), however, translation output was low or even undetectable (for

orf2020 and *i-crel*), suggesting that these transcripts are barely translated in

Chlamydomonas cultures, at least under standard conditions (Figure 4A). The direct TE

ratios of the chloroplast transcripts again show the imbalance between chloroplast

transcript accumulation and translation output, suggesting that the long-living chloroplast

transcripts present a buffering capacity for the fast and efficient adjustment of translation

levels during environmental changes, as for example seen during light and temperature

acclimation and most prominent for PsbA/D1 synthesis (Chotewutmontri and Barkan,

2018; Schuster et al., 2019; Trösch et al., 2022).

We next determined translation output levels of nucleus-encoded transcripts, only

considering the longest transcript variant if multiple splice forms existed. Of the transcripts

with the highest translation output, 51 protein products have a known or predicted

chloroplast localization and 47 are present in the nucleus or cytosol, again showing the

high accumulation levels of chloroplast proteins (Supplemental Figure S3A). Amongst

those, transcripts of the light harvesting complex proteins (LHCs) have the highest

translation output, consistent with their high protein accumulation levels in plant cells

(Dall'Osto et al., 2015). Generally, *LHC* translation showed comparable RPKM translation

output values, whereas transcript accumulation was rather variable. For the transcripts of

the high-light induced photoprotective LHCSR3.1 and LHCSR3.3 proteins, we could not

determine translation profiles (Supplemental Figure S3B), suggesting that these proteins

are not synthesized under the investigated growth conditions (Bonente et al., 2012).

Besides photosynthesis related transcripts, the list with top-most translation contained

transcripts encoding numerous ribosomal proteins of the 80S complex and elongation

factors (Figure 4B). Other abundantly translated proteins are the putative acetate

transporter GYF3 (RPKM 4285, rank 14), the Isocitrate lyase (ICL1, RPKM 3918, rank

16), a key enzyme of the glyoxylate cycle (Plancke et al., 2014), the thiazole biosynthesis

enzyme (THI4, RPKM 3402, rank 22) (Moulin et al., 2013) or the Methionine Adenosyl-

transferase (METM1, RPKM 2438, rank 51), confirming their central role in

Chlamydomonas cells. Plotting of transcripts with the highest TE values showed an over

representation of histone transcripts, suggesting their robust translation in

Chlamydomonas cells under logarithmic growth (Supplemental Figure S3C).

Exploiting individual ribosome profiles to mine for putative regulatory principles

Local elongation rates are known to vary considerably during translation along a transcript,

depending on tRNA availability, mRNA secondary structure, the amino acid sequence of

the resulting polypeptide and co-translational folding (Gloge et al., 2014; Kramer et al.,

2019). With the robust gene body coverage over Chlamydomonas transcripts (Figure 1D),

we aimed to learn more from the dynamics of translation elongation, derived from the RPF

distribution over individual coding sequences (ribosome profiles). Ribosome profiles were

highly comparable between the three RNase I treatment conditions i.-iii., as indicated by

median *Pearson* correlations between data sets ii. and iii. of ~0.62 for nucleus-encoded

transcripts and >0.85 for chloroplast-encoded transcripts. Although RNA-seq read

distributions over individual CDS were even more comparable (median of *r*-value 0.87 for

nucleus-encoded and 0.99 for chloroplast-encoded transcripts) among the RNA-seq data

sets, they showed no correlation (median *r*-value close to zero) to the respective ribosome

profiles (Figure 5A and Supplemental Figure S4). Certainly, methodological-driven

differences between Ribo-seq and RNA-seq cannot be fully excluded, but it can be

assumed that the ribosome profiles reflect transcript-specific translation dynamics and are

likely not the result of sequencing biases.

The correlation coefficients for specific transcripts comparing two experiments

(both *Pearson* and *Spearman*) stably range between 0.7 and 1 for the majority of ribosome

profiles that have an average coverage >100 reads per nucleotide, suggesting that low

correlation coefficients are the consequence of sequencing depth limitations and thus

limited observability of lowly translated transcripts rather than the true absence of a

characteristic translation dynamic of these transcripts (Supplemental Figure S5).

Importantly, more than 10,000 transcripts have *r*-values >0.7 (Figure 5B), again showing

the high reproducibility between the RNase I digest conditions for transcripts with

moderate and high RPF coverage (Supplemental Figure S5). In fact, the *r*-value

distribution shows that these direct comparisons of read coverage per transcript might be

well suited for selecting good cut-off ranges for a deeper analyses of ribosome profiles. In

many previous studies, simply minimum RPKM values (e.g. 1) were taken as cut-off.

Ribosome profiles of the highly translated *LHC* transcripts showed some

remarkable differences between the individual *LHC* species. Efficiently controlling the

expression of *LHC* genes is essential for regulating the antenna size of the photosynthesis

machinery in order to capture photons under low irradiance conditions (increased

expression) or avoid over-excitation and thus photodamage under high irradiance

conditions (reduced expression) (Dall'Osto et al., 2015). Of the transcripts encoding the

21 different LHC members serving Photosystem I or II (LHCA and LHCB, respectively),

20 were found amongst the top 100 highest translated transcripts (not considering splice

variants). Only the transcript encoding LHCB7, a rather newly discovered protein

containing the unusual number of 4 transmembrane domains (Klimmek et al., 2006) had

low RPKM values of ~15 (Supplemental Figure S3B). The variations in *LHC* transcript

levels result in considerably different TE values (Supplemental Figure S3B), suggesting

that translational regulation might be achieved by different strategies for expression of the

LHC species. Indeed, the ribosome profiles are considerably different between transcripts

encoding the different LHCs. For example, *LHCBM6* showed an exceptionally strong peak

of RPFs covering the first 7 codons, which exceeds the average RPF density over the

remaining CDS by more than 4-fold (Figure 5C). While "initiation peaks" are commonly

accumulating over the first few codons of a CDS, their amplitudes are usually within the

dynamic range of the remaining CDS. Such high peaks, in contrast, could point to a

regulative mechanism stalling ribosomes during or shortly after initiation. Indeed,

LHCBM6 was shown to be translationally repressed under moderate light conditions by a

co-translational regulator termed NAB1 (Nucleic Acid Binding protein). NAB1 itself is

tuned by a regulatory circuit sensing carbon dioxide supply levels in Chlamydomonas, for

precisely adjusting the antenna composition of photosystem II (Mussgnug et al., 2005;

Blifernez-Klassen et al., 2021). Similarly, *LHCBM4* is recognized by NAB1, also

presenting a strong RPF peak over the initiation codon, whereas LHCBM1 (not a target of

NAB1) has no over-proportional initiation peak (Figure 5C), despite sharing a 75% amino

acid sequence identity with LHCBM6. Translational regulation during *LHC* expression is

not unique to Chlamydomonas, as *Lhcb4.2*, *Lhcb4.3* and *Lhcb6* are also subject to post-

transcriptional regulation in Arabidopsis (Floris et al., 2013). We searched all transcripts

that had a gene body coverage of at least 70% (8757 nucleus-encoded, 66 chloroplast-

encoded, 6 mitochondria-encoded) for the occurrence of comparable initiation peaks

(average coverage of the first seven codons at least four times higher than the average

coverage of the remaining CDS). Around 38% of selected nucleus-encoded transcripts

had comparable RPF accumulation peaks near the initiation codon, possibly hinting to

more co-translational regulation in the cytosol than previously anticipated. In contrast, only

10% (6 transcripts) of the chloroplast-encoded genes showed such peaks (Figure 5D).

However, it cannot be excluded that these peaks present artifacts of CHX treatment of the

Chlamydomonas cultures during harvest. Upon establishment of the Ribo-seq protocol,

several reports demonstrated that pre-incubation of yeast cultures with CHX may create

a bias by inducing increased RPF accumulation at initiation codons, which could be the

consequence of stalled elongation but unaffected initiation through CHX treatment (Ingolia

et al., 2009; Gerashchenko and Gladyshev, 2014; Sharma et al., 2021). However, a recent

direct comparison of CHX-treated and untreated conditions demonstrated that CHX

causes no initiation-peak bias in human cells, and that even the strong accumulation in

yeast might derive from slightly altered protocols between laboratories (Sharma et al.,

2021). For Chlamydomonas, CHX biases should result in general initiation peaks, but

these peaks are only found in a fraction of transcripts, independent of their translation

activity (e.g. *LHCs*). Thus, varying accumulation of RPFs around the initiation codon may

be an interesting indicator for altered initiation regulation between transcripts in the

cytosol.

We further assayed, if the RPFs of various sizes, ranging from 24 to 34 nt, are

randomly distributed across all chloroplast CDS or if a respective RPF size is transcript

specific. In general, the dominant RPF size is 28 nt long and the trimodal size distribution

appears widely present for translation of most chloroplast transcripts, even if the

overrepresentation of the highly translated transcripts (i.e. *rbcL, psbA)* is normalized out

(Figure 6A). While present for most chloroplast transcripts, accumulation of the different

RPFs sizes varies between transcripts with some over-representation of smaller or larger

RPFs, respectively. For example, *psbA, psbH, psaC, psaJ, petD* and *atpH* have an

additional strong peak at 24 nt. In contrast, *rpl20* and *psbJ* have a shift towards a dominant

population of RPFs with a size >30 nt (Figure 6B-D). Strikingly, the small RPFs within

psbA and *atpH* show in interesting patter with increased accumulation in the regions of

highest RPF coverage. For *pbsA,* the 24 nt RPFs are most prominent in a section between

codon 135 and 150, which is just downstream of the transcript encoding the second

transmembrane segment (TMS, Figure 6E). It appears as if elongation severely slows

down within this region, favoring the extraction of small RPFs, possibly resulting from

ribosomes in an "open-ratchet" formation. Alternatively, the smaller reads may result from

local binding of a regulatory factor to the transcript, although the small RPFs have similar

features as the longer RPFs. Interestingly, we just recently reported that deletion of the

One-Helix-Protein 2, OHP2, locally affects *psbA* translation between initiation and the

coding sequence of the 3rd TMS. However, ribosome occupancy downstream of the 3rd

TMS appeared normal in the mutant. In addition, the absence of OHP2 results in a rapid

degradation of nascent PsbA/D1 protein (Wang et al., 2023). Thus, the observed local

translation slow-down might be an important stage for PsbA/D1 biogenesis and co-factor

integration and OHP2 might be the driver of this regulation. Similarly, RPF accumulation

peaks over the CDS of the first TMS of *atpH,* mostly deriving from fragments with a size

of 24-25 nt, again suggesting that translation elongation is reduced here (Figure 6E). It

will be interesting, to uncover in future experiments if the local translation slowdown is

accompanied or caused by binding of regulatory factors.

Suitability of ribosome profiles for improving Chlamydomonas genome annotation

Lastly, we aimed to exploit our Ribo-seq data for surveying correct CDS

annotations and transcript splice variants within the recently released Chlamydomonas

genome, version 6.1 (Craig et al., 2022). In previous genome versions, several genes

were falsely annotated, ignoring upstream start codons that were possibly the genuine

initiation site of translation (Cross, 2015; Craig et al., 2022). In the new version, the first

in-frame ATG codon was generally considered as the initiation codon, in accordance with

the scanning model of initiating ribosomes (Hinnebusch, 2011). Systematic comparison

of the reference start site with the onset of RPF coverage, confirmed that start codon

assignment has been clearly improved between genome version 5.6 and 6.1. However,

we detected gene models in which the first ATG appears not to be the genuine initiation

site. For example, transcripts encoding LHCA4 (Cre10.g452050) do not initiate from the

first ATG (Figure 7A) and were correctly annotated in the previous genome versions.

Another example is PRPL3 (Cre10.g417700) (Figure 7B). In addition, we determined

some gene models in which RPF accumulation did not correlate with the cognate ATG

start codon, possibly hinting to a translation start from non-cognate start sites as

exemplified by the *PDI2* CDS (Cre01.g033550, Figure 7C). In this case, the ribosome

profile starts with a prominent peak upstream of the annotated start codon located within

the annotated 5'-UTR while the annotated start codon is poorly covered compared with

the remaining CDS. Here, initiation from the non-cognate TTG could be an explanation

(Cao and Slavoff, 2020), which is located in-frame with the annotated coding sequence at

two positions within the area covered by the peak. Ribo-seq was already used before to

mine for non-cognate start sites in other organisms (Brar and Weissman, 2015).

Alternatively, these RPFs could point to the presence of upstream Open Reading Frames

for *PDI2*. *PDI2* represents a good example for detecting alternative splicing. Transcript 1

of *PDI2*, possesses a RPF gap between exon 3 and 4 (379 nt downstream of the

transcription start site), which is spliced out in transcript versions 2-4 (Figure 7C). In

addition, a sudden drop of RPFs is seen at the end of exon 3 in transcript 2 and 4. Based

on distribution of RPFs and the transition of ribosome profiles, transcript 5 has the most

consistent profile and might hence be the prevalently expressed variant in

Chlamydomonas cells, grown under standard laboratory conditions. In future, intelligent

modelling and peak deconvolution algorithms could be potentially applied to Ribo-seq

data to estimate the stoichiometric ratio of different splice variants' translation output.

Concluding remarks

Taken together, ribosome profiling is a fascinating tool for advancing our understanding

of post-transcriptional regulation of gene expression. Our Ribo-seq approach has the

resolution and depth to deeply study translation of the nuclear and chloroplast transcripts

in Chlamydomonas cells. Furthermore, a highly comparable Ribo-Seq protocol was

established for Arabidopsis and tobacco by our team, which will help to directly compare

translation between different land plants and Chlamydomonas (Ting et al., 2023).

Chlamydomonas is a well-suited organism for understanding system level changes of

gene expression throughout the cell cycle or following environmental changes. Ribosome

profiles can be of specific help, for a first understanding of the many transcripts, which

were not experimentally investigated before. To make our data easily accessible in the

research community, we have now linked our Ribo-seq data to the recent

Chlamydomonas genome CC-4532 v6.1 (https://phytozome-next.jgi.doe.gov), thereby

allowing individuals to conveniently browse the ribosome profiles of their genes of interest.

With the new genome release, the number of alternatively spliced transcripts have been

significantly expanded. Visualization of ribosome profiles may help to uncover the

dominant transcript variants, which are translated under standard laboratory conditions.

Certainly, additional experimental approaches are required to understand unexpected

features such as translation initiation from non-cognate start sites, RPFs along upstream

open reading frames, mechanism of translational regulation, and spatial organization of

protein synthesis.

Materials and Methods

Cell growth and harvest

Chlamydomonas reinhardtii CC-1690 cells were grown mixotrophically in Tris-Acetate

Phosphate (TAP) medium (Kropat et al., 2011) to mid-log phase (4 to 5×10^6 cells per mL)

under constant light of 80 mmol m^{-2} s^{-1} (MASTER LEDtube HF 1,200 mm UO 16W830 T8

and 16W840 T8, Philips) and 25°C on a rotary shaker. Immediately before harvest, 100

µg/mL chloramphenicol (CAP) and cycloheximide (CHX) was added and cells were rapidly

chilled by pouring over -80°C cold silicon ice cubes until 4°C culture temperature was

reached. Subsequently, cells were pelleted by 2 min centrifugation at 4,000 g and 4°C,

washed in ice-cold TKM+ buffer (50 mM acetate buffered Tris pH 8.0, 150 mM KCl, 10

mM $MgCl_2$, 100 µg/mL CAP, 100 µg/mL CHX) and pelleted again using the same

centrifugation conditions. Afterwards, the pellet was resuspended in freezing buffer (TKM+

supplemented with 100 mM phenyl methyl sulfonyl fluoride (PMSF) and 16% of 25x Roche

protease inhibitor cocktail) by slow pipetting in 1/2000 of the initial culture volume and

flash frozen by dripping the cell suspension into liquid nitrogen. Frozen cells were stored

at -80°C until further use.

Ribosome profiling and RNA sampling

Frozen cells were ground using a bead mill with nitrogen-cooled steel containers and

beads for two times 2 min at 27 Hz frequency and cooling in liquid nitrogen between

rounds. The cell powder was mixed with equal volumes of 2x concentrated lysis buffer (2x

concentrated TKM+ buffer supplemented with 2 mM dithiotreitol (DTT), 2% Triton X-100

and 20% sucrose) and incubated for 5 min at 4°C before pelleting cell debris at 8,000 g,

4°C for 10 min. For nucleolytic digest conditions i. and ii., lysate was layered on a 2.5 mL

64% sucrose cushion prepared in TKM+ buffer supplemented with 1 mM DTT and

ultracentrifuged for 3 h at 60,000 rpm and 4°C in a Ti70 rotor. The resulting polysome

pellet was briefly rinsed with 500 µL ice-cold diethyl pyro carbonate (DEPC)-treated water

and resuspended overnight in 100 µL ice-cold ribosome buffer (2x concentrated TKM+

buffer diluted with 25x concentrated Roche protease inhibitor cocktail to a total amount of

16%, 1 mM PMSF and 0.1% Triton X-100). Remaining insoluble debris were removed by

centrifugation for 1 min at 1,500 g before addition of 1 U of RNase I (Ambion) per µg RNA

and 0.134 U TURBO DNase (Invitrogen). Nucleolytic digest was performed for 1 h at 4°C

(condition i.) or 23°C (condition ii.) and stopped by addition of 0.4 U per applied unit of

RNase I of SUPERase•In RNase Inhibitor (Invitrogen). For condition iii., 1 U of Ambion

RNase I (Invitrogen) per µg RNA and 0.134 µL per µL lysate of TURBO DNase (Invitrogen)

was directly added to the lysate and incubated for 1 h at 4°C. The digest was stopped by

addition of 0.4 U of SUPERase•In RNase Inhibitor (Invitrogen) per applied unit of RNase

I. The lysate was centrifuged again for 10 min at 8,000 g and 4°C to remove any cell debris

that may have precipitated during the nuclease digest. For all conditions, monosomes

were pelleted through a 750 µL 30% sucrose cushion prepared in TKM+ buffer

supplemented with 1 mM DTT for 30 min in a S150AT rotor at 72,000 rpm and 4°C. The

resulting ribosome pellets were resuspended in 100 µL of ice-cold ribosome buffer, then

supplemented with 15 mM ethylene diamine tetra-acetic acid (EDTA), pH 8.0 and instantly

mixed with 750 µL Invitrogen TRIzol reagent (Invitrogen) and incubated at room

temperature for 5 min. The solution was mixed with 150 µL chloroform and incubated for

2 min at room temperature before 15 min centrifugation at 20,000 g and 4°C. The

supernatant was mixed with 3 µL GlycoBlue co-precipitant (Invitrogen), 1:10 volume of 3

M sodium acetate, pH 5.5, 1 volume isopropanol and incubated overnight at -20°C. The

precipitated RNA was pelleted by 1 h centrifugation at 20,000 g and 4°C, briefly washed

with ice-cold 80% ethanol and centrifuged again for 30 min at the same parameters. The

resulting RNA pellet was briefly dried and dissolved in 11 µL of nuclease-free water.

For RNA-seq samples, 10 mL of culture were harvested by 2 min centrifugation at

4,000 g at room temperature (before harvesting the culture for RPF isolation), shock-

freezing in liquid nitrogen and storage at -80°C. Samples were lysed in prewarmed (50°C)

lysis buffer (50 mM Tris-HCl pH 8.0, 200 mM NaCl, 20 mM EDTA pH 8.0 and 2% SDS)

and incubated for 2 min at 50°C. Immediately after incubation, 500 µL of TRIzol

(Invitrogen) reagent were added and the mixture was incubated for 5 min at room

temperature. The suspension was mixed with 200 µL chloroform and incubated for

another 5 min at room temperature before 3 min centrifugation at 12,000 g and room

temperature. The nucleic acid containing phase was transferred to a fresh tube and mixed

with 1.5 volumes of ethanol. RNA was purified by the Macherey Nagel NucleoSpin RNA

Plant kit.

Ribosome protected fragment extraction and rRNA depletion

50 µg RNA from the ribosomal extractions were diluted to a volume of 20 µL with nuclease-

free water, mixed with 2x formamide RNA loading buffer (90% deionized formamide, 20

mM Tris-HCl, pH 7.5, 0.5 M EDTA, 0.04% bromophenol blue in ethanol) and denatured

for 90 s at 80°C. The samples were instantly put on ice and then loaded onto 12% urea-

TBE-gels (90 mM Tris, 9 mM boric acid, 2 mM EDTA, 12% 19:1 40% acrylamide-

bisacrylamide mix, 8 M urea). Gels were run at 200 V for approximately 1 h and incubated

with SYBR Gold nucleic acid stain (Invitrogen), diluted 1:10,000 in TBE buffer (90 mM

Tris, 9 mM boric acid, 2 mM EDTA). RPF were excised from the gels in a size range from

20 – 40 nt according to a RNA size marker. Gel pieces were crushed and incubated

overnight with 400 µL RPF elution buffer (300 mM sodium acetate, pH 5.5, 0.25% sodium

dodecyl sulfate and 1 mM EDTA, pH 8.0) at room temperature. The eluate was collected,

mixed with 3 µL GlycoBlue co-precipitant, 1 volume of isopropanol and incubated

overnight at -20°C. The precipitated RNA was pelleted by 1 h centrifugation at 20,000 g

and 4°C, briefly washed with ice-cold 70% ethanol and centrifuged again for 30 min at the

same parameters. The resulting RNA pellet was briefly dried and dissolved in 20 µL of

nuclease-free water. For Ribo-seq samples, abundant rRNA fragments were depleted by

incubating 100 - 200 ng of ribosome RPFs with 1.19 µL of 100 µM biotinylated rRNA-oligo

depletion mix (consisting of the oligonucleotides:

AATATGCGTTCAAAGATTCGATGATTCACG,

TAGCTCTAGAATTACTACGGTTATCCGAGTA,

TACCCGACGCTGAGGCAGACATGCTCTTGG,

GATTCGTGAAGTTATCATGATTCACCGCA,

ACGGGATGAATCTCAGTGGATCGTAGCA,

CGATCTAGCCGTCTTAGAGCTAGAAGCAGG) with 4 µL formamide, 1 µL 20x

concentrated hybridization buffer (300 mM sodium citrate, 3 M NaCl), 2 µL 0.5 M EDTA,

pH 8.0 and in a total volume of 20 µL. The mixture was hybridized in a thermocycler by

heating up to 80°C for 5 min and subsequent cooling in 5°C steps for 2 min each until

35°C was reached in the device. Afterwards, 60 µL of hybridization buffer, supplemented

with 20% formamide was added to the tube and the sample was depleted twice for 15 min

using 15 µL of Dynabeads MyOne Streptavidin T1 magnetic beads (Invitrogen) freshly

washed and prepared according to the manufacturer's manual. The supernatant was

mixed with 3 µL GlycoBlue co-precipitant, 2.5 volumes of ethanol and incubated overnight

at -20°C. RPFs were pelleted by 1 h centrifugation at 20,000 g and 4°C, briefly washed

with ice-cold 70% ethanol and centrifuged again for 10 min at the same parameters. The

resulting pellet was briefly dried and dissolved in 41 µL of nuclease-free water. Excess

oligos were removed by 30 min DNase digest at 37°C upon addition of 5 µL 10x TURBO

DNase buffer, 2 µL Invitrogen TURBO DNase and 2 µL SUPERase•In RNase Inhibitor

(Invitrogen). After digest, RPFs were mixed with 150 µL of 100 mM NaCl and purified

again by phenol-chloroform precipitation and final precipitation overnight at -20°C using

2.5 volumes of ethanol and 2 µL GlycoBlue co-precipitant. RPFs were pelleted by 1 h

centrifugation at 20,000 g and 4°C, briefly washed with ice-cold 70% ethanol and

centrifuged again for 10 min at the same parameters. The resulting pellet was briefly dried

and dissolved in 36.5 µL of nuclease-free water.

For sequencing library preparation, ribosome RPFs were denatured at 65°C for 5 min in

a thermocycler and immediately transferred to ice. The sample was then mixed with 5 µL

10 x T4-polynucleotide kinase buffer A, 2.5 µL T4-polynucleotide kinase (Thermo

Scientific), 2 µL SUPERase•In RNase Inhibitor and incubated for 10 min at 37°C.

Subsequently, 5 µL of 10 mM ATP was added and the sample was incubated for additional

30 min at 37°C. After phosphorylation, the RPFs were mixed with 150 µL 100 mM NaCl

and purified again by phenol-chloroform precipitation and final precipitation overnight at -

20°C using 2.5 volumes of ethanol and 2 µL GlycoBlue co-precipitant. RPFs were pelleted

by 1 h centrifugation at 20,000 g and 4°C, briefly washed with ice-cold 70% ethanol and

centrifuged again for 10 min at the same parameters. The resulting pellet was briefly dried

and dissolved in 11.5 µL of nuclease-free water. The whole sample was used as input for

the NEXTFLEX Small RNA-seq Kit v3 (Perkin Elmer) to generate libraries according to

the kit's manual. For RNA-seq, the purified RNA was later used as input for the Zymo-Seq

RiboFree Total RNA library kit for library preparation according to the manufacturer's

manual.

Processing of raw sequencing data

Demultiplexed FASTQ files were derived from an Illumina NextSeq 550 system aiming for

40 M reads per Ribo-seq sample and 20 M reads per RNA-seq sample at a read length

of 75 nt. Samples were processed through a custom-made pipeline in form of a Linux shell

script calling several bioinformatic tools and custom Python scripts. First operation of the

pipeline was removing 3' adapter sequences using cutadapt for RNA-seq and Ribo-seq,

respectively   (Martin,   2011)   with   parameters   [--minimum-length=9,   -

a="TGGAATTCTCGGGTGCCAAGG"]. After removal of 3' adapter sequences, the 3' and

5' unique molecular identifier tags (UMI tags) were removed from each read's sequence

and shifted to the rear of the header row by UMI tools (Smith et al., 2017) with parameters

[extract --bc-pattern=NNNN] for 5' UMI tags and [extract --bc-pattern=NNNN –3prime] for

3' UMI tags. Remaining reads were filtered for sequence length using cutadapt again with

parameters [--minimum length=20 --maximum-length=39] to remove any read that is

unlikely to represent a real RPF. After size filtering, a custom-made Python script

corrected the UMI-tags that were previously shifted to the read headers in the form

[header line_NNNN_NNNN] to [header line_NNNNNNNN], allowing UMI tools in a later

step to deduplicate the data set based on both UMI tags. In a first mapping step, reads

were aligned to a set of *C. reinhardtii* non-coding RNA FASTA sequences provided by

ENSEMBL plants together with the genome version 5.5 to remove contaminating RNA

species using STAR with parameters [--runMode alignReads --outSAMtype BAM

SortedByCoordinate   --outFilterMultimapNmax   20   --outFilterMismatchNmax   3   --

outReadsUnmapped Fastx --alignIntronMax 7000 --twopassMode Basic]. Reads that did

not align to the ncRNA set were copied to a new FASTQ file that was mapped against the

C. reinhardtii genome version 6.1 using STAR (Dobin et al., 2013) with parameters [--

runMode   alignReads   –outSAMtype   BAM

SortedByCoordinate --outFilterMismatchNoverLmax 0.1 --outReadsUnmapped Fastx --

alignIntronMax 8000 –outSAMmultNmax 1 –outMultimapperOrder Random]. After

genome mapping, the resulting BAM file was deduplicated using UMI-tools' dedup

function and afterwards indexed using samtools' index function (Li et al., 2009).

Data processing

All analyses on the NGS data sets including calculations of gene body coverage,

periodicity, P-site offsets, organellar read length distributions, ribosome profiles and gene

expression metrics were performed using a custom-made object-oriented Python module

that makes extensive use of the packages pysam/HTSlib (https://github.com/pysam-

developers/pysam), pandas (McKinney, 2010), NumPy (Harris et al., 2020), SciPy

(Virtanen et al., 2020), matplotlib (Hunter, 2007) and seaborn (Waskom, 2021) among

multiple others. For RPKM calculations in Ribo-seq data sets, only reads mapping to the

annotated CDS of a transcript were considered. Alternatively we calculate counts per

million values for the interested reader (Supplemental Dataset). For any calculations on

RNA-seq data sets, reads mapping to the whole transcript were considered and the full

length of the mature transcript was used for RPKM calculations. Translation efficiency

values were simply calculated by dividing an CDSs RPKM value from a Ribo-seq data set

by the respective transcripts RPKM value from the affiliated RNA-seq data set.

For the 5'-P-site / 3'-A-site offset calculations, all reads fully enclosing either an

annotated start codon (5'-P-site offset) or an annotated stop codon (for 3'-A-site offset)

were taken into consideration separately for the nuclear and the chloroplast genome.

These reads were sorted according to the length of their alignment to the genome, which

was considered as read length to avoid miscalculations due to clipped or masked

sequences. For every read length species, the position of the reads 5' end or 3' end

relative to the position of the first base of the start codon (for 5'-P-site offsets) or the last

base of the stop codon (for 3'-A-site offsets) was determined as the offset. Afterwards, the

most frequently occurring offset for a read length species was determined to be the true

offset of this species and was used in the P-site calculation. To display the results of the

offset calculations, the count matrix of both 5'-P-site offsets and 3'-A-site offsets were

normalized separately to the total number of counts each and plotted as heatmap.

For calculation of the P-sites first position of reads mapping to nucleus-encoded

CDSs, all reads mapping to a CDS were filtered from the data set and for every read, the

5'-most mapping position of the read's alignment + the 5'-P-site offset determined for the

reads alignment length was determined to be the first P-site position in case of transcripts

encoded on the + strand. For reads encoded on the minus strand, the first P-site position

was determined to be the 3'-most mapping position of the RPF alignment minus the

determined 5'-P-site offset. In all cases, splicing was considered and whenever a spliced

read was found, the calculation was adjusted according to the number of genomic

positions spanned by the splice junction of the read. For reads mapping to chloroplast-

encoded transcripts on the + strand, the first P-site position was determined to be the 3'-

most mapping position – (3'-A-site offset + 5). For reads mapping to chloroplast-encoded

transcripts on the - strand, the first P-site position was calculated to be the 5'-most

mapping position + (3'-A-site offset + 5). The metagene P-site profiles of nucleus-encoded

transcripts were calculated considering only the 2,000 most expressed transcripts of a

dataset, according to their RPKM values. If multiple splice variants of the same gene

occurred in the list, only the longest variant (annotated as transcript 1 in the genome) was

considered for the calculation to avoid analyzing the same reads multiple times. The

profiles were constructed by calculating the first position of the P-site for all reads mapping

to the first 30 and the last 30 codons of considered transcript CDSs. For a ribosome RPF

to be in-frame with the annotated CDS, the first position of the P-site must be located on

the first base of a codon. According to this logic, the calculated P-site positions were

categorized into "in-frame", "+1 shifted" and "+2 shifted" and summed up for all codons of

considered transcript CDSs. To calculate the frame preference per codon, the counts in

each category in a codon were divided by the sum of all categories for every codon

separately. For calculation of the chloroplast metagene P-site profile, all protein-coding

chloroplast transcripts were considered.

The extension of gene body coverage versus a respective cut-off was calculated

as the portion of an CDS that was covered by the number of reads equal to or greater

than the respective cut-off. This procedure was applied to all annotated CDSs using

increasing cut-offs from 1 to 10 reads to monitor the stepwise decrease of gene-body

coverage. For every applied cut-off, the CDSs were categorized as being covered over

>10% of their length to >90% of their length in 10% steps and CDSs falling into every

category were counted and displayed in a heat map to estimate both completeness and

strength of CDS coverage in the data set.

To calculate the read length distribution, all reads mapping to the respective

genome were filtered and the length of their alignment to the genome was taken as read

length. After counting the number of all possible read lengths for each genome separately,

these numbers were normalized by the total number of reads mapping to the respective

genome. For read biotype distribution, the reads were further categorized if they map to

an annotated coding sequence, 5'- or 3'-UTR or intergenic region. To handle reads

overlapping two regions of different category, for example 5'-UTR and coding sequence

as seen regularly in initiation peaks, the read was assigned the category that had the

larger share on the read's alignment. In rare cases, when a read mapped to different

categories in two different splice variants of the same transcript, the read was assigned to

both categories.

Supplemental data

Supplemental Figure S1: Features of ribosome protected fragment coverage.

Supplemental Figure S2: RPF length distribution and periodicity.

Supplemental Figure S3: Transcript accumulation, translation output and translation

efficiency of nuclear genes.

Supplemental Figure S4: Correlation between nuclease digestion conditions, transcript

accumulation and translation.

Supplemental Figure S5: Correlation analysis of RPF count between nuclease digest

conditions.

Supplemental Figure S6: Detection of abnormal initiation peaks in the three genomes of

Chlamydomonas reinhardtii.

Supplemental Dataset. List of accession numbers, gene body coverage, RPKM and

CPM of Ribo-seq and RNA-seq experiments.

Acknowledgements

We thank Fabian Ries, Claudia Herkt and Peter Emelin for critical discussion of the data

and the manuscript. Martin Simon, Jaro Schmitt and Günter Kramer for help with next

generation sequencing. We further thank Jeremy Phillips and David Goodstein from the

Joint Genome Institute for making our data available for the public on the Phytozome

website.

Author contribution:

V.L.G. performed ribosome profiling experiments, bioinformatic data analyses and wrote

parts of the manuscript. M.K.Y.T. and R.Z. helped with setting up ribosome profiling, N.H.

and S.R. conducted correlation analyses. F.W. designed experiments, analyzed data, and

wrote the manuscript.

Funding

This work was supported by the Deutsche Forschungsgemeinschaft grant 437345987 (WI

3477/3-1, ZO 302/5-1, RU 2266/2-1 to F.W., R.Z. and S.R., respectively) and the

Forschungsschwerpunkt BioComp to F.W.

Data availability

All gene accession codes are given in the Supplemental Dataset and are in accordance

with the latest Chlamydomonas genome v6.1 annotation (https://phytozome-

next.jgi.doe.gov) or the GenBank/EMBL data libraries. The full Ribo-seq and RNA-seq

data are accessible via the Dataplant consortium.

Competing interests

The authors declare no competing interests.

References

Aird D, Ross MG, Chen WS, Danielsson M, Fennell T, Russ C, Jaffe DB, Nusbaum
C, Gnirke A (2011) Analyzing and minimizing PCR amplification bias in Illumina
sequencing libraries. Genome Biol **12:** R18
Blifernez-Klassen O, Berger H, Mittmann BGK, Klassen V, Schelletter L, Buchholz
T, Baier T, Soleimani M, Wobbe L, Kruse O (2021) A gene regulatory network
for antenna size control in carbon dioxide-deprived *Chlamydomonas reinhardtii*
cells. Plant Cell **33:** 1303-1318
Bonente G, Pippa S, Castellano S, Bassi R, Ballottari M (2012) Acclimation of
Chlamydomonas reinhardtii to different growth irradiances. J Biol Chem **287:** 5833-
5847
Brar GA, Weissman JS (2015) Ribosome profiling reveals the what, when, where and
how of protein synthesis. Nat Rev Mol Cell Biol **16:** 651-664
Calviello L, Mukherjee N, Wyler E, Zauber H, Hirsekorn A, Selbach M, Landthaler M,
Obermayer B, Ohler U (2016) Detecting actively translated open reading frames
in ribosome profiling data. Nat Methods **13:** 165-170

837 **Cao X, Slavoff SA** (2020) Non-AUG start codons: Expanding and regulating the small
838   and alternative ORFeome. Exp Cell Res **391:** 111973
839 **Chartron JW, Hunt KC, Frydman J** (2016) Cotranslational signal-independent SRP
840   preloading during membrane targeting. Nature **536:** 224-228
841 **Chen H, Alonso JM, Stepanova AN** (2022) A Ribo-Seq Method to Study Genome-Wide
842   Translational Regulation in Plants. Methods Mol Biol **2494:** 61-98
843 **Chiu CW, Li YR, Lin CY, Yeh HH, Liu MJ** (2022) Translation initiation landscape profiling
844   reveals hidden open-reading frames required for the pathogenesis of tomato yellow
845   leaf curl Thailand virus. Plant Cell **34:** 1804-1821
846 **Chotewutmontri P, Barkan A** (2016) Dynamics of chloroplast translation during
847   chloroplast differentiation in maize. PLoS Genet **12:** e1006106
848 **Chotewutmontri P, Barkan A** (2018) Multilevel effects of light on ribosome dynamics in
849   chloroplasts program genome-wide and *psbA*-specific changes in translation.
850   PLoS Genet **14:** e1007555
851 **Chotewutmontri P, Barkan A** (2020) Light-induced *psbA* translation in plants is triggered
852   by photosystem II damage via an assembly-linked autoregulatory circuit. Proc Natl
853   Acad Sci U S A **117:** 21775-21784
854 **Chung BY, Hardcastle TJ, Jones JD, Irigoyen N, Firth AE, Baulcombe DC, Brierley I**
855   (2015) The use of duplex-specific nuclease in ribosome profiling and a user-friendly
856   software package for Ribo-seq data analysis. RNA **21:** 1731-1745
857 **Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A,**
858   **Szczesniak MW, Gaffney DJ, Elo LL, Zhang X, Mortazavi A** (2016) A survey of
859   best practices for RNA-seq data analysis. Genome Biol **17:** 13
860 **Costanzo MC, Hogan JD, Cusick ME, Davis BP, Fancher AM, Hodges PE, Kondu P,**
861   **Lengieza C, Lew-Smith JE, Lingner C, Roberg-Perez KJ, Tillberg M, Brooks**
862   **JE, Garrels JI** (2000) The yeast proteome database (YPD) and *Caenorhabditis*
863   *elegans* proteome database (WormPD): comprehensive resources for the
864   organization and comparison of model organism protein information. Nucleic Acids
865   Res **28:** 73-76
866 **Craig RJ, Gallaher SD, Shu S, Salome P, Jenkins JW, Blaby-Haas CE, Purvine SO,**
867   **O'Donnell S, Barry K, Grimwood J, Strenkert D, Kropat J, Daum C, Yoshinaga**
868   **Y, Goodstein DM, Vallon O, Schmutz J, Merchant SS** (2022) The
869   Chlamydomonas Genome Project, version 6: reference assemblies for mating type
870   *plus* and *minus* strains reveal extensive structural mutation in the laboratory. Plant
871   Cell
872 **Cross FR** (2015) Tying Down Loose Ends in the *Chlamydomonas* Genome: Functional
873   Significance of Abundant Upstream Open Reading Frames. G3 (Bethesda) **6:** 435-
874   446
875 **Dall'Osto L, Bressan M, Bassi R** (2015) Biogenesis of light harvesting proteins. Biochim
876   Biophys Acta **1847:** 861-871
877 **Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson**
878   **M, Gingeras TR** (2013) STAR: ultrafast universal RNA-seq aligner. Bioinformatics
879   **29:** 15-21
880 **Döring K, Ahmed N, Riemer T, Suresh HG, Vainshtein Y, Habich M, Riemer J, Mayer**
881   **MP, O'Brien EP, Kramer G, Bukau B** (2017) Profiling Ssb-Nascent Chain
882   Interactions Reveals Principles of Hsp70-Assisted Folding. Cell **170:** 298-311 e220
883 **Eberhard S, Drapier D, Wollman FA** (2002) Searching limiting steps in the expression
884   of chloroplast-encoded proteins: relations between gene copy number,

transcription, transcript abundance and translation rate in the chloroplast of
Chlamydomonas reinhardtii. The Plant journal : for cell and molecular biology **31:**
149-160

Floris M, Bassi R, Robaglia C, Alboresi A, Lanet E (2013) Post-transcriptional control
of light-harvesting genes expression under light stress. Plant Mol Biol **82:** 147-154

Forsythe ES, Grover CE, Miller ER, Conover JL, Arick MA, 2nd, Chavarro MCF, Leal-
Bertioli SCM, Peterson DG, Sharbrough J, Wendel JF, Sloan DB (2022)
Organellar transcripts dominate the cellular mRNA pool across plants of varying
ploidy levels. Proc Natl Acad Sci U S A **119:** e2204187119

Frank J, Agrawal RK (2000) A ratchet-like inter-subunit reorganization of the ribosome
during translocation. Nature **406:** 318-322

Fu Y, Wu PH, Beane T, Zamore PD, Weng Z (2018) Elimination of PCR duplicates in
RNA-seq and small RNA-seq using unique molecular identifiers. BMC Genomics
19: 531

Fujita T, Kurihara Y, Iwasaki S (2019) The Plant Translatome Surveyed by Ribosome
Profiling. Plant Cell Physiol **60:** 1917-1926

Gallaher SD, Fitz-Gibbon ST, Strenkert D, Purvine SO, Pellegrini M, Merchant SS
(2018) High-throughput sequencing of the chloroplast and mitochondrion of
Chlamydomonas reinhardtii to generate improved de novo assemblies, analyze
expression patterns and transcript speciation, and evaluate diversity among
laboratory strains and wild isolates. Plant J **93:** 545-565

Gao Y, Thiele W, Saleh O, Scossa F, Arabi F, Zhang H, Sampathkumar A, Kuhn K,
Fernie A, Bock R, Schottler MA, Zoschke R (2022) Chloroplast translational
regulation uncovers nonessential photosynthesis genes as key players in plant cold
acclimation. Plant Cell **34:** 2056-2079

Gawronski P, Jensen PE, Karpinski S, Leister D, Scharff LB (2018) Plastid ribosome
pausing is induced by multiple features and is linked to protein complex assembly.
Plant Physiol

Gerashchenko MV, Gladyshev VN (2014) Translation inhibitors cause abnormalities in
ribosome profiling experiments. Nucleic Acids Res **42:** e134

Gerashchenko MV, Gladyshev VN (2017) Ribonuclease selection for ribosome profiling.
Nucleic Acids Res **45:** e6

Gloge F, Becker AH, Kramer G, Bukau B (2014) Co-translational mechanisms of protein
maturation. Curr Opin Struct Biol **24:** 24-33

Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D,
Wieser E, Taylor J, Berg S, Smith NJ, Kern R, Picus M, Hoyer S, van Kerkwijk
MH, Brett M, Haldane A, Del Rio JF, Wiebe M, Peterson P, Gerard-Marchant
P, Sheppard K, Reddy T, Weckesser W, Abbasi H, Gohlke C, Oliphant TE
(2020) Array programming with NumPy. Nature **585:** 357-362

Hinnebusch AG (2011) Molecular mechanism of scanning and start codon selection in
eukaryotes. Microbiol Mol Biol Rev **75:** 434-467, first page of table of contents

Hsu PY, Calviello L, Wu HL, Li FW, Rothfels CJ, Ohler U, Benfey PN (2016) Super-
resolution ribosome profiling reveals unannotated translation events in
Arabidopsis. Proc Natl Acad Sci U S A

Hunter J (2007) Matplotlib: A 2D Graphics Environment. Computing in Science &
Engineering **9:** 90-95

Ingolia NT (2014) Ribosome profiling: new views of translation, from single codons to
genome scale. Nat Rev Genet **15:** 205-213

Ingolia NT (2016) Ribosome Footprint Profiling of Translation throughout the Genome.
Cell **165:** 22-33
Ingolia NT, Brar GA, Rouskin S, McGeachy AM, Weissman JS (2012) The ribosome
profiling strategy for monitoring translation in vivo by deep sequencing of ribosome-
protected mRNA fragments. Nat Protoc **7:** 1534-1550
Ingolia NT, Ghaemmaghami S, Newman JR, Weissman JS (2009) Genome-wide
analysis in vivo of translation with nucleotide resolution using ribosome profiling.
Science **324:** 218-223
Ingolia NT, Lareau LF, Weissman JS (2011) Ribosome profiling of mouse embryonic
stem cells reveals the complexity and dynamics of mammalian proteomes. Cell
147: 789-802
Juntawong P, Girke T, Bazin J, Bailey-Serres J (2014) Translational dynamics revealed
by genome-wide profiling of ribosome footprints in *Arabidopsis*. Proc Natl Acad Sci
U S A **111:** E203-212
Klimmek F, Sjodin A, Noutsos C, Leister D, Jansson S (2006) Abundantly and rarely
expressed *Lhc* protein genes exhibit distinct regulation patterns in plants. Plant
Physiol **140:** 793-804
Komine Y, Kwong L, Anguera MC, Schuster G, Stern DB (2000) Polyadenylation of
three classes of chloroplast RNA in *Chlamydomonas reinhadtii*. RNA **6:** 598-607
Kramer G, Shiber A, Bukau B (2019) Mechanisms of Cotranslational Maturation of
Newly Synthesized Proteins. Annu Rev Biochem **88:** 337-364
Kropat J, Hong-Hermesdorf A, Casero D, Ent P, Castruita M, Pellegrini M, Merchant
SS, Malasarn D (2011) A revised mineral nutrient supplement increases biomass
and growth rate in *Chlamydomonas reinhardtii*. Plant J **66:** 770-780
Lareau LF, Hite DH, Hogan GJ, Brown PO (2014) Distinct stages of the translation
elongation cycle revealed by sequencing ribosome-protected mRNA fragments.
Elife **3:** e01257
Lauria F, Tebaldi T, Bernabo P, Groen EJN, Gillingwater TH, Viero G (2018) riboWaltz:
Optimization of ribosome P-site positioning in ribosome profiling data. PLoS
Comput Biol **14:** e1006169
Lei L, Shi J, Chen J, Zhang M, Sun S, Xie S, Li X, Zeng B, Peng L, Hauck A, Zhao H,
Song W, Fan Z, Lai J (2015) Ribosome profiling reveals dynamic translational
landscape in maize seedlings under drought stress. Plant J **84:** 1206-1218
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R, Genome Project Data Processing S (2009) The Sequence
Alignment/Map format and SAMtools. Bioinformatics **25:** 2078-2079
Liu MJ, Wu SH, Wu JF, Lin WD, Wu YC, Tsai TY, Tsai HL, Wu SH (2013) Translational
landscape of photomorphogenic *Arabidopsis*. Plant Cell **25:** 3699-3710
Lukoszek R, Feist P, Ignatova Z (2016) Insights into the adaptive response of
Arabidopsis thaliana to prolonged thermal stress by ribosomal profiling and RNA-
Seq. BMC Plant Biol **16:** 221
Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing
reads. 2011 **17:** 3
McKinney W (2010) Data Structures for Statistical Computing in Python. *In* Proceedings
of the 9th Python in Science Conference, pp 56-61
Merchante C, Brumos J, Yun J, Hu Q, Spencer KR, Enriquez P, Binder BM, Heber S,
Stepanova AN, Alonso JM (2015) Gene-specific translation regulation mediated
by the hormone-signaling molecule EIN2. Cell **163:** 684-697

Moulin M, Nguyen GT, Scaife MA, Smith AG, Fitzpatrick TB (2013) Analysis of
Chlamydomonas thiamin metabolism in vivo reveals riboswitch plasticity. Proc Natl
Acad Sci U S A **110:** 14622-14627
Mussgnug JH, Wobbe L, Elles I, Claus C, Hamilton M, Fink A, Kahmann U,
Kapazoglou A, Mullineaux CW, Hippler M, Nickelsen J, Nixon PJ, Kruse O
(2005) NAB1 is an RNA binding protein involved in the light-regulated differential
expression of the light-harvesting antenna of *Chlamydomonas reinhardtii*. Plant
Cell **17:** 3409-3421
Nickelsen J, Bohne A-V, Westhoff P (2014) Chloroplast gene expression - translation.
49-78
Payne SH (2015) The utility of protein and mRNA correlation. Trends Biochem Sci **40:** 1-
3
Pechmann S, Willmund F, Frydman J (2013) The ribosome as a hub for protein quality
control. Mol Cell **49:** 411-421
Plancke C, Vigeolas H, Hohner R, Roberty S, Emonds-Alt B, Larosa V, Willamme R,
Duby F, Onga Dhali D, Thonart P, Hiligsmann S, Franck F, Eppe G, Cardol P,
Hippler M, Remacle C (2014) Lack of isocitrate lyase in *Chlamydomonas* leads to
changes in carbon metabolism and in the response to oxidative stress under
mixotrophic growth. Plant J **77:** 404-417
Rooijers K, Loayza-Puch F, Nijtmans LG, Agami R (2013) Ribosome profiling reveals
features of normal and disease-associated mitochondrial translation. Nat Commun
4: 2886
Russell JB, Cook GM (1995) Energetics of bacterial growth: balance of anabolic and
catabolic reactions. Microbiol Rev **59:** 48-62
Schneider-Poetsch T, Ju J, Eyler DE, Dang Y, Bhat S, Merrick WC, Green R, Shen
B, Liu JO (2010) Inhibition of eukaryotic translation elongation by cycloheximide
and lactimidomycin. Nat Chem Biol **6:** 209-217
Schroda M, Hemme D, Mühlhaus T (2015) The *Chlamydomonas* heat stress response.
Plant J **82:** 466-480
Schuster M, Gao Y, Schöttler MA, Bock R, Zoschke R (2019) Limited Responsiveness
of Chloroplast Gene Expression during Acclimation to High Light in Tobacco. Plant
Physiol
Sharma P, Wu J, Nilges BS, Leidel SA (2021) Humans and other commonly used model
organisms are resistant to cycloheximide-mediated biases in ribosome profiling
experiments. Nat Commun **12:** 5094
Slobodin B, Dikstein R (2020) So close, no matter how far: multiple paths connecting
transcription to mRNA translation in eukaryotes. EMBO Rep **21:** e50799
Smith T, Heger A, Sudbery I (2017) UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. Genome Res **27:** 491-499
Sun Y, Zerges W (2015) Translational regulation in chloroplasts for development and
homeostasis. Biochim Biophys Acta **1847:** 809-820
Teixeira FK, Lehmann R (2019) Translational Control during Developmental Transitions.
Cold Spring Harb Perspect Biol **11**
Ting MKY, Gao Y, Barahimipour R, Ghandour R, Liu J, Martinez-Seidel F, Smirnova
J, Gotsmann VL, Fischer A, Haydon MJ, Willmund F, Zoschke R (2023)
Improved Plant Ribosome Profiling with Structural Assessment of rRNA
Contaminants. manuscript submitted

Trösch R, Barahimipour R, Gao Y, Badillo-Corona JA, Gotsmann VL, Zimmer D,
Mühlhaus T, Zoschke R, Willmund F (2018) Commonalities and differences of
chloroplast translation in a green alga and land plants. Nat Plants **4:** 564-575
Trösch R, Ries F, Westrich LD, Gao Y, Herkt C, Hoppstädter J, Heck-Roth J, Mustas
M, Scheuring D, Choquet Y, Räschle M, Zoschke R, Willmund F (2022) Fast
and global reorganization of the chloroplast protein biogenesis network during heat
acclimation. Plant Cell **34:** 1075-1099
Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D,
Burovski E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M,
Wilson J, Millman KJ, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E,
Carey CJ, Polat I, Feng Y, Moore EW, VanderPlas J, Laxalde D, Perktold J,
Cimrman R, Henriksen I, Quintero EA, Harris CR, Archibald AM, Ribeiro AH,
Pedregosa F, van Mulbregt P, SciPy C (2020) SciPy 1.0: fundamental algorithms
for scientific computing in Python. Nat Methods **17:** 261-272
Wang F, Dischinger K, Westrich LD, Meindl I, Egidi F, Trosch R, Sommer F, Johnson
X, Schroda M, Nickelsen J, Willmund F, Vallon O, Bohne AV (2023) ONE-
HELIX PROTEIN 2 is not required for the synthesis of photosystem II subunit D1
in Chlamydomonas. Plant Physiol
Wang R, Amoyel M (2022) mRNA Translation Is Dynamically Regulated to Instruct Stem
Cell Fate. Front Mol Biosci **9:** 863885
Waskom M (2021) Seaborn: statistical data visualization. Journal of Open Source
Software **6:** 3021
Westrich LD, Gotsmann VL, Herkt C, Ries F, Kazek T, Trosch R, Armbruster L,
Mühlenbeck JS, Ramundo S, Nickelsen J, Finkemeier I, Wirtz M, Storchova Z,
Räschle M, Willmund F (2021) The versatile interactome of chloroplast ribosomes
revealed by affinity purification mass spectrometry. Nucleic Acids Res **49:** 400-415
Wolin SL, Walter P (1988) Ribosome pausing and stacking during translation of a
eukaryotic mRNA. EMBO J **7:** 3559-3569
Wu CC, Zinshteyn B, Wehner KA, Green R (2019) High-Resolution Ribosome Profiling
Defines Discrete Ribosome Elongation States and Translational Regulation during
Cellular Stress. Mol Cell **73:** 959-970 e955
Wu HL, Song G, Walley JW, Hsu PY (2019) The Tomato Translational Landscape
Revealed by Transcriptome Assembly and Ribosome Profiling. Plant Physiol **181:**
367-380
Yang X, Cui J, Song B, Yu Y, Mo B, Liu L (2020) Construction of High-Quality Rice
Ribosome Footprint Library. Front Plant Sci **11:** 572237
Yang X, Song B, Cui J, Wang L, Wang S, Luo L, Gao L, Mo B, Yu Y, Liu L (2021)
Comparative ribosome profiling reveals distinct translational landscapes of salt-
sensitive and -tolerant rice. BMC Genomics **22:** 612
Zhang W, Dunkle JA, Cate JH (2009) Structures of the ribosome in intermediate states
of ratcheting. Science **325:** 1014-1017
Zoschke R, Bock R (2018) Chloroplast Translation: Structural and Functional
Organization, Operational Control and Regulation. Plant Cell **30:** 745-770
1071

Figure 1: Ribo-seq provides a broad coverage of expressed genes

(A) Illustration of the experimental setup, testing nucleolytic conditions. Polysome purification and subsequent RNase I digest for 1h at 4°C (i.) or 1h at 23°C (ii.) was performed from Chlamydomonas samples, respectively. Alternatively, cell lysate was directly supplemented with RNase I (iii.). (B) Scatter plots representing the reproducibility of Ribo-seq (left panels) and RNA-seq experiments (right panels). Shown are the R-values and RPKM values. (C) Schematic representation of the concept of gene body coverage (left panel: incomplete coverage that strongly declines with an increasing read cut-off; right panel: good coverage that is robust against increasing read cut-offs). (D) and (E) Cumulative heatmaps visualizing the number of genes (given in the respective panel) covered to a certain minimal extent (rows) by a certain minimal number of reads (columns) for ribosome RPFs (C) and transcripts (D). Ribo-seq data shown are derived from the RNase I digest in lysate (condition iii.). The respective data for the other treatments are shown in Supplemental Figure 1.

Figure 2: Tri-nucleotide periodicity of cellular translation

(A) Length distribution of RPFs for each organelle from the three different nuclease treatments tested (i. and ii.: digest of purified ribosomes at 4°C and 23 °C, respectively, iii.: digest in lysate at 4°C). (B) and (C) metagene relative frame preference of nuclear and chloroplast-encoded transcripts per codon for all three nuclease treatments. Codons 1-30 represent the first 30 codons of each transcript while codons 31-60 represent the last 30 codons of each transcript. Frame 0 corresponds to AUG in the ribosomal P-site. (D) Left side: cartoon of the ribosome with small (SSU) and large (LSU) subunits. Heatmaps representing normalized 5'-P-site offset and 3'-A-site offset counts for nucleus- and chloroplast-mapped RPFs that fully enclose a start or stop codon. Columns represent 5'/3'-RPF end positions relative to the start or stop codon while rows represent the RPF lengths. The shades of red are proportional to the number of RPF ends mapping onto a tile. Arrows indicate the triplet-wise movement of ribosomes in early or late elongation state. Asterisks indicate the tiles harboring the most 5'- or 3'-ends of the 21 and 30 nt RPF-species. Dashed boxes indicate the sudden absence of 21 nt RPFs for the last two codons before translation terminates.

Figure 3: Chloroplast-encoded proteins are highly expressed
(A) Violin plots of the RPF data, displaying the RPKM (count per million reads) distribution for genes of each plant cell genome, following the three different RNase I treatments (i. and ii.: digest of purified ribosomes at 4°C and 23°C, respectively, iii.: digest in lysate at 4°C). Values with a RPKM cut-off <2 (log$_2$) are excluded. (B) Violin plots, displaying distribution of RPKM (reads per kilobase of exon per million reads mapped) for RNA values with a cut-off >2 (log$_2$) for each plant cell genome in replicate 3 exemplary. (C) Scatterplots of the averaged Ribo-seq and RNA-seq (RPKM) data for different subcellular-localized proteins. Coloring indicates subcellular localization and origin of the encoded proteins.

Figure 4: Translation output of chloroplast- and nucleus-encoded transcripts
(A) Mean read intensity/TE ratio (per respective coding sequence) of chloroplast-encoded transcripts on the level of RPF, RNA and translation efficiency (TE). The transcripts are grouped by function of the encoded proteins (Cytb_6f= cytochrome b_6f complex, PS I = photosystem I, PS II = photosystem II, polym. = RNA polymerase, chl. synth. = chlorophyll synthesis, misc. = miscellaneous). (B) Values of the 70 most highly expressed nucleus-encoded genes on the translation level. Only the longest splice variants were considered. Respective translation efficiency for these transcripts is given below. For comparison, transcripts with the top 70 highest TE values are given in Supplementary Figure S3C. Error bars indicate standard deviation of the three experiments (treatment i. to iii. for Ribo-seq, and biological replicate 1-3 for RNA-seq), and are shown unidirectional to improve readability.

Figure 5: Putative translational regulation of individual transcripts

(A) Boxplots representing the distribution of Pearson's *r* correlation coefficients comparing the RNA gene body coverage and ribosome profiles of all chloroplast- and nucleus-encoded transcripts between replicates (R1 and R2 for RNA) and between two types of RNase I treatments (ii. and iii. for translation). (B) Average *Pearson* and *Spearman* correlation coefficients depending on the number n of considered transcripts. Transcripts were ordered by descending RPF count per transcript, n most highly covered transcripts were selected, correlation coefficients were computed for each transcript in this subset, and the average was plotted. For correlation coefficients of individual transcripts see Supplemental Figure S5A. (C) Ribosome profiles of LHCBM1 exemplary for a profile with even RPF distribution and LHCBM6 as an example for a profile with an extreme initiation peak. Red parts of the profile represent coverage within the 5'-UTRs, green parts represent 3'-UTRs and alternating blue shades represent exons within the CDS. Black arrows on the x-axis indicate start and stop codons. (D) Stacked column chart representing the fraction of ribosome profiles containing extreme initiation peaks (defined as transcripts with an average coverage in first 7 codons exceeding by at least 4 times the average coverage of the whole CDS). Nucleus-, chloroplast- and mitochondria-encoded transcripts are plotted separately containing only transcripts with a gene body coverage of >70% in the Ribo-seq (transcripts with extreme initiation peak are 3346 out of 8757 for nuclear, 7 out of 66 for chloroplast and 0 out of 7 for mitochondria). Further information on occurrence of abnormal initiation peaks is provided in Supplemental Figure S6.

Figure 6: Chloroplast ribosomes generate characteristic ribosome protected fragments

(A) Size distribution of RPFs that map the CDS of chloroplast-encoded transcripts for nuclease treatment conditions iii. (observed). Equalized size distribution shows the normalized size distribution for each chloroplast transcript, neglecting possible weights from the most abundant transcripts. (B) Heatmap presenting the share of the different RPF sizes for each chloroplast transcript. Transcripts are ranked based on their overall share of RPFs relative to all measured chloroplast RPFs. RMSE (Root Mean Square Error) denotes the deviation of the respective size distribution relative to the average size distribution (top row). (C) Size distribution of chloroplast-encoded transcripts with high deviation relative to the average size distribution. (D) Additional size distribution curves of photosynthesis complex subunits. (E) and (F) Ribosome profile of *psbA* and *atpH,* respectively. Red curve shows the RPF sizes, accumulating within the respective CDS section, black curves indicate RFP coverage per nt. Positions of transmembrane segments (TMS) are given below the graph.

Figure 7: Utilizing ribosome profiles for coding sequence annotation
Exemplary ribosome profiles for determining initiation sites and alternatively spliced transcripts. Red parts of the profiles represent coverage within the 5'-UTRs, green parts represent 3'-UTRs and alternating blue shades represent exons within the CDS. Black arrows on the x-axis indicate start and stop codons. (A) and (B) ribosome profiles of LHCA4 and RPL3 as an example of wrong start codon prediction, respectively. The red arrow indicates the annotated start codon while the black arrows indicate true start and stop codons. (C) Ribosome profiles of PDI2 (Cre01.g033550) transcript variants demonstrating how ribosome profiles can be utilized to identify putative non-cognate start codon initiation and estimate the dominance of transcript variants. Note that AUG was labelled as "true" initiation site, however, an upstream non-cognate start codon might be the correct initiation site.

SUPPLEMENTAL INFORMATION

Utilizing high resolution ribosome profiling for the global investigation of gene expression in *Chlamydomonas*

Vincent Leon Gotsmann[1], Michael Kien Yin Ting[2], Nadin Haase[3], Sophia Rudorf[3], Reimo Zoschke[2], and Felix Willmund[1]*

Author's institution(s)/affiliation(s):

[1] Molecular Genetics of Eukaryotes, University of Kaiserslautern, Paul-Ehrlich-Str. 23, 67663 Kaiserslautern, Germany.

[2] Max Planck Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam-Golm, Germany.

[3] Computational Biology, University of Hannover, Herrenhäuser-Str. 2, 30419 Hannover, Germany.

Supplemental Figures and Legends

Supplemental Figure S1: Features of ribosome protected fragment coverage

Supporting Figure 1. (A) and (B) heatmaps representing transcript numbers categorized by their extent of gene body coverage in dependence of specific read count cut-offs for Ribo-seq data sets i. and ii.

Supplemental Figure S2: RPF length distribution and periodicity

Supporting Figure 2. (A) Length distribution of RPFs for each organelle separated by their match to CDS, 3´- or 5´-UTR or intergenic regions (only data of digest iii. are shown here). (B) metagene RPF accumulation of nuclear and chloroplast-encoded transcripts per codon for all three nuclease treatments. Codons 1-30 represent the first 30 codons of each transcript, whereas codons 31-60 represent the last 30 codons of each transcript. Frame 0 corresponds to AUG in the ribosomal P-site.

Supplemental Figure S3: Transcript accumulation, translation output and translation efficiency of nuclear genes

Supporting Figure 4. (A) Subcellular distribution of the top 100 expressed nucleus-encoded transcripts on the translation level. If multiple transcript variants of the same gene occurred in the list, only the longest occurring variant was considered. (B) Translation output, transcript abundance and translation efficiency of light harvesting protein encoding transcripts across all three Ribo-seq data sets, considering only the longest transcript variant, respectively. Mean values averaging all three Ribo-seq data are plotted, error bars denote standard deviations. (C) Translation efficiency and translation output of the 70 most efficiently translated nucleus-encoded transcripts. Mean values averaging all three Ribo-seq data are plotted, error bars denote standard deviations. If multiple transcript variants of the same gene occurred in the list, only the longest occurring variant was considered, unless stated in the legend.

Supplemental Figure S4: Correlation between nuclease digestion conditions, transcript accumulation and translation

Supporting Figure 5. Boxplots representing the distribution of Pearson correlation coefficients between ribosome profiles and RNA-seq read distribution of the same transcripts each in different data sets. Boxes represent the 0.25 to 0.5 and the 0.5 to 0.75 quartiles, lines represent the median values and whiskers represent the remaining data. Outliers are defined as values being smaller or greater than the respective median ±1.5 times the interquartile range and are depicted as points R1/2/3 indicate different RNA-seq data sets.

Supplemental Figure S5: Correlation analysis of RPF count between nuclease digest conditions

Supporting Figure 5. (A) Relation between correlation coefficient comparing two experimental conditions and average read counts. For each transcript, counts were summed over all positions and then divided by the total number of nucleotides in the transcript. The correlation is high unless the average read counts are low. The distribution of *r*-values might be well suited for selecting good cut-off ranges for a deeper analyses of ribosome profiles. (B) Examples of ribosome profiles with medium and high read counts for all three experimental conditions. Scatter plot of

read counts for each nucleotide for comparison of two experimental conditions. Perfectly identical profiles would correspond to a straight line in the scatter plot.

Supplemental Figure S6: Detection of abnormal initiation peaks in the three genomes of *Chlamydomonas reinhardtii*

Supporting Figure 5. (A) Violin plots representing the distribution of ratios between average per nucleotide coverage within the first seven codons and the average per nucleotide coverage of the remaining CDS for each organelle's transcripts separately. Thick lines within the violins represent the interquartile range while thin lines represent the whiskers with a cut-off of 1.5 times the interquartile range. (B) Scatter plot with lines representing the increase of the percentile borders of the distributions shown in (A). In both cases, only transcripts with a gene-body coverage of at least 70% were considered (8757 nucleus-encoded, 66 chloroplast-encoded, 6 mitochondrial-encoded) to minimize the risk of misinterpretations due to low or absent coverage in either of both regions. Calculation of a 90th-percentile border for mitochondrial transcripts was not possible due to the low number of transcripts in consideration.