Functional convergence of genomic and transcriptomic genetic architecture underlying sociability in a live-bearing fish

Alberto Corral-Lopez*[1,2,3], Natasha I. Bloch[4], Wouter van der Bijl[1], Maria Cortazar-Chinarro[5,6], Alexander Szorkovszky[7], Alexander Kotrschal[8], Iulia Darolti[1,9], Severine D. Buechel[8], Maksym Romenskyy[10], Niclas Kolm[2], Judith E. Mank[1]

Affiliations

- Department of Zoology and Biodiversity Research Centre, University of British Columbia, Vancouver, Canada

- Department of Zoology/Ethology, Stockholm University, Stockholm, Sweden.

- Division of Ecology and Genetics, Uppsala University, Uppsala, Sweden

- Department of Biomedical Engineering, University of Los Andes, Bogota, Colombia

- Department of Earth, Ocean and Atmospheric Sciences, University of British Columbia, Vancouver, Canada

- MEMEG Department of Biology, Lund University, Lund, Sweden.

- RITMO Centre for Interdisciplinary Studies in Rhythm, Time and Motion, University of Oslo, Oslo, Norway

- Behavioural Ecology, Wageningen University & Research, Wageningen, Netherlands.

- Department of Ecology and Evolution, University of Lausanne, Switzerland

- Department of Life Sciences, Imperial College London, London, UK.

*corresponding author: alberto.corral@ebc.uu.se

Abstract

The organization and coordination of fish schools provide a valuable model to investigate the genetic architecture of affiliative behaviors and dissect the mechanisms underlying social behaviors and personalities. We used quantitative genetic methods in replicate guppy selection lines that vary in schooling propensity to investigate the genetic basis of sociability phenotypes. Alignment and attraction, the two major traits forming the sociability personality axis in this species, showed heritability estimates at the upper end of the range previously described for social behaviors, with important variation across sexes. Consistent with findings in collective motion patterns, experimental evolution of schooling propensity increases the sociability of female, but not male, guppies when swimming with unfamiliar conspecifics, a highly relevant trait in a species with fission-fusion social systems in natural populations. Our results from PoolSeq and RNASeq convergently identify genes involved in neuron migration and synaptic function, highlighting a crucial role of glutamatergic synaptic function and calcium-dependent signaling processes in the evolution of schooling.

Introduction

Living in groups, a widespread phenomenon across the animal kingdom, can lead to strikingly complex social behaviors, such as cooperative interactions, subdivision of labor, or collective decision-making[1]. Sociability, the propensity to affiliate with other animals, can also vary across individuals. Sociability represents a fundamental aspect of personality which can influence social interactions and is often subject to strong selective processes[2,3]. Indeed, intra-specific differences in sociability are widespread[e.g 4,5] and often individual genetic variation underlies variability in personality and social behavior phenotypes[6]. However, heritability estimates of social behavior traits are consistent with a complex, polygenic architecture[7]. Human twin and family studies reveal heritability estimates of personality traits generally ≈ 0.40[reviewed in 8]. In non-human animals, a meta-analysis estimated the mean heritability is 0.23 across social behaviors, including personality traits[9], with heritability of affiliative associations ranging substantially, from 0.11 to 0.51[10–13].

Despite this complexity, multiple neural and genetic mechanisms underlying social behavior have been identified[14]. Many of the neural structures and neuromodulators (serotonin, dopamine, vasopressin and oxytocin) are highly conserved within the social-decision making network across vertebrates[15]. Moreover, human personality traits associated with social-decision making have been linked to dopaminergic and serotonergic genes[reviewed in 16], and the regulation of these neuromodulators has been connected to neurodevelopmental disorders that affect affiliative behaviors, such as autism spectrum disorder[6,17,18]. Studies in non-human organisms likewise point towards a major role of genes involved in the regulation of these neurochemical systems. For instance, mouse knock-out mutants for genes involved in dopaminergic signalling exhibit altered sociability phenotypes[19], and changes in sociability in three-spined sticklebacks, *Gasterosteus aculeatus*, are predicted by natural variation in the expression of genes within the dopaminergic and stress pathways[20]. However, while specific groups of genes have been identified for a range of affiliative behaviors, we lack a deeper understanding of their role in inter-individual variation and evolutionary processes underlying sociability.

In fish, group-living often leads to spectacular forms of collective behavior, with members of a school coordinating their movements to increase efficiency in foraging, travelling or predator

avoidance[1]. The extent to which members of a school coordinate their movements is an integral

part of the sociability axis of personality, i.e., how individuals react to the presence or absence of

conspecifics excluding aggressive behaviors[21]. We previously showed that schooling behavior has

a repeatability of 0.43 at the individual level[22], and that can increase substantially over just three

generations of artificial selection in female guppies, *Poecilia reticulata*, generating a 15% increase

in intrinsic schooling propensity compared to controls[22,23]. Selection was based on a group

phenotype, polarization, or the level of alignment between individuals moving together in a group.

Understanding the genetic basis of this schooling phenotype requires linking individual phenotypic

differences to genetic variation. In this study, we phenotyped alignment and attraction of 1496

guppies across 195 families (father, mother, three female and three male offspring from our

replicate experimental selection lines) to estimate the heritability of these two motion

characteristics that previous factor analyses identified to be integral components for the sociability

axis of personality in this species[21]. Because many social interaction patterns in guppies have sex

differences[24–26], and because our selection was performed solely on females, we are able to

examine cross-sex genetic correlations in this ecologically relevant behavioral trait. Genomic and

transcriptomic data from these lines reveal convergence in the genetic architecture of sociability,

highlighting a series of genes with well-defined roles in neurodevelopmental processes. Our results

provide a robust agreement across experiments about the genetic regulation of neural processes in

decision-making and motor control regions of the brain, and its importance for variation of

personality within individuals of this species.

Results

128
129   *Social interactions with unfamiliar individuals and the heritability of sociability in guppies*

130
We first determined whether experimental evolution for higher schooling propensity affected

social interactions with unfamiliar conspecifics. For this, we assessed sociability in 740 females

and 746 males from multiple families of three replicate lines artificially selected for higher

polarization (polarization-selected lines hereon) and three replicate control lines exposed to a

group of non-kin unfamiliar conspecifics in an open field test. Specifically, we quantified their

alignment and nearest neighbor distance (attraction), two measurements of collective motion

characteristics that are demonstrated to capture the most biologically relevant aspects of the
sociability axis of personality in this species[21].

Female guppies from polarization-selected lines presented higher alignment and higher attraction
to an unfamiliar group compared to control lines (Linear Mixed Model for alignment,
$LMM_{alignment}$: line: t = 2.27; df = 9.68; p = 0.047; $LMM_{attraction}$: line: t = -2.34; df = 9.41; p = 0.043;
Fig. 1A; Supplementary Table 1). No differences were observed in these traits between
polarization-selected and control males ($LMM_{alignment}$: line: t = -1.38; df = 9.56; p = 0.20;
$LMM_{attraction}$: line: t = 0.88; df = 9.26; p = 0.40; Fig. 1A; Supplementary Table 2). Our analyses
showed an effect of sex in alignment, with females exhibiting approximately 8% higher alignment
than males ($LMM_{alignment}$: sex: t = -3.02; df = 690.08; p = 0.003), but no difference between sexes
in attraction to a group of unfamiliar conspecifics ($LMM_{attraction}$: sex: t = 0.51; df = 447.05; p =
0.61; Fig. 1A; Supplementary Table 1). There were some differences in sociability between the
parental and offspring generation tested in our experiment, with higher alignment to group average
direction and lower distances to nearest neighbor observed in offspring ($LMM_{alignment}$: generation:
t = -10.13; df = 1141.24; p < 0.001; $LMM_{attraction}$: generation: t = 11.29; df = 992.16; p < 0.001;
Fig. 1A; Supplementary Table 1). Differences in body size between age classes in guppies may
explain these results (see Supplementary Table 3), as the time restrictions involved in testing large
numbers of fish required that we assessed individuals from the parental and offspring generations
at different ages. However, these differences are unlikely to create large biases in our heritability
estimates given that we tested all fish after sexual maturation and that polarization-selected and
control fish were of similar age within parents tested (9 months old) and within offspring tested (5
months-old). In addition, the difference in means between generations is accounted for in our
statistical models (see Methods).

To assess the heritability of sociability in this species, we fitted animal models with alignment and
attraction phenotypes quantified from these 1486 individuals, comprised of parents, three male
and three female offspring for 195 families (99 polarization-selected and 96 control-families).
Given known differences in guppies between the sexes in social interaction patterns[24–26], we
estimated heritability with animal models that only included relationships with same sex

individuals (same-sex pedigree) or that include relationship with individuals from both sexes (full

pedigree).

Using same-sex pedigree animal models, attraction heritability was similar in females ($h^2_{attraction}$,

estimate [95% credible interval] = 0.18 [0.05, 0.34]; Fig. 1B) and males ($h^2_{attraction}$ = 0.19 [0.06,

0.34]; Fig. 1B), however alignment heritability was far higher in females ($h^2_{alignment}$ = 0.34 [0.18,

0.49]; Fig 1B) than males ($h^2_{alignment}$ = 0.06 [0.00, 0.18]; Fig. 1B). Full-pedigree models indicated

lower heritability estimates than those obtained from same-sex pedigree models (Supplementary

Table 5; Fig 1B), except for the heritability estimate of attraction in males ($h^2_{attraction}$ = 0.26 [0.16,

0.37]; Fig 1B). Finally, animal models indicated a positive female-male genetic correlation in

attraction (r_{fm} attraction: 0.68 [0.23, 0.98]; Fig. 1B), although the magnitude of this correlation

contained large credible intervals. For alignment, credible intervals for r_{fm} are also wide and span

zero (r_{fm} alignment: 0.44 [-0.17, 0.95]), and we can only conclude that the cross-sex genetic

correlation is not strongly negative (Supplementary Table 5; Fig 1B).

181
182 *Genetic basis of sociability in guppies*
183
Our quantitative genetic analyses of alignment and attraction suggest an important genetic

influence in sociability phenotypes of guppies. As such, we sequenced DNA pools (Pool-seq) to

identify genome-wide differences in allele frequencies between polarization-selected female

guppies that presented high sociability and polarization-selected females that presented low

sociability. Specifically, we focused on measurements obtained from females in analyses of

alignment to an unfamiliar group. For this, we pooled the DNA from mothers whose family

(normalized mother and daughters alignment score; see methods) were in the top 25% and the

bottom 25% quartiles from each of the three replicated polarization-selected lines (six total pooled

samples with 7 individuals each; Supplementary Figure 1).

DNA reads were aligned to the guppy reference genome (Guppy_female_1.0 + MT, RefSeq

accession: GCA_000633615.2) to compare genome-wide allele frequency differences between

high and low sociability guppies. We ran two independent analyses with these aligned sequences.

For our first analysis, we merged sequences from the three replicates with high sociability pooled

samples and sequences from three replicates with low sociability pooled samples. We filtered

merged sequences to 3,004,974 SNPs (see methods) and performed a Fisher's exact test in
Popoolation2[27] to identify SNPs that significantly differed in their allele frequencies between
guppies with high and low sociability. Using this methodology, we identified 819 SNPs associated
with our sociability phenotype (Fisher's exact test, $p < 10^{-8}$; Fig 2A). SNPs over this standard
genome-wide significance threshold[28] were mostly found in single physically unlinked positions
across the genome, consistent with a polygenic architecture of the trait.
205
Out of these 819 significantly different SNPs, 421 were located within genes or gene promoter
regions of the guppy genome and were used for further functional characterization in association
with Gene Ontology (GO) annotations (273 unique genes). We clustered GO terms based on
semantic similarities and found significant overrepresentation of biological process terms related
to learning and memory, synaptic functioning, response to stimulus, locomotion and growth (Fig.
2B). We likewise found significant overrepresentation of cadherin and calcium-dependent protein
binding annotations (molecular components terms; Supplementary Figure 2) and glutamatergic
synapse annotations (cellular components terms, Supplementary Figure 3).
214
Second, we looked for consistent differences in allele frequencies between high and low sociability
pooled samples in our three replicates by performing the Cochran-Mantel-Haenszel test (cmh-test)
in Popoolation2[27]. Convergent changes in allele frequency likely represent selected sites, and are
less likely the result of genetic drift in any one line. This stringent analysis identified 13 SNPs
from ten different chromosomes with consistent significant differences in allele frequencies across
the three replicates (cmh-test p-value <0.01 with false discovery rate correction). Five of these
SNPs are located within known coding sequence of the guppy genome, of which three are within
well-characterized genes in zebrafish and human homologs with important roles for cognitive
function – ubiquitin specific peptidase 11 (*usp11*), *supt6* histone chaperone and transcription
elongation factor homolog (*supt6h*), and cadherin 13 (*cdh13*; Fig 2A, Table 1). The other two are
classified as novel genes, one of them is matched to an RNA-binding protein *Nova-1-like* gene,
similarly associated with motor function and changes in synaptic function (Fig 2A, Table 1).
227
228 *Neurogenomic response of schooling in guppies*
229

We used transcriptome sequencing to determine differences in gene expression in multiple brain

regions of polarization-selected and control females in response to two different social contexts,

swimming alone (the "Alone" condition) or schooling in a group (groups of eight unfamiliar

females; the "Group" condition). We focused on three separate brain tissues that control distinct

functions. The *optic tectum* is involved in sensory processing of visual signals. The *telencephalon*

is implicated in decision making. The *midbrain* is associated with motor function in response to

auditory and visual stimuli[29,30]. Together, these three brain tissues contain the main components

of the social brain network in fish[31,32].

Differential expression analyses. We identified genes differentially expressed between lines

under each treatment condition and in each brain region separately, to determine the neurogenomic

response triggered by schooling in both lines. Gene expression analyses indicated very little

overlap in differentially expressed (DE) genes between polarization-selected and control lines

(Fig. 3, Supplementary Dataset 1). Specifically, we found that only Adipocyte enhancer-binding

protein 2 gene (*AEBP2;* involved in adipocyte differentiation) in the midbrain and an unknown

gene in the optic tectum were differentially expressed in both polarization-selected and control

lines. Such little overlap suggests that females from different selection lines are activating different

transcriptional cascades and biological pathways in response to social context. In polarization-

selected lines we found an order of magnitude fewer DE genes in the optic tectum than in the other

brain components (n=21 for optic tectum, n=158 for telencephalon, n=109 for midbrain, each p_{adj}

< 0.05). Moreover, in the telencephalon and midbrain, DE genes between Alone and Group

treatment in the polarization-selected lines were enriched for GO annotations associated with

cognition, memory, learning and social behavior (Supplementary Dataset 1). We found enrichment

for these annotation terms for DE genes expressed in the optic tectum, but not in the midbrain or

telencephalon, of control lines.

Hierarchical clustering analyses of DE genes showed that females from polarization-selected lines

in the Group condition clustered uniquely from polarization-selected females in the Alone

condition (Fig. 3, Supplementary Table 6). Similarly, females from polarization-selected lines in

the Group condition clustered uniquely from females from control lines under the Group and Alone

conditions in both the telencephalon and the midbrain, suggesting a unique response in the regions

of the brain associated with behavior to social exposure. This was not observed in samples from
the optic tectum, suggesting that visual processing of social treatments did not differ between
polarization-selected and control females. Hierarchical clustering analyses using all expressed
genes clustered samples by selection line rather than by social context condition (Supplementary
Fig. 4), suggesting that social context affects only a targeted subset of the overall transcriptome
rather than the majority of genes.

Differential coexpression analyses. We used systems biology methods designed to compare the
coexpression networks between conditions to identify genes that change in the way they are
connected to other genes within the coexpression network across conditions, independent of
whether they are differentially expressed[29–31]. Specifically, we used BFDCA[30] to identify
differential coexpressed (DC) gene pairs under the Group and Alone conditions (i.e. pairs of genes
that significantly change in correlation between the two social contexts for each line[30,32]). Similar
to findings in DE analyses, we found little overlap in the genes forming DC gene pairs between
comparisons implemented for control and polarization-selected lines (see Supplementary Tables
7-8, Fig. 3). Together, our results suggest that polarization-selected lines were activating different
biological pathways than control lines to modulate coordinated movement.

We additionally found a group of genes that are both differentially expressed (DE) and
differentially coexpressed (DC) in the same tissue and line, suggesting they might play an
important role mediating coordinated movement (Supplementary Tables 8-9). Specifically, in the
telencephalon, we identified 4 genes that are both DE and DC in polarization-selected lines:
LRRC24, *PTPRS, KHDR2, and PP2BA* (Supplementary Table 9). These genes are part of the
Calcineurin and the Wnt/Oxytocin signalling pathways, known to be involved in modulating social
behavior, learning and memory[33–35]. Enrichment tests confirm the functional relevance of the DC
gene pairs identified, revealing an overrepresentation of genes associated with the glutamatergic
synapse, as well as visual transduction among DC gene pairs in multiple comparisons
(Supplementary Tables 10-11).

Functional characterization of genes of interest across experiments

We combined the information from our genomic and transcriptomic analyses on polarization-
selected and control lines to obtain an intersected delimitation of the gene functions that our
analyses highlighted as important in the development and expression of social interactions with
conspecifics. Specifically, we used functional analyses in the set of genes with differentiated SNPs
between merged sequences of the three replicates with high and low sociability (273 unique genes)
as reference, and compared it to functional analyses of genes differentially expressed in three
different brain tissues of females following exposure to multiple social conditions. We found a
concordance of 79% in the combination of biological processes (BP), cellular components (CC)
and molecular functions (MF) GO terms enriched following analyses of differentially expressed
genes in the telencephalon (n =158). This value represented a 1.7-fold increase in the concordance
of terms in relation to mean values obtained from corresponding enrichment analyses of 1000
random sets of 158 genes (see Methods; mean concordance ± [CIs]: 45% [43,47]). We likewise
found concordances of 64% for differentially expressed genes in the midbrain (n = 109), and of
4.5% for differentially expressed genes in the optic tectum (n = 21). These represented 2.1-fold
and 1.1-fold increases in relation to analyses with 1000 random sets of 109 and 21 genes in
midbrain and optic tectum respectively (mean concordance midbrain: 30% [28,31]; mean
concordance telencephalon: 3.8% [3.6,4.2]). We summarized and visualized GO terms enrichment
lists across experiments and tissues sampled using REVIGO[36]. We found a strong overlap between
enrichment of GO biological process terms associated with learning and memory, synaptic
processes, neuron projection and cell growth, mostly constrained to the telencephalon and
midbrain regions (Fig. 4). We found similar patterns in relation to cellular component GO terms,
with strong overlap in neuronal components, in particular with high enrichment of terms associated
with glutamatergic synapse. Visualization of GO terms associated with molecular functions
suggests a major role of genes with protein binding function across experiments, including a role
of cadherin-binding related genes in the midbrain (Fig. 4).

Discussion

319
320 We used behavioral phenotyping across guppy families, in conjunction with PoolSeq and RNASeq
321 to identify the genetic architecture of coordinated motion. Our broad range of analyses, spanning
322 genomes, transcriptomes, and phenotypes, provides an exceptional evaluation of the molecular

mechanisms underlying sociability in this fish. Our work suggests that genes and gene networks involved in social decision making through neuron migration and synaptic function are key in the evolution of schooling, highlighting a crucial role of glutamatergic synaptic function and calcium-dependent signaling processes.

Our pedigree-based phenotyping analyses of 195 guppy families from polarization-selected and control lines indicate moderate levels of heritability (Alignment: 0.06 - 0.34; Attraction: 0.09 - 0.26), with pronounced sex differences in full pedigree models (Alignment$_{female-male}$: 0.10 ± 0.05; Attraction$_{female-male}$: -0.17 ± 0.05), in estimates for key behavioral traits forming the sociability axis in this species. Our heritability estimates are similar to estimates for affiliative social behavior traits in primates, ungulates and rodents[10–13,37], and to overall estimates of heritability in personality traits across human and non-human animals[8,38]. Given the importance of social behavior in a range of survival and fitness components in natural systems[1,39,40], our results suggest that complex genetic architectures can respond quickly to strong evolutionary pressures, even when only one sex is subject to selection[22], and that our lab population contained significant amounts of standing genetic variation for these traits prior to selection.

The complex genetic architecture makes it difficult to precisely characterize cross sex genetic effects in our study. We nonetheless observed a positive cross sex genetic correlation in attraction (0.68, CI: 0.25 - 0.98), suggesting similarities between males and females in the genetic architecture of this trait. This result is concordant with a study focused on the bold-shy continuum aspect of personality establishing that sex differences in risk-taking behaviors are weak and likely lack sex-specific genetic architecture in this species[41]. Yet, sex differences in heritability estimates of alignment ($\female h^2_{alignment}$ = 0.34 [0.18, 0.49]; $\male h^2_{alignment}$ = 0.06 [0.00, 0.18]) suggest that it is important to account for sex-specific additive genetic variance when inferring the evolvability of personality traits. Additionally, the low cross-sex heritability we observe in these latter traits is particularly interesting, and suggests that selection in one sex for a complex trait need not result in a correlated response in the other sex. Overall, this indicates significant sex-specific genetic variation for sex-specific behaviours, and that sexually dimorphic behaviors need not require decoupling of male and female genetic architecture when sufficient sex-specific genetic variation is present.

We next mapped the genomic and transcriptomic basis of phenotypic differences in polarization in female guppies. Our GWAS experimental design was designed to compare individuals with high- and low-sociability phenotypes from within polarization-selected lines, rather than between polarization-selected and control lines. This may have resulted in compressed phenotypic spread, but carries the important advantage of reducing the incidence of SNP frequencies that vary across alternative selection lines due to drift. As such, our design is conservative. In our most stringent PoolSeq analysis, we identified SNPs in four genes that consistently differed across all three replicate lines, which have previously been associated with cognition and functions relevant to social behavior. The supt6 histone chaperone and transcription elongation factor homolog (*supt6h*) is important in the positive regulation of transcriptional elongation and a substrate of mTOR, a signalling pathway with a role in cognitive function[42,43]. Ubiquitin specific peptidase 11 (*usp11*) homolog in humans has a critical function in the development of neural cortex, and knock-out studies in mice show that the locus protects females from cognitive impairment[44,45]. Similarly, the novel RNA-binding protein *Nova-1-like* gene is associated with a neuron-specific nuclear RNA binding protein in humans and regulates brain-specific splicing related to synaptic function[46].

Finally, our PoolSeq analysis identified Cadherin 13 (*cdh13*), the human homolog of which has a crucial role in GABAergic function[47], with involvement in neural growth and axonal guidance during early development[48,49]. Moreover, deficit of this gene has a major impact in neurodevelopmental disorders including attention-deficit/hyperactivity disorder and autism spectrum disorder[50]. Indeed, *cdh13* knock-out mice display delayed acquisition of learning tasks and a decreased latency in sociability[51]. Interestingly, repeated selection of genes involved in cadherin-signalling pathways[52] has been shown in guppy populations experiencing different predation pressures. Together, natural selection imposed by differences in predation across these populations[53–55], and our combined findings in the genomic background of guppies suggest strong selective pressures for cadherin-signalling genes due to their modulation of affiliative behaviors.

Our expression results revealed differences in regulation in genes associated with learning, behavior, and neural function, mainly in the telencephalon and midbrain, in comparisons of polarization-selected and control lines in different social contexts. Overall, this suggests that the

regulation of highly demanding cognitive processes via synaptic function underlies variation in

sociability. While the integration of visual signals is central in fish schools[56], our results suggest

that higher order cognitive processes are the basis of variation in social affinity. Indeed, the

differences in alignment and attraction observed when swimming with unfamiliar conspecifics are

arguably highly cognitively demanding, as within a collective motion context, the tendency to

copy the directional movements of other individuals implies a direct trade-off between personal

goal-oriented behaviors and the benefits of social conformity[57,58]. Together, our study of

transcriptomic profiles of schooling fish suggests that the regulation of affiliative behaviors in this

species is driven by an intricately linked social decision-making network in the brain[59], with strong

links to functional groups governing social behaviors and personality across species. More

broadly, our results offer insight into important questions about the evolution of behaviour, and

other traits with complex genetic architecture. First, our results of large-scale expression

differences among selection lines are consistent with recent discussions of the role of gene

regulatory networks in co-ordinating large numbers of genes associated with behaviours[60]. It is

highly likely that the genes with convergent expression changes in the selection lines are controlled

via a modular regulatory architecture, as evidenced by our coexpression network analysis

(Supplementary Tables 7-8, Fig. 3).

We find a striking concordance in the functionality of genes independently identified in genomic

and transcriptomic profiling of strongly differentiated experiments assessing social interactions of

polarization-selected female guppies (Fig. 4). The overlap in significantly enriched Gene Ontology

terms, including learning, synaptic processes and neuron projection restricted to brain regions

associated with decision-making and motor control, strongly reinforces the notion that genetic

regulation of these cognitive processes is fundamental for sociability in guppies. Additionally, the

functional concordance we observe between the regulatory and protein differences among our

selection lines is noteworthy in the context of the discussion of whether structural or regulatory

variation is more important in adaptive phenotypes[61,62]. The overlap in functionality in our

genomic and transcriptomic approaches suggests that both are important, with artificial selection

for behavior acting on coding and regulatory variation within the same pathway to achieve adaptive

phenotypes.

Our results indicate that the regulation of glutamatergic synaptic processes is a particularly
promising network for future studies of affiliative behaviors. Interestingly, differential gene
expression of glutamate receptor genes has been identified to regulate female mating preferences
in guppies[63], and is concordantly identified across species of vertebrates in the regulation of long-
term affiliative mating behaviors[64]. Guppies are livebearers, and this has hindered the use of
functional genetic tools such as CRISPR on this species. Although not feasible at this time, future
functional validation via genetic manipulations of guppies of these pathways would prove
extremely interesting.

Our results are also concordant with other comparative transcriptomic studies of behavioral
responses towards conspecific territorial intrusions, which identified calcium ion-binding
regulation across phylogenetically distant species[65]. Together, the consistency in our findings of
specific genes and functional terms associated with calcium-dependent and cadherin binding
molecular functions across our experiments suggest these are promising molecular targets for
future research exploring the evolution and regulation of sociability and affiliative behaviors.

Methods

<u>Study system</u>
To evaluate the genetic architecture of sociability, we performed a series of experiments in guppies
following artificial selection on coordinated motion. The laboratory population of guppies used
originated from a downstream population of the Quare river in Trinidad, which is subject to high
predation levels. The original collection was made in 1998[66], and the laboratory population has
since been kept in several large (>200-litre) tanks of >200 individuals each to avoid inbreeding.
The artificial selection procedure is outlined in detail in[22,23]. In brief, groups of female guppies
were subjected to repeated open field tests, and were subsequently sorted based on their median
polarization, measured by the degree of alignment exhibited by the individuals within the group
when swimming together[22,23]. For three generations, females from groups with higher polarization
were mated with males from those cohorts to generate three lines of guppies that had been selected
for high polarization. In parallel, random females were exposed to the same experimental
conditions and were mated with unselected males to generate three control lines. Analysis of the
third generation of polarization-selection revealed that, on average, females exhibited a 15%

higher level of polarization and a 10% higher level of group cohesiveness compared to control
females[22].
450
Throughout the selection experiment and the completion of experiments described below all fish
were removed from their parental tanks after birth, separated by sex at the first onset of sexual
maturation, and afterwards kept in single-sex groups of eight individuals in 7-L tanks containing
2 cm of gravel with continuously aerated water, a biological filter, and plants for environmental
enrichment. We allowed for visual contact between the tanks. The laboratory was maintained at
26°C with a 12-h light:12-h dark schedule. Fish were fed a diet of flake food and freshly hatched
brine shrimp daily.
458
<u>Social interactions with unfamiliar individuals and the heritability of sociability</u>
To investigate the heritability and cross-sex genetic correlations of sociability in the guppy, we
measured alignment and attraction with unfamiliar groups of conspecifics in parents and offspring
from polarization-selected and control lines. Specifically, using offspring of the F3 generation of
selection we bred 35 families for each of the three polarization-selected and for each of the three
control lines. From our population of F3 generation offspring (kept in single sex groups prior to
the breeding experiments), we used male and female guppies of the same age (approximately 9
months old) and paired them in 3L tanks to generate the parental generation. We collected
offspring for the two first clutches of these pairs and we transferred newborn offspring to 3L tanks
in groups of three siblings. We separated siblings by sex at the first onset of sexual maturation,
and afterwards we kept them in single-sex groups of three individuals until behavioral testing. We
phenotyped sociability of a total of 195 guppy families: mother, father and six offspring (three
females and three males). Any family for which we did not collect at least three female and three
male offspring was disregarded from further behavioral testing. Each of the six selection lines was
represented with a minimum of 30 families in our heritability analyses.
474
Behavioral assays. To phenotype sociability in each member of our guppy families, we measured
alignment and attraction of 1495 guppies from our breeding experiment. For each fish, we
performed an open field assay using white arenas with 55 cm diameter and 3 cm water depth in
which our focal fish (guppies from the breeding experiment) interacted with a group of seven

same-sex conspecifics. Non-focal guppies used in these assays were from a lab wild-type stock

population and of similar age to our focal fish. Prior to the start of the test, focal fish and the seven-

fish group were acclimated in the centre of the arena for one minute in separate opaque white 15

cm PVC cylinders. After this acclimation period, we lifted the cylinders and filmed the arena for

10 minutes using a Point Grey Grasshopper 3 camera (FLIR Systems; resolution, 2048 pixels by

2048 pixels; frame rate, 25 Hz). Three weeks prior to assays, we tagged wild-type fish with small

black elastomere implants (Northwest Marine Technology Inc.) to allow recognition of wild-type

fish after completion of each assay. After completion, we gently euthanized focal fish from the

parental generation with an overdose of benzocaine and kept them in ethanol for future genomic

analyses. Focal fish from the offspring generation were transferred to group tanks for future

experimental use. Groups of seven wild-type fish were transferred to holding tanks and used in a

maximum of seven assays with focal fish.

Data processing. We tracked the movement of fish groups in the collected video recordings using

idTracker[67] and used fine-grained tracking data to calculate the following variables in Matlab

(v2020): i) alignment, the median alignment of the focal fish to the group average direction across

all frames in the assay. This was quantified by the total length of the sum of two-unit vectors, one

representing the heading of the focal fish, and one representing the heading of the group centroid.

Calculations of alignment were only obtained if six out of the eight members of the group presented

tracks following the optimization of our tracking protocol in the setup in ([22,23,68]); ii) attraction, the

median nearest neighbor distance across all frames in the assay; and iii) activity, we obtained the

median speed across all group members and across all frames by calculating the first derivatives

of the x and y time series, then smoothed using a third-order Savitzky–Golay filter. For all

measurements, trials with less than 70% complete tracks (n= 8) were disregarded for further

analyses. The proportion of frames used did not differ between polarization-selected and control

fish for any comparison across different generations and sexes (Supplementary Figure 5). We

calculated these variables for the focal fish and the average for the seven-fish wild-type group. To

recover focal fish id in the tracking data we used idPlayer to visualize trials by projecting the raw

tracking data onto experimental videos. We followed focal individuals for the first two minutes of

the assay and used the stable identity assigned by idTracker in data collection. In trials with less

than 85% complete tracks (n= 8), we followed focal individuals for the total duration of the

recording to verify the consistency of identity assigned by idTracker. This approach has previously

shown strong reliability in individuals that were observed using this protocol for 20-min recordings

in the same experimental setup that quantified sexual behavior of guppies in mixed-sex shoals[69].

Statistical analyses. Analyses were conducted using R Statistical Software (v4.1.3)[70], RStudio

(v2023.3.1.446)[71] and the tidyverse package[72]. We used Linear Mixed Models (LMMs) with

alignment and attraction as dependent variables to test for potential differences between

polarization-selected and control lines in social interactions with unfamiliar individuals. Selection

regime, sex, the interaction between these two factors and generation were included as fixed

effects. The average activity of the wild-type group was coded as a covariate, with a random

intercept for each replicated selection line, the breeding family, and the number of tests previously

performed with the wild-type group as random factors. All models were run using lme4 and

lmerTest packages[73,74]. Model diagnostics showed that residual distributions were roughly normal

with no evidence of heteroscedasticity.

To estimate heritability, the degree of phenotypic variation due to genetic inheritance, and cross-

sex genetic correlations of alignment and attraction we used Bayesian animal models[75]. Animal

models use a matrix of pedigree relationships set as a random effect, to separate phenotypic

variance for each response variable into additive genetic variance and the remaining variance.

Given strong sex differences in social interactions in guppies, we performed three animal models

for each trait: one including the data on the 1495 phenotyped individuals, and two including only

the phenotyped females or males. Parameter values were estimated using the brms interface[76,77] to

the probabilistic programming language Stan[78]. We used normal priors with a mean of 0 and

standard deviation (sd) of 3 for fixed effects, and student-t priors with 5 degrees of freedom, a

mean of 0 and sd of 5 for random effects. The full pedigree model estimated cross-sex correlations

with a Lewandowski-Kurowicka-Joe (LKJ) prior with $\eta = 1$, which is uniform over the range -1

to 1. Posterior distributions for full / same-sex pedigree models were obtained using Stan's no-U-

turn HMC with 24 / 16 independent Markov chains of 2500 / 4000 iterations, discarding the first

1500 / 2000 iterations per chain as warm-up and resulting in 24000 / 32000 posterior samples

overall. Convergence of the chains and sufficient sampling of posterior distributions were

confirmed by a potential-scale-reduction metric (R) below 1.01, and an effective sample size of at

least 1000. For each model, posterior samples were summarized based on the Bayesian point
estimate (posterior median) and posterior uncertainty intervals by HDIs. We calculated estimates
of heritability by taking the ratio of the additive genetic variance to the total phenotypic variance
in each independent model (see Supplementary Tables 5-6).

<u>Genetic basis of sociability in guppies</u>

Pooled DNA sequencing. We extracted DNA of muscle tissue from the caudal peduncle of
polarization-selected females from the parental generation using Qiagen's DNAeasy Blood &
Tissue kit following standard manufacturer's protocol with an additional on-column RNase A
treatment. We quantified DNA concentration using fluorometry (Qubit; ThermoFisher Scientific).
We next pooled samples from the seven females that represented the top and bottom 20%
polarization-selected guppy lines whose families presented higher and lower sociability in six final
pools at equimolar amounts (Supplementary Figure 1). We achieved a minimum of three µg
genomic DNA per pool. We used a Nextera DNA Flex library preparation kit (Illumina) following
the manufacturer protocol. The final library containing six pooled samples was sequenced at
SciLife Lab, Uppsala (Sweden) in one lane of an Illumina NovaSeq 6000. We obtained on average
31.8 million 150bp read pairs per sample (26.9 million read pairs minimum per sample).

Read quality control and trimming. We assessed the quality of reads for each pool using FastQC
v.0.11.4 (www.bioinformatics.babraham.ac.uk/projects/fastqc). After verifying initial read
quality, reads were trimmed with Trimmomatic v0.35[79]. We filtered adaptor sequences and
trimmed reads if the sliding window average Phred score over four bases was <15 or if the
leading/trailing bases had a Phred score <4, removing reads post filtering if either read pair was
<50 bases in length. Quality was verified after trimming with FastQC.

Genome-wide allele frequency analysis. Reads were mapped using default settings to the guppy
reference genome assembly (Guppy_female_1.0 + MT, RefSeq accession: GCA_000633615.2)[80]
with bwa-mem (v0.7.17)[81]. We used Samtools (v. 1.6.0)[82] to convert sam to bam files, to sort bam
files, to remove duplicates, and to make mpileup files. First, to identify SNPs that significantly
differed in their allele frequencies between guppies with high and low sociability, we merged
sequences from high sociability and low sociability pools and used Popoolation2[27] to create a

synchronized file with allele frequencies for high and low sociability (mpileup2sync.pl –min-qual

20), compute allele frequency differences (mpileup2sync.pl --min-count 6 --min-coverage 25 --

max-coverage 200), calculate Fst for every SNP (fst-sliding.pl) and perform a Fisher's exact test

(fisher-test.pl). Second, we similarly used Popoolation2 to detect consistent changes in allele

frequencies of sociability pooled samples for our three replicated artificial selection lines. For this,

we created one sync file per replicate (mpileup2sync.pl –min-qual 20), and perform a Cochran-

Mantel-Haenszel (CMH) test (cmh-test.pl --min-count 18 --min-coverage 25 --max-coverage

200). Using package qqman[83] in R (v 4.1.3)[70] we made Manhattan plots for each chromosome by

plotting the negative log10-transformed p-values of exact fisher and CMH tests as a function of

chromosome position.

Significance tests and functional analyses. We determined SNPs that were significantly different

between high and low sociability merged pools in Fisher's exact tests using the traditional genome-

wide significance threshold [–log10(p) > 8][28]. We next used custom scripts to identify the overlap

between position of these SNPs and genes present in the guppy reference annotated genome[80], and

to find homologous genes of these set in medaka (*Oryzias latipes*). We further used this set of

unique genes (n = 160) to determine associated GO terms between our merged pools. For this, we

performed enrichment tests in PANTHER[84], as implemented in the GO Ontology Consortium

(http://www.geneontology.org/). To test for enrichments of GO terms we performed one-tail

Fisher's exact tests with a Bonferroni corrected p-value threshold of $p < 0.05$ using a full list of

medaka genes orthologous to guppy genes as background. We used Revigo (http://revigo.irb.hr)[36]

to find and visualize representative subsets of terms based on semantic similarity measurements

for our enriched GO terms related to biological processes, cellular components and molecular

functions.

For CMH test results, we determined SNPs that were significantly different between high and low

sociability pools based on FDR corrected p-value < 0.01. We used a custom script to identify the

overlap between position of these SNPs and genes present in the guppy reference annotated

genome[80].

<u>Neurogenomic response of schooling in guppies</u>

Behavioral assays and tissue collection. Using offspring of the F3 generation (6-months-old), we

placed an individual or groups of eight unfamiliar adult control and polarization-selected females

in white 55cm arenas. After 30 min females were euthanized by transfer to ice water. After 30

seconds, with aid of a Leica S4E microscope, we removed the top of the skull and after cutting

transversally posterior of the optic tectum and anterior of the cerebellum, and horizontally through

the optic chiasm, removed the brain from the skull and placed it into ice water. We severed the

telencephalon from the rest of the brain between the ventral telencephalon and thalamus at the

'commissura anterioris', including both the pallium and subpallium regions. Then we cut the

laminated cup-like structures of the *optic tectum*. The remaining part of the brain was the *midbrain*.

Dissections took under 2 minutes and tissue samples were immediately preserved in RNAlater

(Ambion) at 4°C for 24 hours and then at -20°C until RNA extraction.

RNA extraction and sequencing. For each treatment, we pooled tissue from ten individuals into

two non-overlapping pools of five for each replicate line. We used this strategy to reduce noise in

transcript expression data during sample normalization procedures potentially caused by outliers

during behavioral experiments, while maintaining each replicate as a comparable unit. Our

experimental design represents a total of 120 individual females, constituting six pools per

treatment, per selection regime for a total of 24 pools per tissue. Each sample pool was

homogenized, and RNA was extracted using Qiagen's RNAeasy kits following standard

manufacturer's protocol. Libraries for each sample were prepared and sequenced by the Wellcome

Trust Center for Human Genetics at the University of Oxford, UK. All samples were sequenced

across nine lanes on an Illumina HiSeq 4000. We obtained on average 33.9 million 75bp read pairs

per sample (28.9 million read pairs minimum, 39.8 million maximum).

Read quality control and trimming. We assessed the quality of reads for each sample using

FastQC v.0.11.4 (www.bioinformatics.babraham.ac.uk/projects/fastqc). After verifying initial

read quality, reads were trimmed with Trimmomatic v0.35[79]. We filtered adaptor sequences and

trimmed reads if the sliding window average Phred score over four bases was < 15 or if the

leading/trailing bases had a Phred score < 3, removing reads post filtering if either read pair was <

33 bases in length. Quality was verified after trimming with FastQC.

Differential Expression Analysis. We mapped RNAseq reads against the latest release of the
published guppy genome assembly[80], using the HiSat 2.0.5 – Stringtie v1.3.2 suite[81]. For each
individual pool, reads were mapped to the genome and built into transcripts using default
parameters. The resulting individual assemblies were then merged into a single, non-redundant
assembly using the built-in StringTie-merge function. We filtered the resulting assembly for non-
coding RNA using medaka and Amazon molly (*Poecilia formosa*) non-coding RNA sequences as
reference in a nucleotide BLAST (Blastn). After eliminating all sequences matching non-coding
RNAs, we kept only the longest isoform representative for each transcript for further analysis.
Finally, we quantified expression by re-mapping reads to this filtered assembly using RSEM
(v1.2.20)[85].

Lowly expressed genes were removed by filtering transcripts < 2 RPKM (reads per kilobase per
million mapped reads), preserving only those transcripts that have expression above this threshold
in at least half of the samples for each treatment within a line. After this final filter, a total of
26,140 optic tectum transcripts, 25,100 telencephalon transcripts and 26,514 midbrain transcripts
were retained for further analysis. Using sample correlations in combination with MDS plots based
on all expressed transcripts, we determined that none of the 72 pools represented outliers and thus,
all samples were included in the analysis.

We used DESeq2[86] to normalize filtered read counts using standard function to identify
differentially expressed (DE) genes between the alone and the group treatment in control and
polarization-selected lines separately, and then examined the overlap in differentially expressed
genes between them. A transcript was considered differentially expressed if it had an FDR
corrected p-value < 0.05. As behavior could be modulated by small changes in expression, we did
not filter differentially expressed genes based on Log Fold-change in expression between the
treatments.

Differential Coexpression Analysis. We used Bayes approach for Differential Coexpression
Analysis (BFDCA)[30], in order to identify pairs of genes that have different correlation patterns in
two conditions[32,87,88]. Here we compared the Alone and Group treatments within each line for each
tissue separately, in the same manner as the previously described DE analysis. BFDCA is based

on WGCNA and has been shown to be a reliable and accurate method[30]. This untargeted approach
to differential coexpression (DC) analysis uses a combined Bayes factor, a ratio of marginal
likelihood of the data between the two alternative hypotheses, to evaluate which genes are
differentially correlated in two conditions. We controlled for false positives and accounted for
multiple testing by integrating a random permutations approach[32]. In short, we created 1000
permuted datasets and considered a DC gene pair significant if the Bayes factor for the actual
expression data was larger than the 1% tail of the permutated data Bayes factor distribution.

Functional Analyses. To investigate the function of DE genes we performed GO term enrichment
tests. To accomplish this, we initially completed the annotation of the reference genome assembly.
The transcripts without clear gene names from the reference genome, and the de novo transcripts
identified by HiSat were annotated with blastX against the Swissprot non-redundant database. We
then determined which GO terms were associated with differentially expressed genes and
performed Biological Processes (BP), Cellular Components (CC) and Molecular Functions (MF)
enrichment tests in PANTHER[84]. To assess the level of concordance between genes of interest
across experiments, we compared the proportion of BP, CC and MF GO terms that were
significantly enriched in genomic analyses of sociability implemented in polarization-selected
females, and the proportion of BP, CC and MF GO terms enriched in differential expression
analyses in brain tissue of polarization-selected females following exposure to Group and Alone
experimental conditions. To assess their significance, we compared these values to mean
proportions obtained from bootstrap analyses of 1000 random sets of 158 (for comparison with
telencephalon), 109 (midbrain) and 21 (optic tectum) genes from our medaka-guppy orthologous
gene list. All analyses were based on one-tail Fisher's exact tests with a Bonferroni corrected p-
value threshold of $p < 0.05$ using medaka genes orthologous to guppy genes as the background.
Bootstrap analyses with random sets of genes were automated using rbioapi package[89] in R
(v4.1.3)[70]. We next summarized and visualized GO terms enrichment lists across experiments and
tissues using REVIGO[36] (settings: SimRel semantic similarity measure, 0.5 value). To investigate
the function of differentially coexpressed genes, we used g:Profiler[90] to identify the enriched BP
GO terms and pathways that were altered across mating contexts associated with differentially
coexpressed gene pairs. We determined over-represented pathways among DC gene pairs in each

tissue using the human (Homo sapiens) database in g:Profiler. We chose the human database for

its completeness, acknowledging the distant phylogenetic relationship to guppies.

700   Acknowledgements

We thank David Sumpter, Kristiaan Pelckmans and James Herbert-Read for important
contributions to the conceptualization of the artificial selection procedure. We thank three
anonymous reviewers for comments and suggestions to revise a previous version of the
manuscript. We are grateful to Jacelyn Shu for the design of guppy graphics for figures. We thank
Anna Rennie, Emilio Trejo, and Annika Boussard for help with fish husbandry. **Funding**: This
work was supported by the Knut and Alice Wallenberg Foundation (102 2013.0072 to N.K.), the
Canada 150 Research Chair Program, and the European Research Council (680951 to J.E.M), the
Swedish Research Council (2016-03435 to N.K., 2017-04957), The Royal Swedish Academy of
Sciences (BS2019-0046 to A.C-L.), Lars Hiertas Memorial Foundation (FO2019-0477 to A.C-L.),
European Research Council; H2020 Marie Skłodowska-Curie Actions, (654699 to N.I.B);
Universidad de los Andes (FAPA-4700000443 to N.I.B). **Author contributions**: J.E.M., N.K.
and A.C-L. contributed to conceptualization and funding acquisition of the project. N.K., A.K.,
A.S., and M.R. contributed to the design of the selection procedure and behavioral experiments.
A.C-L. conducted research to obtain behavioral data. A.C-L. and W.vdB. performed formal
analyses and visualization of behavioral and heritability data. A.C-L., M.C-C. conducted research
to obtain genomic data. A.C-L., M.C-C. and I.D. perform formal analyses and visualization of
genomic data. N.B. and A.K. conducted research to obtain transcriptomics data. N.B. and. A.C-L.
performed formal analyses and visualization of transcriptomics data. A.C-L., J.E.M., and N.K.
wrote the original draft. All authors contributed to the final version of the manuscript. **Competing**
interests: The authors declare that they have no competing interests. **Data and materials**
availability: Data and code needed to support the conclusions in the paper are deposited in figshare
repository (doi: 10.6084/m9.figshare.23805702). Genomic and transcriptomic data is deposited at
NCBI (accession codes PRJNA994132 and PRJNA504011). Additional data related to this paper
may be requested from the authors.

728   References

730   1.   Krause, J. & Ruxton, G. D. *Living in Groups*. (Oxford University press, 2002).
731   2.   Réale, D., Reader, S. M., Sol, D., McDougall, P. T. & Dingemanse, N. J. Integrating
732   animal temperament within ecology and evolution. *Biological Reviews* **82**, 291–318
733   (2007).

3. Gartland, L. A., Firth, J. A., Laskowski, K. L., Jeanson, R. & Ioannou, C. C. Sociability as
a personality trait in animals: methods, causes and consequences. *Biological Reviews* **97**,
802–816 (2022).
4. Aplin, L. M. *et al.* Individual personalities predict social behaviour in wild networks of
great tits (Parus major). *Ecol Lett* **16**, 1365–1372 (2013).
5. Bevan, P. A., Gosetto, I., Jenkins, E. R., Barnes, I. & Ioannou, C. C. Regulation between
personality traits: individual social tendencies modulate whether boldness and leadership
are correlated. *Proceedings of the Royal Society B* **285**, (2018).
6. Ebstein, R. P., Israel, S., Chew, S. H., Zhong, S. & Knafo, A. Genetics of Human Social
Behavior. *Neuron* **65**, 831–844 (2010).
7. Robinson, G. E., Fernald, R. D. & Clayton, D. F. Genes and social behavior. *Science
(1979)* **322**, 896–900 (2008).
8. Turkheimer, E., Pettersson, E. & Horn, E. E. A Phenotypic Null Hypothesis for the
Genetics of Personality. *Annual Reviews in Psychology* **65**, 515–540 (2014).
9. Dochtermann, N. A. *et al.* The Heritability of Behavior: A Meta-analysis. *Journal of
Heredity* 1–8 (2019).
10. Brent, L. J. N. *et al.* Genetic origins of social networks in rhesus macaques. *Scientific
Reports 2013 3:1* **3**, 1–8 (2013).
11. Lea, A. J., Blumstein, D. T., Wey, T. W. & Martin, J. G. A. Heritable victimization and
the benefits of agonistic relationships. *Proc Natl Acad Sci U S A* **107**, 21587–21592
(2010).
12. Staes, N. *et al.* Bonobo personality traits are heritable and associated with vasopressin
receptor gene 1a variation. *Scientific Reports 2016 6:1* **6**, 1–8 (2016).
13. Knoll, A. T., Jiang, K. & Levitt, P. Quantitative trait locus mapping and analysis of
heritable variation in affiliative social behavior and co-occurring traits. *Genes Brain
Behav* **17**, (2018).
14. Fisher, D. N. Direct and indirect phenotypic effects on sociability indicate potential to
evolve. *J Evol Biol* **36**, 209–220 (2023).
15. Fisher, D. N. & McAdam, A. G. Social traits, social networks and evolutionary biology. *J
Evol Biol* **30**, 2088–2103 (2017).
16. O'Connell, L. A. & Hofmann, H. A. Genes, hormones, and circuits: An integrative
approach to study the evolution of social behavior. *Front Neuroendocrinol* **32**, 320–335
(2011).
17. Balestri, M., Calati, R., Serretti, A. & De Ronchi, D. Genetic modulation of personality
traits: A systematic review of the literature. *Int Clin Psychopharmacol* **29**, 1–15 (2014).
18. Sahin, M. & Sur, M. Genes, circuits, and precision therapies for autism and related
neurodevelopmental disorders. *Science (1979)* **350**, (2015).
19. Shou, J., Tran, A., Snyder, N., Bleem, E. & Kim, S. Distinct Roles of GluA2-lacking
AMPA Receptor Expression in Dopamine D1 or D2 Receptor Neurons in Animal
Behavior. *Neuroscience* **398**, 102–112 (2019).
20. Abbey-Lee, R. N., Kreshchenko, A., Fernandez Sala, X., Petkova, I. & Løvlie, H. Effects
of monoamine manipulations on the personality and gene expression of three-spined
sticklebacks. (2019).
21. Sumpter, D. J. T., Szorkovszky, A., Kotrschal, A., Kolm, N. & Herbert-Read, J. E. Using
activity and sociability to characterize collective motion. *Philosophical Transactions of
the Royal Society B: Biological Sciences* **373**, 20170015 (2018).

22.   Kotrschal, A. *et al.* Rapid evolution of coordinated and collective movement in response
781   to artificial selection. *Sci Adv* **6**, eaba3148 (2020).
782   23.   Szorkovszky, A. *et al.* An efficient method for sorting and quantifying individual social
783   traits based on group-level behaviour. *Methods Ecol Evol* **8**, 1735–1744 (2017).
784   24.   Piyapong, C. *et al.* Sex matters: a social context to boldness in guppies (Poecilia
785   reticulata). *Behavioral Ecology* **21**, 3–8 (2010).
786   25.   Dimitriadou, S., Croft, D. P. & Darden, S. K. Divergence in social traits in trinidadian
787   guppies selectively bred for high and low leadership in a cooperative context.
788   doi:10.1038/s41598-019-53748-4.
789   26.   Griffiths, S. W. & Magurran, A. E. Sex and schooling behaviour in the Trinidadian guppy.
790   *Anim Behav* **56**, 689–693 (1998).
791   27.   Kofler, R., Pandey, R. V. & Schlötterer, C. PoPoolation2: identifying differentiation
792   between populations using sequencing of pooled DNA samples (Pool-Seq).
793   *Bioinformatics* **27**, 3435–3436 (2011).
794   28.   Pe'er, I., Yelensky, R., Altshuler, D. & Daly, M. J. Estimation of the multiple testing
795   burden for genomewide association studies of nearly all common variants. *Genet*
796   *Epidemiol* **32**, 381–385 (2008).
797   29.   Iancu, O. D., Colville, A., Darakjian, P. & Hitzemann, R. Coexpression and Cosplicing
798   Network Approaches for the Study of Mammalian Brain Transcriptomes. *Int Rev*
799   *Neurobiol* **116**, 73–93 (2014).
800   30.   Wang, D., Wang, J., Jiang, Y., Liang, Y. & Xu, D. BFDCA: A Comprehensive Tool of
801   Using Bayes Factor for Differential Co-Expression Analysis. *J Mol Biol* **429**, 446–453
802   (2017).
803   31.   Jiang, Z., Dong, X., Li, Z.-G., He, F. & Zhang, Z. Differential Coexpression Analysis
804   Reveals Extensive Rewiring of Arabidopsis Gene Coexpression in Response to
805   Pseudomonas syringae Infection OPEN. *Nature Publishing Group* (2016)
806   doi:10.1038/srep35064.
807   32.   Bloch, N. I. *et al.* Different mating contexts lead to extensive rewiring of female brain
808   coexpression networks in the guppy. *Genes Brain Behav* **20**, e12697 (2021).
809   33.   Froemke, R. C. & Young, L. J. Oxytocin, Neural Plasticity, and Social Behavior.
810   *https://doi.org/10.1146/annurev-neuro-102320-102847* **44**, 359–381 (2021).
811   34.   Miyakawa, T. *et al.* Conditional calcineurin knockout mice exhibit multiple abnormal
812   behaviors related to schizophrenia. *Proc Natl Acad Sci U S A* **100**, 8987–8992 (2003).
813   35.   Lee, J. *et al.* Opposing Functions of Calcineurin and CaMKII Regulate G-protein
814   Signaling in Egg-laying Behavior of C. elegans. *J Mol Biol* **344**, 585–595 (2004).
815   36.   Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO summarizes and visualizes long
816   lists of gene ontology terms. *PLoS One* **6**, (2011).
817   37.   Albery, G. F. *et al.* Multiple spatial behaviours govern social network positions in a wild
818   ungulate. *Ecol Lett* **24**, 676–686 (2021).
819   38.   Dochtermann, N. A., Schwab, T. & Sih, A. The contribution of additive genetic variation
820   to personality variation: heritability of personality. *Proceedings of the Royal Society B:*
821   *Biological Sciences* **282**, (2015).
822   39.   Couzin, I. D., Krause, J., Franks, N. R. & Levin, S. A. Effective leadership and decision-
823   making in animal groups on the move. *Nature* **433**, 513–516 (2005).
824   40.   Parrish, J. K. & Edelstein-Keshet, L. Complexity, Pattern, and Evolutionary Trade-Offs in
825   Animal Aggregation. *Science (1979)* **284**, 99–101 (1999).

41. White, S. J., Houslay, T. M. & Wilson, A. J. Evolutionary genetics of personality in the
Trinidadian guppy II: sexual dimorphism and genotype-by-sex interactions. *Heredity 2018*
122:1 **122**, 15–28 (2018).

42. Lipton, J. O. & Sahin, M. The Neurology of mTOR. *Neuron* **84**, 275–291 (2014).

43. Zhao, J., Zhai, B., Gygi, S. P. & Goldberg, A. L. MTOR inhibition activates overall
protein degradation by the ubiquitin proteasome system as well as by autophagy. *Proc*
Natl Acad Sci U S A **112**, 15790–15797 (2015).

44. Chiang, S. Y. *et al.* Usp11 controls cortical neurogenesis and neuronal migration through
Sox11 stabilization. *Sci Adv* **7**, 6093–6105 (2021).

45. Yan, Y. *et al.* X-linked ubiquitin-specific peptidase 11 increases tauopathy vulnerability in
women. *Cell* **185**, 3913-3930.e19 (2022).

46. Buckanovich, R. J. & Darnell, R. B. The neuronal RNA binding protein Nova-1
recognizes specific RNA targets in vitro and in vivo. *Mol Cell Biol* **17**, 3194–3201 (1997).

47. Rivero, O. *et al.* Cadherin-13, a risk gene for ADHD and comorbid disorders, impacts
GABAergic function in hippocampus and cognition. *Translational Psychiatry 2015 5:10*
5, e655–e655 (2015).

48. Ciatto, C. *et al.* T-cadherin structures reveal a novel adhesive binding mechanism. *Nature*
Structural & Molecular Biology 2010 17:3 **17**, 339–347 (2010).

49. Hayano, Y. *et al.* The role of T-cadherin in axonal pathway formation in neocortical
circuits. *Development* **141**, 4784–4793 (2014).

50. Hawi, Z. *et al.* The role of cadherin genes in five major psychiatric disorders: A literature
update. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics* **177**,
168–180 (2018).

51. Forero, A. *et al.* Serotonin (5-HT) neuron-specific inactivation of Cadherin-13 impacts 5-
HT system formation and cognitive function. *Neuropharmacology* **168**, 108018 (2020).

52. Whiting, J. R. *et al.* Drainage-structuring of ancestral variation and a common functional
pathway shape limited genomic convergence in natural high- and low-predation guppies.
PLoS Genet **17**, e1009566 (2021).

53. Seghers, B. H. Schooling Behavior in the Guppy (Poecilia reticulata): An Evolutionary
Response to Predation. *Evolution (N Y)* **28**, 489 (1974).

54. Huizinga, M., Ghalambor, C. K. & Reznick, D. N. The genetic and environmental basis of
adaptive differences in shoaling behaviour among populations of Trinidadian guppies,
Poecilia reticulata. *J Evol Biol* **22**, 1860–1866 (2009).

55. Herbert-Read, J. E. *et al.* How predation shapes the social interaction rules of shoaling
fish. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20171126 (2017).

56. Strandburg-Peshkin, A. *et al.* Visual sensory networks and effective information transfer
in animal groups. *Current biology* **23**, R711 (2013).

57. Conradt, L. & Roper, T. J. Conflicts of interest and the evolution of decision sharing.
Philosophical Transactions of the Royal Society B: Biological Sciences **364**, 807–819
(2008).

58. Ioannou, C. C., Singh, M. & Couzin, I. D. Potential Leaders Trade Off Goal-Oriented and
Socially Oriented Behavior in Mobile Animal Groups. *Am Nat* **186**, 284–293 (2015).

59. Bshary, R., Gingins, S. & Vail, A. L. Social cognition in fishes. *Trends Cogn Sci* **18**, 465–
471 (2014).

60. Sinha, S. *et al.* Behavior-related gene regulatory networks: A new level of organization in
the brain. *Proc Natl Acad Sci U S A* **117**, 23270–23279 (2020).

61. Hoekstra, H. E. & Coyne, J. A. THE LOCUS OF EVOLUTION: EVO DEVO AND THE
GENETICS OF ADAPTATION. *Evolution (N Y)* **61**, 995–1016 (2007).
62. Carroll, S. B. Evo-Devo and an Expanding Evolutionary Synthesis: A Genetic Theory of
Morphological Evolution. *Cell* **134**, 25–36 (2008).
63. Bloch, N. I. *et al.* Early neurogenomic response associated with variation in guppy female
mate preference. *Nature Ecology & Evolution 2018 2:11* **2**, 1772–1781 (2018).
64. Young, R. L. *et al.* Conserved transcriptomic profiles underpin monogamy across
vertebrates. *Proc Natl Acad Sci U S A* **116**, 1331–1336 (2019).
65. Rittschof, C. C. *et al.* Neuromolecular responses to social challenge: Common
mechanisms across mouse, stickleback fish, and honey bee. *Proc Natl Acad Sci U S A*
111, 17929–17934 (2014).
66. Pélabon, C. *et al.* The effects of sexual selection on life-history traits: an experimental
study on guppies. *J Evol Biol* **27**, 404–416 (2014).
67. Pérez-Escudero, A., Vicente-Page, J., Hinz, R. C., Arganda, S. & De Polavieja, G. G.
idTracker: tracking individuals in a group by automatic identification of unmarked
animals. *Nature Methods 2014 11:7* **11**, 743–748 (2014).
68. Kotrschal, A. *et al.* Brain size does not impact shoaling dynamics in unfamiliar groups of
guppies (Poecilia reticulata). *Behavioural Processes* **147**, 13–20 (2018).
69. Corral-López, A., Romensky, M., Kotrschal, A., Buechel, S. D. & Kolm, N. Brain size
affects responsiveness in mating behaviour to variation in predation pressure and sex ratio.
J Evol Biol **33**, (2020).
70. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/ (2021).
71. Posit team. RStudio: Integrated Development Environment for R. Posit Software, PBC,
Boston, MA. URL http://www.posit.co/ (2023).
72. Wickham, H. *et al.* Welcome to the Tidyverse. *J Open Source Softw* **4**, 1686 (2019).
73. Bates, D., Sarkar, D., Bates, M. D. & Matrix, L. The lme4 Package. *R package version* **2**,
74 (2007).
74. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in
Linear Mixed Effects Models. *J Stat Softw* **82**, 1–26 (2017).
75. de Villemereuil, P. On the relevance of Bayesian statistics and MCMC for animal models.
Animal Breeding Genetics **136**, 339–340 (2019).
76. Bürkner, P. C. Advanced Bayesian multilevel modeling with the R package brms. *R J* **10**,
395–411 (2018).
77. Bürkner, P. C. brms: An R package for Bayesian multilevel models using Stan. *J Stat*
Softw **80**, 1–28 (2017).
78. Carpenter, B. *et al.* Stan: A probabilistic programming language. *J Stat Softw* **76**, (2017).
79. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).
80. Künstner, A. *et al.* The Genome of the Trinidadian Guppy, Poecilia reticulata, and
Variation in the Guanapo Population. *PLoS One* **11**, e0169087 (2016).
81. Pertea, M., Kim, D., Pertea, G. M., Leek, J. T. & Salzberg, S. L. Transcript-level
expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown.
Nature Protocols 2016 11:9 **11**, 1650–1667 (2016).
82. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

83.   Turner, S. D. qqman: an R package for visualizing GWAS results using Q-Q and
manhattan plots. *J Open Source Softw* **3**, 731 (2018).
84.   Mi, H., Muruganujan, A., Casagrande, J. T. & Thomas, P. D. Large-scale gene function
analysis with the PANTHER classification system. *Nature Protocols 2013 8:8* **8**, 1551–
1566 (2013).
85.   Li, B. & Dewey, C. N. RSEM: Accurate transcript quantification from RNA-Seq data with
or without a reference genome. *BMC Bioinformatics* **12**, 1–16 (2011).
86.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 1–21 (2014).
87.   Amar, D., Safer, H. & Shamir, R. Dissection of Regulatory Networks that Are Altered in
Disease via Differential Co-expression. *PLoS Comput Biol* **9**, e1002955 (2013).
88.   Fukushima, A. DiffCorr: An R package to analyze and visualize differential correlations
in biological networks. *Gene* **518**, 209–214 (2013).
89.   Rezwani, M., Pourfathollah, A. A. & Noorbakhsh, F. rbioapi: user-friendly R interface to
biologic web services' API. *Bioinformatics* **38**, 2952–2953 (2022).
90.   Reimand, J. *et al.* g:Profiler—a web server for functional interpretation of gene lists (2016
update). *Nucleic Acids Res* **44**, W83–W89 (2016).
91.   Karaca, E. *et al.* Genes that Affect Brain Structure and Function Identified by Rare
Variant Analyses of Mendelian Neurologic Disease. *Neuron* **88**, 499–513 (2015).
92.   Takeichi, M. The cadherin superfamily in neuronal connections and interactions. *Nature*
Reviews Neuroscience 2006 8:1 **8**, 11–20 (2006).

Figures and Tables

Figure 1. Heritability of sociability in guppies. (A) Female, but not male, guppies from polarization-selected lines (orange) presented higher alignment to the group direction (left panel) and shorter distances to their nearest neighbor (higher alignment; right panel) than guppies from control lines (gray) when swimming with unfamiliar same-sex conspecifics (see Supplementary Table 1-2). For all boxplots, horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. Boxes in top left position of each facet indicate p-values ($p < 0.05$ in bold) for statistical contrasts by sex in Linear Mixed Model comparing alignment and attraction between selection line treatments (see Supplementary Tables 1-2). **(B)** Animal models using same-sex pedigrees and full pedigrees with alignment and attraction (nearest neighbor distance) phenotypes in 195 families of polarization-selected and control guppy lines indicated a moderate heritability in female guppies of both biologically-relevant aspects of sociability measured, alignment (left) and attraction (right). In males, we found moderate heritability in attraction, but credible intervals in alignment estimates overlapped with 0 suggesting low heritability of this sociability aspect. Our full pedigree animal models provided large credible intervals for male-female correlations in sociability, with estimates overlapping 0 in alignment, but a positive correlation between sexes in attraction (see Supplementary Tables 4-5.

974 **Table 1. Characterization of genes associated with sociability in guppies**
975

SNP location	Gene ID Ensembl	Gene name	Described cognitive function of homologs	References
Chr 7: 30748139	00000004543	*usp11*	Control of cortical neurogenesis and neuronal migration. Mutations of the gene have been associated with neurological disorders.	Chiang et al, 2021[44] Karaca et al. 2015[91]
Chr 13: 31383940	00000018946	*Novel gene (RNA-binding protein Nova-1-like)*	Neuronal RNA binding protein associated with motor function.	Buckanovich and Darnell 1997[46]
Chr 14: 4286109	00000009725	*supt6h*	Substrate of mTOR, a signaling pathway associated with brain function and neurodegenerative disorders.	Lipton and Sahin 2015[42] Zhao et al. 2015[43]
Chr 18: 4286109	00000014318	*Novel gene*	-	-
Chr 19: 3032099	00000019822	*cdh13*	Modulation of brain activity through GABAergic function. Organization of neuronal circuits.	Rivero et al 2015[47] Takeichi 2006[92]

988
Figure 2. Genetic basis of sociability in the guppy. **(A)** Manhattan plot of -log10(p) values across the guppy
genome resulting from a Fisher's exact test comparing allele frequency differences between high and low
sociability female guppies. We merged pooled-DNA sequences of three independent replicates and found 819
SNPs significantly different (above genome-wide threshold highlighted in red), while a highly stringent analyses
of consistent allele frequency differences across our three independent replicates (cmh-test; see methods)
identified 13 SNPs (five of them within genes) associated with sociability in the species (gene names and SNP
location in the genome highlighted in orange). SNPs with -log10(p) values < 2 are omitted. **(B)** Clustering of
statistically significant overrepresented Gene Ontology annotations for biological processes associated with
differences between high and low sociability in guppies. Point size and color provide information on fold
enrichment value from the statistical overrepresentation test performed in PANTHER[84] (see methods). Terms
with fold enrichment lower than 8 are represented but not described in text. Axes have no intrinsic meaning and
are based on multidimensional scaling which clustered terms based on semantic similarities[74].
1001

1002
Figure 3. Neurogenomic response of schooling in guppies. Hierarchical clustering and relative expression
level for all differentially expressed genes between Alone and Group treatments in the (a) optic tectum, (b) the
telencephalon and (c) the midbrain. Differentially expressed genes were identified separately in polarization and
control line samples. Clustering, based on Euclidian distance, represents transcriptional similarity across all
samples, with bootstrap values (1000 replicates) shown at nodes. Venn diagrams summarize the total number of
differentially expressed (DE) genes and differentialy coexpressed (DC) gene pairs in each tissue for polarization-
selected and control lines.

Figure 4. Functional characterization of genes of interest across experiments. Visualization of functional overlap based on Gene Ontology annotations between genes of interest highlighted in strongly differentiated experimental setups evaluating social interactions of female guppies following experimental evolution for higher polarization: 1- genomic analyses of DNA comparing Pool-seq of high and low sociability female guppies (Left Column); 2- transcriptomic analyses evaluating differentially expressed genes in key brain regions of polarization-selected lines of female guppies exposed to two different social contexts, swimming alone or with a group of conspecifics (TEL: telencephalon, MBR: midbrain, OT: optic tectum; Columns 2-4). Shades of green indicate fold enrichment from our statistical overrepresentation tests performed to gene lists obtained from each experiment.

Supplementary Figures for:

Functional convergence of genomic and transcriptomic genetic architecture underlying sociability in a live-bearing fish

Alberto Corral-Lopez*, Natasha I. Bloch, Wouter van der Bijl, Maria Cortazar-Chinarro, Alexander Szorkovszky, Alexander Kotrschal, Iulia Darolti, Severine D. Buechel, Maksym Romenskyy, Niclas Kolm, Judith E. Mank

*corresponding author: alberto.corral@ebc.uu.se

Figure S1. Mean score for alignment to group direction in females for each of the 195 families measured when evaluating the heritability of sociability by means of open field tests in which guppies from three independently replicated polarization-selected (orange) and control (gray) lines were exposed to a group of other seven conspecifics. We obtained combined Z scores by standardizing independently alignment scores obtained in maternal and offspring generations and by adding the score of the mother to the mean score of all female offspring measure in each family. For all boxplots, horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. Colored circles indicate families that were further used for genomic analyses.

Figure S2. Clustering of statistically significant overrepresented Gene Ontology annotations for cellular components associated to differences between high and low sociability in guppies. Point size and color provide information on fold enrichment value from the statistical overrepresentation test performed in PANTHER[1]. Terms with fold enrichment lower than eight are represented but not described in text. Axes have no intrinsic meaning and are based on multidimensional scaling which cluster terms based on semantic similarities[2].

Figure S3. Clustering of statistically significant overrepresented Gene Ontology annotations for molecular functions associated to differences between high and low sociability in guppies. Point size and color provide information on fold enrichment value from the statistical overrepresentation test performed in PANTHER[1]. Terms with fold enrichment lower than eight are represented but not described in text. Axes have no intrinsic meaning and are based on multidimensional scaling which cluster terms based on semantic similarities[2].

Figure S4. Top panels: Hierarchical clustering of gene expression across all significantly expressed genes in the optic tectum (a), telencephalon (b) and midbrain (c) for polarization-selected selection and control lines. Values on top of nodes correspond to Approximately Unbiased bootstrap values. Heatmap represents expression of 500 randomly selected genes. **Bottom panels:** Boxplots of average log2 fold-change for all genes differentially expressed genes between the *Alone* and the *Group* conditions in polarization-selected and control lines for optic tectum (d), telencephalon (e) and midbrain (f). Boxes correspond to 25^{th} - 75th percentiles. Sample sizes on the x-axis indicate number of genes with relative up- or down-regulation.

Figure S5. Proportion of frames obtained from idTracker software in tracking from videos recorded for groups of seven wild-type guppies and one individual from either polarization-selected or control selection lines. Tracking data was used next for the quantification of alignment, attraction and speed in parents and six offspring of 195 families. We found no significant differences for the proportion of frames used for videos used for polarization-selected and control lines in a Linear Mixed Model that included generation, sex and selection line as fixed factors (LMM_{frames}: line: F= 0.82; df = 1464; p = 0.37). Horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range. P-values in top position of each comparison indicate values for statistical contrasts of the model by sex and generation. Consistency in identity assigned by idTracker was verified in trials with less than 85% frames tracked (dashed line).

Supplementary References

1 - Mi, H., Muruganujan, A., Casagrande, J. T. & Thomas, P. D. Large-scale gene function analysis with the PANTHER classification system. *Nature Protocols 2013 8:8* **8**, 1551–1566 (2013).

2 - Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One* **6**, (2011).

Supplementary Tables for:

Functional convergence of genomic and transcriptomic genetic architecture underlying sociability in a live-bearing fish

Alberto Corral-Lopez*, Natasha I. Bloch, Wouter van der Bijl, Maria Cortazar-Chinarro, Alexander Szorkovszky, Alexander Kotrschal, Iulia Darolti, Severine D. Buechel, Maksym Romenskyy, Niclas Kolm, Judith E. Mank

*corresponding author: alberto.corral@ebc.uu.se

Table S1. Statistical results for a Linear Mixed Model comparing the alignment and attraction of polarization-selected and control female guppies when swimming with groups of same-sex non-kin conspecifics.

	Alignment to group				
Predictors	*Estimates*	*CI*	*Statistic*	*p*	*df*
(Intercept)	0.19	0.13 – 0.24	6.78	**<0.001**	1385.81
Selection line [polarization]	0.01	0.00 – 0.01	2.27	**0.047**	9.68
Sex [males]	-0.01	-0.01 – -0.00	-3.02	**0.003**	690.08
Generation [parental]	-0.02	-0.02 – -0.01	-10.13	**<0.001**	1141.24
Average alignment of group	0.79	0.72 – 0.86	22.71	**<0.001**	1436.72
Average speed of group	0.00	0.00 – 0.00	0.20	0.842	1429.55
Selection line [polarization] * Sex [males]	0.00	-0.00 – 0.00	0.20	0.842	1429.55
ICC	0.12				
N_{line}	2				
N_{rep}	3				
N_{testnr}	8				
N_{sex}	2				
N_{family}	195				
Observations	1486				
Marginal R^2 / Conditional R^2	0.657 / 0.697				

	Nearest neighbour distance (attraction)				
Predictors	*Estimates*	*CI*	*Statistic*	*p*	*df*
(Intercept)	2.64	2.56 – 2.71	65.62	**<0.001**	565.12
Selection line [polarization]	-0.04	-0.08 – -0.00	-2.33	**0.044**	9.34
Sex [males]	0.01	-0.02 – 0.04	0.55	0.583	448.53
Generation [parental]	0.12	0.10 – 0.14	11.31	**<0.001**	984.50
Average alignment of group	0.02	0.02 – 0.02	27.31	**<0.001**	1408.27

Average speed of group	0.00	0.00 – 0.00	3.74	**<0.001**	566.82
Selection line [polarization] * Sex [males]	0.03	-0.02 – 0.07	1.24	0.217	371.59
ICC	0.13				
N_{line}	2				
N_{rep}	3				
N_{testnr}	8				
N_{sex}	2				
N_{family}	195				
Observations	1485				
Marginal R^2 / Conditional R^2	0.471 / 0.538				

Table S2. Independent contrasts for comparisons between polarization-selected (P) and control (C) guppies in their alignment and nearest neighbor distance (attraction) when swimming with groups of same-sex non-kin conspecifics.

Alignment

contrast	sex	estimate	SE	df	t.ratio	p.value
C - P	females	-0.006	0.002	9.68	-2.269	**0.047**
C - P	males	-0.003	0.002	9.56	1.378	0.20

Attraction

contrast	treatment	estimate	SE	df	t.ratio	p.value
C - P	females	0.042	0.001	9.34	2.331	**0.043**
C - P	males	0.015	0.002	9.26	0.883	0.40

Table S3. Mean body size of guppies used to estimate the heritability of alignment and attraction at the time of testing across sexes and generations. Values are provided in number of pixels obtained from idTracker data using a custom script implemented in Matlab (v2020a) that extracts body size data accounting for changes in apparent size between the middle and edge of the experimental arena used for video recordings.

Generation	Selection line	Sex	Mean body size	SE
Parents	Control	Females	410	47
		Males	192	29
	Polarization-selected	Females	416	71
		Males	189	37
Offspring	Control	Females	247	55
		Males	158	33
	Polarization-selected	Females	251	60
		Males	160	26

Table S4. Results from animal models only accounting for same-sex pedigree relationships. We obtained heritability estimates by taking the ratio of the additive genetic variance to the total phenotypic variance (additive genetic variance + group variance + population variance + residual variance) in each independent model.

ALIGNMENT

FEMALES					
Parameter group	**Parameter**	**Estimate**	**Est.Error**	**Q2.5**	**Q97.5**
Fixed effects	Intercept	0.504	0.074	0.359	0.648
Fixed effects	linesocial	0.157	0.102	-0.04	0.355
Fixed effects	generationparents	-0.277	0.058	-0.39	-0.163
Fixed effects	speed_std	0.396	0.023	0.35	0.441
Additive genetic variance	Intercept	0.162	0.042	0.086	0.25
Group	Intercept	0.005	0.006	0	0.02
Population	Intercept	0.01	0.027	0	0.064
Residual variance	NA	0.304	0.037	0.234	0.379
Heritability		0.336	0.079	0.183	0.494

MALES					
Parameter group	**Parameter**	**Estimate**	**Est.Error**	**Q2.5**	**Q97.5**
Fixed effects	Intercept	-0.452	0.125	-0.694	-0.213
Fixed effects	linesocial	0.093	0.172	-0.231	0.425
Fixed effects	generationparents	-0.502	0.071	-0.641	-0.362
Fixed effects	speed_std	0.298	0.039	0.221	0.376
Additive genetic variance	Intercept	0.031	0.026	0.000	0.091
Group	Intercept	0.04	0.016	0.013	0.074
Population	Intercept	0.039	0.091	0.001	0.207
Residual variance	NA	0.390	0.033	0.323	0.453
Heritability		0.063	0.052	0.000	0.184

ATTRACTION

FEMALES					
Parameter group	**Parameter**	**Estimate**	**Est.Error**	**Q2.5**	**Q97.5**
Fixed effects	Intercept	-0.614	0.099	-0.811	-0.424
Fixed effects	linesocial	-0.088	0.138	-0.35	0.179
Fixed effects	generationparents	0.251	0.070	0.113	0.390
Fixed effects	speed_std	-0.109	0.024	-0.157	-0.062
Additive genetic variance	Intercept	0.094	0.038	0.025	0.174
Group	Intercept	0.038	0.015	0.011	0.071
Population	Intercept	0.023	0.061	0.000	0.133
Residual variance	NA	0.355	0.039	0.280	0.435
Heritability		0.185	0.072	0.049	0.336

MALES

Parameter group	Parameter	Estimate	Est.Error	Q2.5	Q97.5
Fixed effects	Intercept	0.476	0.134	0.212	0.746
Fixed effects	linesocial	-0.025	0.181	-0.389	0.345
Fixed effects	generationparents	0.614	0.088	0.440	0.785
Fixed effects	speed_std	0.054	0.046	-0.037	0.145
Additive genetic variance	Intercept	0.137	0.049	0.047	0.242
Group	Intercept	0.075	0.023	0.036	0.126
Population	Intercept	0.046	0.097	0.000	0.250
Residual variance	NA	0.454	0.050	0.359	0.556
Heritability		0.194	0.069	0.065	0.337

Table S5 Results from animal models with full pedigree relationships. We obtained heritability estimates for each sex by taking the ratio of the additive genetic variance of each sex to the total phenotypic variance (additive genetic variance + group variance + population variance + residual variance) in each independent model.

ALIGNMENT

Parameter group	Parameter	Estimate	Est.Error	Q2.5	Q97.5
Fixed effects	Intercept	0.52	0.102	0.331	0.718
Fixed effects	sexm	-0.961	0.063	-1.08	-0.838
Fixed effects	linesocial	0.163	0.128	-0.097	0.419
Fixed effects	speed_std	0.36	0.021	0.319	0.4
Fixed effects	sexm:linesocial	-0.075	0.074	-0.22	0.07
Fixed effects	sexf:generationparents	-0.31	0.066	-0.444	-0.181
Fixed effects	sexm:generationparents	-0.479	0.064	-0.604	-0.352
Additive genetic variance	sexf	0.105	0.029	0.051	0.166
Additive genetic variance	sexm	0.048	0.027	0.003	0.105
Additive genetic variance	cross (sexf x sexm)	0.583	0.270	0.021	0.974
Group	Intercept	0.025	0.009	0.009	0.045
Population	Intercept	0.023	0.051	0.001	0.13
Residual variance	NA	0.365	0.026	0.314	0.415
♀ **Heritability**		0.202	0.052	0.102	0.308
♂ **Heritability**		0.103	0.057	0.007	0.224
Δ Heritability (♀ - ♂)		0.098	0.053	0.202	-0.005

ATTRACTION

Parameter group	Parameter	Estimate	Est.Error	Q2.5	Q97.5
Fixed effects	Intercept	-0.627	0.115	-0.847	-0.391
Fixed effects	sexm	1.09	0.075	0.942	1.23
Fixed effects	linesocial	-0.095	0.157	-0.425	0.219
Fixed effects	speed_std	-0.067	0.022	-0.109	-0.024
Fixed effects	sexm:linesocial	0.072	0.082	-0.091	0.235
Fixed effects	sexf:generationparents	0.283	0.079	0.129	0.438
Fixed effects	sexm:generationparents	0.568	0.080	0.411	0.722
Additive genetic variance	sexf	0.050	0.025	0.007	0.104
Additive genetic variance	sexm	0.176	0.041	0.102	0.263
Additive genetic variance	sexm	0.438	0.304	-0.168	0.950
Group	Intercept	0.059	0.013	0.035	0.087
Population	Intercept	0.032	0.056	0.002	0.172
Residual variance	NA	0.406	0.028	0.351	0.460
♀ **Heritability**		0.092	0.044	0.013	0.188
♂ **Heritability**		0.261	0.055	0.157	0.371
Δ Heritability (♀ - ♂)		-0.169	0.051	-0.267	-0.069

Table S6. Differentially expressed genes in analyses of the genomic response in midbrain, optic tectum and telencephalon in polarization-selected and control lines of female guppies when swimming with a group of conspecifics versus when swimming alone.

MIDBRAIN

CONTROL LINES (n=36)				
Transcript	**Gene BLAST**	**Average expression**	**Fold Change**	**P-value (FDR)**
rna32553|gene=LOC103477857	PLCL2	131.212	27.132	<0.001
MSTRG.16482.1|gene=MSTRG.16482	MORN4	37.361	-22.976	<0.001
MSTRG.17365.1|gene=MSTRG.17365	RPGP2	32.181	-22.742	<0.001
rna23930|gene=ubap2l	ubap2l	944.324	-0.346	<0.001
rna14333|gene=trim33	trim33	41.183	19.099	<0.001
rna12705|gene=LOC103466075	MAGI2	1073.054	-0.398	<0.001
rna14220|gene=LOC103466954	FLNA	259.466	8.280	<0.001
rna43613|gene=LOC103458483	SYNE1	7853.703	0.243	<0.001
rna46407|gene=lrfn5	lrfn5	1152.716	-0.665	<0.001
rna12704|gene=LOC103466075	MAGI2	1284.870	0.358	0.001
MSTRG.18176.1|gene=MSTRG.18176	KMCP1	206.043	-0.800	0.001
rna32800|gene=u2af2	u2af2	1910.842	-0.346	0.001
MSTRG.16838.1|gene=MSTRG.16838	K1841	386.297	0.320	0.002
MSTRG.4096.1|gene=MSTRG.4096	AN13C	167.036	1.072	0.002
rna9959|gene=kcnab2	kcnab2	2321.886	-0.139	0.003
MSTRG.14391.4|gene=MSTRG.14391	KMT2A	2251.545	0.175	0.004
rna6890|gene=LOC103462605	IGF1R	151.381	5.464	0.004
MSTRG.16227.1|gene=MSTRG.16227	RIBA2	144.642	-0.623	0.004
rna27802|gene=fndc3a	fndc3a	639.741	-0.517	0.004
rna42558|gene=itpkb	itpkb	253.080	-0.714	0.006
MSTRG.6879.2|gene=MSTRG.6879	AEBP2	492.647	-0.492	0.009
rna15437|gene=LOC103467579	IQEC1	272.493	-0.544	0.010
MSTRG.10425.1|gene=MSTRG.10425	METK2	2522.539	-0.149	0.010
rna39037|gene=tanc2	tanc2	352.805	11.879	0.011
MSTRG.17415.2|gene=MSTRG.17415	FLOT1	1005.390	-0.284	0.018
MSTRG.3778.3|gene=MSTRG.3778	MAP1B	10822.870	0.090	0.022
rna19452|gene=LOC103470078	SCAM1	1692.812	0.214	0.027
rna30679|gene=LOC103476732	ANK3	6992.816	0.149	0.027
rna42097|gene=LOC103457647	NA	245.652	-0.955	0.027
MSTRG.7457.1|gene=MSTRG.7457	FLNA	1583.394	-0.378	0.033
MSTRG.7577.1|gene=MSTRG.7577	F19A5	1752.482	0.375	0.035
rna1276|gene=acsl1	acsl1	1052.049	-0.289	0.037
rna46838|gene=LOC103460240	FBL10	99.185	-0.667	0.038
rna18681|gene=LOC103469777	SMAD2	106.022	-0.567	0.042

Transcript	Gene BLAST	Average expression	Fold Change	P-value (FDR)
rna14493|gene=wdr6	wdr6	151.669	0.864	0.048
rna1738|gene=keap1	keap1	2161.342	-0.225	0.049

POLARIZATION-SELECTED LINES (n=109)

Transcript	Gene BLAST	Average expression	Fold Change	P-value (FDR)
rna7061|gene=LOC103462879	PPR3E	144.810	10.467	< 0.001
rna9051|gene=LOC103463550	TLE4	90.600	-24.816	< 0.001
rna24376|gene=scn1b	scn1b	86.552	24.017	< 0.001
rna44020|gene=srrm1	srrm1	13.933	21.607	< 0.001
MSTRG.24305.1|gene=MSTRG.24305	NA	2510.567	0.748	< 0.001
rna22214|gene=egr1	egr1	30.831	18.839	< 0.001
rna22213|gene=egr1	egr1	30.831	18.839	< 0.001
rna40013|gene=LOC103482261	CPNE3	115.525	-1.047	< 0.001
rna45945|gene=syn3	syn3	2021.198	0.623	< 0.001
MSTRG.17634.6|gene=MSTRG.17634	CHIO	726.382	0.310	< 0.001
MSTRG.11465.4|gene=MSTRG.11465	LAMP2	1545.620	0.231	< 0.001
MSTRG.13269.1|gene=MSTRG.13269	FBCD1	2483.148	0.135	< 0.001
MSTRG.24306.3|gene=MSTRG.24306	SYN3	3100.717	1.061	< 0.001
rna26833|gene=LOC103474460	NCPR	105.368	5.135	0.001
MSTRG.3790.1|gene=MSTRG.3790	NA	6932.462	0.120	0.001
rna46319|gene=rps14	rps14	6939.074	-0.112	0.001
MSTRG.25027.1|gene=MSTRG.25027	NA	440.681	0.287	0.001
rna29486|gene=ubb	ubb	1424.005	-0.204	0.001
MSTRG.3031.1|gene=MSTRG.3031	NRK2	907.778	-0.234	0.001
rna35365|gene=LOC103479452	UN13A	866.972	-0.478	0.001
rna13371|gene=znf746	znf746	1623.600	0.312	0.001
MSTRG.10067.35|gene=MSTRG.10067	KCC2B	7708.760	0.138	0.002
MSTRG.5587.1|gene=MSTRG.5587	NA	387.393	0.325	0.002
rna24078|gene=LOC103472856	SHSA7	1873.094	0.380	0.002
rna4922|gene=LOC103461631	PTPRN	6258.089	0.342	0.002
rna30660|gene=rgs7	rgs7	2386.077	-0.155	0.002
rna46774|gene=abca2	abca2	985.996	0.437	0.002
rna44214|gene=stxbp6	stxbp6	1369.630	0.560	0.002
MSTRG.14206.1|gene=MSTRG.14206	SMBT1	81.425	-8.926	0.002
rna31627|gene=LOC103477235	BTBD3	5069.169	0.085	0.003
MSTRG.22324.1|gene=MSTRG.22324	INT7	201.560	0.525	0.003
rna15437|gene=LOC103467579	IQEC1	262.086	-0.619	0.003
rna30743|gene=LOC103476767	PTPRE	830.200	-0.300	0.003
MSTRG.23588.1|gene=MSTRG.23588	SC5A1	268.214	0.409	0.003
rna18718|gene=sema4c	sema4c	225.613	-0.778	0.003

rna18418|gene=LOC103469455	ZN208	964.520	0.791	0.004
MSTRG.24653.1|gene=MSTRG.24653	ZN665	647.821	-0.301	0.004
rna7261|gene=prc1	prc1	69.432	-0.673	0.004
rna33374|gene=itga8	itga8	177.778	-0.492	0.006
rna14054|gene=LOC103466873	CPNE9	1292.066	-0.186	0.006
rna25858|gene=add1	add1	301.934	-1.200	0.007
rna41611|gene=abracl	abracl	53.205	7.912	0.007
rna8442|gene=tbc1d23	tbc1d23	532.141	0.310	0.009
MSTRG.15056.1|gene=MSTRG.15056	ST32A	275.300	0.492	0.009
rna2302|gene=LOC103472525	NEUL1	298.298	-0.452	0.009
MSTRG.10567.1|gene=MSTRG.10567	EDIL3	901.674	0.304	0.009
rna20215|gene=klf9	klf9	230.772	0.515	0.009
MSTRG.24002.1|gene=MSTRG.24002	KCNA1	5020.815	0.142	0.009
rna43122|gene=npas3	npas3	579.387	0.533	0.010
MSTRG.11745.1|gene=MSTRG.11745	P33MX	3488.184	0.108	0.010
rna28848|gene=gng8	gng8	189.567	-0.795	0.011
MSTRG.14250.1|gene=MSTRG.14250	TMTC3	9253.455	0.190	0.011
rna32670|gene=LOC103477783	PNRC2	6535.494	0.077	0.011
rna37128|gene=LOC103480432	COR1B	96.683	-7.060	0.011
rna836|gene=LOC103465321	PLAK	165.686	-2.434	0.012
MSTRG.22229.1|gene=MSTRG.22229	NA	989.886	0.301	0.012
rna14269|gene=LOC103466973	MYT1	150.362	-8.518	0.012
rna26517|gene=LOC103474357	NA	680.176	0.312	0.013
rna14680|gene=tardbp	tardbp	786.594	-0.256	0.015
rna18417|gene=LOC103469455	ZN208	987.995	-0.992	0.016
rna12053|gene=rpl27a	rpl27a	5598.155	-0.121	0.016
rna3113|gene=fam171b	fam171b	676.632	-0.622	0.017
rna10216|gene=LOC103464678	S12A5	465.767	-0.326	0.017
MSTRG.24178.2|gene=MSTRG.24178	AEBP2	406.228	-0.382	0.018
rna41338|gene=clg20hxorf23	clg20hxorf23	38.754	-5.680	0.018
rna25124|gene=rpl37	rpl37	3795.051	-0.136	0.018
rna21184|gene=dusp1	dusp1	601.109	0.245	0.019
rna22048|gene=LOC103471326	NA	5028.797	0.222	0.021
MSTRG.7705.1|gene=MSTRG.7705	TADBP	659.341	0.309	0.021
rna15823|gene=LOC103467371	LFG2	6610.378	0.128	0.021
rna42117|gene=foxo3	foxo3	4839.429	0.112	0.021
rna15369|gene=baz2a	baz2a	2468.169	0.271	0.023
rna16768|gene=wbp11	wbp11	637.220	0.247	0.023
rna36928|gene=dgkk	dgkk	330.445	-0.856	0.024
rna8044|gene=uchl5	uchl5	1901.123	-0.123	0.025
rna12667|gene=erc1	erc1	1227.615	0.223	0.026

Transcript	Gene BLAST	Average expression	Fold Change	P-value (FDR)
MSTRG.20354.5|gene=MSTRG.20354	DLG5	269.626	0.856	0.029
rna2024|gene=LOC103470494	GRIA2	2812.527	0.166	0.030
rna10139|gene=pdhb	pdhb	2640.130	-0.139	0.030
MSTRG.8108.1|gene=MSTRG.8108	143BA	11583.530	0.122	0.030
MSTRG.11772.3|gene=MSTRG.11772	RNF6	375.432	0.328	0.030
rna35260|gene=wdr47	wdr47	2309.695	-0.136	0.030
MSTRG.1635.3|gene=MSTRG.1635	NLGNX	1082.247	0.316	0.030
MSTRG.6043.23|gene=MSTRG.6043	NFASC	4875.050	0.157	0.031
rna19015|gene=LOC103469839	SNF5	558.868	-0.187	0.031
rna30678|gene=LOC103476732	ANK3	748.739	0.483	0.031
rna26037|gene=LOC103473957	STXB1	55589.294	0.068	0.031
rna27530|gene=LOC103474859	H33	10943.188	-0.110	0.034
rna8930|gene=LOC103463612	RORB	1423.800	0.184	0.034
rna31916|gene=rps27a	rps27a	10460.725	-0.096	0.035
rna41322|gene=eif2s3	eif2s3	2821.758	-0.139	0.035
rna29177|gene=LOC103475891	AIFM1	41.012	5.526	0.035
MSTRG.25212.1|gene=MSTRG.25212	PLXA3	597.785	0.488	0.037
MSTRG.10784.1|gene=MSTRG.10784	RYR1	25.899	1.375	0.039
rna6858|gene=LOC103462628	CHST8	1822.516	0.123	0.039
rna19888|gene=LOC103470302	NA	327.381	-0.746	0.040
MSTRG.11901.1|gene=MSTRG.11901	NA	1157.565	0.231	0.044
rna28113|gene=actn4	actn4	1503.002	-0.194	0.044
MSTRG.24804.1|gene=MSTRG.24804	ABCA2	1009.735	-0.353	0.045
MSTRG.15875.1|gene=MSTRG.15875	ARHGC	511.656	0.253	0.045
rna1297|gene=fbxo41	fbxo41	2701.441	0.134	0.046
rna11314|gene=LOC103465330	AGAP2	872.129	0.343	0.047
rna25393|gene=cdc37l1	cdc37l1	253.320	-0.270	0.047
rna43115|gene=akap6	akap6	659.763	4.534	0.047
MSTRG.9250.1|gene=MSTRG.9250	MFS6L	901.045	0.193	0.048
rna22955|gene=gatad2b	gatad2b	5809.079	0.156	0.048
MSTRG.1327.2|gene=MSTRG.1327	G137C	1064.576	0.197	0.048
MSTRG.17605.2|gene=MSTRG.17605	CEP72	4481.235	0.158	0.048
MSTRG.9514.1|gene=MSTRG.9514	KPCA	2230.300	0.158	0.049

OPTIC TECTUM

CONTROL LINES (n=187)				
Transcript	Gene BLAST	Average expression	Fold Change	P-value (FDR)
rna31156|gene=LOC103476975	ERLN1	44.054	-23.947	< 0.001
rna45754|gene=kcnc2	kcnc2	862.523	0.336	< 0.001
MSTRG.3467.33|gene=MSTRG.3467	MADD	535.023	-0.426	< 0.001
MSTRG.15801.1|gene=MSTRG.15801	FEZ1	709.036	0.450	< 0.001

rna47175|gene=fsd1l	fsd1l	57.585	-6.841	< 0.001
rna8739|gene=LOC103463733	PAIRB	184.831	-8.877	0.001
rna11571|gene=nfasc	nfasc	204.340	-10.374	0.001
rna21659|gene=gsr	gsr	152.518	-0.791	0.001
MSTRG.14289.2|gene=MSTRG.14289	CLPB	860.114	-0.369	0.004
MSTRG.18014.5|gene=MSTRG.18014	SAC1A	583.077	-0.301	0.004
rna38083|gene=eme2	eme2	298.137	-0.519	0.004
MSTRG.5196.2|gene=MSTRG.5196	HEMK1	387.510	-1.001	0.004
rna28725|gene=proser1	proser1	414.492	-0.419	0.004
MSTRG.15730.1|gene=MSTRG.15730	MAP6	1424.326	-0.361	0.004
rna28934|gene=LOC103475715	NPAS2	236.645	-1.310	0.004
MSTRG.917.1|gene=MSTRG.917	MYG	2925.534	0.261	0.004
MSTRG.9656.1|gene=MSTRG.9656	MFS11	246.930	-0.343	0.005
rna12640|gene=sephs1	sephs1	61.114	-8.344	0.005
rna9975|gene=timp4	timp4	4018.568	0.133	0.005
rna21336|gene=sparc	sparc	2184.843	0.289	0.005
rna18695|gene=LOC103469766	XNIF	6926.725	-0.137	0.005
MSTRG.13682.4|gene=MSTRG.13682	DYN1	902.700	-0.436	0.005
rna39368|gene=LOC103481895	PHIPL	1250.523	0.275	0.006
MSTRG.24446.1|gene=MSTRG.24446	NA	129.841	-0.756	0.006
rna20093|gene=LOC103470431	rsmG	56982.959	0.171	0.006
MSTRG.10409.5|gene=MSTRG.10409	HIP1R	229.374	-1.111	0.007
MSTRG.24774.24|gene=MSTRG.24774	E41L3	222.455	-0.670	0.008
rna17061|gene=cacna1i	cacna1i	967.348	-0.420	0.009
rna19619|gene=pde4d	pde4d	389.720	-0.416	0.009
rna23861|gene=clk2	clk2	1467.038	-0.164	0.009
rna20745|gene=LOC103470831	S12A2	1144.392	0.736	0.009
MSTRG.22864.5|gene=MSTRG.22864	MABP1	646.420	-0.385	0.009
rna39712|gene=cacna1g	cacna1g	915.637	-0.347	0.010
MSTRG.10346.1|gene=MSTRG.10346	PERF	664.481	0.237	0.010
MSTRG.24190.1|gene=MSTRG.24190	NYNRI	734.469	-0.722	0.010
MSTRG.13834.2|gene=MSTRG.13834	NMDZ1	4878.659	-0.246	0.010
rna46689|gene=LOC103460147	E41L3	326.656	-0.451	0.010
rna1478|gene=suclg1	suclg1	310.395	0.424	0.010
rna18243|gene=LOC103469344	RHG23	663.109	0.551	0.010
MSTRG.16227.1|gene=MSTRG.16227	RIBA2	133.475	-0.668	0.010
rna11521|gene=LOC103465433	NA	3055.567	-0.322	0.010
rna32056|gene=kiaa0319	kiaa0319	425.023	-0.831	0.011
rna33697|gene=LOC103478428	FEZ2	157.395	-0.446	0.011
rna39112|gene=LOC103481688	ANK3	1151.108	-0.739	0.011
MSTRG.7552.1|gene=MSTRG.7552	GBRD	924.733	-0.409	0.011

rna6818|gene=chd9	chd9	506.594	1.474	0.012
MSTRG.6879.2|gene=MSTRG.6879	AEBP2	755.567	-0.594	0.013
rna8820|gene=bsg	bsg	5022.883	0.117	0.013
rna32896|gene=rbp5	rbp5	700.965	0.314	0.013
rna9941|gene=LOC103464464	NA	6700.276	0.142	0.013
rna4264|gene=s100b	s100b	44245.997	0.203	0.013
rna10292|gene=LOC103464714	SYT2	951.110	-0.683	0.013
rna32087|gene=LOC103477466	PI4KB	1081.824	-0.409	0.013
rna24933|gene=LOC103473353	SMCA2	1265.118	-0.322	0.013
rna8437|gene=gtpbp6	gtpbp6	407.444	-0.307	0.013
MSTRG.20719.13|gene=MSTRG.20719	SHSA6	1502.578	-0.408	0.013
rna43889|gene=LOC103458562	CO052	769.649	-0.511	0.013
rna31795|gene=rims1	rims1	871.867	-0.267	0.013
rna39949|gene=arhgef4	arhgef4	391.940	-0.310	0.013
MSTRG.14169.1|gene=MSTRG.14169	APBP2	489.681	-0.306	0.013
MSTRG.17087.27|gene=MSTRG.17087	SYGP1	2095.222	-0.290	0.013
MSTRG.18474.3|gene=MSTRG.18474	GDE	1577.365	-0.197	0.014
rna24455|gene=ago4	ago4	805.347	0.399	0.014
MSTRG.15686.3|gene=MSTRG.15686	SRRM3	873.289	-0.377	0.014
rna24998|gene=fnbp1	fnbp1	733.773	-0.604	0.015
rna27624|gene=serpine2	serpine2	3522.939	0.124	0.015
rna35717|gene=LOC103479568	PSA7	1970.107	0.113	0.015
rna5025|gene=LOC103460921	PDXK	3641.298	0.200	0.016
MSTRG.24191.1|gene=MSTRG.24191	EFC1	758.821	-0.790	0.016
MSTRG.19248.1|gene=MSTRG.19248	NALP1	250.455	-0.743	0.016
rna14471|gene=LOC103467107	VWA1	291.946	0.459	0.016
rna27096|gene=sik2	sik2	1497.974	-0.340	0.016
rna4809|gene=dapl1	dapl1	529.088	0.361	0.016
rna20030|gene=LOC103470390	LCAP	1033.226	0.321	0.017
rna385|gene=LOC103474313	HEBP2	6881.290	0.112	0.018
rna14585|gene=LOC103467181	GSTM1	383.607	0.332	0.018
MSTRG.1958.1|gene=MSTRG.1958	LRRF2	1052.046	-0.222	0.021
rna16860|gene=pick1	pick1	224.937	-0.839	0.021
MSTRG.9169.3|gene=MSTRG.9169	TTYH3	243.157	0.339	0.021
MSTRG.3724.24|gene=MSTRG.3724	SHAN2	960.854	-0.308	0.022
rna29486|gene=ubb	ubb	1938.270	0.196	0.023
rna35679|gene=LOC103479583	DPP6	226.201	-0.421	0.024
rna1353|gene=pgm2	pgm2	298.878	1.051	0.024
rna4718|gene=ndufs1	ndufs1	704.549	0.317	0.024
rna25185|gene=LOC103473483	SCRB2	1298.625	0.227	0.024
MSTRG.11218.1|gene=MSTRG.11218	CC142	510.828	-0.225	0.024

rna6428|gene=LOC103462451	TRI66	691.559	0.489	0.024
rna39783|gene=atad1	atad1	224.854	-0.367	0.024
rna39263|gene=LOC103481783	ZN235	294.711	-0.637	0.025
rna31608|gene=LOC103477221	CYT	2957.013	0.211	0.027
rna8962|gene=LOC103463594	ZFYV9	2108.108	-0.195	0.027
rna30529|gene=LOC103476656	GDS1	1925.695	0.332	0.027
MSTRG.4676.4|gene=MSTRG.4676	DOT1L	1214.736	-0.288	0.027
rna25497|gene=LOC103473681	NA	2400.413	0.321	0.027
rna33379|gene=LOC103478617	HEY1	707.379	0.232	0.028
rna42980|gene=gja1	gja1	24024.747	0.156	0.029
rna1352|gene=pgm2	pgm2	430.082	-0.541	0.029
MSTRG.4713.3|gene=MSTRG.4713	ZFYV9	3966.457	-0.290	0.029
rna5711|gene=mrpl46	mrpl46	342.311	0.276	0.029
MSTRG.6387.2|gene=MSTRG.6387	AASS	519.728	-0.348	0.029
rna9652|gene=ccdc51	ccdc51	89.289	0.739	0.029
MSTRG.4313.1|gene=MSTRG.4313	LORF2	40.998	1.225	0.029
MSTRG.14276.7|gene=MSTRG.14276	HIP1	502.571	-0.272	0.029
MSTRG.6781.1|gene=MSTRG.6781	NA	176.637	-0.375	0.029
rna34295|gene=aldh2	aldh2	948.974	0.186	0.029
MSTRG.8952.3|gene=MSTRG.8952	CAC1I	2257.156	-0.189	0.030
rna16138|gene=LOC103468055	L1CAM	2054.282	-0.230	0.030
rna17402|gene=rhbdf1	rhbdf1	461.449	-0.505	0.030
rna45|gene=fbxl18	fbxl18	1526.888	-0.201	0.030
MSTRG.21801.2|gene=MSTRG.21801	NA	161.643	-11.290	0.030
MSTRG.11045.2|gene=MSTRG.11045	STAG2	696.471	-0.327	0.030
MSTRG.18008.6|gene=MSTRG.18008	RBM42	981.920	-0.292	0.030
MSTRG.5146.19|gene=MSTRG.5146	CAPS1	2211.292	-0.368	0.031
rna554|gene=LOC103464924	NA	1135.002	0.275	0.031
MSTRG.22927.5|gene=MSTRG.22927	PANK2	199.419	-0.358	0.031
MSTRG.4337.1|gene=MSTRG.4337	NA	163.807	-0.398	0.031
MSTRG.6421.1|gene=MSTRG.6421	DNM1L	284.253	-0.331	0.032
MSTRG.20883.51|gene=MSTRG.20883	CAC1G	2176.919	-0.280	0.032
rna17397|gene=hbz	hbz	2654.700	0.777	0.032
rna9952|gene=pax7	pax7	2715.333	0.252	0.033
MSTRG.12285.7|gene=MSTRG.12285	CSPG5	498.929	-0.448	0.033
MSTRG.4508.3|gene=MSTRG.4508	ANR29	148.773	-0.558	0.034
rna15287|gene=LOC103467653	MYPT2	161.547	-0.529	0.034
MSTRG.21917.2|gene=MSTRG.21917	AIG1	812.919	-0.226	0.034
rna31978|gene=cul9	cul9	815.724	-0.454	0.034
MSTRG.5079.1|gene=MSTRG.5079	CCD66	158.970	-0.655	0.034
rna33477|gene=hdac9	hdac9	1208.808	-0.204	0.034

rna32208|gene=LOC103477566	NA	136.825	-8.874	0.034
MSTRG.23173.1|gene=MSTRG.23173	M3K9	393.992	-0.443	0.034
rna44660|gene=rcan3	rcan3	1297.437	0.176	0.034
MSTRG.22663.6|gene=MSTRG.22663	CNIH3	1399.089	-0.365	0.035
rna2408|gene=LOC103472140	STRN	534.708	-0.374	0.035
MSTRG.12275.1|gene=MSTRG.12275	NA	289.805	-0.465	0.035
MSTRG.11276.1|gene=MSTRG.11276	NA	808.305	0.259	0.035
MSTRG.6086.4|gene=MSTRG.6086	SETD5	646.909	-0.482	0.035
rna20492|gene=ncs1	ncs1	1548.351	0.135	0.035
rna44860|gene=mtpn	mtpn	3939.993	0.096	0.035
rna9563|gene=LOC103464301	SC6A6	3463.148	0.093	0.035
MSTRG.20597.113|gene=MSTRG.20597	ANK3	3978.012	-0.257	0.035
rna15537|gene=LOC103467937	ST1S3	92.866	0.506	0.035
MSTRG.2919.2|gene=MSTRG.2919	KPBB	1982.178	-0.275	0.035
MSTRG.8196.8|gene=MSTRG.8196	CDK17	523.422	-0.384	0.035
MSTRG.10756.1|gene=MSTRG.10756	CACP	73.826	1.141	0.036
MSTRG.21665.4|gene=MSTRG.21665	ASAP1	2262.217	-0.338	0.036
rna9074|gene=LOC103463535	SAFB2	486.492	0.877	0.036
MSTRG.6893.1|gene=MSTRG.6893	KCND2	751.620	-0.249	0.037
rna37092|gene=gabarap	gabarap	9554.826	0.122	0.037
MSTRG.21609.4|gene=MSTRG.21609	MYPT1	538.478	-1.474	0.037
MSTRG.17335.7|gene=MSTRG.17335	KPCA	1293.373	-0.233	0.037
MSTRG.14062.6|gene=MSTRG.14062	NCAM1	671.251	-0.400	0.038
rna48393|gene=LOC103461400	PPAL	460.373	0.240	0.039
rna43205|gene=ckb	ckb	568.475	0.225	0.039
rna46432|gene=ddx46	ddx46	2735.921	-0.244	0.039
rna10428|gene=LOC103464779	HES5	738.767	0.267	0.039
rna13605|gene=LOC103466450	RFX7	1889.887	0.384	0.039
rna1698|gene=LOC103468483	ELOF1	256.797	0.280	0.039
rna37063|gene=bcl6b	bcl6b	201.983	0.501	0.039
rna36246|gene=smg7	smg7	2826.022	-0.149	0.039
MSTRG.8764.2|gene=MSTRG.8764	DDX5	9210.083	-0.235	0.039
rna20266|gene=LOC103470511	DNM1L	1990.942	-0.169	0.039
MSTRG.8986.28|gene=MSTRG.8986	CAC1A	5587.949	-0.207	0.039
MSTRG.12290.2|gene=MSTRG.12290	CTNB1	2910.930	-0.206	0.039
rna564|gene=gde1	gde1	785.583	0.260	0.041
MSTRG.16875.3|gene=MSTRG.16875	S22AF	106.886	-0.823	0.041
MSTRG.11867.7|gene=MSTRG.11867	GRIA2	7579.132	-0.254	0.041
rna28723|gene=thsd1	thsd1	124.740	-0.707	0.042
rna32852|gene=casp2	casp2	194.575	0.487	0.042
MSTRG.19932.2|gene=MSTRG.19932	ERF3A	147.054	-0.532	0.042

Transcript		Gene BLAST	Average expression	Fold Change	P-value (FDR)
MSTRG.14143.1|gene=MSTRG.14143		WSB1	2604.092	-0.329	0.043
MSTRG.1695.1|gene=MSTRG.1695		WBP4	118.747	-0.790	0.043
rna12783|gene=cib1		cib1	172.780	0.314	0.045
rna39046|gene=LOC103481669		RHG27	219.443	0.377	0.045
rna4111|gene=etv5		etv5	177.296	0.799	0.045
rna45179|gene=LOC103459268		CAC1C	1903.168	-0.854	0.046
rna43863|gene=zbtb42		zbtb42	95.269	-10.618	0.047
rna43182|gene=tyro3		tyro3	1426.998	0.392	0.047
rna26654|gene=snx19		snx19	2368.539	-0.171	0.047
rna16854|gene=LOC103468448		BORG5	259.020	0.308	0.048
MSTRG.8096.18|gene=MSTRG.8096		WNK2	3178.569	-0.321	0.049
MSTRG.17167.1|gene=MSTRG.17167		ARL4A	107.263	-9.441	0.049
MSTRG.10976.1|gene=MSTRG.10976		NA	339.637	-0.260	0.049
rna18354|gene=LOC103469494		EIF1	480.176	-0.425	0.049
rna12804|gene=slc1a2		slc1a2	38882.612	0.102	0.049
MSTRG.5171.1|gene=MSTRG.5171		RBM5	1463.233	-0.303	0.049
MSTRG.18686.1|gene=MSTRG.18686		RHG12	753.715	-0.292	0.049
MSTRG.24504.1|gene=MSTRG.24504		RAE1	743.520	-0.294	0.049
MSTRG.3297.5|gene=MSTRG.3297		DGLA	461.380	-0.328	0.049

POLARIZATION-SELECTED LINES (n=21)

Transcript	Gene BLAST	Average expression	Fold Change	P-value (FDR)
MSTRG.21801.2|gene=MSTRG.21801	NA	102.640	24.315	< 0.001
rna20241|gene=entpd4	entpd4	13.232	21.501	< 0.001
rna5604|gene=cdh8	cdh8	1553.154	0.643	< 0.001
rna24376|gene=scn1b	scn1b	92.098	19.494	< 0.001
rna22214|gene=egr1	egr1	39.104	18.875	< 0.001
rna22213|gene=egr1	egr1	39.104	18.875	< 0.001
MSTRG.2915.2|gene=MSTRG.2915	CADH8	292.560	-7.961	< 0.001
rna15769|gene=LOC103467406	DSLE	156.024	-0.735	0.002
MSTRG.22698.1|gene=MSTRG.22698	LAMA4	269.181	-8.539	0.004
rna29823|gene=wwc1	wwc1	283.825	-9.054	0.008
rna1269|gene=gucy1a3	gucy1a3	206.417	-0.773	0.010
rna28422|gene=LOC103475388	CTR3	1020.217	0.605	0.015
rna35022|gene=LOC103479254	KC1G2	174.252	-1.351	0.020
rna16375|gene=LOC103468166	MPP2	287.359	0.758	0.020
rna29120|gene=LOC103475822	SPTB2	245.255	-8.404	0.032
rna12010|gene=slc6a15	slc6a15	4110.120	0.134	0.034
rna23231|gene=ahdc1	ahdc1	875.602	-0.421	0.034
rna35400|gene=LOC103479433	P66B	129.015	-0.435	0.034

Transcript	Gene BLAST	Average expression	Fold Change	P-value (FDR)
rna39880|gene=tmem94	tmem94	138.800	8.215	0.040
rna7779|gene=LOC103463266	RAB3C	271.999	-0.557	0.040
				0.040
rna41072|gene=LOC103456849	ESCO1	357.308	11.958	

TELENCEPHALON

<div align="center">

CONTROL LINES (n=29)

</div>

Transcript	Gene BLAST	Average expression	Fold Change	P-value (FDR)
rna16008|gene=zmynd12	zmynd12	131.990	3< 0.001	< 0.001
rna45464|gene=slc37a3	slc37a3	38.721	-22.872	< 0.001
rna14004|gene=LOC103466852	CNGB1	40.820	22.427	< 0.001
rna36260|gene=LOC103480019	PTPR2	26.631	-22.537	< 0.001
MSTRG.16482.1|gene=MSTRG.16482	MORN4	15.553	-22.396	< 0.001
rna9271|gene=ncln	ncln	35.830	8.337	< 0.001
rna1296|gene=fbxo41	fbxo41	364.240	-0.593	< 0.001
rna10792|gene=plxna1	plxna1	299.891	1.086	0.001
rna34474|gene=LOC103478864	HHATL	249.894	0.719	0.003
MSTRG.16496.1|gene=MSTRG.16496	LDB1	437.207	-0.439	0.006
rna8510|gene=cc2d2a	cc2d2a	231.497	-0.488	0.007
rna45291|gene=grip1	grip1	1272.284	0.304	0.009
rna564|gene=gde1	gde1	945.463	0.225	0.009
rna752|gene=tsc2	tsc2	572.522	-0.466	0.014
rna33839|gene=LOC103478362	COBL	316.960	0.373	0.015
rna38088|gene=LOC103481105	VATL	241.247	-0.797	0.015
MSTRG.10409.5|gene=MSTRG.10409	HIP1R	136.542	-1.063	0.016
MSTRG.20355.9|gene=MSTRG.20355	KCMA1	1362.688	-0.259	0.017
MSTRG.21801.2|gene=MSTRG.21801	NA	148.516	-8.653	0.018
rna13396|gene=islr2	islr2	11284.671	0.081	0.020
MSTRG.22908.2|gene=MSTRG.22908	ATRN	287.114	-0.510	0.020
MSTRG.22993.1|gene=MSTRG.22993	SSBP3	1070.362	0.254	0.020
rna2770|gene=map4k4	map4k4	1829.434	0.204	0.020
MSTRG.205.3|gene=MSTRG.205	IFT27	76.967	2.301	0.021
rna45311|gene=ptprb	ptprb	221.601	-10.953	0.027
rna28655|gene=LOC103475549	TRIM3	234.203	0.738	0.027
rna26416|gene=LOC103474211	ROBO2	1300.445	-0.301	0.044
MSTRG.23960.3|gene=MSTRG.23960	RB6I2	643.993	-0.251	0.046
rna16561|gene=LOC103468305	FRIM	3458.490	-0.117	0.049

<div align="center">

POLARIZATION-SELECTED LINES (n=158)

</div>

Transcript	Gene BLAST	Average expression	Fold Change	P-value (FDR)
rna18687|gene=tia1	tia1	401.711	0.422	< 0.001

MSTRG.23945.1|gene=MSTRG.23945	CAC1C	1299.068	0.768	< 0.001
rna31577|gene=LOC103477205	TACC2	2488.829	0.178	< 0.001
MSTRG.21916.1|gene=MSTRG.21916	ZEP2	6873.343	0.187	< 0.001
rna9693|gene=abhd14a	abhd14a	1497.664	-0.168	0.001
MSTRG.18769.2|gene=MSTRG.18769	KIF1A	7524.576	0.217	0.001
rna20419|gene=rictor	rictor	558.369	0.353	0.001
rna42366|gene=LOC103457350	MAX	355.081	0.441	0.001
MSTRG.7449.1|gene=MSTRG.7449	AT2B3	4327.516	0.172	0.001
MSTRG.11745.1|gene=MSTRG.11745	P33MX	3551.885	-0.160	0.002
rna15494|gene=LOC103467551	SLMAP	1057.196	-0.220	0.002
rna10910|gene=LOC103464892	SORT1	2240.260	0.317	0.002
rna46476|gene=ppm1e	ppm1e	1440.302	0.315	0.003
rna39999|gene=LOC103482255	PFKAP	882.019	-1.088	0.003
rna39992|gene=LOC103482255	PFKAP	1367.223	0.413	0.003
rna9416|gene=LOC103464216	LRC24	586.747	0.403	0.003
MSTRG.9631.2|gene=MSTRG.9631	ADA11	7045.660	0.126	0.003
rna42290|gene=gphn	gphn	1236.892	0.239	0.006
rna139|gene=pcm1	pcm1	415.579	0.513	0.006
MSTRG.26314.1|gene=MSTRG.26314	LIMS1	248.767	-0.421	0.006
MSTRG.9224.4|gene=MSTRG.9224	GFAP	2397.131	-0.223	0.008
rna21649|gene=atp8a1	atp8a1	223.183	7.673	0.008
rna30694|gene=pgbd5	pgbd5	1308.949	0.466	0.008
MSTRG.14919.1|gene=MSTRG.14919	KLC1	317.188	-0.494	0.009
rna44843|gene=magi2	magi2	333.647	-6.397	0.009
rna4264|gene=s100b	s100b	8575.720	-0.294	0.011
rna8601|gene=LOC103463821	REM2	1266.508	0.215	0.011
MSTRG.5319.1|gene=MSTRG.5319	MTG8R	289.321	0.659	0.011
MSTRG.8385.2|gene=MSTRG.8385	CHDH	613.368	-0.337	0.011
rna35142|gene=LOC103479283	PTPRS	3184.232	0.228	0.011
MSTRG.25460.5|gene=MSTRG.25460	DPYL2	1490.984	-0.159	0.011
rna10217|gene=LOC103464678	S12A5	292.625	0.760	0.011
rna3294|gene=LOC103477739	T22D1	942.977	0.314	0.013
MSTRG.16563.2|gene=MSTRG.16563	KHDR2	596.821	-0.263	0.014
MSTRG.904.5|gene=MSTRG.904	SMCA4	825.667	0.940	0.014
rna7625|gene=LOC103463153	KCC4	813.768	0.306	0.015
rna23887|gene=LOC103472687	RS27	8841.463	-0.132	0.015
rna40585|gene=atp6v1a	atp6v1a	3486.538	0.279	0.015
rna28612|gene=LOC103475568	NBEA	1738.565	0.230	0.015
rna30893|gene=eif4ebp2	eif4ebp2	4206.554	-0.129	0.015
rna22733|gene=LOC103472023	SIA4A	329.212	-0.484	0.016
rna22398|gene=gria2	gria2	1501.733	0.394	0.018

MSTRG.13007.1|gene=MSTRG.13007	DEN1A	523.827	-0.258	0.018
MSTRG.604.1|gene=MSTRG.604	PCD10	853.615	-0.214	0.020
rna20270|gene=LOC103470513	BMP1	704.668	0.216	0.020
MSTRG.8142.2|gene=MSTRG.8142	GRM7	2211.218	0.219	0.020
rna26271|gene=apbb1	apbb1	616.593	-0.529	0.021
rna7160|gene=LOC103463007	SHAN1	1246.491	0.303	0.021
rna38233|gene=jph3	jph3	1481.033	0.154	0.021
rna9994|gene=atp2b2	atp2b2	2968.692	0.160	0.021
MSTRG.7852.1|gene=MSTRG.7852	TM269	642.667	-0.332	0.021
rna41217|gene=u2surp	u2surp	260.347	0.496	0.021
rna16889|gene=smcr8	smcr8	505.442	0.242	0.021
rna20696|gene=LOC103470760	ODO1	3680.969	0.128	0.022
rna6714|gene=madd	madd	581.771	0.781	0.024
rna28842|gene=LOC103475446	PP2BA	1272.204	0.483	0.025
rna18171|gene=crebbp	crebbp	477.352	0.698	0.025
MSTRG.24874.5|gene=MSTRG.24874	LHPL5	1008.636	-0.240	0.025
rna7423|gene=LOC103463224	DDAH1	614.691	-0.294	0.025
rna38076|gene=LOC103481097	MPRIP	1004.516	0.694	0.026
rna15448|gene=bicd2	bicd2	802.184	0.257	0.026
rna21869|gene=mid2	mid2	858.588	-0.264	0.028
rna1133|gene=LOC103463598	S14L1	402.061	0.463	0.028
rna18056|gene=tom1l2	tom1l2	1234.140	-0.201	0.028
rna26413|gene=LOC103474211	ROBO2	957.985	0.457	0.029
rna3034|gene=tmem47	tmem47	663.543	-0.408	0.029
MSTRG.26426.1|gene=MSTRG.26426	AT2B3	509.885	0.223	0.029
rna14675|gene=kif1b	kif1b	6926.073	0.221	0.029
rna28593|gene=pds5b	pds5b	4058.133	0.116	0.029
rna6897|gene=LOC103462737	RGRF1	3742.185	0.223	0.029
rna12565|gene=cacna2d1	cacna2d1	307.267	0.642	0.029
rna25879|gene=LOC103473860	SHC3	917.756	0.542	0.030
MSTRG.20400.1|gene=MSTRG.20400	NETR	2933.854	-1.107	0.030
rna10085|gene=r3hdm2	r3hdm2	882.792	0.293	0.030
rna2232|gene=sh3gl2	sh3gl2	1153.102	0.336	0.030
rna3256|gene=LOC103477319	NA	369.700	-0.551	0.030
rna7150|gene=cttn	cttn	1432.342	0.154	0.030
MSTRG.23582.7|gene=MSTRG.23582	UNC79	2182.014	-0.299	0.031
rna28372|gene=LOC103475354	SEP8A	1479.812	0.221	0.031
rna40511|gene=stau2	stau2	3751.772	0.101	0.031
rna18955|gene=rasgef1b	rasgef1b	314.812	0.469	0.031
rna40847|gene=cnksr2	cnksr2	2782.045	0.236	0.031
rna32682|gene=LOC103477774	NA	5094.213	-0.150	0.031

rna11361|gene=LOC103465345	LRP1	1464.591	0.319	0.031
rna12226|gene=ptprz1	ptprz1	939.568	-0.197	0.032
MSTRG.24892.1|gene=MSTRG.24892	JUN	1541.880	0.159	0.032
rna48421|gene=LOC103461426	PACA	6547.169	0.114	0.032
MSTRG.1426.7|gene=MSTRG.1426	MCF2L	1541.030	0.216	0.032
rna19642|gene=smarcad1	smarcad1	374.477	0.279	0.033
rna45583|gene=LOC103459444	PRIC1	953.356	0.188	0.033
rna9152|gene=ttll7	ttll7	351.793	0.344	0.033
rna9558|gene=rims4	rims4	2057.525	0.123	0.033
rna26824|gene=baz1b	baz1b	250.301	0.419	0.033
rna27920|gene=LOC103475082	CUL5	478.350	0.422	0.034
rna30614|gene=LOC103476702	PKHA1	1179.339	-0.144	0.034
MSTRG.10701.2|gene=MSTRG.10701	DPYL2	293.373	-0.561	0.035
rna22264|gene=LOC103471194	SFXN1	1299.021	-0.161	0.035
rna45277|gene=cand1	cand1	2417.395	0.115	0.035
rna44024|gene=srrm1	srrm1	719.011	0.472	0.035
rna14390|gene=gnb1	gnb1	2102.920	0.133	0.035
rna19155|gene=camsap1	camsap1	366.586	-0.849	0.035
rna11115|gene=LOC103465107	BSN	3020.560	0.361	0.038
rna13371|gene=znf746	znf746	1535.108	0.216	0.038
rna39374|gene=LOC103481898	KGP1	3810.363	0.112	0.038
rna45925|gene=LOC103459684	CDK17	1635.928	0.146	0.038
rna22088|gene=LOC103471305	SSH	1433.568	0.179	0.038
rna9540|gene=LOC103464285	AT2B4	966.284	0.325	0.038
rna7314|gene=LOC103462926	GSLG1	2352.158	0.169	0.039
rna25580|gene=dgcr2	dgcr2	862.995	0.353	0.039
rna19014|gene=ralgds	ralgds	289.546	0.335	0.039
rna46771|gene=rad17	rad17	289.675	-0.260	0.040
rna21459|gene=LOC103471609	ZIC3	848.137	0.211	0.041
rna33285|gene=LOC103478674	T200B	768.169	0.236	0.041
rna29284|gene=klc2	klc2	1027.397	0.157	0.041
rna21130|gene=LOC103471057	BRD8	579.688	0.296	0.041
rna47838|gene=LOC103460885	ATRN1	749.916	0.186	0.042
rna43237|gene=jade1	jade1	486.768	0.251	0.042
rna39189|gene=srcap	srcap	3117.647	0.130	0.043
rna5118|gene=LOC103460319	TM163	2283.059	0.153	0.044
rna40679|gene=arfgef1	arfgef1	1093.914	0.292	0.044
rna36138|gene=LOC108167071	MIB1	579.766	-0.245	0.044
rna33009|gene=gsk3a	gsk3a	390.156	0.245	0.045
rna33558|gene=kmt2b	kmt2b	1100.465	0.334	0.045
rna7409|gene=LOC103463058	LRP8	248.208	0.326	0.045

rna36173|gene=dnajb4	dnajb4	1321.044	0.178	0.045
rna2990|gene=LOC103474271	SCN2A	797.677	0.430	0.045
rna12431|gene=LOC103465918	S6A13	261.539	-0.709	0.045
rna39333|gene=LOC103481878	GRID1	688.931	0.397	0.045
MSTRG.11445.1|gene=MSTRG.11445	TSN7	913.167	-0.359	0.045
rna4111|gene=etv5	etv5	603.824	0.555	0.045
rna14581|gene=LOC103467180	AMPD2	4309.644	0.105	0.045
rna17113|gene=sox8	sox8	1165.511	0.165	0.045
rna6646|gene=LOC103462596	TLN2	1334.136	0.289	0.045
rna1884|gene=acox1	acox1	922.029	-0.142	0.045
MSTRG.11781.2|gene=MSTRG.11781	ATRX	5501.649	0.144	0.045
rna41153|gene=ptprn2	ptprn2	11939.245	0.175	0.045
rna4922|gene=LOC103461631	PTPRN	8049.927	0.206	0.045
rna2054|gene=LOC103470599	TEN3	754.778	-0.460	0.045
rna37531|gene=LOC103480722	PP2BA	3022.883	0.160	0.045
MSTRG.2884.1|gene=MSTRG.2884	CD81	3884.961	-0.100	0.045
rna10431|gene=LOC103464782	HES5	297.115	-0.279	0.045
rna35326|gene=LOC103479394	SCG2	38239.656	0.110	0.045
rna19050|gene=LOC103469848	KCC2B	398.238	0.440	0.046
MSTRG.9859.3|gene=MSTRG.9859	TIA1	386.806	-0.289	0.046
rna2730|gene=app	app	1253.390	0.129	0.046
rna43935|gene=kbtbd11	kbtbd11	3946.524	0.180	0.048
rna22259|gene=gabrb2	gabrb2	2528.544	0.149	0.048
rna28126|gene=hnrnpl	hnrnpl	1361.666	0.189	0.048
rna4001|gene=sumo1	sumo1	1338.464	-0.129	0.049
rna19299|gene=slc20a2	slc20a2	629.265	-0.668	0.049
MSTRG.10579.6|gene=MSTRG.10579	MEF2C	2175.170	-0.305	0.049
rna32394|gene=cspg5	cspg5	7576.875	0.148	0.049
rna12473|gene=syt12	syt12	4226.735	0.156	0.049
rna11522|gene=LOC103465434	CDK18	613.584	0.944	0.049
MSTRG.21665.4|gene=MSTRG.21665	ASAP1	1872.921	-0.200	0.049
MSTRG.8757.1|gene=MSTRG.8757	UBP43	232.268	-0.358	0.049
rna43050|gene=ndufaf4	ndufaf4	394.548	-0.229	0.049
rna28590|gene=fry	fry	756.197	0.748	0.049

Table S7. Summary of differentially co-expressed gene pair results

	OPTIC TECTUM			TELENCEPHALON			MIDBRAIN		
	Control line		Schooling line	Control line		Schooling line	Control line		Schooling line
DC gene pairs	29		65	228		35	67		60
DC unique genes	27		61	161		36	61		55
DC genes in common (between CT and SCH lines)		NA			ELAV2			MKL2 IGF1R TRIM39 RPL19 POLR1B	
Genes that are DC in more than one comparison	TNIKB GBA2		ELAVL2 MSTRG.8823 RANBP3	NTMT1 RL19 TRI39	LOC103468557 ARL4A FACR1	GRIA2 GOLGA5 PARVA	KCAB1 IGF1R PFKFB2	LBH MKL2 PTGR2	SURF2 POLR1B

Table S8. List of differentially coexpressed genes identified by BFDCA by brain region and line with number of differentially correlated connections (DC gene pairs) for each gene. Genes highlighted in green are DC in more than one comparison, and those highlighted in blue are also Diferentially Expressed.

OPTIC TECTUM				TELENCEPHALON				MIDBRAIN			
Control lines		Polarization-selected lines		Control lines		Polarization-selected lines		Control lines		Polarization-selected lines	
ADPRH	4	TLK2	7	mea1	13	slc17a6	4	MAGI1	5	LOC103461370	6
CGAS	4	AP1B1	6	KAT7	7	tenm2	4	csk	4	PLAK	6
ABCA1	3	rtn4	5	ndufs4	7	tm2d1	4	IGF1R	4	mmp16	5
CENPS	3	ZN226	5	apmap	6	clmn	3	PPARA	4	CNTNAP5	4
MSTRG.7285	3	PARVA	4	arhgef4	6	FACR1	3	CL17A	3	CPSF3	4
LOC103468557	3	pfkfb2	4	NIPLB	6	KHDR2	3	F264	3	pou4f1	4
NRIP3	3	SHC3	4	pkn1	6	LRC24	3	hmga2	3	abracl	3
pcdh19	3	tom1l2	4	rpl17	6	PTPRS	3	LOC103468544	3	CCDC127	3
STIMA	3	ASIC1	3	TRIM69	6	SUIS_HUMAN	2	LOC103468557	3	DNAI2	3
SURF2	3	CCDC51	3	ARIP4	5	grm1	2	MILK1	3	HES6	3
BC11B	2	CDH6	3	cndp2	5	LBH	2	polr1b	3	LOC103461363	3
DLGAP2	2	ERLN1	3	ldhb	5	MACF1	2	RASSF5	3	NUP205	3
ENAH	2	GRID2	3	LRRT4	5	MSTRG.5809	2	rbm33	3	otp	3
GBA2	2	KCC2B	3	rps8	5	mthfs	2	rbm45	3	pdgfrb	3
KCNH7	2	KLHL17	3	smarca4	5	nrbp2	2	S4A7	3	ABRAL	2
mapk6	2	mphosph8	3	tbc1d13	5	PNN	2	ABCD1	2	ARI1B	2
rbms1	2	ntmt1	3	AT8B2	4	RAB3A	2	dctn1	2	ARL4A	2
slc29a2	2	r3hdm2	3	CYTB	4	rheb	2	dnal1	2	ARL6	2
tmbim6	2	TSPAN8	3	dvl1	4	rpl7	2	DNPH1	2	CALB2	2
ADHX	1	cdk14	2	gatad2b	4	RXRBA	2	DOCK9	2	clg20hxorf23	2
ARL4A	1	CROL2	2	htr7	4	TLK1B	2	eaf1	2	CORO2A	2
clns1a	1	EAF1	2	ier2	4	TM205	2	GRIA2	2	ctps1	2
SCD5	1	FOXP2	2	itgb5	4	ABLIM2	1	grip1	2	gapdhs	2
sephs1	1	HAPLN1	2	jarid2	4	CSK21	1	HMCS1	2	golga5	2
slc39a7	1	HDGFL3	2	NUCB2	4	elavl2	1	IPKA	2	GTDC1	2
tnikb	1	HOME3	2	PCDHGA2	4	LYSC	1	itpkb	2	IGF1R	2
ZN235	1	LOC103459553	2	rabgap1	4	MAPK9	1	KCAB1	2	KCNJ9	2
		lrrtm4l1	2	rpl14	4	MSTRG.8823	1	LBH	2	LOC103460906	2
		NPAS2	2	SATT	4	ntmt1	1	LOC103476863	2	mkl2	2
		NRP1A	2	scai	4	PP2BA	1	LORF2	2	nrg2	2
		NSG2	2	SCN5A	4	ppp1r13l	1	LRRF2	2	NTM1B	2
		PHF1	2	tbl1xr1	4	PTGR2	1	map6d1	2	polr1b	2
		ptprd	2	tmem145	4	thumpd2	1	mkl2	2	ranbp3	2
		st3gal4	2	tnrc6b	4	TRAPPC1	1	nfe2l1	2	RL19	2
		ube2v2	2	ZN865	4	TSP1L	1	nlgn2	2	rpl37a	2
		ALDH18A1	1	aars2	3	zmynd12	1	NPRL3	2	SEPTIN12	2
		APBA1	1	ache	3			pfkp	2	SFMBT1	2
		cfap99	1	AGRN	3			PKIA	2	sin3a	2
		CHD8	1	AMT	3			PRC1	2	TLE4	2
		cpeb4	1	ANK2	3			prickle2	2	TRI39	2
		EIF4G3	1	ASAP2	3			rexo1	2	abcg4	1
		GLSK	1	ATX1	3			rpl15	2	cdc42bpb	1
		GLUL	1	bdnf	3			SURF2	2	COR1A	1
		GNG7	1	bicd2	3			thap5	2	FACR1	1
		heatr5b	1	cacna1a	3			tnikb	2	gdpd2	1
		ITPR1	1	cdipt	3			TRI39	2	IL1A	1
		kiaa2026	1	COX18	3			TSPAN7	2	MRPL55	1
		LOC103464370	1	cpsf6	3			TUFT1	2	MSTRG.24943	1
		LOC103477566	1	csgalnact2	3			ZFP2	2	ncoa4	1
		NCAM2	1	DDX42	3			atp2b2	1	orc4	1
		NR1AA	1	dnmbp	3			CNGB1	1	PARVA	1
		nt5e	1	dtnb	3			DCNL1	1	pfkfb2	1
		orc5	1	fau	3			elavl4	1	pla1a	1
		ptprm	1	FLRT1	3			GBA2	1	PTGR2	1
		ranbp3	1	GRIA2	3			grk4	1	RHG22	1
		rpl8	1	HACE1	3			MSTRG.8823	1		
		RTase	1	HDAC4	3			MYO1E	1		
		SRSF5	1	HS6ST2	3			ran	1		
		TSC22D2	1	hypk	3			RL19	1		

vps45	1	KCD15	3	*rundc3a*	1
XPC	1	MAGI2	3	*SCHIP1*	1
		nell1	3	*shroom2*	1
		NTRK2	3	*SYPH*	1
		NYNRIN	3	*tanc2*	1
		P85B	3	*znf512*	1
		PMM2	3		
		psmc3	3		
		RAC3	3		
		RADIL	3		
		RL19	3		
		rpl10	3		
		rpl11	3		
		rpl30	3		
		rpl7a	3		
		rps3a	3		
		SCAM5	3		
		slc1a1	3		
		SPAG9	3		
		sptbn4	3		
		SYGP1	3		
		TACC1	3		
		tmem19	3		
		TRPC4AP	3		
		UBA3	3		
		USP9X	3		
		wdr18	3		
		zcchc6	3		
		zyg11b	3		
		AN13C	2		
		AN32E	2		
		atg3	2		
		B3GL2	2		
		CHD5	2		
		csmd2	2		
		EF1A1	2		
		EVL	2		
		GNAQ	2		
		golga5	2		
		GRM5	2		
		KCNH8	2		
		kdm4a	2		
		LAP2	2		
		LOC103460099	2		
		LOC103467299	2		
		lrrc75a	2		
		LRRT2	2		
		marcksl1	2		
		ncln	2		
		NLGN1	2		
		NOVA1	2		
		NTL9	2		
		PAK1IP1	2		
		pfdn6	2		
		PHLPP2	2		
		ppargc1b	2		
		PPP2R5C	2		
		prkca	2		
		RL12	2		
		rpl31	2		
		rplp0	2		
		S4A11	2		
		SCN8A	2		
		SI1L1	2		
		slc1a2	2		
		slc6a11	2		
		spen	2		
		SPTBN4	2		
		TESK2	2		
		TMTC3	2		

	TRPM6	2
	uba52	2
	UBC2	2
	UNC13A	2
	XPO1	2
	ZN665	2
	ZNF521	2
	znf839	2
	AHNK	1
	BMTI6	1
	BSN	1
	ctsa	1
	elavl2	1
	GP155	1
	GPR37	1
	hspd1	1
	KCAB1	1
	LOC103460110	1
	LOC103460981	1
	lrig1	1
	lrrc8a	1
	MYT1	1
	ndufa11	1
	PLXA2	1
	ptprb	1
	radil	1
	SHSA6	1
	snrk	1
	snrpa	1
	SNTG1	1
	tnikb	1
	TRI46	1

Table S9. Summary of genes found to be differentially co-expressed and differentially expressed in our transcriptomic analyses

COMMON DE & DC GENES					
OPTIC TECTUM		TELENCEPHALON		MIDBRAIN	
CONTROL	POLARIZATION	CONTROL	POLARIZATION	CONTROL	POLARIZATION
3	0	2	4	3	0
SEPHS1 *ZN235* *ARL4A*		*NCLN* *PTPRB*	*LRRC24* *PTPRS* *KHDR2* *PP2BA*	*IGF1R* *ITPKB* *TANC2*	

Table S10. KEGG pathways of interest associated in BFDCA differentially coexpressed gene pairs. Only showing pathways and select genes relevant to neural processes associated with behaviour that presented a FDR corrected p-value <0.05

OPTIC TECTUM

Polarization-selection lines

Pathways of interest (enriched)	Genes of interest
Glutamatergic synapse	GLUL,GNG7,ITPR1
Long-term depression	GRID2,ITPR1
Dopaminergic synapse	GNG7,ITPR1
GABAergic synapse	GLUL,GNG7
Cholinergic synapse	GNG7,ITPR1
Serotonergic synapse	GNG7,ITPR1

Control lines

Signal Transduction of S1P Receptor	MAPK6

TELENCEPHALON

Polarization-selection lines

Pathways of interest (enriched)	Genes of interest
FoxO signaling pathway	GRM1,MAPK9
Phospholipase D signaling pathway	GRM1,RHEB
Synaptic vesicle cycle	SLC17A6,RAB3A
Glutamatergic synapse	SLC17A6,GRM1
JNK (c-Jun kinases) phosphorylation and activation	MAPK9
mediated by activated human TAK1	RAB3A
GABA synthesis, release, reuptake and degradation	RAB3A
Glutamate Neurotransmitter Release Cycle	

Control lines

Glutamatergic synapse	CACNA1A,GRIA2,SLC1A1,GNAQ,GRM5,PRKCA,SLC1A2
Synaptic vesicle cycle	CACNA1A,SLC1A1,SLC1A2,SLC6A11,UNC13A
PI3K-Akt signaling pathway	PKN1,ITGB5,BDNF,MAGI2,NTRK2,PHLPP2,PPP2R5C,PRKCA
Long-term depression	CACNA1A,GRIA2,GNAQ,PRKCA
Dopaminergic synapse	CACNA1A,GRIA2,GNAQ,PPP2R5C,PRKCA
Long-term potentiation	GRIA2,GNAQ,GRM5,PRKCA
Cholinergic synapse	ACHE,CACNA1A,GNAQ,PRKCA
Ras signaling pathway	HTR7,BDNF,NTRK2,RAC3,PRKCA

MIDBRAIN

Polarization-selection lines

EGFR tyrosine kinase inhibitor resistance	PDGFRB,IGF1R,NRG2
Ras signaling pathway	PDGFRB,IGF1R,PLA1A
MAPK signaling pathway	PDGFRB,IGF1R,IL1A
AMPK signaling pathway	IGF1R,PFKFB2

Control lines

cAMP signaling pathway	PPARA,GRIA2,ATP2B2,CNGB1
Rap1 signaling pathway	MAGI1,IGF1R,RASSF5
Long-term depression	IGF1R,GRIA2
AMPK signaling pathway	IGF1R,PFKP
cGMP-PKG signaling pathway	ATP2B2,CNGB1

| Glutamate binding, activation of AMPA receptors and synaptic plasticity | GRIA2,GRIP1,TSPAN7 |

Table S10: Gene Ontology (GO) term of interest associated BFDCA DC gene pairs

OPTIC TECTUM	
Polarization-selected lines	
GO terms of interest (enriched)	Genes of interest
response to stimulus signaling response to stress signal transduction cell-cell signaling response to external stimulus synapse postsynaptic density asymmetric synapse postsynaptic specialization postsynapse	TLK2,RTN4,PARVA,PFKFB2,SHC3,TOM1L2,ASIC1,CDH6,GRID2,TSPAN8,CDK14,FOXP2,HDGFL3,NPAS2,NSG2,PHF1,PTPRD,ST3GAL4,UBE2V2,CHD8,CPEB4,GLUL,GNG7,ITPR1,NT5E,PTPRM,SRSF5,TSC22D2,VPS45,XPC TLK2,RTN4,PFKFB2,SHC3,TOM1L2,ASIC1,CDH6,GRID2,CDK14,HDGFL3,NSG2,PTPRD,APBA1,CHD8,CPEB4,GLUL,GNG7,ITPR1,NT5E,PTPRM,XPC TLK2,RTN4,ASIC1,TSPAN8,NPAS2,PHF1,ST3GAL4,UBE2V2,CPEB4,GLUL,ITPR1,NT5E,TSC22D2,VPS45,XPC TLK2,RTN4,SHC3,TOM1L2,ASIC1,CDH6,GRID2,CDK14,HDGFL3,NSG2,PTPRD,CHD8,CPEB4,GNG7,ITPR1,NT5E,PTPRM,XPC PFKFB2,ASIC1,GRID2,CDK14,PTPRD,APBA1,CHD8,GLUL,ITPR1 PARVA,SHC3,GRID2,TSPAN8,ST3GAL4,CHD8,CPEB4,GLUL,NT5E,PTPRM,XPC RTN4,ASIC1,GRID2,KLHL17,HAPLN1,PTPRD,APBA1,CPEB4,GLUL,ITPR1,RPL8,VPS45 RTN4,GRID2,KLHL17,CPEB4,ITPR1,RPL8 RTN4,GRID2,KLHL17,CPEB4,ITPR1,RPL8 RTN4,GRID2,KLHL17,CPEB4,ITPR1,RPL8 RTN4,GRID2,KLHL17,APBA1,CPEB4,ITPR1,RPL8
Control lines	
response to stimulus	CGAS,ABCA1,CENPS,STIMATE,BCL11B,DLGAP2,ENAH,MAPK6,TMBIM6,ADH5
positive regulation of cGMP-mediated signaling	CGAS
neurogenesis	BCL11B,ENAH,GBA2,MAPK6
response to L-glutamate	TMBIM6
response to external stimulus	CGAS,ABCA1,BCL11B,ENAH,ADH5
regulation of signal transduction	CGAS,ABCA1,STIMATE,DLGAP2,TMBIM6
neurotransmitter reuptake	SLC29A2
TELENCEPHALON	
Polarization-selected lines	

L-glutamate transmembrane transporter activity	SLC17A6, GRM1
neurotransmitter transmembrane transporter activity	SLC17A6
	GRM1
glutamate receptor activity	GRM1
postsynaptic neurotransmitter receptor activity	SLC17A6, MACF1, RAB3A,
L-glutamate transmembrane transport	RAB3A, MACF1
evoked neurotransmitter secretion	MAPK9
protein localization to tricellular tight junction	MAPK9
JUN phosphorylation	PTPRS
regulation of cell junction assembly	SLC17A6
neurotransmitter loading into synaptic vesicle	GRM1
adenylate cyclase-inhibiting G protein-coupled glutamate receptor signaling pathway	SLC17A6
neurotransmitter transport	GRM1, MACF1
L-glutamate import across plasma membrane	GRM1
G protein-coupled glutamate receptor signaling pathway	MAPK9
protein localization to cell-cell junction	PTPRS, TENM2, PTPRS, GRM1, MACF1, PNN, RAB3A, RHEB, RPL7, PPP1R13L
cell junction assembly	MTHFS
	GRM1
glutamate metabolic process	
glutamate receptor signaling pathway	
clathrin-sculpted glutamate transport vesicle membrane	RAB3A
	RAB3A
clathrin-sculpted glutamate transport vesicle	PNN
exon-exon junction complex	PTPRS
glutamatergic synapse	

Control lines

high-affinity glutamate transmembrane transporter activity	SLC1A1,SLC1A2
	SLC1A1,SLC1A2
glutamate:sodium symporter activity	KAT7,PKN1,SMARCA4,DVL1,GATAD2B,IER2,JARI
aromatic compound biosynthetic process	D2,SCAI,TBL1XR1,TNRC6B,AGRN,HDAC4,PMM2,P SMC3,RPL10,TACC1,UBA3,USP9X,CHD5,KDM4A,P PARGC1B,SPEN,UBA52,XPO1,ZNF521,ELAVL2,GP R37,HSPD1,MYT1
synapse organization	DVL1,ACHE,AGRN,BDNF,NTRK2,SLC1A1,GRM5,N
glutamate receptor signaling pathway	LGN1,UNC13A,BSN
neurotransmitter transport	GRIA2,SLC1A1,GNAQ,GRM5
	DVL1,SLC1A1,NLGN1,SLC1A2,SLC6A11,UNC13A

MIDBRAIN

Polarization-selected lines

protein tyrosine kinase activity	PDGFRB,IGF1R
cellular aromatic compound metabolic process	CPSF3,POU4F1,HES6,NUP205,OTP,PDGFRB,CTPS1, GAPDHS,MRTFB,POLR1B,RPL19,RPL37A,SFMBT1,
aromatic compound biosynthetic process	SIN3A,TLE4,IL1A,NCOA4,ORC4,PFKFB2
	CPSF3,POU4F1,HES6,OTP,PDGFRB,CTPS1,MRTFB,
inactivation of MAPKK activity	POLR1B,SFMBT1,SIN3A,TLE4,IL1A,NCOA4
signal transduction	IGF1R
	POU4F1,PDGFRB,ARL6,CORO2A,IGF1R,NRG2,TLE
equilibrioception	4,TRIM39,CDC42BPB,IL1A,NCOA4
regulation of signal transduction	POU4F1
suckling behavior	POU4F1,PDGFRB,IGF1R,NRG2,TLE4,TRIM39,IL1A
	POU4F1
Control lines	
signaling receptor activity	CSK,IGF1R,PPARA,GRIA2,NLGN2
signal transduction	MAGI1,CSK,IGF1R,PPARA,HMGA2,RASSF5,DCTN1
behavior	,DOCK9,GRIA2,GRIP1,ITPKB,LBH,NLGN2,NPRL3,P
glutamatergic synapse	RICKLE2,TUFT1,CNGB1,GRK4,MYO1E,RUNDC3A,
AMPA glutamate receptor complex	SCHIP1
	PPARA,DCTN1,NLGN2,ELAVL4
	GRIP1,ATP2B2,ELAVL4
	GRIA2

a Optic tectum **b** Telencephalon **c** Midbrain

Polarization
Group
Alone

Control
Group
Alone

Expression level
High
Low

DE genes

DC genes