Developmental Glucocorticoid exposure in zebrafish alters adult behavior and primes

differential expression of genes linked to human psychiatric disorders

Min-Kyeung Choi[1], Alex Cook[2], Helen Eachus[1], Anna Tochwin[1], Sara Kuntz[2], Siyong Kim[2],

Soojin Ryu*[1,2]

[1] Living Systems Institute, Faculty of Health and Life Sciences, University of Exeter, Stocker

Road, Exeter, EX4 4QD, United Kingdom

[2] University Medical Center, Johannes Gutenberg University Mainz, Duesbergweg 6, 55128

Mainz, Germany

* Correspondence: s.ryu@exeter.ac.uk

Running Title: Zebrafish Glucocorticoid primed genes

Abstract

Chronic or severe stress during development increases psychiatric disease susceptibility. One

putative mechanism involves priming of stress-sensitive gene networks by early life

Glucocorticoids (GCs) exposure, which modifies responses to stress later in life. However,

brain-wide GC-primed molecular alterations are currently unknown. Here, we used an

optogenetic zebrafish model to increase endogenous GCs during development and identified

comprehensive transcriptomic alterations in the whole brain of GC-exposed fish across the

life course. When exposed to acute stress in adulthood, developmental GC-exposed animals

show highly exaggerated and pervasive brain-wide transcriptional changes which include

gene sets associated with axon development, neuronal signaling, and epigenetic modulators.

Strikingly, these altered gene sets showed an overrepresentation of human psychiatric

disorder risk genes. We identify in developmental GC-exposed individuals: 1) altered

biological processes, 2) hitherto uncharacterized novel GC-primed genes, 3) altered adult

behaviors, and 4) subsets of primed genes and putative transcriptional regulators associated

with changes in adult social behavior including oxytocin and *myt1la*. Thus, our study

establishes a new translationally relevant zebrafish model and suggests potential molecular

mechanisms by which developmental GC-exposure exerts its life-long impact.

Introduction

Early life stress (ELS) can lead to long-term alterations in neural plasticity, functional

connectivity, epigenetic dynamics, and behavior[1-4]. Reflecting these far-reaching effects, ELS

is a key factor affecting healthy development and aging as well as susceptibility to

psychiatric disorders in humans[5-8]. The molecular response to stress in vertebrates is

primarily mediated by a highly conserved neuroendocrine stress regulatory system called the

Hypothalamo-Pituitary-Adrenal axis (HPA-axis) in humans[9] and the Hypothalamic-Pituitary-

Interrenal axis (HPI-axis) in fish[10]. Like the mammalian adrenal gland, the fish interrenal

gland is composed of two cell types: aminergic chromaffin cells and steroidogenic interrenal

cells, the latter of which produce glucocorticoids (GCs), effector stress hormones that are

highly conserved across the vertebrate kingdom[9]. While GCs have pleiotropic effects on

many aspects of animal physiology, exposure to elevated GCs during brain development is

known to cause long-term alteration of behavior, physiology, and stress regulation[1]. However,

how high GC exposure during development predisposes individuals to adulthood dysfunction

is poorly understood. One potential mechanism is long-lasting epigenetic changes, which

modify how animals respond to stress in adulthood. For example, altered DNA methylation at

the glucocorticoid receptor (GR) gene promoter by poor maternal care is associated with
adulthood alterations in histone acetylation, DNA methylation, transcription factor binding,
GR expression, and HPA response to stress[11]. Moreover, a recent study identified
differentially expressed genes (DEGs) "primed" by ELS in specific regions of ELS-exposed
mouse brains[12]. These genes showed enhanced differential expression in adulthood upon re-
exposure to stress and may be associated with behavioral changes in adulthood. Another
study identified GC-primed molecular alterations using a GC (dexamethasone)-exposed
human hippocampal progenitor cell line and identified long-lasting DNA methylation
alterations[13]. Lastly, genetic differences in the immediate transcriptome response to GR
activation are shown to predict risk for depression and co-heritable psychiatric diseases[14].
Together, the above studies establish that ELS or developmental GC exposure results in
distinct responses to stress later in life by priming and maintaining longer-term alteration of
specific genes which facilitate increased psychiatric disease susceptibility. However, so far,
ELS or GC-primed genes have only been identified in specific regions of the brain including
reward circuitry or cell lines, and brain-wide GC-primed molecular alterations have not been
investigated.
In this study, we utilized optogenetic manipulation of steroidogenic interrenal cells in a
zebrafish model to increase the endogenous GC level during larval and juvenile stages.
Although the mammalian counterpart of the fish interrenal, the adrenal gland, is known to
produce three types of steroid hormones, GCs, mineralocorticoids (MCs), and androgens[15],
MCs and androgens are not detectable in larval zebrafish interrenal gland[16-19]. Therefore, our
manipulation is suitable to assess specific effects of elevating endogenous GCs. As a first
step in determining the effects of developmental GC exposure, the zebrafish were exposed to
high levels of GCs throughout development starting from the postembryonic stage.
Here we report brain-wide transcriptomic alteration of GC-exposed fish across the life course.
Strikingly, when subjected to acute stress in adulthood, developmental GC-exposed fish show
highly exaggerated endocrine and transcriptional responses, indicating that adult stress is
processed differently depending on GC exposure history. Remarkably, the genes primed by
GC-exposure were enriched in gene sets associated with human neuropsychiatric disorders
indicating high levels of evolutionary conservation in how early GC exposure leads to adult
dysfunction. We further identified 1) biological processes altered by developmental GC
exposure, 2) adult behaviors altered by developmental GC exposure, 3) hitherto
uncharacterized novel GC-primed genes, and 4) subsets of primed genes and putative

transcriptional regulators associated with changes in adult social behavior induced by

developmental GC exposure including oxytocin and *myt1la*.

Methods and materials

88 The detailed protocol for all the procedures used in this study is available as step-wise online

89 methods[20].

Zebrafish husbandry and maintenance

The tübingen (TU) strain was kept at 28☐ on a 12:12 hour light/dark cycle and housed at a

maximum density of 5 fish/L. For experiments with adult fish, only females were used. All

animal procedures were carried out in compliance with the ethical guidelines of the national

animal welfare law and approved by relevant authorities (Landesuntersuchungsamt

Rheinland-Pfalz, Germany, Project number 23 177-07/G20-1-033 or UK Home Office PPL

number PEF291C4D).

Generation of transgenic line

100 Linearized pT2-2kbStAR:bPAC-2A-tdTomato[21] (ZFIN ID: ZDB-GENE-001120-3)

101 recombinant plasmids were injected into 1-cell stage wild-type embryos of TU strain at 10-50

102 ng/mL in the presence of 10-50 ng/mL Tol2 transpose mRNA and 0.05% phenol red. The

103 progenies of injected fish were screened for interrenal-specific tdTomato expression and the

104 stable transgenic line, *Tg(star:bPAC-2A-tdTomato)*[uex300] was established.

Sample collection

107 All samples were collected within a 2-hour window in the morning (08:30 to 10:30) without

108 feeding.

Whole-body Cortisol assay

111 The competitive cortisol assay (Cisbio HTRF® Cortisol Kit, 62CRTPEG) was performed

112 following the manufacturer's protocol. ELISA signal was detected by CLARIO star plate

113 reader (BMG Labtech, Ortenberg, Germany).

Behavior tests

Social behavior tests and acute stressor delivery using looming dot (LD) presentation were

carried out as described in Cook *et al.*[22]. Basal locomotion, food consumption and fear

conditioning assay were performed following the protocol in online methods[20]. For all

behavioral tests, female adult zebrafish (6-9 months old) were moved into a designated

behavior experimental room one week prior to testing and housed in groups of 10 fish in 3L

tanks. Experiments were carried out between 09:00 and 14:00 and fish were not fed in the

morning of testing. Fish were not used in any repeat experiments. The trials were recorded at

30 FPS. A camera (acA1300-60gm, Basler ace) was positioned at the front and/or top side of

the tank. The test fish was tracked using Ethovision (EthoVision XT 13, Noldus Information

Technology).

RNA preparation and quantification

Each sample for the larval stage (6 and 13 dpf) contained 25-30 larval whole brains whilst for

the adult stage (120 dpf) three brains constituted one sample. Transcripts were quantified

using real-time qPCR and next-generation mRNA sequencing. Primer information for qPCR

is described in Supplementary Table 1. A total of 60 samples was sequenced, consisting of 5

biological replicates at four different time points (6, 13, 120 dpf, and acute-stressed at 120

dpf) for each genotype (wild type, bPAC+ and bPAC-).

Immunohistochemistry

Immunohistochemistry was performed as described with minor modifications[23]. Custom-

made primary rabbit anti-oxt antibodies (1:200 dilution)[23] and Alexa 488 anti-rabbit

antibodies (A-11008, Invitrogen, USA) as secondary antibodies (1:1000 dilution) were used

for the visualization of oxytocinergic neurons.

Imaging and analysis

Confocal stacks through the larval head were recorded using a Zeiss LSM 880 with Airyscan

FAST with PApo 10X/0.45 – dry objective with Airyscan mode. Stacks were analyzed using

ImageJ (FIJI, 1.53c)[24]. Corrected Total Cell Fluorescence (CTCF) was calculated as

integrated density − (Area of cells × Mean fluorescence of background reading).

Statistics

Statistical analyses were performed using R and Prism 9 (Graphpad Software Inc, San Diego,

CA, USA). A detailed description of the statistical analyses is provided in online methods[20]

Data availability

All sequenced reads for RNA-seq were deposited in European Nucleotide Archive (ENA,
PRJEB53713).
154
155 **Code availability**
156 The detailed data processing and R scripts for bioinformatic analysis are available at the
157 GitHub repository (https://github.com/minkechoi/tx_star-bPAC_brain).
158
Results
Optogenetic elevation of endogenous GC during development
We have previously reported that elevation of endogenous cortisol (GC in fish) is induced in
a transgenic zebrafish line by blue light present in ambient light which activates *beggiatoa*
Photoactivated Adenylate Cyclase (bPAC), expressed specifically in steroidogenic interrenal
cells[21, 25]. For the present study, we generated a new transgenic line, *Tg(star:bPAC-2A-*
tdTomato)[uex300], using the same construct in a different strain to obtain a higher cortisol level
than what was previously feasible and raised the fish under ambient light (hitherto referred to
as bPAC+) (Fig. 1a,b). Indeed, we observed a strong cortisol increase at 5 days post
fertilization (dpf) in bPAC+ compared to bPAC-, siblings of bPAC+ which do not carry the
transgene (Fig. 1c). bPAC+ larvae raised under a filter, which restricts blue light penetration,
did not show a cortisol increase, supporting that the cortisol elevation was due to optogenetic
activation of bPAC (Fig. 1c). Moreover, the expression of *fkbp5*, which regulates the
responsiveness of the glucocorticoid receptor (GR)[26], was increased in bPAC+ brains
compared to bPAC- and to wild type, confirming the activation of the GR signaling cascade
(Fig. 1d). Although bPAC- fish did not carry the transgene, since transgenerational
inheritance of stress-induced phenotypes is reported in mammals and fish[27-29], we compared
bPAC+ and bPAC- to each other, as well as to wild type in this study. Consistent with the
reported onset of endogenous *star* gene expression[30], an increased cortisol level was observed
starting around 2 dpf (Fig. 1e). The high cortisol level was sustained during larval and
juvenile stages in bPAC+ fish. However, by adulthood (three months post-fertilization (mpf)),
no significant difference was detected between bPAC+ and wild type (Figs. 1e,f). These
results indicate that by using bPAC+ we can achieve a robust and persistent increase in
endogenous cortisol levels throughout development.
183
Alteration of adult behavior in developmental GC-exposed fish

We assessed whether bPAC+ fish exhibited behavioral changes as adults using a set of tests

aimed at assessing behaviors commonly affected in neuropsychiatric diseases[31]. We first

established that bPAC+ fish did not show significant differences in basal locomotor activity

compared to wild-type and bPAC- fish (Figs. 2a,b). We then focused on social behavior as

social processes are translatable across multiple species and are often perturbed in a range of

disorders associated with ELS[32]. To assess social behavior, we employed a new test that we

recently established, which allows the assessment of distinct phases of social behavior,

namely a drive to approach a social cue, and an ability to engage and maintain social

interaction (Fig. 2c)[22]. In this test, after the fish was placed in a test tank and before the

presentation of conspecifics, wild type, bPAC- and bPAC+ all displayed preference for the

blue-colored bottom zone with no significant genotype differences (Figs. 2d,e). When the

social cue was revealed, within 10 seconds the vast majority of wild type, bPAC-, and

bPAC+ entered the social interaction zone (shaded green in Fig. 2f) adjacent to the

conspecifics, within which wild types display a stereotypical C-shaped pattern of swimming

(Fig. 2h). No difference in social approach between the groups was found using a Log-rank

test (P=0.77), indicating that bPAC+ did not show deficits in recognition nor approach to

social cues (Fig. 2g). In contrast, we found significant differences between wild type and

bPAC+ during the 2-minute social phase in the time spent in the interaction zone and the

number of social interactions performed (Figs. 2h-j). bPAC- also spent less time in the

interaction zone and performed fewer interactions than the wild type (Figs. 2i,j).

Next, we tested food consummatory behavior by counting the number of floating food pellets

eaten by a fish during a 10-minute interval (Supplementary Fig. 1a). Wild-type adult fish

consumed on average 14.5 pellets while bPAC+ fish consumed on average 2.33 pellets out of

a total of 25 (Supplementary Figs. 1b,c). This was not due to the lack of recognition nor

interest in the food, as bPAC+ fish approached the food pellets and actually spent a longer

time in the zone near the pellets (feeding zone) than wild types (Supplementary Fig. 1d).

To assess cognitive processes, we tested associative learning using a Pavlovian avoidance

learning test where fish learn to associate the color green with a mild electric shock in the

testing chamber (Supplementary Figs. 1e,f). Fish were first habituated in the chamber for 30

minutes and tracked to determine the baseline preference for green or blue color. Wild-type

and bPAC+ fish showed no significant differences in baseline color preference

(Supplementary Fig. 1g). However, after the conditioning, wild-type fish spent on average

89.2% of their time in the blue zone, significantly more than bPAC+ fish who spent on

average 62.68% of their time in the blue zone (Supplementary Fig. 1h), suggesting a deficit

in associative learning in bPAC+ fish.

Lastly, we tested the behavior of bPAC+ fish in a novel environment using a novel tank test

(Supplementary Figs. 1i-m). In the novel tank test, wild-type fish first dive to the bottom and

demonstrate a sustained depth-preference whilst performing bouts of erratic high-speed

swimming[33, 34]. In contrast, bPAC+ fish showed reduced depth-preference (Supplementary

Fig. 1j) and generally swam slower than wild-type controls (Supplementary Fig. 1k). Time

spent performing fast-swimming bouts was also significantly decreased in bPAC+ fish

(Supplementary Fig. 1l). However, there was no significant difference in freezing duration

between groups (Supplementary Fig. 1m).

Together these results show that social interaction, food consumption, associative learning,

and adaptation to a novel environment are altered in bPAC+ fish compared to wild types. Our

results point to modification of a broad range of goal-directed behavior performance without

changes in basal locomotion parameters in bPAC+.

Whole-brain transcriptional alterations in bPAC+ fish across the life course

Whole-brain RNA sequencing was carried out using larval (6 and 13 dpf) and adult (120 dpf)

brains to identify and compare transcriptomic differences in wild type, bPAC+, and bPAC-.

A large number of DEGs was identified in pairwise comparisons of bPAC+ and wild-type

brains at 6 dpf (up/down=602/1253), 13 dpf (up/down=777/1566), and 120 dpf

(up/down=684/1078) (Fig. 3, Supplementary Table 2). At all-time points, the number of

downregulated genes was greater than the number of upregulated genes, indicating that

developmental GC-exposure tends to repress rather than activate genes (Fig. 3a). Further, the

number of DEGs consistently downregulated from early life until adulthood (273 DEGs) was

greater than the upregulated DEGs (48 DEGs) (Fig. 3a). We identified fewer DEGs when

comparing bPAC- to wild type, with 571 (up/down=329/242), 1026 (up/down=372/654), and

74 (up/down=42/32) DEGs at 6, 13, and 120 dpf, respectively, and many of these DEGs

overlapped with those identified in the bPAC+/wild type comparison (Supplementary Fig.

2a). To compare the overall transcriptomic landscape of all identified DEGs, we performed

unsupervised hierarchical clustering, which revealed large differences in the gene expression

landscape between the bPAC+ and bPAC- compared to wild type (Fig. 3b). At each time

point, bPAC+ and bPAC- transcriptome profiles clustered together separately from those of

wild type. Next, to confirm an increase in GR signaling, we measured the expression level of

GR target genes, *nr3c1, nr3c2, fkbp5, tsc22d3,* and *hsd11b2* at 6, 13, and 120 dpf. The

expression levels of *nr3c1, fkbp5, tsc22d3,* and *hsd11b2* were altered at multiple time points

in bPAC+ (Fig. 3c). bPAC- also showed differences to wild type, albeit generally to a lesser

degree (Fig. 3c). Both the alteration of GR signaling and similarity of DEGs identified in

bPAC- and bPAC+ point to potential transgenerational effects present in bPAC- resulting

from high GC-exposure in the parental generation. Supporting this notion, even though

bPAC- was not exposed to a high level of GC, some of the DEGs at 6 dpf in bPAC- are

involved in processes that potentially regulate GC inactivation (Supplementary Fig. 2b,

Supplementary Table 3).

Developmental GC exposure leads to downregulation of transcripts involved in nervous

system development

To identify molecular consequences of developmental GC exposure, we characterized DEGs

in bPAC+ compared to the wild type. Upregulated DEGs in bPAC+ fish brains at 6 dpf were

enriched in the GO terms that include potential regulatory enzymes for GC such as

cytochrome monooxygenases, 11β-hydroxysteroid dehydrogenase or sulfotransferases[35-37]

(left panel of Fig. 3d). At 13 dpf, the top-ranked enriched biological processes for

upregulated DEGs were cell cycle, chromosome segregation, and DNA metabolic process

which can cause genomic instability by replication stress[38] (left panel of Fig. 3d). At 120 dpf,

bPAC+ brains showed upregulation of genes involved in ribonucleoprotein complex

biogenesis, peptide metabolic, translation, protein modification, RNA processing and more,

suggesting effects of developmental GC exposure on gene expression regulation and post-

transcriptional modification in adulthood (left panel of Fig. 3d, Supplementary Table 3). In

contrast, downregulated DEGs across all time points were mainly enriched in processes

related to cell adhesion and cell migration, as well as nervous system development including

neuron differentiation, cell morphogenesis, and axon development (right panel of Fig. 3d,

Supplementary Table 3). Brain-wide analyses using qPCR and immunohistochemistry

confirmed alteration of neurogenesis in larval bPAC+ fish. These results are reported in detail

in a separate report[39]. During larval stages, downregulated DEGs were also enriched in

anatomical structure formation, muscle structure development, vasculature development,

external encapsulating structure organization, and more. We also identified functions

associated with time-point-specific downregulated DEGs (Supplementary Table 3). For

example, DEGs at 6 dpf were enriched in brain development while DEGs at 13 dpf were

enriched in the cellular response to organic substance (right panel of Fig. 3d). At 120 dpf, we

identified downregulated DEG enrichments in synaptic signaling, regulation of molecular

function, and regulation of GTPase activity and more (right panel of Fig. 3d).

These results highlight long-lasting downregulation of neuronal development and signaling

processes, and adult-onset alteration of transcription and translation regulation following

developmental GC exposure.

bPAC+ fish show an enhanced endocrine response to acute stress in adulthood

To assess whether high GC exposure during development affects the stress response in

adulthood, we subjected wild-type, bPAC+, and bPAC- fish to a looming dot stimulus (LD),

which mimics an approaching predator[22] (Figs. 4a,b). Interestingly, bPAC+ adults exhibited a

much greater cortisol increase after LD exposure compared to wild-type adults (Fig. 4c),

indicative of a sensitized endocrine response to acute stress in bPAC+. Similarly, we found

an altered transcriptional response of stress-associated genes in bPAC+ after LD exposure

(Fig. 4d). For example, the expression of *nr3c2*, the gene coding for the mineralocorticoid

receptor (MR), was further downregulated only in bPAC+ adults following LD exposure,

although basal expression of *nr3c2* was already decreased in bPAC+ compared to wild type.

In contrast, the GC-inactivating enzyme coding gene, *hsd11b2,* was significantly upregulated

in the wild type after LD exposure, but upregulated to a lesser degree in bPAC+ following

LD exposure. Together, the significant increase in LD-induced cortisol level and altered

expression of *nr3c2*, *fkbp5,* and *hsd11b2* indicate an enhanced endocrine response following

acute stress exposure in adult bPAC+.

Developmental GC exposure primes large-scale differential gene expression following

acute stress in adulthood

To identify the transcriptional response to acute stress following developmental GC exposure,

we carried out whole-brain RNA sequencing of wild type, bPAC+ and bPAC- that had been

exposed to LD in adulthood and made three separate comparisons (Figs. 4e,f). First, we

compared the wild-type transcriptome at basal (pre-LD) and after LD exposure (post-LD), to

identify the normal acute stress response in wild-type fish (referred to as DEG1). Secondly,

we compared pre-LD and post-LD in bPAC+ (referred to as DEG2), and finally the post-LD

transcriptome of wild type and bPAC+ was also compared (referred to as DEG3). DEG3

includes LD-induced DEGs (overlapping with DEG1 and DEG2) as well as those DEGs that

were already altered in bPAC+ prior to LD exposure and therefore the number of DEG3

genes (2034 genes) was greater than that of DEG1 and DEG2 (Fig. 4g). Strikingly, bPAC+

showed more than 20-fold greater number of DEGs in response to LD exposure (DEG2, 1617 genes) compared to wild type (DEG1, 64 genes) (Fig. 4g). Altogether we identified 2515 genes that appeared to be altered in the adult brain by developmental GC exposure in bPAC+, and refer to them collectively as adult bPAC+ DEGs.

The exaggerated transcriptional LD response in bPAC+ indicates that developmental GC exposure in zebrafish primes enhanced alteration of gene expression in adulthood, similar to observations in previous mammalian ELS or GC-primed gene studies[12,13]. To define GC-primed genes, we took the entire DEG2 set and excluded 44 genes that overlapped with DEG1 as these represent normal acute stress-induced genes present also in the wild type. To this gene set, we combined the 18 genes in DEG1 that do not overlap with DEG2 as these genes are reflective of wild-type responses that are missing or altered in bPAC+ (Fig. 4g, Supplementary Table 4). This set of 1591 genes is collectively referred to as GC-primed genes here and represents those genes that are primed by developmental GC exposure to produce an altered transcriptional response to acute stress in adulthood. Among the 1591 GC-primed genes, 1337 downregulated genes were enriched in neuron projection development, synaptic signaling, receptor complex, synapse, metal ion transmembrane transport, and more (Fig. 4h, Supplementary Table 5).

To compare GC-primed genes in zebrafish with those identified in humans, we carried out a comparison with a recent study that reported 702 GC-primed DEGs linked with long-lasting differentially methylated sites following re-exposure to GC (dexamethasone)[13]. These DEGs were identified in GC-treated mature neurons which were exposed to GC during neurogenesis in a human hippocampal progenitor cell (hpc) line. Through the comparison between 702 DEGs (hpc_ELS-primed genes, up:368, down:334)[13] and our DEG3, which represents a similar GC re-exposed condition, we found 163 overlapping genes (up/down=99/64).

Together, these results reveal for the first time a brain-wide set of genes primed by developmental GC exposure. Further, we found a significant overlap between subsets of GC-primed genes identified in zebrafish with those from mammalian studies, suggesting a high degree of evolutionary conservation of molecular mechanisms underlying reprogramming by GC.

Adult bPAC+ DEGs are associated with neuropsychiatric disorders in humans

Since stress exposure during development is a risk factor for the development of psychiatric disorders in humans[6-8], GC-primed genes were compared with known psychiatric disease risk

factors using the curated DisGeNET database[40]. Strikingly, human homologs of GC-primed genes exhibit a significant enrichment of known psychiatric disease-associated genes (Supplementary Table 8). Top-ranked psychiatric disorders were schizophrenia, bipolar disorder, and autistic disorder (Supplementary Fig. 4). Further, the entire adult bPAC+ DEG set was strongly enriched in known psychiatric disease risk factors and in this case depressive disorder was also included in the top four ranked diseases (Figs. 5a,b, Supplementary Table 9). Upon LD exposure, the difference in expression of schizophrenia-associated genes in bPAC+ was greatly enhanced, as represented by the size of nodes in Fig. 5c. Further, schizophrenia-associated adult bPAC+ DEGs were enriched in either of two KEGG pathways: neuroactive ligand-receptor interaction (FDR=2.01e-08), or MAPK signaling pathway (FDR=9.74e-08). Overall, across all four disease categories, the top-ranked disease-associated adult bPAC+ DEGs were mainly enriched in the downregulation of synaptic signaling, and multicellular organismal process (Fig. 5d).

Adult bPAC+ DEGs associated with schizophrenia, bipolar disorder, depressive disorders, and autistic disorders were predominantly downregulated following LD exposure (Fig. 5e). Interestingly, we noted that some of the disease-associated adult bPAC+ DEGs were already differentially expressed from the larval stage (trans-temporal, 60 genes) while some others showed alteration in their expression only upon exposure to acute stress in adulthood (adult-onset, 65 genes) (Fig. 5f). Other non-disease-associated adult bPAC+ DEGs maintained differential expression from the larval stage (540 genes) or exhibited an enhanced alteration upon adulthood stress (adult-onset, 463 genes) (Supplementary Table 10). Trans-temporal adult bPAC+ DEGs were mostly enriched in axon development (FDR=1.33e-08, 31 DEGs), whilst adult-onset adult bPAC+ DEGs were enriched in ion transport (FDR=7.26eE-05, 44 DEGs) (Supplementary Table 11). Interestingly, a cell-type enrichment test using the human homologs of adult bPAC+ DEGs showed that 24 trans-temporal adult bPAC+ DEGs were associated with non-microglial oligodendrocyte lineages. In contrast, the downregulated adult-onset adult bPAC+ DEG set was associated with non-microglial neuron-enriched genes including GABAergic receptor-associated genes (*GABRA1*, *GABRG2*), other ion channel activity genes (*SLC12A5*, *CACNA1E*, *CACNA1I*, *UNC80*, *KCNH1*, *SCN2B*), and genes associated with synapse and neuronal cell signaling (*DNM1*, *DLGAP2*, *RPH3A*, *GPR158*) (Supplementary Table 12). More than half of those genes were GC-primed genes. These results indicate that oligodendrocyte lineage cells are consistently suppressed following developmental GC-exposure, and developmental GC primes suppression of neuronal signaling primarily in adulthood when encountering stressful stimuli.

Altered gene expression associated with epigenetic modifications and RNA processing in bPAC+ fish

To identify underlying mechanisms by which developmental GC exposure regulates gene expression in the brain, we analyzed the expression of epigenetic modifiers. We identified 70 adult bPAC+ DEGs associated with epigenetic modifications (Supplementary Figs. 5 a,b, Supplementary Table 13). These genes are associated with RNA modification (up/down=3/3), RNA processing including mRNA splicing (up/down=27/17), histone modification (up/down=4/18), and DNA modification (up/down=0/3). Among them, the DNA methyltransferase (DNMT) 3aa gene and the histone modification-associated genes, *chd3*, *hdac4*, *hdac5*, *kdm2ba*, *kdm4aa*, *mta3*, *nsd3*, *padi2*, *suv39h1b*, and *yeats2,* showed consistent downregulation across time-points.

We also found significant differences in the expression of 4 out of 7 *DNMTs* in the zebrafish genome including *dnmt1*, *dnmt3aa*, *dnmt3ab,* and *dnmt3bb.3* across time points and conditions in bPAC+ compared to the wild type (Supplementary Fig. 5c). In the wild type, *dnmt1* is highly expressed at 6 dpf and expression then gradually decreases over time, but in bPAC+ expression was significantly reduced at 6 dpf. Expression patterns of *dnmt3aa* and *dnmt3ab*, orthologous to human *DNMT3A,* known as one of the writers of *de novo* methylation, mainly showed differences in bPAC+ adult brains. Further, low expression of *dnmt3aa* and upregulation of *dnmt3ab* was observed upon LD exposure in adulthood in bPAC+. To determine whether alteration of the most differentially regulated epigenetic modulators (labeled with * in Supplementary Fig. 5a) lasts throughout adulthood, we analyzed their expression at 8 mpf, with and without LD exposure. Downregulated expression of those genes was continuously observed in the later adult stage (8 mpf) mainly upon acute stress in bPAC+ adults (Supplementary Fig. 5d). These results suggest that developmental GC exposure leads to pervasive, long-lasting changes to epigenetic modulators.

Social behavior-associated genes are a part of the GC-primed genes

To search for a potential link between the observed molecular changes and social behavior, we analyzed the expression patterns of 101 known social behavior-associated genes from the Gene Ontology Annotation database (henceforth referred to as social genes). Interestingly, 54 of these social genes were differentially expressed in bPAC+ brains, where 44 were downregulated at least at a single time point ($|FC|>1.5$ and FDR<0.05) and only 11 were

upregulated (*npas4a* was double counted because downregulated at 6 dpf and upregulated at 13 dpf, Fig. 6a, upper panel). In the basal condition, 6 dpf brains showed the largest change (up/down=2/31). The most significant downregulated social genes over time were *nlgn2b*, *mapk8ip2*, *brinp1*, *slc6a4a,* as well as oxytocin (*oxt*), a well-known social gene[41] (Fig. 6a). bPAC- showed a similar trend in the downregulated gene expression pattern following LD exposure as in bPAC+, albeit the level of difference was lower though. (Fig. 6a, bottom panel). We confirmed the downregulation of *oxt* both in larval and adult bPAC+ brains using qPCR (Fig. 6b). Downregulated *oxt* expression and the number of oxt+ cells in larval brains were also confirmed using immunohistochemistry which showed preferential reduction of the posterior cluster of oxt+ cells which is likely to be similar to mammalian parvocellular oxt+ neurons based on their projection[42, 43]. (Figs. 6c-e).

Using differentially expressed social genes at 6 dpf, we performed network-based analyses using the String database[44] (Fig. 6f). The identified protein network was enriched in modulation of chemical synaptic transmission, synaptic signaling, learning and memory, pre-synapse assembly, and social behavior. Notably, *oxt* regulators including *myt1la*, *med12*, and *thrab* were downregulated at 6 dpf and two of them (*myt1la*, *med12*) remained downregulated until adulthood (Fig. 6g). Further, transcription factor (TF) enrichment analysis of human orthologues of the social genes identified *myt1la* as one of the top regulators of 47 differentially expressed social genes, based on the ChEA3 database[45] (Fig. 6h). These results suggest that downregulation of *myt1la* in bPAC+ may result in the altered expression of social genes including *oxt*[46].

Strikingly, a subset of social genes was also part of the GC-primed genes indicating that the regulation of their adult expression is highly sensitive to the history of developmental GC exposure (red label in Fig. 6a). Next, we performed a TF enrichment test of GC-primed genes using the ChEA3 platform[45] and identified common potential upstream regulators of GC-primed genes (Figs. 6i,j, Supplementary Table 6). For the upregulated GC-primed genes, immediate early genes were top-ranked TFs. Among the top-ranked TFs for downregulated GC-primed genes, *camta2* and *myt1l* may be involved in the regulatory process of 149 and 139 downregulated genes, respectively. Expression of *myt1l* was consistently downregulated in bPAC+ adults at both pre-LD and post-LD condition (Fig. 6k). Together, these results suggest developmental GC-induced regulation of *myt1la* as one potential mechanism by which stress in early life alters adult social behavior and primes expression of brain-wide gene networks.

Discussion

This study examines the effects of chronic developmental GC exposure in shaping adult phenotypes at both behavioral and molecular levels. We report altered goal-directed behaviors in adults and molecular consequences of GC-exposure in both larvae and adults. Further, we report a comprehensive identification of genes differentially expressed in adulthood following developmental GC exposure, many of which correspond to genes associated with neuropsychiatric disorders in humans. These findings suggest that developmental GC exposure primes differences in GR (acute stress)-induced transcriptional activation in adulthood and may increase the risk for neuropsychiatric disorders by affecting a network of stress-sensitive genes.

Our bPAC+ model involves the optogenetic activation of *star*-expressing interrenal cells, leading to an increase in endogenous cortisol levels in transgenic animals. Previous studies in fish using LC-MS have not detected MCs in any significant quantities in either larval nor adult samples[16-19], suggesting that the zebrafish interrenal produces no/negligible amounts of MC. Similarly, androgens are not detectable in zebrafish larvae[16], but are detectable in adult male whole-body samples [17-19], suggesting that androgens are only produced in the testes in zebrafish. As such, the phenotypes observed in our bPAC+ model are primarily due to the effects of GC and its downstream effectors, rather than due to MC or androgen over-exposure. The increase in cortisol in our model is observed from 2 dpf onwards, consistent with the onset of *star* gene expression. 2 dpf is considered to represent an early postembryonic development stage comparable to fetal and neonatal stages in mammals[47, 48]. As zebrafish develop externally, it is not possible to distinguish pre- vs. post-natal stress exposure. Elevated cortisol level is maintained until 2 months of development and decreases with the onset of adulthood. This may be due to the decrease in blue-light penetration following the reduced transparency of the skin as fish grows.

We report enhanced endocrine sensitivity of adult bPAC+ fish in response to acute stress. Further, we observed a significant decrease in the expression of *nr3c2,* which encodes the high GC-affinity MC receptor (MR), both at pre- and post-LD exposure. Given the higher affinity of MR to GC, MR can be a regulator of GR signaling in MR and GR co-expressing cells in basal conditions[49,50]. Moreover, reduced transcriptional reactivity of *hsd11b2,* a key enzyme for cortisol inactivation[51,52], in bPAC+ after LD may lead to an increased accumulation of active cellular cortisol. These results suggest that developmental GC-

exposure may lead to a sensitized response to stress by decreasing MR expression at pre- and

post-LD and attenuating the increase of *hsd11b2* expression.

We identified brain-wide alterations in bPAC+ through whole-brain transcriptome analyses

using both larval and adult zebrafish. Downregulated genes in bPAC+ are mainly enriched in

cell morphogenesis, neuronal cellular signaling, and neuropsychiatric disorders-associated

gene sets. Despite the differences in species, stressor type, and brain region, the results of our

GO enrichment test and observed phenotypes are broadly similar to previous findings

obtained from ELS rodent models which report alterations in axon development, synaptic

density and plasticity, altered hippocampal mossy fibres[4, 5, 53]. However, we identified a

greater number of DEGs when compared to previously reported mammalian ELS paradigms

consistent with the fact that the model described here represents a stronger and longer stress

exposure.

Interestingly, we identified a large set of adult-onset upregulated DEGs which were enriched

in DNA metabolism, various RNA processing processes, and posttranslational modifications.

This may indicate that developmental GC-exposure can lead to genome instability by DNA

replication stress and altered gene expression regulation via alternative splicing, non-coding

RNAs, and posttranslational modifications in adulthood. Moreover, in adult bPAC+, we

identified oligodendrocyte lineage cell markers in downregulated trans-temporal DEGs while

neuron-enriched genes were enriched in downregulated adult-onset DEGs. This result

suggests that developmental GC-exposure has a cell-type-specific impact.

Identified adult bPAC+ DEGs were enriched in neuropsychiatric disorders when compared to

a publicly available catalogue of disease-associated genes based on gene expression network,

eQTLs and GWAS[40]. The majority of schizophrenia, bipolar disorder, depressive disorder,

and autistic disorder-associated adult bPAC+ DEGs showed an exaggerated transcriptional

response to stress in adult bPAC+ brains. Those DEGs were overrepresented in synaptic

signaling processes including genes for GABA receptors (*gabrg2*, *gabbr1a*, *gabbr1b*, *gabbr2*,

gabra1, *gabra2a*, *gabra4*, *gabra5*, *gabrg1*, *gabrg2*), glutamate receptors and transporter

(*grm1b*, *grm3*, grm5b, *slc17a7a*), dopamine receptors and transporter (*drd1b*, *drd2a*, *drd2b*,

slc6a3), serotonin receptor (*htr1b*), cholinergic receptor (*chrm2a*), opioid receptor (*pnoca*),

and neurexins (*nlgn1*, *nlgn2b*, *nlgn4xb*). In a human study, genetic variants associated with

GR-mediated immediate transcript response were able to predict risk for psychiatric disorders

including depression[14]. Consistent with this, our findings suggest developmental GC
exposure leads to differences in GR-induced transcriptional activation in adult brains and
may mediate the risk for psychiatric disorders by altering a network of neuronal signaling-
related stress-sensitive genes.

Growing evidence indicates that GC signaling can lead to modifications to DNA methylation,
histone codes, and non-coding RNAs[54-56]. Similarly, we identified significantly altered
expression of epigenetic modulators. One of the most significantly altered epigenetic
regulators in bPAC+ brains are *DNMTs* (*dnmt1* and *dnmt3aa*) at the larval stage (6 dpf),
which may result in a global or local reduction in DNA methylation. DNMT1 mainly plays a
role in the maintenance of DNA methylation[57] while accumulation or gain of DNA
methylation is achieved by DNMT3a and -3b[58]. Given such far-reaching effects of each
DNMT, alteration of expression levels of dnmt genes that we observe in bPAC+ is predicted
to change the global DNA methylation landscape across the life course. In our study, we
identified consistently downregulated adult bPAC+ DEGs associated with histone
modifications across the life course. Temporal regulation of chromatin structure via the
"histone code" in neurons has been regarded as a crucial factor for dynamic plasticity in the
mature nervous system and long-lasting neuronal responses[59]. Among the histone modulator
genes, *hdac4* and *hdac5* were consistently downregulated in bPAC+ fish, and have been
implicated in depression, fear learning, PTSD, eating disorder (Anorexia nervosa), and
schizophrenia[60-65]. Furthermore, many epigenetic-associated adult bPAC+ DEGs are involved
in RNA processing and RNA editing. Among them, we found a downregulation of *adarb2*
and *pum2* in bPAC+ fish after exposure to LD. Chen et al. suggest that ADARB2 has a
regulatory role in RNA editing by inhibiting the binding of other adenosine deaminases
acting on RNA (ADARs)[66] and ADARs are associated with the development of epileptic
seizures in mice and depressive phenotypes[67,68]. Further, *pum2* is involved in axon growth
and branching[69]. However, further investigation is required to reveal direct mechanisms
between the identified adult bPAC+ DEGs and alterations of epigenetic modulators.

As early exposure to excess GC is considered a global risk factor for the development of
mental disorders, our behavioral pipeline was tailored to assess performance in key
behavioral manifestations rather than disease-specific tasks. bPAC+ fish did not show defects
in aspects of perception as they can recognize visual and olfactory stimuli in the form of
social cues and food pellets. However, bPAC+ fish showed deficits in fear learning using a

pavlovian conditioning paradigm that pairs electric shock with the color green. It is possible that perceived salience of electric shock as a stressor rather than fear learning is diminished in bPAC+. However, we consider this possibility less likely as bPAC+ can respond robustly to acute stress presented in the form of LD. This phenotype is consistent with what has been reported in rodent ELS models, which showed deficits in fear learning, learning impairment, and decreased cognitive flexibility[70-72]. In zebrafish, fear-conditioned avoidance learning is a robust phenotype whereby fish exhibit freezing and erratic behavior upon re-exposure to the conditioned stimulus (CS)[73, 74]. Although one study found that chronic unpredictable stress did not have any effect on fear conditioning in adult zebrafish[75], a different ELS protocol found improvement of short-term memory by early life stress when testing in the unconditioned y-maze[76]. Differences in how zebrafish learn fear association are dependent on whether they are reactive or proactive responders to stress, raising the question of how bPAC+ fish's response fits into that spectrum[77].

To assess how bPAC+ responds in a novel environment, we utilized the novel tank test which has been compared to the open field test in rodents. A well-reported typical response of wild-type fish in the novel tank assay consists of immediate dive to the bottom, sustained depth-preference, and erratic bouts of fast swimming[78,79]. Both depth-preference and erratic bouts are enhanced with increased levels of stress and can be ameliorated by treatment with anxiolytic compounds[80-82]. bPAC+ fish showed less depth preference throughout the trial and fewer erratic swimming bouts compared to wild types, suggesting altered novelty perception and processing in bPAC+ fish. This outcome further supports the findings in the fear conditioning assay, that bPAC+ fish may possess dysfunctional fear processing and/or novelty adaption mechanisms.

bPAC+ fish exhibited a reduction in the maintenance of social interactions in the absence of social recognition or approach deficits using our new social approach and maintenance assay[22]. Both a reduction in the time spent in the interaction zone and the number of interactions performed by bPAC+ fish were observed. Similarly in previous zebrafish ELS studies, diminished shoaling behavior was observed following early-life social isolation and was associated with both motivational deficiencies linked to dopaminergic systems[83] and increased sensitivity to stimuli which was reversed by anxiolytic treatment[84]. Impairments in social behavior or social withdrawal are common among different psychiatric disorders,

including autism[85] and schizophrenia[86], and have been described in rodent models of ELS in

which deficits in social interaction in the absence of social recognition and approach have

been observed[87]. Interestingly, neonatal handling and short bouts of maternal separation in

male rats resulted in reduced oxytocin neurons in the parvocellular region of the

paraventricular nucleus (PVN) of the hypothalamus and a reduction in the time spent

interacting and sniffing, with no differences in recognition or approach reported[87]. We

observed reduced expression of oxytocin, in bPAC+ brains during the larval stage, and this

downregulation persisted until adulthood, although the extent of the difference decreased

over time. Consistent with our observation that posterior oxt+ cluster is selectively reduced in

bPAC+, a subset of parvocellular oxt neurons enriched genes identified in recent scRNA-seq

results of mice were downregulated in bPAC+ (such as *brinp1*, *baiap3*, *pacsin1a*, *syt17*, and

rcan2)[88]. This may suggest that subgroups of oxt+ neurons are differentially controlled by

ELS during development and play distinct roles in ELS-induced modification of stress

responses and social behavior.

In conclusion, we show that developmental GC exposure leads to altered behaviors and

brain-wide transcriptional changes that were enriched in the processes of DNA metabolism,

nervous system development and neuronal signaling across the life course. We identified

brain-wide adult bPAC+ DEGs that are overrepresented in gene sets associated with

neuropsychiatric disorders and are enriched in epigenetic modulators. Further, developmental

GC exposure led to exaggerated endocrine responses and primed a set of genes to exhibit

altered transcriptomic responses to subsequent challenges in adult fish. Lastly, we identified

reduced expression of *oxt* and *mytl1a* in bPAC+ which may contribute to the altered social

behavior phenotype in bPAC+. We propose that the brain-wide DEG set identified here will

be useful for guiding future studies dissecting the mechanism underlying developmental GC-

exposure and behavioral dysfunction and neuropsychiatric disease susceptibility in adulthood.

The zebrafish model presented here provides an opportunity to dissect the long-term effects

of endogenous GC elevation during development and can be leveraged in future studies to

identify critical period, intensity, and duration of developmental GC exposure that leads to

adult dysfunction.

Acknowledgment

This project was funded by the German Federal Office for Education and Research (BMBF)

grant number 01GQ1404 and Mireille and Dennis Gillings Foundation Grant to SR, Basic

Science Research Program (2020R1A6A3A03037828) of National Research Foundation
(NRF) of Korea to MC and Institutional Strategic Support Fund 3 scheme (ISSF3) to
Translational Research Exchange @ Exeter (ISSF3-TREE-Choi, 2022) funded by Wellcome
Trust to MC. We acknowledge the support of TRON gGmbH (Mainz, Germany) for the
RNA-sequencing, Bioimaging Center (University of Exeter, UK) for confocal imaging and
Aquatic Resources Center (University of Exeter, UK) and Kathrin Domdera (University of
Mainz, DE) for expert fish care, and High-Performance Computing (HPC) facility
(University of Exeter, UK) for the server for informatic analyses.
632

633 **Author contributions**

634 SR conceived the project. MC and SR designed the project. MC acquired and analyzed
635 transcriptomic data. AC acquired and analyzed behavior data. SK, HE, AT and SK
636 contributed to reagent generation or data acquisition. MC, SR, and AC drafted the article.
637 MC, HE, and SR critically revised the manuscript. All authors approved the submission.
638

639 **Competing interests**

640 SR holds a patent, European patent number 2928288 and US patent number 10,080,355: "A
641 novel inducible model of stress.". The remaining authors declare no known competing
642 interests.
643

Figure legends

Figure 1. Ambient light-exposed bPAC+ fish show elevation of cortisol during development. (a) An image of *Tg(star:bPAC-2A-tdTomato)*[+/-] larva at 5 dpf. The transgene is expressed solely in the interrenal gland (arrow). The schematic represents the intracellular events in the steroidogenic interrenal cells of transgenic fish (dotted box). Endogenous GC production is stimulated by ACTH signaling. Similarly, bPAC mimics ACTH signaling by increasing cAMP when exposed to blue light. (b) Fish were raised under ambient aquarium lighting conditions with white fluorescent light (12-hour light and 12-hour dark). Elevation of basal GC level was observed at larval and juvenile stages. (c) White ambient light (blue+) elevates the basal GC level of 6 dpf bPAC+ larvae. ***=$P<0.001$, two-tailed unpaired t-test. However, there was no significant difference in GC level when fish were raised under filter restricting blue light spectrum within the ambient light (blue-). (d) One of the well-known GR signaling markers, *fkbp5*, was significantly upregulated in bPAC+ larvae ($P = 0.016$) compared to wild-type control, two-tailed unpaired t-test. * = $P<0.05$. Basal GC level at larval stages (e) and from juvenile to adult (f). Abbreviations: GC, glucocorticoid; ACTH, Adrenocorticotropic Hormone; AC, adenylyl cyclase; cAMP, Cyclic adenosine monophosphate

Figure 2. Reduced social interaction in adult bPAC+ fish. No significant differences were observed in average speed (a) nor turn frequency (b) among wild type, bPAC+ and bPAC-. (c) Schematic figures for the social approach and maintenance test. During the pre-social phase, no significant difference in color preference (d) nor depth (e) was observed. (f) Representative traces of wild-type fish during the social phase. The green shade indicates the interaction zone. (g) During the social phase, no significant difference in approach rate to conspecific fish between groups. Log-rank test (Mantel-Cox, $P = 0.77$). (h) Representative traces of wild-type and bPAC+ fish during the social phase. bPAC+ and bPAC- showed a significantly reduced duration (i) and frequency of social interactions (j) compared to the wild type. Dunnett's multiple comparisons followed one-way ANOVA, a) $F_{(2, 27)}=0.2248$, $P=0.80$; b) $F_{(2, 27)}=2.332$, $P=0.12$; d) $F_{(2, 98)}=0.6224$, $P=0.54$; e) $F_{(2, 98)}=1.801$, $P=0.1706$; i) $F_{(2, 98)}=6.273$, $P=0.0027$; j) $F_{(2, 98)}=10.93$, $P<0.0001$.

Figure 3. Developmental GC exposure leads to differential gene expressions across the life course (a) Venn Diagram for up- and down-regulated DEGs between wild type and bPAC+; criteria for DEG identification: |FC|>2 and FDR <0.01. A lower FC threshold, |FC|>1.5, was applied for samples at 120 dpf. (b) Average read counts of genes by genotype and time points are represented by z-score in the heat map. Bars on the right side indicate the paternity of fish. (c) Expression profiles of GC-associated genes. *=$P<0.05$, **=$P<0.01$, ***=$P<0.001$, ****=$P<0.0001$. Tukey's multiple comparisons test followed two-way repeated measures ANOVA. (d) Top enriched biological processes for up- and down-regulated DEGs in bPAC+. d6: 6 dpf, d13: 13 dpf, d120: 120 dpf, wt: wild type, FC: fold change, GC: glucocorticoid, *adj.p*: adjusted *P*-value, ratio: identified DEGs/the number of genes included in the term. Complete DEG-enriched GO terms are described in Supplementary Table 3.

Figure 4. Developmental GC exposure primes differential gene expression following acute stress in bPAC+ adults. (a) Schematic of the acute stress paradigm. Looming dots (LD) mimic approaching predators from above (Cook et al. 2023). (b) Schematic for acute-stressed and control sample collection (arrows). Acute stress is induced by looming dot (LD)-exposure in adults (120 dpf). post-LD samples were collected after 15 min of LD exposure.

Yellow striped box: bPAC+ activation; yellow stars: LD. (c) bPAC+ showed significantly
higher GC levels in post-LD conditions, compared with basal. Tukey's multiple comparisons
followed one-way ANOVA; F(5, 54)=19.31, $P<0.0001$. (d) Expression profiles of GC-
induced genes following LD exposure. *= $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$ from
multiple two-tailed unpaired t-tests. Error bars indicate mean+SEM. (e) Schematic for three
comparisons to identify transcriptional changes in response to LD exposure in adults. DEGs
were determined by FC and FDR. |FC|>1.5 and FDR<0.01 for DEG1 and 2; |FC|>2 and
FDR<0.01 for DEG3. (f) z-score for mean CPM of adult bPAC+ DEGs is visualized in a
heatmap. (g) Venn diagram for adult bPAC+ DEGs. The list of DEGs is described in
Supplementary Table 4. The adult bPAC+ DEGs = DEG1 + DEG2 + DEG3 - (DEG1 ∩
DEG2). GC-primed genes = DEG1 + DEG2 - (DEG1 ∩ DEG2). The red asterisk represents
the overlap (2 genes) between DEG1 and DEG3. (h) The top 3 GO terms enriched in
downregulated GC-primed genes. The enrichment score = -$\log_{10}$(adjusted P-value). The
complete results of the functional enrichment test are described in Supplementary Table 5.
708
Figure 5. Adult bPAC+ DEGs are enriched in gene sets associated with human
neuropsychiatric disorders. (a) Result of human disorders enrichment test based on the
DisGeNET database. The ratio represents overlapping/input genes. Top ranked disorders are
schizophrenia (FDR= 1.61e-04, ratio (adult bPAC+ DEGs/disease genes)=170/883), bipolar
disorder (FDR= 7.48e-04, ratio=100/477), depressive disorder (FDR= 2.89e-03,
ratio=63/289), mental depression (FDR= 2.87e-03, ratio=59/254) and autistic disorder (FDR=
1.61e-04, ratio=65/261). (b) Venn diagram for adult bPAC+ DEGs that are associated with
the top five human psychiatric disorders in panel a. Depressive disorder and Mental
Depression-associated genes (as shown in a), are combined and referred to as depressive
disorders in b and d. (c) STRING protein-protein interaction (PPI) networks of
schizophrenia-associated adult bPAC+ DEGs genes in pre- and post-LD conditions indicate
LD-sensitive gene expression. The size and color of the node represent comparative fold
change, and gene expression; up (orange) or downregulation (blue), respectively. DEGs
enriched in KEGG pathways Neuroactive ligand-receptor interaction (FDR = 2.01e-08) are
labeled in yellow whilst those enriched in MAPK signaling pathway (FDR=9.74e-08) are
labeled in green. (d) Top GO terms enriched in the top-ranked disease-associated adult
bPAC+ DEGs. Adult bPAC+ DEGs that are associated with schizophrenia, depression,
autism, and bipolar disorder are mainly enriched in the downregulated functions in synaptic
signaling, and multicellular organismal process. The enrichment score = -log10(adjusted P-
value). (e) The heatmap depicts the $\log_2$FC of GC-altered genes associated with the top five
human psychiatric disorders in panel a (Depressive disorder and mental depression are
merged). The numbers in the brackets represent the number of adult bPAC+ DEGs. 6, 13, and
120 denote dpf, and LD denotes post-LD at 120 dpf. (f) Subsets of adult bPAC+ DEGs
associated with the top five diseases in a show consistent differential expression across the
life course (significant at all time points, labeled trans-temporal) or only in adulthood
(significant only in adulthood, labeled adult-onset). Genes that exhibited expression patterns
outside of these conditions were excluded.
736
Figure 6. Social behavior-associated genes tend to be downregulated in bPAC+ fish
across the life course. (a) The expression of social behavior-associated genes, log2FC from
two comparisons, bPAC+ vs. wild type, and bPAC- vs. wild type are represented by a
heatmap. Asterix in each cell denotes that FDR < 0.05. Red labels indicate GC-primed genes.
(b) Decreased *oxt* expression in bPAC+ was confirmed both in larval and adult brains by real-
time qPCR. * = $P<0.05$, multiple unpaired t-tests. Error bar indicates mean±SEM. (c)
Expression of *oxt* was decreased especially in the medial posterior cluster of oxt+ cells

(yellow arrows) in the neurosecretory preoptic area (NPO) of 5 dpf bPAC+ larvae. a: anterior,
p: posterior. scale bar represents 25µm. (d) the signal intensity of *oxt* in NPO was
significantly reduced in bPAC+ NPO. (e) the number of oxt+ cells in bPAC+ larvae were
significantly less than those of wild type. Mann-Whitney test, d) n=8; e) n=70. (f) Protein-
protein interaction network of significant DEGs that are implicated in social behavior at 6 dpf.
The color and size of nodes represent log2FC. This protein network was enriched in
modulation of chemical synaptic transmission (strength=1.38, FDR=5.44e-12), synaptic
signaling (strength=1.33, FDR=1.12e-11), learning and memory (strength=1.67, FDR=1.60e-
12), pre-synapse assembly (strength=2.41, FDR=5.44e-11) and social behavior
(strength=2.44, FDR=1.17e-30). (g) Log2FC of known oxt regulatory elements. *= FDR <
0.05. (h) Top enriched transcription factors for differentially expressed social behavior-
associated genes. Light green represents TFs that are significantly differentially expressed in
bPAC+. (i-j) Top enriched transcription factors (TFs) for upregulated (i) or downregulated (j)
GC-primed genes are presented as a bar plot. The light green color indicates TFs that are
significantly differentially expressed in bPAC+. (k) Expression of *myt1la* is significantly
downregulated in in bPAC+ both in pre- and post-LD. ns= $P>0.05$, *=$P<0.05$, **=$P<0.01$,
****=$P<0.0001$, Tukey's multiple comparisons test followed by one-way ANOVA.
761
762
763

764   **References**

766   1.   Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan
767   on the brain, behaviour and cognition. *Nat Rev Neurosci* 2009; **10**(6): 434-445.
768

769   2.   Teicher MH, Samson JA, Anderson CM, Ohashi K. The effects of childhood
770   maltreatment on brain structure, function and connectivity. *Nature Reviews*
771   *Neuroscience* 2016; **17**(10): 652-666.
772

773   3.   Chen Y, Baram TZ. Toward Understanding How Early-Life Stress Reprograms
774   Cognitive and Emotional Brain Networks. *Neuropsychopharmacology* 2016; **41**(1):
775   197-206.
776

777   4.   Sandi C, Haller J. Stress and the social brain: behavioural effects and neurobiological
778   mechanisms. *Nat Rev Neurosci* 2015; **16**(5): 290-304.
779

780   5.   Brunson KL, Kramár E, Lin B, Chen Y, Colgin LL, Yanagihara TK *et al.* Mechanisms of
781   late-onset cognitive decline after early-life stress. *J Neurosci* 2005; **25**(41): 9328-9338.
782

783   6.   Kessler RC, McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM *et al.*
784   Childhood adversities and adult psychopathology in the WHO World Mental Health
785   Surveys. *Br J Psychiatry* 2010; **197**(5): 378-385.
786

787   7.   McEwen BS, Gianaros PJ. Central role of the brain in stress and adaptation: links to
788   socioeconomic status, health, and disease. *Ann N Y Acad Sci* 2010; **1186**: 190-222.
789

790   8.   Heim C, Nemeroff CB. The role of childhood trauma in the neurobiology of mood and
791   anxiety disorders: preclinical and clinical studies. *Biol Psychiatry* 2001; **49**(12): 1023-
792   1039.
793

794   9.   Charmandari E, Tsigos C, Chrousos G. Endocrinology of the stress response. *Annu Rev*
795   *Physiol* 2005; **67**: 259-284.
796

797   10.   Wendelaar Bonga SE. The stress response in fish. *Physiol Rev* 1997; **77**(3): 591-625.
798

799   11.   Weaver IC, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR *et al.*
800   Epigenetic programming by maternal behavior. *Nat Neurosci* 2004; **7**(8): 847-854.
801

802   12.   Peña CJ, Smith M, Ramakrishnan A, Cates HM, Bagot RC, Kronman HG *et al.* Early life
803   stress alters transcriptomic patterning across reward circuitry in male and female
804   mice. *Nat Commun* 2019; **10**(1): 5098.
805

806   13.   Provençal N, Arloth J, Cattaneo A, Anacker C, Cattane N, Wiechmann T *et al.*
807   Glucocorticoid exposure during hippocampal neurogenesis primes future stress
808   response by inducing changes in DNA methylation. *Proc Natl Acad Sci U S A* 2020;
809   **117**(38): 23280-23285.

811 14. Arloth J, Bogdan R, Weber P, Frishman G, Menke A, Wagner Klaus V *et al.* Genetic
812   Differences in the Immediate Transcriptome Response to Stress Predict Risk-Related
813   Brain Function and Psychiatric Disorders. *Neuron* 2015; **86**(5): 1189-1202.
814

815 15. Bacila I, Cunliffe VT, Krone NP. Interrenal development and function in zebrafish.
816   *Molecular and Cellular Endocrinology* 2021; **535**: 111372.
817

818 16. Eachus H, Zaucker A, Oakes JA, Griffin A, Weger M, Güran T *et al.* Genetic Disruption
819   of 21-Hydroxylase in Zebrafish Causes Interrenal Hyperplasia. *Endocrinology* 2017;
820   **158**(12): 4165-4173.
821

822 17. Oakes JA, Li N, Wistow BRC, Griffin A, Barnard L, Storbeck K-H *et al.* Ferredoxin 1b
823   Deficiency Leads to Testis Disorganization, Impaired Spermatogenesis, and
824   Feminization in Zebrafish. *Endocrinology* 2019; **160**(10): 2401-2416.
825

826 18. Li N, Oakes JA, Storbeck K-H, Cunliffe VT, Krone NP. The P450 side-chain cleavage
827   enzyme Cyp11a2 facilitates steroidogenesis in zebrafish. *Journal of Endocrinology*
828   2020; **244**(2): 309-321.
829

830 19. Oakes JA, Barnard L, Storbeck K-H, Cunliffe VT, Krone NP. 11β-Hydroxylase loss
831   disrupts steroidogenesis and reproductive function in zebrafish. *Journal of*
832   *Endocrinology* 2020; **247**(2): 197-212.
833

834 20. Choi M, Cook A, Eachus H, Tochwin A, Kuntz S, Ryu S. Comprehensive analysis
835   methods for developmental GC exposed zebrafish. *protocolsio* 2023.
836

837 21. Gutierrez-Triana JA, Herget U, Castillo-Ramirez LA, Lutz M, Yeh CM, De Marco RJ *et al.*
838   Manipulation of Interrenal Cell Function in Developing Zebrafish Using Genetically
839   Targeted Ablation and an Optogenetic Tool. *Endocrinology* 2015; **156**(9): 3394-3401.
840

841 22. Cook A, Beckmann H, Azap R, Ryu S. Acute stress modulates social approach and
842   social maintenance in adult zebrafish. *bioRxiv* 2023: 2023.2002.2010.528021.
843

844 23. Herget U, Wolf A, Wullimann MF, Ryu S. Molecular neuroanatomy and
845   chemoarchitecture of the neurosecretory preoptic-hypothalamic area in zebrafish
846   larvae. *J Comp Neurol* 2014; **522**(7): 1542-1564.
847

848 24. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T *et al.* Fiji: an
849   open-source platform for biological-image analysis. *Nature Methods* 2012; **9**(7): 676-
850   682.
851

852 25. Nagpal J, Eachus H, Lityagina O, Ryu S. Optogenetic induction of chronic
853   glucocorticoid exposure in early-life impairs stress-response in larval zebrafish.
854   *bioRxiv* 2022: 2022.2009.2009.507267.
855

856   26.   Binder EB. The role of FKBP5, a co-chaperone of the glucocorticoid receptor in the
857   pathogenesis and therapy of affective and anxiety disorders.
858   *Psychoneuroendocrinology* 2009; **34 Suppl 1:** S186-195.

860   27.   Meaney MJ. Maternal Care, Gene Expression, and the Transmission of Individual
861   Differences in Stress Reactivity Across Generations. *Annual Review of Neuroscience*
862   2001; **24**(1): 1161-1192.

864   28.   Vera-Chang MN, St-Jacques AD, Gagné R, Martyniuk CJ, Yauk CL, Moon TW *et al.*
865   Transgenerational hypocortisolism and behavioral disruption are induced by the
866   antidepressant fluoxetine in male zebrafish Danio rerio. *Proceedings of the National*
867   *Academy of Sciences* 2018; **115**(52): E12435-E12442.

869   29.   Ord J, Heath PR, Fazeli A, Watt PJ. Paternal effects in a wild-type zebrafish implicate
870   a role of sperm-derived small RNAs. *Molecular Ecology* 2020; **29**(14): 2722-2735.

872   30.   White RJ, Collins JE, Sealy IM, Wali N, Dooley CM, Digby Z *et al.* A high-resolution
873   mRNA expression time course of embryonic development in zebrafish. *Elife*, vol.
874   62017, p e30860.

876   31.   Griffiths KR, Morris RW, Balleine BW. Translational studies of goal-directed action as
877   a framework for classifying deficits across psychiatric disorders. *Frontiers in Systems*
878   *Neuroscience* 2014; **8**.

880   32.   Ike KGO, de Boer SF, Buwalda B, Kas MJH. Social withdrawal: An initially adaptive
881   behavior that becomes maladaptive when expressed excessively. *Neuroscience &*
882   *Biobehavioral Reviews* 2020; **116:** 251-267.

884   33.   Levin ED, Bencan Z, Cerutti DT. Anxiolytic effects of nicotine in zebrafish. *Physiology*
885   *& Behavior* 2007; **90**(1): 54-58.

887   34.   Egan RJ, Bergner CL, Hart PC, Cachat JM, Canavello PR, Elegante MF *et al.*
888   Understanding behavioral and physiological phenotypes of stress and anxiety in
889   zebrafish. *Behavioural Brain Research* 2009; **205**(1): 38-44.

891   35.   Gulliver LS. Xenobiotics and the Glucocorticoid Receptor. *Toxicol Appl Pharmacol*
892   2017; **319:** 69-79.

894   36.   Draper N, Stewart PM. 11beta-hydroxysteroid dehydrogenase and the pre-receptor
895   regulation of corticosteroid hormone action. *J Endocrinol* 2005; **186**(2): 251-271.

897   37.   Balyan R, Cai M, Zhao W, Dai Z, Zhai Y, Chen G. Repeated restraint stress upregulates
898   rat sulfotransferase 1A1. 2019; **30**(2): 265-273.

900   38.   Zheng D-Q, Zhang K, Wu X-C, Mieczkowski PA, Petes TD. Global analysis of genomic
901   instability caused by DNA replication stress in *Saccharomyces cerevisiae*. *Proceedings*
902   *of the National Academy of Sciences* 2016; **113**(50): E8114-E8121.

904 39. Eachus H, Choi MK, Tochwin A, Kaspareit J, Ho M, Ryu S. Elevated Glucocorticoid
905 alters the trajectory of hypothalamic development and function. *bioRxiv* 2023**:**
906 2023.2001.2027.525966.
907

908 40. Piñero J, Saüch J, Sanz F, Furlong LI. The DisGeNET cytoscape app: Exploring and
909 visualizing disease genomics data. *Computational and Structural Biotechnology*
910 *Journal* 2021; **19:** 2960-2967.
911

912 41. Neumann ID. Brain Oxytocin: A Key Regulator of Emotional and Social Behaviours in
913 Both Females and Males. *Journal of Neuroendocrinology* 2008; **20**(6)**:** 858-865.
914

915 42. Herget U, Gutierrez-Triana JA, Salazar Thula O, Knerr B, Ryu S. Single-Cell
916 Reconstruction of Oxytocinergic Neurons Reveals Separate Hypophysiotropic and
917 Encephalotropic Subtypes in Larval Zebrafish. *eNeuro* 2017; **4**(1).
918

919 43. Wee CL, Nikitchenko M, Wang W-C, Luks-Morgan SJ, Song E, Gagnon JA *et al.*
920 Zebrafish oxytocin neurons drive nocifensive behavior via brainstem premotor
921 targets. *Nature Neuroscience* 2019; **22**(9)**:** 1477-1492.
922

923 44. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J *et al.* STRING v11:
924 protein–protein association networks with increased coverage, supporting functional
925 discovery in genome-wide experimental datasets. *Nucleic Acids Research* 2019;
926 **47**(D1)**:** D607-D613.
927

928 45. Keenan AB, Torre D, Lachmann A, Leong AK, Wojciechowicz ML, Utti V *et al.* ChEA3:
929 transcription factor enrichment analysis by orthogonal omics integration. *Nucleic*
930 *Acids Research* 2019; **47**(W1)**:** W212-W224.
931

932 46. Blanchet P, Bebin M, Bruet S, Cooper GM, Thompson ML, Duban-Bedu B *et al.* MYT1L
933 mutations cause intellectual disability and variable obesity by dysregulating gene
934 expression and development of the neuroendocrine hypothalamus. *PLoS Genet* 2017;
935 **13**(8)**:** e1006957.
936

937 47. Parichy DM, Elizondo MR, Mills MG, Gordon TN, Engeszer RE. Normal table of
938 postembryonic zebrafish development: Staging by externally visible anatomy of the
939 living fish. *Developmental Dynamics* 2009; **238**(12)**:** 2975-3015.
940

941 48. Mueller T, Wullimann MF. Anatomy of neurogenesis in the early zebrafish brain.
942 *Developmental Brain Research* 2003; **140**(1)**:** 137-155.
943

944 49. de Kloet ER, Joëls M, Holsboer F. Stress and the brain: from adaptation to disease.
945 *Nature Reviews Neuroscience* 2005; **6**(6)**:** 463-475.
946

50. Reul JM, van den Bosch FR, de Kloet ER. Differential response of type I and type II
corticosteroid receptors to changes in plasma steroid level and circadian rhythmicity.
Neuroendocrinology 1987; **45**(5)**:** 407-412.

51. Seckl JR. 11beta-hydroxysteroid dehydrogenases: changing glucocorticoid action.
Curr Opin Pharmacol 2004; **4**(6): 597-602.

52. Wyrwoll CS, Holmes MC, Seckl JR. 11β-hydroxysteroid dehydrogenases and the brain:
from zero to hero, a decade of progress. *Front Neuroendocrinol* 2011; **32**(3): 265-286.

53. Wang XD, Rammes G, Kraev I, Wolf M, Liebl C, Scharf SH *et al.* Forebrain CRF_1
modulates early-life stress-programmed cognitive deficits. *J Neurosci* 2011; **31**(38):
13625-13634.

54. Matthews SG, McGowan PO. Developmental programming of the HPA axis and
related behaviours: epigenetic mechanisms. *Journal of Endocrinology* 2019; **242**(1):
T69-T79.

55. Wang HT, Huang FL, Hu ZL, Zhang WJ, Qiao XQ, Huang YQ *et al.* Early-Life Social
Isolation-Induced Depressive-Like Behavior in Rats Results in Microglial Activation
and Neuronal Histone Methylation that Are Mitigated by Minocycline. *Neurotox Res*
2017; **31**(4): 505-520.

56. Thomassin H, Flavin M, Espinás M-L, Grange T. Glucocorticoid-induced DNA
demethylation and gene memory during development. *The EMBO Journal* 2001;
20(8): 1974-1983.

57. Hermann A, Goyal R, Jeltsch A. The Dnmt1 DNA-(cytosine-C5)-methyltransferase
Methylates DNA Processively with High Preference for Hemimethylated Target Sites.
Journal of Biological Chemistry 2004; **279**(46): 48350-48359.

58. Okano M, Bell DW, Haber DA, Li E. DNA methyltransferases Dnmt3a and Dnmt3b are
essential for de novo methylation and mammalian development. *Cell* 1999; **99**(3):
247-257.

59. Borrelli E, Nestler EJ, Allis CD, Sassone-Corsi P. Decoding the epigenetic language of
neuronal plasticity. *Neuron* 2008; **60**(6): 961-974.

60. Hobara T, Uchida S, Otsuki K, Matsubara T, Funato H, Matsuo K *et al.* Altered gene
expression of histone deacetylases in mood disorder patients. *J Psychiatr Res* 2010;
44(5): 263-270.

61. Maddox SA, Kilaru V, Shin J, Jovanovic T, Almli LM, Dias BG *et al.* Estrogen-dependent
association of HDAC4 with fear in female mice and women with PTSD. *Mol*
Psychiatry 2018; **23**(3): 658-665.

62. Saha P, Gupta R, Sen T, Sen N. Histone Deacetylase 4 Downregulation Elicits Post-
Traumatic Psychiatric Disorders through Impairment of Neurogenesis. *Journal of*
Neurotrauma 2019; **36**(23): 3284-3296.

63.   Renthal W, Maze I, Krishnan V, Covington HE, 3rd, Xiao G, Kumar A *et al.* Histone
deacetylase 5 epigenetically controls behavioral adaptations to chronic emotional
stimuli. *Neuron* 2007; **56**(3): 517-529.
1000

1001 64.   Cui H, Moore J, Ashimi SS, Mason BL, Drawbridge JN, Han S *et al.* Eating disorder
1002   predisposition is associated with ESRRA and HDAC4 mutations. *J Clin Invest* 2013;
1003   **123**(11): 4706-4713.
1004

1005 65.   Wu Y, Hou F, Wang X, Kong Q, Han X, Bai B. Aberrant Expression of Histone
1006   Deacetylases 4 in Cognitive Disorders: Molecular Mechanisms and a Potential Target.
1007   *Front Mol Neurosci* 2016; **9**: 114.
1008

1009 66.   Chen CX, Cho DS, Wang Q, Lai F, Carter KC, Nishikura K. A third member of the RNA-
1010   specific adenosine deaminase gene family, ADAR3, contains both single- and double-
1011   stranded RNA binding domains. *Rna* 2000; **6**(5): 755-767.
1012

1013 67.   Higuchi M, Maas S, Single FN, Hartner J, Rozov A, Burnashev N *et al.* Point mutation
1014   in an AMPA receptor gene rescues lethality in mice deficient in the RNA-editing
1015   enzyme ADAR2. *Nature* 2000; **406**(6791): 78-81.
1016

1017 68.   Bombail V, Qing W, Chapman KE, Holmes MC. Prevention of 5-hydroxytryptamine2C
1018   receptor RNA editing and alternate splicing in C57BL/6 mice activates the
1019   hypothalamic-pituitary-adrenal axis and alters mood. *Eur J Neurosci* 2014; **40**(11):
1020   3663-3673.
1021

1022 69.   Martínez JC, Randolph LK, Iascone DM, Pernice HF, Polleux F, Hengst U. Pum2
1023   Shapes the Transcriptome in Developing Axons through Retention of Target mRNAs
1024   in the Cell Body. *Neuron* 2019; **104**(5): 931-946.e935.
1025

1026 70.   Lehmann J, Pryce CR, Bettschen D, Feldon J. The maternal separation paradigm and
1027   adult emotionality and cognition in male and female Wistar rats. *Pharmacol Biochem
1028   Behav* 1999; **64**(4): 705-715.
1029

1030 71.   Aisa B, Tordera R, Lasheras B, Del Río J, Ramírez MJ. Effects of maternal separation
1031   on hypothalamic-pituitary-adrenal responses, cognition and vulnerability to stress in
1032   adult female rats. *Neuroscience* 2008; **154**(4): 1218-1226.
1033

1034 72.   Thomas AW, Caporale N, Wu C, Wilbrecht L. Early maternal separation impacts
1035   cognitive flexibility at the age of first independence in mice. *Dev Cogn Neurosci* 2016;
1036   **18**: 49-56.
1037

1038 73.   Maximino C, do Carmo Silva RX, Dos Santos Campos K, de Oliveira JS, Rocha SP,
1039   Pyterson MP *et al.* Sensory ecology of ostariophysan alarm substances. *J Fish Biol*
1040   2019; **95**(1): 274-286.
1041

1042 74. Blank M, Guerim LD, Cordeiro RF, Vianna MR. A one-trial inhibitory avoidance task to
1043   zebrafish: rapid acquisition of an NMDA-dependent long-term memory. *Neurobiol*
1044   *Learn Mem* 2009; **92**(4)**:** 529-534.
1045

1046 75. Fontana BD, Gibbon AJ, Cleal M, Norton WHJ, Parker MO. Chronic unpredictable
1047   early-life stress (CUELS) protocol: Early-life stress changes anxiety levels of adult
1048   zebrafish. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 2021;
1049   **108:** 110087.
1050

1051 76. Fontana BD, Gibbon AJ, Cleal M, Sudwarts A, Pritchett D, Miletto Petrazzini ME *et al.*
1052   Moderate early life stress improves adult zebrafish (Danio rerio) working memory
1053   but does not affect social and anxiety-like responses. *Dev Psychobiol* 2021; **63**(1)**:** 54-
1054   64.
1055

1056 77. Baker MR, Wong RY. Contextual fear learning and memory differ between stress
1057   coping styles in zebrafish. *Sci Rep* 2019; **9**(1)**:** 9935.
1058

1059 78. Gerlai R, Lahav M, Guo S, Rosenthal A. Drinks like a fish: zebra fish (Danio rerio) as a
1060   behavior genetic model to study alcohol effects. *Pharmacol Biochem Behav* 2000;
1061   **67**(4)**:** 773-782.
1062

1063 79. Cachat JM, Canavello PR, Elkhayat SI, Bartels BK, Hart PC, Elegante MF *et al.* Video-
1064   Aided Analysis of Zebrafish Locomotion and Anxiety-Related Behavioral Responses.
1065   In: Kalueff AV, Cachat JM (eds). *Zebrafish Neurobehavioral Protocols*. Humana Press:
1066   Totowa, NJ, 2011, pp 1-14.
1067

1068 80. Wong K, Elegante M, Bartels B, Elkhayat S, Tien D, Roy S *et al.* Analyzing habituation
1069   responses to novelty in zebrafish (Danio rerio). *Behav Brain Res* 2010; **208**(2)**:** 450-
1070   457.
1071

1072 81. Parra KV, Adrian JC, Jr., Gerlai R. The synthetic substance hypoxanthine 3-N-oxide
1073   elicits alarm reactions in zebrafish (Danio rerio). *Behav Brain Res* 2009; **205**(2)**:** 336-
1074   341.
1075

1076 82. Duarte T, Fontana BD, Müller TE, Bertoncello KT, Canzian J, Rosemberg DB. Nicotine
1077   prevents anxiety-like behavioral responses in zebrafish. *Prog Neuropsychopharmacol*
1078   *Biol Psychiatry* 2019; **94:** 109655.
1079

1080 83. Shams S, Amlani S, Buske C, Chatterjee D, Gerlai R. Developmental social isolation
1081   affects adult behavior, social interaction, and dopamine metabolite levels in
1082   zebrafish. *Developmental Psychobiology* 2018; **60**(1)**:** 43-56.
1083

1084 84. Tunbak H, Vazquez-Prada M, Ryan TM, Kampff AR, Dreosti E. Whole-brain mapping
1085   of socially isolated zebrafish reveals that lonely fish are not loners. *Elife* 2020; **9:**
1086   e55863.
1087

1088   85.   Chevallier C, Kohls G, Troiani V, Brodkin ES, Schultz RT. The social motivation theory
1089   of autism. *Trends in Cognitive Sciences* 2012; **16**(4): 231-239.

1090

1091   86.   Miyakawa T, Leiter LM, Gerber DJ, Gainetdinov RR, Sotnikova TD, Zeng H *et al.*
1092   Conditional calcineurin knockout mice exhibit multiple abnormal behaviors related
1093   to schizophrenia. *Proceedings of the National Academy of Sciences* 2003; **100**(15):
1094   8987-8992.

1095

1096   87.   Todeschin AS, Winkelmann-Duarte EC, Jacob MHV, Aranda BCC, Jacobs S, Fernandes
1097   MC *et al.* Effects of neonatal handling on social memory, social interaction, and
1098   number of oxytocin and vasopressin neurons in rats. *Hormones and Behavior* 2009;
1099   **56**(1): 93-100.

1100

1101   88.   Lewis EM, Stein-O'Brien GL, Patino AV, Nardou R, Grossman CD, Brown M *et al.*
1102   Parallel Social Information Processing Circuits Are Differentially Impacted in Autism.
1103   *Neuron* 2020; **108**(4): 659-675.e656.

1104
1105

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6