Direct long read visualization reveals metabolic interplay between two antimalarial drug targets

Short title: metabolic connection drives adaptive evolution

Shiwei Liu[1, 2], Emily R. Ebel[3], Aleksander Luniewski[1], Julia Zulawinska[1], Mary Lewis Simpson[1], Jane Kim[1], Nnenna Ene[1], Thomas Werner Anthony Braukmann[4], Molly Congdon[5], Webster Santos[5], Ellen Yeh[4], Jennifer L. Guler[1]*

[1]University of Virginia, Department of Biology, Charlottesville, VA, USA

[2]Current affiliation: Indiana University School of Medicine, Indianapolis, IN, USA

[3]Stanford, Departments of Pediatrics and Microbiology & Immunology, Stanford, CA, USA

[4]Stanford University, Departments of Pathology and Microbiology & Immunology, Stanford, CA, USA

[5]Virginia Tech, Department of Chemistry, Blacksburg, VA, USA

*Corresponding Author

Email: jlg5fw@virginia.edu

Abstract

Increases in the copy number of large genomic regions, termed genome amplification, are an important adaptive strategy for malaria parasites. Numerous amplifications across the *Plasmodium falciparum* genome contribute directly to drug resistance or impact the fitness of this protozoan parasite. During the characterization of parasite lines with amplifications of the *dihydroorotate dehydrogenase* (*DHODH*) gene, we detected increased copies of an additional genomic region that encompassed 3 genes (~5 kb) including *GTP cyclohydrolase I* (*GCH1* amplicon). While this gene is reported to increase the fitness of antifolate resistant parasites, *GCH1* amplicons had not previously been implicated in any other antimalarial resistance context. Here, we further explored the association between *GCH1* and *DHODH* copy number. Using long read sequencing and single read visualization, we directly observed a higher number of tandem *GCH1* amplicons in parasites with increased *DHODH* copies (up to 9 amplicons) compared to parental parasites (3 amplicons). While all *GCH1* amplicons shared a consistent structure, expansions arose in 2-unit steps (from 3 to 5 to 7, etc copies). Adaptive evolution of *DHODH* and *GCH1* loci was further bolstered when we evaluated prior selection experiments; *DHODH* amplification was only successful in parasite lines with pre-existing *GCH1* amplicons. These observations, combined with the direct connection between metabolic pathways that contain these enzymes, lead us to propose that the *GCH1* locus is beneficial for the fitness of parasites exposed to *DHODH* inhibitors. This finding highlights the importance of studying variation within individual parasite genomes as well as biochemical connections of drug targets as novel antimalarials move towards clinical approval.

Author Summary

Malaria is caused by a protozoan parasite that readily evolves resistance to drugs that are used to

treat this deadly disease. Changes that arise in the parasite genome, including extra copies of

important genes, directly contribute to this resistance or improve how well the resistant parasite

competes. In this study, we identified that extra copies of one gene (*GTP cyclohydrolase* or *GCH1*)

were more likely to be found in parasites with extra copies of another gene on a different

chromosome (*dihydroorotate dehydrogenase* or *DHODH*). A method that allows us to view long

pieces of DNA from individual genomes was especially important for this study; we were able to

assess gene number, arrangement, and boundary sequences, which provided clues into how extra

copies evolved. Additionally, by analyzing previous experiments, we identified that extra *GCH1*

copies improved resistance to drugs that target DHODH. The relationship between these two loci

is supported by a direct connection between the folate and pyrimidine biosynthesis pathways that

the parasite uses to make DNA. Since *GCH1* amplicons are common in clinical parasites

worldwide, this finding highlights the need to study metabolic connections to avoid resistance

evolution.

Introduction

Malaria is a disease caused by the protozoan *Plasmodium* parasite and *P. falciparum* is the leading

cause of human malaria deaths (1). Due to the lack of effective vaccines against malaria infection,

antimalarial drugs are the primary approach for malaria treatment (2). However, drug efficacy is

mitigated by the frequent emergence of antimalarial resistant parasites (3).

Changes in the copy number of large genomic regions, termed copy number variations or CNVs,

are an important adaptive strategy for malaria parasites (4–12). Amplification, one type of CNV

with increased gene copy number, plays an essential role in the evolution of resistance to various

antimalarials (6,7,9,11,13–16). As one example, amplification of the *dihydroorotate*

dehydrogenase (*DHODH*) gene directly confers resistance to DHODH inhibitors (i.e. DSM1) in

parasites propagated in vitro (11). DHODH is an important enzyme in the *P. falciparum*

pyrimidine biosynthesis pathway that contributes resources for nucleic acid synthesis (17,18).

DHODH amplicons increase transcription and presumably translation of the drug target to directly

impact drug sensitivity (11).

In another example, amplification of the *GTP cyclohydrolase 1* (*GCH1*) gene increases the fitness

of clinical parasite populations that are resistant to antifolates (i.e. pyrimethamine and sulfadoxine)

(4,7,8,16). GCH1 is the first enzyme in the folate biosynthesis pathway and increased flux through

this pathway compensates for fitness costs of resistance-conferring mutations in *dihydropteroate*

synthase (*DHPS*) and *dihydrofolate synthase* (*DHFS*) genes (4,8,15). Although the contribution of

GCH1 amplicons to antifolate resistance is well studied, this locus has not been reported to

contribute to resistance to other antimalarials (15,19,20).

Typically, gene copy number is studied using widely accessible high coverage short read

sequencing (11,21–24). However, this approach has limitations including non-unique mapping of

repetitive sequences, the inability to resolve complex genomic rearrangements, and the overall

difficulty of detecting structural variations (25–27). These challenges are exacerbated by the high

AT-content of the *P. falciparum* genome (28,29). Long read technologies such as Oxford

Nanopore sequencing have the potential to span low complexity and repetitive regions and more

accurately represent structural variation (Cretu Stancu et al., 2017; Sedlazeck et al., 2018a, 2018b;

Ho et al., 2020). Moreover, this single molecule sequencing approach allows the examination of

clonal heterogeneity within cellular populations (34,35).

In this study, we identified an unprecedented association between *GCH1* and *DHODH* copy

numbers from a previously acquired short read data set. To explore this relationship further, we

performed long read sequencing and directly observed the step-wise expansion of the *GCH1*

amplicon in parasites with elevated *DHODH* copies. Using single long read visualization, we

determined that the structure and orientation of the amplicon were preserved during expansion and

amplicons increased in a step-wise manner. The positive correlation between *GCH1* and *DHODH*

copy number presented here, combined with the recognition that resistance to DHODH inhibitors

has only been selected in parasite lines with *GCH1* expansions, suggest the adaptive evolution of

these two genomic loci. In addition, the direct connection between folate and pyrimidine

biosynthesis supports metabolic interplay between the two pathways. Further study of this

relationship is important considering the potential use of DHODH inhibitors to treat clinical

malaria with preexisting *GCH1* amplicons.

Results

Through expanded analysis of short read data from a family of parasites selected with DSM1,

originally presented in (11) (**Figure 1A**), we observed a positive association between *GCH1* and

DHODH copy number (**Figure 1B**). Using droplet digital PCR on the same parasite lines that had

recently been propagated in our laboratory, we confirmed that *GCH1* copy number trended higher

as *DHODH* copy number increased (**Table 1**).

Fig 1. *GCH1* copy number increase is positively correlated with *DHODH* copy number in one family of DSM1 selected parasites. **(A)** DSM1 selection schematic, as presented previously (11). Blue text: Illumina short read sequenced lines. Underline: modern lines confirmed by ddPCR analysis (Table 1). Asterisk (*): lines subjected to long read sequencing in this study. Wild-type (WT1, *Dd2*) *P. falciparum* was selected with DSM1 in two steps; the first step selected for low-level (L) resistance and the second step selected for moderate- (M) or high-level (H) resistance. DSM1 EC50 values are as follows: L1 (1 µM), L2 (0.9 µM), M1 (7.2 µM), H2 (85 µM), H6 (56 µM), H4 (49 µM). All values were previously reported and clone names were adapted as previously (11,36). **(B)** Relationship between *GCH1* and *DHODH* copy number in DSM1 selected parasites as quantified using short read data from Guler et al. 2013. A trend line was added to show the relationship between *GCH1* and *DHODH* copy numbers but a correlation coefficient could not be calculated due to the small sample size (n=5) and dependence among the lines.

Table 1. Positive *GCH1*:*DHODH* association is validated using Droplet Digital PCR on modern parasite lines.

Line	Sample type	Average copy number (SD)*			*GCH1* CN relative to parent	*DHODH* CN relative to parent
		GCH1	*DHODH*	*ATP6*		
WT1	Parent	2.5 (0.2)	0.8 (0.1)	1.1 (0.0)	1.0	1.0
L1	Round 1 selection	3.9 (0.3)	3.2 (0.1)	1.0 (0.1)	1.6	4
M1	Round 1 selection	3.9 (0.1)	5.2 (0.3)	1.0 (0.1)	1.6	6.5
H2	Round 2 selection	4.6 (0.2)	6.0 (0.3)	1.1 (0.1)	1.8	7.5
H6	Round 2 selection	4.3 (0.3)	7.0 (0.4)	1.0 (0.1)	1.7	8.8

*The average copy number is calculated by comparing the ddPCR signal to a single copy gene signal (*Seryl tRNA synthetase*, PF3D7_0717700). N=4. *ATP6* copy number is a single copy gene expected to be 1 in all parasite lines.

We conducted long read sequencing to more precisely define *GCH1* copy number and amplicon

structure in the parental line versus two DSM1-selected lines (WT1, M1, and H2, **Table S1**). We

directly visualized single reads using an application that represents gene segments of individual

long reads (*Materials and Methods*). Small amplicons, like those including *GCH1*, are especially

conducive to this approach because long reads span multiple copies of the amplicon as well as

both up and downstream regions. Read visualization showed that the general structure of two 3-

gene amplicons, separated by an inversion of the same 3-genes, was conserved between the

parental and selected lines (**Figure 2A-C**). This amplicon structure was reported previously in the

WT1 (*Dd2*) parental line (**Figure 2D**) (4).

Fig 2. Long-read visualization shows conserved boundaries and structure of *GCH1* amplicon from parental and resistance parasites. (A-C) Representative images from the Shiny application comparing the *GCH1* amplicon in WT1 (**A**), M1 (**B**), and H2 (**C**) spanning reads. Red dashed square: copies of *GCH1* amplicon covering 3 genes. Each gene sequence from the *3D7* reference genome (no *GCH1* amplicon represented) was split into <=500 bp fragments and blasted against individual Nanopore reads (dark gray: genic regions, light grey: intergenic regions, more details

in *Materials and Methods*). **(D)** Orientation of *GCH1* amplicons from each parasite line. The three
genes within the *GCH1* amplicon unit include PF3D7_1224000 (*GTP cyclohydrolase I, GCH1*),
PF3D7_1223900 (*50S ribosomal protein L24*, putative, *50S RPL24)*, and PF3D7_1223800
(*citrate/oxoglutarate carrier protein*, putative, *YHM2*). Note: The gene order (*GCH1-50S RPL24-*
YHM2) is reversed compared to the 3D7 reference genome (*YHM2-50S RPL24-GCH1*) in order to
facilitate comparison with the read images in panels A-C. Red circles: amplicon junction
sequences that act as potential secondary recombination sites.

Using direct read visualization, we also recorded the number of *GCH1* amplicon units (**Figure 2D**)

from both spanning and non-spanning reads (**Figure 3A, Table S2**) Spanning reads cover genes

both upstream and downstream of the amplicon while non-spanning reads end within the amplicon.

We consistently detected up to 3 amplicon units on reads from the parental line. Although the

overall mean copy number was similar for all parasite lines (**Figure 3A**, WT1, 2.5 copies; M1, 3.6

copies; H2; 3.2 copies, $F(2,95)=[2.61]$, $p = 0.07$), we observed more reads that encompassed a

higher number of *GCH1* amplicon units in the selected lines (up to 9 units, **Figure 3A**). When we

restricted our analysis to spanning reads, or those that displayed the entire amplified region, we

detected a significant difference in mean copy number between the parasite lines (WT1 vs M1,

$p=0.0003$ and WT1 vs H2, $p=0.01$; **Figure S1**). The presence of longer reads was not contributing

to this observation; there was not a significant difference in the length of reads used for *GCH1*

analysis from each parasite line ($F(2,96)=[2.51]$, $p = 0.09$, **Figure 3B**, median read lengths: WT1,

51144bp; M1, 49018bp; H2, 26429bp). The larger variation observed for the M1 sample (**Figure**

3B) was due to an improvement in ONT technology and methods leading to an increase in the

number of longer reads (i.e. WT1 N50 of 15.6kb on R9.4.1 flow cell and M1 N50 of 99.7kb on

R10.4.1 flow cell, **Table S1,** see *Materials and Methods*).

167
Fig 3. Quantification of long reads displays an increase in *GCH1* amplicon number in DSM1
selected parasites that is not dependent on overall read length. (**A**) *GCH1* copy number from
WT1 and selected (M1 and H2) parasite lines. Spanning and non-spanning reads are included;
spanning reads were grouped with their corresponding non-spanning count (e.g. a read showing 2
copies of the *GCH1* amplicon is counted as a 2+ read). Bars represent mean, error bars represent
standard deviation (dotted lines, values that represent detected spanning reads, see **Figure S1**). (**B**)
Read length distribution from long reads (>=10kb) covering the *GCH1* amplicon. Thick line
represents median, thin line represents quartiles.

During this analysis, we identified that >50% of reads from M1 and H2 lines depicted amplicon

copy numbers greater than the expected WT copy number (**Figure 3A**, **Table S2**). To assess

whether variation in the copy number of the *GCH1* locus was common in other laboratory-adapted

parasite lines, we evaluated several additional long read datasets (**Table S2**, *Supporting*

Information, Vembar et al., 2016). In general, parasite lines exhibited the expected *GCH1*

amplicon size (i.e. *Dd2* versus *3D7*, as reported previously (4) and visualized in **Figures 2D** and

S2) and the amplicon copy number for each line was relatively stable across diverse datasets

(limited to <25% of reads above the expected copy number).

When we evaluated the amplicon structure, all M1 and H2 reads covering the *GCH1* region began

with a set of 3 amplicon units, as observed in the parental line (**Figure 2A**). This basic structure

was followed by groups of 2 units, where one copy was inverted and the other was in normal

orientation (**Figure 2D**). This suggests that increases in copy number were arising as two unit

steps. We identified further evidence of the two unit amplicon expansion when we limited our

analysis to only spanning reads, which allow direct quantification of amplicon copy number. Here,

we did not detect any spanning reads that carried *even GCH1* copy numbers (only 3, 5, 7, or 9

copies, **Figure S1**).

To investigate whether an increase in *GCH1* amplicons was contributing generally to resistance to

DHODH inhibitors, we evaluated the copy number of *GCH1* amplicons using short read data

collected from parasites selected with other DSM compounds in two studies (18,37). Contrary to

DSM1-selected parasites (**Figure 1B**), we did not detect increases in *GCH1* amplicon number in

parasites selected with DSM265, DSM267, or DSM705 compared to parental *Dd2* or *3D7* parasite

lines (**Figure S3**, **Table S3**). For this analysis, we only included parasite lines that acquired

DHODH amplicons to confer resistance.

The absence of a correlation between *DHODH* and *GCH1* copy number in these other datasets

suggests that *GCH1* may not play a direct role in resistance. However, parental lines used in these

two studies harbored pre-existing *GCH1* amplicons (~3-6 copies, **Figure 4A**), indicating that this

locus may support the evolution of resistance involving *DHODH* amplification. Therefore, we

compiled information on all previous resistance selections conducted in parasite lines that naturally

vary in *GCH1* copy number (7 total studies, **Table 2**). *3D7* and *Dd2* parasite lines, which carry the

highest *GCH1* copy numbers (3-6 copies) with the smallest overall amplicon sizes of common

laboratory parasites (1 and 3 genes, respectively, **Figure 4B**), were most commonly used in

successful resistance selections (6 and 4 studies, respectively, **Table 2**) and were capable of

acquiring *DHODH* amplicons (**Figure 4A**). *K1* parasites harbor a larger low copy *GCH1* amplicon

(anticipated as 2 copies of 7 genes, **Figure 4B**) and although resistance could be selected in one

study, it was 1000-fold less efficient than *Dd2* selections run in parallel (38). The *DHODH* copy

number status of this K1-derived selected line was not investigated. Finally, *Hb3*, which harbors

one *GCH1* copy (**Table 2, Figure 4B**), was incapable of developing resistance in two independent

studies (11,38) (**Figure 4A**).

215
Fig 4. Preexisting *GCH1* amplicons confer resistance competence. (**A**) Relationship between
GCH1 and *DHODH* copy number in parental (black outline) and DHODH inhibitor-selected
parasite lines as quantified using short read data (see **Table 2** for source data). Each data point
represents a copy number from an individual parasite clone or line from a distinct selection (see
Tables S3 and **S4** for copy number assessment). *3D7*, yellow squares (2 selections); *Dd2*, blue
circles (2 selections); *K1*, red triangle (1 selection, unknown *DHODH* copy number post-
selection); *HB3*, grey diamonds (2 selections). (**B**) *GCH1* amplicon size and gene number vary
between parasite lines used for DHODH inhibitor selections (see **Table 2** for selection details).
3D7, 4 copy, single gene (PF3D7_1224000, 2kb, this study); *Dd2*, 3 copy, three genes
(PF3D7_1223800- PF3D7_1224000, 5kb, this study); *K1*, unknown copy number (grey dash,
anticipated 2 copy), 7 genes (PF3D7_1223500 - PF3D7_1224100, 19kb) (39); *HB3*, 1 copy of
region (40).

228
229

Table 2. Parental line *GCH1* copy number determines resistance competence

Study (author, year)	Selection Compounds	Success of Resistance Selection				*DHODH* amplicon
		Dd2 (*GCH1*=3)	*3D7* (*GCH1*=4-6)	*Hb3* (*GCH1*=1*)	Other lines	
Guler, 2013 (11)	DSM1	**Yes**	N/A	**No**	N/A	Yes
Palmer, 2021 (37)	DSM705	**Yes**	**Yes**	N/A	N/A	Yes
Phillips, 2015 (38)	DSM265	**Yes**	N/A	**No**	K1-Yes**	Yes
Mandt, 2019 (18)	DSM265, DSM267	**Yes**	**Yes**	N/A	N/A	^Yes
White, 2019 (41)	DSM265	**Yes**	N/A	N/A	N/A	No
Ross, 2014 (42)	DSM74	**Yes**	**Yes**	N/A	N/A	No
Lukens, 2014 (43)	Genz-666136, DSM74	N/A	**Yes**	N/A	N/A	No

*The *GCH1* copy number for the *HB3* parasite line is single copy based on whole genome sequencing (40). Prior studies detected two *GCH1* copies using quantitative PCR but did not detect an increase in expression from this region (8,15).

**The *K1* parasite line has a *GCH1* amplicon that is ~19kb (961,455–980,630, (39)) but the exact copy number has not been reported. Phillips et al. reported that it was 100x more difficult to select for DSM265 resistance using *K1* than *Dd2* (MIR- 2×10^8 versus 2×10^6, respectively, (38). Resistance mechanisms (i.e. mutation or amplification) were not evaluated for the resistant *K1* line in this study.

^Although both *Dd2* and *3D7* were used in selections, only two *3D7* clones acquired *DHODH* amplifications in this study (18) and were evaluated for *GCH1* copy number in our study (**Tables S3 and S4**).

Discussion

Using both short-read sequencing as well as an accurate PCR-based method, ddPCR, we uncovered that *GCH1* copy number is positively associated with *DHODH* copy number in malaria parasites selected with DSM1 (**Figures 1B and 4, Table 1**). However, limitations in each of these methods contribute to inaccuracies in copy number results. Most importantly, both of these methods provide an average value for the parasite population and do not reflect variation from individual genomes. Long-read sequencing combined with a custom visualization tool (44) allowed us to directly quantify the copy number of the *GCH1* locus and assess the structure of all amplicons (**Figures 2** and **3**). In the context of this haploid, single cellular parasite, each read represents a distinct DNA strand from a single genome; this approach is an accurate way to visualize copy number and junction heterogeneity across a population (**Figure 2**, **Figure S2**). We showed that the detection of more *GCH1* amplicons on reads from DSM1 selected parasite lines

(**Figure 3A**) was not caused by sample preparation (**Figure 3B**, **Table S2**) or natural variation during in vitro culture (**Table S2**). The visualization of tandem amplicons on single reads (**Figure 2A-C**) and the step-wise increase by 2 amplicon units (from 3 to 5 to 7 copies, **Figures 2D** and **S1**) provides direct evidence for *GCH1* amplicon expansion in the DSM1 resistance context.

The two-unit steps suggest that there is a secondary recombination site that leads to facile re-amplification of the region (**Figure 2D**). In the high coverage M1 sample, we identified long AT-rich sequences at the end of the amplicon unit (**Figure S4**) that may participate in further DNA breakage (AT dinucleotides at the *GCH1* end, **Figure S4C**) and recombination after the initial event (homopolymeric A/T tracks at the *YHM2* end, **Figure S4A**)(24). Multiple junctions at these positions provide evidence of the initial event that formed the 3 unit amplicon as well as a secondary event resulting in a two-unit step-wise expansion.

As recently recognized in organisms from bacteria to humans (34,35,45–47), long read sequencing is an exceptional tool to investigate genomic heterogeneity within populations. This method is not without its limitations; reads that end within the amplicon region lead to an underestimation in the total copy number; in our study, 65/98 (66%) of reads were non-spanning, which contributed to a similar overall mean copy number between parasite lines (**Figure 3A**). However, the visualization of spanning reads overcame this limitation, clearly showing the two-unit expansion of *GCH1* amplicons (**Figures 2A-C** and **S1**). The quantification of spanning reads also confirms that variation in *GCH1* amplicon numbers likely arises stochastically in the parasite population; while most spanning reads from the high coverage M1 line showed 5 amplicon copies (15/21 reads), we observed higher amplicon copies on a minority of reads (7 and 9 copies, 3/21 reads). In line with our previous model for malaria genome evolution (11,24), parasites with higher copy numbers are present within the population, poised to be selected under specific conditions.

The *GCH1* locus is a proven hotspot for genetic change (8,39). *GCH1* amplicons have previously

been associated with antifolate resistance; an increase in flux through the folate biosynthesis

pathway via *GCH1* alleviates fitness effects of mutations that confer pyrimethamine and

sulfadoxine resistance (4,8,16,19,20). A similar contribution to resistance to DHODH inhibitors

would not be surprising, given the close connection of the folate and pyrimidine biosynthesis

pathways (**Figure 5**); they both contribute to nucleotide biosynthesis and converting dUMP to

dTMP in pyrimidine biosynthesis requires reciprocal conversion of N^5, N^{10}-methylene-THF to

DHF by the folate pathway. Strong evidence for this metabolic connection comes from the

observation that parasite lines without *GCH1* amplicons are not able to develop resistance to

DHODH inhibitors (**Table 2**) (11,38). Assessment of resistance competence, combined with our

observations in the DSM1 context and folate-pyrimidine metabolic interplay, lead us to propose

that increases in *GCH1* copy number are also beneficial for the fitness of parasites carrying

DHODH amplicons. During resistance evolution, increased DHODH protein levels interact with

more of the inhibitor to cause resistance. When the inhibitor is not present, higher DHODH levels

increase flux through pyrimidine biosynthesis, which in turn requires more THF for the completion

of the pathway (**Figure 5**).

Fig 5. The connection between pyrimidine and folate biosynthesis pathways. Enzymes with
gene copy number variations are indicated in blue text (DHODH: dihydroorotate dehydrogenase;
GCH1: GTP cyclohydrolase 1) and connections between the two pathways in blue arrows. Gln:
Glutamine; DHO: Dihydroorotate; UMP: Uridine monophosphate; dUMP: Deoxyuridine
monophosphate; dTMP: deoxythymidine monophosphate; GTP: Guanosine-5'-triphosphate;
DHPS: Dihydropteroate synthase; DHFR: Dihydrofolate reductase; DHF: Dihydrofolate; THF:
Tetrahydrofolate; HMDP-P2: 6-hydroxymethyl-7, 8-dihydropterin diphosphate; pABA: para-
amino-benzoic acid; 5, 10-CH2-THF: 5,10-Methylenetetrahydrofolate. *$Dd2$ carries 3 mutations
in both DHPS and DHFR (6 total) and HB3 and 3D7 have a single mutation each in DHFR and
DHPS, respectively (15,48).

There are a few factors that may impact parasite reliance on folate biosynthesis including parasite

genetic background and the presence of folate precursors in the host environment. Antifolate

resistant parasites from different regions of the world that harbor specific mutations rely more

heavily on increased GCH1 flux to alleviate fitness effects (i.e. DHFR and DHPS mutants,

(4,8,15,19), **Figure 5**). While $Dd2$ and $3D7$ have distinct origins and mutational patterns (49,50)

(**Figure 5**), the fact that they both maintain the ability to develop DHODH inhibitor resistance

excludes the possibility that DHFR and DHPS mutations matter in this context. Additionally,

levels of folate precursors, such as para-aminobenzoic acid (pABA), vary widely in host versus in

vitro environments (51–55) and exert different selective pressures on parasites during drug

selection (56). Higher pABA levels in human serum, as were used during DSM1 selections (11),

may drive the accumulation of higher *GCH1* copy numbers and hold extra benefits by bolstering

both folate and pyrimidine biosynthesis (**Figure 5**). Our observation of some level of variation in

GCH1 amplicon number in standard parasite lines (**Table S2**) suggests that differing in vitro media

formulations may drive copy number changes at this locus.

Importantly, DHODH inhibitors have entered trials for the treatment of clinical malaria (18) and

resistance-conferring *DHODH* amplicons can evolve in vivo (57). Meanwhile, antifolate

resistance and *GCH1* amplifications are widespread in clinical *P. falciparum* isolates, especially

in Southeast Asia (4,7,8,39) and now Africa (16,58). Additionally, a zoonotic malaria species, *P.*

knowlesi, also acquires extra *GCH1* copies after adaptation to human blood (59). The potential use

of DHODH inhibitors on clinical parasites carrying *GCH1* amplicons necessitates further

evaluation of the interplay between these two important metabolic pathways.

Materials and Methods

DSM1 and parasite clones

DSM1 inhibits dihydroorotate dehydrogenase enzyme of *Plasmodium* pyrimidine biosynthesis

(60). DSM1-resistant parasites were selected during a previous study and cloned by limiting

dilution at the time of isolation (11) (**Figure 1A**). The naming scheme representing low (L),

moderate (M), and high (H) levels of DSM1 resistance follows that used previously by our group

(36).

Parasite culture

We thawed erythrocytic stages of *P. falciparum* (*Dd2*, MRA-150; *3D7*, MRA-102, Malaria

Research and Reference Reagent Resource Center, BEI Resources and DSM1 selected clones as

highlighted in **Figure 1A**, a generous gift from Pradipsinh Rathod, University of Washington)

from frozen stocks and maintained them as previously described (61). Briefly, we grew parasites

at 37°C in vitro at 3% hematocrit (serotype A positive human erythrocytes, Valley Biomedical,

VA or BioIVT, NY) in RPMI 1640 medium (Invitrogen, USA) containing 24 mM NaHCO3 and

25 mM HEPES, and supplemented with 20% human type A positive heat inactivated plasma

(Valley Biomedical, VA or BioIVT, NY) in sterile, sealed flasks flushed with 5% O_2, 5% CO_2,

and 90% N_2 (11). We maintained the cultures with media changes every other day and sub-cultured

them as necessary to keep parasitemia below 5%. We determined all parasitemia measurements

and staging using SYBR green-based flow cytometry (62). We routinely tested cultures using the

LookOut Mycoplasma PCR Detection Kit (Sigma-Aldrich, USA) to confirm negative

Mycoplasma status.

Short read sequencing analysis and CNV detection

We analyzed amplifications in Illumina short read datasets of *P. falciparum* parasites selected by

different DSM antimalarial drugs (DSM1, DSM265, DSM267, and DSM705, **Table S3**)

(11,18,37). We first processed and mapped the reads to the reference genome as previously

described (24,63). Briefly, we trimmed Illumina adapters from reads with the BBDuk tool in

BBMap (version 38.57) (64). We aligned each fastq file to the *3D7* reference genome with

Speedseq (version 0.1.0) through BWA-MEM alignment (65). We discarded the reads with low-

mapping quality scores (below 10) and removed duplicated reads using Samtools (version 1.10)

(66). We analyzed split and discordant reads from the mapped reads using LUMPY in Speedseq

to determine the location and length of the previously reported *GCH1*, *DHODH*, and multidrug

resistance protein 1 (*MDR1*) amplicons (**Table S4**) (67). For read-depth analysis, we further

filtered the mapped reads using a mapping quality score of 30. To determine the copy number of

the *GCH1*, *DHODH*, and *MDR1* amplicons, we used CNVnator (version 0.4.1) with a bin size of

100 bp; the optimal bin size was chosen to detect GCH1 amplicons in all analyzed samples (68).

Droplet Digital PCR

Prior to Droplet Digital (dd) PCR, we digested DNA with restriction enzyme RsaI (Cut Site:

GT/AC) following the manufacturer's instructions (New England Biolabs, Ipswich, MA, USA) in

37°C incubation for one hour. We selected the restriction enzyme RsaI to cut outside of the ddPCR

amplified regions of desired genes and separate amplicon copies to be distributed into droplets.

We diluted the digested DNA for ddPCR reactions. We performed ddPCR using ddPCR Supermix

for Probes (no dUTP, Bio-Rad Laboratories, California, USA) with DNA input 0.1 ng (in duplicate

per sample), 0.025 ng (in duplicate per sample) as previously described (36). The primers and

probes used in reactions are included in **Table S5**. The PCR protocol for probe-based assay was

95°C for 10 min, followed by 40 rounds of 95°C for 30 sec and 60°C for 1 min. *Seryl-tRNA*

synthetase and *calcium-transporting ATPase* (*ATP6*) served as single copy reference genes on

chromosome 7 and chromosome 1 respectively; *dihydroorotate dehydrogenase* (*DHODH*) and

GTP cyclohydrolase 1 (*GCH1*) are multi-copy genes (**Table S5**). We performed droplet generation

and fluorescence readings per the manufacturer's instructions. For each reaction, we required a

minimum number of 10,000 droplets to proceed with analysis. We calculated the ratio of positive

droplets containing a single- (*ATP6*) or multi-copy gene (*GCH1*, *DHODH*) versus a single-copy

gene (*seryl-tRNA synthetase*) using the Quantasoft analysis software (QuantaSoft Version 1.7,

BioRad Laboratories) and averaged between independent replicates.

DNA extraction for long read sequencing

We lysed asynchronous *P. falciparum*-infected erythrocytes with 0.15% saponin (Sigma-Aldrich,

USA) for 5 min at room temperature and washed them three times with 1× PBS (diluted from 10×

PBS Liquid Concentrate, Gibco, USA). We then lysed parasites with 0.1% Sarkosyl Solution

(Bioworld, bioPLUS, USA) in the presence of 1 mg/ml proteinase K (from *Tritirachium album*,

Sigma-Aldrich, USA) overnight at 37°C. We first extracted nucleic acids with

phenol/chloroform/isoamyl alcohol (25:24:1) pH 8.0 (Sigma-Aldrich, USA) three times using

1.5ml light Phase lock Gels (5Prime, USA), then further extracted nucleic acids with chloroform

twice using 1.5ml light Phase lock Gels (5Prime, USA). Lastly, we precipitated the DNA with

ethanol using the standard Maniatis method (69). To obtain high molecular weight genomic DNA,

we avoided any pipetting during the extraction, transferred solutions by direct pouring from one

tube to another, and mixed solutions by gently inverting the tubes.

Oxford Nanopore long read sequencing and analysis

For WT1 and H2 samples, we subjected ~1μg of high molecular weight genomic DNA to library

preparation for Oxford Nanopore sequencing following the Nanopore Native barcoding genomic

DNA protocol (version: NBE_9065_v109_revAB_14Aug2019) with 1x Ligation Sequencing kit

(SQK-LSK109, Oxford Nanopore Technologies, Oxford, UK). We performed DNA repair and

end preparation using NEBNext FFPE DNA Repair Mix (New England Biolabs, Ipswich, MA,

USA) and NEBNext End Repair/dA-Tailing Module (New England Biolabs, Ipswich, MA, USA).

We cleaned the A-tailed fragments using 0.9X AMPure XP beads (Beckman Coulter, High

Wycombe, UK). We then ligated barcodes to the end-prepped DNA using the Native Barcoding

Expansion 1–12 kit (EXP-NBD104, Oxford Nanopore Technologies, Oxford, UK) and Blunt/TA

Ligase Master Mix (New England Biolabs, Ipswich, MA, USA). We cleaned the barcoded samples

using 0.9X AMPure XP beads. Then we pooled barcoded samples in equimolar ratios and

subjected them to an adaptor ligation step, using the Adapter Mix II from the Native Barcoding

Expansion 1–12 kit and NEBNext Quick Ligation Reaction Buffer (New England Biolabs,

Ipswich, MA, USA) as well as Quick T4 DNA Ligase (New England Biolabs, Ipswich, MA, USA).

After adaptor ligation, we cleaned the WT1 and H2 libraries using AMPure XP beads.

For M1 samples, we isolated high molecular weight genomic DNA from 10 flasks of parasites (1-

4% parasitemia, ~30-40% late stage parasites) and prepared libraries as above except that we used

the cleanup and precipitation reagents provided in the Ultra-long DNA Sequencing kit (Oxford

Nanopore Technologies, Oxford, UK, V14). We quantified the adapter-ligated and barcoded DNA

using a Qubit fluorimeter (Qubit 1X dsDNA High Sensitivity Assay Kit, Life Technologies,

Carlsbad, CA).

We sequenced the WT1 and initial H2 libraries using the R9.4.1 flow cell (FLO-MIN106D),

another H2 library using the R10 flow cell (FLO-MIN111), and M1 libraries using the R10.4.1

flow cell (FLO-MIN114) on MinION (Oxford Nanopore Technologies, Oxford, UK). For WT and

H2 samples, we ran flow cells for 48 hours (controlled and monitored using the MinKNOW

software 3.6.5). For the M1 samples, we ran one-third of the library sequentially on the flow cell

for 24 hours each (up to 72 hours in total).

For base calling and demultiplexing of the Nanopore sequencing reads, we used Guppy (version

3.4.5+fb1fbfb) with the parameter settings "-c dna_r9.4.1_450bps_hac.cfg --barcode_kits "EXP-

NBD104" -x auto" for samples sequenced with R9.4.1 flow cell and "-c dna_r10_450bps_hac.cfg

-x auto" for the sample sequenced with R10 flow cell. We checked the read length and read quality

using "Nanoplot" (version 1.0.0) (see **Table S1**). We trimmed the adapters with "qcat" (version

1.1.0) (70) and filtered the reads with a cutoff "length ≥ 500 and Phred value ≥ 10" using the

program " filtlong version 0.2.0" (https://github.com/rrwick/Filtlong). To estimate the coverage of

sequencing reads in each sample, we aligned the filtered reads to *3D7* reference genome

(PlasmoDB) using "minimap2" (version 2.17) (71). "QualiMap" (version 2.2.1) (72) was used to

calculate the coverage of the aligned reads (see **Table S1**).

For junction analysis, we aligned Q10-filtered M1 reads (.bam) to the *3D7* reference genome

(PlasmdDB) and visualized read sequence at junctions of the GCH1 amplicon using Integrated

Genome Viewer (73,74).

Direct visualization of long reads

To visualize structural variants in the parasite genome, we used a custom R Shiny script to plot the

arrangement of reference gene segments on individual Nanopore reads (44). Briefly, we defined a

target region in the *3D7* reference genome (chromosome 12: 932916bp – 999275bp) that contained

3 genes in the *GCH1* amplicon and 11 flanking genes. We extracted the reference sequences of

these genes and subsequent intergenic regions, then split these sequences into fragments of 500-

1000 base pairs. We compared these fragments to individual Nanopore reads (>10kb) using

BLAST (75). We used the BLAST output as input for a custom R script, which drew blocks

representing homology between the defined genes (y-axis) and each individual read (x-axis). If

genes on the read were arranged in the same order as the reference genome then the blocks would

create a diagonal line on the plot. If there was a CNV that altered the gene order or direction then

the diagonal line is disrupted. We allowed the percent identity required to draw a homologous

rectangle to vary between reads, which varied in quality, using a slider in the Shiny application.

To filter out long reads with potentially spurious hits to gene fragments, we also used BLAST to

compare Nanopore reads to the reference genome and removed reads with <90% identity to

chromosome 12. To compare the mean copy number of GCH1 amplicon and read lengths between

WT1, M1, and H2 reads, we performed a one-way ANOVA in Graphpad Prism with an alpha

value of 0.05. If significant, differences were further explored using Tukey's multiple comparisons

testing.

Acknowledgments

Our thanks to Michelle Warthan, Noah Brown, and Ali Guler (University of Virginia) for

experimental, analysis, and statistical support, Martin Wu and Mercedes Campos-Lopez

(University of Virginia) for Oxford Nanopore sequencing support, and the laboratory of Dr.

William Petri Jr (University of Virginia) for their helpful discussions and insight. We are grateful

to David Fidock (Columbia University) and Dyann F. Wirth (Harvard University) for sharing

sequencing data and information on DSM-selected cell lines.

References

1. Rich SM, Leendertz FH, Xu G, LeBreton M, Djoko CF, Aminake MN, et al. The origin of
malignant malaria. Proc Natl Acad Sci U S A. 2009 Sep 1;106(35):14902–7.

2. Casares S, Brumeanu TD, Richie TL. The RTS,S malaria vaccine. Vaccine. 2010 Jul
12;28(31):4880–94.

3. Blasco B, Leroy D, Fidock DA. Antimalarial drug resistance: linking Plasmodium
falciparum parasite biology to the clinic. Nature medicine. 2017 Aug 4;23(8):917–28.

4. Kidgell C, Volkman SK, Daily J, Borevitz JO, Plouffe D, Zhou Y, et al. A Systematic Map
of Genetic Variation in Plasmodium falciparum. PLOS Pathogens. 2006;2(6):e57.

5. Conway DJ. Molecular epidemiology of malaria. Clin Microbiol Rev. 2007 Jan;20(1):188–
204.

6. Hyde JE. Drug-resistant malaria - an insight. FEBS J. 2007 Sep;274(18):4688–98.

7. Ribacke U, Mok BW, Wirta V, Normark J, Lundeberg J, Kironde F, et al. Genome wide
gene amplifications and deletions in Plasmodium falciparum. Molecular and Biochemical
Parasitology. 2007 Sep 1;155(1):33–44.

8. Nair S, Miller B, Barends M, Jaidee A, Patel J, Mayxay M, et al. Adaptive Copy Number
Evolution in Malaria Parasites. PLOS Genetics. 2008;4(10):e1000243.

9. Cheeseman IH, Gomez-Escobar N, Carret CK, Ivens A, Stewart LB, Tetteh KKA, et al.
Gene copy number variation throughout the Plasmodium falciparum genome. BMC
Genomics. 2009 Aug 4;10:353–353.

10. Bopp SER, Manary MJ, Bright AT, Johnston GL, Dharia NV, Luna FL, et al. Mitotic
Evolution of Plasmodium falciparum Shows a Stable Core Genome but Recombination in
Antigen Families. PLOS Genetics. 2013 Feb 7;9(2):e1003293.

11. Guler JL, Freeman DL, Ahyong V, Patrapuvich R, White J, Gujjar R, et al. Asexual
Populations of the Human Malaria Parasite, Plasmodium falciparum, Use a Two-Step
Genomic Strategy to Acquire Accurate, Beneficial DNA Amplifications. PLOS Pathogens.
2013 May 23;9(5):e1003375.

12. Menard D, Dondorp A. Antimalarial Drug Resistance: A Threat to Malaria Elimination.
Cold Spring Harb Perspect Med. 2017;7(7):a025619.

13. Foote SJ, Thompson JK, Cowman AF, Kemp DJ. Amplification of the multidrug resistance
gene in some chloroquine-resistant isolates of P. falciparum. Cell. 1989 Jun 16;57(6):921–
30.

14. Wilson CM, Volkman SK, Thaithong S, Martin RK, Kyle DE, Milhous WK, et al.
Amplification of pfmdr1 associated with mefloquine and halofantrine resistance in
Plasmodium falciparum from Thailand. Molecular and Biochemical Parasitology. 1993 Jan
1;57(1):151–60.

15. Heinberg A, Siu E, Stern C, Lawrence EA, Ferdig MT, Deitsch KW, et al. Direct evidence
for the adaptive role of copy number variation on antifolate susceptibility in Plasmodium
falciparum. Molecular microbiology. 2013 May;88(4):702–12.

16. Osei M, Ansah F, Matrevi SA, Asante KP, Awandare GA, Quashie NB, et al. Amplification
of GTP-cyclohydrolase 1 gene in Plasmodium falciparum isolates with the quadruple mutant
of dihydrofolate reductase and dihydropteroate synthase genes in Ghana. PLOS ONE.
2018;13(9):e0204871.

17. Phillips MA, Rathod PK. Plasmodium dihydroorotate dehydrogenase: a promising target for
novel anti-malarial chemotherapy. Infect Disord Drug Targets. 2010;10(3):226–39.

18. Mandt REK, Lafuente-Monasterio MJ, Sakata-Kato T, Luth MR, Segura D, Pablos-Tanarro
A, et al. In vitro selection predicts malaria parasite resistance to dihydroorotate
dehydrogenase inhibitors in a mouse infection model. Science Translational Medicine.
2019;11(521):eaav1636.

19. Kümpornsin K, Modchang C, Heinberg A, Ekland EH, Jirawatcharadech P, Chobson P, et al.
Origin of Robustness in Generating Drug-Resistant Malaria Parasites. Mol Biol Evol. 2014
Jul;31(7):1649–60.

20. Heinberg A, Kirkman L. The molecular basis of antifolate resistance in Plasmodium
falciparum: looking beyond point mutations. Ann N Y Acad Sci. 2015/02/18 ed.
2015;1342(1):10–8.

21. Herman JD, Rice DP, Ribacke U, Silterra J, Deik AA, Moss EL, et al. A genomic and
evolutionary approach reveals non-genetic drug resistance in malaria. Genome Biol.
2014;15(11):511–511.

22. Manary MJ, Singhakul SS, Flannery EL, Bopp SER, Corey VC, Bright AT, et al.
Identification of pathogen genomic variants through an integrated pipeline. BMC
Bioinformatics. 2014 Mar 3;15(1):63.

23. Cowell AN, Istvan ES, Lukens AK, Gomez-Lorenzo MG, Vanaerschot M, Sakata-Kato T, et
al. Mapping the malaria parasite druggable genome by using in vitro evolution and
chemogenomics. Science. 2018;359(6372):191–9.

24. Huckaby AC, Granum CS, Carey MA, Szlachta K, Al-Barghouthi B, Wang YH, et al.
Complex DNA structures trigger copy number variation across the Plasmodium falciparum
genome. Nucleic Acids Research. 2018 Dec 21;47(4):1615–27.

25. Alkan C, Coe BP, Eichler EE. Genome structural variation discovery and genotyping. Nature
Reviews Genetics. 2011 May 1;12(5):363–76.

26. Treangen TJ, Salzberg SL. Repetitive DNA and next-generation sequencing: computational
challenges and solutions. Nature Reviews Genetics. 2012 Jan 1;13(1):36–46.

27. Kosugi S, Momozawa Y, Liu X, Terao C, Kubo M, Kamatani Y. Comprehensive evaluation
of structural variation detection algorithms for whole genome sequencing. Genome Biology.
2019 Jun 3;20(1):117.

28. Beghain J, Langlois AC, Legrand E, Grange L, Khim N, Witkowski B, et al. Plasmodium
copy number variation scan: gene copy numbers evaluation in haploid genomes. Malaria
Journal. 2016 Apr 12;15(1):206.

29. Miles A, Iqbal Z, Vauterin P, Pearson R, Campino S, Theron M, et al. Indels, structural
variation, and recombination drive genomic diversity in Plasmodium falciparum. Genome
Res. 2016/08/16 ed. 2016;26(9):1288–99.

30. Cretu Stancu M, van Roosmalen MJ, Renkens I, Nieboer MM, Middelkamp S, de Ligt J, et
al. Mapping and phasing of structural variation in patient genomes using nanopore
sequencing. Nature Communications. 2017 Nov 6;8(1):1326.

31. Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, von Haeseler A, et al.
Accurate detection of complex structural variations using single-molecule sequencing. Nat
Methods. 2018/04/30 ed. 2018;15(6):461–8.

32. Sedlazeck FJ, Lee H, Darby CA, Schatz MC. Piercing the dark matter: bioinformatics of
long-range sequencing and mapping. Nature Reviews Genetics. 2018 Jun 1;19(6):329–46.

33. Ho SS, Urban AE, Mills RE. Structural variation in the sequencing era. Nature Reviews
Genetics. 2020 Mar 1;21(3):171–89.

34. Belikova D, Jochim A, Power J, Holden MTG, Heilbronner S. "Gene accordions" cause
genotypic and phenotypic heterogeneity in clonal populations of Staphylococcus aureus.
Nature Communications. 2020 Jul 14;11(1):3526.

35. Rugbjerg P, Dyerberg ASB, Quainoo S, Munck C, Sommer MOA. Short and long-read ultra-
deep sequencing profiles emerging heterogeneity across five platform Escherichia coli
strains. Metabolic Engineering. 2021 May 1;65:197–206.

36. McDaniels JM, Huckaby AC, Carter SA, Lingeman S, Francis A, Congdon M, et al.
Extrachromosomal DNA amplicons in antimalarial-resistant Plasmodium falciparum. Mol
Microbiol. 2020/11/19 ed. 2021;115(4):574–90.

37. Palmer MJ, Deng X, Watts S, Krilov G, Gerasyuto A, Kokkonda S, et al. Potent
Antimalarials with Development Potential Identified by Structure-Guided Computational
Optimization of a Pyrrole-Based Dihydroorotate Dehydrogenase Inhibitor Series. Journal of
Medicinal Chemistry. 2021 May 13;64(9):6085–136.

38. Phillips MA, Lotharius J, Marsh K, White J, Dayan A, White KL, et al. A long-duration
dihydroorotate dehydrogenase inhibitor (DSM265) for prevention and treatment of malaria.
Sci Transl Med. 2015 Jul 15;7(296):296ra111.

39. Turkiewicz A, Manko E, Sutherland CJ, Benavente ED, Campino S, Clark TG. Genetic
diversity of the Plasmodium falciparum GTP-cyclohydrolase 1, dihydrofolate reductase and
dihydropteroate synthetase genes reveals new insights into sulfadoxine-pyrimethamine
antimalarial drug resistance. PLOS Genetics. 2020 Dec 31;16(12):e1009268.

40. Sepúlveda N, Campino SG, Assefa SA, Sutherland CJ, Pain5 A, Clark TG. A Poisson
hierarchical modelling approach to detecting copy number variation in sequence coverage
data. BMC Genomics. 2013;14:128.

41. White J, Dhingra SK, Deng X, El Mazouni F, Lee MCS, Afanador GA, et al. Identification
and Mechanistic Understanding of Dihydroorotate Dehydrogenase Point Mutations in
Plasmodium falciparum that Confer in Vitro Resistance to the Clinical Candidate DSM265.
ACS Infect Dis. 2019 Jan 11;5(1):90–101.

42. Ross LS, Gamo FJ, Lafuente-Monasterio MJ, Singh OMP, Rowland P, Wiegand RC, et al. In
vitro resistance selections for Plasmodium falciparum dihydroorotate dehydrogenase
inhibitors give mutants with multiple point mutations in the drug-binding site and altered
growth. J Biol Chem. 2014 Jun 27;289(26):17980–95.

43. Lukens AK, Ross LS, Heidebrecht R, Javier Gamo F, Lafuente-Monasterio MJ, Booker ML,
et al. Harnessing evolutionary fitness in Plasmodium falciparum for drug discovery and
suppressing resistance. Proc Natl Acad Sci U S A. 2014 Jan 14;111(2):799–804.

44. Ebel ER, Kim BY, McDew-White M, Egan ES, Anderson TJC, Petrov DA. Antigenic
diversity in malaria parasites is maintained on extrachromosomal DNA. bioRxiv. 2023 Jan
1;2023.02.02.526885.

45. Girgis HS, DuPai CD, Lund J, Reeder J, Guillory J, Durinck S, et al. Single-molecule
nanopore sequencing reveals extreme target copy number heterogeneity in arylomycin-
resistant mutants. Proc Natl Acad Sci U S A. 2021 Jan 5;118(1):e2021958118.

46. Grigorev K, Foox J, Bezdan D, Butler D, Luxton JJ, Reed J, et al. Haplotype diversity and
sequence heterogeneity of human telomeres. Genome Res. 2021 Jul;31(7):1269–79.

47. Dixon K, Shen Y, O'Neill K, Mungall KL, Chan S, Bilobram S, et al. Defining the
heterogeneity of unbalanced structural variation underlying breast cancer susceptibility by
nanopore genome sequencing. Eur J Hum Genet. 2023 May;31(5):602–6.

48. Pholwat S, Liu J, Stroup S, Jacob ST, Banura P, Moore CC, et al. The Malaria TaqMan
Array Card Includes 87 Assays for Plasmodium falciparum Drug Resistance, Identification
of Species, and Genotyping in a Single Reaction. Antimicrobial Agents and Chemotherapy
[Internet]. 2017 May 1 [cited 2020 Jul 29];61(5). Available from:
https://aac.asm.org/content/61/5/e00110-17

49. Walliker D, Quakyi IA, Wellems TE, McCutchan TF, Szarfman A, London WT, et al.
Genetic analysis of the human malaria parasite Plasmodium falciparum. Science. 1987 Jun
26;236(4809):1661–6.

50. Wellems TE, Panton LJ, Gluzman IY, do Rosario VE, Gwadz RW, Walker-Jonah A, et al.
Chloroquine resistance not linked to mdr-like genes in a Plasmodium falciparum cross.
Nature. 1990 May 17;345(6272):253–5.

51. Chulay JD, Watkins WM, Sixsmith DG. Synergistic Antimalarial Activity of Pyrimethamine
and Sulfadoxine against Plasmodium falciparum In Vitro. The American Journal of Tropical
Medicine and Hygiene. 1984 May 1;33(3):325–30.

52. Watkins WM, Sixsmith DG, Chulay JD, Spencer HC. Antagonism of sulfadoxine and
pyrimethamine antimalarial activity in vitro by p-aminobenzoic acid, p-
aminobenzoylglutamic acid and folic acid. Molecular and Biochemical Parasitology. 1985
Jan 1;14(1):55–61.

53. Wang P, Brobey RKB, Horii T, Sims P, Hyde JE, Schapira A, et al. Utilization of exogenous
folate in the human malaria parasite Plasmodium falciparum and its critical role in antifolate
drug synergy The Susceptibility of Plasmodium Falciparum to Sulfadoxine and
Pyrimethamine: Correlation of in Vivo and in Vitro Results Synergistic Antimalarial
Activity of Pyrimethamine and Sulfadoxine against Plasmodium falciparum In Vitro.
Molecular Microbiology. 1986 Mar 1;32(2):239–45.

54. Salcedo-Sora JE, Ochong E, Beveridge S, Johnson D, Nzila A, Biagini GA, et al. The
Molecular Basis of Folate Salvage in Plasmodium falciparum: CHARACTERIZATION OF
TWO FOLATE TRANSPORTERS*. Journal of Biological Chemistry. 2011 Dec
30;286(52):44659–68.

55. Valenciano AL, Fernández-Murga ML, Merino EF, Holderman NR, Butschek GJ, Shaffer
KJ, et al. Metabolic dependency of chorismate in Plasmodium falciparum suggests an
alternative source for the ubiquinone biosynthesis precursor. Scientific Reports. 2019 Sep
26;9(1):13936.

56. Kumar S, Li X, McDew-White M, Reyes A, Delgado E, Sayeed A, et al. A Malaria Parasite
Cross Reveals Genetic Determinants of Plasmodium falciparum Growth in Different Culture
Media. Front Cell Infect Microbiol. 2022 May 30;12:878496.

57. Llanos-Cuentas A, Casapia M, Chuquiyauri R, Hinojosa JC, Kerr N, Rosario M, et al.
Antimalarial activity of single-dose DSM265, a novel plasmodium dihydroorotate
dehydrogenase inhibitor, in patients with uncomplicated Plasmodium falciparum or
Plasmodium vivax malaria infection: a proof-of-concept, open-label, phase 2a study. Lancet
Infect Dis. 2018 Aug;18(8):874–83.

58. Ravenhall M, Benavente ED, Mipando M, Jensen ATR, Sutherland CJ, Roper C, et al.
Characterizing the impact of sustained sulfadoxine/pyrimethamine use upon the Plasmodium
falciparum population in Malawi. Malar J. 2016 Nov 29;15:575.

59. Moon RW, Sharaf H, Hastings CH, Ho YS, Nair MB, Rchiad Z, et al. Normocyte-binding
protein required for human erythrocyte invasion by the zoonotic malaria parasite
Plasmodium knowlesi. Proc Natl Acad Sci U S A. 2016 Jun 28;113(26):7231–6.

60. Phillips MA, Gujjar R, Malmquist NA, White J, El Mazouni F, Baldwin J, et al.
Triazolopyrimidine-based dihydroorotate dehydrogenase inhibitors with potent and selective
activity against the malaria parasite Plasmodium falciparum. J Med Chem. 2008/06/04 ed.
2008;51(12):3649–53.

61. Haynes JD, Diggs CL, Hines FA, Desjardins RE. Culture of human malaria parasites
Plasmodium falciparum. Nature. 1976 Oct 1;263(5580):767–9.

62. Bei AK, Desimone TM, Badiane AS, Ahouidi AD, Dieye T, Ndiaye D, et al. A flow
cytometry-based assay for measuring invasion of red blood cells by Plasmodium falciparum.
Am J Hematol. 2010 Apr;85(4):234–7.

63. Liu S, Huckaby AC, Brown AC, Moore CC, Burbulis I, McConnell MJ, et al. Single-cell
sequencing of the small and AT-skewed genome of malaria parasites. Genome Medicine.
2021 May 4;13(1):75.

64. Bushnell B. BBMap [Internet]. [cited 2021 Dec 1]. Available from:
http://sourceforge.net/projects/bbmap/ (2019)

65. Chiang C, Layer RM, Faust GG, Lindberg MR, Rose DB, Garrison EP, et al. SpeedSeq:
ultra-fast personal genome analysis and interpretation. Nat Methods. 2015 Oct;12(10):966–8.

66. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078–9.

67. Layer RM, Chiang C, Quinlan AR, Hall IM. LUMPY: a probabilistic framework for
structural variant discovery. Genome Biology. 2014 Jun 26;15(6):R84.

68. Abyzov A, Urban AE, Snyder M, Gerstein M. CNVnator: an approach to discover, genotype,
and characterize typical and atypical CNVs from family and population genome sequencing.
Genome Res. 2011/02/07 ed. 2011;21(6):974–84.

69. Maniatis T, Sambrook J, Fritsch EF. Molecular cloning: a laboratory manual. Cold Spring
Harbor, NY: Cold Spring Harbor Laboratory Press; 1989. xxxviii + 1546 pp.

70. Oxford Nanopore Technologies. qcat (version 1.1.0) [Internet]. [cited 2021 Jun 1]. Available
from: https://github.com/nanoporetech/qcat

71. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34(18):3094–100.

72. García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, et al.
Qualimap: evaluating next-generation sequencing alignment data. Bioinformatics. 2012 Aug
22;28(20):2678–9.

73. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nat Biotechnol. 2011 Jan;29(1):24–6.

74. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. Briefings in Bioinformatics. 2013
Mar 1;14(2):178–92.

75. Ye J, McGinnis S, Madden TL. BLAST: improvements for better sequence analysis. Nucleic
Acids Research. 2006;34(suppl_2):W6–9.

Supporting Information Captions

Figure S1: Spanning reads carry odd copy numbers of *GCH1* amplicons

Figure S2: Long-read visualization of *GCH1* amplicon from *3D7* parasite line

Figure S3: Parasites lines with *DHODH* amplicons are selected from parent lines with *GCH1*

amplicons

Figure S4: *GCH1* amplicon junctions show signs of primary and secondary expansion events

Supplemental Methods: Procedures used for the preparation of long read data by Ellen Yeh's

group, Stanford University, as presented in Table S2, including in vitro cultures, mutagenesis,

DNA extraction, and sequencing.

Table S1: Oxford Nanopore sequencing datasets new to this manuscript

Table S2: Quantification of long reads covering the *GCH1* amplicon displays greater variability in

amplicon number from DSM1 resistant parasites

Table S3: *DHODH* and *GCH1* copy number in parasites resistant to DHODH inhibitors

Table S4: Detection of known amplifications by Illumina short reads from parasites resistant to

DHODH inhibitors

Table S5: ddPCR primer and probe sequences