JUMP: statistical analysis of replicability across two studies

Pengfei Lyu[1†], Yan Li[2†] and Hongyuan Cao[1*]

Abstract

Motivation: Replicability is the cornerstone of scientific research. The current statistical method for high-dimensional replicability analysis either cannot control the false discovery rate (FDR) or is too conservative.

Results: We propose a statistical method, JUMP, for the high-dimensional replicability analysis of two studies. The input is a high dimensional paired sequence of p-values from two studies and the test statistic is the maximum of p-values of the pair. JUMP uses four states of the p-value pairs to indicate whether they are null or non-null. Conditional on the hidden states, JUMP computes the cumulative distribution function of the maximum of p-values for each state to conservatively approximate the probability of rejection under the composite null of replicability. JUMP estimates unknown parameters and uses a step-up procedure to control FDR. By incorporating different states of composite null, JUMP achieves a substantial power gain over existing methods while controlling the FDR. Analyzing two pairs of spatially resolved transcriptomic datasets, JUMP makes biological discoveries that otherwise cannot be obtained by using existing

[1]Department of Statistics, Florida State University, Tallahassee, FL 32306, USA. [2]School of Mathematics, Jilin University, Changchun, Jilin 130012, China.
[†]The first two authors should be regarded as Joint First Authors.
[*]Corresponding author: hongyuancao@gmail.com

methods.

Availability: The proposed method is implemented in the R package JUMP available at `https://github.com/hongyuan-cao/JUMP`.

1 Introduction

Spatially resolved transcriptomics (SRT) links the transcriptomes to their cellular locations, providing a comprehensive understanding of gene expression profiles in the spatial context of biological systems (Kleino et al., 2022). Multiple spatial transcriptomics technologies emerged in the past few years. These spatial transcriptomic technologies include those that are based on single molecular fluorescent *in situ* hybridization (smFISH) (Raj et al., 2008), those that are based on microdissection techniques (Bidarimath et al., 2015), and those that are based on *in situ* mRNA capturing followed by high-throughput sequencing techniques (Lubeck and Cai, 2012) such as spatial transcriptome (ST) (Ståhl et al., 2016), Slide-seq (Rodriques et al., 2019), and high-definition spatial transcriptomics (Vickovic et al., 2019).

An important first step toward characterizing the spatial transcriptomic landscape of complex tissues is identifying genes that display spatial expression patterns. Several computational methods have been developed for transcriptome-wide identification of spatially variable genes (SVGs) from SRT data. Examples include Trendsceek (Edsgärd et al., 2018), SpatialDE (Svensson et al., 2018), SPARK (Sun et al., 2020), SPARK-X (Zhu et al., 2021), and SpaGCN (Hu et al., 2021), among others. These methods get p-values for each gene and adjust for multiple comparisons to claim statistical significance. A classic method for multiple comparisons is the false discovery rate (FDR) control approach proposed in Benjamini and Hochberg (1995) (BH). The false discovery rate is defined as the expected value of the ratio of false rejections over total rejections. Suppose we have m hypotheses. The BH procedure works as follows. First, order p-values $p_{(1)} \leq \ldots \leq p_{(m)}$. Second, find the largest i_0 such that

$p_{(i_0)} \leq \alpha i_0 / m$. Third, reject hypotheses corresponding to $p_{(1)}, \ldots, p_{(i_0)}$. Under the assumption that p-values from the null are independent and follow standard uniform distribution, it is shown that FDR is controlled at level $\alpha \pi_0$, where π_0 is the proportion of null hypotheses. The BH procedure is robust under positive dependence among p-values under the null (Benjamini and Yekutieli, 2001).

To ensure the quality and rigor of scientific findings, replicability is indispensable. One definition of replicability is to obtain consistent results across different studies to answer the same scientific question (Hung and Fithian, 2020; Bogomolov and Heller, 2022). For SVGs detection in SRT studies, an *ad hoc* approach to have replicability is to first compute p-values for each study. A multiple comparison method such as BH can be used to claim significance in each study. Replicable SVGs are those that are significant in both studies. This seems to be conservative. Yet it does not have FDR control. Intuitively speaking, if there is no danger that a multiple testing procedure produces false positives, then this *ad hoc* approach would work. However, multiple testing procedures have a non-zero probability of making false positives unless the procedure does not reject. Thus, an approach that provides control over false positives in each study separately does not guarantee control of overall false positives to test replicability (Bogomolov and Heller, 2013). The partial conjunction approach in Benjamini et al. (2009) suggests applying a multiple testing procedure, such as BH, on the maximum of p-values across two studies. This provides effective FDR control yet the power is low.

In this paper, we propose a joint super-uniform maximum p-value (JUMP) method that improves on the maximum of p-values for replicability analysis of SVGs in two SRT studies. The null hypothesis for replicability is composite and consists of three states. The vanilla version of the maximum of p-values does not incorporate different states of the null and has a cumulative distribution function far smaller than that of standard uniform distribution, causing power loss. We use four states for the p-value pairs indicating whether they are null or non-null (Chung et al., 2014). Conditional on the hidden states, we compute the cumulative

distribution function of the maximum of p-values for each state separately. Combining with the proportion estimation for each state, we get a conservative approximation of the cumulative distribution function of the maximum of p-values under replicability null. To estimate the proportion of each state, we extend the proportion of null estimation in Storey et al. (2004) from the two-group model to a four-group model. We require p-values in each sequence to follow a standard uniform distribution under the null. This assumption is needed for the BH procedure and Storey et al. (2004)'s method of the proportion of null estimation. A step-up procedure is developed on the maximum of p-values to control FDR. By incorporating the composite null feature of replicability analysis, we achieve a substantial power gain. The computation is scalable with a computational cost similar to that of BH.

As a proof-of-concept, we applied JUMP to two pairs of SRT datasets collected from different tissue sections to identify replicable SVGs: mouse olfactory bulb data measured with ST technology (Ståhl et al., 2016) and 10X Visium technology; mouse cerebellum data measured with Slide-seq technology (Rodriques et al., 2019) and Slide-seqV2 technology (Stickels et al., 2021). We show that at the same FDR level, JUMP identifies important replicable SVGs that would otherwise be missed by using existing methods.

2 Methods

2.1 Background and notations

Suppose we have m genes common to two SRT studies. We are interested in identifying genes that display replicable spatial expression patterns. The input of the replicability analysis is a pair of p-values from two studies $(p_{1i}, p_{2i}), i = 1, \ldots, m$. The null hypothesis for ith gene states that it does not show any replicable spatial expression pattern. Let θ_{ji} denote underlying state of ith gene in study j ($j = 1, 2$), where $\theta_{ji} = 1$ indicates ith gene is an SVG in study j and

$\theta_{ji} = 0$ otherwise. A four-group model is assumed for the paired p-value sequence, i.e.,

$$p_{1i} \mid \theta_{1i} \sim (1 - \theta_{1i})f_0 + \theta_{1i}f_1,$$

$$p_{2i} \mid \theta_{2i} \sim (1 - \theta_{2i})f_0 + \theta_{2i}f_2, \quad i = 1, \ldots, m,$$

where f_0 is the density function of p-values under the null, and f_1 and f_2 denote the non-null density functions of p-values from study 1 and study 2, respectively. Let $\tau_i = (\theta_{1i}, \theta_{2i}), i = 1, \ldots, m$ denote the joint hidden states across two studies. Then $\tau_i \in \{(0,0), (0,1), (1,0), (1,1)\}$ with $\mathbb{P}(\tau_i = (k,l)) = \xi_{kl}$ for $k, l \in \{0,1\}$, and $\sum_{k,l} \xi_{kl} = 1$. Based on this four-group model, the replicability null hypothesis for ith gene can be defined as

$$H_{i0} : \tau_i \in \{(0,0), (0,1), (1,0)\}, i = 1, \ldots, m,$$

where the ith gene is replicable if it shows spatial expression patterns in both study 1 and study 2. For simplicity, we denote the hidden state of H_{i0} as h_i, where $h_i = 0$ indicates H_{i0} is true and $h_i = 1$ otherwise. Hence $\mathbb{P}(h_i = 0) = \xi_{00} + \xi_{01} + \xi_{10}$ and $\mathbb{P}(h_i = 1) = \xi_{11}$.

2.2 JUMP for replicability analysis

The schematic of JUMP for identifying replicable SVGs from two SRT studies is shown in Fig. 1. After obtaining paired p-values from two SRT studies, we define the maximum p-values as

$$p_i^{\max} = \max\{p_{1i}, p_{2i}\}, \quad i = 1, \ldots, m.$$

Figure 1: Schematic of JUMP for identifying replicable SVGs from two SRT studies

Under the assumption that f_0 follows the standard uniform distribution, we have

$$\mathbb{P}(p_i^{\max} \le t | \tau_i = (0,0)) = t^2,$$

$$\mathbb{P}(p_i^{\max} \le t | \tau_i = (0,1)) \le t,$$

$$\mathbb{P}(p_i^{\max} \le t | \tau_i = (1,0)) \le t, \quad i = 1, \dots, m,$$

where we use the assumption that conditional on the joint hidden states, two p-value sequences are independent.

Under the replicable null, $p_i^{\max}$ follows a two-group mixture model

$$p_i^{\max} \mid h_i \sim (1 - h_i)g_0 + h_i g_1,$$

where g_0 and g_1 denote the density function of $p_i^{\max}$ under replicable null and non-null,

respectively. For any $t \in (0, 1)$, we compute the cumulative distribution function of $p_i^{\max}$ under replicable null as follows.

$$
\begin{aligned}
&\mathbb{P}(p_i^{\max} \leq t \mid h_i = 0) \\
&= \frac{\mathbb{P}(p_i^{\max} \leq t, h_i = 0)}{\mathbb{P}(h_i = 0)} \\
&= \frac{\xi_{00} \mathbb{P}\left(p_i^{\max} \leq t | \tau_i = (0, 0)\right)}{\xi_{00} + \xi_{01} + \xi_{10}} + \frac{\xi_{01} \mathbb{P}\left(p_i^{\max} \leq t | \tau_i = (0, 1)\right)}{\xi_{00} + \xi_{01} + \xi_{10}} + \frac{\xi_{10} \mathbb{P}\left(p_i^{\max} \leq t | \tau_i = (1, 0)\right)}{\xi_{00} + \xi_{01} + \xi_{10}} \\
&\leq \frac{\xi_{00} t^2 + (\xi_{01} + \xi_{10}) t}{\xi_{00} + \xi_{01} + \xi_{10}}
\end{aligned}
$$

Denote

$$
G(t) = \frac{\xi_{00} t^2 + (\xi_{01} + \xi_{10}) t}{\xi_{00} + \xi_{01} + \xi_{10}}, \tag{1}
$$

we have

$$
G(t) - t = \frac{\xi_{00}}{\xi_{00} + \xi_{01} + \xi_{10}} t(t - 1) \leq 0.
$$

Hence $p_i^{\max}$ follows a super-uniform distribution under the replicable null. This verifies that the vanilla maximum p-value method has valid FDR control.

For $t \in (0, 1)$, the number of rejections is $R(t) = \sum_{i=1}^{m} I\{p_i^{\max} \leq t\}$, and the number of false rejections is bounded by $m(\xi_{00} + \xi_{01} + \xi_{10}) G(t)$. At threshold t, we have a conservative estimate of FDR by

$$
\mathrm{FDR}^*(t) = \frac{m(\xi_{00} + \xi_{01} + \xi_{10}) G(t)}{R(t) \vee 1}.
$$

In the oracle case that we know ξ_{00}, ξ_{01} and ξ_{10}, at FDR level α, let

$$
t_m = \sup\{t \in (0, 1) : \mathrm{FDR}^*(t) \leq \alpha\}. \tag{2}
$$

101   Reject H_i if $p_i^{\max} \leq t_m$.

2.3   Estimation of unknowns

As ξ_{00}, ξ_{01}, and ξ_{10} are unknown in practice, we provide their estimates in this section. Following Storey (2002) and Storey et al. (2004), in study j, the proportion of null hypotheses, $\pi_0^{(j)}$, can be estimated by

$$\hat{\pi}_0^{(j)}(\lambda_j) = \frac{\sum_{i=1}^m I\{p_{ji} \geq \lambda_j\}}{m(1 - \lambda_j)}, j = 1, 2.$$

Similarly, we estimate ξ_{00} as

$$\hat{\xi}_{00}(\lambda_3) = \frac{\sum_{i=1}^m I\{p_{1i} \geq \lambda_3, p_{2i} \geq \lambda_3\}}{m(1 - \lambda_3)^2},$$

103   where λ_1, λ_2 and λ_3 are tuning parameters. We use the smoothing method provided in Storey

104   and Tibshirani (2003) to select tuning parameters. Please see Supplementary Notes A for

105   more details. We have

$$\hat{\xi}_{01} = \hat{\pi}_0^{(1)} - \hat{\xi}_{00},$$

$$\hat{\xi}_{10} = \hat{\pi}_0^{(2)} - \hat{\xi}_{00}.$$

We estimate $\mathrm{FDR}^*(t)$ by

$$\widehat{\mathrm{FDR}}^*(t) = \frac{m(\hat{\xi}_{00}t^2 + \hat{\xi}_{01}t + \hat{\xi}_{10}t)}{R(t) \vee 1}.$$

106   The data-adaptive thresholding is

$$\hat{t}_m = \sup\{t \in (0, 1) : \widehat{\mathrm{FDR}}^*(t) \leq \alpha\}. \tag{3}$$

We claim the replicability of ith gene if $p_i^{\max} \le \hat{t}_m$.

This is equivalent to the following step-up procedure. First, order the maximum p-values $p_{(1)}^{\max} \le \cdots \le p_{(m)}^{\max}$. Second, find

$$\hat{k} := \max\left\{ 1 \le i \le m : \widehat{\mathrm{FDR}}^* \left(p_{(i)}^{\max} \right) \le \alpha \right\}. \tag{4}$$

Reject $H_{(i)}$ for $i = 1, \ldots, \hat{k}$, where $H_{(i)}$ corresponds to $p_{(i)}^{\max}$.

The key to the power gain is to incorporate different states in the composite null. This is similar in spirit to plugging in the proportion of the null hypothesis with a single p-value sequence in Storey (2002).

3   Results

In this section, we evaluate the FDR control and power of JUMP via realistic simulations and conduct data analysis to identify replicable SVGs from two pairs of SRT datasets measured with different technologies.

3.1   Simulation studies

We conducted extensive simulations based on Replicate 9 and Replicate 12 of ST datasets measured in mouse olfactory bulb (Ståhl et al., 2016). The parameters used for data generation were inferred from SPARK (Sun et al., 2020). Power is defined as the expectation of true replicable discoveries over the total number of non-null hypotheses. We compare JUMP with two other methods for replicability analysis, BH and MaxP. Here BH refers to a naïve method that applies the BH procedure (Benjamini and Hochberg, 1995) on each study and identifies genes significant in both studies as replicable SVGs. MaxP (Benjamini et al., 2009) directly applies the BH procedure (Benjamini and Hochberg, 1995) to the maximum p-values across

two studies, $p_i^{\max}$.

In each simulation, a pair of count data is generated for m genes following the simulation

strategy in Sun et al. (2020). States of genes in two SRT studies, θ_{1i} and θ_{2i}, were generated

from a multinomial distribution with probabilities, $\mathbb{P}(\theta_{1i} = k, \theta_{2i} = l) = \xi_{kl}, k, l \in \{0, 1\}$,

for pre-specified $\xi_{00}, \xi_{01}, \xi_{10}$ and ξ_{11}. After obtaining θ_{ji} for $i = 1, \ldots, m$ and $j = 1, 2$, we

simulated gene count matrices based on corresponding ST data and parameters with different

signal strengths (moderate or strong) and different standard deviations for the error term (τ_j

$= 0.3$ or 0.5). Details of the data generation process can be found in Supplementary Notes B.2.

Figure 2: FDR control and power comparison of different methods in realistic simulations. Simulations were performed under the setting of $m = 10,000, \xi_{11} = 0.05$ and $\xi_{01} = \xi_{10}$. The signal strengths were set to be strong for study 1 and moderate for study 2. Each column corresponds to a different ξ_{00} setting. Each row corresponds to a different spatial expression pattern on which the paired count data were generated. Patterns I-III for two studies are shown in Supplementary Fig. S3. In each panel, the empirical FDR and power of different methods were calculated at a target FDR level of 0.05 (horizontal dashed line in the plots of FDR) for different standard deviations (left: $\tau_1 = 0.3, \tau_2 = 0.5$; right: $\tau_1 = 0.5, \tau_2 = 0.3$).

135   Let $m = 10,000, \xi_{11} = 0.05$ and $\xi_{01} = \xi_{10}$. We set a strong signal for study 1 and a

moderate signal for study 2. For a given value of ξ_{00}, corresponding ξ_{01} and ξ_{10} can be obtained by $\xi_{01} = \xi_{10} = (1 - \xi_{00} - \xi_{11})/2$. Fig. 2 shows the FDR control and power comparison of different methods across different settings. We observe that MaxP and JUMP controlled the FDR at the nominal level across all settings, and JUMP is more powerful than maxP. BH is not valid in practice since it failed to control the FDR in some settings (e.g., $\xi_{00} = 0.5$). The power increased for all methods from Pattern I to Pattern III. By examining the three spatial expression patterns on which the corresponding data were generated (Supplementary Fig. S3), we speculate that this might be due to the increasing spatial variability from Pattern I to Pattern III. More results of simulation studies can be found in Supplementary Fig. S1, S2, and S4.

3.2 Analysis of mouse olfactory bulb data

We first analyzed the SRT data from mouse olfactory bulb measured with ST technology (Ståhl et al., 2016) and 10X Visium technology. Ståhl et al. (2016) published 12 replicates of the mouse olfactory bulb ST datasets on the Spatial Research Website (`https://www.spatialresearch.org/`). We used Replicate 9 for our analysis, which includes $15,284$ genes on 237 spots. The 10X Visium dataset was downloaded from the 10X Visium spatial gene expression repository (`https://www.10xgenomics.com/resources/datasets`) and contains $32,285$ genes on $1,185$ spots. We filtered out genes that are expressed in less than 10% spatial spots and selected spots with at least ten total read counts, resulting in $9,547$ genes on 236 spots for the ST dataset and $10,680$ genes on $1,185$ spots for the 10X Visium dataset, respectively. We separately analyzed the two datasets using SPARK (Sun et al., 2020) to produce p-values. Paired p-values of $8,547$ genes common to both studies is the input to the replicability analysis. As can be seen in Fig. 3a, JUMP has higher power than the other two methods. At the FDR level of 0.05, MaxP identified 618 replicable SVGs, which were all

Figure 3: Analysis and validation results of the mouse olfactory bulb data measured with ST technology and 10X Visium technology. (a) Venn diagram shows the number of replicable SVGs identified by different methods at FDR level 0.05 and the intersection of discoveries. (b) Three distinct spatial expression patterns based on the 807 replicable SVGs identified by JUMP in the ST study (top) and 10X Visium study (bottom). Color represents relative expression levels (antique white: low; navy blue: high). The corresponding hematoxylin and eosin staining (HE) images for the two studies are shown in the right panel. (c) Spatial expression patterns of three representative genes identified by JUMP, corresponding to Pattern I-III, respectively. *In situ* hybridization images of corresponding genes obtained from the Allen Brain Atlas are shown in the top panel. (d) The box plot shows Moran's *I* statistic of the 189 replicable SVGs additionally identified by JUMP and that of all genes based on the ST study (left) and the 10X Visium study (right). (e) The bar chart displays the number of replicable SVGs additionally identified by JUMP and BH compared to that identified by all three methods. They were validated in two reference gene lists from the Harmonizome database: one from the Allen Brain Atlas dataset and the other from the BioGPS dataset. (f) The bubble plot shows the GO enrichment analysis result of JUMP. The horizontal dashed line represents the FDR level 0.01. The color of a bubble shows different GO term categories: BP (tan), CC (green), and MF (blue). The size of a bubble represents the counts of corresponding gene sets.

detected by BH and JUMP. JUMP identified 807 replicable SVGs, including 637 of the 638 SVGs detected by BH.

We clustered the 807 replicable SVGs identified by JUMP into three groups using the hierarchical agglomerative clustering algorithm implemented in the R package *amap* (v0.8-18) and summarized the spatial expression patterns based on the expression level of the genes in corresponding groups. In both studies, the summarized patterns were consistent with three main cell layers in mouse olfactory bulb. In Fig. 3b and Fig. S5a, Pattern I corresponds to the glomerular layer, Pattern II corresponds to the mitral layer and Pattern III corresponds to the granular layer. Spatial expression patterns of three representative SVGs (*Kctd12*, *Plcxd2*, *Ctxn1*) identified by JUMP are presented in Fig. 3c, which corresponds to Pattern I-III, respectively and consistent with the *in situ* hybridization images obtained from the Allen Brain Atlas. We also randomly selected 30 genes from the 189 additional findings of JUMP compared to the overlaps of three methods and showed their spatial expression patterns in Supplementary Fig. S6a, b. Moreover, we calculated Moran's I statistic (Moran, 1950) of the 189 replicable SVGs additionally identified by JUMP and compared it with that using all $8,547$ genes to show the autocorrelations of the additional findings by JUMP (Fig. 3d).

To further validate the identified replicable SVGs, we downloaded two published gene sets from the Harmonizome database (Rouillard et al., 2016) consisting of genes related to mouse olfactory bulb as a reference (Fig. 3e). The first gene set was summarized based on three layers (glomerular, mitral, and granular) of the main olfactory bulb from the Allen Brain Atlas adult mouse brain tissue gene expression profiles dataset (Sunkin et al., 2012), including $3,485$ genes with high or low expression in main olfactory bulb relative to other tissues. 33% of the 618 replicable SVGs that were identified by all three methods were validated. Among the 189 replicable SVGs additionally identified by JUMP, 26% were validated in the reference list, whereas only 4 of the 20 SVGs additionally identified by BH were in the same list. The second reference gene set includes $2,031$ genes differentially expressed in mouse olfactory bulb

relative to other cell types and tissues from the BioGPS mouse cell type and tissue gene expression profiles dataset (Wu et al., 2013). In addition to the replicable SVGs identified by all three methods (34% validated), 38 of the 189 replicable SVGs only detected by JUMP were in the list, whereas 4 of the 20 replicable SVGs only detected by BH were validated in the same list. These results provide additional biological evidence for the improved power of JUMP.

Additionally we performed Gene Ontology (GO) enrichment analysis on replicable SVGs identified by JUMP and BH (Fig. 3f). At the FDR level of 0.01, JUMP enriched 846 GO terms and BH enriched 764 GO terms (708 overlaps). Many of the 138 GO terms only identified by JUMP are related to nervous system development and olfactory bulb organization. For instance, transmission of nerve impulse (GO:0019226), neuronal action potential (GO:0019228), and forebrain neuron differentiation (GO:0021879) are closely related to the establishment of axodendritic and dendrodendritic synaptic contacts within the olfactory bulb (Belluzzi et al., 2003); GABA-ergic synapse (GO:0098982) plays a key role in the organization of olfactory bulb (Hanson et al., 2020); GO terms GO:0045744 and GO:0002029 are related to G protein-coupled receptor, which can be encoded by odorant receptor genes differentially expressed at conserved positions in the olfactory bulb (Katidou et al., 2018).

3.3 Analysis of mouse cerebellum data

We next analyzed the SRT data from mouse cerebellum measured with Slide-seq technology (Rodriques et al., 2019) and Slide-seqV2 technology (Stickels et al., 2021). The two datasets were obtained from Broad Institute's single-cell repository (`https://singlecell.broadinstitute.org/single_cell`) with IDs SCP354 and SCP948, respectively. The Slide-seq dataset (file "Puck_180430_6") contains $18,671$ genes on $25,551$ beads. we filtered out beads with total counts less than 50. The Slide-seqV2 dataset contains $23,096$ genes on $39,496$

beads. We cropped regions of interest by filtering out beads with total counts less than 100.

Mitochondrial genes and genes that are not expressed in any locations were filtered out from

the two datasets, resulting in $17,481$ genes on $14,667$ beads for the Slide-seq data and $20,117$

genes on $11,626$ beads for the Slide-seqV2 data. After applying the SPARK-X method on the

two datasets separately, we obtained two sequences of p-values and matched them by gene.

We used the paired p-values for $16,519$ genes common to both studies as input for the repli-

cability analysis of SVGs. At FDR level 0.05, MaxP identified 279 replicable SVGs, which

were all identified by BH and JUMP. JUMP detected all BH findings (394) and identified 54

additional replicable SVGs.

We first examined the spatial expression patterns of the 448 replicable SVGs identified

by JUMP by clustering them into three groups showing distinct spatial expression patterns

(Fig. 4a). In both Slide-seq (left) and Slide-seqV2 (right) datasets, the 448 SVGs showed con-

sistent patterns: Pattern I and Pattern III correspond to the purkinje cell layer and granular

cell layer, respectively; and Pattern II corresponds to other cell layers. Three representative

genes identified by JUMP (*Pcp2*, *Mbp*, and *Snap25*) exhibited corresponding spatial expres-

sion patterns, which were validated by *in situ* hybridization in the Allen Brain Atlas (Fig. 4b).

Fig. S7a,b display spatial expression patterns of 24 genes randomly selected from the 169 repli-

cable SVGs identified by JUMP in addition to genes identified by all three methods. As shown

in Fig. 4d, the additional spatial autocorrelations revealed by JUMP were further confirmed

by Moran's I (Moran, 1950).

We provided two gene sets obtained from published literature to validate the results. First,

we obtained a list of genes that are highly differentially expressed across all clusters in mouse

cerebellar cortex from Kozareva et al. (2021). We further filtered out genes with absolute

log fold change smaller than 0.05 and obtained $3,976$ genes for the validation. In addition

to genes identified by all three methods (77% validated), 107 of the 169 replicable SVGs

additionally identified by JUMP were in the list, whereas 73 of the 115 findings by BH were

Figure 4: Analysis and validation results of the mouse cerebellum data measured with Slide-seq technology and Slide-seqV2 technology. (a) Three distinct spatial expression patterns based on the 448 replicable SVGs identified by JUMP in the Slide-seq study (left) and Slide-seqV2 study (right). Color represents relative expression levels (antique white: low; navy blue: high). (b) Spatial expression patterns of three representative genes identified by JUMP, corresponding to Pattern I-III, respectively. *In situ* hybridization images of corresponding genes obtained from the Allen Brain Atlas are shown in the top panel. (c) The bar chart displays the number of replicable SVGs additionally identified by JUMP and BH compared to that identified by all three methods. They were validated in two reference gene lists: one from Kozareva et al. (2021) and the other from the Allen Brain Atlas dataset summarized in the Harmonizome database. (d) The box plot shows Moran's *I* statistic of the 169 replicable SVGs additionally identified by JUMP and that of all genes based on the Slide-seq study (left) and the Slide-seqV2 study (right). (e) The bubble plot shows the GO enrichment analysis result of JUMP. The horizontal dashed line represents the FDR level 0.01. The color of a bubble shows different GO term categories: BP (tan), CC (green), and MF (blue). The size of a bubble represents the counts of corresponding gene sets.

validated. Second, we downloaded three gene sets related to mouse cerebellum in the Allen Brain Atlas datasets (Sunkin et al., 2012) from the Harmonizome database (Rouillard et al., 2016) and summarized them to a list of 3, 000 genes that are differentially expressed in mouse cerebellum, cerebellar cortex, and cerebellar hemisphere. Among the 279 SVGs identified by all three methods, 30% were in this gene list. 39 of the 169 replicable SVGs additionally identified by JUMP were validated, whereas 28 of the 115 SVGs additionally identified by BH were in the same list.

Finally, we performed GO enrichment analysis on the replicable SVGs identified by different methods to evaluate the biological significance. At the FDR level of 0.01, JUMP enriched 452 GO terms and BH identified 418 GO terms (394 overlaps). Among the 58 GO terms only enriched by JUMP, many are closely related to the structural constitution and functional development of mouse cerebellum, such as GO terms of cerebellar cortex development (GO:0021695), cerebellum development (GO:0021549), central nervous system neuron development (GO0021954), dendritic transport (GO:0098935), and photoreceptor ribbon synapse (GO:0098684), among others.

4 Discussion

We present a new method, JUMP, for identifying replicable SVGs from two SRT studies. JUMP models the joint hidden states and calculates rejection probabilities under each state. This provides a better approximation to the probability of rejection under a composite replicable null hypothesis. Analysis of different SRT studies identifies important replicable SVGs that might otherwise be missed by existing methods. JUMP is simple to implement and computationally scalable to tens of thousands of genes (Supplementary Table S1 and S2). Moreover, JUMP does not require two SRT studies to have the same spatial alignment or the same tissue thickness. In practice, using alignments, rotations and normalizations may distort

important biological information contained in the data. In addition, JUMP is flexible and can accommodate data from other modalities, such as scRNA-seq, ATAC-seq, and CITE-seq, among others.

One limitation of JUMP is that it only identifies replicable SVGs from two SRT studies. If we want to extend it to more than two studies, say n studies, we require a 2^n-group model for the data, and the composite null is composed of $(2^n - 1)$ hidden joint states whose proportions need to be estimated. We leave this for future research.

Data availability

The data underlying this article are publicly available in corresponding repositories. The mouse olfactory bulb ST dataset can be obtained with file 'MOB Replicate 9' in the Spatial Research Website at `https://www.spatialresearch.org/resources-published-datasets/doi-10-1126science-aaf2403/`. The mouse olfactory bulb 10X Visium dataset is available in the 10X Visium spatial gene expression repository at `https://www.10xgenomics.com/resources/datasets/adult-mouse-olfactory-bulb-1-standard-1`. The mouse cerebellum Slide-seq dataset and Slide-seqV2 dataset are provided in the Broad Institute's single-cell repository at `https://singlecell.broadinstitute.org/single_cell/` under IDs SCP354 (file 'Puck_180430_6') and SCP948, respectively.

Acknowledgements

This work was partially supported by the China Postdoctoral Science Foundation (No. 801212021410) to Y. L.

Competing interests

The authors declare that they have no competing interests.

References

O. Belluzzi, M. Benedusi, J. Ackman, and J. J. LoTurco. Electrophysiological differentiation of new neurons in the olfactory bulb. *Journal of Neuroscience*, 23(32):10411–10418, 2003.

Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300, 1995.

Y. Benjamini and D. Yekutieli. The control of the false discovery rate in multiple testing under dependency. *Annals of statistics*, pages 1165–1188, 2001.

Y. Benjamini, R. Heller, and D. Yekutieli. Selective inference in complex research. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 367(1906):4255–4271, 2009.

M. Bidarimath, A. K. Edwards, and C. Tayade. Laser capture microdissection for gene expression analysis. *Methods Mol Biology*, pages 115–137, 2015.

M. Bogomolov and R. Heller. Discovering findings that replicate from a primary study of high dimension to a follow-up study. *Journal of the American Statistical Association*, 108(504): 1480–1492, 2013.

M. Bogomolov and R. Heller. Replicability across multiple studies. *arXiv preprint arXiv:2210.00522*, 2022.

D. Chung, C. Yang, C. Li, J. Gelernter, and H. Zhao. Gpa: a statistical approach to prioritizing gwas results by integrating pleiotropy and annotation. *PLoS genetics*, 10(11):e1004787, 2014.

D. Edsgärd, P. Johnsson, and R. Sandberg. Identification of spatial expression trends in single-cell gene expression data. *Nature methods*, 15(5):339–342, 2018.

E. Hanson, J. Swanson, and B. R. Arenkiel. Gabaergic input from the basal forebrain promotes the survival of adult-born neurons in the mouse olfactory bulb. *Frontiers in Neural Circuits*, 14:17, 2020.

J. Hu, X. Li, K. Coleman, A. Schroeder, N. Ma, D. J. Irwin, E. B. Lee, R. T. Shinohara, and M. Li. Spagcn: Integrating gene expression, spatial location and histology to identify spatial domains and spatially variable genes by graph convolutional network. *Nature methods*, 18 (11):1342–1351, 2021.

K. Hung and W. Fithian. Statistical methods for replicability assessment. *The Annals of Applied Statistics*, 14(3):1063–1087, 2020.

M. Katidou, X. Grosmaitre, J. Lin, and P. Mombaerts. G-protein coupled receptors mc4r and drd1a can serve as surrogate odorant receptors in mouse olfactory sensory neurons. *Molecular and Cellular Neuroscience*, 88:138–147, 2018.

I. Kleino, P. Frolovaitė, T. Suomi, and L. L. Elo. Computational solutions for spatial transcriptomics. *Computational and Structural Biotechnology Journal*, 2022.

V. Kozareva, C. Martin, T. Osorno, S. Rudolph, C. Guo, C. Vanderburg, N. Nadaf, A. Regev, W. G. Regehr, and E. Macosko. A transcriptomic atlas of mouse cerebellar cortex comprehensively defines cell types. *Nature*, 598(7879):214–219, 2021.

E. Lubeck and L. Cai. Single-cell systems biology by super-resolution imaging and combinatorial labeling. *Nature methods*, 9(7):743–748, 2012.

P. A. Moran. Notes on continuous stochastic phenomena. *Biometrika*, 37(1/2):17–23, 1950.

A. Raj, P. Van Den Bogaard, S. A. Rifkin, A. Van Oudenaarden, and S. Tyagi. Imaging individual mrna molecules using multiple singly labeled probes. *Nature methods*, 5(10): 877–879, 2008.

S. G. Rodriques, R. R. Stickels, A. Goeva, C. A. Martin, E. Murray, C. R. Vanderburg, J. Welch, L. M. Chen, F. Chen, and E. Z. Macosko. Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution. *Science*, 363(6434):1463–1467, 2019.

A. D. Rouillard, G. W. Gundersen, N. F. Fernandez, Z. Wang, C. D. Monteiro, M. G. Mc-Dermott, and A. Ma'ayan. The harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes and proteins. *Database*, 2016, 2016.

P. L. Ståhl, F. Salmén, S. Vickovic, A. Lundmark, J. F. Navarro, J. Magnusson, S. Giacomello, M. Asp, J. O. Westholm, M. Huss, et al. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science*, 353(6294):78–82, 2016.

R. R. Stickels, E. Murray, P. Kumar, J. Li, J. L. Marshall, D. J. Di Bella, P. Arlotta, E. Z. Macosko, and F. Chen. Highly sensitive spatial transcriptomics at near-cellular resolution with slide-seqv2. *Nature biotechnology*, 39(3):313–319, 2021.

J. D. Storey. A direct approach to false discovery rates. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 64(3):479–498, 2002.

J. D. Storey and R. Tibshirani. Statistical significance for genomewide studies. *Proceedings of the National Academy of Sciences*, 100(16):9440–9445, 2003.

J. D. Storey, J. E. Taylor, and D. Siegmund. Strong control, conservative point estimation and simultaneous conservative consistency of false discovery rates: a unified approach. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 66(1):187–205, 2004.

S. Sun, J. Zhu, and X. Zhou. Statistical analysis of spatial expression patterns for spatially resolved transcriptomic studies. *Nature methods*, 17(2):193–200, 2020.

S. M. Sunkin, L. Ng, C. Lau, T. Dolbeare, T. L. Gilbert, C. L. Thompson, M. Hawrylycz, and C. Dang. Allen brain atlas: an integrated spatio-temporal portal for exploring the central nervous system. *Nucleic acids research*, 41(D1):D996–D1008, 2012.

V. Svensson, S. A. Teichmann, and O. Stegle. Spatialde: identification of spatially variable genes. *Nature methods*, 15(5):343–346, 2018.

S. Vickovic, G. Eraslan, F. Salmén, J. Klughammer, L. Stenbeck, D. Schapiro, T. Äijö, R. Bonneau, L. Bergenstråhle, J. F. Navarro, et al. High-definition spatial transcriptomics for in situ tissue profiling. *Nature methods*, 16(10):987–990, 2019.

C. Wu, I. MacLeod, and A. I. Su. Biogps and mygene. info: organizing online, gene-centric information. *Nucleic acids research*, 41(D1):D561–D565, 2013.

J. Zhu, S. Sun, and X. Zhou. Spark-x: non-parametric modeling enables scalable and robust detection of spatial expression patterns for large spatial transcriptomic studies. *Genome biology*, 22(1):1–25, 2021.