Model-based frontal cortex analysis reveals origins of human non-verbal communication

Leonardo Ceravolo[1,2*†], Coralie Debracque[1,2†], Eva Pool[2,3], Thibaud Gruber[1,2,4‡] & Didier Grandjean[1,2‡]

[1] Neuroscience of Emotions and Affective Dynamics lab, Department of Psychology and Educational
Sciences, University of Geneva, Geneva, Switzerland

[2] Swiss Center for Affective Sciences, University of Geneva, Geneva, Switzerland

[3] E3 Lab, Department of Psychology and Educational Sciences, University of Geneva, Geneva,
Switzerland

[4] eccePAN lab, Department of Psychology and Educational Sciences, University of Geneva, Geneva,
Switzerland

† shared first authors

‡ shared last authors

Corresponding author:

Leonarado Ceravolo: Leonardo.Ceravolo@unige.ch

ORCID: https://orcid.org/0000-0003-0638-3981

Abstract

The ability to process verbal language seems unique to humans and relies not only on semantics but on

other forms of communication such as affective vocalisations, that we share with other primate

species—particularly great apes (*Hominidae*). To better understand these processes at the behavioural

and brain level, we asked human participants to categorize vocalizations of four primate species

including human, great apes (chimpanzee and bonobo), and monkey (rhesus macaque) during MRI

acquisition. Classification was above chance level for all species but bonobo vocalizations. Imaging

analyses were computed using a participant-specific, trial-by-trial fitted probability categorization value

in a model-based style of data analysis. Model-based analyses revealed the implication of the bilateral

orbitofrontal cortex and inferior frontal gyrus *pars triangularis* (IFG_{tri}) respectively correlating and anti-

correlating with the fitted probability of accurate species classification. Further conjunction analyses

revealed enhanced activity in a sub-area of the left IFG_{tri} specifically for the accurate classification of

chimpanzee calls compared to human voices. Our data therefore reveal distinct frontal mechanisms that

shed light on how the human brain evolved to process non-verbal language.

Author contributions

CD and LC were involved in every steps of the study including experimental design, programming, data

acquisition, data analysis and redaction of the first draft of the manuscript and subsequent editing. EP

helped design the model-based MRI analyses and interpret the results. TG and DG were involved in the

experimental design and study planification. All authors wrote and edited the manuscript.

Introduction

Processing and understanding non-verbal language are fundamental aspects of everyday life in human (1) and animal (2, 3) communication. Their importance can sometimes even be greater than that of verbal communication since a short, powerful in-context vocal burst, for instance, an urgency signal, can easily be noticed and understood by other individuals (2). A critical part of non-verbal language can therefore be represented by 'affective' communication, especially through affective vocalizations, which are of particular interest for cross-species research. Indeed, the ability to express and understand such vocalizations is shared by several primate species, particularly Hominids, a taxonomic family that includes humans and other great apes (4, 5). Such commonalities therefore open a critical window into the investigation of human brain structures recruited by non-verbal communication, for instance by studying these crucial abilities to perceive, process and subsequently categorize and classify affective vocalizations expressed by primate species. These abilities are known to recruit superior temporal, orbitofrontal and inferior frontal cortices in humans (6-9) while these brain areas evolved a lot in primates (10, 11). We will now detail the particular importance of studying cross-species non-verbal language processing and categorization taking place in the frontal cortex of humans.

The *Hominidae* clade appeared between 13 and 18 million years ago (12). Encompassing all living great apes including humans (*Homo sapiens*), chimpanzees (*Pan troglodytes*), bonobos (*Pan Paniscus*), gorillas (*Gorilla subs*), and orangutans (*Pongo subs*), as well as our extinct ancestors, this unique primate taxon is key to understanding the evolution of human behaviour, physiology, communication and cognitive abilities. Very few studies have used bonobo vocalizations as stimuli (13), despite sharing the same phylogenetic proximity to *Homo sapiens* with chimpanzees, their behaviour repertoire—of which gestures can be decoded by humans (14)—including their vocal communication are noticeably different (15-18). Despite a larger and more folded brain compared to other great ape species, human neuroanatomical traits are mainly considered to belong to a continuum in the primate brain evolution (11, 19). For instance, findings in anatomical magnetic resonance imaging (MRI) have demonstrated the existence of a large frontal cortex in all great ape species—humans included(11)—emphasizing the particular interest of a comparative approach to investigate anatomical structures and related functions

of the frontal regions. Moreover, the functions of the frontal lobe, often associated to problem solving,

emotion processing and especially evaluative judgment, communication and language or even motor

and sexual behaviours (6, 20-25), as well as their related brain structures are shared by most primate

species. This includes two critical brain structures that are the focus of this article: the orbitofrontal

cortex (OFC) and the inferior frontal gyrus (IFG).

Existing in phylogenetically distant non-human primate species such as rhesus macaques (*Macaca*

mulatta) (26), the OFC has similar neuroanatomical architecture and functions across the primate family

tree. From humans to monkeys, studies have highlighted the roles of OFC in emotional memory (20,

21), emotional expression (9, 27), executive functions (20) and integration of contextual information in

emotional evaluation (8). Interestingly, despite these shared evolutionary OFC functions in primate

species, its role in heterospecific affective recognition, such as affective vocalizations, has rarely been

investigated. In fact, recent functional MRI data have pointed out a greater enhancement of activity in

OFC, in particular of its medial part, in the human identification of emotional voices compared to

affective animal vocalizations including positive and negative calls expressed by chimpanzees,

macaques and cats (*Felis catus*) (28, 29). More investigations on the role of the human OFC in this

context are thus crucially needed.

Functionally and anatomically connected to the OFC (30), the IFG is also present in great apes as well

as in monkey species (31). Yet, unlike the OFC, the left IFG is most prominently linked to speech

production (Broca's area) in humans (32), the only species 'capable' of a complex and sophisticated

language. Hence, in addition to likely shared functions such as decision formation (6, 33), inhibition

(34, 35) and emotional judgement (8, 36-38) shaped by natural selection, the IFG also became

specialized in semantic auditory processing in the human lineage in the course of evolution (7, 39, 40).

Because the current roles of human IFG are linked to both the *Hominidae* and the specific *Homo sapiens*

evolutionary histories (41), this frontal region, together with the OFC, is of high interest to investigate

conspecific and heterospecific identification mechanisms. Noteworthy is the fact that these specific

frontal regions have strong anatomical ties with temporal lobe regions that are crucial in processing

vocalizations, especially when they are biologically relevant (23, 30, 36, 42). Besides the importance of

OFC and IFG regions in the evolution of primate brains, comparative and integrative approaches to

investigate the current functions of these specific frontal regions are still missing. This is especially true

for auditory affective processing (43), species categorization and by extension non-verbal

communication comprehension in humans—excluding sign language.

The present study focused on the processing and subsequent categorization of primate vocalizations in

human OFC and IFG using fMRI. To do so, we asked participants to categorize four primate species

(human, chimpanzee, bonobo, and macaque) through their affective vocalizations. Non-verbal sounds—

namely affective vocal bursts—were selected as human vocalizations to match as well as possible the

nature of our primates' vocalizations stimuli. Research on vocal production and perception has indeed

demonstrated that non-verbal vocalizations expressed by humans are similar to those expressed by other

mammalian species, non-human primates included, especially in affiliative (e.g., play) or agonistic

contexts (e.g., aggression) (44). Interestingly, as most animal vocalizations are expressed in a

motivational or affective context, existing literature has remained focused on the human recognition of

affects in heterospecific vocalizations while neglecting humans' ability to identify other species

independently of affect (13, 29, 45-49). Hence, in light of previous literature as well as taking into

account the commonalities and differences of the selected four primate species in their evolutionary

path, we predicted: i) a behavioural gradient reflecting phylogenetic distance from humans for the

species with a higher probability of correctly categorizing human, then chimpanzee, bonobo and finally

macaque affective vocalizations (13, 46, 48); for similar reasons, we predicted that chimpanzee and

bonobo vocalizations should be more confused with each other than with macaque calls; ii) neural

correlates of the probability of correctly categorizing the four primate species in the OFC and IFG,

especially in the medial OFC (28, 29) and $IFG_{tri/op(50)}$, respectively; iii) a gradient of activity in the *pars*

triangularis and *opercularis* of the IFG when comparing the processing and correct categorization of

each non-human primate species' vocalizations to human voices using conjunction analyses, both

subparts being crucially involved in this process (37, 50).

Results

The present study aimed at assessing the ability of humans to accurately classify human and more

importantly other primate calls, in the present case chimpanzee, bonobo, and macaque vocalizations

(Fig.1**a**) and how frontal brains areas may underlie such successful non-verbal stimuli categorization.

Behavioural analyses were performed using a Species factor (human, chimpanzee, bonobo, macaque)

interacting with the Affective context factor for each vocalization (affiliative, agonistic 'threat',

agonistic 'distress') in a generalized linear mixed-effects modelling of the accuracy data ('1', correct

response; '0', incorrect response) and a linear mixed-effects modelling of the confusion data. This

statistical procedure was used to better assess the probability of correctly categorizing each species'

vocalizations as well as the pattern of confusion between each species, respectively. Following similar

reasoning, we used the trial-level probability of correct species categorization in model-based fMRI

methodology. We tested the specific frontal mechanism underlying the combined processing and

accurate classification of each non-human primate species vocalizations compared to human voices

using conjunction analyses. Both analyses were conducted in our regions of interest (ROI), namely in

the bilateral IFG (*pars opercularis*, *triangularis*, *orbitalis*) and OFC (superior, middle, medial) using

masked second-level analyses with voxelwise corrected statistics ($p<.05$ False Discovery Rate

correction 'FDR', k>5 voxels). Selected low-level acoustics (mean and standard deviation of stimulus

energy and fundamental frequency) were used as covariates of no interest for behavioural and

neuroimaging analyses to better control our data for low-level vocal fluctuations. Statistical modelling

power is reported for fixed effects ('R^2c') and for both fixed and random effects taken together ('R^2m').

Behavioural results

Accuracy data

As predicted, the generalized linear mixed-effects on accuracy revealed a significant effect of Species

factor ($\chi^2(3)=222.78$, $p<.0001$, $R^2c=56.38\%$ and $R^2m=61.10\%$). All complementary results of this model

can be found in Supplementary Table 1. We then tested the contrasts of interest according to our

behavioural hypothesis namely the impact of phylogenetic proximity on accurate species categorization:

our results highlighted a higher probability of correctly categorizing human compared to the

vocalizations of each non-human primate ($\chi^2(1)$=128.96, p_{Bonf} <.0001; human > chimpanzee:

$\chi^2(1)$=95.89, p_{Bonf}<.0001; human > bonobo: $\chi^2(1)$=199.51, p_{Bonf}<.0001; human > macaque: $\chi^2(1)$=79.40,

p_{Bonf}<.0001). We additionally compared each non-human primate species with each other and found a

higher probability of correct categorization for chimpanzee compared to bonobo ($\chi^2(1)$=52.64, p_{Bonf}

<.0001), and macaque compared to bonobo vocalizations ($\chi^2(1)$=18.99, p_{Bonf}<.0001). No difference was

observed when comparing the probability of correct species categorization between chimpanzee and

macaque vocalizations ($\chi^2(1)$=1.84, p_{Bonf}=.17, Fig. 1b).

Confusion data analyses were computed by considering species miscategorization, namely one species'

vocalization being categorized as belonging to another species (Fig.1**c**). The confusion for human voices

was not significant ($\chi^2(3)$=1.65, p=.65), meaning that these were not significantly miscategorised with

any other species. Confusion concerning chimpanzee vocalizations showed a significant main effect of

Species ($\chi^2(3)$=265.79, p<.0001; R^2c=16.41% and R^2m=24.64%), showing that these were mostly

miscategorised as bonobo vocalizations ($\chi^2(1)$=149.52, p_{Bonf}<.0001), and then as macaque ($\chi^2(1)$=47.11,

p_{Bonf}<.0001) vocalizations (for chimpanzee confusions, bonobo > macaque: $\chi^2(1)$=15.34, p_{Bonf}<.0001),

but not as human voices ($\chi^2(1)$=0.21, p_{Bonf} >.99). Confusion for bonobo vocalizations showed a

significant main effect of the Species factor ($\chi^2(3)$=107.96, p<.0001; R^2c=24.50% and R^2m=28.95%)

and specific confusion was the highest with macaque ($\chi^2(1)$=114.70, p_{Bonf}<.0001) and then chimpanzee

($\chi^2(1)$=111.28, p_{Bonf}<.0001) vocalizations (for bonobo confusions, macaque > chimpanzee: $\chi^2(1)$=5.33,

p_{Bonf} <.05) but no significant confusion with human voices was observed ($\chi^2(1)$=0.06, p_{Bonf} >.99).

Confusion for macaque vocalizations showed a main effect of the Species factor ($\chi^2(3)$=469.68, p<.0001;

R^2c=29.01 % and R^2m=30.65%) with highest confusion with bonobo ($\chi^2(1)$=565.04, p_{Bonf} <.0001)

followed by chimpanzee ($\chi^2(1)$=145.96, p_{Bonf}<.0001) vocalizations (for macaque confusions, bonobo >

chimpanzee: $\chi^2(1)=87.34$, $p_{Bonf} <.0001$). No confusion with the human voice reached significance

($\chi^2(1)=0.04$, $p_{Bonf} >.99$). All other specific confusion effects can be found in Supplementary Table 2.

Figure 1. Experimental procedure and behavioural results of the probability of correct species

categorization and confusions. a, Example of four consecutive experimental task trials (inter-trial

interval of 3-5s) for which the participants had to categorize the species' stimulus (750ms duration) for

each trial (pseudo-randomized species order) and spectrogram example for each species. **b,** Probability

of the correct categorization of each species (*y* axis) modelled across all participants, trials and species

(i.e., human, chimpanzee, bonobo, macaque); Each circle represents a participant's averaged value per

species, full circles represent the general average per species across participants ± the standard error of

the mean, chance level is 25% (grey dotted line). **c,** Percentage of categorization confusion (*y* axis)

specific to each species (*x* axis); Each circle represents a participant's averaged value, full circles

represent the general average per species across participants ± the standard error of the mean. Hum:

human; Chimp: chimpanzee; Bon: bonobo; Mac: macaque. ***$p<.001$, *$p<.05$, n.s.: non-significant.

Neuroimaging results using IFG and OFC as regions of interest

We first used a model-based approach in order to characterize the general frontal mechanism underlying

the accurate categorization of primate vocalizations by human participants (see model 1 in the methods

section). To do so, we modelled all trials with the sample-specific fitted probability of a correct vs.

incorrect species categorization as a covariate of interest for each trial's onset. This covariate was then

taken as the main regressor of interest to masked group-level analyses to reveal frontal patterns of

activity related to the general human neural mechanism of classifying human, chimpanzee, bonobo, and

macaque vocalizations. These fitted probability values were obtained across participants and species by

predicting accuracy data at the trial level, therefore including all trials, correct and incorrect, in a logistic

regression analysis. Fixed effects included the Species and Affective context factors and covariates of

no interest (mean and standard deviation of vocalization energy and fundamental frequency) with

random effects characterizing the identity of each participant (random intercept) and each stimulus

(random slope).

Activity correlating with the probability of correctly categorizing each species' vocalizations was

revealed in the right IFG$_{tri}$ (Fig.2**b**) and bilateral IFG$_{orb}$ (Fig.2**abd**) as well as in the bilateral OFC$_{med}$

(Fig.2**c**). Anti-correlated activity was revealed in a large left IFG$_{tri}$, IFG$_{op}$ cluster (Fig.2**ef**) and to a lesser

extent in the left IFG$_{orb}$ (Fig.2**e**), right IFG$_{op}$ and IFG$_{orb}$ (Fig.2**fh**) and in the right OFC$_{mid}$ (Fig.2**b**) and

left OFC$_{sup}$ (Fig.2**e**). See Supplementary Table 3 for cluster coordinates.

General neural mechanism for categorizing Human,Chimpanzee,Bonobo & Macaque species
Controlled for: vocalization context(agonistic,affiliative), energy,f0(mean, std)
p<.05 FDR, k>5, ROI: IFG, OFC

Figure 2. Model-based neural correlates and anti-correlates of the probability of correctly categorizing each species' vocalizations in the IFG and OFC. abcd, Trial-by-trial correlates of the probability of correct species categorization. **efgh,** Trial-by-trial anti-correlates of the probability of correct species categorization. Data of this analysis were computed by using a modelling of species categorization accuracy across all participants, species (Species factor) and trials including the Affective context factor and trial-level covariates of no interest (mean and standard deviation of the vocalization energy and fundamental frequency). Subsequently, the fitted probability values of this model were used as a parametric modulator of interest for each trial's onset, further taken as main regressor in second-

level analyses. Colorbars illustrate t-value statistics. All activations thresholded at a voxelwise $p<.05$

FDR, k>5 voxels, masked by IFG and OFC regions of interest (k=9635 voxels in total). IFG: inferior

frontal gyrus; tri: pars triangularis; op: pars opercularis; orb: pars orbitalis; OFC: orbitofrontal cortex;

med: medial; mid: middle; sup: superior.

We then computed two conjunction analyses with each non-human primate species compared to human

voice categorization. In these analyses, enhanced activations for each non-human primate species was

tested while driving the conjunction but only the [(chimpanzee > human) > (bonobo > human) >

(macaque > human) vocalizations] contrast showed significant results (see model 2 in the methods

section). This conjunction revealed left-lateralized IFG$_{tri}$ and IFG$_{op}$ as well as right IFG$_{tri}$ activity for the

processing of the vocalizations (i.e., general processing displayed as white-to-pink activations,

Fig.3**abd**) while enhanced activity for correctly categorized species (see model 3 in the methods section)

was only found in left IFG$_{tri}$ (i.e., correct categorization displayed as black-to-white activations, Fig.3**a**).

No enhanced signal was observed in the OFC. Other combinations of non-human primate vs. human

species conjunction contrasts did not yield to any above-threshold voxels. Mean and standard deviation

of each vocalization's energy and fundamental frequency were used as trial-level covariates of no

interest in these statistical analyses as well. See Supplementary Table 4 for cluster coordinates.

Figure 3. Conjunction results of the processing and correct categorization of non-human primates
compared to human vocalizations in the IFG and OFC. abcd, Conjunction analysis for the
processing (white-to-pink activations, all trials of each species) and correct species categorization
(black-to-white activations with pink outline, correct trials only) of the contrasts [(chimpanzee > human)
> (bonobo > human) > (macaque > human) vocalizations]. No above-threshold voxels were found for
[bonobo > human] or [macaque > human] contrasts driving the conjunction analyses. For both models
(species processing; correct species categorization), trial-level covariates of no interest included the
mean and standard deviation of the vocalization energy and fundamental frequency. Colorbars illustrate
t-value statistics. All activations thresholded at a voxelwise $p<.05$ FDR, k>5 voxels, masked by IFG and
OFC regions of interest (k=9635 voxels in total). IFG: inferior frontal gyrus; tri: pars triangularis; op:
pars opercularis; orb: pars orbitalis; OFC: orbitofrontal cortex; med: medial; mid: middle; sup: superior.

The comparisons between the processing (all trials) or the correct categorization (correct trials only) of
the vocalizations of each non-human species compared to those of humans, and of human voices
compared to the vocalizations of each non-human primate species are reported in Fig.S1 and Fig.S2,
respectively. Activations specific to the processing and correct categorization of each non-human
primate species or between each non-human primate species are reported in Fig.S3 and Fig.S4,
respectively.

Discussion

The present study was designed to clarify the importance of primate vocalizations processing and classification by humans as a proxy for the more general treatment of human non-verbal communication. In fact, the processing of primate vocalizations by humans might boost our understanding of non-verbal communication in today's *Homo sapiens* and inform our reconstruction of our last common ancestor with other Hominids. To operationalize such a research objective, we investigated the behavioural and frontal brain mechanisms in modern humans underlying the vocal recognition of non-human primates' vocalizations, considering their phylogenetic proximity to the human species. Our data highlighted the evolutionary shared history of our species with the communication system of other primates, particularly other great apes, with both the recognition and classification processes of hominid vocalizations involving the medial orbitofrontal and bilateral inferior frontal cortex, especially the *pars triangularis*. Behavioural data emphasized the astonishing ability of human participants to categorize chimpanzee and macaque calls above chance level while it was not the case for bonobo vocalizations. Despite the close phylogenetic proximity of this species with *Homo sapiens*, the peculiar acoustical structure of bonobo calls may have prevented modern humans from recognizing them, yielding to miscategorization. Indeed, confusion matrices showed that, despite their close phylogenetic relationship to chimpanzees, bonobo vocalizations were in fact often confused for those of macaques. This peculiarity in the processing of bonobo vocalizations has also been reported in previous literature showing that human adults are highly sensitive to pitch information for discriminating the arousal of primate babies' cries (13). In fact, human participants rated bonobo infant calls as more distressful in comparison to those of humans or chimpanzees and after further acoustical analyses, the authors found that bonobo infants' calls had higher pitched vocalizations compared to those of human and chimpanzee babies, both species sharing more similar acoustical features (13). Such acoustical divergence of bonobos among non-human primates and especially the increased fundamental frequency of their calls may well have its origin in the shorter length of the larynx of this species (15). Interestingly in our study, chimpanzee calls were not better categorized by human participants than those of a more evolutionary distant species such as

rhesus macaques. This was especially surprising considering that previous studies have mostly found a

better recognition rate for chimpanzee vocalizations compared to the calls of other non-human primates

(46, 51). Hence, our behavioral analysis suggests that the capacity of modern humans to accurately

identify non-human primate calls could rely more heavily on acoustic similarity rather than phylogenetic

proximity alone (52).

To assess the frontal brain mechanisms involved in the general processing of human voice and non-

human primate vocalization recognition, we turned to computational modelling and more precisely to

model-based fMRI analyses. These data pointed to a frontal network including the bilateral medial OFC

for the correct recognition and categorization of human voices and non-human primate vocalizations.

This first result suggests that the medial OFC has a central role in the accurate categorization of primate

calls, extending previous fMRI results that showed an enhancement of activity in OFC regions—

especially of the medial part—for the discrimination of emotions in human, chimpanzee, macaque, and

cat vocalizations (28, 29). Other frontal brain areas underlying the probability of accurate species

categorization included small clusters of the bilateral IFG_{orb} as well as the right IFG_{op} and IFG_{tri}. Frontal

brain areas underlying miscategorization and/or what could be labelled as 'hesitation' were found in

large portions of the bilateral IFG, especially in the IFG_{tri} which is coherent with existing literature (53).

This result is also supported by our conjunction analyses, which allowed a more concise and focused

multi-contrast approach to pinpoint the specific frontal brain mechanisms that may underlie the

processing as well as accurate recognition and categorization of non-human primate vocalizations

compared to the human voice. These analyses revealed large, similar bilateral IFG_{tri} clusters of activity

when creating a hierarchy for processing chimpanzee, bonobo and macaque vocalizations each

compared to the human voice—namely, contrasting (chimpanzee > human) > (bonobo > human) >

(macaque > human) at the same time. Interestingly, this conjunction contrast also revealed a difference

in the hierarchy between the accurate recognition of chimpanzee vocalizations and those of bonobos and

macaques when each was compared to the human voice in a subpart of the left IFG_{tri}. In the literature,

the IFG_{tri} was repeatedly involved when human participants have to make a decision based on vocal

signals, for instance when human participants discriminate or categorize human vocal emotion (37, 50)

or when they implicitly or explicitly process human emotional voices (7). Causal links between IFG$_{tri}$

functioning and emotion recognition were also revealed in the literature (54). Noteworthy is the fact that

activity in the left IFG$_{tri}$ also translates a level of uncertainty when decisional processes occur (53, 55-

57). Combining our behavioural and neuroimaging results, it seems extremely plausible that decisional

uncertainty was peaking when human participants were categorizing non-human primate vocalizations.

We therefore think that according to our results and to the existing literature, activity in the IFG$_{tri}$ might

underlie both categorization uncertainty as revealed by anti-correlates of the probability of accurate

species recognition in the model-based neuroimaging results, as well as decisional processes that lead

to an accurate species recognition in a very concise region of the left IFG$_{tri}$, as highlighted by our

conjunction results. Therefore, OFC and IFG brain areas—especially IFGtri activity—may be the

necessary biological and functional 'hardware' allowing *Homo sapiens* to classify and integrate non-

verbal communication, be it human voice or other primate vocalizations. Such ability may then have

been co-opted for the treatment of the proto-language used by our human ancestors, although precisely

how close this proto-language was from to today's non-human primates vocalizations remains an open

question , with a question mark regarding a potential interaction with between-species acoustical

differences (52). We contend that the common activity we evidenced in the OFC and IFG across stimuli

suggests that such proto-language may have shared acoustic properties with non-human primate

vocalizations.

Even though our results shed new light on the origins of human non-verbal communication, our study

suffered from several limitations. Such limitations include the absence of higher temporal resolution,

the absence of other great apes as stimuli (for instance gorillas and/or orangutans) and the absence of

highly specific acoustic distance assessment. Also, the time-course of such frontal mechanisms

underlying non-verbal communication in humans should be investigated.

To conclude, the present study revealed the existence of distinct mechanisms at play in the human frontal

cortex for the integration and further categorization of non-human primate vocalizations. In fact,

computational modelling of the fMRI data allowed a clear distinction between accurate categorization

on the one hand and miscategorization processes on the other hand, as revealed by bilateral medial OFC

and left IFG$_{tri}$, respectively. Combined with our behavioural data, evidence converges toward a

preserved ability of *Homo sapiens* to recognize the calls of species close to his ancestors that were most likely very similar to those of today's non-human primate vocalizations. Natural selection might also have led to specific anatomical adaptations in some non-human primates across time, and the shorter larynx of bonobos leading to the significantly increased fundamental frequency of their calls is a well-fitting example, especially in the context of the present study. Therefore, a combination of phylogenetic and acoustical proximity may be necessary for humans to recognize and successfully categorize the calls of non-human primates, including those of great apes. Overall, the present study substantially contributes to the understanding of primate calls' recognition by modern humans as a proxy for more general non-verbal communication mechanisms in *Homo sapiens*.

Material and Methods

Experimental procedure and paradigm

Participants laid comfortably in a 3T scanner and heard a total of seventy–two stimuli in pseudo-randomized order through MRI-compatible earphones (Sensimetrics Corporation, Gloucester, MA, USA) at a sound pressure level of 70dB. Participants were instructed to identify the species that expressed the vocalizations using a response box (Current Designs, Inc., Philadelphia, PA, USA). For instance, the instructions could be "Human – press 1, Chimpanzee – press 2, Bonobo – press 3 or Macaque – press 4". There were four different mappings of the keys that were randomly assigned across participants. In a 3 – 5 second interval (jittering of 400 ms) after each stimulus, participants were asked to categorize the species. If the participant did not respond during this interval—namely a missed trial, the next stimulus followed automatically and the value of the response was '0' while '1' coded a correct response, '2' an incorrect response (Fig.1**a**).

Stimuli

Seventy-two vocalizations of four primate species (human, chimpanzee, bonobo and rhesus macaque) were used in this study. The eighteen human voices were expressed by two male and two female actors, obtained from a non-verbal (affective vocal burst) validated set of Belin and collaborators (58). Non-

linguistic vocalizations were selected as human voice stimuli due to their acoustic and perceptual

proximity with the other mammalian calls, non-human primates included (44). The eighteen selected

chimpanzee, bonobo and rhesus macaque vocalizations contained single calls or two call sequences

produced by 6 to 8 different individuals in their natural environment. For each non-human primate

species, the stimuli were either produced in an affiliative context or produced in an agonistic context

(threating or distressful). All vocal stimuli were standardized to 750 milliseconds using PRAAT

(www.praat.org) but were not normalized in order to preserve the 'naturality' of the sounds (59).

Participants

Twenty-five right-handed, healthy, either native or highly proficient French-speaking participants took

part in the study. One participant was excluded because he had no correct response at all and may have

fallen asleep, while another participant was excluded due to incomplete scanning and technical issues,

leaving us with twenty-three participants (10 female, 13 male, mean age 24.65 years, SD 3.66). All

participants were naive to the experimental design and study, had normal or corrected-to-normal vision,

normal hearing and no history of psychiatric or neurologic incidents. Participants gave written informed

consent for their participation in accordance with ethical and data security guidelines of the University

of Geneva. The study was approved by the Ethics Cantonal Commission for Research of the Canton of

Geneva, Switzerland (CCER) and was conducted according to the Declaration of Helsinki.

Statistical analysis

Behavioural data were analysed using R studio software (60). In order to test our hypotheses regarding

the impact of phylogenetic distance on species categorization, we performed mixed-effects models

analyses of the 'lme4' package (61) on accuracy and confusion (one accuracy column: '1', correct

response; '0', incorrect response; four confusion columns: '1', confusion with the species, '0' no

confusion with the species, for each of the four species column) and reaction times data (for correct,

incorrect trials and specific confusion for each species) with Species (human, chimpanzee, bonobo,

macaque) and Affective context (affiliative and agonistic) as interacting fixed factors, four covariates

(mean energy of each species, mean fundamental frequency of each species, energy standard deviation

of each species, fundamental frequency standard deviation of each species) and the participants' and the

stimuli identity as random factors (random intercept and random slope, respectively). The inclusion of

the sex and age of the participants for which we had no specific hypothesis in data modelling did not

yield to a significantly better model neither for accuracy ($\chi^2(2)$=.81, p=.66) nor for reaction times data

($\chi^2(2)$=.01, p=1) and these random effects were therefore discarded from the analyses. The same model

was used to analyse accuracy, confusion and reaction times data, except that a logistic regression

(generalized linear mixed-effects models, 'glmer') was used for accuracy and a linear regression (linear

mixed-effects models, 'lmer') for specific pairs of species confusion (four models in total) and for

reaction times. Then, we used specific contrasts computed using the 'phia' package (62) to compare

between-species categorization, testing our hypothesis of an impact of phylogenetic proximity on

species categorization. Contrasts therefore included human vs. non-human primate planned

comparisons: [human > chimpanzee, bonobo, macaque vocalizations], [human > chimpanzee

vocalizations], [human > bonobo vocalizations], [human > macaque vocalizations]; non-human primate

vs. human comparisons: [chimpanzee > human vocalizations], [bonobo > human vocalizations],

[macaque > human vocalizations]. These inverse contrasts actually yield to a sign-reversal of the

difference estimate of the original contrast for accuracy data. For confusion data, contrasts were

computed to uncover any difference between one human or non-human primate species compared to the

remaining species: [Confusion for human vocalizations: chimpanzee response ≠ bonobo response ≠

macaque response], [Confusion for chimpanzee vocalizations: human response ≠ bonobo response ≠

macaque response], [Confusion for bonobo vocalizations: human response ≠ chimpanzee response ≠

macaque response], [Confusion for macaque vocalizations: human response ≠ chimpanzee response ≠

bonobo response]. All contrasts were corrected for multiple comparisons using a Bonferroni adjustment

method implemented in the 'phia' package (adjustment="Bonf") and p-value is accordingly reported as

'p_{Bonf}'. Reaction times data were analysed using the same contrasts and are reported in Supplementary

Material.

Image acquisition

Structural and functional brain imaging data were acquired by using a 3T scanner Siemens Trio, Erlangen, Germany with a 32-channel coil. A 3D GR\IR magnetization-prepared rapid acquisition gradient echo sequence was used to acquire high-resolution ($0.35 \times 0.35 \times 0.7$ mm^3) T1-weighted structural images (TR = 2400 ms, TE = 2.29 ms). Functional images were acquired by using fast fMRI, with a multislice echo planar imaging sequence 79 transversal slices in descending order, slice thickness 3 mm, TR = 650 ms, TE = 30 ms, field of view = 205 x 205 mm2, 64 x 64 matrix, flip angle = 50 degrees, bandwidth 1562 Hz/Px.

Wholebrain image analysis

Functional images were analysed with Statistical Parametric Mapping software (SPM12, Wellcome Trust Centre for Neuroimaging, London, UK). Preprocessing steps included realignment to the first volume of the time series, slice timing, normalization into the Montreal Neurological Institute (MNI) space (63) using the DARTEL toolbox (64) and spatial smoothing with an isotropic Gaussian filter of 8 mm full width at half maximum. To remove low-frequency components, we used a high-pass filter with a cut-off frequency of 128s. Three general linear models were used to compute first-level statistics.

Model 1: model-based categorization probability of each species, including all trials

In this model, the probability of correctly categorizing each species in each production context was modelled using the 'fitted' function of the 'stats' package (65) to obtain the fitted probability values from the behavioural accuracy data described in the behavioural data analysis section. This modelling was performed by using each trial (N=72) of each fMRI session (N=1) of each participant (N=23) with a binary '1' and '0' coding for the correct and incorrect species categorization, respectively, as dependent variable. As mentioned above, the model included fixed effects, namely the interaction between the Species factor (human, chimpanzee, bonobo, macaque) and the Affective context factor (affiliative, agonistic) and four covariates of interest (in this order: mean energy of each species, mean fundamental frequency of each species, energy standard deviation of each species, fundamental frequency standard deviation of each species). Random effects included the participants' and the stimuli

identity (random intercept and random slope, respectively). The sample-specific matrix (participants per

trials=23*72=1656 lines) was then cut to the corresponding participant-specific matrix (72 lines). Model

1 therefore included two 72-lines regressors, namely the onsets of each trial in its order of appearance—

modelled by using a boxcar function and convolved with the hemodynamic response function time-

locked to the onset of each stimulus—and the corresponding modelled probability of correctly

categorizing each species as a covariate of interest. Six motion parameters were included as regressors

of no interest to account for movement in the data and our design matrix therefore included a total of 8

columns plus the constant term. The regressor of interest (modelled probability of correctly categorizing

each species) was used to compute a simple contrast for each participant, leading to separate main effect

of the probability of correctly categorizing each species. This simple contrast was then taken to a one-

sample t-test second-level analysis in which variance was set to 'unequal'.

Model 2: processing of all species, including all trials

In this model, each event was modelled by using a boxcar function and was convolved with the

hemodynamic response function, time-locked to the onset of each stimulus. Separate regressors were

created for all trials of each species (Species factor: human, chimpanzee, bonobo, macaque) and

production context (Affective context factor: affiliative, agonistic) and four covariates each (mean

energy of each species, mean fundamental frequency of each species, energy standard deviation of each

species, fundamental frequency standard deviation of each species) for a total of 47 regressors. The

means and standard deviations of the fundamental frequency and the energy of the vocalizations were

extracted using the extended Geneva Acoustic parameters set, which is defined as the optimal acoustic

indicators related to human voice analysis (GeMAPS (66)). This set of acoustical parameters was

selected based on i) their potential to index affective physiological changes in voice production, ii) their

proven value in former studies as well as their automatic extractability, and iii) their theoretical

significance.

Six motion parameters were included as regressors of no interest to account for movement in the data

and our design matrix therefore included a total of 66 columns plus the constant term. The species

regressors were used to compute simple contrasts for each participant, leading to separate main effects

of human, chimpanzee, bonobo and macaque vocalizations for each production context (12 simple contrasts). Covariates were set to zero in order to model them as no-interest regressors. These simple contrasts were then taken to a flexible factorial second-level analysis in which there were three factors: the Participants factor (data independence set to 'yes', variance set to 'unequal'), the Species factor and the Affective context factor (for both factors: data independence set to 'no', variance set to 'unequal'). For this first model, the main effect of the Species factor was computed in the flexible factorial modelling while the Affective context factor was modelled but excluded from the results because we did not have any hypothesis regarding this factor and because each context modality contained only 6 trials per species.

Model 3: correct categorization of each species, including only correct trials

Separate regressors were created for each correctly categorized species (Correct Species factor: human hits, chimpanzee hits, bonobo hits, macaque hits vocalizations) for each production context (Affective context factor: affiliative, agonistic) and four covariates each (mean energy of each species, mean fundamental frequency of each species, energy standard deviation of each species, fundamental frequency standard deviation of each species) as well as a regressor containing a concatenation of categorization errors across species for a total of 44 regressors. Regressors were modelled by using a boxcar function and convolved with the hemodynamic response function, time-locked to the onset of each stimulus. Six motion parameters were included as regressors of no interest to account for movement in the data and our design matrix therefore included a total of 51 columns plus the constant term. The regressors of interest (Correct Species factor) were used to compute 8 simple contrasts for each participant, leading to separate main effects of human hits, chimpanzee hits, bonobo hits and macaque hits vocalizations for each production context. Covariates were set to zero in order to model them as no-interest regressors. These simple contrasts were then taken to a flexible factorial second-level analysis in which there were three factors: the Participants factor (independence set to 'yes', variance set to 'unequal'), the Correct Species factor and the Affective context factor (for both factors: independence set to 'no', variance set to 'unequal'). For this third model, the main effect of the Correct Species factor was computed in the flexible factorial modelling while the Affective context factor was modelled but

excluded from the results because we did not have any hypothesis regarding this factor and because each

modality contained only 6 trials per species.

For all three models and to be consistent in our analyses, neuroimaging activations were masked by a

binary image including all subparts of the IFG (*pars opercularis*, *triangularis* and *orbitalis*) and OFC

(superior, middle and medial) for a total of twelve subregions, 9635 voxels and thresholded in SPM12

by using a voxel-wise false discovery rate (FDR) correction at $p<.05$ and an arbitrary cluster extent of

k>5 voxels to remove extremely small clusters of activity. Region masks were extracted using the

'automated anatomical labelling' ('aal') atlas (67) implemented in the 'WFU_Pickatlas' toolbox

(https://www.nitrc.org/projects/wfu_pickatlas/). Contrast activations were rendered on semi-inflated

brains from the CONN toolbox (68) with black outlines delineating each of our IFG-OFC regions of

interest.

Acknowledgements

We warmly thank Prof. Katie Slocombe and Dr. Zanna Clay for providing the non-human primates

vocalizations as well as Prof. Daphne Bavelier for her valuable advice on the experimental task. We

would like also to acknowledge the staff of the Brain and Behaviour Laboratory at The University of

Geneva where all data were acquired.

Competing interests

The authors declare no competing interests whatsoever.

References

1.   Tracy JL, Randles D, Steckler CM. The nonverbal communication of emotions. Current opinion

in behavioral sciences. 2015;3:25-30.

2.   Leger DW. Contextual sources of information and responses to animal communication signals.

Psychol Bull. 1993;113(2):295.

3.   Marler P. Animal Communication Signals: We are beginning to understand how the structure

of animal signals relates to the function they serve. Science. 1967;157(3790):769-74.

4.   Gagneux P, Varki A. Genetic differences between humans and great apes. Molecular

phylogenetics and evolution. 2001;18(1):2-13.

5.   Prado-Martinez J, Sudmant PH, Kidd JM, Li H, Kelley JL, Lorente-Galdos B, et al. Great ape

genetic diversity and population history. Nature. 2013;499(7459):471-5.

6.   Binder JR, Liebenthal E, Possing ET, Medler DA, Ward BD. Neural correlates of sensory and

decision processes in auditory object identification. Nature Neuroscience. 2004;7(3):295-301.

7.   Frühholz S, Ceravolo L, Grandjean D. Specific brain networks during explicit and implicit

decoding of emotional prosody. Cereb Cortex. 2012;22(5):1107-17.

8.   Grandjean D. Brain Networks of Emotional Prosody Processing. Emotion Review. 2020.

9.   Murray EA, Izquierdo A. Orbitofrontal Cortex and Amygdala Contributions to Affect and

Action in Primates. Annals of the New York Academy of Sciences. 2007;1121(1):273-96.

10.   Damasio AR, Anderson SW. The frontal lobes. 2003.

11.   Semendeferi K, Lu A, Schenker N, Damasio H. Humans and great apes share a large frontal

cortex. Nature Neuroscience. 2002;5(3):272-6.

12.   Perelman P, Johnson WE, Roos C, Seuánez HN, Horvath JE, Moreira MAM, et al. A Molecular

Phylogeny of Living Primates. PLOS Genetics. 2011;7(3):e1001342.

13.   Kelly T, Reby D, Levréro F, Keenan S, Gustafsson E, Koutseff A, et al. Adult human perception

of distress in the cries of bonobo, chimpanzee, and human infants. Biological Journal of the Linnean

Society. 2017;120(4):919-30.

14.   Graham KE, Hobaiter C. Towards a great ape dictionary: Inexperienced humans understand

common nonhuman ape gestures. PLOS Biology. 2023;21(1):e3001939.

15.   Grawunder S, Crockford C, Clay Z, Kalan AK, Stevens JMG, Stoessel A, et al. Higher

fundamental frequency in bonobos is explained by larynx morphology. Current biology: CB.

2018;28(20):R1188-R9.

16.   Gruber T, Clay Z. A Comparison Between Bonobos and Chimpanzees: A Review and Update.

Evolutionary Anthropology: Issues, News, and Reviews. 2016;25(5):239-52.

17.   Hare B, Wobber V, Wrangham R. The self-domestication hypothesis: evolution of bonobo

psychology is due to selection against aggression. Animal Behaviour. 2012;83(3):573-85.

18.   Staes N, Smaers JB, Kunkle AE, Hopkins WD, Bradley BJ, Sherwood CC. Evolutionary

divergence of neuroanatomical organization and related genes in chimpanzees and bonobos. Cortex.

2018.

19.   Herculano-Houzel S. The human brain in numbers: a linearly scaled-up primate brain. Frontiers

in Human Neuroscience. 2009;3.

20.   Barbas H. Connections underlying the synthesis of cognition, memory, and emotion in primate

prefrontal cortices. Brain Res Bull. 2000;52(5):319-30.

21.   Barbas H, Zikopoulos B, Timbie C. Sensory Pathways and Emotional Context for Action in

Primate Prefrontal Cortex. Biological Psychiatry. 2011;69(12):1133-9.

22.   Davidson RJ. Anterior cerebral asymmetry and the nature of emotion. Brain and Cognition.

1992;20(1):125-51.

23.   Frühholz S, Grandjean D. Processing of emotional vocalizations in bilateral inferior frontal

cortex. Neuroscience and Biobehavioral Reviews. 2013;37(10 Pt 2):2847-55.

24.   Kambara T, Brown EC, Silverstein BH, Nakai Y, Asano E. Neural dynamics of verbal working

memory in auditory description naming. Scientific Reports. 2018;8(1):15868.

25.   LeDoux J. Rethinking the emotional brain. Neuron. 2012;73(4):653-76.

26.   Rolls ET. Convergence of sensory systems in the orbitofrontal cortex in primates and brain

design for emotion. The Anatomical Record Part A: Discoveries in Molecular, Cellular, and

Evolutionary Biology. 2004;281A(1):1212-25.

27.   Sander D, Grandjean D, Scherer KR. A systems approach to appraisal mechanisms in emotion.

Neural Networks. 2005;18(4):317-52.

28.   Belin P, Fecteau S, Charest I, Nicastro N, Hauser MD, Armony JL. Human cerebral response to

animal affective vocalizations. Proc Biol Sci. 2008;275(1634):473-81.

29.   Fritz T, Mueller K, Guha A, Gouws A, Levita L, Andrews TJ, et al. Human behavioural

discrimination of human, chimpanzee and macaque affective vocalisations is reflected by the neural

response in the superior temporal sulcus. Neuropsychologia. 2018;111:145-50.

30.   Hsu C-C, Rolls ET, Huang C-C, Chong ST, Zac Lo C-Y, Feng J, et al. Connections of the

Human Orbitofrontal Cortex and Inferior Frontal Gyrus. Cerebral Cortex. 2020;30(11):5830-43.

31.   Petrides M, Pandya DN. Comparative cytoarchitectonic analysis of the human and the macaque

ventrolateral prefrontal cortex and corticocortical connection patterns in the monkey. Eur J Neurosci.

2002;16(2):291-310.

32.   Guenther FH, Tourville JA, Bohland JW. Speech Production. In: Toga AW, editor. Brain

Mapping. Waltham: Academic Press; 2015. p. 435-44.

33.   Brück C, Kreifelts B, Kaza E, Lotze M, Wildgruber D. Impact of personality on the cerebral

processing of emotional prosody. NeuroImage. 2011;58(1):259-68.

34.   Hampshire A, Chamberlain SR, Monti MM, Duncan J, Owen AM. The role of the right inferior

frontal gyrus: inhibition and attentional control. NeuroImage. 2010;50(3-3):1313-9.

35.   Tops M, Boksem MAS. A Potential Role of the Inferior Frontal Gyrus and Anterior Insula in

Cognitive Control, Brain Rhythms, and Event-Related Potentials. Frontiers in Psychology. 2011;2.

36.   Ethofer T, Anders S, Erb M, Herbert C, Wiethoff S, Kissler J, et al. Cerebral pathways in

processing of affective prosody: a dynamic causal modeling study. NeuroImage. 2006;30(2):580-7.

37.   Gruber T, Debracque C, Ceravolo L, Igloi K, Marin Bosch B, Frühholz S, et al. Human

Discrimination and Categorization of Emotions in Voices: A Functional Near-Infrared Spectroscopy

(fNIRS) Study. Frontiers in Neuroscience. 2020;14.

38.   Zhang D, Zhou Y, Yuan J. Speech Prosodies of Different Emotional Categories Activate

Different Brain Regions in Adult Cortex: an fNIRS Study. Scientific Reports. 2018;8(1):218.

39.   Belyk M, Brown S, Lim J, Kotz SA. Convergence of semantics and emotional expression within

the IFG pars orbitalis. NeuroImage. 2017;156:240-8.

40.   Schirmer A, Kotz SA. Beyond the right hemisphere: brain mechanisms mediating vocal

emotional processing. Trends in Cognitive Sciences. 2006;10(1):24-30.

41.   Stout D, Chaminade T. Stone tools, language and the brain in human evolution. Philosophical

Transactions of the Royal Society B: Biological Sciences. 2012;367(1585):75-87.

42.   Ethofer T, Bretscher J, Gschwind M, Kreifelts B, Wildgruber D, Vuilleumier P. Emotional voice

areas: anatomic location, functional properties, and structural connections revealed by combined

fMRI/DTI. Cereb Cortex. 2012;22(1):191-200.

43.   Gruber T, Grandjean DM. A comparative neurological approach to emotional expressions in

primate vocalizations. Neuroscience and Biobehavioral Reviews. 2017;73:182-90.

44.   Anikin A, Bååth R, Persson T. Human non-linguistic vocal repertoire: Call types and their

meaning. Journal of Nonverbal Behavior. 2018;42(1):53-80.

45.   Belin P. Voice processing in human and non-human primates. Philosophical Transactions of the

Royal Society B: Biological Sciences. 2006;361(1476):2091-107.

46.   Filippi P, Congdon JV, Hoang J, Bowling DL, Reber SA, Pašukonis A, et al. Humans recognize

emotional arousal in vocalizations across all classes of terrestrial vertebrates: evidence for acoustic

universals. Proceedings of the Royal Society B: Biological Sciences. 2017;284(1859):20170990.

47.   Kamiloğlu RG, Slocombe KE, Haun DBM, Sauter DA. Human listeners' perception of

behavioural context and core affect dimensions in chimpanzee vocalizations. Proceedings of the Royal

Society B: Biological Sciences. 2020;287(1929):20201148.

48.   Scheumann M, Hasting AS, Kotz SA, Zimmermann E. The voice of emotion across species:

how do human listeners recognize animals' affective states? PLOS ONE. 2014;9(3):e91192.

49.   Scheumann M, Hasting AS, Zimmermann E, Kotz SA. Human Novelty Response to Emotional

Animal Vocalizations: Effects of Phylogeny and Familiarity. Frontiers in Behavioral Neuroscience.

2017;11.

50.   Dricu M, Ceravolo L, Grandjean D, Frühholz S. Biased and unbiased perceptual decision-

making on vocal emotions. Scientific Reports. 2017;7(1):16274.

51.   Linnankoski I, Laakso M, Aulanko R, Leinonen L. Recognition of emotions in macaque

vocalizations by children and adults. Language & Communication. 1994;14(2):183-92.

52.   Debracque C, Clay Z, Grandjean D, Gruber T. Humans recognize affective cues in primate

vocalizations: Acoustic   and   phylogenetic   perspectives.   2022:Preprint   at

https://www.biorxiv.org/content/10.1101/2022.01.26.477864v1.

53.   Moss H, Abdallah S, Fletcher P, Bright P, Pilgrim L, Acres K, et al. Selecting among competing

alternatives: selection and retrieval in the left inferior frontal gyrus. Cerebral Cortex. 2005;15(11):1723-

35.

54.   Keuken MC, Hardie A, Dorn BT, Dev S, Paulus MP, Jonas KJ, et al. The role of the left inferior

frontal gyrus in social perception: an rTMS study. Brain Res. 2011;1383:196-205.

55.   Ceravolo L, Moisa M, Grandjean D, Ruff C, Fruhholz S. Disrupting inferior frontal cortex

activity alters affect decoding efficiency from clear but not from ambiguous affective speech.

2021:Preprint at https://www.biorxiv.org/content/10.1101/2021.12.15.472758v2.

56.   Nastase S, Iacovella V, Hasson U. Uncertainty in visual and auditory series is coded by

modality-general and modality-specific neural systems. Human Brain Mapping. 2014;35(4):1111-28.

57.   Toelch U, Bach DR, Dolan RJ. The neural underpinnings of an optimal exploitation of social

information under uncertainty. Social Cognitive and Affective Neuroscience. 2014;9(11):1746-53.

58.   Belin P, Fillion-Bilodeau S, Gosselin F. The Montreal Affective Voices: A validated set of

nonverbal affect bursts for research on auditory affective processing. ResearchGate. 2008.

59.   Ferdenzi C, Patel S, Mehu-Blantar I, Khidasheli M, Sander D, Delplanque S. Voice

attractiveness: Influence of stimulus duration and type. Behavior Research Methods. 2013;45(2):405-

13.

60.   Team R. RStudio: Integrated Development for R. RStudio. Boston MA: RStudio, Inc.; 2020.

61.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4.

Journal of Statistical Software. 2015;67(1):1-48.

62.   De Rosario-Martinez H, Fox J, Team R, De Rosario-Martinez MH. Package 'Phia'.

2015;Retrieved 1.

63.   Collins DL, Neelin P, Peters TM, Evans AC. Automatic 3D intersubject registration of MR

volumetric data in standardized Talairach space. J Comput Assist Tomogr. 1994;18(2):192-205.

64.   Ashburner J. A fast diffeomorphic image registration algorithm. NeuroImage. 2007;38(1):95-

113.

65.   Team RC. R: A language and environment for statistical computing. 2013.

66.   Eyben F, Scherer K, Schuller B, Sundberg J, André E, Busso C, et al. The Geneva Minimalistic

Acoustic Parameter Set (GeMAPS) for Voice Research and Affective Computing. IEEE transactions on

affective computing. 2016;7(2):190-202.

67.   Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, et al.

Automated Anatomical Labeling of Activations in SPM Using a Macroscopic Anatomical Parcellation

of the MNI MRI Single-Subject Brain. NeuroImage. 2002;15(1):273-89.

68.   Whitfield-Gabrieli S, Nieto-Castanon A. Conn: a functional connectivity toolbox for correlated

and anticorrelated brain networks. Brain Connect. 2012;2(3):125-41.