Decision-Making Under Effortful Constraints: The Impact of Acute High-Intensity Activity

on Cognitive Processes

Davranche Karen[1], Giraud Dorian[2],[3], Hays Arnaud[2], Gajdos Thibault[1]

[1]Aix Marseille Univ, CNRS, CRPN, Marseille, France

[2]Aix Marseille université, CNRS, Université de Toulon, Institut Paoli-Calmettes, HIPE

HumanLab, N°202324378C, Marseille, France

[3]Aix Marseille Univ, CNRS, IUSTI, Marseille, France

Corresponding author:

Dr Karen Davranche, Aix-Marseille Université, 3 place Victor Hugo,

13331 Marseille Cedex 03, France

Phone: +33(0)4 13 55 11 35

E-mail: karen.davranche@univ-amu.fr

ORCID: 0000-0003-4348-1265

Abstract

This study investigates the acute impact of high-intensity activity on decision-

making processes, using computational modeling to assess changes induced during and

after an effortful activity. Participants completed a perceptual decision-making task at rest,

both before and after a series of eight 5-minute cycling bouts (totaling 47 minutes) at 86 ±

7% of their maximum heart rate (HRmax). They also performed the decision-making task in

a dual-task condition (cognitive and physical effort) at various time points during cycling,

specifically during six of the eight 5-minute bouts. Cognitive processes were assessed by

fitting a drift diffusion model (DDM) to accuracy and reaction time data, allowing us to

assess changes in perceptual discrimination, decision bias, encoding of relevant

information, and decision execution. Findings suggest an improvement in cognitive

performance at rest immediately after the effortful activity, with significant changes in DDM

parameters indicating enhanced perceptual discrimination and more efficient non-

decisional processing compared to pre-activity. During the effortful activity, we observed a

decrease in decision thresholds over the course of the demanding dual-task, reflecting

reduced caution, alongside a dynamic adjustment in non-decisional time. These findings

provide new insights into how sustained high-intensity activity modulates decision-making

mechanisms, contributing to our understanding of effort's impact on cognitive processing.

Introduction

April 15, 2019. Notre-Dame is ablaze. After a strenuous climb, during which a firefighter analyzes and interprets the situation, she finally reaches the top of the tower. Now, she must decide which side of the raging fire to attack first. Is her decision-making affected by sustained physical and mental effort? And if so, which specific cognitive processes are altered during and after this demanding, effortful activity?

It is well established that acute moderate-intensity activity can enhance cognitive performance across various cognitive tasks, including memory, vigilance, and conflict resolution (e.g., Morris, 2021; Pontifex et al., 2019). However, recent narrative reviews have underscored the well-established effects of moderate-intensity aerobic activity on cognitive performance, contrasting with the less clarified impacts of high-intensity aerobic activity (Sudo, Costello, McMorris & Ando, 2022; Zheng et al., 2021; Pontifex et al., 2019). Despite theoretical expectations, the literature regarding the effects of high-intensity activity on cognitive performance is inconclusive and contradictory (McMorris, 2021; Moreau and Chou, 2019). While some studies support the notion of impaired cognitive performance following high-intensity activity, others suggest no significant effects or even improvements. This ambiguity highlights the complexity of the relationship between high-intensity effortful activity and cognitive function, indicating that further research is needed to fully understand these effects. The emphasis on moderate-intensity activity, primarily driven by methodological considerations, has

left a gap in our comprehension of the interaction between effortful activity and cognitive

performance, especially when physiological demands are substantial.

The impact of high-intensity activity, defined as physiological demand exceeding 80%

maximal oxygen uptake (McMorris et al., 2016), or equivalent [e.g., exceeding 80% maximal

heart rate (HRmax)], warrants deeper investigation. This is particularly crucial because it is

associated with more pronounced cellular and molecular changes than conventional aerobic

activity (Herold et al., 2019). Consequently, its influence on cognitive processes, likely mediated

by specific peripheral biomarkers associated with high-intensity effortful activity, such as

circulating lactate (Hashimoto et al., 2021), blood glucose, circulating cortisol levels, or

neurotrophic factors (Singh & Staines, 2015), may differ from the commonly reported findings in

existing literature. These systemic physiological adaptations could have widespread effects on the

organism, potentially altering cognitive functioning. For instance, they may enhance motor cortex

sensitivity to upstream influences, increase sensory sensitivity (facilitating stimulus encoding),

elevate arousal levels (accelerating processing speed), and modulate corticospinal excitability

(promoting response execution).

Quantifying the influence of effortful activity on cognitive processes poses a challenge

because these processes cannot be directly observed. The cognitive computational approach

provides a way to tackle this issue. It offers theoretically grounded models, whose parameters

represent latent cognitive processes and can be inferred from behavioral data. In this study, we

relied on the Drift Diffusion Model (DDM, Ratcliff, 1978) to decipher the impact of high-

intensity intermittent exercise on cognitive processes involved in a perceptual decision-making

task. This model has proven to be particularly fruitful (Ratcliff et al., 2016). It belongs to the

wide class of Evidence Accumulation Models (EAM, Bogacz et al., 2006), which postulate that

decisions result from the continuous accumulation of a stochastic evidence until a given amount

of evidence is attained. The basic DDM contains four parameters, each related to a different

cognitive process: the starting point of the accumulation process (z), which represents the bias of

the decision; the drift rate of the accumulation process (v), which depends on the difficulty of the

task or on the subject's perceptual ability; the distance between the bounds corresponding to each

response (a), which represent the subject's cautiousness (the larger a, the more accurate and

slower the responses); and a non-decision time (Ter), that encompass the encoding of relevant

information and the decision execution processes. Specifically, participants performed a random

dot motion task, a task for which the DDM consistently offers excellent explanations for

behavioral data, both before, during, and after engaging in an effortful high-intensity activity

consisting in eight 5-minutes bouts cycling exceeding 80% HRmax. Cognitive processes were

assessed by determining the best-fitting parameters of a model based on behavioral data

(accuracy and reaction times).

Materials and methods

Participants

Nineteen young adults (one woman, age range: 18–34; mean age: 25 years) voluntarily

participated in the experiment. All participants were healthy and practiced regular physical

activities (peak oxygen uptake: 55 ± 8 mL kg-1 min-1; maximal heart rate: 184 ± 9 bpm). All

participants were fully informed about the study and signed a written informed consent before

inclusion. The study was approved by the Ethics Committee for Research in Science and

Technology of Physical Sports Activities (IRB00012476-2022-21-01-148).

Random dot motion task (RDK)

We used a perceptual decision-making task, the random dot motion task (RDK), for

which the DDM has consistently provided excellent accounts of behavioral data. The experiment

was programmed in Python, utilizing components of the PsychoPy toolbox (Peirce et al., 2019),

and was executed on a PC computer operating natively on Windows 10 with a 19.5-inch screen.

Participant responses were recorded using two buttons (one for each hand).

The task requires participants to determine the global direction of a random dot

kinematogram featuring a proportion of dots moving coherently in the left or right signal

direction. Each trial started with the presentation of the random dot motion stimulus, which

remained on the screen until the participant responded. A response time deadline was set to 5 s.

The interval between the response to the stimulus and the next trial was 1.5 s. The coherence

parameter (proportion of dots moving in the same direction) was set for each participant at the

beginning of the experiment according to a 2 up and 1 down staircase method lasting 30 trials.

White dots were presented within a virtual 12.6° circular aperture centered on a 24.8° 3 13.9°

black field. Each dot was a 4 x 4 pixel (0.05° square), moving at a speed of 8°/s. Dot density was

fixed at 16.7 dots/deg2/s. Random dot motion was controlled by a white noise algorithm (Pilly &

Seitz, 2009).

Procedure

Participants made three separate visits to the laboratory, each spaced at least 48 hours

apart: a characterization session, a familiarization session, and the experimental session.

Participants were positioned on an upright cycle ergometer (Lode Excalibur Sport, Groningen,

The Netherlands), equipped with two thumb response buttons on the right and left handle grips. A

computer screen was positioned at eye level, approximately 1 meter in front of the participant.

The preliminary session was conducted to individually adjust the difficulty of the

cognitive task, assess $\dot{V}O2max$, and determine ventilatory thresholds (Wasserman, 2012) (see

Table 1 for details). Participants were seated on the cycle ergometer, fitted with a facemask to

measure gas exchange data (Innocor CO, Cosmed, Italia) and a heart rate monitor. The

incremental cycling activity to exhaustion test started at an initial power level defined according

to the participant's level (100 watts for non-cyclists and 150 watts for cyclists) and increased by

30 watts every 2 minutes until exhaustion. The end of the test was determined by volitional

cessation of exercise or failure to maintain pedal cadence above 60 rpm despite strong verbal

encouragement. During the first step of this maximal test, 30 trials of the RDK task were

performed to set up the consistency parameter according to a staircase procedure.

Table 1. *Anthropometric and physiological characteristics of participants.*

Mean ± standard deviation	
Variables	**All**
Sample size	19
Age (years)	25±5
Height (cm)	177±7
Weight (kg)	67±9
Fat mass (%)	15±4
$\dot{V}O_2$max (ml.kg^{-1}.min^{-1})	55±8
Heart rate max (bpm)	184±9
Maximal aerobic power (W)	295±58
P_{SV1} (W)	208±55
P_{SV2} (W)	254±52
Heart rate HIE (% max)	86±6
Central temperature HIE (°C)	38,2±0,4

Note. P_{SV}: Power corresponding to the first SV1 and second SV2 ventilatory thresholds, HIE:

high-intensity exercise, $\dot{V}$O2max: maximal oxygen consumption.

Before the experimental session, a familiarization session (about 1,250 trials), identical in

all respects to the experimental session, was conducted to stabilize performance and control for

potential learning effects, as well as to assess the physiological load of the dual task performed

while cycling. The data of the familiarization session were not included in the analyses.

The experimental protocol involved high-intensity interval activity, a discontinuous

mode of endurance exercise characterized by short bouts of high-intensity workloads interspersed

with periods of rest. Participants completed a perceptual decision-making task at rest both before

(PRE) and after (POST) eight 5-minute bouts of cycling. Each 5-minute cycling bout was

followed by a 1-minute rest period, resulting in a total session duration of 47 minutes. During six

of the eight cycling bouts, participants performed the perceptual decision-making task while

cycling at a workload fixed at 90% of SV1—an intensity determined during the incremental test

of the preliminary session—placing them in a dual-task situation involving both cognitive and

physical effort. Cardiorespiratory parameters were continuously recorded, and capillary blood

samples from the fingertip were collected before and immediately after each of the 5-minute

cycling bouts for the measurement of lactate and glucose levels. The skin was prepared by

cleaning with alcohol, allowed to dry, and then punctured with an automated lancet. Blood

samples were analyzed using a Biosen blood analyzer (EKF Diagnostics, UK).

Data Analyses

The experiment was designed to evaluate the exercise-induced effects on cognitive

processes throughout high-intensity activity and immediately afterward the effortful protocol.

Due to the interplay between physical demands and cognitive performance in the dual-task

condition, cognitive data recorded at rest (before and after, 4916 trials) and during the dual-task

condition (6 blocks while cycling, 12402 trials) were analyzed separately.

During the effortful activity, we anticipated that the impacts on decision-making

processes may vary according to time spent on task. To prevent the loss of detailed aspects of

how effort affects cognitive performance, we conduct analyses on three time-windows: the

Beginning of the sustained physical effort spanning from 1 to 6 minutes (B1); the Middle of

physical effort, covering minutes 19 to 30 (B4-B5); and the End of physical effort, encompassing

minutes 37 to 48 (B7-B8). The cognitive performance recorded during the first 5-minute physical

effort block (B1) was used as a baseline. This decision was made because the cognitive task is

performed in a dual-task condition (while cycling), and at that time, the physiological effect

induced by exercise had not yet been fully established.

Anticipations (RTs < 200 ms; 0.13 %) were discarded from all analyses. Five subjects did

not perform the perceptual task appropriately (i.e., exhibiting a percentage of accuracy below

50% or employing a stereotyped response strategy). They were excluded from the initial sample

of 24 participants. We conducted a mixed-effects generalized logistic regression analysis on

accuracy and performed a linear mixed-effects model analysis on correct reaction time, including

'block' as a fixed effect, with 'subject' and 'block' intercepts treated as random effects.

Regarding physiological measures, we analyzed glucose and lactate levels within linear

mixed-effects models. These models included the 'sample' (PRE and B1 to B8) expressed in

mmol/L as a fixed effect, with 'subject' as random effect.

The Drift Diffusion Model parameters were estimated using a hierarchical Bayesian

procedure implemented with the DMC package (Heathcote et al., 2019) for the R software (R

Core Team, 2016). Specifically, we considered the DDM with inter-trial variability in the drift

rate (sv) and the starting point (sz), and let the drift (v), threshold (a), non-decision time (Ter) and

drift rate variability vary across conditions. We used broad truncated normal distributions for the

means and uniform distribution on [0,1] for the standard-deviations. In order to fit the model to

the data, we used the two-steps procedure described by Heathcote et al. (2019). In a first step, we

fitted each participant's data individually. The averaged resulting parameters were used in a

second step as starting values for hierarchical sampling. The quality of the fit was evaluated by

visually inspecting Markov chain Monte Carlo chains, comparing predicted data according to

best fitting parameters to actual data, and computing Brooks and Gelman's (1998) R statistics

(see Appendix). Finally, we tested the presence of an effect of the condition on the parameter

values by computing Bayesian p values (Heathcote et al., 2019).

Results

Physiological measures

The workload set for the high-intensity activity corresponded to 86% ± 7% of HRmax,

ranging from 71% to 99%. Linear mixed model demonstrated a significant effect of block ($F_{(7,}$

$_{126)}$ = 15.114, $p < .001$). Heart rate was lower during the first two blocks and then remained

stable from the 3rd block to the end of exercise (Figure 1). Specifically, block B1 differed from

the blocks B3, B4, B5, B6, B7 and B8; and the block B2 differed from the blocks B5, B6, B7 and

B8.

Figure 1. Heart rate as function of exercise duration (bmp/min)

The linear mixed model performed on blood sample analyses confirmed a significant

impact of the effortful activity on glucose levels (F(8, 143.03) = 6.2595, p < 0.001) and on

lactate levels (F(8, 143.01) = 21.21, p < 0.001) collected at the end of each 5-minute cycling

bout. Post-hoc comparisons revealed lower glucose levels after each bout than before sustained

physical effort (PRE = 4.70 mmol/L ± 0.76 mmol/L). No other comparisons were found to be

statistically significant on glucose levels (Figure 2).

Figure 2. Glucose as function of exercise duration (mmol/L)

For the lactate concentration, higher lactate levels were observed after each 5-minute

cycling bout than before physical effort (PRE = 1.66 ± 0.56 mmol/L). The lactate level after the

first bout (B1 = 6.05 ± 2.22 mmol/L) was also significantly higher from those collected after B4

(4.66 ± 2.47 mmol/L), B5 (4.42 ± 2.29 mmol/L), B6 (4.49 ± 2.50 mmol/L), B7 (4.47 ± 2.66

mmol/L) and B8 (4.14 ± 2.73 mmol/L). Higher lactate level was also observed after the second

bout (B2 = 5.66 ± 2.59 mmol/L) compared to B7 and B8. No other comparisons were found to be

statistically significant (Figure 3).

Figure 3. Lactate as function of exercise duration (mmol/L). Note: 4 mmol/L is a classical
marker used to define the onset of blood lactate accumulation (OBLA), considered a valid
indicator for transition from aerobic to anaerobic performance.

Behavioral data

Accuracy. The mixed-effects generalized logistic regression analysis conducted on

accuracy did not reveal any significant effort-induced effects. The accuracy was not significantly

different following physical effort compared to before (PRE = 87.67% vs POST = 86.64%, p =

0.891), nor during the sustained physical effort compared to the baseline (Baseline = 85.91%;

Middle of physical effort = 86.28%, p = 0.917), nor to the End of physical effort (86.21%, p =

0.525).

Correct reaction time (RT). The linear mixed model conducted on correct RTs collected

at rest before the effortful activity (PRE = 996 ms ± 634 ms) and after (POST = 929 ms ± 625

ms) revealed a trend toward an effect that did not reach the level of significance $(F(1, 17.316) =$

$3.9154, p = 0.064)$. The linear mixed model conducted on correct RTs collected during the high-

intensity activity, in the dual task situation, revealed a significant effect of block $(F(2, 16.597) =$

$8.0106, p < 0.01)$. Post-hoc comparisons showed that RTs were faster at the End of effort

(covering minutes 37 to 48, B7-B8: 910 ms ± 562 ms, $p < 0.001$) compared to the baseline in the

dual task situation (covering minutes 1 to 6, B1: 1001 ms ± 631 ms). Reaction times in the

Middle of effort (covering minutes 19 to 29, B4-B5: 942 ms ± 602 ms) were not significantly

different from those in the baseline $(p = 0.269)$ (Figure 4).

Figure 4. Reaction time (ms) as function of exercise duration Error bar represent SD.

DDM parameters. We investigated which component of the perceptual decision-making process

was impact by effort by comparing the best-fitting parameters of the DDM across conditions at

the group level (detailed results of the fitting procedure are available in the Supplementary Data).

Post-effort Effects. We observed significant changes in the parameters of DDM after the

effortful activity compared to before. The results indicated a general improvement in cognitive

processes at rest associated with perceptual decision-making. The drift rate (v) increased

significantly after effort compared to before, the boundary (a) increased, and the non-decision

time (Ter) decreased by approximately 64 milliseconds (Figure 5 and Table 2, for details).

Figure 5. Diffusion model best-fitting parameters before and after high-intensity activity

Table 2. *Diffusion model best-fitting parameters before and after exercise, with 95% credible intervals and p values.*

Parameters	Before	After	Proba
v: drift rate	1.780	1.982	Delta : +0.202
	[1.695, 1.867]	[1.896, 2.068]	p<0.001
a: upper decision threshold	1.943	2.040	Delta : +0.096
	[1.880, 2.011]	[1.973, 2.111]	p < 0.05
Ter: non-decision time	0.504	0.445	Delta : -0.058
	[0.489, 0.516]	[0.431, 0.457]	p<0.001

In order to quantify the effect of the modulation of the v and a, we performed simulations, using best fitting parameters before the effortful activity (as a baseline) and varying each parameter separately to its best-fitting value after effort. According to the results of these simulations, the increase of the drift rate led to a decrease in RT by approximately 48 ms and an increase in accuracy by 1.1 percentage points. Conversely, the boundary increase resulted in an RT increase of 31 ms while improving accuracy by 0.48 percentage points.

During-effort Effects. We did not observe any modulation of the drift rate (v) in either

the first or second part of the effortful activity. The boundary (a) decreased gradually as time

passed, and the non-decision time (Ter) increased in the Middle of physical effort by 18 ms and

decreased afterward by 14 ms (see Figure 6 and Table 3, for details).

Figure 6. Diffusion model best-fitting parameters during high-intensity activity

Table 3. *Diffusion model best-fitting parameters during exercise, with 95% credible intervals*

and p values.

Parameters	Baseline	Middle	End	Proba
v: drift rate	1.862	1.860	1.926	Baseline vs End (-0.064):
	[1.758,	[1.790,	[1.852,	p=0.163
	1.968]	1.934]	2.001]	Middle vs End (-0.066):
				p=0.1
				Baseline vs Middle
				(+0.002): p=0.488
a: upper decision threshold	2.158	2.027	1.959	Baseline vs End (+0.198): p<0.001

	[2.077, 2.243]	[1.973, 2.086]	[1.911, 2.011]	Middle vs End (+0.067): p=0.038 Baseline vs Middle (+0.131): p=0.005
Ter: non decision time	0.451 [0.438, 0.463]	0.469 [0.459, 0.478]	0.455 [0.447, 0.462]	Baseline vs End (-0.003): p=0.314 Middle vs End (+0.014): p=0.006 Baseline vs Middle (-0.018): p=0.008

Discussion

Summary

The present study aimed to investigate the impact of high-intensity activity on cognitive

processes involved in perceptual decision-making. To achieve this, participants completed a

perceptual decision task before, after, and during a dual-task condition involving both cognitive

and physical effort. Cognitive processes were assessed by fitting a drift diffusion model (DDM)

to behavioral data collected across these time periods.

Behavioral data indicated reduced RTs during the high-intensity activity in the dual task

condition and suggested a trend toward enhanced cognitive performance at rest immediately

afterward. Moreover, the DDM provided new insights into the mechanisms underlying this

enhancement by revealing changes in main diffusion model parameters (drift rate, boundary

separation, and non-decision time) following effortful activity compared to before. In the dual-

task condition, the impact of effortful activity on cognitive processes followed a more complex

pattern, as discussed below. These findings highlight the need for further investigation to better

understand the relationship between high-intensity physical effort and cognition, offering

promising directions for future research.

282   *Post-effort modulation of cognitive processes.*

283   Reaction times recorded immediately after the effortful activity tended to be shorter (-70

ms) without a change in accuracy. This trend was further supported by computational modeling,

which revealed significant post-effort changes in diffusion model parameters. Specifically, the

results suggested an improvement in perceptual discrimination, more efficient non-decisional

processes (perceptual encoding and motor execution), and a more cautious decision strategy.

These findings align with previous research demonstrating the impact of physiological states on

cognitive processes. For instance, Ratcliff & Van Dongen (2011) showed that vigilance states

induced by sleep deprivation modulate the drift rate in DDM, while Du Rietz et al. (2019)

reported a positive association between high-intensity activity and changes in brain indices

related to arousal and attentional allocation. In line with this, elevated neuronal activity and

metabolism induced by effortful activity have been proposed to contribute to cognitive

enhancement (Ide & Secher, 2000).

Beyond these neural mechanisms, systemic physiological adaptations may also play a

crucial role. Cognitive improvements following high-intensity activity have been observed

even after exercise cessation (Tsukamoto et al., 2016; Moreau & Chou, 2019; McMorris et

al., 2016), suggesting a lasting effect of metabolic and neurochemical changes. Notably,

increased blood lactate has been shown to enhance motor system reactivity and sensory

perception (Coco et al., 2009, 2016), which could partially explain the observed

improvements in decision-making efficiency.

One possible underlying mechanism is the increase in cerebral blood flow and oxygenation,

which has been linked to cognitive benefits in response to exercise (Rooks et al., 2010).

Additionally, the metabolism of catecholamines and the role of lactate—continuously

produced even under fully aerobic conditions—warrant further exploration as potential

contributors to these cognitive enhancements. Investigating these physiological pathways

could provide deeper insight into the complex interactions between high-intensity effort

and cognitive performance.

Ongoing physical effort impact on cognitive processes.

Behavioral data indicated that reaction times (RTs) were shorter at the end of the physical

effort (minutes 37 to 48) compared to the first five minutes of exercise (-91 ms), with no

significant modulation of accuracy. This suggests a facilitating effect of prolonged physical effort

on response speed. To better understand the cognitive mechanisms underlying this effect, we

examined the parameters of the model.

First, we observed a progressive decrease in the decision threshold over the course of the physical

effort, indicating that participants became less cautious as time passed. Notably, while this

reduction in caution led to a substantial time gain (56 ms between the beginning and the end of

physical effort), the associated decrease in accuracy remained relatively small (0.73 percentage

points). This pattern suggests that physical exertion promotes a strategic shift in decision-making,

where participants prioritize speed over absolute accuracy. Such an adaptation could be linked to

increased arousal and enhanced neural efficiency due to heightened catecholaminergic activity

(McMorris et al., 2016; Moreau & Chou, 2019) or metabolic changes such as increased cerebral

lactate availability, which has been shown to influence sensorimotor processing (Coco et al.,

2009; 2016). The minimal trade-off between speed and accuracy may indicate that the

physiological adaptations induced by sustained effort enable participants to maintain performance

efficiency while adopting a more rapid decision strategy.

Second, analysis of the non-decision time (Ter) revealed a dynamic pattern: the duration of

processes unrelated to the perceptual decision increased during the middle phase of the physical

effort (minutes 19 to 30) compared to the first five minutes but then diminished toward the end of

the effort (minutes 37 to 48). We formulated two hypotheses. On the one hand, the initial rise of

Ter may be attributed to competition for biomechanical, physiological, and/or attentional

resources (e.g., McMorris, 2021; Sudo et al., 2022). On the other hand, as blood lactate

concentration appears to follow a similar pattern, the decline in lactate levels/production in

the second half of the activity compared to the first half might fail to adequately support

elevated brain neuronal activity and metabolism, thereby impacting cognitive performance.

Indeed, the implementation of physiological adaptations, potentially coupled with improved

lactate clearance and a shift in dominant energy metabolism in the later stages of physical effort,

could explain the subsequent decrease in Ter (Herold et al., 2022; Hashimoto et al., 2018;

Tsukamoto et al., 2016).

Finally, we found no modulation of the drift rate. This indicates that we did not observe any

impact of physical effort on subjects' perceptual discrimination abilities.

At this point, it is important to acknowledge a limitation of this study: the substantial

inter-individual variability in physiological responses to exercise at 90% of the first ventilatory

threshold. This variability, reflected in heart rates ranging from 71% to 99% of HRmax, suggests

that the relative contributions of aerobic and anaerobic metabolism differed across participants.

Given that effort-induced cognitive effects may depend on metabolic demands, such

heterogeneity could partly explain inconsistencies in our findings.

Post-hoc examination of individual lactate levels further confirmed this variability, as

participants were roughly evenly split between those above and below the 4 mmol/L threshold,

commonly associated with the onset of blood lactate accumulation (OBLA). Since OBLA marks

the transition from predominantly aerobic to anaerobic energy production, future studies should

account for this factor by systematically assessing metabolic thresholds to better capture the

relationship between exercise intensity, energy metabolism, and cognitive performance.

Additionally, individual differences in fitness levels may play a critical role in shaping

post-exercise cognitive effects. Higher fitness levels are associated with more efficient lactate

clearance, which could influence the duration and magnitude of post-exercise cognitive

enhancements. Furthermore, tolerance to elevated lactate concentrations varies across

individuals, potentially affecting the extent to which metabolic byproducts contribute to cognitive

modulation. This underscores the need for future research to incorporate measures of aerobic

capacity and lactate tolerance to refine our understanding of the interplay between physical effort,

metabolic responses, and cognitive function.

By applying DDM to a perceptual decision-making task, this study provides new insights

into the cognitive mechanisms underlying high-intensity exercise-induced changes. Our findings

suggest that sustained physical effort facilitates a shift toward faster, albeit slightly less cautious,

decision-making, potentially mediated by metabolic and neurophysiological adaptations.

Furthermore, post-exercise improvements in decision efficiency highlight the complex interplay

between physiological states and cognitive performance. Given the observed inter-individual

variability, future research should systematically assess metabolic thresholds, aerobic fitness, and

lactate clearance efficiency to refine our understanding of these effects. High-intensity exercise,

being time-efficient and ecologically valid, emerges as a promising alternative to traditional

moderate-intensity protocols for cognitive enhancement, warranting further investigation.

Acknowledgements

We would like to thank the Filière Fastspor'in for its support in carrying out this study,

and Anita Vergnaud et Laurine Stefanuto for their assistance in data collection.

Declarations of interest statement

The authors declare that they have no competing interests. No funds, grants, or other

support was received. The authors have no relevant financial or non-financial interests to

disclose.

References

Ando, S., Komiyama, T., Tanoue, Y., Sudo, M., Costello, J. T., Uehara, Y., & Higaki, Y.

(2022). Cognitive Improvement After Aerobic and Resistance Exercise Is Not Associated With

Peripheral Biomarkers. *Frontiers in Behavioral Neuroscience*, 16, 853150.

https://doi.org/10.3389/fnbeh.2022.853150

Bogacz, R., Brown, E., Moehlis, J., Holmes, P., & Cohen, J. D. (2006). The physics of

optimal decision making: A formal analysis of models of performance in two-alternative forced-

choice tasks. *Psychological Review*, 113, 700-765. https://doi.org/10.1037/0033-295X.113.4.700

Brooks, S. P., & Gelman, A. (1998). General Methods for Monitoring Convergence of

Iterative Simulations. *Journal of Computational and Graphical Statistics*, 7(4), 434-455.

https://doi.org/10.1080/10618600.1998.10474787

Du Rietz, E., Barker, A. R., Michelini, G., Rommel, A.-S., Vainieri, I., Asherson, P., &

Kuntsi, J. (2019). Beneficial effects of acute high-intensity exercise on electrophysiological

indices of attention processes in young adult men. *Behavioural Brain Research*, 359, 474-484.

https://doi.org/10.1016/j.bbr.2018.11.024

Hashimoto, T., Tsukamoto, H., Ando, S., & Ogoh, S. (2021). Effect of Exercise on Brain

Health: The Potential Role of Lactate as a Myokine. *Metabolites*, 11(12), 813.

https://doi.org/10.3390/metabo11120813

Hashimoto, T., Tsukamoto, H., Takenaka, S., Olesen, N. D., Petersen, L. G., Sorensen,

H., et al. (2018). Maintained exercise-enhanced brain executive function related to cerebral

lactate metabolism in men. *FASEB Journal*. 32, 1417–1427. doi: 10.1096/fj.201700381RR

Heathcote, A., Lin, Y.-S., Reynolds, A., Strickland, L., Gretton, M., & Matzke, D. (2019).

Dynamic models of choice. *Behavior Research Methods*, 51(2), 961-985.

https://doi.org/10.3758/s13428-018-1067-y

Herold F, Behrendt T, Meißner C, Müller NG, Schega L. The Influence of Acute Sprint

Interval Training on Cognitive Performance of Healthy Younger Adults. *International Journal of*

Environmental Research and *Public Health*. 2022 Jan 5;19(1):613. doi:

10.3390/ijerph19010613. PMID: 35010873; PMCID: PMC8745010.

Herold, F., Müller, P., Gronwald, T., & Müller, N. G. (2019). Dose–Response Matters! A

Perspective on the Exercise Prescription in Exercise–Cognition Research. *Frontiers in*

Psychology, 10, 2338. https://doi.org/10.3389/fpsyg.2019.02338

McMorris, T., Turner, A., Hale, B. J., & Sproule, J. (2016). *Chapter 4 - Beyond the*

Catecholamines Hypothesis for an Acute Exercise–Cognition Interaction: A Neurochemical

Perspective. In T. McMorris (Ed.), Exercise-Cognition Interaction (p. 65-103). Academic Press.

https://doi.org/10.1016/B978-0-12-800778-5.00004-9

McMorris, T. (2021). The acute exercise-cognition interaction: From the catecholamines

hypothesis to an interoception model. *International Journal of Psychophysiology,* 170, 75–88.

doi: 101016/j.ijpsycho.2021.10.005

Moreau, D., & Chou, E. (2019). The Acute Effect of High-Intensity Exercise on

Executive Function: A Meta-Analysis. *Perspectives on Psychological Science*, 14(5), 734-764.

https://doi.org/10.1177/1745691619850568

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman,

E., & Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior Research*

Methods, 51(1), 195-203. https://doi.org/10.3758/s13428-018-01193-y

Pilly, P. K., & Seitz, A. R. (2009). What a difference a parameter makes: A

psychophysical comparison of random dot motion algorithms. *Vision Research*, 49(13),

1599-1612. https://doi.org/10.1016/j.visres.2009.03.019

Pontifex, M. B., McGowan, A. L., Chandler, M. C., Gwizdala, K. L., Parks, A. C., Fenn,

K., & Kamijo, K. (2019). A primer on investigating the after effects of acute bouts of physical

activity on cognition. *Psychology of Sport and Exercise*, 40, 1-22.

https://doi.org/10.1016/j.psychsport.2018.08.015

Ramdani, C., Sagui, E., Schmid, B., Castagna, O., Davranche, K., Vidal, F., &

Hasbroucq, T. (2021). Action monitoring fails when motor execution is too fast: no time for

correction. *Journal of Systems and Integrative Neuroscience*, 7(3).

https://doi.org/10.15761/JSIN.1000246

Ratcliff, R. (1978). A theory of memory retrieval. Psychological review, 85(2), 59.

Ratcliff, R., Smith, P. L., Brown, S. D., & McKoon, G. (2016). Diffusion Decision Model:

Current Issues and History. *Trends in Cognitive Sciences*, 20(4), 260-281.

https://doi.org/10.1016/j.tics.2016.01.007

Ratcliff, R., & Van Dongen, H. P. A. (2011). Diffusion model for one-choice reaction-

time tasks and the cognitive effects of sleep deprivation. *Proceedings of the National Academy of

Sciences*, 108(27), 11285-11290. https://doi.org/10.1073/pnas.1100483108

Singh, A. M., & Staines, W. R. (2015). The effects of acute aerobic exercise on the

primary motor cortex. *Journal of Motor Behavior*, 47(4), 328–339.

Sudo, M., Costello, J. T., McMorris, T., & Ando, S. (2022). The effects of acute high-

intensity aerobic exercise on cognitive performance: A structured narrative review. *Frontiers in

Behavioral Neuroscience*, 16, 957677. https://doi.org/10.3389/fnbeh.2022.957677

Wasserman, K. (2012). *Principles of Exercise Testing and Interpretation: Including

Pathophysiology and Clinical Applications.* Wolters Kluwer Health/Lippincott Williams &

Wilkins.

450   Zheng, K., Zou, L., Wei, G., & Huang, T. (2021). Concurrent Performance of Executive

451   Function during Acute Bouts of Exercise in Adults: A Systematic Review. *Brain Sciences*,

452   11(10), 1364. https://doi.org/10.3390/brainsci11101364