Histone variant H2B.Z acetylation is necessary for maintenance of *Toxoplasma gondii* biological fitness

Laura Vanagas[1], Daniela Muñoz[1], Constanza Cristaldi[1], Agustina Ganuza[1], Rosario Nájera[1], Mabel C. Bonardi[1],

Valeria R. Turowski[2], Fanny Guzman[3], Bin Deng[4], Kami Kim[5], William J. Sullivan Jr.[6] and Sergio O. Angel[1]

[1]Laboratorio de Parasitología Molecular, INTECH, CONICET-UNSAM, Av. Intendente Marino Km. 8.2, C.C 164, (B7130IIWA),

Chascomús, Prov. Buenos Aires, Argentina.

[2]Laboratorio de Bioquímica y Biología Celular de Parásitos, INTECH, CONICET-UNSAM, Av. Intendente Marino Km. 8.2, C.C

164, (B7130IIWA), Chascomús, Prov. Buenos Aires, Argentina.

[3]Núcleo de Biotecnología Curauma, Pontificia Universidad Católica de Valparaiso. Av. Universidad 330 Curauma, Valparaiso

[4]Department of Biology and VBRN, University of Vermont, Vermont, USA.

[5] Department of Internal Medicine, Morsani College of Medicine, University of South Florida, Tampa, FL 33612

[6] Department of Pharmacology and Toxicology, Indiana School of Medicine, Indianapolis, Indiana 46202, USA

Corresponding author: Sergio O. Angel. Laboratorio de Parasitología Molecular, IIB-INTECH, CONICET-UNSAM, Av.
Intendente Marino Km. 8.2, C.C 164, (B7130IIWA), Chascomús, Prov. Buenos Aires, Argentina. Tel.: +542241424303

E-mail address: sangel@intech.gov.ar (SO Angel)

Abstract

Through regulation of DNA packaging, histone proteins are fundamental to a wide array of biological processes. A variety of post-translational modifications (PTMs), including acetylation, constitute a proposed histone code that is interpreted by "reader" proteins to modulate chromatin structure. Canonical histones can be replaced with variant versions that add an additional layer of regulatory complexity. The protozoan parasite *Toxoplasma gondii* is unique among eukaryotes in possessing a novel variant of H2B designated H2B.Z. The combination of PTMs and the use of histone variants is important for gene regulation in *T. gondii,* offering new targets for drug development. In this work, *T. gondii* parasites were generated in which the 5 N-terminal acetylatable lysines in H2B.Z were mutated to either alanine (c-Myc-A) or arginine (c-Myc-R). c-Myc-A mutant only displayed a mild effect in its ability to kill mice. c-Myc-R mutant presented an impaired ability to grow and an increase in differentiation to latent bradyzoites. This mutant line was also more sensitive to DNA damage, displayed no virulence in mice, and provided protective immunity against future infection. While nucleosome composition was unaltered, key genes were abnormally expressed during *in vitro* bradyzoite differentiation. Our results show that the N-terminal positive charge patch of H2B.Z is important for these procceses. Pull down assays with acetylated N-terminal H2B.Z peptide and unacetylated one retrieved common and differential interactors. Acetylated peptide pulled down proteins associated with chromosome maintenance/segregation and cell cycle, opening the question of a possible link between H2B.Z acetylation status and mitosis.

Key words: *Positive charge patch – cell cycle – differentiation – virulence – mitosis*

Introduction

Histones proteins function to form nucleosomes that regulate packaging of DNA, thereby affecting

DNA transcription, replication, and repair. In addition to the four canonical histones (H2A, H2B, H3 and H4),

variant histones exist that can be substituted into the nucleosome, affecting its properties. Canonical and

variant histones are also subject to a wide array of post-translational modifications (PTMs) that can alter

the nucleosome or recruit chromatin remodeling machinery. The combination of PTMs and exchange of

histone variants is believed to be important for gene regulation in the protozoan parasite *Toxoplasma*

gondii, offering new targets for drug development [1-3].

T. gondii is a member of phylum Apicomplexa and infects between 10-90% of the population

depending on the country [4], presumably influenced by dietary habits and environmental conditions [5].

T. gondii converts from rapidly growing tachyzoites, which cause acute infection, to slow or non-growing

bradyzoites, which cause chronic infection and typically reside in the brain [6]. Although *T. gondii* infection

is asymptomatic in healthy people, untreated clinical toxoplasmosis can be lethal in immunocompromised

individuals such as transplant or HIV patients, and can produce severe disease during congenital infection,

especially in early stages of development [7, 8]. Chronic toxoplasmosis has also been linked to brain tumor

predisposition, attention-deficit/hyperactivity disorder, obsessive-compulsive disorder and schizophrenia

[9-13].

The *T. gondii* life cycle is comprised of sexual and asexual stages. While sexual replication takes

place only in felids, *T. gondii* is able to replicate asexually as tachyzoites in any nucleated cell in any warm-

blooded vertebrate. Following infection, tachyzoites convert into bradyzoites, which are house in tissue

cysts that are impervious to immunity and current drug treatments. Recently, a myb-like transcription

factor (BFD1) was described to be necessary and sufficient to induce bradyzoite differentiation [14]. How

BFD1 is recruited to stage-specific promoters remains to be determined, but likely involves interplay with

chromatin remodelers and epigenetic processes that include histone PTMs and/or variant histone exchange

[15]. In support of this idea, treatment of tachyzoites with inhibitors of histone deacetylase 3 (HDAC3)

initiates bradyzoite differentiation [16]. The roles of histone variants in *T. gondii* remain poorly understood.

T. gondii expresses histone variants that we have previously shown mark functional regions of the

genome [17]. H2A.Z is a ubiquitous variant that has been implicated in both transcriptional activation and

gene silencing in eukaryotes, and in fine regulation of important processes as DNA damage repair [18-20],

which is also present in *T. gondii* [21]. Unlike H2A.Z, only specialized isoforms of the H2B histone family

have been described in nature, except for apicomplexan parasites [22-25] and trypanosomatids [26], in

which a H2B.Z and a H2Bv variants, were identified, respectively.

Histones H2A.Z and H2B.Z form a dimer that localizes with the transcriptional activation mark

H3K4me3 in promoter/transcriptional start site (TSS) regions surrounding the nucleosome-free region

upstream of the transcription start site [17]. In addition, H2B.Z and H2A.Z histones localize to the gene

bodies of silent genes, including repressed stage-specific genes, suggesting a role in the regulation of stage

transitions. In *T. gondii*, both H2A.Z (TGGT1_300200) and H2B.Z (TGGT1_209910) are essential during the

lytic cycle with CRISPR fitness scores of -5.08 and -4.05, respectively [27]. By mass spectrometry, both

histone variants were found to be hyperacetylated at the N-terminal domain, whereas few or no acetylation

marks were identified on canonical H2A, H2B and H2A.X [28]. *T. gondii* H2B.Z was shown to be acetylated

at 5 lysine residues in its N-terminal tail. The H2A.Z N-terminal tail has 10 acetylatable lysines, in which

lysine 18 can also be methylated [28]. H2A.Z N-terminal tail acetylation has been widely shown to be a

hallmark for active chromatin whereas N-terminal methylation is associated with gene silencing [29-32],

suggesting it is essential for changes in gene expression during cell differentiation [33-37]. While the role

of N-terminal lysine acetylation in H2B.Z has not yet been studied, N-terminal acetylated canonical H2B is

associated with some, but not all, active genes in vertebrates [38]. Interestingly, H2A.Z in most eukaryotes

only bears 4-5 acetylatable lysine residues in the N-terminus, while in protists like *T. gondii, Plasmodium* or

Tetrahymena termophila, this number is considerably higher (between 5 and 16). Moreover, the presence

of a *T. gondii* double variant nucleosome that altogether carries 15 acetylatable lysines that could be

regulated is highly intriguing.

In the present work we studied the role of the five *T. gondii* H2B.Z N-terminal tail lysines, using a

mutagenesis strategy; we also pursued a gene knockout strategy for H2B.Z. In an RH strain background, we

generated *T. gondii* that possess mutated versions of H2B.Z: c-Myc-R, in which the five N-terminal

acetylatable lysines were replaced by arginines, and c-Myc-A, in which they were replaced by alanines.

These lines were analyzed for their ability to grow *in vitro*, differentiate to bradyzoites *in vitro* and produce

virulence in mice. In addition, expression of key genes was studied, as well as nucleosome composition and

DNA damage sensitivity. Our results suggest that regulation of the N-terminal positive charge patch by

lysine acetylation, rather than the histone code, is essential for the aforementioned biological processes.

We have also found some differential interactors for an acetylated N-terminal H2B.Z peptide, compared to

a unacetylated one, opening new questions on the function of this modification and this histone variant.

The implications of these data on the putative role of H2B.Z is discussed.

Results

Sequence alignment of Apicomplexa H2B.Z N-tail

H2B.Z is a variant histone of the H2B family that appears early in the phylum Apicomplexa.

Previously, it was observed that *T. gondii* and *Plasmodium falciparum* H2B.Z N-terminal tails contain five

acetylated lysines (K4,K9,K13,K14 and K18) [28, 39, 40]. The conservation of the five acetylatable lysines is

clear from the alignment of H2B.Z N-terminal tail sequences from different Apicomplexa (Fig. 1A). This

profuse number of modifications in the N-terminal tail, concomitant with the nucleosome partner, H2A.Z,

in which several acetylatable lysines are present, make this double variant nucleosome a likely feature for

chromatin modification relevant to gene expression, replication and/or DNA damage responses.

Overexpression of tagged wild type and mutant versions of H2B.Z and Knock out of endogenous *H2B.Z*

Attempts to obtain a mutant cell line by deletion of the *h2b.z* gene were unsuccessful (data not

shown). The failure to obtain an H2B.Z. knockout is consistent with its CRISPR fitness score of -4.05, strongly

suggesting it is essential [27] (ToxoDB, TGGT1_209910).

Given the high number of modifiable lysines in the H2B.Z variant unique to apicomplexan parasites,

we analyzed their role by using an over-expression strategy of myc-tagged wild-type or mutant forms. We

generated parasites over-expressing wild-type H2B.Z (c-Myc-WT-OE) or a mutated form in which all the

acetylatable lysines in the N-terminal tail were changed to alanine, which mimics how acetylation ablates the

positive charge of the lysine residues. We also over-expressed a mutated version in which the lysines were

replaced with arginines, which would generate a constitutive positive charge patch but no longer support

acetylation (Fig. 1B). Neither mutant form would be acetylated in this region, therefore would be incapable

of interacting with bromodomain containing proteins. Correct localization of each c-Myc-H2B.Z protein was

checked by immunofluorescence (IFA) (Fig. S1A) and by western blotting, observing two bands in every over-

expressing line (Fig. S1B). To note, c-Myc-WT-OE was already used in a ChIP seq screen, showing the same

genomic localization as endogenous H2B.Z when the experiments were performed using α-H2B.Z antibody

[17].

After that, endogenous *H2B.Z* gene was deleted by using CRISPR methodology in the previously

generated *T. gondii* lines overexpressing mutant H2B.Z versions (Fig 2A). We took advantage of the number

of mutations in the N-terminal tail of the tagged version to design sgRNA specific for endogenous *H2B.Z* gene.

We obtained *T. gondii* RH lines only expressing c-Myc-tagged H2B.Z with N-terminal mutations, which were

named c-Myc-A and c-Myc-R (Fig. 2A-C, Fig. S1C-D). Each form of c-Myc H2B.Z in these parasites localized

properly to the nucleus (Fig. 2B). In both CRISPR manipulated clones, a single band corresponding to c-Myc-

tagged H2B.Z can be observed by western blot with anti-H2B.Z and anti-c-Myc antibodies (Fig. 2C).

Quantification of the band corresponding to c-Myc H2B.Z relativized to Sag1 as a charge control, was 1.43 ±

0.77 and 1.65 ± 0.82 for c-Myc-R and c-Myc-A, respectively, indicating comparable levels of expression

between both mutants.

In vitro **growth analysis in c-Myc-R and c-Myc-A** *T. gondii* **lines**

We used a competitive growth assay to determine differences in parasite growth between parental

wild-type and mutant parasites (c-Myc-A and c-Myc-R) obtained, which could be distinguished by the c-Myc

tag. Approximately equal amounts of c-Myc positive and c-Myc negative (parental) tachyzoites were mixed

and allowed to invade hTert cells, which were analyzed by IFA on different days post-infection. While the c-

Myc-A parasites showed no change in growth relative to the parental line over time, the c-Myc-R line showed

a marked growth deficiency compared to the parental line (Fig. 3A). As arginines conserve the positive charge

but cannot be acetylated, these results indicate that the constitutive positive charge patch on the H2B.Z N-

terminal tail, but not the lack of acetylation, impairs tachyzoite growth *in vitro*. Therefore, wild-type *T. gondii*

must regulate the charge patch on H2B.Z N-terminal tail by PTMs such as acetylation for normal growth.

In vitro **differentiation analysis in c-Myc-R and c-Myc-A** *T. gondii* **lines**

We next studied the effect of the mutant H2B.Z variants on *in vitro* tachyzoite to bradyzoite

differentiation using alkaline stress to trigger stage conversion and *Dolichos biflorus* lectin (DBL) to monitor

tissue cyst wall formation. Strikingly, we found that c-Myc-R, but not c-Myc-A, showed an increase in DBL-

positive vacuoles compared to the wild-type control (Fig. 3B), suggesting that the impossibility to regulate

the presence of a positive charge patch in the H2B.Z N-terminal tail promotes an *in vitro* differentiation.

We also checked whether key stage-specific genes were altered in parasites expressing mutant

H2B.Z. We obtained RNA from freshly lysed tachyzoites, early bradyzoites (48 hours post-differentiation),

and late bradyzoites (96 hours post-differentiation) of the parental RH strain and c-Myc-A or c-Myc-R

parasites. We analyzed the expression of *Sag1* (tachyzoite), *Bag1* (early bradyzoite)*, LDH2* (early bradyzoite),

AP2IX-9 (mature bradyzoites) [41], *AP2 IV-3* (in which expression is maximal at 48h post- differentiation,

decreasing in mature bradyzoites) [42] and *BFD1* [14] mRNA by RT-qPCR. Actin and Tubulin were used as

housekeeping genes, and data were normalized to actin. The amount of expression relative to tachyzoite in

each parasite line is plotted, showing a trend of expression in c-Myc-R similar to what would be expected for

a cystogenic strain with the treatment, and different from RH expression profile (Fig. 3C). Also, the expression

of *BFD1* showed an increase in c-Myc-R, compared to RH (Fig. 3C). Notably, the expression profile of BFD1 in

RH type I strain has not been published to date. These results are consistent with the heightened bradyzoite

differentiation rate seen in c-Myc-R parasites.

Sensitivity to DNA damaging agents in c-Myc-A and c-Myc-R lines

In other species, H2A.Z has been linked to double strand break repair driven by the non-homologous

end joining (NHEJ) pathway [20, 43, 44]. Consequently, H2A.Z is associated as a responder to HU or MMS

genotoxic effects [44]. We have previously characterized the anti-*T.gondii* effect of MMS, HU, and

campothecin [45]. Topotecan is a genotoxic drug analog to campthotecin that produces fork collapse, and

ulterior DNA double strand break repaired by HRR [46].

Given the roles of histone variants in DNA repair, we analyzed c-Myc-A and c-Myc-R parasites for

sensitivity to topotecan at its IC_{50} value (20 μM) and MMS (50 μM) using the competitive growth assay. We

observed a significative decrease in growth compared to the parental after treatment only with MMS for c-

Myc-R tachyzoites (Fig. 4A). By contrast, topotecan did not produce a significant defect in growth, although

c-Myc-R seems to be more sensitive (Fig. 4A). In addition, treatment with MMS produced a disorganization

of vacuoles, rounded shapes and a loss of genetic material in both mutants as well as parental (Fig. 4B). *T.*

gondii normally presents a synchronic replication in the parasitophorous vacuole (PV). As a result, PVs usually

show 2, 4, 8, 16 and so on, tachyzoites (Tz) per PV. Treatment with MMS and Topotecan also caused a loss

of synchronization, with an altered number of Tz per PV, which was also counted as "anomalous vacuole".

Anomalous vacuoles and loss of genetic material were also observed with topotecan treatment (Fig. 4B).

However, quantification showed that anomalous PV are significatively more abundant in c-Myc-R than

parental and c-Myc-A, with both drugs tested (Fig. 4C). Taken together, the constitutive positive charge patch

generated in the c-Myc-R parasites alter the DNA damage response after MMS treatment, suggesting that

acetylation status of H2B.Z N-terminal tail is likely to be modulated for proper DNA repair.

***In vivo* analysis in c-Myc-R and c-Myc-A *T. gondii* lines**

Proper gene regulation is required for tachyzoite propagation in different tissue environments and

differentiation *in vivo*. Progression of tachyzoite infection of these mutants was analyzed in C57BL/6 mice by

inoculating 100 tachyzoites per *T. gondii* line. As expected, parental parasites displayed high virulence, killing

mice within 7 days (Fig. 5A). c-Myc-A tachyzoites were also lethal to mice, but with a delay in the time of

death (Fig. 5A). According to Mantel-Cox and Gehan-Breslow-Wilcoxon tests, the survival curves are

significantly different ($p > 0.0001$), suggesting that acetylation of at least some of the 5 N-terminal tail lysines

is relevant for parasite progression *in vivo*.

In contrast, c-Myc-R and Me49 infected mice, as well as non-infected (PBS group) survived until

sacrifice on day 30-35 (Fig. 5A). c-Myc-WT-OE was used a control that the tagged version is not responsible

per se of any affect observed. *T. gondii* line expressing both endogenous H2B.Z and c-Myc-H2B.Z-R (c-Myc-R-

OE) previous to CRISPR procedure, were as lethal as the parental line (data not shown), suggesting that the

randomly integrated ptub-*H2B.Z-R* gene is not responsible for the loss of virulence. Infection with 1,000 and

10,000 c-Myc-R tachyzoites were also nonlethal to mice (Fig S2A).

To confirm acute infection took place, hematoxylin and IFA staining of the peritoneal fluids at day 4

post-infection were performed (Fig. 5B). Intraperitoneal tachyzoites could be detected only with parental

and c-Myc-A *T. gondii* lines infections, but not with c-Myc-R parasites (Fig. 5B). Surviving animals (c-Myc-R

and Me49 *T. gondii* lines) were analyzed to detect antibodies against *T. gondii*. An increase in total IgG in

mice infected with c-Myc-R tachyzoites was observed, comparable to Me49, which indicated that an initial

infection had taken place (Fig. 5C, Fig. S2B). The analysis of IgG subtypes showed a Th1-like humoral response

in both cases, associated to IgG2a and higher for IgG2b (normal for C57/BL6 mice), typical for an intracellular

parasite infection (Fig. 5C, Fig. S2B). Following sacrifice, brains of surviving mice were processed for cyst

detection by direct observation with optical microscopy and DBL staining. Tissue cysts were observed in

brains of mice infected with Me49 strain, but not in those infected with c-Myc-R parasites (data not shown).

To determine if infection with c-Myc-R parasites elicits an adequate immune response, mice

previously infected with 1,000 c-Myc-R tachyzoites were reinfected with 100 tachyzoites of the highly virulent

RH parental line. c-Myc-R infected mice survived until sacrifice, while all mice in the control group died at day

38 (day 8 after re-infection) (Fig. 5D). These results suggest that c-Myc-R tachyzoites must have been capable

of invading cells and producing a stimulation of the cellular immune response adequate to control a challenge

infection.

H2B.Z acetylation status is involved in regulation of ROP proteins expression

The ability of the c-Myc-R line to establish a protective immune response suggest these parasites can

invade the host cells but are efficiently controlled by the immune system. Among the *T. gondii* virulence

proteins that exert protection from PV establishment are the polymorphic rhoptry proteins Rop5 and Rop18,

which disrupt the γIFN-inducible IGR pathway [47-51]. Here, we used antibodies against *T. gondii* Rop5

(TGGT1_411430) and Rop18 (TGGT1_205250) recombinant proteins for western blot analysis. Rop5 was

present in parental, c-Myc-A, and c-Myc-R parasites, whereas Rop18 only in the parental line (Fig. 6A).

However, quantification of the band intensities showed a significative decrease in Rop5 abundance in c-Myc-

R parasites and a lack of Rop18 protein in both c-Myc-A and c-Myc-R (Fig. 6B and Fig. S3A). To test whether

the absence of Rop18 was a consequence of the insertion of the H2B.Z construction in that locus, we

performed the same experiment using the over-expressing tachyzoites obtained prior to CRISPR

manipulation. As shown in Fig.S3 (B, C) both Rop5 and Rop18 are detectable in every clone similar to parental.

Co-localization analysis showed a correct localization of Rop5 in all *T. gondii* lines whereas Rop18 could only

be detected in the parental line (Fig. 6C). Of note, we also detected a decrease in the fluorescence intensity

of Rop5 in c-Myc-R parasites (Fig. 6D). RT-PCR analysis indicated that both genes were expressed in every *T.*

gondii line (Fig. S3D), suggesting that alteration of Rop5 and Rop18 expression in the c-Myc-R line is likely at

the post-transcriptional level.

Double variant nucleosome and transcription is not altered in *T. gondii* mutant lines in normal conditions

Since H2BZ is part of the double variant nucleosome with H2AZ, which is in the promoter region of

active genes, it was expected that expression alterations would occur at the transcriptional level rather than

post-trancriptionally. To determine if the observations for Rop5 and Rop18 were a specific situation, we

looked closer at the role of H2B.Z in transcription. We first analyzed the double variant (H2A.Z/H2B.Z)

formation in both mutant lines by co-immunoprecipitation (co-IP) assays by using commercial anti-c-Myc

agarose. Mononucleosomes of c-Myc-A and c-Myc-R lines were obtained after MNAse digestion before co-

IP (Fig. 7A). In figure 7B it can be observed that both c-Myc-A and c-Myc-R H2B.Z interact with H2A.Z,

confirming the assembly of the double variant nucleosome already found in *T. gondii*. We also decided to

study the interaction of the double variant nucleosome with acetylated H3, a known mark of active

chromatin. Both c-Myc-A and c-Myc-R H2B.Z interacted with this acetylated histone, which indicates that lack

of acetylation of H2B.Z does not alter this PTM crosstalk (Fig. 7B and Fig. S4). Quantification of the IP band

intensities relative to the respective input band showed that both *T. gondii* lines exhibit similar interaction

levels with H2A.Z and acetylated H3 (Fig. 7C). These data indicate that transcriptional expression was not

altered in c-Myc-R parasites with the double variant nucleosome remaining functional.

We next decided to explore if transcription was altered in tachyzoites of the mutant lines. We

analyzed expression of *Sag1* (tachyzoite), *tubulin* (constitutive), *Bag1* and *LDH2* (bradyzoite) and *BFD1*

(master differentiation factor). Only BFD1 expression was statistically higher in c-Myc-R that in the parental

or c-Myc-A (Fig. 7D), consistent with our observation that c-Myc-R parasites are more prone to differentiate

into bradyzoites.

Chromatin compaction does not explain differences observed in *T. gondii* mutant lines

The positive charge patch on H2B.Z could be associated with a more compact chromatin structure.

To analyze chromatin compaction, we measured the size of the nuclei of these lines compared to the parental

in IFAs stained with α-H2B.Z antibody. As shown in Fig. 7E, nuclei sizes are significantly smaller that the

parental in c-Myc-R parasites in accordance with the hypothesis. However, this significant difference is also

seen in c-Myc-A tachyzoites, indicating that the inability of this histone to be post-translationally modified is

enough to lead to this state. The sizes of c-Myc-WT-OE parasite nuclei were similar to parental nuclei,

suggesting that the c-Myc tag is not responsible for the effect observed (Fig. 7E). Taken together, acetylation

of the H2B.Z N-terminal tail is required for proper control of DNA compaction status.

H2B.Z interacting proteins

While results largely implicate the positive charge patch of H2B.Z N-terminal tail as responsible for the

observed phenotypes, a role for acetylation cannot be ruled out. Acetylation may explain why the c-Myc A

line showed a mild defect in virulence *in vivo* and defects in chromatin compaction. These defects could be

associated with the interaction with acetylation "reader" proteins. We therefore examined if there are

differential proteins interacting with H2B.Z N-terminal region depending on its acetylation state. We

designed two peptides, one containing two acetylated lysines (K14 and K18) and the other unacetylated;

both peptides contained a biotin tag towards the C-terminal end (Fig. 8). Peptide design was performed

according to Wysocka 2006 [52]. In this work, it was recommended to perform the synthesis of peptides

about 20 amino acids long, with biotin conjugated through a linker on the C-terminus for N-terminal histone

peptides, and with the modification positioned close to the center of the peptide. In addition, two

consecutive modifications were not recommended, explaining our choice of K14 and K18.

We used extracellular tachyzoites (RH) to perform the pull-down assays. After resolving on SDS-

PAGE, samples were analyzed by mass spectrometry. We detected a total of 48 interactors, 18 exclusive of

non-acetylated oligo, 17 exclusive of acetylated oligo and 14 present in both (Table S1). While many

interactors have nuclear localization, other localizations were detected, and many typical contaminants (e.g.

ribosomal subunits). Among the nuclear proteins, we can select a few that can be interacting with the H2B.Z

N-terminal tail independent of acetylation status; among them, GCN5-A, histone H4 and a PHD protein (Fig.

8). Interestingly, only three nuclear proteins were differentially pulled down at high score or with strong

evidence (2 independent pull downs), one of them TGME49_262620, a RRM protein (unacetylated peptide),

H3.3 and a putative CDK protein (both with the acetylated peptide). Considering the nuclear genes that only

appeared in 1 pull down, the acetylated peptide retrieved only genes that code for proteins associated with

the maintenance and segregation of chromosomes during the cell cycle, as peptidase c50 (TGME49_262825),

RecF/RecN/SMC N terminal domain-containing protein (TGME49_231170) and meiotic recombination

protein DMC1 family (TGME49_216400) (Table S1). Among the retrieved peptides in both acetylated and

non-acetylated pull-downs, we also detected acetyl-CoA carboxylase ACC1 (TGME49_221320) (Fig. 8 and

Table S1). It is interesting to observe that an apicoplast protein displayed a significant number of peptides,

indicative of a consistent interaction. We discuss the implications of these findings in the next section.

Discussion

In *T. gondii* gene expression is associated to the cell cycle, lytic cycle and differentiation [15, 53-58].

In addition to gene expression, the presence of histone variants as well as the turnover of PTMs on histones

are also associated to chromosomal organization, DNA repair or DNA replication [59]. The Apicomplexa

H2A.Z/H2B.Z double variant histone nucleosome is unique and is extensively modified by PTMs including N-

terminal tail acetylation. *T. gondii* epigenome has been studied, and it has been observed that this parasite

uses a set of different histone modifications as a complex set of tools to control its gene expression [60].

Acetylation and methylation are associated to opposite impacts in gene expression, the first favoring

expression, while the second is often a repressive mark. As an example, acetylation of lysine 31 on histone

H4 (H4K31) was associated to the promoter of a nearby active gene, while in the core body of these genes

H4K31me1 was detected [59]. Until now, the role of PTMs in *T. gondii* H2BZ has not been studied. However,

T. gondii H2B.Z/H2A.Z double variant nucleosomes were found in the promoter of active genes and in the

gene body of inactive genes [17]. This change in genomic location could be regulated by the different PTMs

of H2B.Z and/or H2A.Z. The present work is the first attempt to decipher the role of the N-terminal

acetylation of H2B.Z.

According to the results obtained in this work, the presence of a constitutive positive charge patch

in the N-terminal tail of the histone (c-Myc-R mutant) causes an increase in differentiation, a decrease in

growth, and an impaired DNA damage response *in vitro*. On the other hand, virulence was severely affected

in vivo indicating some sort of regulation at least in those processes for this H2B.Z PTM. Those effects were

not observed when lysine was replaced by alanine, an uncharged amino acid that mimics acetylated lysine,

except a mild effect on virulence *in vivo*. While these results argue in favor of control by the positive charge

patch as a crucial event in *T. gondii* tachyzoite, a minor or fine role supported by acetylation marks on the N-

terminal tail of H2B.Z cannot be ruled out. For *T. termophila* when engineering a constitutively positive

charge patch on H2A.Z, this was highly toxic, suggesting that its regulation has an essential function [61].

Moreover, the presence of a single acetylation site in this region was sufficient to avoid the toxic effect of

the positive charge patch [61, 62]. In our case, some phenotypical alterations were observed *in vitro*, but the

T. gondii c-Myc-R tachyzoites were viable except during mouse infection. This indicates that at least *in vitro*,

if there is a toxicity due to the constitutive positive charge patch of H2B.Z, it could be partly compensated by

H2A.Z, which presents 10 acetylatable lysines [28]. Together, H2A.Z/H2B.Z, would represent a "combined N-

terminal tail" of 15 lysines. Studies involving the role of acetylation on H2A.Z N-terminal tail of *T. gondii* are

needed to understand this.

An interesting aspect is that the *T. gondii* c-Myc-R line can grow responding to DNA damage *in vitro*,

with significant but surmountable defect. This would indicate a role in DNA damage repair for this histone

variant in *T.gondii*, with acetylation as a central player. However, this sensitivity was only significant for c-

Myc-R tachyzoites, arguing once again in favor of the positive charge patch toxicity hypothesis in this mutant.

Since H2B.Z does not exist in nature beyond Apicomplexa, we can only infer the putative role of this variant

histone being part of H2A.Z/H2B.Z nucleosome. The participation of H2A.Z in DNA damage repair pathways

has been studied in mammals and yeast [20, 43, 44]. More experiments should be conducted to analyze more

deeply the role of H2B.Z in DNA damage response.

However, the contributions of effects observed *in vitro* are less difficult to ascertain in the *in vivo*

model. Clearly c-Myc-R line is capable of invasion, generating an adequate immune response, but it is quickly

neutralized during the infective process. This suggests that the tachyzoite is exposed *in vivo* to stronger

environmental pressures than those observed *in vitro*. These pressures would include adaptation to new

environments, including different cell types and organs, which may present differences in metabolism,

nutrient availability, or immune responses. Under these conditions, where the tachyzoite must respond

quickly, fine reprograming of gene expression and modifications in metabolic pathways would be required

and the impossibility of modulating the positive charge patch of H2B.Z could hamper the ability of parasites

to adapt to changing environments *in vivo*.

A possible explanation for the changes provoked by the positive charges in the N-tail could be a

chromatin compaction due to interaction with the negative charges in DNA. Genome size expansion in

eukaryote evolution has led to an increase in histone H2A N-terminal tail arginines. Arginine-rich histones

bind more tightly to DNA, compacting chromatin and, in some cases, the nuclei are also smaller [63].

However, in this work we showed that nuclei size was smaller in both clones compared to parental and to c-

Myc-WT-OE tachyzoites, indicating that the charges in the N-terminal tail are not responsible for chromatin

compaction. Chromatin compaction has also been described to be regulated by linker histone H1, that

stabilizes the nucleosome structure. A candidate H1 has been recently identified in *T. gondii*, and interaction

of H1 with H2B.Z was observed [64]. The acetylation status of H2B.Z may be important for this interaction,

leading to a more compact chromatin in both alanine and arginine mutants because of linker histone H1.

Further experiments are needed to test this hypothesis.

It could be expected that the phenotypic alterations are given by defects in the formation of the

nucleosome with an impact on the expression of the genes. In our analysis, both situations are not altered.

Although a massive analysis of gene expression is necessary to detect specific changes, part of the study

shows that the expression of Rop5 and Rop18 does not occur at the transcriptional level, but rather post-

transcriptionally, affecting their translation. This regulation could be given by the presence of non-coding

RNAs (ncRNAs) and there is increasing evidence for a significant role for pos-transcriptional mechanisms [65,

66]. Future studies should take this possibility into account, considering a possible role of H2B.Z acetylation,

to analyze ncRNAs expression in *T. gondii*. With the results shown in this work it is only possible to correlate

the decrease in one of the more virulent isoforms of Rop5 together with the absence of Rop18 protein to the

loss of virulence of c-Myc-R mutant, although it cannot be ruled out that other virulence factors could be

altered in these parasites. On the other hand, the absence of Rop18 protein in c-Myc-A correlates with the

significative delay in mice death with c-Myc-A tachyzoites infection as reported by Resse et al [49].

Although we could not detect significative changes in most of the genes analyzed by RT-qPCR in

normal conditions for tachyzoites, *BFD1* was significatively higher in c-Myc-R. Interestingly, we observe a

higher differentiation rate *in vitro* for these parasites. The differentiation process of c-Myc-R under alkaline

stress was also accompanied by the expression of bradyzoite-specific genes. It is important to note that the

expression of the BFD1 gene is post-transcriptionally regulated [14]. Therefore, the increase in transcription

per se is not enough to induce differentiation, but the higher levels of BFD1 mRNA could favor differentiation

under alkaline stress. There is also another possibility of inducing differentiation based on epigenetic

regulation. HDACi FR235222 treatment was shown to favor acetylation of H4K31ac in bradyzoite gene

promoters, promoting transcription of bradyzoite genes and therefore, bradyzoite development [59]. The

fact that we only observe a mis-regulation in gene expression for c-Myc-R could be related to the constitutive

positive charge patch generated in this mutant. This could imply that the lack in modulation the positive

charges on H2B.Z could alter the correct role of this variant histone in the regulation of gene expression,

under alkaline stress.

The positive charge patch seems to be responsible for most of the differences shown in this work.

However, c-Myc-A mutant could be mimicking a situation where all H2B.Z N-terminal lysines were neutralized

by a PTM such as acetylation. This situation could be the most physiological one for the parasites, and why

c-Myc-A tachyzoites have a similar phenotype as the parental. The results obtained in the pull-down

experiments may be in consonance. While acetylated and unacetylated peptides pull-downs share 3 nuclear

proteins: GCN5A, PHD and clathrin, unacetylated peptide recognized one protein specifically, a RNA

recognition motif (RRM)-containing protein (TGME49_262620). This RRM-containing protein was also pulled

down with GCN5b acetyltransferase [55]. RRM-containing proteins are involved in post-transcriptional gene

expression processes, such as mRNA and rRNA processing, RNA export, translation, localization, stability, and

turnover [67]. In the case of acetylated peptide, it pulled down 2 specific proteins: a cyclin-dependent kinase

regulator (CDK_ TGME49_219832) and H3.3 histone variant (TGME49_218260). CDK is ortholog to a

Neospora caninum putative high mobility group protein (HMG_NCLIV_060770) and could be related to

transcription and cell cycle according to GO term analysis. CDKs have undergone an extraordinary degree of

evolutionary divergence and specialization and in yeast there is a number of different CDKs that

phosphorylate multiple substrates, thus regulating the cell-division cycle in response to different cellular cues

[68]. Also associated with the acetylated peptide H3.3 histone was found. H3.3 and H2A.Z are associated to

promoter regions flanking the nucleosome depleted region (NDR) of active genes [69]. Therefore, it may be

likely that acetylated H2A.Z and H2B.Z would associate to this variant histone in active promoters. However,

this association was not found in our genome wide analysis on tachyzoite stage [17]. Among the proteins

pulled down independent of the acetylation is the HAT GCN5A. GCN5A was shown to impact on H2A.Z

acetylation via crosstalk with H4 acetylation by means of bromodomain in p300 [70].   Future studies should

confirm the association of H2B.Z with GCN5A and stablish in this way a possible role of this histone in

regulation of gene expression.

In addition, CDKs have also been linked to mitosis [71, 72] and H3.3 is related to chromosome

segregation, nuclear structure, and the maintenance of genome integrity [73]. In this direction, it is

interesting that our pull-down assay has also retrieved, only for the acetylated peptide, other genes, all of

them associated to chromosome segregation (Table S1). Among them, peptidase c50 (TGME49_262825) a

separase or caspase-like protease. Separases are known to play a central role in the chromosome segregation

in yeast [74]. Structural maintenance of chromosomes (SMC) proteins -like RecF/RecN/SMC N terminal

domain-containing protein (TGME49_231170) found in our study- function together with other proteins in a

range of chromosomal transactions, including chromosome condensation, sister-chromatid cohesion,

recombination, DNA repair and epigenetic silencing of gene expression [75-77]. Finally, meiotic

recombination protein DMC1 family (TGME49_216400) was also retrieved in this pull down. This family of

proteins participate in meiotic recombination, specifically in homologous strand assimilation, which is

required for the resolution of meiotic double-strand breaks [78]. These findings are interesting, linking H2B.Z

and its N-tail acetylation to the mitotic and meiotic processes. Further studies are needed to shed light in

these subjects.

The decrease in acetyl-CoA in *T. gondii* leads to acetylation defects of several proteins including H2A.Z

and H2B.Z and could link histone PTM to nutritional stress [79]. An acetylation deficit could be a possible

physiological explanation for a positive charge patch in H2B.Z. Although the retrieval of acetyl-CoA

carboxylase ACC1 (TGME49_221320) by our peptides is tempting to speculate on the role of H2B.Z in sensing

this nutrient availability through lysine acetylation, the apicoplast localization makes it difficult to understand

this interaction. One possible explanation would be the presence of the histone in this organelle. The extra-

nuclear localization of histones was highly evidenced in other models [80-83]. Deficiencies in *Trypanosoma*

cruzi H2Bv histone variant amount present similar phenotypes as observed here, and some are also explained

in a putative extra-nuclear or extra-nucleosome function [84]. Interestingly, *T. gondii* H2B.Z peptides were

detected in extra-nuclear locations in ToxoDB: mitochondria, conoid and membrane fraction. To note, H2B.Z

abundance makes it the major H2B family histone [22], compared to H2A.Z variant.

It is relevant that these tachyzoites were able to generate a complete immune response, which only

happens when cells are invaded by tachyzoites and not with an infection with dead or lysed tachyzoites [85-

88]. In this sense, Toxovac® is a live attenuated vaccine used to prevent sheep congenital toxoplasmosis [89].

Recently, a Ca^{2+}-dependent protein kinase (CDPK) knock-out in Me49 type II strain was evaluated in mice as

a potential vaccine candidate [90]. The protection we observed in this work generated in mice facing an

infection with virulent RH tachyzoites is encouraging to continue with the proper assays needed for the

formulation of a vaccine.

Conclusion

In this work, we have shown that N-terminal tail lysine acetylation of histone H2B.Z is important for

the normal progression of *Toxoplasma gondii* life cycle. The presence of 5 positive charges in the N-terminal

of H2B.Z is most likely to be detrimental for the parasite. This toxicity is not very high, allowing the parasite

to replicate, but conferring a lower fitness to the tachyzoite. Moreover, this lower fitness is traduced in a

higher sensitivity to DNA damage, a mis-regulation of gene expression conducing to a higher differentiation

rate *in vitro*, and also a complete lack of virulence when facing the immune response in mice. All these effects

are not explained by an heterochromatinization of DNA due to the interaction with the positive charges,

because we observed that nuclei are also smaller in c-Myc-A tachyzoites. Besides, the effects cannot be

explained by a dislocalization of the histone in the genome because the nucleosome is still constituted

normally. However, it is quite unlikely to find a physiologically a situation where the histone would be

completely unacetylated, and c-Myc-A tachyzoites may reflect the normal condition observed *in vivo*.

Therefore, the similarity observed between c-Myc-A and parental tachyzoites in most experiments

performed in this work, would confirm that in normal conditions H2B.Z is acetylated in *T. gondii*.

The findings in this work are not exclusively explained by regulation of gene expression. This is not

surprising, as it is known that histone proteins may exert other functions, related for example to cell division.

Even extra-nuclear functions are attributed to histones in many models, and research in this area could bring

new insights for *T. gondii* as well.

Having a better understanding of how chromatin and gene expression are modulated in *T. gondii* will

allow inferring about chromatin modulation in other parasites of the phylum as well as the possibility of

suggesting a novel mechanism of epigenetic regulation in organisms with complex life cycles that are

evolutionarily distant from fungi and vertebrates.

470 Materials and methods

Parasite culture and manipulation

RHΔ*hxgprt* strain was used in all cases and grown in standard tachyzoite conditions *in vitro*: hTERT (ATCC®

CRL-4001, USA) monolayers were infected with tachyzoites and incubated with Dulbecco's modified Eagle

medium (DMEM, Invitrogen) supplemented with 1% fetal bovine serum (FBS, Internegocios S.A.,

Argentina) and penicillin (10,000 units/ml)-streptomycin (10 mg/ml) solution (Gibco, Argentina) at 37°C and

5% CO_2.

Cloning over-expression strategy

To generate the c-Myc-H2B.Z WT construct, the open reading frame (ORF) was amplified using the primers

F: 5'-<u>ATGCAT</u>TCAGGGAAAGGTCCGGCACAG-3' and R: 5'-<u>TTAATTAA</u>CTATGCACCAGAAGTCGTG-3'. For the

mutant   constructs,   reverse   primer   was   the   same,   and   forward   was:

<u>ATGCAT</u>TCAGGG**GCA**GGTCCGGCACAG**GCA**TCTCAGGCGGCG**GCGGCG**ACCGCCGGG**GCG**TCTCTGGGAC for c-

Myc-A,   <u>ATGCAT</u>TCAGGG**CGC**GGTCCGGCACAG**CGC**TCTCAGGCGGCG**CGCCGC**ACCGCCGGG**CGC**TCTCTGGGAC

for c-Myc-R, where lysine coding codons were replaced for alanine and arginine coding codons, respectively

(in bold type). NsiI and PacI restriction sites (underlined) were engineered into the 5' end of primers F and R,

respectively, to clone into *T. gondii* expression vector PTUB8mycGFPPfTailTy-HX (a kind gift of Dr. Dominique

Soldati, Université de Geneve) with a strong tubulin promoter. The resulting PCR product of the expected

size was removed using a Qiaex II Gel Extraction kit (Qiagen), cloned in pGEM-T vector (Promega) and

sequenced (Macrogen). The cloned ORF was digested with NsiI and PacI and ligated into the plasmid. The

final plasmids were electroporated into RH strain tachyzoites lacking hypoxanthine–xanthine guanine

phosphoribosyl transferase (RH∆*hxgprt*). Transgenic parasites were selected in mycophenolic acid (25 µg/ml,

Sigma) and xanthine (50 µg/ml, Sigma) and cloned by limiting dilution.

CRISPR/Cas9 gene deletion strategy

Single guide RNAs were designed directed to the N-terminal portion of the *h2bz* gene: F:

<u>AAGTT</u>GAGGCGGCGAAGAAGACCGCCG and R: <u>AAAAC</u>GGCGGTCTTCTTCGCCGCCTCA. Bsa I restriction sites

were included (underlined). Both oligos were annealed and cloned into the pU6 CRISPR Universal plasmid

(kindly provided by Lourido´s Lab, Whitehead Institute), by using the Bsa I restriction site. The plasmids were

confirmed by PCR, using the gRNA as primers, were sequenced (Macrogen) and prepared for transfection in

Toxoplasma. In order to obtain a clean KO, oligos were also designed to amplify the DHFR selection cassette

from the pUPRT_DHFR plasmid available in the laboratory, with homology regions (20 bp underlined) to the

3´and   5´UTR   of   the   *h2bz*   gene   (F:

<u>GAAAGGTCCGGCACAGAAATCTCAGGCGGCGAAGAAGACC</u>AAGCTTCGCCAGGCTGTAAATCC,   and   R:

GAATTCCCTCTCAACATCAAATTTCACCCCGTGGTTTATTCCTCATCCTGCAAGTGCATAGAAGGA).   The   PCR

amplification product was purified from agarose gel and used to co-transfect the parasites along with the

CRISPR plasmid. c-Myc-R and c-Myc-A tachyzoites (one clone for each one was selected) were transfected

and Pyrimethamine (1 μm, Sigma) was used for selection. Cas9 was detected in the nucleus of the parasites

after transfection by using α-Flag antibody (Sigma, F1804). Selection was continued until three passages and

cloning by limiting dilution was performed. Positive clones were selected by PCR with primers designed for

that   purpose: TgH2Bv-upUTR-F: GTTGTCATGCGCATTTGCATCAC;   TgH2Bv-downUTR-R:

GTGTGCACGCGTTATAATGAGCAC;   Val-TgH2Bv-R: CATGCTCTTCTTCGACACACCAG;   DHFRCX-F:

GTGGCATTTCACACAGTCTCACCTC; DHFRCX-R: AGACGCAGACGCATACAACGTTAG.

Immunofluorescence assay (IFA)

Intracellular tachyzoites grown in cover slips were fixed using cold methanol 100% for 8 min, washed with

PBS and blocked with 1% BSA. Primary antibody αrH2B.Z (rabbit), αrSag1 (mouse), αrHSP90 (rabbit), αrROP5

(rabbit), αrROP18 (rabbit) produced in the laboratory and αc-Myc (rabbit, abCam ab9106 or mouse, Santa

Cruz sc-42), αAcTubulin (mouse, Millipore MABT868) were incubated for 1 h at room temperature. After

several washes with PBS cover slips were incubated with secondary antibodies Alexa fluor goat anti-mouse

488 and anti-rabbit 594 (Invitrogen). DAPI was used to stain nuclei. Axio Imager.M2 Microscope Carl Zeiss

(Germany) with objective Plan-Apochromat 63x/1.40 Oil M27 and Video digital camera Zeiss 503

monochromatic 2.8 megapixeles, was used. Image J 1.53q (Fiji) software was used to process images.

Western blot analysis

Tachyzoites were collected, filtered, counted and lysed by 6 cycles of rapid freezing/defreeze in hypotonic

buffer, and boiled with LB for 5 minutes. 0.5 to 1×10^7 parasites were loaded per well and resolved by 15%

SDS-PAGE. Proteins were transferred to PVDF membrane for 1h at 100V. Western blot was then performed

as described [91]. The primary antibodies: αrH2B.Z [22], αrH2A.Z [23], αrROP5 (rabbit) and αrROP18 (rabbit)

were used at 1/5000, whereas α-c-Myc (rabbit, abCam ab9106) was used at 1/2000 and αrSag1 [88] 1/200

for 1 h at room temperature. αH3ac (rabbit, Millipore, 06-599B) 1/200, was incubated overnight. Appropriate

secondary antibodies were used: phosphatase alkaline-conjugated goat anti-mouse or anti-rabbit (Sigma)

along with the NBT and BCIP (Promega) detection system.

Phenotypical assays

For growth/competition assays, coverslips seeded with Htert cells were infected with a mix of 50% parental

(RHΔ*hxgprt*) and 50% of each clone. 3 coverslips per clone were infected with 0,05 to 0,1 tachyzoite per cell,

incubated 10 minutes on ice, and 2 hours at 37°C in incubator for invasion. Slides were washed twice with

PBS to remove tachyzoites that did not enter the cells, and fresh DMEM supplemented with 1% SFB was

added. Slides were fixed at different times, up to 96 hours and IFA was performed with α-c-Myc and α-Sag1

antibodies to distinguish between parental and c-Myc positive vacuoles. Number of c-Myc positive and total

vacuoles in at least 100 vacuoles, randomly choosing different fields of each slide were counted. In the case

of competition assays with genotoxic drugs treatment, the experiment was modified as follows: 3 slides per

clone mixture were fixed after 24 h, to stablish the initial percentage of c-Myc positive vacuoles. At this time

point, media was changed in the rest of the slides for DMEM supplemented with DMSO or drugs at the

concentrations indicated. These slides were fixed after 96 h, and IFA was performed and counted as explained

before.

For *in vitro* differentiation assays, in a similar way using Htert confluent slides, media was changed by DMEM

HEPES pH 8.1, after invasion with 1 parasite every 10 cells, accompanied with deprivation of CO_2. Media was

changed every day, and differentiation rate was estimated after 96 h by using *Dolichos Biflorus* Lectin

conjugated to Fluorescein (DLB 1:200, FL-1031; Vector Laboratories) to stain the wall of the cysts in formation

and α-Sag1 antibody to stain the tachyzoites. In the case of collection of parasites for RNA extraction,

differentiation assay was performed in a similar way, but in t25 dishes. After 48 or 96 h parasites were forced

out of the cells by 23G, 25G and 27G syringe passages.

For *in vivo* survival, virulence and differentiation assays, C57BL/6 female mice were used. For the use of

animals, C.I.C.U.A.E-UNSAM 10/22 was approved. Mice were maintained in optimal conditions in the

biotherium, with controlled temperature and free access to sterilized water and food. Infection was carried

out intra-peritoneal in 10 mice per group with 100, 1000 or 10000 tachyzoites of parental lines or clones,

depending on the assay. PBS was injected as negative control in five mice per experiment. Deceases were

registered along 30-35 days in each experiment, and signs of illness were monitored daily. Two mice from

each group (except PBS) were sacrificed at day 5 (when RH infected mice have symptoms of illness), and

intraperitoneal fluids were collected in order to detect acute *Toxoplasma* infection. Blood samples were

taken at days 0, 14, 21 and 35 and sera was frozen for ELISA assays. After sacrifice, surviving mice brains were

processed for cyst observance by optical microscopy. Also, a sample was stained using DLB and checked by

fluorescence microscopy.

In every phenotypical assay, three independent experiments were performed. One representative

experiment is shown.

RT-PCR and RT-qPCR

Tachyzoites or parasites exposed to 48 or 96 h differentiation stress as explained before were conserved in

TriZol (Invitrogen) solution at -80°C until use. All experiments were performed in three independent

replicates. RNA extraction was performed according to the manufacturer's instructions and cDNA was

obtained by means of MMLV reverse transcriptase (Promega) using oligo dT with the protocol provided with

the enzyme. For RT-PCR, primers used were: SAG1: Fw: TGAGAACCCGTGGCAGGGTAA; Rv:

GCTTTTTGACTCGGCTGGAA; TUB: Fw: ATGTTCCGTGGTCGCATGT; Rv: TGGGAATCCACTCAACGAAGT; ACTIN:

Fw: GGGCGGTTTCATGACCTAAA; Rv: ACGTATGATGCGCGAGAAAA; ROP5: Fw:

CTAGCTAGCATGGCGACGAAGCTCG;   Rv: CCCAAGCTTTCAAGCGACTGAGGGCGCA;   ROP18:   Fw:

CTAGCTAGCATGTTTTCGGTACAGCGGCCA; Rv: GCCAAGCTTTTATTCTGTGTGGAGATGTTC.

For RT-qPCR primers used were: SAG1: Fw: TGAGAACCCGTGGCAGGGTAA; Rv: GCTTTTTGACTCGGCTGGAA;

BAG1:   Fw:CAACGGAGCCATCGTTATCAAAGG;   Rv: TAGAACGCCGTTGTCCATTG;   LDH2:

Fw: ACAATGGCCCAGGCATTCT;   Rv: CAATAAACATATCGTGAAGCCCATA;   AP2IX-9:

Fw: GGGCGTTCTCAGCGTTCACT;   Rv: GGCGCGTCTCATCTGTTTCA;   AP2IV-3:

Fw: GAGCCCATTGACCCCATGAA;   Rv: GGCTTCGCTTCTTTCCGTGA;   BFD1:   Fw:

ATGTCGGGAACGATGGTTTA;   Rv: TCTTCACGGCATTCTCTGTG;   TUB:

Fw: ATGTTCCGTGGTCGCATGT; Rv: TGGGAATCCACTCAACGAAGT; ACTIN: Fw: GGGCGGTTTCATGACCTAAA; Rv:

ACGTATGATGCGCGAGAAAA.

The primers were first assayed for efficiency using a pooled cDNA in 1 to 0.001 dilutions, and the threshold

was defined for each set of primers. Melt curves were also obtained for each set. For each set of experiments,

SybrGreen master solution (Roche) was used and qPCR was run in StepOne Real time equipment (Applied

Biosystems). Actin and tubulin were used as housekeeping genes and data was normalized to those genes

by Infostat software. For the experiments in figure 3, all sets of data were relativized to tachyzoite

amplification. For experiment shown in figure 7, data from the different clones was relativized to the

parental.

ELISA assay

Blood samples were extracted from surviving mice at days 0 (Pi), 14, 21 and 35. Sera was frozen until

processing for ELISA. TLA (Toxoplasma lysate antigen, obtained from fresh RH tachyzoites) in buffer

Carbonate (7,13g/L NaHCO3 + 1,59 g/L Na2CO3 pH 9,5) was used to immobilize 96 well plaques (Nunc

Immuno™ MicroWell™ 96 well solid plates, Sigma), o.n. at 4°C. After bocking with 5% non-fat milk in PBS-T

they were incubated with sera diluted 1:100 in the same solution. Total IgG (IgGt) and subtypes (IgG1, IgG2a

and IgG2b) HRP-conjugated anti-mouse, made in rat, were used 1:5000. Tetramethyl-bencidine (TMB,

Invitrogen) was used to reveal, and plaques were read after 20 min reaction at 655 nm in a Synergy H1. Data

was analyzed in GraphPad Prism v7. Sera from mice inoculated with PBS (negative control) or infected with

Me49 (positive control) were assayed.

Co-Immunoprecipitation (Co-IP)

Approximately 5×10^8 RHΔ*hxgprt* tachyzoites were used for each immunoprecipitation. Tachyzoites were

treated with micrococcal endonuclease (MNase) in order to obtain mononucleosomes as described [23] and

Co-IP was performed as described in Dalmasso et al [23] with minor modifications. Protease inhibitor cocktail

(Sigma) was added in every step. Mononucleosomes were incubated with Agarose c-Myc beads (abCam,

ab1253) overnight at 4°C. Immunocomplexes were washed twice with washing buffer 1 (50 mM Tris, pH8,

200 mM NaCl and 0.05 % Igepal100), twice with washing buffer 2 (50 mM Tris, pH8, 300 mM NaCl and 0.05

% Igepal100), and twice with buffer TE (10 mM Tris, pH8, 1 mM EDTA); then resuspended in 60 µl of SDS-

PAGE loading buffer. Samples were boiled for 5 min and 20 µl were loaded per well in a 15% SDS-PAGE gel

for immunoblotting. The absence of contaminating proteins was corroborated by Western blot with murine

anti-SAG1antibody [88]. Quantification of the bands was done with Image J 1.53q (Fiji) software.

Tachyzoites nuclei size assay

Three slides with confluent hTert cells were infected as explained before with parental or c-Myc-A/c-Myc-

R/c-Myc-WT-OE clones and let replicate for 24 hours. Slides were fixed and IFA was performed as explained

before using αrH2B.Z to stain nuclei. Images were processed with Image J 1.53q (Fiji) software to measure

the size of the nuclei, by delimiting them with the appropriate tool in at least 50 vacuoles in fields randomly

chosen, for each slide and each independent experiment. Data was graphed and statistical analyzed with

GraphPad Prism 8 software.

Peptide synthesis and characterization

Peptides were synthesized by solid-phase multiple peptide system using tea-bags with Fmoc/tBu strategy

according Guzmán [92], briefly: Rink amide resin (0.55 meq/g) was used as solid support and Fmoc amino acids,

including acetylated lysine (Iris Biotech). Peptides were couplet to biotin for their detection The peptide cleavage

was performed with a solution of Trifluoracetic acid/triisopropylsilane/1.2-ethandithiol /water (92.5 /2.5

/2.5/2.5), washed with cold ether dried and lyophilized. Peptides were characterized by HPLC and electrospray

ionization mass spectrometry, and purified by C18 cartridges (United Chemical Technologies, Bristol, PA, USA)

before use.

Peptides used in the pull-down assays were: Un-acetylated: GPAQKSQAAKKTAGKSLGPRK(Biotin)

Acetylated: GPAQKSQAAK(Ac)K(Ac)TAGKSLGPR-K(Biotin)

Pull-down assays

Pull-down experiments were performed as detailed in Wysocka [52] after nuclear extraction with NE-PER

commercial kit following manufacturer´s instructions (Thermo Scientific #78833) and run in SDS-PAGE for a

short time in order to avoid separation. Protein gels were fixed by soaking for 3 hr in 30% methanol, 2%

phosphoric acid. Then the gel was washed with deionized water 3 times, 5 minutes each. After removing the

deionized water, the staining solution (0.5 g/L Coomasie blue brilliant R250, 18% methanol, 17% $(NH_4)2SO_4$,

2% phosphoric acid) was added until covering the gel. It was stained for 1 h in gentle shaking. Then the gel

was rinsed in deionized water 3 times for 5 minutes each. The background staining was removed with 30%

methanol. The gel was stored in deionized water until use. Each lane of the gel was cut into individual slices.

Each band was then cut into 1 mm^3 cube and further treated with three washes of 50 mM NH_4HCO_3 in 50%

CH_3CN with 10 min incubations. Each group of gel cubes was then dehydrated in CH_3CN for 10 min and dried

in a Speed Vac. Protein samples were reduced by dithiothreitol (DTT) and alkylated by iodoacetamide [93]. A

solution of 10 ng/µL trypsin in 50 mM NH_4HCO_3 was used to re-swell the gel pieces completely at 4°C for 30

min, followed by a 37°C digestion overnight. A small amount of 10% formic acid was then added to stop the

digestion. The sample was then centrifuged at 2,800 x g, and the supernatant was collected for LC-MS/MS.

LC-MS/MS Analysis

A fused silica microcapillary LC column (15-cm long x 75-μm inside diameter) packed with Halo C18 reversed-

phase resin (2.7 μm particle size, 90 nm pore size, MichromBioresources.) was used with EASY-nLC 1200

system (Thermo Fisher). The nanospray ESI was fitted onto the Thermo Q-Exactive plus mass spectrometer

(Thermo Electron, San Jose, CA) that was operated in a Higher-energy C-trap dissociation mode to obtain

both MS and tandem MS (MS/MS) spectra. Two μL of tryptic peptide samples were loaded onto the

microcapillary column and separated by applying a gradient of 3-40% acetonitrile in 0.1% formic acid at a

flow rate of 300 nL/min for 80 min. Mass spectrometry data were acquired in a data-dependent top-10

acquisition mode, which uses a full MS scan from m/z 350-1700 at 70,000 resolution (automatic gain control

[AGC] target, 1e6; maximum ion time [max IT], 100 ms; profile mode). Resolution for dd-MS2 spectra was set

to 17,500 (AGC target: 1e5) with a maximum ion injection time of 50 ms. The normalized collision energy

was 27 eV.

Protein Identification

Obtained MS spectra were searched against the *T. gondii* ToxoDB-42 _TgondiiME49 protein database using

Proteome Discoverer 2.2 (Thermo Electron, San Jose, CA). The search parameters permitted a 10 ppm

peptide MS tolerance and a 0.02 Da MS/MS tolerance. Carboxymethylation of cysteines was set as a fixed

modification, and oxidation of methionine as a dynamic modification. Up to two missed tryptic peptide

cleavages were considered. The proteins for which the False Discovery Rate was less than 1% at the peptide

level were included in the following analysis. The raw data of all mass spectra had been submitted to the

MassIVE (https://massive.ucsd.edu/; Project accession: MassIVE MSV000091226).

Author Contributions: L.V. participated in all the experiments and analyses and wrote the first draft; D.M.

performed Crispr/Cas9 and some of the characterization experiments along with the pull-down experiments

with L.V. supervision; C.C. performed the DNA damage experiments and some of the transfections; A.G. was

involved in many of the experiments with technical support, and was in charge of RNA and cDNA preparation;

R.N. performed RT-PCR experiments and ELISAs with LV supervision; M.C.B. performed Co-IP experiments with

L.V. supervison; V.T. perfomed Rop5 and Rop18 antibodies and provided the respective primers for RT-PCR; F.G.

synthesized biotynilated peptides for pull down experiments; B.D. performed mass spectrometry analysis; K.K.

and W.J.S. participated in design of the study and manuscript final corrections; S.O.A. designed the study and

participated in its supervision. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Ministerio Nacional de Ciencia y Tecnología (MINCyT): PICT 2015 1288

(S.O.A.), PICT 2018 2434 (L.V.), Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET): PIP

11220150100145CO and 11220210100572CO (S.O.A., L.V.) and by National Institute of Health: NIH-NIAID

1R01AI129807 (S.O.A.) and AI152583 (to W.J.S.).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data that support the findings of this study are available from the

corresponding author upon reasonable request.

Acknowledgments: L. Vanagas (Researcher), D. Muñoz (Fellow), C. Cristaldi (Fellow), D. Ganuza (Technician), V.

Turowski (Researcher), and S.O. Angel (Researcher) are members of CONICET. S.O. Angel (Full) and L. Vanagas

(Adjunct) are Professors at Universidad Nacional General de San Martin (UNSAM). The Vermont Biomedical

Research Network (VBRN) Proteomics Facility is supported through NIH grant P20GM103449 from the INBRE

Program of the National Institute of General Medical Science (BD).

Conflicts of interest: The authors declare that there are no conflict of interest.

Figure 1. H2B.Z N-tail acetylation and mutagenesis strategy. A. Representative Apicomplexa H2B.Z N-tail

alignment. *Toxoplasma gondii* (TGME49_209910), *Neospora caninum* (NCLIV_004160), *Eimeria tenella*

(ETH_00030960), *Plasmodium falciparum* (PF3D7_0714000.1), *Cryptosporidium hominis* (cgd7_1700-RA),

Babesia bovis (BBOV_IV006840) and *Theileria equi* (BEWA_028110) H2B.Z protein sequences were aligned

by Clustal W. The N-tail region is shown. **B.** Scheme of the constructions generated in RHΔ*hxgprt* strain

tachyzoites. The N-terminal region is shown with the aminoacidic sequence, with or without mutations. All

constructions were performed associated to a tubulin promoter, in frame with a c-Myc tag in the N-

terminal.

Figure 2. H2B.Z gene deletion strategy. **A.** CRISPR/Cas9 deletion of endogenous *h2b.z,* **s**trategy. Single guide

RNA was directed to the N-terminal region of H2B.Z. Tachyzoites of one clone c-Myc-R-OE and c-Myc-A-OE

were co-transfected with Crispr plasmid and a PCR product obtained with primers to amplify DHFR selection

cassette from pUPRT_DHFR plasmid with homology arms (20 pb) to 5´and 3´UTR of *h2b.z*. The scheme shows

the result of selection with Pyrimethamine and cloning, where parasites obtained replaced *h2b.z* by DHFR

cassette, and c-Myc-H2B.Z construction (with the corresponding mutations) remained. gDNA: genomic DNA;

Pr$_{end}$: endogenous promoter; Pr$_{tub}$: tubulin promoter. **B.** Immunofluorescence assay: coverslips seeded with

Htert cells at confluency were infected with 1 tachyzoite per cell of each obtained clone and fixed after 18-

24 hours. IFA was performed with α-cMyc (abCam), and α-Sag1 (mouse). α-IgG-mouse Alexa 488, and α-IgG

rabbit Alexa 564 were used 1:4000 for 30 minutes. DAPI was used to stain nuclei. Scale bar=5 μm. **C.** Western

blot assay. Clones confirmed by PCR were grown and lysed to run 0.5-1 x10^7 parasites per lane, transferred

to PVDF and assayed with indicated antibodies: α-c-Myc, α-H2B.Z, secondary α-rabbit IgG (negative control).

Over-expressing clones (c-Myc-R-OE and c-Myc-A-OE) used for transfection, as well as the parental (RH) were

loaded as controls. Mw: molecular weight marker (17 kDa band). eH2B.Z: endogenous H2B.Z.

Figure 3. *In vitro* growth and differentiation analysis on c-Myc-A and c-Myc-R. **A.** Growth assay, performed

as explained in Materials and Methods. The percentage of c-Myc positive vacuoles is expressed as relative to

the total (α-Sag1), in at least 100 vacuoles per slide. Three slides per clone were counted. Representative

assay of three independent experiments. **B.** *In vitro* differentiation assay was performed as explained.

Percentage of DLB positive vacuoles was obtained after IFA with DLB and α-Sag1 to detect tachyzoites. PAR:

parental (RHΔ*hxgprt)*. Data was analyzed with Graphpad, and one-way Anova was performed. **: p< 0.01. **C.**

Tachyzoites (Tz) of parental (RHΔ*hxgprt*); c-Myc-A or c-Myc-R were freshly collected and conserved in Trizol

until processing. *In-vitro* differentiation was performed in dishes for 48 or 96 hours, and parasites were

collected and conserved in Trizol. RNA was extracted from each sample and cDNA was obtained by reverse

transcription with MMLV as explained in Materials and Methods. RT-qPCR was performed in Real time PCR

equipment (Applied Biosystems) with Sybr Green reagent with sets of primers indicated in each graph, using

tubulin and actin as housekeeping controls. Data was normalized to each of the housekeeping, and relative

quantities to tachyzoite in each sample is plotted for each of the genes studied. GraphPad Prism was used to

statisticaly analyze data, by two-way Anova. ***: p< 0.0001; **: p< 0.01; *: p< 0.05; ns: not significative.

Figure 4. H2B.Z acetylation status is important for DNA damage sensitivity. A. Competition assay after DNA damage induced by Topotecan (20 μM) or MMS (50 μM). Competition assay was performed as explained, and IFA with anti-Sag1 for total vacuoles count and anti-cMyc for clones count was performed in three independent experiments in triplicate. Graph shown corresponds to one representative assay. Initial: % of c-Myc vacuoles of the total after 24 hours of infection with the mixture. Final: % of c-Myc vacuoles of the total after 72 hours of treatment with each drug or DMSO as control. c-Myc-A/Par: mixture of c-Myc-A with parental (RHΔ*hxgprt*) tachyzoites; c-Myc-R/Par: mixture of c-Myc-R with parental (RHΔ*hxgprt*) tachyzoites. GraphPad Prism 8 statistical ANOVA analysis, multiple comparison where DMSO final is compared to initial in each case and drugs data is compared to DMSO: a: ns; b: p<0.05; c: p<0.01. **B.** Vacuoles affected by drugs.

Representative images of most affected parasites (c-Myc-R) with both drug treatments. DMSO images are

included as control. Yellow arrowheads point to parasites with loss of genetic material. **C.** Quantification of

anomalous vacuoles. Anomalous and normal vacuoles were counted in at least 50 fields randomly chosen,

per clone, experiment and drug treatment. Results were plotted in GraphPad Prism 8 and statistical analysis

by Multiple comparisions was performed. a: ns; b: p<0.05; c: p<0.001.

Figure 5. c-Myc-R tachyzoites are not virulent i*n vivo* and elicit a protective immune response. A. Survival

assay. 10 mice C57BL/6 were intra-peritoneal infected with either 100 tachyzoites of parental (RHΔ*hxgprt*),

and one clone of each KO line: c-Myc-R and c-Myc-A. As controls we also infected mice with c-Myc-WT-OE,

vehicle (PBS) and with Me49 tachyzoites (type II). The survival curve shown corresponds to one assay,

representative of three independent experiments. **B.** Acute infection. At day 5, 2 mice of each group were

sacrificed, and intra-peritoneal liquid was extracted and analyzed in the microscope, either by hematoxylin

staining (upper panel), or by IFA with α-HSP90 antibody (lower panel). Arrowheads in the upper panel show

tachyzoites. **C.** Evaluation of immune response by ELISA. Blood samples were extracted from surviving mice

at days 0 (Pi), 14, 21 and 35. Pooled sera from mice inoculated with PBS (negative control) or infected with

Me49 (positive control) were assayed. Total IgG and subtypes (IgG2a, IgG2b and IgG1) were analyzed. In the

graph only day 35 is shown. **D.** Re-infection assay. The indicated number of mice were infected with 1000

parental (Par) or c-Myc-R tachyzoites or inoculated with PBS. After 30 days, surviving mice were re-infected

with 100 parental RH tachyzoites. The survival curve shown corresponds to one assay, representative of three

independent experiments.

Figure 6. Expression of ROP proteins is affected by H2B.Z acetylation. A. Western blot identification of ROP5

and ROP18. The image is representative of three independent experiments, in which tachyzoite lysates of

parental (RHΔ*hxgprt*), c-Myc-A or c-Myc-R were run in SDS-PAGE, transferred to PVDF membrane and

assayed with anti Sag1 (charge control), anti Rop5 and anti-Rop18 antibodies. **B.** Quantification of band

intensities, relative to Sag1 band intensity in each lane. Image J software was employed to quantify relative

intensities and graphed with GraphPad Prism 8. Average plus SD of three independent assays is represented.

*: $p<0.05$. **C.** Immunofluorescence assay. Htert confluent slides were infected with tachyzoites of parental

(RHΔ*hxgprt*), c-Myc-A and c-Myc-R and fixed after 24 h for immunofluorescence analysis using anti Rop5 and

anti Rop18 antibodies (green). AcTubulin in red, was used to stain parasites and DAPI for the nuclei. **D.** Rop5

intensity quantification. Image J software was used to quantify the antibodies intensities in three

independent experiments, by triplicate, at least 10 vacuoles per slide. The graph shows the average intensity

in relative units; statistical analysis was performed by GraphPad Prism 8 software. *: $p<0.05$.

Figure 7. Nucleosome composition and chromatin structure is maintained in c-Myc-A and c-Myc-R. A. Mnase digestion of tachyzoites for co-immunoprecipitation analysis. Tachyzoites were MNase digested as explained in Materials and Methods, and an aliquot of the total tachyzoites and the digestion was precipitated and run in 1% agarose gel. A representative image of digestion is shown. **B.** Co-immunoprecipitation-WB analysis. Assays were performed in triplicate as explained in Materials and

Methods. A representative image of Western-blot analysis is shown for c-Myc-A and c-Myc-R. IP:

immunoprecipitation; IN: input. H3ac: western blot revealed with commercial anti H3acetylated antibody.

H2A.Z: western blot revealed with anti H2A.Z antibody performed in the lab. Sag1: western blot revealed

with anti Sag1 antibody. Color key: Orange for c-Myc-A; lilac for c-Myc-R. **C.** Quantification of western blot

band intensities. Image J software was used to quantificate intensities of H3ac and H2A.Z IP bands and

relativized to respective IN band intensities. Graph shows the relative intensities of three independent

experiments as the average plus SD for each antibody and clone, with the same key color as before. Graphpad

Prism 8 statistical analysis showed no significative differences between clones for both H3ac and H2A.Z

quantification. **D.** Tachyzoites gene expression. Tachyzoites (Tz) of parental (RHΔ*hxgprt*); c-Myc-A or c-Myc-

R were freshly collected in triplicate and conserved in Trizol until processing. RNA was extracted from each

sample and cDNA was obtained by reverse transcription with MMLV as explained in Materials and Methods.

RT-qPCR was performed in Real time PCR equipment (Applied Biosystems) with Sybr Green reagent with sets

of primers indicated in the graph, using actin as housekeeping control. Data was normalized to actin, and

relative quantities to RH tachyzoites in each sample is plotted for each of the genes studied. GraphPad Prism

was used to statisticaly analyze data, by two-way Anova. ***: $p < 0.0001$; ns: not significative. **E.** Tachyzoite

nuclei size. Tachyzoites of parental (RHΔ*hxgprt*), c-Myc-A, c-Myc-R or c-Myc-WT-OE were allowed to invade

hTert confluent slides and replicate for 20 h, and IFA was performed with α-H2B.Z antibody to detect the

nuclei. Image J software was used to measure the average nuclei size of at least 100 tachyzoites per slide, in

three independent experiments, and this was plotted and analyzed with Graphpad Prism 8, by two-way

ANOVA. ****: $p < 0.0001$.

Figure 8. Identification of H2B.Z interacting proteins by pull-down assay. Scheme of the two different

peptides synthesized is graphed above. Ac: acetylation. Most representative peptides retrieved in two

independent assays for each peptide. Proteins are separated according to localization in nucleus or

cytosol/organelles.

Supplementary figures

Figure S1

Fig. S1. Controls for tachyzoite lines over-expressing H2B.Z WT or mutants and CRISPR/Cas9 deletion of

endogenous H2B.Z gene. A. Immunofluorescence assay. Htert confluent slides were infected with the

different clones obtained and immunofluorescence was performed with anti c-Myc (green) and anti-H2B.Z

(red); DAPI was used to stain nuclei. Images show a representative IFA of one clone of each over-expressing

line obtained. **B.** Western-blot assay. One clone of each line and the parental were lysed and run in SDS-PAGE

for WB assay with anti-H2B.Z, anti-c-Myc and anti-Sag1 (charge control). **C.** Immunofluorescence after

transfection showing flag-CAS9 signal in the nucleus of one vacuole (green arrow). **B.** PCR selection of positive

clones. The scheme shows the primers designed for PCR detection of positive and negative clones. The image

corresponds to a representative PCR where a positive and a negative clone with the parental as control are

shown. PCR1-5: primers combinations for PCR shown in the scheme.

Figure S2

Fig. S2. c-Myc-R tachyzoites up to 10,000 are unable to kill mice but immune response is triggered. A.

Survival assay. 10 mice C57BL/6 were intra-peritoneal infected with either 1000 or 10000 tachyzoites of Me49

or c-Myc-R tachyzoites. Vehicle (PBS) was inoculated in 5 mice as control. The survival curve shown

corresponds to one assay, representative of three independent experiments. **B.** Evaluation of immune

response by ELISA. Assay shown in Fig. 5C, now showing data for 14, 21 and 35 days samples from mice

infeceted with c-Myc-R tachyzoites. Total IgG and subtypes (IgG2a, IgG2b and IgG1) were analyzed.

Figure S3

Fig. S3. Western blot analysis of ROP proteins. A. Complete image of Western blot shown in Fig. 5. **B.**

Representative Western-blot of over-expressing lines. Antibodies against Rop5, Rop18 and Sag1 were used.

WT: c-Myc-WT-OE; A: c-Myc-A-OE; R: c-Myc-R-OE. The upper bands detected with αRop18 are of higher

molecular weight than Rop18 full length protein, being unspecific. However, the intensities are similar to

Sag1, being a second charge control. **C.** Quantification of Rop5 and Rop18 bands, relativized to Sag1 band

intensity in each lane, in two independent WB experiments with the OE lines, and statistical analysis by

GraphPad Prism 8. ns: not significative. **D.** RT-PCR. Tachyzoites of parental (RHΔ*hxgprt*), c-Myc-A and c-Myc-

R were collected by triplicate and conserved in TriZol until RNA extraction, and cDNA preparation. PCR was

run with primers for the genes indicated and ran in agarose gels. Image is representative of three

independent experiments.

Figure S4

Fig. S4. Co-immunoprecipitation assay. A. Complete image of the Pounceau (left) and WB (right) shown in

Fig. 7 for c-Myc-A. **B.** Complete image of the Pounceau (left) and WB (right) shown in Fig. 7 for c-Myc-R. For

both images in lanes 1-11 was run: IN-M-IP-IP-space-IN-IN-T-NU-space-IP. IN: input; IP: immunoprecipitation;

T: total; NU: after immunoprecipitation, not bound to agarose beads. MW markers are indicated at the left.

References

[1] Vanagas L, Jeffers V, Bogado SS, Dalmasso MC, Sullivan WJ, Jr., Angel SO. Toxoplasma histone acetylation remodelers as novel drug targets. Expert review of anti-infective therapy. 2012;10:1189-201.

[2] Mouveaux T, Rotili D, Boissavy T, Roger E, Pierrot C, Mai A, et al. A potent HDAC inhibitor blocks Toxoplasma gondii tachyzoite growth and profoundly disrupts parasite gene expression. Int J Antimicrob Agents. 2022;59:106526.

[3] Kim K. The Epigenome, Cell Cycle, and Development in Toxoplasma. Annual review of microbiology. 2018;72:479-99.

[4] Pappas G, Roussos N, Falagas ME. Toxoplasmosis snapshots: global status of Toxoplasma gondii seroprevalence and implications for pregnancy and congenital toxoplasmosis. Int J Parasitol. 2009;39:1385-94.

[5] Attias M, Teixeira DE, Benchimol M, Vommaro RC, Crepaldi PH, De Souza W. The life-cycle of Toxoplasma gondii reviewed using animations. Parasit Vectors. 2020;13:588.

[6] Weiss LM, Kim K. The development and biology of bradyzoites of Toxoplasma gondii. Front Biosci. 2000;5:D391-405.

[7] Luft BJ, Remington JS. Toxoplasmic encephalitis in AIDS. Clin Infect Dis. 1992;15:211-22.

[8] Carlier Y, Truyens C, Deloron P, Peyron F. Congenital parasitic infections: a review. Acta Trop. 2012;121:55-70.

[9] Brynska A, Tomaszewicz-Libudzic E, Wolanczyk T. Obsessive-compulsive disorder and acquired toxoplasmosis in two children. Eur Child Adolesc Psychiatry. 2001;10:200-4.

[10] Miman O, Mutlu EA, Ozcan O, Atambay M, Karlidag R, Unal S. Is there any role of Toxoplasma gondii in the etiology of obsessive-compulsive disorder? Psychiatry Res. 2010;177:263-5.

[11] Yolken RH, Dickerson FB, Fuller Torrey E. Toxoplasma and schizophrenia. Parasite Immunol. 2009;31:706-15.

[12] Vittecoq M, Elguero E, Lafferty KD, Roche B, Brodeur J, Gauthier-Clerc M, et al. Brain cancer mortality rates increase with Toxoplasma gondii seroprevalence in France. Infect Genet Evol. 2012;12:496-8.

[13] de Haan L, Sutterland AL, Schotborgh JV, Schirmbeck F, de Haan L. Association of Toxoplasma gondii Seropositivity With Cognitive Function in Healthy People: A Systematic Review and Meta-analysis. JAMA Psychiatry. 2021;78:1103-12.

[14] Waldman BS, Schwarz D, Wadsworth MH, 2nd, Saeij JP, Shalek AK, Lourido S. Identification of a Master Regulator of Differentiation in Toxoplasma. Cell. 2020;180:359-72 e16.

[15] Jeffers V, Tampaki Z, Kim K, Sullivan WJ, Jr. A latent ability to persist: differentiation in Toxoplasma gondii. Cellular and molecular life sciences : CMLS. 2018;75:2355-73.

[16] Bougdour A, Maubon D, Baldacci P, Ortet P, Bastien O, Bouillon A, et al. Drug inhibition of HDAC3 and epigenetic control of differentiation in Apicomplexa parasites. The Journal of experimental medicine. 2009;206:953-66.

[17] Nardelli SC, Silmon de Monerri NC, Vanagas L, Wang X, Tampaki Z, Sullivan WJ, Jr., et al. Genome-wide localization of histone variants in Toxoplasma gondii implicates variant exchange in stage-specific gene expression. BMC genomics. 2022;23:128.

[18] Santisteban MS, Kalashnikova T, Smith MM. Histone H2A.Z regulats transcription and is partially redundant with nucleosome remodeling complexes. Cell. 2000;103:411-22.

[19] Oberdoerffer P, Miller KM. Histone H2A variants: Diversifying chromatin to ensure genome integrity. Semin Cell Dev Biol. 2022.

[20] Xu Y, Ayrapetov MK, Xu C, Gursoy-Yuzugullu O, Hu Y, Price BD. Histone H2A.Z controls a critical
chromatin remodeling step required for DNA double-strand break repair. Mol Cell. 2012;48:723-33.
[21] Fenoy IM, Bogado SS, Contreras SM, Gottifredi V, Angel SO. The Knowns Unknowns: Exploring
the Homologous Recombination Repair Pathway in Toxoplasma gondii. Front Microbiol. 2016;7:627.
[22] Dalmasso MC, Echeverria PC, Zappia MP, Hellman U, Dubremetz JF, Angel SO. Toxoplasma
gondii has two lineages of histones 2b (H2B) with different expression profiles. Molecular and
biochemical parasitology. 2006;148:103-7.
[23] Dalmasso MC, Onyango DO, Naguleswaran A, Sullivan WJ, Jr., Angel SO. Toxoplasma H2A
variants reveal novel insights into nucleosome composition and functions for this histone family. J
Mol Biol. 2009;392:33-47.
[24] Dalmasso MC, Sullivan WJ, Jr., Angel SO. Canonical and variant histones of protozoan parasites.
Frontiers in bioscience. 2011;16:2086-105.
[25] Petter M, Selvarajah SA, Lee CC, Chin WH, Gupta AP, Bozdech Z, et al. H2A.Z and H2B.Z double-
variant nucleosomes define intergenic regions and dynamically occupy var gene promoters in the
malaria parasite Plasmodium falciparum. Molecular microbiology. 2013;87:1167-82.
[26] Lowell JE, Kaiser F, Janzen CJ, Cross GA. Histone H2AZ dimerizes with a novel variant H2B and is
enriched at repetitive DNA in Trypanosoma brucei. J Cell Sci. 2005;118:5721-30.
[27] Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, et al. A Genome-wide CRISPR
Screen in Toxoplasma Identifies Essential Apicomplexan Genes. Cell. 2016;166:1423-35 e12.
[28] Nardelli SC, Che FY, Silmon de Monerri NC, Xiao H, Nieves E, Madrid-Aliste C, et al. The histone
code of Toxoplasma gondii comprises conserved and unique posttranslational modifications. mBio.
2013;4:e00922-13.
[29] Giaimo BD, Ferrante F, Herchenrother A, Hake SB, Borggrefe T. The histone variant H2A.Z in
gene regulation. Epigenetics & chromatin. 2019;12:37.
[30] Talbert PB, Henikoff S. Histone variants--ancient wrap artists of the epigenome. Nature reviews
Molecular cell biology. 2010;11:264-75.
[31] Bruce K, Myers FA, Mantouvalou E, Lefevre P, Greaves I, Bonifer C, et al. The replacement
histone H2A.Z in a hyperacetylated form is a feature of active genes in the chicken. Nucleic Acids
Res. 2005;33:5633-9.
[32] Binda O, Sevilla A, LeRoy G, Lemischka IR, Garcia BA, Richard S. SETD6 monomethylates H2AZ
on lysine 7 and is required for the maintenance of embryonic stem cell self-renewal. Epigenetics.
2013;8:177-83.
[33] Law C, Cheung P. Expression of Non-acetylatable H2A.Z in Myoblast Cells Blocks Myoblast
Differentiation through Disruption of MyoD Expression. J Biol Chem. 2015;290:13234-49.
[34] Valdes-Mora F, Song JZ, Statham AL, Strbenac D, Robinson MD, Nair SS, et al. Acetylation of
H2A.Z is a key epigenetic modification associated with gene deregulation and epigenetic remodeling
in cancer. Genome research. 2012;22:307-21.
[35] Whittle CM, McClinic KN, Ercan S, Zhang X, Green RD, Kelly WG, et al. The genomic distribution
and function of histone variant HTZ-1 during C. elegans embryogenesis. PLoS Genet.
2008;4:e1000187.
[36] Updike DL, Mango SE. Temporal regulation of foregut development by HTZ-1/H2A.Z and PHA-
4/FoxA. PLoS Genet. 2006;2:e161.
[37] Ridgway P, Brown KD, Rangasamy D, Svensson U, Tremethick DJ. Unique residues on the H2A.Z
containing nucleosome surface are important for Xenopus laevis development. J Biol Chem.
2004;279:43815-20.
[38] Myers FA, Chong W, Evans DR, Thorne AW, Crane-Robinson C. Acetylation of histone H2B
mirrors that of H4 and H3 at the chicken beta-globin locus but not at housekeeping genes. J Biol
Chem. 2003;278:36315-22.

[39] Miao J, Fan Q, Cui L, Li J, Li J, Cui L. The malaria parasite Plasmodium falciparum histones:
organization, expression, and acetylation. Gene. 2006;369:53-65.
[40] Trelle MB, Salcedo-Amaya AM, Cohen AM, Stunnenberg HG, Jensen ON. Global histone analysis
by mass spectrometry reveals a high content of acetylated lysine residues in the malaria parasite
Plasmodium falciparum. Journal of proteome research. 2009;8:3439-50.
[41] Radke JB, Lucas O, De Silva EK, Ma Y, Sullivan WJ, Jr., Weiss LM, et al. ApiAP2 transcription factor
restricts development of the Toxoplasma tissue cyst. Proc Natl Acad Sci U S A. 2013;110:6871-6.
[42] Hong DP, Radke JB, White MW. Opposing Transcriptional Mechanisms Regulate Toxoplasma
Development. mSphere. 2017;2.
[43] Gursoy-Yuzugullu O, Ayrapetov MK, Price BD. Histone chaperone Anp32e removes H2A.Z from
DNA double-strand breaks and promotes nucleosome reorganization and DNA repair. Proc Natl
Acad Sci U S A. 2015;112:7507-12.
[44] Morillo-Huesca M, Clemente-Ruiz M, Andujar E, Prado F. The SWR1 histone replacement
complex causes genetic instability and genome-wide transcription misregulation in the absence of
H2A.Z. PloS one. 2010;5:e12143.
[45] Munera Lopez J, Ganuza A, Bogado SS, Munoz D, Ruiz DM, Sullivan WJ, Jr., et al. Evaluation of
ATM Kinase Inhibitor KU-55933 as Potential Anti-Toxoplasma gondii Agent. Frontiers in cellular and
infection microbiology. 2019;9:26.
[46] Angel SO, Vanagas L, Ruiz DM, Cristaldi C, Saldarriaga Cartagena AM, Sullivan WJJ. Emerging
Therapeutic Targets Against Toxoplasma gondii: Update on DNA Repair Response Inhibitors and
Genotoxic Drugs. Front Cell Infect Microbiol. 2020;10.
[47] Behnke MS, Fentress SJ, Mashayekhi M, Li LX, Taylor GA, Sibley LD. The polymorphic
pseudokinase ROP5 controls virulence in Toxoplasma gondii by regulating the active kinase ROP18.
PLoS pathogens. 2012;8:e1002992.
[48] Etheridge RD, Alaganan A, Tang K, Lou HJ, Turk BE, Sibley LD. The Toxoplasma pseudokinase
ROP5 forms complexes with ROP18 and ROP17 kinases that synergize to control acute virulence in
mice. Cell host & microbe. 2014;15:537-50.
[49] Reese ML, Zeiner GM, Saeij JP, Boothroyd JC, Boyle JP. Polymorphic family of injected
pseudokinases is paramount in Toxoplasma virulence. Proc Natl Acad Sci U S A. 2011;108:9625-30.
[50] Saeij JP, Boyle JP, Coller S, Taylor S, Sibley LD, Brooke-Powell ET, et al. Polymorphic secreted
kinases are key virulence factors in toxoplasmosis. Science. 2006;314:1780-3.
[51] Fentress SJ, Behnke MS, Dunay IR, Mashayekhi M, Rommereim LM, Fox BA, et al.
Phosphorylation of immunity-related GTPases by a Toxoplasma gondii-secreted kinase promotes
macrophage survival and virulence. Cell host & microbe. 2010;8:484-95.
[52] Wysocka J. Identifying novel proteins recognizing histone modifications using peptide pull-
down assay. Methods. 2006;40:339-43.
[53] Croken MM, Nardelli SC, Kim K. Chromatin modifications, epigenetics, and how protozoan
parasites regulate their lives. Trends Parasitol. 2012;28:202-13.
[54] Croken MM, Qiu W, White MW, Kim K. Gene Set Enrichment Analysis (GSEA) of Toxoplasma
gondii expression datasets links cell cycle progression and the bradyzoite developmental program.
BMC genomics. 2014;15:515.
[55] Wang J, Dixon SE, Ting LM, Liu TK, Jeffers V, Croken MM, et al. Lysine acetyltransferase GCN5b
interacts with AP2 factors and is required for Toxoplasma gondii proliferation. PLoS pathogens.
2014;10:e1003830.
[56] Jeffers V, Gao H, Checkley LA, Liu Y, Ferdig MT, Sullivan WJ, Jr. Garcinol Inhibits GCN5-Mediated
Lysine Acetyltransferase Activity and Prevents Replication of the Parasite Toxoplasma gondii.
Antimicrobial agents and chemotherapy. 2016;60:2164-70.

[57] Huang S, Holmes MJ, Radke JB, Hong DP, Liu TK, White MW, et al. Toxoplasma gondii AP2IX-4
Regulates Gene Expression during Bradyzoite Development. mSphere. 2017;2.
[58] Khelifa AS, Guillen Sanchez C, Lesage KM, Huot L, Mouveaux T, Pericard P, et al. TgAP2IX-5 is a
key transcriptional regulator of the asexual cell cycle division in Toxoplasma gondii. Nature
communications. 2021;12:116.
[59] Sindikubwabo F, Ding S, Hussain T, Ortet P, Barakat M, Baumgarten S, et al. Modifications at
K31 on the lateral surface of histone H4 contribute to genome structure and expression in
apicomplexan parasites. eLife. 2017;6.
[60] Gissot M, Kelly KA, Ajioka JW, Greally JM, Kim K. Epigenomic modifications predict active
promoters and gene structure in Toxoplasma gondii. PLoS pathogens. 2007;3:e77.
[61] Ren Q, Gorovsky MA. Histone H2A.Z acetylation modulates an essential charge patch. Mol Cell.
2001;7:1329-35.
[62] Ren Q, Gorovsky MA. The nonessential H2A N-terminal tail can function as an essential charge
patch on the H2A.Z variant N-terminal tail. Mol Cell Biol. 2003;23:2778-89.
[63] Macadangdang BR, Oberai A, Spektor T, Campos OA, Sheng F, Carey MF, et al. Evolution of
histone 2A for chromatin compaction in eukaryotes. eLife. 2014;3.
[64] Severo V, Souza R, Vitorino F, Cunha J, Avila A, Arrizabalaga G, et al. Previously Unidentified
Histone H1-Like Protein Is Involved in Cell Division and Ribosome Biosynthesis in Toxoplasma gondii.
mSphere. 2022;7:e0040322.
[65] Sarah Sokol Borrelli SMR, Katherine G. Sharp, Leah F. Cabo, Hisham S. Alrubaye, Bruno
Martorelli Di Genova and Jon P. Boyle. A transcriptional network required for Toxoplasma gondii
tissue cyst formation is dispensable for long-term persistence. bioRxiv. 2022.
[66] M. Haley Licon CJG, Sundeep Chakladar, Lindsey Shallberg, Benjamin S. Waldman, Christopher
A. Hunter, Sebastian Lourido. A positive feedback loop controls Toxoplasma chronic differentiation.
Biorxvi. 2022.
[67] Afroz T, Cienikova Z, Clery A, Allain FHT. One, Two, Three, Four! How Multiple RRMs Read the
Genome Sequence. Methods in enzymology. 2015;558:235-78.
[68] Malumbres M. Cyclin-dependent kinases. Genome biology. 2014;15:122.
[69] Jin C, Zang C, Wei G, Cui K, Peng W, Zhao K, et al. H3.3/H2A.Z double variant-containing
nucleosomes mark 'nucleosome-free regions' of active promoters and other regulatory regions.
Nature genetics. 2009;41:941-5.
[70] Colino-Sanguino Y, Cornett EM, Moulder D, Smith GC, Hrit J, Cordeiro-Spinetti E, et al. A
Read/Write Mechanism Connects p300 Bromodomain Function to H2A.Z Acetylation. iScience.
2019;21:773-88.
[71] Rahal R, Amon A. Mitotic CDKs control the metaphase-anaphase transition and trigger spindle
elongation. Genes & development. 2008;22:1534-48.
[72] Lukasik P, Zaluski M, Gutowska I. Cyclin-Dependent Kinases (CDK) and Their Role in Diseases
Development-Review. International journal of molecular sciences. 2021;22.
[73] Ors A, Papin C, Favier B, Roulland Y, Dalkara D, Ozturk M, et al. Histone H3.3 regulates mitotic
progression in mouse embryonic fibroblasts. Biochemistry and cell biology = Biochimie et biologie
cellulaire. 2017;95:491-9.
[74] Page MJ, Di Cera E. Evolution of peptidase diversity. J Biol Chem. 2008;283:30010-4.
[75] Harvey SH, Krien MJ, O'Connell MJ. Structural maintenance of chromosomes (SMC) proteins, a
family of conserved ATPases. Genome biology. 2002;3:REVIEWS3003.
[76] Laflamme G, Tremblay-Boudreault T, Roy MA, Andersen P, Bonneil E, Atchia K, et al. Structural
maintenance of chromosome (SMC) proteins link microtubule stability to genome integrity. J Biol
Chem. 2014;289:27418-31.

[77] Serrano D, Cordero G, Kawamura R, Sverzhinsky A, Sarker M, Roy S, et al. The Smc5/6 Core
Complex Is a Structure-Specific DNA Binding and Compacting Machine. Mol Cell. 2020;80:1025-38
e5.
[78] Borgogno MV, Monti MR, Zhao W, Sung P, Argarana CE, Pezza RJ. Tolerance of DNA Mismatches
in Dmc1 Recombinase-mediated DNA Strand Exchange. J Biol Chem. 2016;291:4928-38.
[79] Kloehn J, Oppenheim RD, Siddiqui G, De Bock PJ, Kumar Dogga S, Coute Y, et al. Multi-omics
analysis delineates the distinct functions of sub-cellular acetyl-CoA pools in Toxoplasma gondii. BMC
Biol. 2020;18:67.
[80] Attar N, Campos OA, Vogelauer M, Cheng C, Xue Y, Schmollinger S, et al. The histone H3-H4
tetramer is a copper reductase enzyme. Science. 2020;369:59-64.
[81] Singh A, Verma S, Modak SB, Chaturvedi MM, Purohit JS. Extra-nuclear histones: origin,
significance and perspectives. Mol Cell Biochem. 2022;477:507-24.
[82] Choi YS, Hoon Jeong J, Min HK, Jung HJ, Hwang D, Lee SW, et al. Shot-gun proteomic analysis of
mitochondrial D-loop DNA binding proteins: identification of mitochondrial histones. Molecular
bioSystems. 2011;7:1523-36.
[83] Zanin MK, Donohue JM, Everitt BA. Evidence that core histone H3 is targeted to the
mitochondria in Brassica oleracea. Cell Biol Int. 2010;34:997-1003.
[84] Roson JN, Vitarelli MO, Costa-Silva HM, Pereira KS, Pires DDS, Lopes LS, et al. H2B.V demarcates
divergent strand-switch regions, some tDNA loci, and genome compartments in Trypanosoma cruzi
and affects parasite differentiation and host cell invasion. PLoS pathogens. 2022;18:e1009694.
[85] Denkers EY, Gazzinelli RT. Regulation and function of T-cell-mediated immunity during
Toxoplasma gondii infection. Clin Microbiol Rev. 1998;11:569-88.
[86] Blader IJ, Saeij JP. Communication between Toxoplasma gondii and its host: impact on parasite
growth, development, immune evasion, and virulence. APMIS. 2009;117:458-76.
[87] Elsaid MM, Vitor RW, Frezard FJ, Martins MS. Protection against toxoplasmosis in mice
immunized with different antigens of Toxoplasma gondii incorporated into liposomes. Mem Inst
Oswaldo Cruz. 1999;94:485-90.
[88] Sanchez-Lopez EF, Corigliano MG, Oliferuk S, Ramos-Duarte VA, Rivera M, Mendoza-Morales
LF, et al. Oral Immunization With a Plant HSP90-SAG1 Fusion Protein Produced in Tobacco Elicits
Strong Immune Responses and Reduces Cyst Number and Clinical Signs of Toxoplasmosis in Mice.
Front Plant Sci. 2021;12:726910.
[89] Buxton D, Innes EA. A commercial vaccine for ovine toxoplasmosis. Parasitology. 1995;110
Suppl:S11-6.
[90] Wu M, Liu S, Chen Y, Liu D, An R, Cai H, et al. Live-attenuated ME49Deltacdpk3 strain of
Toxoplasma gondii protects against acute and chronic toxoplasmosis. NPJ Vaccines. 2022;7:98.
[91] Echeverria PC, Matrajt M, Harb OS, Zappia MP, Costas MA, Roos DS, et al. Toxoplasma gondii
Hsp90 is a potential drug target whose expression and subcellular localization are developmentally
regulated. J Mol Biol. 2005;350:723-34.
[92] Guzman F, Gauna A, Roman T, Luna O, Alvarez C, Pareja-Barrueto C, et al. Tea Bags for Fmoc
Solid-Phase Peptide Synthesis: An Example of Circular Economy. Molecules. 2021;26.
[93] Spiess PC, Deng B, Hondal RJ, Matthews DE, van der Vliet A. Proteomic profiling of acrolein
adducts in human lung epithelial cells. J Proteomics. 2011;74:2380-94.

1071