Title: CXCR6 promotes dermal CD8+ T cell survival and transition to long-term tissue residence

Authors: Taylor A. Heim[1], Ziyan Lin[2], Maria M. Steele[1,3], Tenny Mudianto[1], Amanda W. Lund[1,3-5]*

Affiliations:

[1]Ronald O. Perelman Department of Dermatology, NYU Grossman School of Medicine, New York, NY, USA

[2]Applied Bioinformatics Laboratories, NYU Langone Health, New York, NY, USA

[3]Department of Cell, Developmental & Cancer Biology and Knight Cancer Institute, Oregon Health & Science University, Portland, OR, USA

[4]Department of Pathology, NYU Grossman School of Medicine, New York, NY, USA

[5]Laura and Isaac Perlmutter Cancer Center, NYU Grossman School of Medicine, New York, NY, USA

*Corresponding author. Email: amanda.lund@nyulangone.org

Abstract: Tissue resident memory T cells (T_{RM}) provide important protection against infection, and yet the interstitial signals necessary for their formation and persistence remain incompletely understood. Here we show that antigen-dependent induction of the chemokine receptor, CXCR6, is a conserved requirement for T_{RM} formation in peripheral tissue after viral infection. CXCR6 was dispensable for the early accumulation of antigen-specific CD8$^+$ T cells in skin and did not restrain their exit. Single cell sequencing indicated that CXCR6$^{-/-}$ CD8$^+$ T cells were also competent to acquire a transcriptional program of residence but exhibited deficiency in multiple pathways that converged on survival and metabolic signals necessary for memory. As such, CXCR6$^{-/-}$ CD8$^+$ T cells exhibited increased rates of apoptosis relative to controls in the dermis, leading to inefficient T_{RM} formation. CXCR6 expression may therefore represent a common mechanism across peripheral non-lymphoid tissues and inflammatory states that increases the probability of long-term residence.

One Sentence Summary: CXCR6 promotes mechanisms of cellular adaptation to tissue infiltration that support local survival and the transition to tissue residence.

Main Text:

INTRODUCTION

Resident memory T cells (T_{RM}) provide rapid, localized protection due to their position in barrier tissues. T_{RM} formation depends first upon the recruitment of effector T cells to sites of inflammation where local cues then direct the acquisition of a transcriptional program required for tissue adaptation and persistence. One key feature of this program is the downregulation of a transcriptional module associated with tissue exit via the lymphatic vasculature, including *Klf2*, *Ccr7*, and *S1pr1*, which presumably helps fix T_{RM} in a sedentary state. Consistent with this idea, *Ccr7* deficiency prevents CD4$^+$ T cell egress from the skin via dermal lymphatic vessels following viral infection (*1*) and boosts CD8$^+$ T_{RM} formation (*2*). Similarly, constitutive expression of *S1pr1* limits CD8$^+$ T_{RM} formation in skin (*3*) and the conserved T_{RM} marker CD69, is thought to promote residency by sequestering S1PR1 and inhibiting T cell egress (*4*). While these mechanisms may limit the exit of T_{RM} progenitors from inflamed tissues, the signals that instruct T cell position within tissue along the T_{RM} differentiation trajectory, and the interstitial cues that switch between these migratory modules, remain incompletely understood.

Interestingly, one of the highly expressed and conserved chemokine receptors in both CD4$^+$ and CD8$^+$ T_{RM} across mice and humans is CXCR6 (*5–7*). The sole ligand for CXCR6, CXCL16 (*8*), is expressed by multiple cell types including macrophages, dendritic cells, keratinocytes, and endothelial cells (*9–12*). Induced by TNFα and IFNγ, CXCL16 is expressed in a transmembrane form and acts in trans at the cell-cell interface, but may also be cleaved from the cell surface to signal as a classical chemoattractant (*13*). CXCR6 is functionally required for T_{RM} formation or maintenance in lung (*14*), skin (*15*), and liver (*16*), however, the specific mechanism by which CXCR6 guides CD8$^+$ T_{RM} formation remains incompletely understood. In a mouse model of influenza infection, CXCR6 is dispensable for CD8$^+$ T cell trafficking to the lung parenchyma but needed for localization to the CXCL16-expressing lung epithelium (*14*). In the liver, however, CXCR6 has been shown to be necessary for tissue infiltration during a graft vs host response (*17*), but also dispensable for early accumulation following malaria infection (*16*). Therefore, across peripheral, non-lymphoid tissues, when, exactly where, and how CXCR6 promotes CD8$^+$ T cell accumulation remains unclear.

Here we used cutaneous, murine viral infection to define the kinetics of CXCR6 expression and its impact on the transition of effector CD8$^+$ T cells to long-term tissue residence. We find that recirculating CD8$^+$ T cells in the spleen and within non-lymphoid peripheral tissues are distinguished from interstitial CD8$^+$ T$_{RM}$ in part by CXCR6 expression. CXCR6 expression was conserved across anatomical compartments where it was necessary for optimal CD8$^+$ T$_{RM}$ formation. In skin, rather than facilitating early CD8$^+$ T cell accumulation or directly preventing egress, we demonstrate that local antigen encounter upregulates CXCR6 to promote CD8$^+$ T cell survival in the dermis allowing for more efficient seeding of the epidermis over time. These data add to a mechanistic model for how CD8$^+$ T cell position in tissue optimizes receipt of the signals necessary for T$_{RM}$ formation and implicates CXCR6 as a conserved feature of tissue adaptation and survival.

RESULTS
CXCR6 is conserved on CD8$^+$ T$_{RM}$ across peripheral, non-lymphoid tissues

In mice and humans *Cxcr6* is a part of the core cluster of genes associated with residence (*5–7*), and is associated with exacerbated peripheral tissue inflammation (*18, 19*). Still, the mechanisms that regulate *Cxcr6* expression and its role in mediating peripheral tissue CD8$^+$ T cell behavior remain unclear. To track CXCR6 expression and determine its functional contribution to T$_{RM}$ formation we infected C57Bl/6 mice with vaccinia virus (VACV) by scarification on the ear pinna (10^6 PFU). CXCR6 was highly expressed on the surface of endogenous VACV-specific CD8$^+$ T cells (H2-K^b-B8R$_{20-27}$) in skin 30 days post infection (d.p.i). but was relatively low in VACV-specific and naive CD8$^+$ T cells in the spleen at matched time points (**Fig 1A and B**). While CD103 labels a subset (38±7% of CD69$^+$) of antigen-specific T$_{RM}$ in skin 28 d.p.i., CXCR6 was highly expressed on a larger percentage of CD69$^+$ cells (85±4%) (**Fig 1A and C**). Further, CD69$^+$ cells, which are considered to be bona fide T$_{RM}$ (*20*), had significantly higher levels of CXCR6 compared to CD69$^-$ T cells in the same skin (**Fig 1C**). These data may indicate that CXCR6 expression is not absolutely necessary for entry into the skin but increases upon tissue infiltration.

Given that T$_{RM}$ phenotype is highly dependent upon pathogenic insult and anatomical location (*21*) and therefore that residence mechanisms may be tissue specific, we asked whether high CXCR6 expression was restricted to CD8$^+$ T$_{RM}$ formed following cutaneous VACV infection. Vesicular

stomatitis virus (VSV) establishes CD8$^+$ T$_{RM}$ in diverse non-lymphoid tissues when delivered intravenously (10^6 PFU) (22). To seed antigen-specific T$_{RM}$ we transferred naïve CD45.1$^+$ OT-1 T cells, specific for H2K^b-OVA$_{257\text{-}264}$ (SIINFEKL), into mice and infected them with VSV-expressing ovalbumin (VSV-OVA). In all tissues tested (liver, brain, kidney, salivary gland, and female reproductive tract), CXCR6 expression was higher on tissue CD8$^+$ T$_{RM}$ (at least 30 d.p.i.) than antigen-specific CD8$^+$ T cells in the spleen (**Fig 1D**). Furthermore, as seen in skin, CD69$^+$ OT-1 T$_{RM}$ expressed significantly more CXCR6 compared to CD69$^-$ T cells in the same tissue (**Fig 1E and F**), suggesting that CXCR6 expression is a conserved feature of anti-viral, CD8$^+$ T$_{RM}$.

These data indicated that even with a fixed TCR and at a static time point, both circulating and resident CD8$^+$ T cells could be captured that were stratified by their expression of CXCR6. To gain higher resolution on the transcriptional CD8$^+$ T cell states captured within tissue at a single time point and to understand where CXCR6 is turned on in the T$_{RM}$ differentiation trajectory, we performed single-cell RNA sequencing (scRNAseq) on interstitial CD8$^+$ T cells. Congenically distinct, OT-1 T cells were transferred intravenously into C57Bl/6 mice. The following day, mice were infected with VACV expressing the immunodominant CD8$^+$ T cell epitope of OVA (OVA$_{257\text{-}264}$; VACV-OVA). Live, extravascular OT-1 T cells (negative for intravascular stain; IV$^-$) were sorted from infected skin 21 d.p.i. and submitted for scRNAseq to capture cells at the beginning of memory formation (20). Transcriptional analysis and visualization using Uniform Manifold Approximation Projection (UMAP) identified six clusters (**Fig. 1G and Fig S1A**). Cluster 1 expressed high levels of *Tcf7, Klf2,* and *S1pr1* and scored for a core circulating gene signature (T$_{CIRC}$, **Fig. 1H and S1B**), while clusters 2-5 expressed high levels of *Itgae* and *Cd69* and enriched for a transcriptional program associated with residence (T$_{RM}$, **Fig. 1H, S1C and S1D**). Cluster 6, interestingly expressed high levels of *Zeb2*, a marker of terminal differentiation that is absent from long-lived T$_{RM}$ memory following lymphocytic choriomeningitis virus (LCMV) infection (23). Cluster 6 also differentially expressed genes associated with RNA splicing and apoptotic cleavage of cellular proteins, perhaps indicating an increased rate in cell death (**Fig. S1E**). As previously reported, T$_{RM}$ CD8$^+$ T cells exhibited lower expression of *S1pr1* relative to T$_{CIRC}$, consistent with its proposed role in tissue egress (3). Of note, little *Ccr7* expression was seen in any subset, instead T$_{CIRC}$ cells expressed elevated levels of *Cxcr4,* which facilitates egress from melanoma through CXCL12-producing dermal lymphatic vessels (24). In addition to several receptors including,

S1pr4, *Ccr2*, *Ccr5,* and *Cxcr3, Cxcr6* was significantly enriched in T_{RM} relative to T_{CIRC} (**Fig 1I and S1F),** indicating that the high surface expression observed across T_{RM} transcriptional states likely depended upon a transcriptional event that occurred after tissue entry.

5   The heterogeneity observed within the T_{RM} clusters suggested that even at a single time point and with a fixed TCR, CD8$^+$ T cells could be captured in distinct transcriptional states that might reflect either functional or positional heterogeneity within the tissue. The T_{RM} clusters we defined (2-5) varied in their expression of effector molecule transcripts and several transcription factors related to T cell activation and differentiation. Cluster 2 cells were defined by high expression of

10   chemokines *Ccl3*, *Ccl4* and transcription factors *Fos* and *Jun*, which may be indicative of recent activation (*25*). Cells in cluster 3, interestingly, did not express transcripts high enough to meet our criteria for marker genes (fold change of 2, relative to average expression in other clusters), but expressed the highest levels of *Klf2*, *Tcf7* and *S1pr1* of the T_{RM} clusters. This cluster may therefore represent an intermediate state, not yet fully differentiated toward residency. Cluster 5

15   was defined by high expression of *Gzma*, *Gzmf* and *Krt83*, a keratin primarily found in the hair follicle, and marker genes for cluster 5 also included *Lmna* and *Anxa.* High *Cxcr6* expression was seen in all T_{RM} subclusters, including the intermediate cluster 3, pointing towards *Cxcr6* expression as an early event in T_{RM} differentiation (**Fig. S1G**).

20   **CXCR6 is required for optimal T_{RM} formation in skin**

Given the highly conserved expression of CXCR6 by T_{RM} across peripheral tissues, we asked whether CXCR6 was functionally required for CD8$^+$ T_{RM} formation. To this end, we co-transferred equal numbers of congenically mismatched, WT and CXCR6$^{-/-}$ (*Cxcr6*$^{GFP/GFP}$) (*26*) OT-1 T cells into mice and infected ear skin with VACV-OVA the following day (**Fig 2A and Fig S2A**). We

25   compared the ratio of WT and CXCR6$^{-/-}$ OT-1 T cells more than 45 d.p.i. and observed significant enrichment for WT over CXCR6$^{-/-}$ OT-1 T cells in the skin interstitium relative to the spleen (**Fig 2B and C**). These data indicated that CXCR6 was required for optimal CD8$^+$ T_{RM} formation in the skin following VACV infection, consistent with observations made using HSV (*15*). Interestingly, of the OT-1 T cells within the skin, WT cells exhibited more robust CD69 and CD103 expression

30   than CXCR6$^{-/-}$ OT-1 T cells (**Fig 2D and E**), perhaps indicating a less mature T_{RM} phenotype. We next asked if CXCR6 expression was also required for optimal T_{RM} formation in other non-

lymphoid tissues. Again, equal numbers of WT and CXCR6$^{-/-}$ OT-1 T cells were co-transferred into mice prior to infection with VSV-OVA (**Fig 2F**). At a memory time point (at least 30 d.p.i.) we compared the ratio of WT and CXCR6$^{-/-}$ OT-1 T cells in the spleen to that of other tissues. While we found that the differences in expression of CD69 and CD103 seen in skin were not replicated across all tissues (**Fig S2B-F**), we again observed enrichment for WT OT-1 T cells in all non-lymphoid tissues examined, including the brain, kidney and female reproductive tract (**Fig 2G**). As such, CXCR6 expression is necessary for CD8$^+$ T$_{RM}$ formation in disparate anatomical locations and in response to diverse viral infections.

Peripheral tissue antigen encounter upregulates CXCR6 expression

The low levels of CXCR6 expression on circulating anti-viral CD8$^+$ T cells within the skin (**Fig 1**) suggested that CXCR6 expression might not be strictly necessary for infiltration into inflamed skin. To directly test whether CXCR6 is required for tissue infiltration and the early accumulation of anti-viral effector CD8$^+$ T cells, we quantified the ratio of co-transferred WT and CXCR6$^{-/-}$ OT-1 T cells at 5 and 14 d.p.i. with VACV-OVA (**Fig 3A**). At 5 d.p.i. in the draining LN, WT and CXCR6$^{-/-}$ OT-1 T cells were present at similar levels, indicating that CXCR6 deficiency does not impair the magnitude of priming and expansion (**Fig S2G**). Both WT and CXCR6$^{-/-}$ OT-1 T cells infiltrated infected skin and were present at a similar ratio to that found in spleen at early (day 5) to late effector (day 14) time points (**Fig 3B**). Further, 5 d.p.i. CXCR6 expression by WT OT-1 T cells in skin was almost indistinguishable from expression on WT OT-1 T cells found in the spleen (**Fig 3C**) indicating that CXCR6 was likely upregulated after entry into the skin. Consistent with this hypothesis, CXCR6 surface expression increased in skin relative to spleen over time, with established CD8$^+$ T$_{RM}$ (day 49) expressing almost 6 times as much surface CXCR6 as their circulating memory counterparts (**Fig 3C**). Using the same co-transfer system and VSV-OVA infection, we next asked if CXCR6 facilitated early accumulation of T cells in tissues besides skin. Consistent with cutaneous VACV infection, CXCR6 was dispensable for early T cell accumulation in the skin following systemic VSV (**Fig S2H**), and also dispensable for accumulation in the small intestine epithelium, and brain 7 d.p.i with VSV-OVA. WT OT-1 T cells, however, outcompeted CXCR6$^{-/-}$ T cells in the liver, salivary gland, female reproductive tract, and kidney (**Fig S2I**). These differences may indicate either that low levels of CXCR6 on circulating CD8$^+$ T cells directly promotes transendothelial migration in a tissue-specific manner, or that CD8$^+$ T cells entering these

unique tissue environments are more sensitive to the effects of rapid, interstitial CXCR6 expression.

These data indicated that the induction of CXCR6 within the tissue parenchyma was a necessary event for T_{RM} formation in diverse peripheral tissues. Antigen encounter in peripheral non-lymphoid tissue induces CD69 expression and $CD8^+$ T_{RM} formation in both skin (*20*) and lung (*27, 28*) following infection. Given the concordance between CXCR6 expression and CD69, we hypothesized that CXCR6 enrichment in skin was also antigen dependent. To test this, we transferred naïve OT-1 $CD8^+$ T cells into mice and the next day infected one ear with VACV and the contralateral ear with VACV-OVA. VACV-OVA primed OT-I T cells infiltrate both infected sites independent of antigen expression (*20*), allowing us to evaluate CXCR6 levels on a fixed $CD8^+$ T cell population in the presence or absence of cognate antigen (**Fig 3D**). We observed significantly higher levels of CXCR6 on OT-1 T cells in the presence of cognate antigen (VACV-OVA) compared to VACV infected ears or the spleen 14 d.p.i. (**Fig 3E and F**). Interestingly, while cognate antigen clearly boosted surface CXCR6 expression on anti-viral $CD8^+$ T cells, there was a significant increase in CXCR6 on bystander OT-1 T cells in VACV-infected skin relative to spleen, indicating an additional role for inflammatory cytokines in surface CXCR6 expression in skin (e.g. IL-2 and IL-15) (*26*).

In order to further confirm a role for antigen encounter in CXCR6 expression and account for potential cytokine differences in VACV and VACV-OVA infected skin, we used mice that report TCR stimulation via transgenic expression of green fluorescent protein (GFP) under control of the *Nr4a1*(Nur77) promoter (Nurr77-GFP) (*29*). In VACV-OVA infected skin 7 d.p.i., surface CXCR6 was enriched on Nurr77-GFP expressing OT-1 T cells compared to Nurr77-GFP negative OT-1 T cells (**Fig 3G and H),** indicating that recent antigen recognition in skin is associated with increased CXCR6 levels. Furthermore, *ex vivo* activation of $CD8^+$ effector T cells (7 d.p.i. listeria monocytogenes-OVA) with cognate SIINFEKL antigen resulted in significant CXCR6 upregulation relative to unstimulated controls over 24 hours (**Fig 3I**). TGF-β signaling is required for the transition of circulating $CD8^+$ T cells towards residence, in part by downregulating S1PR1 and upregulating CD103 (*30–32*), however, the addition of TGF-β did not affect CXCR6 expression in the presence of antigen stimulation *ex vivo* (**Fig 3I**). These data, together with the

observation that CD103$^+$ T$_{RM}$ are a subset of CD69$^+$CXCR6$^+$ T$_{RM}$ (**Fig 1E**), indicate that antigen-dependent upregulation of CXCR6 likely precedes TGF-β driven changes and is therefore consistent with CXCR6 as an early feature of CD8$^+$ T cell adaptation to the infected dermis.

CXCR6/ CXCL16 interactions in the dermis do not act to restrain T cell exit

To begin to define the anatomical niche within which CXCR6 acts on CD8$^+$ T cell differentiation we analyzed expression of its ligand. CXCL16 was present in skin at all time points analyzed but appeared to peak in expression 7 d.p.i., returning to basal-like levels at late memory time points (**Fig 4A**). At effector time points (14 d.p.i.), CXCL16 was surprisingly absent from the epidermis (**Fig 4B**), unlike the lung where epithelial expression is notable (*14*). Instead, depots of CXCL16 were found within the dermis of infected skin (**Fig 4B**). Given that effector CD8$^+$ T cells first enter the dermis through activated capillaries and then transition with time to the epidermal compartment as they acquire a residence program (**Fig 4C**), the enrichment of CXCL16 in the dermis might suggest a role for CXCR6 in the transition to epidermal residence.

One hypothesis was that a failure to retain CD8$^+$ T cells in the dermis in a CXCL16/CXCR6-dependent manner could lead to increased rates of recirculation and exit through the proximal dermal lymphatic vessels. Dermal lymphatic vessels facilitate CD4$^+$ and CD8$^+$ T cell egress out of skin (*33*), and the egress of CD8$^+$ T cells out of skin might limit the number of CD8$^+$ T cells that transition to residence. We therefore asked if we could rescue CXCR6$^{-/-}$ T$_{RM}$ in the absence of dermal lymphatic vessels, where a physical loss of vessels would prevent T cell exit. Mice in which a VEGFR3-Immunogloubulin fusion protein is expressed under control of the K14 promoter (K14-VEGFR3-Ig), fail to form mature dermal lymphatic vessels (*34*), which restricts both fluid and cellular efflux out of skin (*35*). We again co-transferred equal numbers of congenically distinct, WT and CXCR6$^{-/-}$ OT-1 T cells into naïve K14-VEGFR3-Ig mice and WT controls (**Fig 4C**). Skin was infected with VACV-OVA and more than 45 d.p.i. we compared the ratio of WT and CXCR6$^{-/-}$ OT-1 T cells in the spleen to that of the skin (**Fig 4D**), expecting that if CXCR6 was a retention signal, the removal of the egress route would be sufficient to rescue their accumulation in skin. The lack of dermal lymphatic vasculature in K14-VEGFR3 mice, however, failed to normalize the ratio between WT and CXCR6$^{-/-}$ OT-1 T cells in the skin (**Fig 4E**). In line with this finding,

CXCR6$^{-/-}$ OT-1 T cells were as likely to egress through functional infection-associated lymphatic vessels as their WT counterparts when directly tracked using a photoconvertible transgenic mouse (Kaede-tg) (*33*, *36*). Kaede-tg OT-1 CD8$^+$ T cells and Kaede-tg CXCR6$^{-/-}$ OT-1 CD8$^+$ T cells were transferred into mice one day prior to infection with VACV-OVA. At 6 d.p.i, ear skin was photoconverted and the number of Kaede-red$^+$CD8$^+$ OT-I T cells were quantified in draining LNs 24 hours later (**Fig 4F, Fig S3A and B**). We found no difference in the ratio of photoconverted WT to CXCR6$^{-/-}$ CD8$^+$ T cells in the skin compared to those that had egressed to the draining LN (**Fig 4G and H**), indicating that despite being a marker of residence, CXCR6 did not appear to be acting as a physical retention cue. These data pointed to a dermally-restricted, CXCR6-dependent mechanism of CD8$^+$ T cell accumulation in infected skin.

CXCR6$^{-/-}$ T cells acquire the full spectrum of transcriptional T$_{RM}$ phenotypes

Our data indicated that CXCR6 likely acted in the dermis to facilitate the transition from effector T cell to T$_{RM}$ over time, though exactly how this chemokine receptor might do this remained unclear. At 21 d.p.i. *Cxcr6* is expressed in all WT OT-1 that bear a transcriptional program consistent with residency (**Fig 1**), indicating its expression is an early event in T$_{RM}$ differentiation. We reasoned that CXCR6 might influence the differentiation potential of effector CD8$^+$ T cells in skin. We therefore integrated our WT OT-1 scRNAseq with CXCR6$^{-/-}$ OT-1 T cells extracted from the same skin at 21 d.p.i., when WT and CXCR6$^{-/-}$ OT-1 CD8$^+$ T cells are still at roughly equal ratios (**Fig S4A**). Integrated analysis revealed a similar clustering pattern as seen for the WT cells alone (**Fig 1A**), again with a circulating cluster (T$_{CIRC}$: cluster 1), 4 clusters of T$_{RM}$ (T$_{RM}$: clusters 2,3,4 and 5) (**Fig 5A**), and a sixth cluster expressing high amounts of *Mcm9* (Mini chromosome maintenance 9) and the transcription factor *Zeb2,* associated with terminal differentiation (*37*).

Interestingly, both WT and CXCR6$^{-/-}$ OT-1 T cells were present in all clusters (**Fig 5A**) and proportional analysis revealed similar distribution of WT and CXCR6$^{-/-}$ OT-1 T cells in T$_{CIRC}$ and the T$_{RM}$ clusters (**Fig 5B**) as well as in each of the individual T$_{RM}$ subclusters (**Fig S4B**). Despite the relatively equal representation, we asked if WT and CXCR6$^{-/-}$ cells had similarly adopted core circulating and resident memory transcriptional programs. We scored T$_{CIRC}$ and T$_{RM}$ with published gene signatures of circulating and resident memory CD8$^+$ T cells (*38*). While WT and CXCR6$^{-/-}$ T$_{CIRC}$ clusters scored similarly for the established circulatory signature, CXCR6$^{-/-}$ OT-1

T cells in T_{RM} clusters scored significantly lower than WT for a resident memory gene signature (**Fig 5C**). Differentially expressed genes in CXCR6[-/-] T_{RM} included several genes (e.g. *Icos*, *Il21r*, and *Nr4a3*) (**Fig 5D**) that are functionally implicated in T_{RM} formation within various tissue compartments (*38–40*). Pathway analysis of the differentially expressed genes revealed that WT cells expressed higher levels of genes involved in leukocyte activation, cytokine signaling and regulation of apoptosis (**Fig 5E**). Genes enriched in these pathways included negative regulators of TCR and NFκB signaling (*Grap, Tnfaip3, Traf1, Nfkbia, Nfkbie*) (*41*) and components of the immunoproteasome (*Psmb8, Psmb9, Psma6*) important for regulating the transition to memory and cell survival (*42*). Referencing the TRRUST database confirmed transcriptional regulation of *Nfkb1* (p = 8.232e-8) and *Nfkbia* (p = 3.476e-5). In contrast, the pathway significantly affected in CXCR6[-/-] cells was oxidative phosphorylation (**Fig 5E**), largely driven by genes associated with mitochondrial electron transport complex (*Cox6c, Cox7a2, Cox8a, Ndufa4, Ndufb11*). Given that cellular metabolism is a common target of these pathways and since the formation of T cell memory is dependent upon mitochondrial biogenesis, which increases spare respiratory capacity to promote survival and rapid response to recall (*43*), we reasoned that these changes could signal sub-optimal cellular survival as a function of reduced cytokine and TCR-mediated metabolic reprogramming. We, therefore, directly quantified mitochondrial biomass and membrane potential. We noted that CXCR6[-/-] OT-1 T cells exhibited reduced mitochondrial mass (Mito Deep Red, **Fig 5F**) and membrane potential (CMX-Ros, **Fig. S4C**) 14 d.p.i. relative to WT OT-1 T cells in the same tissue. Taken together, these data indicate that while CXCR6[-/-] CD8[+] T cells could adopt a T_{RM}-like transcriptional state, they exhibited deficiencies across a range of pathways known to be necessary for T_{RM} formation and memory CD8[+] T cell survival*,* which may reduce the efficiency with which they transition to long-term residence.

Interestingly, though CXCR6[-/-] OT-1 T cells were equally represented in the T_{CIRC} and T_{RM} clusters, they were overrepresented (60%) in cluster 6. Cluster 6 exhibited signs of terminal differentiation, expressing high levels of *Zeb2,* and sat at the end of the predicted pseudotime trajectory defined by RNA velocity (**Fig. 5G**) (*44*). Following LCMV infection, a *Zeb2*-expressing CD8[+] T_{RM} cluster did not persist to late memory time points (60-90 d.p.i.) (*23*), in contrast to a cluster expressing *Id3*, *Jun*, *Fos*, *Ccl3*, and *Ccl4*, genes enriched in our T_{RM} cluster 1. We therefore hypothesized that cluster 6 captured cells falling off of the T_{RM} trajectory. Consistent with this

hypothesis, cluster 6 scored highest for an apoptotic gene signature (R-MMU-109581) (**Fig 5H and I**), with elevated expression of pro-apoptosis genes such as *Casp8, Bcl2l1,* and *Acin1* (**Fig 5J**). Even within the T_{RM} clusters, we found that transcripts for the anti-apoptotic proteins, *Bcl2a1b* and *Bcl2a1d,* were significantly reduced in CXCR6$^{-/-}$ relative to WT T cells (**Fig S4D**), perhaps indicating these cells were poised for apoptosis. *Bcl2a1* is an NF-κB target gene with four isoforms that promote lymphocyte survival and differentiation (*45*). BCL2 protein was indeed lower in CXCR6$^{-/-}$ OT-1 T cells in the skin 21 d.p.i (**Fig 5K**), and CXCR6$^{-/-}$ OT-1 T cells exhibited reduced viability in skin relative to matched WT T cells (**Fig 5L**), consistent with recent evidence that CXCR6 promotes T cell survival in the liver and tumor (*10*, *16*)

CXCR6 promotes dermal T cell survival during memory formation in infected skin

The transcriptional data pointed toward a loss of CXCR6$^{-/-}$ T cells as a function of poor survival likely driven by a combination of effects including reduced cytokine signaling, metabolic dysfunction, and a failure to fully activate a resident memory transcriptional program. Given the high expression of CXCL16 in the dermis, we asked whether CXCR6 preferentially impacted dermal CD8^{+} T cell viability and thereby transition to long-term maintenance in the epidermis. At 14 d.p.i. with VACV-OVA, when OT-1 T cells are present in both the dermis and epidermis, CXCR6$^{-/-}$ T cells exhibited similar viability in the epidermis but reduced viability in the dermis relative to WT (**Fig 6A and B**). We next asked if the reduced viability of CXCR6$^{-/-}$ T cells in the dermis impacted their frequency in the dermis and epidermis over time. Using immunofluorescent imaging of skin 14 d.p.i., we observed that WT OT-1 T cells were more prevalent than CXCR6$^{-/-}$ OT-1 T cells in the dermis, while in the epidermis, WT and CXCR6$^{-/-}$ OT-1 T cells were present at roughly equal levels (**Fig 6C and D**). This suggested that WT cells were outcompeting CXCR6$^{-/-}$ in the dermis at effector time points, prior to acquiring epidermal residence. We therefore hypothesized that CXCR6 supported tissue adaptation and survival during the transition to residency. Consistent with this hypothesis, at later time points, 42-45 d.p.i, CXCR6$^{-/-}$ OT-1 T cells were now less prevalent in both the dermis and epidermis. These data indicate that CXCR6$^{-/-}$ CD8^{+} T cells form T_{RM} at suboptimal levels due to reduced survival in the dermis during a critical window of T_{RM} differentiation. Our data therefore positions CXCR6 as a TCR-dependent, interstitial signal necessary for optimal survival and transition to residence.

DISCUSSION

The formation of T_{RM} in peripheral non-lymphoid tissues depends on mechanisms of tissue adaptation, cellular positioning, and differentiation. While the transcriptional programs that are enriched during transition from effector to memory are well described, the functional implications of key markers remain unclear. Transcriptional analyses of CD8[+] T cells in diverse anatomical locations of mice and humans identified the chemokine receptor CXCR6 as a highly expressed marker of T_{RM} (5–7). In this study, we investigated the functional significance of CXCR6 for T_{RM} formation. Using competitive co-transfer experiments of CXCR6[-/-] and WT controls, we find that CXCR6 expression is needed for optimal T_{RM} formation in peripheral non-lymphoid tissues after viral infection. Surprisingly, we found that CXCR6 has no effect on expansion in the draining lymph node, early accumulation of CD8[+] T cells in the skin, nor does it prevent egress out of skin after infection. Instead, CXCR6 deficiency diminished T_{RM} quantity by reducing CD8[+] T cell viability in inflamed skin, specifically within the dermis, leading to a failure to transition to epidermal residence.

Using adoptive transfer experiments and multiple viral models we show that CXCR6-deficiency diminished the expected quantity of T_{RM} in diverse tissues including brain, kidney and skin. While chemokines and their receptors are known to play critical roles in shaping the spatial distribution of T cells between and within tissues, here we identify a migration-independent mechanism of action. Across various non-lymphoid peripheral tissues and two infections, CXCR6 was not absolutely required for tissue infiltration. This is similar to observations made in mouse models of influenza infection, where CXCR6 is not needed for T cell entry into the lung parenchyma (14), and HSV infection of skin, where CXCR6[-/-] CD8[+] T cells still form suboptimal T_{RM} numbers even after direct intradermal injection (46). Furthermore, CXCR6 did not appear to help anchor T cells in tissue, as deficient T cells did not exhibit increased rates of lymphatic egress and loss of the lymphatic vasculature was not sufficient to boost T_{RM} formation. Further, we find CXCL16 to be most highly expressed in the dermis at effector time points after infection, which pointed towards a functional contribution prior to acquisition of epidermal residence. This observation was interestingly in contrast to the lung where CXCL16 is expressed by the airway epithelium and thought to promote antigen-specific CD8[+] T cell homing to that tissue compartment (14). Interestingly, in VACV-infected skin, CXCR3-deficient T cells fail to properly migrate towards

and kill infected cells in the epidermis (*47*), which may hint at tissue and or viral-specific mechanisms of target cell homing.

Our findings therefore point to a model where CD8$^+$ T cell antigen encounter in the dermis and concomitant CXCR6 upregulation precedes migration to the epidermis. Surface expression of CXCR6 by CD8$^+$ T cells in skin is upregulated between 5 and 7 d.p.i. and depends on local antigen recognition. We recently demonstrated in tumors that local antigen encounter downregulates CXCR4, which acts as a mechanism of tumor egress (*24*), and in concordance *Cxcr4* was transcriptionally downregulated in cells transitioning to residence following infection. In contrast, and as expected based on data presented here, CXCR6 was enriched on tumor-retained CD8$^+$ T cells that exhibited signs of chronic antigen stimulation (*24*). Interestingly, tissue entry and initial antigen presentation may be tightly linked in infected skin presenting the possibility that *Cxcr6* upregulation occurs rapidly in a dermal, perivascular niche (*48*), where at least in tumors, the CXCL16-expressing dendritic cells reside (*10*). We found that *Cxcr6* was expressed across transcriptional states in resident memory precursors and followed a similar kinetic as antigen-dependent induction of CD69 (*20*). Further consistent with our model, CXCR6 expression on restimulated effector CD8$^+$ T cells was not affected by the presence of TGF-β and CXCR6 upregulation precedes TGF-β-dependent CD103 upregulation by several days. CD103 permits adhesion to E-cadherin expressed in epidermal keratinocytes and TGF-β is activated at the epidermal interface (*49*, *50*). While TGF-β may facilitate residence in part through its role in downregulating *S1pr1, S1pr5* and *Cxcr4* (*32*), our data suggests CXCR6 function precedes TGF-β dependent mechanisms of residency. Taken together, these findings indicate that CXCR6 upregulation likely takes place in the dermis, early after tissue entry and before CD8$^+$ T cell transition to the epidermis.

Our data reveal that CXCR6 plays an important role for CD8$^+$ T cells in the transition from circulating effector to resident memory but CXCR6 deficiency does not place an absolute block on T$_{RM}$ differentiation. CXCR6$^{-/-}$ OT-I T cells at an early memory time points adopt the same transcriptional states seen in WT OT-1 T cells taken from skin, which is consistent with the observation that while loss of CXCR6 impairs T$_{RM}$ formation it does not completely prevent it. These observations may suggest that it's the efficiency of the process, not the ability to

differentiate, that limits T_{RM} formation. Interestingly, we did find that CXCR6-deficient T_{RM} in the skin score lower for a resident memory gene signature. The differences in this resident memory gene signature, while modest, were driven by reduced expression for over a dozen genes including several genes with known roles in T_{RM} formation including, *Il21r, Nr4a3* and *Icos* (*38–40*).

Decreased expression of *Il21r* is particularly interesting as IL-21 promotes memory T cell survival and proliferation, in a similar manner to IL-15 (*51*). In addition to these known T_{RM}-related transcripts, CXCR6$^{-/-}$ T cells appear to exhibit dysregulation of NFκ-B signaling, the immunoproteasome, oxidative phosphorylation and apoptosis, all of which may indicate a cell not activating the appropriate cues to efficiently transition to memory. Based on these data, it seems likely, that rather than a single transcriptional change being the driver of impaired T_{RM} formation, it's the effects on multiple survival and activation pathways that converge to decrease the probability of T cell survival and therefore long-term residence.

Previous studies evaluating the importance of CXCR6 expression by CD8$^+$ T cells in anti-tumor responses shed further light on why CXCR6 is crucial for T_{RM} development. In mouse models of melanoma and pancreatic cancer, CXCR6 expression is necessary for tumor control by CD8$^+$ T cells (*10, 52*). In these systems, CXCR6 optimizes interactions with CXCL16 and IL-15 expressing DCs in perivascular niches of the tumor stroma (*10*). T_{RM} dependency on IL-15 varies by anatomical location (*53*) but IL-15 is essential for the maintenance of CD103$^+$ T_{RM} in skin (*2*). IL-15 regulates oxidative phosphorylation for CD8$^+$ T cells by inducing mitochondrial biogenesis and also increases BCL-2 expression (*43*). Our data associated loss of BCL-2 with a reduction in mitochondrial biogenesis and membrane potential, which could lead to increased production of reactive oxygen species and ultimately cell death, and consistently we observe increased rates of antigen-specific T cell death in the dermis. These data are consistent with a hypothesis that reduced exposure to IL-15 in the dermis impairs the metabolic reprogramming necessary for survival and transition to memory. We cannot, however, exclude the possibility that CXCR6 might directly activate NF-κB through Akt (*54*), thereby improving CD8$^+$ memory T cell fitness in part through regulation of BCL2 (*55*).

Interestingly, in other peripheral tissues, such as the kidney, where T cells experience elevated levels of hypoxia (*56*), these stress-adaptation mechanisms may have more significant effects on

interstitial persistence. While CXCR6$^{-/-}$ T cells were equally competent to reside and accumulate in skin at early time points (7 d.p.i.), they were already significantly impaired in the kidney, salivary gland, liver, and female reproductive tract. While additional work would be needed to rule out a direct effect of CXCR6 on tissue homing and transendothelial migration, the relatively low levels of expression on circulating CD8$^+$ T cells and robust upregulation upon tissue infiltration potentially points to a tissue-restricted survival failure. Further speaking to its broad impact on tissue residency, CXCR6 is expressed on diverse resident lymphocyte populations including CD4$^+$ T$_{RM}$ in the liver lung and spleen . Furthermore, resident NKT cells in the liver, along with unconventional T cells in the LN, also show high levels of CXCR6 (*5, 6, 57–59*). It is unclear if CXCR6 expression by these lymphocyte populations also promotes survival in the same manner as in the skin after viral infection. Nevertheless, the highly conserved expression of CXCR6 suggests it likely facilitates a critical mechanism of cellular adaptation adopted across peripheral non-lymphoid tissues and inflammatory states.

Taken all together, our findings demonstrate that CXCR6 instigates mechanisms of cellular adaptation to tissue and promotes local survival by facilitating signals that boost mitochondrial biogenesis and protect cells from death. Here, we have demonstrated that CXCR6 contributes to the process of T$_{RM}$ formation by facilitating CD8$^+$ T cell survival and transition to residency during phases of immune resolution 14-30 days post infection. It will be interesting to determine if CXCR6 remains required to maintain T$_{RM}$ survival in the epidermis or if it is only needed during these inflammatory periods when CXCL16 expression is elevated, dermal effector cells are transitioning to epidermal memory, and survival cytokines might be limiting. Experimental models allowing for spatial and temporal control of CXCR6 or its ligand, CXCL16, will be needed to determine their role in maintenance and potential to regulate established T$_{RM}$ populations. Since T$_{RM}$ are potent sources of localized immunity that can mediate tissue-specific autoimmune responses, mechanisms of cellular adaptation to tissue stressors that are required for T$_{RM}$ formation and maintenance may represent targets to limit T cell-induced injury across tissue sites (*56*). In sum, our data provide mechanistic insight into the parameters that tune CD8$^+$ T cell transition to residence and contributes to an emerging hypothesis that targeting T cell position is a viable strategy for tissue-specific immunotherapy across disease indications.

MATERIALS AND METHODS
Mice

C57BL/6J, B6.SJL-PtprcaPepcb/BoyJ (CD45.1), C57BL/6-Tg(TcraTcrb)1100Mjb/J (OT-1), B6.129P2-Cxcr6tm1Litt/J (CXCR6$^{-/-}$) (*26*), B6.PL-Thy1a/CyJ (CD90.1), C57BL/6-Tg(Nr4a1-EGFP/cre)820Khog/J (Nur77-GFP) mice were purchased from Jackson Laboratory and bred in specific pathogen-free conditions at Oregon Health and Sciences University or New York University. B6.Cg-Tg(CAG-tdKaede)-15Utr (Kaede-Tg) (*60*) were obtained via D.J. Fowell in agreement with RIKEN BioResource Research Center. K14-VEGFR3-Ig mice (*34*) were a generous gift from Dr. Kari Alitalo and obtained from Dr. Melody A. Swartz. Mice were age and sex matched and female mice were used unless otherwise stated. OT-1 chimeras were created by adoptively transferring 15,000 CD8^{+} T cells from OT-1 mice into naïve mice. For co-transfer experiments, 15,000 wildtype OT-1 T cells and 15,000 CXCR6$^{-/-}$ OT-1 T cells were co-transferred into naïve mice. For experiments with K14-VEGFR3-Ig mice, both male and female recipients were used and 30,000 wildtype OT-1 T cells and 30,000 CXCR6$^{-/-}$ OT-1 T cells were co-transferred into naïve mice. Mice were used between 8-14 weeks of age. All animal procedures were approved and performed in accordance with the Institutional Animal Care and Use Committees at OHSU and NYU Langone Health.

Pathogens and Infections

Mice were infected the following day after adoptive transfer, with vaccinia virus (VACV), VACV expressing SIINFEKL (VACV-OVA) or VSV expressing full length ovalbumin (VSV-OVA). VACV, and VACV-OVA were propagated in BSC-40 cells as per standard protocols. Mice were infected by administering 1-5 x10^{6} PFU of VACV, or VACV-OVA in 10 µl of PBS to the ventral side of the ear pinna followed by 25 pokes with a 29-G needle (skin scarification). For VSV-OVA infections, mice were injected intravenous with 1 x 10^{7} PFU. To generate effector CD8^{+} T cells for in vitro studies, ActA deficient *Listeria monocytogenes* expressing SIINFEKL (LM-OVA) was grown in tryptic soy broth supplemented with 50 µg/ml streptomycin at 37°C and until it reached 1 x 10^{8} CFU/ml (OD600=0.1). 1 x 10^{7} CFU was administered intravenously and spleens harvested 7 days later.

In vitro Restimulation

Splenocytes were cultured *in vitro* for 24 hours in 1nm SIINFEKL peptide or with 10ng/mL of TGF-β in RPMI containing 10% FBS.

In vivo T cell egress

To label Kaede expressing T cells, ear skin was photoconverted 6 days after infection using 405-nm light (Steele 2019) for 2 minutes (1 minute each side) at 100mW. Mice were euthanized 24 hours later.

Immunofluorescent microscopy

After euthanizing mice, ears were frozen in OCT on an isopentane bath. 7μm sections were cut and fixed in acetone for 15 minutes. Slides were blocked for 10 minutes with (1%BSA in PBS) and stained with antibodies in 1%BSA in PBS for 1-2 hours at room temperature. Antibodies were obtained from Biolegend, Invitrogen, BD Biosciences, Bioss, and Tonbo and included CXCL16 (pAb Bioss), K14 (POLY9060), CD8 (53-6.7), CD45.1 (A20), CD90.1 (OX-7). Nuclei were detected and cover slips were attached using 4′,6-diamidino-2-phenylindole dihydrochloride (DAPI) containing Prolong Diamond Anti-Fade (Life Technologies). Slides were imaged on a Keyence BX-X810 microscope.

Leukocyte isolation

Ears of mice were removed and the dorsal and ventral sides separated and incubated for 45 minutes at 37 C in 1ml of HBSS (Hyclone) containing $CaCl_2$ and $MgCl_2$ supplemented with 125 U/ml of collagenase D (Invitrogen) and 60 U/ml of Dnase-I (Sigma-Aldrich). Tissue was smashed on a scored plate and poured through a 70 μm filter. For some experiments, leukocytes were purified by resuspending pellets in 35% Percoll (GE Healthcare) and HBSS followed by room temperature centrifugation at 1800 RPM. Single cell suspensions of leukocytes from lymph nodes and spleens were made by smashing the tissue through a 70μm filter. Spleens were resuspended for 2 minutes in 2ml of ammonium-chloride-potassium lysis buffer. Lymphocyte isolation from all other non-lymphoid tissues was performed as previously described (*61*)(Steinert 2015) using 35% Percoll and HBSS followed by room temperature centrifugation at 1800 RPM.

Flow Cytometry

For intravenous (IV) labeling, mice were injected intravenous with 2-3 μg of anti-CD45 antibody and euthanized 3 minutes later. For surface stains, single-cell suspensions were stained with antibodies in FACS buffer (1%BSA in PBS) for 30 minutes at 4C. Antibodies were obtained from Biolegend, Invitrogen, BD Biosciences, Bioss, and Tonbo and included BCL2 (BCL/10C4 or 3F11), CD8 (53-6.7), CD45.1 (A20), CD90.1 (OX-7), CD90.2 (30-H12), CXCR6 (SA051D1), CD69 (H1.2F3), CD103 (2E7), CD44 (IM7), CD45 (30-F11), and CD16/32 (2.4G2). Ghost dye 780 (ZombieRed) was used to measure cell viability. For some flow cytometry analysis, $CXCR6^{-/-}$ $CD90.1^{+}$ OT-1 T cells were identified via GFP expression and AF488-CD90.1 antibody. Staining of intracellular BCL2 was performed with True-Nuclear™ Transcription Factor Buffer Set (Biolegend) as per manufacturer protocol. Samples were analyzed on a BD LSR II, or BD FACSymphony. FlowJo 10.8.1. was used to analyze flow cytometry data.

ELISA

Ears from naïve or previously infected mice were weighed and snap frozen in liquid nitrogen. Ears were homogenized and CXCL16 concentration was quantified using Mouse CXCL16 DuoSet ELISA (R&D Systems) as per the manufacturers protocol.

Sorting and single-cell RNA-seq

Ears of C57Bl/6J mice were infected with VACV-OVA the day after receiving 15,000 WT OT-1 T cells and 15,000 $CXCR6^{-/-}$ OT-1 T cells. At 21 days post infection, lymphocytes were isolated from skin after intravenous injection of anti-CD45-PE antibody as previously described. WT OT-1 T cells ($CD8^{+}CD90.2^{+}CD44^{+}CD45$ IV$^{-}CD45.1^{+}CD90.1^{-}$) and $CXCR6^{-/-}$ OT-1 T cells ($CD8^{+}CD90.2^{+}CD44^{+}CD45$ IV$^{-}CD45.1^{-}CD90.1^{+}$) were sorted. The sorted cellular suspensions were loaded on a 10x Genomics Chromium instrument to generate single-cell gel beads in emulsion. Libraries were prepared using Single cell 3'Reagent kits v3.1 (Chromium Next GEM Single Cell 3' GEM, Library & Gel Bead Kit v3.1, 16 rxns PN-1000121;10x Genomics) and were sequenced using Illumina Novaseq 6000.

Bioinformatics and pathway analysis

Quality control was performed on sequenced cells to calculate the number of genes, UMIs and the proportion of mitochondrial genes for each cell. Cells with low number of covered genes (gene-

count < 200) and high mitochondrial counts (mt-genes > 0.08) were filtered out. There were 8634 cells in the Cxcr6$^{-/-}$ sample and 7409 cells in the WT sample after filtering. The matrix was normalized based on their library sizes. A general statistical test was performed to calculate gene dispersion, base mean and cell coverage to use to build a gene model for performing Principal Component Analysis (PCA). Genes with high coverage (top 500) and high dispersion (dispersion > 1.5) were chosen to perform PCA and batch alignment using iCellR R package (v1.6.5). T-distributed Stochastic Neighbor Embedding (t-SNE) and Uniform Manifold Approximation and Projection (UMAP) were performed on the top 10 PCs. PhenoGraph (*62*) clustering was then performed on the UMAP results. Marker genes were found for each cluster and visualized on heatmaps, bar plots and box plots. The marker genes were used to determine cell types. Proportion (percentage) of the cell communities in each condition were calculated. The Tirosh scoring method (*63*) was used to calculate gene signature scores for circulating and resident memory (*38*). RNA velocity results were performed by using velocyto (v0.17) (*64*) https://doi.org/10.1038/s41586-018-0414-6) and scVelo (v0.2.3) (https://scvelo.readthedocs.io/). Pathway analysis was performed using marker genes in Metascape v.3.5 (http://metascape.org).

Statistics

Statistical analysis was performed using GraphPad Prism 9 software. Comparisons between two groups were conducted using paired or unpaired students t tests. Comparisons between three or more unpaired groups were conducted using ANOVA with Tukeys multiple comparison test. Comparisons between three or more paired groups were conducted using repeated measures ANOVA with Greenhouse-Geisser correction. Proper sample size was based off of prior experience. Statistical analysis of sequencing data was performed with pairwise Wilcoxon rank test and Bonferroni correction in R. P values are reported as follows: * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$. Error bars show the mean $\pm$SD.

Supplementary Materials

Figs. S1 to S4

Table S1 to S2

References

1. S. K. Bromley, S. Yan, M. Tomura, O. Kanagawa, A. D. Luster, Recirculating Memory T Cells Are a Unique Subset of CD4+ T Cells with a Distinct Phenotype and Migratory Pattern. *J. Immunol.* **190**, 970–976 (2013).

2. L. K. Mackay, A. Rahimpour, J. Z. Ma, N. Collins, A. T. Stock, M.-L. Hafon, J. Vega-Ramos, P. Lauzurica, S. N. Mueller, T. Stefanovic, D. C. Tscharke, W. R. Heath, M. Inouye, F. R. Carbone, T. Gebhardt, The developmental pathway for CD103+CD8+ tissue-resident memory T cells of skin. *Nat. Immunol.* **14**, 1294–1301 (2013).

3. C. N. Skon, J.-Y. Lee, K. G. Anderson, D. Masopust, K. A. Hogquist, S. C. Jameson, Transcriptional downregulation of S1pr1 is required for establishment of resident memory CD8+ T cells. *Nat. Immunol.* **14**, 1285–1293 (2013).

4. L. K. Mackay, A. Braun, B. L. Macleod, N. Collins, C. Tebartz, S. Bedoui, F. R. Carbone, T. Gebhardt, Cutting edge: CD69 interference with sphingosine-1-phosphate receptor function regulates peripheral T cell retention. *J. Immunol. Baltim. Md 1950*. **194**, 2059–2063 (2015).

5. B. V. Kumar, W. Ma, M. Miron, T. Granot, R. S. Guyer, D. J. Carpenter, T. Senda, X. Sun, S.-H. Ho, H. Lerner, A. L. Friedman, Y. Shen, D. L. Farber, Human Tissue-Resident Memory T Cells Are Defined by Core Transcriptional and Functional Signatures in Lymphoid and Mucosal Sites. *Cell Rep.* **20**, 2921–2934 (2017).

6. L. K. Mackay, M. Minnich, N. A. M. Kragten, Y. Liao, B. Nota, C. Seillet, A. Zaid, K. Man, S. Preston, D. Freestone, A. Braun, E. Wynne-Jones, F. M. Behr, R. Stark, D. G. Pellicci, D. I. Godfrey, G. T. Belz, M. Pellegrini, T. Gebhardt, M. Busslinger, W. Shi, F. R. Carbone, R. A. W. van Lier, A. Kallies, K. P. J. M. van Gisbergen, Hobit and Blimp1 instruct a universal transcriptional program of tissue residency in lymphocytes. *Science*. **352**, 459–463 (2016).

7. P. Hombrink, C. Helbig, R. A. Backer, B. Piet, A. E. Oja, R. Stark, G. Brasser, A. Jongejan, R. E. Jonkers, B. Nota, O. Basak, H. C. Clevers, P. D. Moerland, D. Amsen, R. A. W. van Lier, Programs for the persistence, vigilance and control of human CD8+ lung-resident memory T cells. *Nat. Immunol.* **17**, 1467–1478 (2016).

8. M. Matloubian, A. David, S. Engel, J. E. Ryan, J. G. Cyster, A transmembrane CXC chemokine is a ligand for HIV-coreceptor Bonzo. *Nat. Immunol.* **1**, 298–304 (2000).

9. B. B. Ural, S. T. Yeung, P. Damani-Yokota, J. C. Devlin, M. de Vries, P. Vera-Licona, T. Samji, C. M. Sawai, G. Jang, O. A. Perez, Q. Pham, L. Maher, P. Loke, M. Dittmann, B. Reizis, K. M. Khanna, Identification of a nerve-associated, lung-resident interstitial macrophage subset with distinct localization and immunoregulatory properties. *Sci. Immunol.* **5**, eaax8756 (2020).

10. M. Di Pilato, R. Kfuri-Rubens, J. N. Pruessmann, A. J. Ozga, M. Messemaker, B. L. Cadilha, R. Sivakumar, C. Cianciaruso, R. D. Warner, F. Marangoni, E. Carrizosa, S.

Lesch, J. Billingsley, D. Perez-Ramos, F. Zavala, E. Rheinbay, A. D. Luster, M. Y. Gerner, S. Kobold, M. J. Pittet, T. R. Mempel, CXCR6 positions cytotoxic T cells to receive critical survival signals in the tumor microenvironment. *Cell*. **184**, 4512-4530.e22 (2021).

11. F. Scholz, A. Schulte, F. Adamski, C. Hundhausen, J. Mittag, A. Schwarz, M.-L. Kruse, E. Proksch, A. Ludwig, Constitutive Expression and Regulated Release of the Transmembrane Chemokine CXCL16 in Human and Murine Skin. *J. Invest. Dermatol.* **127**, 1444–1455 (2007).

12. B. Linke, S. Meyer Dos Santos, B. Picard-Willems, M. Keese, S. Harder, G. Geisslinger, K. Scholich, CXCL16/CXCR6-mediated adhesion of human peripheral blood mononuclear cells to inflamed endothelium. *Cytokine*. **122**, 154081 (2019).

13. S. Abel, C. Hundhausen, R. Mentlein, A. Schulte, T. A. Berkhout, N. Broadway, D. Hartmann, R. Sedlacek, S. Dietrich, B. Muetze, B. Schuster, K.-J. Kallen, P. Saftig, S. Rose-John, A. Ludwig, The Transmembrane CXC-Chemokine Ligand 16 Is Induced by IFN-γ and TNF-α and Shed by the Activity of the Disintegrin-Like Metalloproteinase ADAM10. *J. Immunol.* **172**, 6362–6372 (2004).

14. A. N. Wein, S. R. McMaster, S. Takamura, P. R. Dunbar, E. K. Cartwright, S. L. Hayward, D. T. McManus, T. Shimaoka, S. Ueha, T. Tsukui, T. Masumoto, M. Kurachi, K. Matsushima, J. E. Kohlmeier, CXCR6 regulates localization of tissue-resident memory CD8 T cells to the airways. *J. Exp. Med.* **216**, 2748–2762 (2019).

15. A. Zaid, J. L. Hor, S. N. Christo, J. R. Groom, W. R. Heath, L. K. Mackay, S. N. Mueller, Chemokine Receptor–Dependent Control of Skin Tissue–Resident Memory T Cell Formation. *J. Immunol.* **199**, 2451–2459 (2017).

16. S.-W. Tse, A. J. Radtke, D. A. Espinosa, I. A. Cockburn, F. Zavala, The Chemokine Receptor CXCR6 Is Required for the Maintenance of Liver Memory CD8+ T Cells Specific for Infectious Pathogens. *J. Infect. Dis.* **210**, 1508–1516 (2014).

17. T. Sato, H. Thorlacius, B. Johnston, T. L. Staton, W. Xiang, D. R. Littman, E. C. Butcher, Role for CXCR6 in Recruitment of Activated CD8+ Lymphocytes to Inflamed Liver. *J. Immunol.* **174**, 277–283 (2005).

18. W. G. McMaster, A. Kirabo, M. S. Madhur, D. G. Harrison, Inflammation, Immunity, and Hypertensive End-Organ Damage. *Circ. Res.* **116**, 1022–1033 (2015).

19. A. Wehr, C. Baeck, F. Heymann, P. M. Niemietz, L. Hammerich, C. Martin, H. W. Zimmermann, O. Pack, N. Gassler, K. Hittatiya, A. Ludwig, T. Luedde, C. Trautwein, F. Tacke, Chemokine Receptor CXCR6-Dependent Hepatic NK T Cell Accumulation Promotes Inflammation and Liver Fibrosis. *J. Immunol.* **190**, 5226–5236 (2013).

20. T. N. Khan, J. L. Mooster, A. M. Kilgore, J. F. Osborn, J. C. Nolz, Local antigen in nonlymphoid tissue promotes resident memory CD8+ T cell formation during viral infection. *J. Exp. Med.* **213**, 951–966 (2016).

21. D. Masopust, A. G. Soerens, Tissue-Resident T Cells and Other Resident Leukocytes. *Annu. Rev. Immunol.* **37**, 521–546 (2019).

22. D. Masopust, V. Vezys, A. L. Marzo, L. Lefrançois, Preferential Localization of Effector Memory Cells in Nonlymphoid Tissue. *Science*. **291**, 2413–2417 (2001).

23. N. S. Kurd, Z. He, T. L. Louis, J. J. Milner, K. D. Omilusik, W. Jin, M. S. Tsai, C. E. Widjaja, J. N. Kanbar, J. G. Olvera, T. Tysl, L. K. Quezada, B. S. Boland, W. J. Huang, C. Murre, A. W. Goldrath, G. W. Yeo, J. T. Chang, Early precursors and molecular determinants of tissue-resident memory CD8+ T lymphocytes revealed by single-cell RNA sequencing. *Sci. Immunol.* **5**, eaaz6894 (2020).

24. M. M. Steele, I. D. Dryg, D. Murugan, J. Femel, H. du Bois, C. Hill, S. A. Leachman, Y. H. Chang, L. M. Coussens, A. W. Lund, "T cell Egress via Lymphatic Vessels Limits the Intratumoral T cell Repertoire in Melanoma" (preprint, Immunology, 2022), , doi:10.1101/2022.05.30.494080. in press *Nature Immunology*

25. J. Jain, P. G. McCaffrey, V. E. Valge-Archer, A. Rao, Nuclear factor of activated T cells contains Fos and Jun. *Nature*. **356**, 801–804 (1992).

26. D. Unutmaz, W. Xiang, M. J. Sunshine, J. Campbell, E. Butcher, D. R. Littman, The Primate Lentiviral Receptor Bonzo/STRL33 Is Coordinately Regulated with CCR5 and Its Expression Pattern Is Conserved Between Human and Mouse. *J. Immunol.* **165**, 3284–3292 (2000).

27. S. R. McMaster, A. N. Wein, P. R. Dunbar, S. L. Hayward, E. K. Cartwright, T. L. Denning, J. E. Kohlmeier, Pulmonary antigen encounter regulates the establishment of tissue-resident CD8 memory T cells in the lung airways and parenchyma. *Mucosal Immunol.* **11**, 1071–1078 (2018).

28. J. M. Stolley, T. S. Johnston, A. G. Soerens, L. K. Beura, P. C. Rosato, V. Joag, S. P. Wijeyesinghe, R. A. Langlois, K. C. Osum, J. S. Mitchell, D. Masopust, Retrograde migration supplies resident memory T cells to lung-draining LN after influenza infection. *J. Exp. Med.* **217**, e20192197 (2020).

29. A. E. Moran, K. L. Holzapfel, Y. Xing, N. R. Cunningham, J. S. Maltzman, J. Punt, K. A. Hogquist, T cell receptor signal strength in Treg and iNKT cell development demonstrated by a novel fluorescent reporter mouse. *J. Exp. Med.* **208**, 1279–1289 (2011).

30. A. P. Nath, A. Braun, S. C. Ritchie, F. R. Carbone, L. K. Mackay, T. Gebhardt, M. Inouye, Comparative analysis reveals a role for TGF-β in shaping the residency-related transcriptional signature in tissue-resident memory CD8+ T cells. *PLOS ONE*. **14**, e0210495 (2019).

31. K. A. Casey, K. A. Fraser, J. M. Schenkel, A. Moran, M. C. Abt, L. K. Beura, P. J. Lucas, D. Artis, E. J. Wherry, K. Hogquist, V. Vezys, D. Masopust, Antigen-independent differentiation and maintenance of effector-like resident memory T cells in tissues. *J. Immunol. Baltim. Md 1950*. **188**, 4866–4875 (2012).

32. S. N. Christo, M. Evrard, S. L. Park, L. C. Gandolfo, T. N. Burn, R. Fonseca, D. M. Newman, Y. O. Alexandre, N. Collins, N. M. Zamudio, F. Souza-Fonseca-Guimaraes, D. G. Pellicci, D. Chisanga, W. Shi, L. Bartholin, G. T. Belz, N. D. Huntington, A. Lucas, M. Lucas, S. N. Mueller, W. R. Heath, F. Ginhoux, T. P. Speed, F. R. Carbone, A. Kallies, L. K. Mackay, Discrete tissue microenvironments instruct diversity in resident memory T cell function and plasticity. *Nat. Immunol.* **22**, 1140–1151 (2021).

33. M. Tomura, T. Honda, H. Tanizaki, A. Otsuka, G. Egawa, Y. Tokura, H. Waldmann, S. Hori, J. G. Cyster, T. Watanabe, Y. Miyachi, O. Kanagawa, K. Kabashima, Activated regulatory T cells are the major T cell type emigrating from the skin during a cutaneous immune response in mice. *J. Clin. Invest.* **120**, 883–893 (2010).

34. T. Mäkinen, L. Jussila, T. Veikkola, T. Karpanen, M. I. Kettunen, K. J. Pulkkanen, R. Kauppinen, D. G. Jackson, H. Kubo, S.-I. Nishikawa, S. Ylä-Herttuala, K. Alitalo, Inhibition of lymphangiogenesis with resulting lymphedema in transgenic mice expressing soluble VEGF receptor-3. *Nat. Med.* **7**, 199–205 (2001).

35. C. P. Loo, N. A. Nelson, R. S. Lane, J. L. Booth, S. C. Loprinzi Hardin, A. Thomas, M. K. Slifka, J. C. Nolz, A. W. Lund, Lymphatic Vessels Balance Viral Dissemination and Immune Activation following Cutaneous Viral Infection. *Cell Rep.* **20**, 3176–3187 (2017).

36. M. M. Steele, M. J. Churchill, A. P. Breazeale, R. S. Lane, N. A. Nelson, A. W. Lund, Quantifying Leukocyte Egress via Lymphatic Vessels from Murine Skin and Tumors. *J. Vis. Exp.*, 58704 (2019).

37. K. D. Omilusik, J. A. Best, B. Yu, S. Goossens, A. Weidemann, J. V. Nguyen, E. Seuntjens, A. Stryjewska, C. Zweier, R. Roychoudhuri, L. Gattinoni, L. M. Bird, Y. Higashi, H. Kondoh, D. Huylebroeck, J. Haigh, A. W. Goldrath, Transcriptional repressor ZEB2 promotes terminal differentiation of CD8+ effector and memory T cell populations during infection. *J. Exp. Med.* **212**, 2027–2039 (2015).

38. J. J. Milner, C. Toma, B. Yu, K. Zhang, K. Omilusik, A. T. Phan, D. Wang, A. J. Getzler, T. Nguyen, S. Crotty, W. Wang, M. E. Pipkin, A. W. Goldrath, Runx3 programs CD8+ T cell residency in non-lymphoid tissues and tumours. *Nature.* **552**, 253–257 (2017).

39. C. Peng, M. A. Huggins, K. M. Wanhainen, T. P. Knutson, H. Lu, H. Georgiev, K. L. Mittelsteadt, N. N. Jarjour, H. Wang, K. A. Hogquist, D. J. Campbell, H. Borges da Silva, S. C. Jameson, Engagement of the costimulatory molecule ICOS in tissues promotes establishment of CD8+ tissue-resident memory T cells. *Immunity.* **55**, 98-114.e5 (2022).

40. H. M. Ren, E. M. Kolawole, M. Ren, G. Jin, C. S. Netherby-Winslow, Q. Wade, Shwetank, Z. S. M. Rahman, B. D. Evavold, A. E. Lukacher, IL-21 from high-affinity CD4 T cells drives differentiation of brain-resident CD8 T cells during persistent viral infection. *Sci. Immunol.* **5**, eabb5590 (2020).

41. R. Shen, Y.-B. Ouyang, C.-K. Qu, A. Alonso, L. Sperzel, T. Mustelin, M. Kaplan, G.-S. Feng, Grap negatively regulates T-cell receptor-elicited lymphocyte proliferation and interleukin-2 induction. *Mol. Cell. Biol.* **22**, 3230–3236 (2002).

42. C. E. Widjaja, J. G. Olvera, P. J. Metz, A. T. Phan, J. N. Savas, G. de Bruin, Y. Leestemaker, C. R. Berkers, A. de Jong, B. I. Florea, K. Fisch, J. Lopez, S. H. Kim, D. A. Garcia, S. Searles, J. D. Bui, A. N. Chang, J. R. Yates, A. W. Goldrath, H. S. Overkleeft, H. Ovaa, J. T. Chang, Proteasome activity regulates CD8+ T lymphocyte metabolism and fate specification. *J. Clin. Invest.* **127**, 3609–3623 (2017).

43. G. J. W. van der Windt, B. Everts, C.-H. Chang, J. D. Curtis, T. C. Freitas, E. Amiel, E. J. Pearce, E. L. Pearce, Mitochondrial Respiratory Capacity Is a Critical Regulator of CD8+ T Cell Memory Development. *Immunity*. **36**, 68–78 (2012).

44. V. Bergen, M. Lange, S. Peidli, F. A. Wolf, F. J. Theis, Generalizing RNA velocity to transient cell states through dynamical modeling. *Nat. Biotechnol.* **38**, 1408–1414 (2020).

45. M. Vogler, BCL2A1: the underdog in the BCL2 family. *Cell Death Differ.* **19**, 67–74 (2012).

46. A. Zaid, J. L. Hor, S. N. Christo, J. R. Groom, W. R. Heath, L. K. Mackay, S. N. Mueller, Chemokine Receptor–Dependent Control of Skin Tissue–Resident Memory T Cell Formation. *J. Immunol.* **199**, 2451–2459 (2017).

47. H. D. Hickman, G. V. Reynoso, B. F. Ngudiankama, S. S. Cush, J. Gibbs, J. R. Bennink, J. W. Yewdell, CXCR3 Chemokine Receptor Enables Local CD8+ T Cell Migration for the Destruction of Virus-Infected Cells. *Immunity*. **42**, 524–537 (2015).

48. H. Prizant, N. Patil, S. Negatu, N. Bala, A. McGurk, S. A. Leddon, A. Hughson, T. D. McRae, Y.-R. Gao, A. M. Livingstone, J. R. Groom, A. D. Luster, D. J. Fowell, CXCL10+ peripheral activation niches couple preferred sites of Th1 entry with optimal APC encounter. *Cell Rep.* **36**, 109523 (2021).

49. T. Hirai, Y. Zenke, Y. Yang, L. Bartholin, L. K. Beura, D. Masopust, D. H. Kaplan, Keratinocyte-Mediated Activation of the Cytokine TGF-β Maintains Skin Recirculating Memory CD8+ T Cells. *Immunity*. **50**, 1249-1261.e5 (2019).

50. T. Hirai, Y. Yang, Y. Zenke, H. Li, V. K. Chaudhri, J. S. De La Cruz Diaz, P. Y. Zhou, B. A.-T. Nguyen, L. Bartholin, C. J. Workman, D. W. Griggs, D. A. A. Vignali, H. Singh, D. Masopust, D. H. Kaplan, Competition for Active TGFβ Cytokine Allows for Selective Retention of Antigen-Specific Tissue- Resident Memory T Cells in the Epidermal Niche. *Immunity*. **54**, 84-98.e5 (2021).

51. R. Zeng, R. Spolski, S. E. Finkelstein, S. Oh, P. E. Kovanen, C. S. Hinrichs, C. A. Pise-Masison, M. F. Radonovich, J. N. Brady, N. P. Restifo, J. A. Berzofsky, W. J. Leonard, Synergy of IL-21 and IL-15 in regulating CD8+ T cell expansion and function. *J. Exp. Med.* **201**, 139–148 (2005).

52. S. Lesch, V. Blumenberg, S. Stoiber, A. Gottschlich, J. Ogonek, B. L. Cadilha, Z. Dantes, F. Rataj, K. Dorman, J. Lutz, C. H. Karches, C. Heise, M. Kurzay, B. M. Larimer, S. Grassmann, M. Rapp, A. Nottebrock, S. Kruger, N. Tokarew, P. Metzger, C. Hoerth, M.-R. Benmebarek, D. Dhoqina, R. Grünmeier, M. Seifert, A. Oener, Ö. Umut, S. Joaquina, L.

Vimeux, T. Tran, T. Hank, T. Baba, D. Huynh, R. T. A. Megens, K.-P. Janssen, M. Jastroch, D. Lamp, S. Ruehland, M. Di Pilato, J. N. Pruessmann, M. Thomas, C. Marr, S. Ormanns, A. Reischer, M. Hristov, E. Tartour, E. Donnadieu, S. Rothenfusser, P. Duewell, L. M. König, M. Schnurr, M. Subklewe, A. S. Liss, N. Halama, M. Reichert, T. R. Mempel, S. Endres, S. Kobold, T cells armed with C-X-C chemokine receptor type 6 enhance adoptive cell therapy for pancreatic tumours. *Nat. Biomed. Eng.* **5**, 1246–1260 (2021).

53. J. M. Schenkel, K. A. Fraser, K. A. Casey, L. K. Beura, K. E. Pauken, V. Vezys, D. Masopust, IL-15–Independent Maintenance of Tissue-Resident and Boosted Effector Memory CD8 T Cells. *J. Immunol.* **196**, 3920–3926 (2016).

54. B. Chandrasekar, S. Bysani, S. Mummidi, CXCL16 Signals via Gi, Phosphatidylinositol 3-Kinase, Akt, IκB Kinase, and Nuclear Factor-κB and Induces Cell-Cell Adhesion and Aortic Smooth Muscle Cell Proliferation. *J. Biol. Chem.* **279**, 3188–3196 (2004).

55. K. M. Knudson, C. J. Pritzl, V. Saxena, A. Altman, M. A. Daniels, E. Teixeiro, NFκB–Pim-1–Eomesodermin axis is critical for maintaining CD8 T-cell memory quality. *Proc. Natl. Acad. Sci.* **114** (2017), doi:10.1073/pnas.1608448114.

56. P.-M. Chen, P. C. Wilson, J. A. Shyer, M. Veselits, H. R. Steach, C. Cui, G. Moeckel, M. R. Clark, J. Craft, Kidney tissue hypoxia dictates T cell–mediated injury in murine lupus nephritis. *Sci. Transl. Med.* **12**, eaay1620 (2020).

57. M. A. Ataide, K. Knöpper, P. Cruz de Casas, M. Ugur, S. Eickhoff, M. Zou, H. Shaikh, A. Trivedi, A. Grafen, T. Yang, I. Prinz, K. Ohlsen, M. Gomez de Agüero, A. Beilhack, J. Huehn, M. Gaya, A.-E. Saliba, G. Gasteiger, W. Kastenmüller, Lymphatic migration of unconventional T cells promotes site-specific immunity in distinct lymph nodes. *Immunity.* **55**, 1813-1828.e9 (2022).

58. G. Liu, O. Abas, A. B. Strickland, Y. Chen, M. Shi, CXCR6+CD4+ T cells promote mortality during Trypanosoma brucei infection. *PLOS Pathog.* **17**, e1009968 (2021).

59. A. E. Oja, B. Piet, C. Helbig, R. Stark, D. van der Zwan, H. Blaauwgeers, E. B. M. Remmerswaal, D. Amsen, R. E. Jonkers, P. D. Moerland, M. A. Nolte, R. a. W. van Lier, P. Hombrink, Trigger-happy resident memory CD4+ T cells inhabit the human lungs. *Mucosal Immunol.* **11**, 654–667 (2018).

60. M. Tomura, N. Yoshida, J. Tanaka, S. Karasawa, Y. Miwa, A. Miyawaki, O. Kanagawa, Monitoring cellular movement in vivo with photoconvertible fluorescence protein "Kaede" transgenic mice. *Proc. Natl. Acad. Sci.* **105**, 10871–10876 (2008).

61. E. M. Steinert, J. M. Schenkel, K. A. Fraser, L. K. Beura, L. S. Manlove, B. Z. Igyártó, P. J. Southern, D. Masopust, Quantifying Memory CD8 T Cells Reveals Regionalization of Immunosurveillance. *Cell.* **161**, 737–749 (2015).

62. J. H. Levine, E. F. Simonds, S. C. Bendall, K. L. Davis, E. D. Amir, M. D. Tadmor, O. Litvin, H. G. Fienberg, A. Jager, E. R. Zunder, R. Finck, A. L. Gedman, I. Radtke, J. R.

Downing, D. Pe'er, G. P. Nolan, Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like Cells that Correlate with Prognosis. *Cell*. **162**, 184–197 (2015).

63. I. Tirosh, B. Izar, S. M. Prakadan, M. H. Wadsworth, D. Treacy, J. J. Trombetta, A. Rotem, C. Rodman, C. Lian, G. Murphy, M. Fallahi-Sichani, K. Dutton-Regester, J.-R. Lin, O. Cohen, P. Shah, D. Lu, A. S. Genshaft, T. K. Hughes, C. G. K. Ziegler, S. W. Kazer, A. Gaillard, K. E. Kolb, A.-C. Villani, C. M. Johannessen, A. Y. Andreev, E. M. Van Allen, M. Bertagnolli, P. K. Sorger, R. J. Sullivan, K. T. Flaherty, D. T. Frederick, J. Jané-Valbuena, C. H. Yoon, O. Rozenblatt-Rosen, A. K. Shalek, A. Regev, L. A. Garraway, Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science*. **352**, 189–196 (2016).

64. G. La Manno, R. Soldatov, A. Zeisel, E. Braun, H. Hochgerner, V. Petukhov, K. Lidschreiber, M. E. Kastriti, P. Lönnerberg, A. Furlan, J. Fan, L. E. Borm, Z. Liu, D. van Bruggen, J. Guo, X. He, R. Barker, E. Sundström, G. Castelo-Branco, P. Cramer, I. Adameyko, S. Linnarsson, P. V. Kharchenko, RNA velocity of single cells. *Nature*. **560**, 494–498 (2018).

Acknowledgments: The authors would like to acknowledge Austin Schultz and all other members of the Lund Lab for critical feedback and technical support.

Funding:

National Institutes of Health grant R01CA238163 (AWL)

National Institutes of Health grant T32AI100853 (TAH)

National Institutes of Health grant T32CA106195 (MMS)

Cancer Research Institute, Lloyd J. Old STAR Award (AWL)

American Cancer Society, RSG-18-169-01-LIB (AWL)

National Institutes of Health grant P30-CA016087 (Laura and Isaac Perlmutter Cancer Center supporting the Flow Cytometry and Cell Sorting Core and the Genome Technology Center (RRID: SCR_017929)

National Institutes of Health grant P30CA069533 (OHSU Knight Cancer Center supporting the Flow Cytometry and Cell Sorting Core)

Author contributions:

Conceptualization: TAH, AWL

Methodology: TAH, ZL

Investigation: TAH, ZL, MMS, TM

Funding acquisition: AWL

Supervision: AWL

Writing – original draft: TAH, AWL

Writing – review & editing: TAH, ZL, MMS, TM, AWL

Competing interests: AWL reports consulting services for AGS Therapeutics. All other authors declare that they have no competing interests.

Data and materials availability: scRNAseq data is deposited in GEO, GSE223727. All sequence data, code, and materials used in the analysis will be made publicly available upon manuscript acceptance.

Figure 1. CXCR6 expression is a conserved feature of resident CD8⁺ T cells after viral infection. (**A**) Representative flow of CD44⁺ vaccinia virus (VACV) specific (H2-K^{b}B8R$_{20-27}$) CD8⁺ T cells 28 days post infection (d.p.i.) (VACV, scarification) and expression of CD69, CD103, and CXCR6. (**B,C**) Quantification of CXCR6 geometric mean fluorescence intensity (GMFI) on CD44⁺CD8⁺ B8R-specific T cells from (A), as a function of location (B) and CD69 expression (C). (**D**) CXCR6 expression on CD45.1⁺ OT-1 T cells in non-lymphoid tissues 31 d.p.i. with vesicular stomatitis virus expressing OVA (VSV-OVA, intraperitoneal) (gated on CD8⁺CD45.1⁺CD44⁺CD69⁺, and IV⁻ in all tissues except liver and spleen) (SG = salivary gland and FRT = Female reproductive tract). (**E**) Representative flow plot and (**F**) quantification of CXCR6 GMFI on IV⁻CD45.1⁺ OT-1 T cells from kidney as a function of CD69 expression. (**A-F**) Data are representative of at least two experiments with n=4 each. Statistical significance determined with one way ANOVA (**B** and **D**), and paired t test (**C** and **F**). (**G**) UMAP representation of single-cell RNA-seq data from extravascular (IV⁻) OT-1 CD8⁺ T cells isolated out of skin 21 d.p.i.

with VACV expressing SIINFEKL (VACV-OVA). Circulating (T_{circ}) and resident (T_{RM}). (**H**) Scoring T_{RM} and T_{CIRC} with published gene signatures of circulating memory (left) or resident memory (right). (**I**) Expression of chemokine receptors in T_{CIRC} or T_{RM} clusters. (**J**) Scaled and normalized expression of *Cxcr6* in T_{CIRC} and T_{RM} clusters. (**G-J**) Data represent a total of 7409 cells across all clusters. (**H** and **J**) statistical significance determined using pairwise Wilcoxon rank test. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$

Figure 2. CXCR6 is required for optimal formation of T$_{RM}$ across non-lymphoid peripheral tissues. (**A**) Experimental design where 15,000 wildtype (WT) OT-1 T cells (CD45.1$^+$CD90.1$^-$) and 15,000 CXCR6$^{-/-}$ OT-1 T cells (CD45.1$^-$CD90.1$^+$) were co-transferred into naïve mice (CD45.1$^-$CD90.1$^-$) and the following day, ear skin was infected with vaccinia virus expressing OVA$_{257-264}$ (VACV-OVA, scarification). (**B**) Representative flow plots at least 49 days post infection (d.p.i.). Gated on CD8$^+$CD44$^+$IV$^-$ in the skin. (**C**) Quantification of the ratio of WT and CXCR6$^{-/-}$ OT-1 T cells isolated from the spleen and ear skin from (**B**). (**D** and **E**) Representative FACS analysis and quantification of CD69 (**D**) and CD103 (**E**). (**A-E**) Data are cumulative from 3 experiments for a total of n=12. (**F**) Experimental design schematic. Same as (**A**) but mice were infected with vesicular stomatitis virus expressing full length ovalbumin (VSV-OVA, intravenous). (**G**) Quantification of the ratio of WT and CXCR6$^{-/-}$ OT-1 T cells isolated 31 d.p.i.. Gated on CD8$^+$CD44$^+$IV$^-$ in kidney, FRT, brain and salivary gland; CD8$^+$CD44$^+$ in liver and spleen. Data are cumulative of two experiments for a total of n=8. Bars represent average + SD. Statistical significance was determined using paired student's t test (**C-E**) or one way ANOVA (**G**). * p ≤ 0.05, ** p ≤ 0.01, *** p ≤ 0.001, **** p ≤ 0.0001

Figure 3. T cell encounter with cognate antigen in skin drives CXCR6 upregulation. (**A**) Experimental design where 15,000 wildtype (WT) and 15,000 CXCR6$^{-/-}$ OT-1 T cells were co-transferred into naïve mice and the following day, skin was infected with vaccinia virus expressing OVA$_{257-264}$ (VACV-OVA, scarification). (**B**) Quantification of the ratio of WT and CXCR6$^{-/-}$ OT-1 T cells in spleen and skin and 5 and 14 days post infection (d.p.i). Data are cumulative from two experiments with n=4 (**C**) Expression of CXCR6 by WT OT-1 T cells in skin relative to spleen over time. Data are representative of at least two experiments with n=3 or 4. (**D**) Experimental design where OT-1 T cells were transferred to naïve mice, which were subsequently infected on the left ear VACV-OVA and the right with VACV (scarification). (**E**) Representative histogram and quantification (**F**) of CXCR6 by OT-1 T cells in spleen or ear skin 14 d.p.i.. Geometric mean fluorescence intensity (GMFI). Data are representative of at least two experiments with n=3 or 4. (**G** and **H**) Nurr77-GFP OT-1 T cells were transferred to naïve mice and the skin was infected with VACV-OVA the next day. (**G**) Representative histograms and (**H**) quantification of CXCR6 by Nurr77-GFP$^+$ and Nurr77-GFP$^-$ OT-1 T cells in skin 7 d.p.i.. Data are representative of two experiments with n=5. (**I**) Expression of CXCR6 by effector CD8$^+$ T cells from spleens of listeria monocytogenes (LM-OVA, 7 d.p.i) infected mice stimulated ex vivo with SIINFEKL and TGF-β. Bars represent average +SEM. Data are representative of two experiments with n=3. Cells in skin (**B,C,G** and **H**) are gated on IV-. Statistical significance determined using paired student's t test (**B,H**) or one way ANOVA (**C,F,I**). * p ≤ 0.05, ** p ≤ 0.01, *** p ≤ 0.001, **** p ≤ 0.0001

Figure 4. Dermal CXCL16 does not restrain T cell egress out of skin. (**A**) Quantification of CXCL16 in vaccinia virus (VACV) infected skin at various days post infection (d.p.i) by scarification; n=3-5 per group. (**B**) Representative immunofluorescence images of skin 14 d.p.i. with VACV. Scale bar = 100μm. (**C**) The percentage of OT-1 T cells in the epidermis after infection with $OVA_{257-264}$ -expressing VACV (VACV-OVA) calculated from immunofluorescent images. Data are cumulative of at least two experiments with n=5 (day 14) or n=6 (day 20 and 42-45). (**D**) Experimental design where 15,000 WT and 15,000 CXCR6$^{-/-}$ OT-1 T cells were transferred to K14-VEGFR3-Ig mice and the following day infected with VACV-OVA via skin scarification. (**E**) Quantification of the ratio of WT and CXCR6$^{-/-}$ OT-1 T cells in the spleen and skin of K14 VEGFR3-Ig mice 49 d.p.i. with VACV-OVA. Data are representative from at least two experiments with n=4. (**F**) Experimental design where congenically distinct 15,000 WT Kaede^{+} OT-1 T cells and 15,000 CXCR6$^{-/-}$ Kaede^{+} OT-1 T cells were co-transferred to C57Bl/6J mice and infected the following day with VACV-OVA via skin scarification. Kaede expressing cells in skin were photoconverted 6 d.p.i. and harvested 1 day later from skin and draining lymph node (dLN). (**G**) Representative flow of skin and dLN. (**H**) The ratio of photoconverted WT Kaede red^{+} to photoconverted CXCR6$^{-/-}$ Kaede red^{+} in skin and dLN following photoconversion. Data are representative from at least two experiments with n=5. Statistical significance determined using unpaired student's t test (**A,C**) or paired student's t test (**E,H**). Error bars represent SD. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$

Figure 5. CXCR6$^{-/-}$ T cells fail to fully acquire a residence transcriptional program and display signs of metabolic dysfunction and apoptosis. (**A**) UMAP showing integrated wildtype (WT) and CXCR6$^{-/-}$ OT-1 T cells isolated from skin 21 days post infection (d.p.i.) with vaccinia virus expressing OVA$_{257-264}$ (VACV-OVA). (**B**) Relative abundance of WT and CXCR6$^{-/-}$ in resident (T$_{RM}$), or circulating (T$_{CIRC}$) clusters. (**C**) Cells in T$_{CIRC}$ cluster and T$_{RM}$ clusters scored for published circulating memory (left) or resident memory (right) gene signatures. (**D**) Volcano plot of differentially expressed genes (DEG) between WT and CXCR6$^{-/-}$ cells in T$_{RM}$ clusters. (**E**) Pathway analysis of DEG between WT and CXCR6$^{-/-}$ cells in T$_{RM}$ clusters. p value calculated by hypergeometric distribution test. (**F**) MitoTracker deep red in WT and CXCR6$^{-/-}$ OT-1 T cells in skin 14 d.p.i.. Geometric mean fluorescence

intensity (GMFI). (**G**) Relative abundance of WT and CXCR6$^{-/-}$ cells in cluster 6. (**H**) Projected RNA velocity (scVelo) laid over UMAP. (I) Integrated (WT and CXCR6$^{-/-}$) clusters were scored for apoptosis signature (R-MMU-109581). (J) Expression of select apoptosis-related transcripts in integrated clusters (Bioplanet 2019). (**K-M**) WT and CXCR6$^{-/-}$ OT-1 T cells were isolated from VACV-OVA infected ear skin 21 d.p.i. (**K**) Representative flow and (**L**) quantification of BCL2 GMFI. (**M**) Ex vivo uptake of Zombie red viability dye. Statistical significance determined using paired student's t test (**F,K,L**) or pairwise Wilcoxon rank test (**C,I**). * $p \leq 0.05$, *** $p \leq 0.001$, **** $p \leq 0.0001$

Figure 6. CXCR6 promotes dermal T cell survival during memory formation in infected skin. (**A**) Representative flow plots and (**B**) quantification of ex vivo uptake of ZombieRed viability dye in wildtype (WT) and Cxcr6$^{-/-}$ OT-1 T cells in the dermis and epidermis 14 days post infection (d.p.i) with vaccinia virus expressing OVA$_{257-264}$ (VACV-OVA). Data are representative of at least two experiments with n=4. (**C**) Representative immunofluorescence image and (**D**) quantification of WT and CXCR6$^{-/-}$ (KO) OT-1 T cells in skin at 14 d.p.i. with VACV-OVA. (**E**) Representative immunofluorescence and (**F**) quantification of WT and CXCR6$^{-/-}$ OT-1 T cells in skin 42-45 d.p.i. with VACV-OVA. White line delineates the epidermis. Scale bar = 100μm. Data are cumulative of two experiments with n=5 (**C,D**) or n=6 (**E,F**). Statistical significance determined using paired student's t test. * p ≤ 0.05, *** p ≤ 0.001