Common roles for serotonin in rats and humans for computations underlying flexible decision-making

Qiang Luo[1,2,3,4,*,+], Jonathan W. Kanen[3,4,+], Andrea Bari[5], Nikolina Skandali[6,7,8], Christelle Langley[4,6], Gitte Moos Knudsen[9,10], Johan Alsiö[3,4], Benjamin U. Phillips[3,4], Barbara J. Sahakian[1,4,6], Rudolf N. Cardinal[6,11], Trevor W. Robbins[1,3,4,*]

[1]National Clinical Research Center for Aging and Medicine at Huashan Hospital, State Key Laboratory of Medical Neurobiology and Ministry of Education Frontiers Center for Brain Science, Institutes of Brain Science and Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai 200433, PR China

[2]Center for Computational Psychiatry, Ministry of Education Key Laboratory of Computational Neuroscience and Brain-Inspired Intelligence, Human Phenome Institute, Fudan University, Shanghai 200433, China

[3]Department of Psychology, University of Cambridge, Cambridge CB2 3EB, UK

[4]Behavioural and Clinical Neuroscience Institute, University of Cambridge, Cambridge CB2 3EB, UK

[5]Aelis Farma, 33077 Bordeaux, France

[6]Department of Psychiatry, University of Cambridge, Cambridge CB2 0SZ, UK

[7]Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, UK

[8]NIHR Biomedical Research Centre, University of Cambridge, UK

[9]Neurobiology Research Unit, the Neuroscience Centre, Copenhagen University Hospital Rigshospitalet, Denmark

[10]Faculty of Health and Medical Sciences, University of Copenhagen, Denmark

[11]Liaison Psychiatry Service, Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge CB2 0QQ, UK

[*]Correspondence to: qluo@fudan.edu.cn or twr2@cam.ac.uk

[+]joint first author

Abstract

Serotonin is critical for adapting behavior flexibly to meet changing environmental demands. Cognitive flexibility is important both for successful attainment of goals, as well as for social interactions, and is frequently impaired in neuropsychiatric disorders, including obsessive-compulsive disorder (OCD). However, a unifying mechanistic framework accounting for the role of serotonin in behavioral flexibility has remained elusive. Here, we demonstrate common effects of manipulating serotonin function across two species (rats and humans) on latent processes supporting choice behavior during probabilistic reversal learning using computational modelling. The findings support a role of serotonin in behavioral flexibility and plasticity, indicated, respectively, by increases or decreases in choice repetition ('stickiness') or reinforcement learning rates depending upon manipulations intended to increase or decrease serotonin function. More specifically, the rate at which expected value increased following reward and decreased following punishment (reward and punishment 'learning rates') was greatest after sub-chronic administration of the selective serotonin reuptake (SSRI) citalopram (5 mg/kg for 7 days followed by 10 mg/kg twice a day for 5 days) in rats. Conversely, humans given a single dose of an SSRI (20mg escitalopram), which can decrease post-synaptic serotonin signalling, and rats that received the neurotoxin 5,7-dihydroxytryptamine (5,7-DHT), which destroys forebrain serotonergic neurons, exhibited decreased reward learning rates. A basic perseverative tendency ('stickiness'), or choice repetition irrespective of the outcome produced, was likewise increased in rats after the 12-day SSRI regimen and decreased after single dose SSRI in humans and 5,7-DHT in rats. These common effects of serotonergic manipulations on rats and humans – identified via computational modelling – suggest an evolutionarily conserved role for serotonin in plasticity and behavioral flexibility and have clinical relevance transdiagnostically for neuropsychiatric disorders.

Introduction

Humans and other animals alike must maximise rewards and minimise punishments to survive and thrive. Across phylogeny this involves learning about cues or locations that inform whether an action is likely to result in a good or bad outcome. Adaptive behavior, however, must also be flexible: the ability to disengage from previously learned actions that are no longer useful or appropriate to the situation is fundamental to well-being. Indeed, behavior can become abnormally stimulus-bound and perseverative in compulsive disorders [1-5]. Furthermore, learning the best course of action can require withstanding occasional negative feedback, which should sometimes be ignored if rare. Indeed, inappropriately switching behavior away from an adaptive action following misleading or even negative feedback ('lose-shift') has been reported across several traditional psychiatric diagnostic categories [6-10].

The neurotransmitter serotonin (5-hydroxytryptamine; 5-HT) is widely implicated in behavioral flexibility[11-18]. Perturbing 5-HT function can affect both perseveration and lose-shift behavior, which are commonly assessed using probabilistic reversal learning (PRL) paradigms (Figure 1 A-B): a subject learns through trial and error the most adaptive action in a choice procedure, the contingencies of which eventually reverse, sometimes repeatedly [12, 19-21]. A unifying framework for 5-HT in these processes has, however, remained elusive. To this end, we proposed to use a mechanistic modelling framework to align behavioral changes in PRL following serotonergic manipulations in rats [19] and humans [22].

Reinforcement learning (RL) is a well-established computational mechanism for the analysis of latent mechanisms underlying choice behavior as it unfolds dynamically over time [23].

Standard RL models typically conceptualise choice in relation to an action's value, derived

from an accumulated reinforcement history, and incorporate parameters that estimate how

quickly action values are learned ('learning rate') and the extent to which that value is acted

upon (often termed 'inverse temperature' in relation to the mathematical softmax function

typically used; here, termed 'reinforcement sensitivity') [24]. Stickiness parameters, by contrast,

track the extent to which behavioral tendencies are shaped by engagement with discrete cues

(stimuli) or locations, irrespective of an action's outcome. Stickiness can therefore be

considered a value-free component of behavior [25, 26]. Across six previously published

experiments in rats and humans and a recently published computational modelling study in

humans, we examined whether stickiness or other RL parameters (learning rates or

reinforcement sensitivity) contributed meaningfully to behavior, and examined whether 5-HT

function would consistently modulate any of these parameters across species.

The stickiness parameter has recently emerged as important for understanding compulsivity:

stickiness was significantly high in stimulant use disorder (SUD) but abnormally low in

obsessive–compulsive disorder (OCD) during PRL performance [7]. Meanwhile, value-free

influences have been notably absent from prominent computational accounts of goal-directed

(or 'model-based') versus habitual (or 'model-free') controllers of behavior [26]. These have

traditionally revolved around environmental features relevant to outcomes [27, 28]. This has

hindered contextualisation within the rich literature on the neural basis of habits (reinforcer-

independent perseveration) [29]. A traditional view of stimulus–response habits is that they are

created and strengthened by reinforcement, acting to enhance direct links between

environmental stimuli and responses [30]; they are thus "model-free" in that they do not involve

representations of the expected consequences of behavior, but are "value-based" in that they

are created by valenced reinforcement. However, there are other aspects of behavior that are

independent of reinforcement or value. Indeed, value-free (action outcome-irrelevant) factors

similar to stickiness were recently shown to be important for understanding goal-directed

decision-making [28]. Accounting for stickiness – value-free perseveration – may therefore aid

in better dissecting the nature of imbalanced goal-directed versus habitual behavior seen in

OCD, SUD, and other conditions [31-33], a balance that is sensitive to serotonergic disruption in

humans and rodents [34-36].

Two common methods for studying serotonin are through serotonin depletion and treatment

with selective serotonin reuptake inhibitors (SSRIs). In non-human animals, depletion can be

achieved using the neurotoxin 5,7-dihydroxytryptamine (5,7-DHT) which produces a

profound loss of serotonergic fibers [37]. SSRIs, meanwhile, are first-line pharmacological

treatments for several psychiatric conditions including major depressive disorder (MDD) [38],

anxiety disorders [39], post-traumatic stress disorder (PTSD) [40], and OCD [41], yet both the

computational and neural mechanisms underlying their efficacy remain poorly understood.

SSRIs block the 5-HT transporter and thus reuptake of 5-HT, which increases extracellular

serotonin levels; however, this occurs not only in projection areas but also in the vicinity of 5-

HT_{1A} somatodendritic autoreceptors, activation of which leads to decreased firing rates of 5-

HT neurons [42]. SSRIs can thus paradoxically lower 5-HT concentrations in projection regions

when given acutely, especially at low doses [43], and firing rates return to baseline after $5-HT_{1A}$

autoreceptors are desensitised by repeated administration [44]. This mechanism might be

reflected in a delayed clinical onset of the treatment effect of SSRI on mood [45]. For this

reason, effects of both acute and chronic SSRIs in rats were studied, with the prediction that a

higher acute dose and a chronic use could overcome these feedback effects of a low acute

dose and produce an increase in serotonin transmission [19]. The 20mg used in the acute study

with healthy humans [22], while within the therapeutic range, is a lower acute dose than used in

some experimental animal studies.

Here, the primary question was whether serotonergic manipulations would cause similar

perturbations of model parameters across both rats and humans, thereby demonstrating the

evolutionary significance of the role of serotonin in cognitive flexibility. As an increased

tendency for lose-shift behavior induced by acute SSRI has been conceptualised as

hypersensitivity to negative feedback [19, 22], we asked whether this would be reflected in

elevated punishment learning rates. Selective 5-HT depletion via 5,7-DHT of the orbitofrontal

cortex (OFC) or amygdala in marmoset monkeys, meanwhile, reduced reinforcement learning

rates (for rewards or punishments), and modulated stickiness [46]; we hypothesised that changes

in learning rate or stickiness parameters would occur following global 5-HT manipulations in

rats and humans. We predicted that incorporating stickiness parameters would be central to

capturing effects of 5-HT on behavioral flexibility and would increase or decrease depending

on changes in serotonin transmission.

Materials and Methods

Probabilistic reversal learning task: humans

The task used in the human SSRI experiment [22] is shown in Figure 1A, and contained 80

trials: 40 during acquisition and 40 following reversal. In other words, there was a fixed

number of trials and a single reversal. For the first 40 trials, one option yielded positive

feedback on 80% of trials, the other option on 20% of trials. These contingencies reversed for

the latter 40 trials. Positive feedback was given in the form of the word "CORRECT" on the

touchscreen computer and a high tone, negative feedback was conveyed by the word

"WRONG" and a low tone. The task was self-paced.

Probabilistic reversal learning task: rats

Following training and determination of stable levels of accuracy and a lack of side bias [19] in

operant chambers controlled by the Whisker control system [47], rats were presented with two

apertures illuminated simultaneously to the left and right of a central (inactive) aperture

(Figure 1B). Responding at the 'correct' location was associated with an 80% probability food

reward (and 20% probability of a time-out punishment), whereas responding at the 'incorrect'

location yielded reward on only 20% of trials (and punishment on 80%). Reward was in the

form of a 45 mg food pellet (Noyes dustless pellets; Sandown Scientific, Middlesex, UK)

delivered to a food magazine positioned on the opposite wall of the operant chamber.

Punishment was given in the form of a 2.5-second time-out. The left and right apertures were

illuminated for 30 seconds signifying the response window. The next trial was triggered by

retrieval of the pellet from the magazine. If no response was made, the trial was categorised as

an omission and resulted in a 5-second time-out. Responding to an unlit aperture had no

programmed consequence. Reversals occurred after the animal made eight consecutive correct

responses, at which point the correct aperture became the incorrect aperture and vice versa. A

session consisted of 200 trials to be completed during a 40-minute period. One session was

conducted per day.

5,7-DHT forebrain 5-HT depletion: rats

Sixteen rats were included in the final analysis. Rats were pre-treated intraperitoneally (i.p.)

with 20 mg/kg of desipramine hydrochloride (Sigma, Poole, UK) in order to preserve

noradrenergic neurons. Half of the rats were randomly assigned to receive bilateral

intracerebroventricular (i.c.v.) infusions of 80 μg 5,7-DHT creatinine sulfate diluted in 10 μg

of 10% ascorbic acid in saline, guided by a stereotaxic frame, whilst the other half received a

sham infusion of 10 µg 0.01 M phosphate-buffered saline (PBS) – vehicle [19]. Post-mortem

neurochemistry confirmed that 5,7-DHT infusions produced a near-total depletion of brain

serotonin and decreased levels of the serotonin metabolite 5-hydroxyindoleacetic acid (5-

HIAA) relative to controls in all regions examined: OFC, prelimbic cortex, anterior cingulate

cortex, nucleus accumbens, dorsomedial striatum, dorsolateral striatum, amygdala, dorsal

hippocampus (all $p<.05$)[19]. Levels of dopamine, norepinephrine, and the dopamine metabolite

dihydroxyphenylacetic acid (DOPAC) were not significantly different from controls in any of

these regions (all $p > .05$)[19]. Data were analysed from seven consecutive sessions conducted

following surgery in the previous report [19]. Computational model convergence was achieved

when modelling behavior from all seven sessions collectively, which is reported in the current

study. Conversely, computational model convergence could not be achieved when modelling

the seven sessions separately.

SSRI administration: rats

Animals were divided into groups matched for task accuracy and then randomly assigned via

a Latin square design to receive injections i.p. of either citalopram hydrobromide (1 mg/kg or

10 mg/kg; Tocris, Bristol, UK). Citalopram, dissolved in 0.01 M PBS, or vehicle was

administered 30 minutes before the task [19]. Eleven rats were included in the final analysis

after receiving vehicle, 1 mg/kg, or 10 mg/kg citalopram [19]. Fourteen rats were included in the

repeated and sub-chronic citalopram experiment. The citalopram group was administered 5

mg/kg citalopram 30 min before testing, for seven consecutive days (*n*=7). The vehicle group

(*n*=7), instead, received the same number of daily injections of 0.01 M phosphate-buffered

saline [19]. After seven days, the citalopram group received 10 mg/kg of citalopram twice a day

(about 4 h before the testing) for five consecutive days, to study the long-lasting effects of

sub-chronic dosing [19].

SSRI administration: humans

The protocol was ethically approved (Cambridge Central NHS Research Ethics Committee,

reference 15/EE/0004). Volunteers gave informed consent and were paid. Participants were

healthy and without a personal or family history of psychiatric or neurological disorders [22]. In

a randomised, double-blind, placebo-controlled, between-groups design [22], healthy volunteers

received either escitalopram (n=32) or placebo (n=33). The PRL task was conducted

following a 3-hour waiting period after oral drug administration to attain peak plasma

escitalopram concentration [48]. Plasma analysis (n=59) verified increased escitalopram

concentration [22] at 2.5 hours after the dose (t_{54} = 18.835, p < 0.001, mean = 14 ng/ml, standard

deviation [SD] = 5.72) just before the task administration, and at 5.5 hours (t_{54} = 20.548, p <

0.001, mean = 17.24 ng/ml, SD = 4.27). Mood ratings were unaffected by single dose

escitalopram administration (p > .05). There were no differences between groups in age, sex,

years of education, depressive symptoms, or trait anxiety (all p > .05).

Computational modelling of behavior

Overview

These methods are based on Kanen *et al.* [7]. Four RL models were fitted to the behavioral data,

which incorporated parameters that have been studied previously using a hierarchical

Bayesian method[7, 49]. Models were fitted via Hamiltonian Markov chain Monte Carlo

sampling implemented in Stan 2.17.2[50]. Convergence was checked according to $\hat{R}$, the

potential scale reduction factor measure[51, 52], which approaches 1 for perfect convergence.

Values below 1.1 are typically used as a guideline for determining model convergence and 1.1

as a stringent criterion [51]. In the current study, most of the models had an $\hat{R} < 1.1$, except for

Model 4 in the sub-chronic 10 mg/kg experiment in rats ($\hat{R} = 1.7$) and Model 1 in the 5,7-

DHT experiment in rats ($\hat{R} = 1.5$). We assumed the four models examined had the same prior

probability (0.25). Models were compared via a bridge sampling estimate of the likelihood [53],

using the "bridgesampling" package in R [54]. Bridge sampling directly estimates the marginal

likelihood, and therefore the posterior probability of each model given the data (and prior

model probabilities), under the assumption that the models represent the entire group of those

to be considered. Posterior distributions were interpreted using the highest density interval

(HDI) of posterior distributions, which is the Bayesian "credible interval", at different

significance levels including 75%, 80%, 85%, 90% and 95%. Together with the HDI, the

group mean difference (MD) was also reported. The priors used for each parameter are shown

in Supplemental Table 1. For the human experiments, trials were sequenced across all 80

trials of the PRL task, and on each trial the computational model was supplied with the

participant's identification number and condition, whether the trial resulted in positive or

negative feedback, and which visual stimulus was selected. For the rat experiments, trials

were sequenced across all sessions conducted under a given manipulation, and the

computational model was supplied with the same information, but instead with the location of

the aperture selected rather than the identification of the stimulus selected. Omissions were

rare and they were not included in the computational analysis.

Models

Model 1 incorporated three parameters and was used to test the hypothesis that 5-HT would

affect how positive versus negative feedback guides behavior. Separate learning rates for

positive feedback (reward) α^{rew} and negative feedback (nonreward/punishment) α^{pun} were

implemented. Positive reinforcement led to an increase in the value V_i of the stimulus i that

was chosen, at a speed governed by the *reward learning rate* α^{rew}, via $V_{i,t+1} \leftarrow V_{i,t} + \alpha^{rew}(R_t -$

$V_{i,t})$. R_t represents the outcome on trial t (defined as 1 on trials where positive feedback

occurred), and $(R_t - V_{i,t})$ the prediction error. On trials where negative feedback occurred $R_t =$

0, which led to a decrease in value of V_i at a speed governed by the *punishment learning rate*

α^{pun}, according to $V_{i,t+1} \leftarrow V_{i,t} + \alpha^{pun}(R_t - V_{i,t})$. Stimulus value was incorporated into the final

quantity controlling choice according to $Q^{reinf}_t = \tau^{reinf}V_t$. The additional parameter τ^{reinf}, termed

reinforcement sensitivity, governs the degree to which behavior is driven by reinforcement

history. The quantities Q associated with the two available choices, for a given trial, were then

input to a standard softmax choice function to compute the probability of each choice:

264
$$P(\text{action}_a) = \text{softmax}^a_\beta(Q_1 \ldots Q_n) = \frac{e^{\beta Q_a}}{\Sigma^n_{k=1} e^{\beta Q_k}},$$

for $n=2$ choice options. The probability values for each trial emerging from the softmax

function (*i.e.*, the probability of choosing stimulus 1) were fitted to the subject's actual

choices (*i.e.*, did the subject choose stimulus 1?). Softmax inverse temperature was set to $\beta =$

1, and as a result the reinforcement sensitivity parameter (τ^{reinf}) directly represented the weight

given to the exponents in the softmax function.

Model 2 was as model 1 but for the human experiments incorporated a "stimulus stickiness"

parameter τ^{stim}, which measures the tendency to repeat a response to a specific perceptual

stimulus, irrespective of the action's outcome. For the rat experiments a "side (location)

stickiness" parameter τ^{loc} was substituted, which measures the tendency to repeat a response

to a specific aperture in the operant chamber. Incorporating these two different stickiness

parameters, depending on the species, accounts for task differences between the human and

rat PRL experiments. This four-parameter model served to test whether accounting for

stimulus-response learning, in addition to learning about action-outcome associations, would

best characterise behavior. The stimulus stickiness effect was modelled as $Q^{stim}_t = \tau^{stim} s_{t-1}$,

where s_{t-1} was 1 for a stimulus that was chosen on the previous trial and was otherwise 0. The

final quantity controlling choice incorporated this additional parameter as $Q_t = Q^{reinf}_t + Q^{stim}_t$.

Quantities Q, corresponding to the two choice options on a given trial, were then fed into the

softmax function as above.

Model 3 incorporated three parameters and served to test whether a single learning rate α^{reinf},

rather than separate learning rates for rewards and punishments, optimally characterised

behavior. Reward led to an increase in the value V_i of the stimulus i that was chosen, at a

speed controlled by the *reinforcement rate* α^{reinf}, via $V_{i,t+1} \leftarrow V_{i,t} + \alpha^{reinf}(R_t - V_{i,t})$. R_t represents

the outcome on trial t (defined as 1 on trials where reward occurred), and $(R_t - V_{i,t})$ the

prediction error. On trials where punishment occurred $R_t = 0$, which led to a decrease in value

of V_i. Model 3 also included the stimulus stickiness parameter. The final quantity controlling

choice was determined by $Q_t = Q^{reinf}_t + Q^{stim}_t$.

Model 4 took a different approach, and had three parameters: φ (phi), ρ (rho), and β (beta).

Derived from the experienced-weighted attraction model (EWA) of Camerer and Ho [55], here

it was implemented as in den Ouden *et al.* [14] a study in which the EWA model best described

behavior best on a nearly identical human task. A key difference to the other reinforcement

learning models tested in this study is that here the learning rate can decline over time,

governed by a decay factor ρ (rho). The EWA model weighs the value of new information

against current expectations or beliefs, accumulated from previous experience.

Learning from reinforcement is modulated by an "experience weight", $n_{c,t}$, which is a measure

of how often the subject has chosen a stimulus (*i.e.* experienced the action), and is updated

every time the stimulus is chosen (where c is choice and t is trial) according to the experience

decay factor ρ (range $0 < \rho < 1$) and can increase without bounds [14]:

$$n_{c,t} \leftarrow n_{c,t-1}\, \rho + 1.$$

The value of a choice is updated according to the outcome, λ, and the decay factor for

previous payoffs, φ (range $0 < \varphi < 1$) [14]

$$v_{c,t} \leftarrow (v_{c,t-1}\, \varphi\, n_{c,t-1} + \lambda_{t-1}) / n_{c,t}.$$

The payoff decay factor φ (phi) is related to a Rescorla–Wagner-style [56] learning rate α (as in

Models 1-3), by $\alpha = 1 - \varphi$. A high value of φ means that stimuli keep a high fraction of their

previous value and thus learning from reinforcement is slow. When ρ is high, then "well-

known" actions (with high n) are updated relatively little by reinforcement, by virtue of the

terms involving n, whilst reinforcement has a proportionately larger effect on novel actions

(with low n). For comparison to Models 1-3, when $\rho = 0$, the experience weight n, is 1, which

reduces to a learning rate α controlling the influence of learning from prediction error. Choice

in the EWA model is also governed by a softmax process, only here the softmax inverse

temperature β was also a parameter able to vary, in contrast to Models 1-3.

320   Results

321   Choice of model

Behavior in all experiments was best described by reinforcement learning models

incorporating parameters for stickiness, reinforcement sensitivity, and learning rates,

consistent with previous work [7, 49]. Convergence was good with most models having $\hat{R} < 1.1$

(see Methods). Model comparison metrics are shown in Supplemental Table 2. For all

experiments, the winning model had separate learning rates for reward (α^{rew}) and punishment

(α^{pun}). The reward learning rate (α^{rew}) indexed how quickly action value representation

increased following a reward prediction error (when action outcome was better than

predicted). Punishment learning rate (α^{pun}) is an assay of the speed at which action value

decreased following a punishment prediction error (outcome was worse than predicted).

Stickiness measures a basic perseverative tendency: whether or not an action chosen on the

previous trial was repeated, irrespective of its outcome. For rats, stickiness indexed the side

(or location; τ^{loc}) of responding whereas for humans, stickiness referred to (visual) stimulus

stickiness (τ^{stim}). Reinforcement sensitivity (τ^{reinf}) measures the degree to which the values

learned through reinforcement impact on choice behavior. Reinforcement sensitivity can be

viewed as a value-based inverse temperature; stickiness as a value-free inverse temperature.

Low values of stickiness or reinforcement sensitivity can be thought of as two different types

of exploratory behavior; low reinforcement sensitivity represents exploration away from the

more highly valued choice whereas low stickiness represents exploration away from the

previously chosen stimulus or location irrespective of value. The accuracy of the parameter

recovery was confirmed for this modelling approach previously [7] and also confirmed by

simulations for those parameter values estimated here in each experiment (Supplementary

Table 3).

Serotonin depletion by intraventricular 5,7 dihydroxytryptamine (5,7-DHT): rats

Results are shown in Figure 1C and Table 1. Post-mortem neurochemistry confirmed that 5,7-

DHT infusions produced a near-total depletion of brain serotonin (for more details see the

Methods and also Bari *et al.* 2010). The conventional analysis in the previous publication [19]

found a decreased win-stay rate, an increased lose-shift rate and a reduced number of

reversals completed in the group of depletion-operated rats ($n = 8$) compared with the group

of sham-operated rats ($n = 8$). After computational modelling, we found that the depletion

decreased the side (location) stickiness parameter (τ^{loc}; MD = -0.2938 [95% HDI, -0.4635 to -

0.1134]) and the reward learning rate (α^{rew}; MD = -0.0401 [85% HDI, -0.0757 to -0.0033]).

There was no effect of 5,7-DHT on the punishment learning rate (α^{pun}) or reinforcement

sensitivity (τ^{reinf}) [0 ∈ 75% HDI]. The decreased lose-shift rate was retrodicted in the

simulation of the computational model (Supplementary Result 1). Furthermore, because

reinforcement sensitivity was also unaffected in Model 1, which did not contain the stickiness

parameter, the effect of 5,7-DHT on stickiness was unlikely to be a misattribution of

reinforcement sensitivity.

Acute SSRI: rats

Results for acute citalopram administered to rats (n = 11 with a cross-over design for vehicle,

1mg/kg, and 10mg/kg) are shown in Figure 2 and Table 1. The conventional analysis showed

the number of reversals completed was significantly lower following a low dose of 1 mg/kg

SSRI compared with a high dose of 10 mg/kg SSRI [19]. After computational modelling of the

behavior, we found a single dose of 1 mg/kg citalopram in rats diminished the side (location)

stickiness parameter (MD = -0.1862 [95% HDI, -0.3330 to -0.0441]), as seen following 5,7-

DHT. The reward learning rate was enhanced by the 1 mg/kg dose in rats (MD = 0.2098 [95%

HDI, 0.0184 to 0.3959]). There was no effect of 1 mg/kg on the punishment learning rate or

reinforcement sensitivity (0 ∈ 75% HDI). A single high dose of citalopram in rats (10 mg/kg)

decreased the reward learning rate (MD = -0.1489 [85% HDI, -0.2888 to -0.0009]) and

enhanced reinforcement sensitivity (MD = 0.2900 [85% HDI, 0.0346 to 0.5590]). However,

there was no effect of 10 mg/kg on the punishment learning rate or side (location) stickiness

(0 ∈ 75% HDI). Simulation of the wining model retrodicted the significant difference in the

number of reversals completed between the low-dose group and the high-dose group

(Supplementary Result 1).

Repeated and sub-chronic SSRI: rats

Results for 'repeated' 5 mg/kg citalopram administered for consecutive 7 days to rats (the Cit

group; $n = 7$) compared with the vehicle group (the Veh group; $n = 7$) are shown in Figure 3A

and Table 1. After 7 days, the Cit group received 10 mg/kg of citalopram twice a day for 5

consecutive days to study the longer-lasting effects of 'sub-chronic' dosing. Results for sub-

chronic dosing are shown in Figure 3B and Table 1. The conventional analyses showed the

win-stay rate increased by repeated citalopram treatment and the number of reversals was

increased by sub-chronic dosing [19]. Following computational modelling of the behavior, we

found that repeated citalopram enhanced both the punishment learning rate (MD = 0.3299 [95%

HDI, 0.0432 to 0.6404]) and side (location) stickiness (MD = 0.1581 [75% HDI, 0.0135 to

0.3054]). There was no effect of repeated citalopram on the reward learning rate and

reinforcement sensitivity (0 ∈ 75% HDI). The sub-chronic dosing enhanced the reward

learning rate (MD = 0.4769 [95% HDI, 0.2699 to 0.6780]), the punishment learning rate (MD

= 0.4762 [95% HDI, 0.2172 to 0.7323]), and the side (location) stickiness (MD = 0.1676 [75%

HDI, 0.0075 to 0.3414]), but decreased the reinforcement sensitivity (MD = -0.9972 [95%

HDI, -1.7233 to -0.2540]). Simulation of the winning model retrodicted the significant

increase of the win-stay rate for repeated citalopram compared with the vehicle, but did not

show a significant increase in the number of reversals for sub-chronic dosing (Supplementary

Result 1).

Acute SSRI: humans

Modelling results ($n = 32$ escitalopram, $n = 33$ placebo) are shown in Figure 4 and Table 1.

The prior conventional analysis suggested that the impaired reversal learning after acute SSRI

mainly resulted from an elevated lose-shift rate [22]. After computational modelling, we found

that the administration of a single 20 mg dose of escitalopram to healthy humans decreased

the reward learning rate (MD = -0.2019 [95% HDI, -0.3612 to -0.0392]), stimulus stickiness

(MD = -0.1841 [85% HDI, -0.3476 to -0.0045]) and reinforcement sensitivity (MD = -1.6848

[80% HDI, -3.1501 to -0.1553]), but had no effect on the punishment learning rate ($0 \in 75\%$

HDI). Simulation of the computational model retrodicted a significantly increased lose-shift

rate (Supplementary Result 1).

Chronic SSRI treatment in humans

As reported in our recent publication for the effect of chronic use of SSRI on behavioral

flexibility by a double-blind, placebo-control, semi-randomized study [57], the computational

modelling approach was applied to the behavioral data of the same probabilistic reversal

learning task in healthy volunteers. The participants were semi-randomized into the treatment

group ($n = 32$) receiving 20 mg escitalopram or the control group receiving the placebo for 3

to 5 weeks. The conventional analysis identified no significant group differences[57]. After

computational modelling, we found that the chronic use of SSRI reduced reinforcement

sensitivity compared to placebo ($n = 34$) in healthy volunteers (MD = -2.7673 [90% HDI,

−5.2846 to −0.3959]), but had no effect on reward/punishment learning rates or stimulus

stickiness ($0 \in 75\%$ HDI) [57].

Relationship between model parameters and conventional behavioral measures

Next, we conducted correlational analyses to demonstrate how our modelling results compared

with traditional metrics of PRL. There were converging effects across species involving

stickiness. Results were corrected for multiple comparisons by false discovery rate (FDR) and

are summarised in Supplemental Tables 5-7. The conventional measures examined for the rat

experiments were win-stay (proportion of trials where the subject stayed with the same choice

following a reward), lose-shift (proportion of trials where the subject shifted choice following

punishment), and number of reversals completed [19]. Win-stay and lose-shift were also

examined in the human studies, as was perseveration [18]. In the human SSRI acute experiment,

stimulus stickiness was positively correlated with the win-stay rate (r = .51, p = .0066 on

placebo; r = .62, p = .0005 following escitalopram) and also negatively correlated with the

lose-shift rate (r = -.63, p = .0003 on placebo; r = -.78, p = 7.95×10^{-7} following escitalopram).

In rats, side (location) stickiness was negatively correlated with the lose-shift rate following an

acute 1 mg/kg dose of citalopram (r = -.89, p = .006), and positively correlated with the win-

stay rate in the vehicle group with daily injections of 0.01 M phosphate-buffered saline for 7

days (r = .95, p = .0065). Side (location) stickiness was also positively correlated with the

number of reversals achieved during the repeated administration (r = .89, p = .0205 following

5 mg/kg citalopram per day and r = .97, p = .0049 with the same number of daily injections of

vehicle). Further correlations with other model parameters are reported in the Supplementary

Tables 5-7.

Summary of results

In rats, stickiness was decreased after 5,7-DHT and acute 1 mg/kg citalopram, whereas

stickiness was increased after repeated 5 mg/kg citalopram and sub-chronic 10 mg/kg

citalopram. In humans, stickiness was decreased following 20 mg escitalopram, similar to the

effects of 5,7-DHT and low dose citalopram in rats. Also in cross-species alignment, the

reward learning rate was decreased following 5,7-DHT and acute 10 mg/kg citalopram in rats

as well as in humans following 20 mg escitalopram. The reward learning rate in rats was

additionally increased following acute 1 mg/kg citalopram and sub-chronic 10mg/kg

citalopram. The punishment learning rate was increased for both repeated 5 mg/kg citalopram

and sub-chronic citalopram in rats only. Reinforcement sensitivity was increased following 10

mg/kg of citalopram and decreased during sub-chronic treatment in rats, agreeing with our

own recent analysis of chronic escitalopram treatment in humans [57], although this parameter

was also shown to be decreased in the present analysis following acute 20mg escitalopram in

humans.

Discussion

We have demonstrated converging effects of a range of bidirectional 5-HT manipulations

across both rats and humans which bolsters its evolutionarily conserved role in behavioral

flexibility and plasticity. Computational modelling of choice behavior indicated increases or

decreases in choice repetition ('stickiness') or reinforcement learning rates depending upon

manipulations intended to increase or decrease serotonin function, respectively. Stickiness, a

basic tendency to persevere versus 'explore', was modulated in five serotonergic

manipulations examined across both rats and humans. Stickiness was decreased by neurotoxic

5-HT depletion in rats and by acute 1 mg/kg SSRI in rats (citalopram) and healthy humans

(20 mg escitalopram), treatments presumably reducing 5-HT signalling. By contrast,

stickiness was increased following both repeated (5 mg/kg for 7 days) and sub-chronic (10

mg/kg twice a day for 5 days) dosing of SSRI in rats, treatments probably boosting 5-HT

function. Learning rates were also modulated by five serotonergic manipulations across

species. The reward learning rate increased the most after sub-chronic administration of the

SSRI citalopram (5 mg/kg for 7 days followed by 10 mg/kg twice a day for 5 days) compared

with the vehicle group. Conversely, humans given a single dose of an SSRI (20mg

escitalopram), which can decrease post-synaptic serotonin signalling, and rats that received

5,7-DHT demonstrated decreased reward learning rates. This in turn parallels the reduction of

reinforcement learning rates following 5,7-DHT infused directly in the marmoset amygdala or

OFC to produce local 5-HT depletion [46]. Collectively, the present and the previous results

show that serotonin has common effects on latent computational mechanisms supporting

flexible decision-making and plasticity in rats, marmoset monkeys and humans.

The neural substrates of PRL are relatively well understood [46, 58, 59] and involve interactions in

particular among the orbitofrontal cortex (OFC), amygdala, and striatum. Administration of

5,7-DHT directly to either the marmoset OFC or amygdala produced changes in both

stickiness and reinforcement learning rates [46]. Marmosets that received 5,7-DHT in the OFC

repeated choices to recently chosen stimuli across a longer timescale, whereas 5,7-DHT in the

amygdala produced a more ephemeral tendency to repeat choices [46]. Dietary depletion of

tryptophan, serotonin's biosynthetic precursor, in humans, also modulated stickiness and

corresponding activity in frontopolar cortex during a four-choice probabilistic task [60].

Stickiness, the only value-free parameter in our reinforcement learning model, contributed to

a core feature of complex behavior, *i.e.* exploration. Lower stickiness, even negative

stickiness, is generally associated with more exploratory behavior. However, exploratory

behavior is not a unitary construct [61]. At one level, exploratory behavior can reflect directed

information gathering, but on another level it can be mechanistic or rigid, resulting from

'decisional noise', producing apparently flexible behavior but, in fact, representing a

fundamental performance heuristic recruited in volatile settings that evokes a primitive form

of exploration. Another potential measure of exploratory behavior is reflected in

reinforcement sensitivity, as a value-based parameter in our model, which can be interpreted

as reflecting the balance between exploiting and exploring tendencies (low reinforcement

sensitivity is sometimes referred to as 'random exploration') [62].

Whilst the effects of serotonin on reinforcement sensitivity revealed by the present analyses

were ostensibly more difficult to interpret – underscoring that stickiness is a distinct

mechanism – there is an intriguing parallel with a recent study. Langley *et al.* [57] have recently

shown diminished reinforcement sensitivity in healthy humans following chronic – at least 21

days – of 20 mg escitalopram performing the same PRL task and modelled in an identical

fashion – this reduction is hence the same direction as for the acute dose in humans and sub-

chronic dosing in rats. Although this parallel between single and chronic dosing in humans

was unexpected, it is notable that reinforcement sensitivity in rats following sub-chronic

dosing was also decreased. These effects of reduced reinforcement sensitivity (value-based)

may relate to what has been termed "emotional blunting" or "SSRI-induced apathy

syndrome" in patients with MDD [57, 63-65]. The reduction in inverse temperature can also be

interpreted as a reduction in "maximisation" of reinforcement and this a shift in the balance

between "exploitation" and "exploration" [61]. However, it is evident that this drift to

exploration is not always accompanied by reduced "stickiness", suggesting different processes

underlying choice variability.

The present analyses focusing on behavioral flexibility are relevant to current hypotheses of

effects of psychedelic agents such as psilocybin and LSD and their hypothetical actions on

neuronal plasticity and cognitive flexibility [66, 67]. There are in fact intriguing parallels between

the present global manipulations of serotonin and the effects of LSD on latent mechanisms

underlying PRL in humans. Whilst LSD is mostly known for its 5-HT$_{2A}$ agonist properties, it

is also a 5-HT$_{1A}$ agonist and suppresses dorsal raphe serotonin neuron activity [68]. Indeed, LSD

was recently shown to reduce stickiness during PRL performance of healthy humans [69, 70],

which aligns with 5-HT$_{1A}$ somatodendritic autoreceptor effects associated with the reduced

stickiness shown here following acute SSRI in humans and low dose SSRI in rats. At the

same time, LSD markedly increased the reinforcement learning rates for both reward and

punishment [70], which were also increased following sub-chronic SSRI dosing in rats. The

parallel with our sub-chronic SSRI results from rats with the effects of LSD on learning rates

in humans agrees with the literature showing that optogenetic stimulation of 5-HT neurons in

the dorsal raphe increased reinforcement learning rates[71]. Given the well-established role of

the $5\text{-}HT_{2A}$ receptor in reversal learning, and its involvement in SSRI-related reversal

improvements [72], a $5\text{-}HT_{2A}$ mechanism may well be implicated in the present data. Indeed, the

$5\text{-}HT_{2A}$ receptor is involved in plasticity [73, 74] and associative learning [75]. Furthermore, during

initial learning (pre-reversal), LSD decreased reinforcement sensitivity [70], in line with the

acute and chronic [57] SSRI effects in humans and sub-chronic effect in rats.

Other studies have investigated other forms of exploratory behavior, sometimes assessed with

a four-choice, rather than two-choice, task as here. For example, directed exploration – where

the goal is to explore uncertain options to maximise information gained – was modulated by

dopamine [76] and attenuated in gambling disorder [77]. *Tabula rasa* exploration (disregarding

history), meanwhile, ignores all prior knowledge (*e.g.* choice history, reinforcement history,

and estimates of uncertainty, respectively), has been associated with norepinephrine but not

dopamine function [78] and may be enhanced in individuals with attention-deficit/hyperactivity

disorder (ADHD) symptoms [79]. Understanding distinct types of exploratory behavior and their

neurochemical modulation is therefore relevant transdiagnostically. We posit that low

stickiness is a fundamental form of exploration, and have shown here that serotonin

modulates it; this is likely by affecting a neural network that includes the dorsomedial PFC,

OFC, and amygdala [46].

Manifestation of high or low stickiness may bear on the neural representation of discrete

states of the world. In the context of PRL, for example, one state would be "option A is

mostly correct" (pre-reversal) whilst another state would be "option B is mostly correct"

(post-reversal). To perform well during PRL, in this view, veridical state representations

inferred by the brain are critical as are veridical probabilities of transitions between states.

Indeed, the OFC is implicated in representing states [80, 81]. One possibility, therefore, is that

these results concerning stickiness collectively reflect an influence of serotonin on inferring

states or state transitions. This would align with recent theorising on OCD (where stickiness is

low during PRL) [7], which posits that the disorder can be characterised by excessive statistical

uncertainty (variance, or inverse precision) about the probability of transitions between states

(*e.g.* from the state of dirty hands to clean hands after washing), particularly those that are

action-dependent [82]. The optimal response to uncertainty about the current state would be

exploratory behavior to continue gathering information [82]. SUD (where stickiness is high) [7],

meanwhile, may be characterised by over-encoding of state-specific rules and information [83].

The model of state transition uncertainty can explain excessive behavioral switching (*i.e.* low

stickiness) as well as heightened perseveration (*i.e.* high stickiness) and can be extended to

account for other conditions including generalised anxiety disorder, autism spectrum disorder

(ASD), and schizophrenia [82]. Indeed, reversal learning deficits have been documented in ASD

[6] and schizophrenia [84, 85].

Dose-dependent effects of SSRIs are key to understanding serotonin function in this cross-

species analysis. Acute low- and high-dose SSRI administration lowered and increased

stickiness, respectively, which likely reflected sensitive measures of opposite effects on 5-HT

activity. Evidence from positron emission tomography (PET) imaging has shown that acute

SSRI in humans, at the dose used here, lowers 5-HT concentrations in projection regions [86],

although there can be considerable individual differences in this action[87] - which may relate to

the considerable variability in the reinforcement sensitivity parameter evident in Figure 4. The

reduction in 5-HT levels in terminal projection areas is believed to reflect the activation of 5-

HT_{1A} autoreceptors by increases in extracellular serotonin following reuptake inhibition,

which in turn leads to decreased firing rates of 5-HT neurons [42, 44]. We posit that the high

acute dose of SSRI used in rats, which heightened stickiness, overcame $5-HT_{1A}$ autoreceptor-

mediated regulation.

The dose-dependent effects on stickiness may have implications for the treatment of OCD, in

particular, one of numerous conditions for which SSRIs are first-line pharmacotherapy [38-41].

One puzzle has been why doses up to three times higher than those used in MDD are optimal

for reducing symptoms of OCD [88]. In fact, guidelines for OCD recommend titrating to the

maximum approved dose [89], yet using these high doses in MDD does not improve efficacy

and instead increases side-effects [88]. That both the repeated 5 mg/kg SSRI and the sub-chronic

10 mg/kg treatments in rats increased stickiness in the present study may be relevant for

understanding this clinical phenomenon.

Conclusion

It is imperative to overcome the challenge of relating animal and human experiments in order

to advance models of psychiatric disorder and drug development [90-92]. Here, we have provided

evidence across rats and humans that serotonin modulates fundamental components of

learning important for plasticity (reinforcement learning rates) and behavioral flexibility

(stickiness), bidirectionally. Stickiness, a basic perseverative tendency less commonly studied

in conjunction with RL, may be a fundamental mechanism involved in choice. Moreover, we

have shown a consistent role for serotonin in affecting basic tendencies to persevere or

explore in comparable decision-making tasks in rats and humans. These results demonstrate

that the role of serotonin in cognitive flexibility is preserved across species and are thus of

evolutionary significance. In addition, this role of serotonin is of clinical relevance for

neuropsychiatric disorders where SSRIs are the first line of treatment. The translational

results of this study are of particular relevance for the pathophysiology and treatment of OCD

and SUD, where parallel learning processes have been perturbed [7], and have implications for

a wide range of other neuropsychiatric disorders, including depression [8, 9] and schizophrenia

[27, 93].

Competing Interests Statement

T.W.R. discloses consultancy with Cambridge Cognition and Supernus; he receives research grants from Shionogi & Co and Sirgartan and editorial honoraria from Springer Verlag and Elsevier. B.J.S. consults for Cambridge Cognition and receives royalties from PopReach. R.N.C. consults for Campden Instruments and receives royalties from Cambridge Enterprise, Routledge, and Cambridge University Press. All other authors declare no conflicts of interest.

Acknowledgements

This research was funded by the Wellcome Trust (Grant 104631/Z/14/Z to T.W.R) and the Lundbeck Foundation (Grant R281-2018-131 to B.J.S and G.M.K). Q.L. was supported by the National Key Research and Development Program of China (Grant 2019YFA0709502), the National Natural Science Foundation of China (Grant 82272079), the Natural Science Foundation of Shanghai (Grant 20ZR1404900), the Shanghai Municipal Science and Technology Major Project (Grant 2018SHZDZX01). Most of the analyses of this study had been conducted when Q.L. was a Visiting Fellow at the Clare Hall, University of Cambridge, Cambridge, UK. J.W.K. was supported by a Gates Cambridge Scholarship and an Angharad Dodds John Bursary in Mental Health and Neuropsychiatry. R.N.C was funded by the UK Medical Research Council (MC_PC_17213). N.S. was supported by an Academic Clinical Fellowship (University of Cambridge/ Cambridgeshire and Peterborough NHS Foundation Trust). B.U.P was supported by an Angharad Dodds John Bursary in Mental Health and Neuropsychiatry and is a current employee of AstraZeneca plc.

Author Contributions
TWR, QL and JWK made substantial contributions to the conception or design of the work; AB, NS and CL contributed substantially to the acquisition of the data; QL, JWK, JA, BUP and RNC contributed substantially to the analysis of the data; QL, JWK, GMK, BJS, RNC and TWR contributed substantially to the interpretation of data; JK and QL wrote the first draft; AB, NS, CL, GMK, JA, BUP, BJS, RNC and TWR made critical revisions.

Reference

639
640
1.   Berlin, G.S. & Hollander, E. Compulsivity, impulsivity, and the DSM-5 process. *Cns*
Spectrums **19**, 62-68 (2014).
2.   Jentsch, J.D. & Taylor, J.R. Impulsivity resulting from frontostriatal dysfunction in drug
abuse: implications for the control of behavior by reward-related stimuli.
Psychopharmacology **146**, 373-390 (1999).
3.   Koob, G.F. & Volkow, N.D. Neurobiology of addiction: a neurocircuitry analysis.
Lancet Psychiatry **3**, 760-773 (2016).
4.   Robbins, T.W., Vaghi, M.M. & Banca, P. Obsessive-Compulsive Disorder: Puzzles and
Prospects. *Neuron* **102**, 27-47 (2019).
5.   Tiffany, S.T. A COGNITIVE MODEL OF DRUG URGES AND DRUG-USE BEHAVIOR - ROLE
OF AUTOMATIC AND NONAUTOMATIC PROCESSES. *Psychological Review* **97**, 147-168 (1990).
6.   Brolsma, S.C.A.*, et al.* Challenging the negative learning bias hypothesis of
depression: reversal learning in a naturalistic psychiatric sample. *Psychological Medicine* **52**,
303-313 (2022).
7.   Kanen, J.W., Ersche, K.D., Fineberg, N.A., Robbins, T.W. & Cardinal, R.N.
Computational modelling reveals contrasting effects on reinforcement learning and cognitive
flexibility in stimulant use disorder and obsessive-compulsive disorder: remediating effects
of dopaminergic D2/3 receptor agents. *Psychopharmacology* **236**, 2337-2358 (2019).
8.   Mukherjee, D., Filipowicz, A.L.S., Vo, K., Satterthwaite, T.D. & Kable, J.W. Reward and
Punishment Reversal-Learning in Major Depressive Disorder. *Journal of Abnormal Psychology*
129, 810-823 (2020).
9.   Murphy, F.C., Michael, A., Robbins, T.W. & Sahakian, B.J. Neuropsychological
impairment in patients with major depressive disorder: the effects of feedback on task
performance. *Psychological Medicine* **33**, 455-467 (2003).
10.   Taylor Tavares, J.V.*, et al.* Neural basis of abnormal response to negative feedback in
unmedicated mood disorders. *Neuroimage* **42**, 1118-1126 (2008).
11.   Alsio, J.*, et al.* Serotonergic Innervations of the Orbitofrontal and Medial-prefrontal
Cortices are Differentially Involved in Visual Discrimination and Reversal Learning in Rats.
Cerebral Cortex **31**, 1090-1105 (2021).
12.   Brown, H.D., Amodeo, D.A., Sweeney, J.A. & Ragozzino, M.E. The selective serotonin
reuptake inhibitor, escitalopram, enhances inhibition of prepotent responding and spatial
reversal learning. *Journal of Psychopharmacology* **26**, 1443-1455 (2012).
13.   Clarke, H.F., Dalley, J.W., Crofts, H.S., Robbins, T.W. & Roberts, A.C. Cognitive
inflexibility after prefrontal serotonin depletion. *Science* **304**, 878-880 (2004).
14.   den Ouden, H.E.M.*, et al.* Dissociable Effects of Dopamine and Serotonin on Reversal
Learning. *Neuron* **80**, 1090-1100 (2013).
15.   Lapiz-Bluhm, M.D.S., Soto-Pina, A.E., Hensler, J.G. & Morilak, D.A. Chronic
intermittent cold stress and serotonin depletion induce deficits of reversal learning in an
attentional set-shifting test in rats. *Psychopharmacology* **202**, 329-341 (2009).
16.   Matias, S., Lottem, E., Dugue, G.P. & Mainen, Z.F. Activity patterns of serotonin
neurons underlying cognitive flexibility. *Elife* **6** (2017).
17.   Kanen, J.W.*, et al.* Effect of lysergic acid diethylamide (LSD) on reinforcement
learning in humans. *bioRxiv*, 2020.2012.2004.412189 (2021).

18.   Kanen, J.W.*, et al.* Probabilistic reversal learning under acute tryptophan depletion in
healthy humans: a conventional analysis. *Journal of Psychopharmacology* **34**, 580-583
(2020).
19.   Bari, A.*, et al.* Serotonin Modulates Sensitivity to Reward and Negative Feedback in a
Probabilistic Reversal Learning Task in Rats. *Neuropsychopharmacology* **35**, 1290-1301
(2010).
20.   Chamberlain, S.R.*, et al.* Neurochemical modulation of response inhibition and
probabilistic learning in humans. *Science* **311**, 861-863 (2006).
21.   Lawrence, A.D., Sahakian, B.J., Rogers, R.D., Hodges, J.R. & Robbins, T.W.
Discrimination, reversal, and shift learning in Huntington's disease: mechanisms of impaired
response selection. *Neuropsychologia* **37**, 1359-1374 (1999).
22.   Skandali, N.*, et al.* Dissociable effects of acute SSRI (escitalopram) on executive,
learning and emotional functions in healthy humans. *Neuropsychopharmacology* **43**, 2645-
2651 (2018).
23.   Daw, N.D. Trial-by-trial data analysis using computational models. *Decis Making,*
Affect Learn Atten Perform **XXIII**, 1–26 (2011).
24.   Sutton, R.S. & Barto, A.G. Reinforcement Learning: An Introduction, 2nd edn. *The MIT*
Press (2012).
25.   Bennett, D., Niv, Y. & Langdon, A.J. Value-free reinforcement learning: policy
optimization as a minimal model of operant behavior. *Current Opinion in Behavioral Sciences*
41, 114-121 (2021).
26.   Miller, K.J., Shenhav, A. & Ludvig, E.A. Habits Without Values. *Psychological Review*
126, 292-311 (2019).
27.   Deserno, L. & Hauser, T.U. Beyond a Cognitive Dichotomy: Can Multiple Decision
Systems Prove Useful to Distinguish Compulsive and Impulsive Symptom Dimensions? *Biol*
Psychiatry **88**, e49-e51 (2020).
28.   Shahar, N.*, et al.* Credit assignment to state-independent task representations and its
relationship with model-based decision making. *Proceedings of the National Academy of*
Sciences of the United States of America **116**, 15871-15876 (2019).
29.   Balleine, B.W. & O'Doherty, J.P. Human and Rodent Homologies in Action Control:
Corticostriatal Determinants of Goal-Directed and Habitual Action.
Neuropsychopharmacology **35**, 48-69 (2010).
30.   Cardinal, R.N., Parkinson, J.A., Hall, J. & Everitt, B.J. Emotion and motivation: the role
of the amygdala, ventral striatum, and prefrontal cortex. *Neuroscience and Biobehavioral*
Reviews **26**, 321-352 (2002).
31.   Ersche, K.D.*, et al.* Carrots and sticks fail to change behavior in cocaine addiction.
Science **352**, 1468-1471 (2016).
32.   Gillan, C.M.*, et al.* Disruption in the Balance Between Goal-Directed Behavior and
Habit Learning in Obsessive-Compulsive Disorder. *American Journal of Psychiatry* **168**, 718-
726 (2011).
33.   Voon, V.*, et al.* Disorders of compulsivity: a common bias towards learning habits.
Molecular Psychiatry **20**, 345-352 (2015).
34.   Ohmura, Y.*, et al.* Disruption of model-based decision making by silencing of
serotonin neurons in the dorsal raphe nucleus. *Current Biology* **31**, 2446-+ (2021).
35.   Worbe, Y.*, et al.* Valence-dependent influence of serotonin depletion on model-
based choice strategy. *Molecular Psychiatry* **21**, 624-629 (2016).

36.   Worbe, Y., Savulich, G., de Wit, S., Fernandez-Egea, E. & Robbins, T.W. Tryptophan
Depletion Promotes Habitual over Goal-Directed Control of Appetitive Responding in
Humans. *Int J Neuropsychopharmacol* **18**, pyv013 (2015).
37.   Bjorklund, A., Baumgarten, H.G. & Rensch, A. 5,7-DIHYDROXYTRYPTAMINE -
IMPROVEMENT OF ITS SELECTIVITY FOR SEROTONIN NEURONS IN CNS BY PRETREATMENT
WITH DESIPRAMINE. *Journal of Neurochemistry* **24**, 833-835 (1975).
38.   Gelenberg AJ, F.M., Markowitz JC, et al. Practice guideline for the treatment of
patients with major depressive disorder. *American Psychiatric Association* (2010).
39.   Baldwin, D.S.*, et al.* Evidence-based pharmacological treatment of anxiety disorders,
post-traumatic stress disorder and obsessive-compulsive disorder: A revision of the 2005
guidelines from the British Association for Psychopharmacology. *Journal of*
Psychopharmacology **28**, 403-439 (2014).
40.   Benedek DM, F.M., Zatzick D, Ursano RJ. Practice guideline for the treatment of
patients with acute stress disorder and posttraumatic stress disorder. *Am Psychiatr Assoc*
(2009).
41.   Fineberg NA, D.L., Reid J, et al. Management and treatment of OCD. In: Geddes JR,
Andreasen NC, Goodwin GM (eds) New Oxford Textbook of Psychiatry, 3rd edn. *Oxford*
University Press, Oxford, England (2020).
42.   Richardson-Jones, J.W.*, et al.* 5-HT1A Autoreceptor Levels Determine Vulnerability to
Stress and Response to Antidepressants. *Neuron* **65**, 40-52 (2010).
43.   Nord, M., Finnema, S.J., Halldin, C. & Farde, L. Effect of a single dose of escitalopram
on serotonin concentration in the non-human and human primate brain. *International*
Journal of Neuropsychopharmacology **16**, 1577-1586 (2013).
44.   Fischer, A.G., Jocham, G. & Ullsperger, M. Dual serotonergic signals: a key to
understanding paradoxical effects? *Trends Cogn Sci* (2014).
45.   Taylor, M.J., Freemantle, N., Geddes, J.R. & Bhagwagar, Z. Early onset of selective
serotonin reuptake inhibitor antidepressant action - Systematic review and meta-analysis.
Archives of General Psychiatry **63**, 1217-1223 (2006).
46.   Rygula, R.*, et al.* Role of Central Serotonin in Anticipation of Rewarding and Punishing
Outcomes: Effects of Selective Amygdala or Orbitofrontal 5-HT Depletion. *Cerebral Cortex*
25, 3064-3076 (2015).
47.   Cardinal, R.N. & Aitken, M.R.F. Whisker: A client-server high-performance multimedia
research control system. *Behavior Research Methods* **42**, 1059-1071 (2010).
48.   Rao, N. The clinical pharmacokinetics of escitalopram. *Clinical Pharmacokinetics* **46**,
281-290 (2007).
49.   Alsio, J.*, et al.* Dopamine D2-like receptor stimulation blocks negative feedback in
visual and spatial reversal learning in the rat: behavioural and computational evidence.
Psychopharmacology **236**, 2307-2323 (2019).
50.   Carpenter, B.*, et al.* Stan: A Probabilistic Programming Language. *Journal of Statistical*
Software **76**, 1-29 (2017).
51.   Brooks, S.P. & Gelman, A. General methods for monitoring convergence of iterative
simulations. *Journal of Computational and Graphical Statistics* **7**, 434-455 (1998).
52.   Gelman, A., Hill, J. & Yajima, M. Why We (Usually) Don't Have to Worry About
Multiple Comparisons. *Journal of Research on Educational Effectiveness* **5**, 189-211 (2012).
53.   Gronau, Q.F.*, et al.* A tutorial on bridge sampling. *Journal of Mathematical*
Psychology **81**, 80-97 (2017).

54.   Gronau, Q.F., Singmann, H. & Wagenmakers, E.-J. bridgesampling: An R Package for
Estimating Normalizing Constants. *Journal of Statistical Software* **92**, 1-29 (2020).
55.   Camerer, C. & Ho, T.H. Experience-weighted attraction learning in normal form
games. *Econometrica* **67**, 827-874 (1999).
56.   Rescorla, R.A. & Wagner, A.R. A theory of classical conditioning: Variations in the
effectiveness of reinforcement and nonreinforcement. In: Black AH, Prokasy WF (eds)
Classical Conditioning II Current Research and Theory. *Appleton-Century-Crofts* (1972).
57.   Langley, C.*, et al.* Chronic escitalopram in healthy volunteers has specific effects on
reinforcement sensitivity: a double-blind, placebo-controlled semi-randomised study.
Neuropsychopharmacology (2023).
58.   O'Doherty, J., Kringelbach, M.L., Rolls, E.T., Hornak, J. & Andrews, C. Abstract reward
and punishment representations in the human orbitofrontal cortex. *Nature Neuroscience* **4**,
95-102 (2001).
59.   Cools, R., Clark, L., Owen, A.M. & Robbins, T.W. Defining the Neural Mechanisms of
Probabilistic Reversal Learning Using Event-Related Functional Magnetic Resonance Imaging.
The Journal of Neuroscience **22**, 4563 (2002).
60.   Seymour, B., Daw, N.D., Roiser, J.P., Dayan, P. & Dolan, R. Serotonin Selectively
Modulates Reward Value in Human Decision-Making. *Journal of Neuroscience* **32**, 5833-5842
(2012).
61.   Wilson, R.C., Geana, A., White, J.M., Ludvig, E.A. & Cohen, J.D. Humans Use Directed
and Random Exploration to Solve the Explore-Exploit Dilemma. *Journal of Experimental
Psychology-General* **143**, 2074-2081 (2014).
62.   Gershman, S.J. Deconstructing the human algorithms for exploration. *Cognition* **173**,
34-42 (2018).
63.   Barnhart, W.J., Makela, E.H. & Latocha, M.J. SSRI-induced apathy syndrome: a clinical
review. *J Psychiatr Pract* **10**, 196-199 (2004).
64.   Price, J., Cole, V. & Goodwin, G.M. Emotional side-effects of selective serotonin
reuptake inhibitors: qualitative study. *Br J Psychiatry* **195**, 211-217 (2009).
65.   Marazziti, D.*, et al.* Emotional Blunting, Cognitive Impairment, Bone Fractures, and
Bleeding as Possible Side Effects of Long-Term Use of SSRIs. *Clin Neuropsychiatry* **16**, 75-85
(2019).
66.   Carhart-Harris, R.L. & Friston, K.J. REBUS and the Anarchic Brain: Toward a Unified
Model of the Brain Action of Psychedelics. *Pharmacol Rev* **71**, 316-344 (2019).
67.   Vollenweider, F.X. & Preller, K.H. Psychedelic drugs: neurobiology and potential for
treatment of psychiatric disorders. *Nature Reviews Neuroscience* **21**, 611-624 (2020).
68.   Nichols, D.E. Hallucinogens. *Pharmacology & Therapeutics* **101**, 131-181 (2004).
69.   Kanen, J.W.*, et al.* Probabilistic reversal learning under acute tryptophan depletion in
healthy humans: a conventional analysis. *J Psychopharmacol* **34**, 580-583 (2020).
70.   Kanen, J.W.*, et al.* Effect of lysergic acid diethylamide (LSD) on reinforcement
learning in humans. *Psychol Med*, 1-12 (2022).
71.   Iigaya, K., Fonseca, M.S., Murakami, M., Mainen, Z.F. & Dayan, P. An effect of
serotonergic stimulation on learning rates for rewards apparent after long intertrial
intervals. *Nature Communications* **9**, 2477 (2018).
72.   Furr, A., Lapiz-Bluhm, M.D. & Morilak, D.A. 5-HT2A receptors in the orbitofrontal
cortex facilitate reversal learning and contribute to the beneficial cognitive effects of chronic
citalopram treatment in rats. *The international journal of neuropsychopharmacology* **15**,
1295-1305 (2012).

73.   Barre, A., *et al.* Presynaptic serotonin 2A receptors modulate thalamocortical
plasticity and associative learning. *Proc Natl Acad Sci U S A* **113**, E1382-1391 (2016).
74.   Vaidya, V.A., Marek, G.J., Aghajanian, G.K. & Duman, R.S. 5-HT2A receptor-mediated
regulation of brain-derived neurotrophic factor mRNA in the hippocampus and the
neocortex. *J Neurosci* **17**, 2785-2795 (1997).
75.   Harvey, J.A. Role of the serotonin 5-HT(2A) receptor in learning. *Learn Mem* **10**, 355-
362 (2003).
76.   Chakroun, K., Mathar, D., Wiehler, A., Ganzer, F. & Peters, J. Dopaminergic
modulation of the exploration/exploitation trade-off in human decision-making. *Elife* **9**
(2020).
77.   Wiehler, A., Chakroun, K. & Peters, J. Attenuated Directed Exploration during
Reinforcement Learning in Gambling Disorder. *Journal of Neuroscience* **41**, 2512-2522
(2021).
78.   Dubois, M., *et al.* Human complex exploration strategies are enriched by
noradrenaline-modulated heuristics. *Elife* **10** (2021).
79.   Dubois M, B.A., Moses-Payne M, et al. Tabula-rasa exploration decreases during
youth and is linked to ADHD symptoms. *bioRxiv* (2020).
80.   Schuck, N.W., Cai, M.B., Wilson, R.C. & Niv, Y. Human Orbitofrontal Cortex
Represents a Cognitive Map of State Space. *Neuron* **91**, 1402-1412 (2016).
81.   Wilson, R.C., Takahashi, Y.K., Schoenbaum, G. & Niv, Y. Orbitofrontal Cortex as a
Cognitive Map of Task Space. *Neuron* **81**, 267-279 (2014).
82.   Fradkin, I., Adams, R.A., Parr, T., Roiser, J.P. & Huppert, J.D. Searching for an Anchor
in an Unpredictable World: A Computational Model of Obsessive Compulsive Disorder.
Psychological Review **127**, 672-699 (2020).
83.   Mueller, L.E., Sharpe, M.J., Stalnaker, T.A., Wikenheiser, A.M. & Schoenbaump, G.
Prior Cocaine Use Alters the Normal Evolution of Information Coding in Striatal Ensembles
during Value-Guided Decision-Making. *Journal of Neuroscience* **41**, 342-353 (2021).
84.   Deserno L, B.R., Mathys C, et al. Volatility Estimates Increase Choice Switchingand
Relate to Prefrontal Activity in Schizophrenia. *Biol Psychiatry Cogn Neurosci Neuroimaging*
(2020).
85.   Waltz, J.A. & Gold, J.M. Probabilistic reversal learning impairments in schizophrenia:
Further evidence of orbitofrontal dysfunction. *Schizophrenia Research* **93**, 296-303 (2007).
86.   Nord, M., Finnema, S.J., Halldin, C. & Farde, L. Effect of a single dose of escitalopram
on serotonin concentration in the non-human and human primate brain. *Int J*
Neuropsychopharmacol **16**, 1577-1586 (2013).
87.   da Cunha-Bang, S., *et al.* Measuring endogenous changes in serotonergic
neurotransmission with [11C]Cimbi-36 positron emission tomography in humans.
Translational Psychiatry **9**, 134 (2019).
88.   Derksen, M., Feenstra, M., Willuhn, I. & Denys, D. The serotonergic system in
obsessive-compulsive disorder. in *Handbook of the Behavioral Neurobiology of Serotonin,*
2nd Edition (ed. C.P. Muller & K.A. Cunningham) 865-891 (2020).
89.   Koran, L.M., Hanna, G.L., Hollander, E., Nestadt, G. & Simpson, H.B. Practice guideline
for the treatment of patients with obsessive-compulsive disorder. *American Journal of*
Psychiatry **164**, 5-53 (2007).
90.   Benzina, N., N'Diaye, K., Pelissolo, A., Mallet, L. & Burguiere, E. A cross-species
assessment of behavioral flexibility in compulsive disorders. *Communications Biology* **4**
(2021).

91.   Robbins, T.W. & Cardinal, R.N. Computational psychopharmacology: a translational
and pragmatic approach. *Psychopharmacology* **236**, 2295-2305 (2019).
92.   Robble, M.A.*, et al.* Concordant neurophysiological signatures of cognitive control in
humans and rats. *Neuropsychopharmacology* **46**, 1252-1262 (2021).
93.   Frith, C.D. & Done, D.J. STEREOTYPED RESPONDING BY SCHIZOPHRENIC-PATIENTS ON
A 2-CHOICE GUESSING TASK. *Psychological Medicine* **13**, 779-786 (1983).

Tables and Figure Captions

Table 1. Summary of learning parameter effects.

	Stickiness τ^{stim} (humans) τ^{loc} (rats)	Reward learning rate α^{rew}	Punishment learning rate α^{pun}	Reinf. Sensitivity τ^{reinf}
Rats: neurotoxic depletion of 5-HT	↓***	↓*	–	–
Rats: 1 mg/kg citalopram	↓**	↑**	–	–
Humans: 20 mg escitalopram	↓*	↓**	–	↓..
Rats: 10 mg/kg citalopram	–	↓*	–	↑*
Rats: 5mg/kg citalopram chronic	↑.	–	↑***	–
Rats: 10mg/kg citalopram sub-chronic	↑.	↑***	↑***	↓***
Humans: 20 mg escitalopram chronic [57]	–	–	–	↓**

rew reward, *pun* punishment, *reinf* reinforcement, *stim* stimulus, *loc* location

*** stands for 0 ∉ 95% HDI, ** for 0 ∉ 90% HDI, * for 0 ∉ 85% HDI, **..** for 0 ∉ 80% HDI,

. for 0 ∉ 75% HDI

Figure 1. Task schematics for probabilistic reversal learning and effects of serotonin depletion on model parameters in rats.

A) Experiment in humans (example trial on touchscreen computer) and **B)** Experiment in rats (two apertures illuminated simultaneously to the left and right of a central aperture with reinforcement contingencies 80% : 20% for left : right or right : left, and a food pellet was given to a food magazine positioned on the opposite wall of the operant chamber if the rewarding location was chosen). **C)** Side (location) stickiness was diminished by neurotoxic 5-HT depletion, *i.e.*, 5,7- dihydroxytryptamine. Reinf. = reinforcement. Red signifies a difference between the parameter per-condition mean according to the Bayesian "credible interval", 0 ∉ 95% HDI. Blue signifies a significance by the 85% HDI. The inner interval represents the 85% HDI, while the outer interval represents the 95% HDI.

Figure 2. Effects of acute SSRI (citalopram) at two doses on model parameters in rats.

A) for 1 mg/kg and **B**) for 10 mg/kg. Reinf. = reinforcement. mg/kg = milligrams per kilogram. Red signifies a difference between the parameter per-condition mean according to the Bayesian "credible interval", $0 \notin 95\%$ HDI. Blue signifies a significance by the 85% HDI. The inner interval stands for the 90% HDI in A), and 85% HDI in B), while the outer interval represents the 95% HDI.

Figure 3. Effects of repeated and sub-chronic SSRI on model parameters in rats.

A) for the repeated SSRI (5 mg/kg citalopram) experiment, and **B**) for the sub-chronic SSRI (10 mg/kg citalopram) experiment. Reinf. = reinforcement. Red signifies a difference between the parameter per-condition mean according to the Bayesian "credible interval", $0 \notin 95\%$ HDI, and orange signifies a significance by the 75% HDI. All outer intervals represent the 95% HDI. The inner intervals represent the 75% HDI for side stickiness and the 90% HDI for the other 3 parameters.

Figure 4. Effects of acute SSRI (20 mg escitalopram) on model parameters in humans.

Stimulus stickiness was decreased following acute SSRI. Reinf. = reinforcement. Red signifies a difference between the parameter per-condition mean according to the Bayesian "credible interval", $0 \notin 95\%$ HDI. Similarly, blue and purple signify the significance levels by 85% and 80% HDI's, respectively. All outer intervals represent the 95% HDI.