Research Article

De novo fatty-acid synthesis protects invariant NKT cells from cell death, thereby promoting their homeostasis and pathogenic roles in airway hyperresponsiveness

Jaemoon Koh[1,2,*], Yeon Duk Woo[2,*], Hyun Jung Yoo[3], Jun-Pyo Choi[4], Sae Hoon Kim[4,5], Yoon-Seok Chang[4,5], Kyeong Cheon Jung[1], Ji Hyung Kim[3], Yoon Kyung Jeon[1], Hye Young Kim[2], Doo Hyun Chung[1,2]

[1]Department of Pathology, Seoul National University College of Medicine, Seoul, Korea

[2]Laboratory of Immune Regulation in Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul, Korea

[3]Laboratory of Immunology and Vaccine Innovation, Department of Biotechnology, College of Life Sciences and Biotechnology, Korea University, Seoul, Korea

[4]Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam, Korea

[5]Institute of Allergy and Clinical Immunology, Seoul National University Medical Research Council, Seoul, Korea

Corresponding author:

Doo Hyun Chung

Department of Pathology, Seoul National University College of Medicine,

Laboratory of Immune Regulation, Department of Biosciences, Seoul National University College of Medicine

101 Daehak-ro, Jongno-gu,

Seoul 03080, Korea

Tel: 82-2-740-8915

Fax: 82-2-743-5530

E-mail: doohyun@snu.ac.kr

*J. Koh and Y. D. Woo contributed equally to this work

Abstract

Invariant natural-killer T (*i*NKT) cells play pathogenic roles in allergic asthma in murine models and possibly also humans. While many studies show that the development and functions of innate and adaptive immune cells depend on their metabolic state, the evidence for this in *i*NKT cells is very limited. It is also not clear whether such metabolic regulation of *i*NKT cells could participate in their pathogenic activities in asthma. Here, we showed that acetyl-coA-carboxylase 1 (ACC1)-mediated *de novo* fatty-acid synthesis is required for the survival of *i*NKT cells and their deleterious functions in allergic asthma. ACC1, which is a key fatty-acid synthesis enzyme, was highly expressed by lung *i*NKT cells from WT mice that were developing asthma. *Cd4*-Cre*Acc1*^{fl/fl} mice failed to develop OVA-induced and HDM-induced asthma. Moreover, *i*NKT cell-deficient mice that were reconstituted with ACC1-deficient *i*NKT cells failed to develop asthma, unlike when WT *i*NKT cells were transferred. ACC1 deficiency in *i*NKT cells associated with reduced expression of fatty acid-binding proteins (FABPs) and peroxisome proliferator-activated receptor (PPAR)γ, but increased glycolytic capacity that promoted *i*NKT-cell death. Furthermore, circulating *i*NKT cells from allergic-asthma patients expressed higher *ACC1* and *PPARG* levels than the corresponding cells from non-allergic-asthma patients and healthy individuals. Thus, *de novo* fatty-acid synthesis prevents *i*NKT-cell death *via* an ACC1-FABP-PPARγ axis, which contributes to their homeostasis and their pathogenic roles in allergic asthma.

Introduction

Invariant natural-killer T (iNKT) cells are a subset of innate T cells that express a restricted T-cell receptor (TCR) repertoire (Vα24-Jα18 in humans and Vα14-Jα18 in mice) and respond to glycolipids presented on the CD1d molecule. One such glycolipid is α-galactosylceramide (α-GalCer), which originates from a marine sponge and strongly activates iNKT cells *in vitro* and *in vivo* [1, 2]. Once activated, iNKT cells secrete a diverse array of cytokines, including interferon (IFN)γ, interleukin (IL)-4, IL-13, IL-17A, and IL-10 [3]. These iNKT cell-derived cytokines play important roles in the pathogenesis of various immunological diseases, including sepsis, arthritis, tumors, and asthma [4, 5, 6, 7]. With regard to the latter, iNKT cells are one of the key effectors that promote allergic asthma in murine models. Specifically, by secreting IL-4, iNKT cells stimulate the differentiation of T-helper (Th)2 cells and their responses. Moreover, by secreting IL-13, iNKT cells promote the contraction of airway smooth muscle, which is a cardinal feature of asthma [8] [9]. In addition, we recently reported that iNKT cells contribute to the development of asthma by secreting X-C motif chemokine ligand-1, which recruits conventional X-C motif chemokine receptor 1-expressing type-1 dendritic cells into the lungs: this then stimulates the Th2 responses that drive asthma [10]. Thus, iNKT cell-derived soluble factors play multiple critical roles in the development of allergic asthma in murine models. It should be noted that it is somewhat unclear what role iNKT cells play in human asthma. Several studies suggest that human asthmatic airways contain large numbers of iNKT cells, but this was not observed by other studies [11, 12, 13, 14, 15, 16]. However, a recent review describes additional lines of evidence that suggest that iNKT cells do contribute to the development and exacerbation of human asthma development, although the mechanisms may not necessarily involve increased iNKT cell frequencies. Thus, further research on the roles of iNKT cells in mouse models and humans with asthma is needed to improve our understanding of the pathogenesis of this disease.

Many studies show that the development and functions of innate and adaptive immune

cells are dependent on the metabolic states of these cells [17]. In particular, the metabolic

balance between glycolysis and oxidative phosphorylation (OXPHOS) shapes the

differentiation and functions of CD4[+] Th cells, Foxp3[+] regulatory T cells (Tregs), cytotoxic

CD8[+] T cells, and macrophages [18]. The development and functions of iNKT cells may also

depend on their metabolism: while the evidence to date is quite limited [19], the thymic

development of iNKT cells and their inflammatory cytokine production as mature cells

associate with changes in their glycolysis:OXPHOS balance [20]. Moreover, peripheral iNKT-

cell functions are regulated by their levels of reactive oxygen species, which are byproducts

of aerobic metabolism [21]. It is also possible that immune-cell metabolism may participate in

asthma since a recent study showed that fatty acid-oxidation inhibitors alleviate house-dust

mite (HDM)-induced airway inflammation [22]. Similarly, the airway Th2 cells that mediate

allergic responses against HDM depend on glycolysis and lipid metabolism [23].

Several studies show that lipid biogenesis, which is a major metabolic pathway, also

affects the differentiation and functions of conventional CD4[+] T cells [24, 25, 26]: for example, a

CD5-like molecule enforces low cholesterol biosynthesis in Th17 cells, which limits ligand

availability for Rorγt, the Th17 master transcription factor, thereby preventing Th17 cells

from becoming pathogenic and inducing autoimmunity [23]. Similarly, several studies show

that acetyl-CoA-carboxylase 1 (ACC1) affects multiple T-cell subsets. ACC1 is a key fatty-

acid synthesis enzyme in the cytosol that catabolizes the ATP-dependent carboxylation of

acetyl-CoA to malonyl-CoA [27], and its downregulation impairs the peripheral persistence and

homeostatic proliferation of CD8[+] T cells, augments Treg development from naïve CD4[+] T

cells and their suppressive functions, blocks the development of Th17 cells from naïve CD4[+]

T cells [24, 28], impairs the formation of IL-5-producing CD4[+] T cells [29], and promotes CD4[+] T

cell development [30]. The T-cell changes caused by inhibiting ACC1 reduce disease severity in

murine models of listeria infection, chronic graft-versus-host disease, multiple sclerosis[31],

and T cell-mediated asthma [29]; it also promotes resistance to parasites [30]. Thus, by shaping

fatty-acid synthesis, ACC1 can alter pathogenic and protective T-cell responses.

To our knowledge, the role(s) of ACC1-mediated fatty-acid synthesis in other immune

cells, including *i*NKT cells, and in asthma have not been assessed. In this study, we used

ACC1-deficient (*Cd4*-Cre*Acc1*^fl/fl^) mice and adoptive transfer experiments to determine

whether ACC1-mediated fatty-acid metabolism in *i*NKT cells participates in the development

of asthma. Indeed, we found that ACC1 promotes the survival of *i*NKT cells by regulating a

fatty acid-binding protein (FABP)-PPARγ axis that can downregulate glycolysis. The

improved ACC1-mediated *i*NKT-cell survival shapes the homeostasis and functions of these

cells, which in turn promote the development of ovalbumin (OVA)- or HDM-induced allergic

asthma in mice. We also observed that the ACC1-FABP-PPARγ axis may be active in

patients with allergic asthma but not controls.

Results

Activated _i_NKT cells express high levels of a fatty-acid synthesis-related gene whose deletion blocks allergen-induced asthma

To determine whether intracellular metabolic processes in _i_NKT cells play important roles in allergic asthma, _i_NKT cells sorted from the lungs of OVA-induced asthma model mice were assessed for the expression of key enzymes involved in glycolysis, the tricarboxylic acid (TCA) cycle, β-oxidation, and fatty-acid synthesis (Fig. 1). Indeed, OVA challenge significantly increased _i_NKT cell expression of β-oxidation and fatty-acid synthesis enzymes and tended to elevate TCA-cycle enzyme expression. However, glycolysis-enzyme expression remained low (Fig. 1A, B). The significantly upregulated fatty-acid synthesis enzymes were Acc1, Fasn, and Oxsm: Acc1 and Fasn play a particularly key role in _de novo_ fatty-acid synthesis. Notably, when purified murine _i_NKT cells and conventional CD4$^+$ and CD8$^+$ T cells were activated _in vitro_ with anti-CD3/CD28, the _i_NKT cells expressed much higher levels of _Acc1_ and _Fasn_ than the other cells. By contrast, the CD4$^+$ and CD8$^+$ T cells expressed higher levels of TCA cycle (_Sdh2b_ and _Idh1a_) and glycolysis (_Hk2_ and _Pkm2_) enzymes, respectively (Fig. 1C). This is consistent with the previous Gene Set Enrichment Analysis of Oh et al.: when they compared unstimulated murine _i_NKT and CD4$^+$ T cells to each other, they found that genes involved in the 'lipid, fatty acid and steroid metabolism' pathway were enriched in _i_NKT cells (Fig. S1A) [32]. Thus, _i_NKT cells express _Acc1_ and _Fasn_ when they are stimulated, including in asthma, whereas conventional CD4$^+$ and CD8$^+$ T cells do not.

We then focused on ACC1. To test whether ACC1-mediated _de novo_ fatty-acid synthesis in _i_NKT cells shapes AHR and allergic immune responses, we generated _Cd4_-Cre_Acc1_$^{fl/fl}$ mice: the CD4-positive cells (_i.e._ conventional CD4$^+$ T and _i_NKT cells) in these mice cannot express ACC1. Since conventional CD4$^+$ T cells did not upregulate ACC1 when stimulated _in vitro_ or _in vivo_ (Fig. 1B, C), the knockout is likely to be physiologically

relevant for iNKT cells only. When the *Cd4*-Cre*Acc1*$^{fl/fl}$ and *Acc1*$^{fl/fl}$ control mice were

induced to undergo OVA-mediated asthma, we indeed observed that the loss of ACC1

expression attenuated AHR (Fig. 1D) and the total immune-cell numbers in the lung and the

bronchoalveolar lavage fluid (BALF) (Fig. S1B). These changes associated with significantly

decreased total lymphocyte, neutrophil, eosinophil, and macrophage frequencies in the BALF

(Fig. S1C) and lower eosinophil frequencies in the lung (Fig. S1D). The lung CD8$^+$ T cell

and alveolar- and interstitial-macrophage frequencies did not change. Importantly, the total

lung CD4$^+$ T cell population (excluding iNKT cells) also did not change significantly (Fig.

S1D). By contrast, there was a sharp drop in iNKT cells in the lung; this was also associated

with a significant increase in PI staining (Fig. 1E). qRT-PCR analysis of the conventional

lung CD4$^+$ T cell population (excluding iNKT cells) showed that loss of ACC1 expression

associated with lower *Gata3*, *Il4*, *Il5*, and *Il13* expression but similar *Tbx21*, *Rorc*, *Foxp3*,

Ifng, *Il-17a*, and *Il-10* expression (Fig. S1E, F). Indeed, when the lung iNKT cells and

conventional CD4$^+$ T cells were separately isolated, both iNKT cells and the conventional

CD4$^+$ T cells demonstrated dramatically lower *Il4* and *Il13* expression but similar *Ifng* and

Il17a expression (Fig. 1F and Fig. S1F). Notably, all of these findings were recapitulated

when we conducted the same analyses in mice that underwent HDM-induced asthma (Fig.

1G–I and Fig. S1G–K). Thus, the inability of lung iNKT cells to express ACC1 induced

iNKT-cell death during allergen-induced asthmogenesis. This also reduced their expression

of *Gata3*, *Il4*, *Il5*, and *Il13* in lung conventional CD4$^+$ T cells: these effects markedly reduced

the eosinophilia and AHR. These effects were independent of the inability of conventional

lung CD4$^+$ T cells to express ACC1 in the mice.

ACC1-deficient iNKT cells cannot generate α-GalCer-mediated AHR and their adoptive

transfer into iNKT cell-deficient mice cannot restore allergen-induced asthma

The experiments above suggest that iNKT cells alone mediate ACC1-mediated

regulation of airway inflammation. To test this further, we challenged *Cd4*-Cre*Acc1*[fl/fl] and

Acc1[fl/fl] mice intranasally with α-GalCer, which has been shown previously to induce *i*NKT

cell activation-mediated AHR [33]. Consistent with the results from OVA- and HDM-induced

asthma models, the *Cd4*-Cre*Acc1*[fl/fl] mice showed the following: less AHR and lung and

BALF inflammation; significantly fewer lymphocytes, neutrophils, and macrophages in the

BALF; less eosinophilia in the BALF and lungs; no difference in lung CD4$^+$ and CD8$^+$ T cell

frequencies but significantly fewer lung *i*NKT cells and greater PI staining of these cells;

lower expression by the lung conventional CD4$^+$ T cell population of *Gata3, Il4, Il5,* and *Il13*

but not *Tbx21, Rorc, Foxp3, Ifng, IL17a, or Il10*; and similar *Ifng* and *Il17a* expression but

dramatically reduced *Il4* and *Il13* expression by the lung *i*NKT cells (Fig. 2A–H). Thus,

ACC1-deficient *i*NKT cells also failed to promote α-GalCer-induced AHR.

To further confirm that it is the ACC1 in *i*NKT cells that regulated AHR, we adoptively

transferred an equal number of WT *i*NKT cells (*i.e.* from *Acc1*[fl/fl] mice) or ACC1-deficient

*i*NKT cells into Jα18 KO mice, which lack NKT cells [34]. The mice were then subjected to

OVA-induced asthma. The WT *i*NKT cells, but not the ACC1-deficient *i*NKT cells, restored

the following features: AHR (Fig. 3A); the high total immune-cell numbers in the BALF and

lungs (Fig. 3B); the high lymphocyte, neutrophil, eosinophil, and macrophage frequencies in

the BALF and lungs (Fig. 3C, D); the high *Gata3*, *Il4*, *Il5*, and *Il13* expression by the lung

CD4$^+$ T cell population (Fig. 3E, F). As expected, neither adoptively transferred cell type

affected the *Tbx21*, *Rorc*, *Foxp3*, *Ifng*, *Il17a*, and *Il10* expression by the lung conventional

CD4$^+$ T cell population (Fig. 3E, F). Moreover, the transfer of WT *i*NKT cells restored the

*i*NKT cell-numbers in the lungs and their expression of *Il4* and *Il13* without changing their

Ifng and *Il17a* expression; by contrast, transfer of ACC1-deficient *i*NKT cells led to very low

*i*NKT-cell frequencies in the lung and low expression of all cytokines (Fig. 3G, H). In

addition, the adoptively transferred ACC1-deficient *i*NKT cells in the lung displayed more

apoptosis than the transferred WT *i*NKT cells (Fig. 3I). Lastly, transfer of the WT *i*NKT cells

associated with significantly higher histological lung scores than transfer of the ACC1-

deficient *i*NKT cells (Fig. 3J). Thus, ACC1-deficient *i*NKT cells cannot promote AHR in

allergic asthma models.

ACC1-deficient *i*NKT cells exhibit defects in thymic development, homeostasis, and

activation

To investigate the effects of ACC1-mediated metabolic programming on the

development and function of *i*NKT cells, we examined the thymic development and

peripheral homeostasis of *i*NKT cells in *Cd4*-Cre*Acc1*$^{fl/fl}$ mice. Mature *Cd4-Acc1*$^{fl/fl}$ mice had

fewer *i*NKT cells in the thymus, spleen, and liver than *Acc1*$^{fl/fl}$ mice (Fig. S2A). By contrast,

the conventional CD4$^+$ and CD8$^+$ T cell numbers in the thymus and spleen were not affected

by *Cd4-Acc1*$^{fl/fl}$ (Fig. S2B). However, the ACC1 deletion did not alter the frequencies of the

NKT1, NKT2, and NKT17 subsets in the *i*NKT cell population at any of the developmental

stages, namely, stage S1 (CD24$^+$c-Kit$^-$CD44$^-$), stage S2 (CD24$^+$c-Kit$^-$CD44$^+$), and stage S3

(CD24$^+$c-Kit$^+$CD44$^+$) (Fig. S2C). Thus, ACC1 deficiency appeared to reduce *i*NKT-cell

numbers.

This was tested further by generating mixed bone marrow (BM) chimera mice: thus, a

1:1 ratio of CD45.1$^+$ WT BM cells and CD45.2$^+$ *Cd4-Acc1*$^{fl/fl}$ BM cells were transferred into

lethally irradiated CD45.2$^+$ WT recipient mice (Fig. S2D). The ACC1-deficient BM cells

generated markedly fewer *i*NKT cells in the thymus, spleen, and liver than the WT BM cells

(Fig. S2E, F). This confirmed that ACC1 deficiency induced an intrinsic defect in *i*NKT-cell

thymic development and peripheral homeostasis. Moreover, when *i*NKT cells sorted from the

livers of *Cd4-Acc1*$^{fl/fl}$ and *Acc1*$^{fl/fl}$ mice underwent TCR ligation with anti-CD3/CD28 *in vitro*,

the ACC1-deficient cells produced less CD69 activation marker and cytokine (Fig. S2G, H).

Thus, ACC1 plays a cell-intrinsic role in the regulation of *i*NKT-cell thymic

development, peripheral homeostasis, and activation.

Enhanced glycolysis promotes the cell death of ACC1-deficient *i*NKT cells

To investigate the mechanism(s) by which ACC1-mediated metabolic programming of *i*NKT cells promotes allergic asthma, we subjected the sorted unstimulated *i*NKT cells from *Cd4*-Cre*Acc1*[fl/fl] and *Acc1*[fl/fl] mice to transcriptome and metabolome analyses (Fig. 4A–D). The ACC1-deficient *i*NKT cells displayed distinct metabolic reprogramming compared to WT *i*NKT cells. Specifically, they exhibited low fatty-acid synthesis but high glycolysis. Indeed, compared to WT *i*NKT cells, their extracellular acidification rate (ECAR), the level of 2-NBDG, which is an indicator of glucose uptake, and expression of Glut1 and other glycolysis-related genes were higher while their expression of gluconeogenesis genes were lower. However, their oxygen consumption rate (OCR) was similar (Fig. 4E–H). Furthermore, the transcriptome analysis revealed significantly altered levels of cell death-related genes in ACC1-deficient *i*NKT cells (Fig. 4C). These observations suggest that ACC1-deficient *i*NKT cells are prone to dying and that this may correlate positively with glycolysis. Indeed, the ECAR and cell death in ACC1-deficient *i*NKT cells correlated positively (Fig. 4I). This relationship was confirmed by treating the ACC1-deficient *i*NKT cells with 2-deoxy-D-glucose (2-DG), which inhibits glycolysis [35]: this decreased their cell death and expression of the pro-apoptotic genes while increasing their expression levels of anti-apoptotic genes. These effects were not observed in WT *i*NKT cells (Fig. 4J, K). Thus, the high levels of glycolysis in ACC1-deficient *i*NKT cells promote their cell death.

ACC1-mediated metabolic programming promotes PPARγ expression and the subsequent survival of *i*NKT cells

To identify the target genes that regulate the metabolic reprogramming and cell death in ACC1-deficient *i*NKT cells, we analyzed the transcription factors in the transcriptome data described above. This showed that sorted unstimulated ACC1-deficient *i*NKT cells expressed

Pparg at significantly lower levels than WT *i*NKT cells (Fig. 5A, B). This was also observed

when the cells were stimulated with anti-CD3/CD28: the treatment elevated *Pparg*

expression in both cell types but much less in the ACC1-deficient cells (Fig. 5B). Thus,

PPAR-γ may play a critical role in ACC1-mediated modification of *i*NKT-cell functions.

This is consistent with the fact that PPAR-γ is important for activating genes involved in lipid

biosynthesis and the maintenance of metabolic homeostasis[36, 37]. Indeed, when anti-

CD3/CD28-stimulated ACC1-deficient *i*NKT cells were treated with the PPARγ agonist

pioglitazone, their expression of *Pparg* was not only significantly restored, their expression

of the gluconeogenesis genes *Cebpa*, *Fbp1*, and *G6pase* rose while their 2-NBDG uptake,

expression of Glut1 and the glycolysis genes *Hk2* and *Pkm2*, and cell death dropped (Fig.

5B–D).

ACC1 catalyzes the carboxylation of acetyl-CoA to malonyl-CoA, which is then

converted to the long-chain fatty acid palmitate [38]. We found that treatment with palmitate

had similar effects on CD3/CD28-stimulated ACC1-deficient *i*NKT cells as pioglitazone,

namely, it increased *Pparg* and gluconeogenesis gene expression and downregulated

glycolysis (Fig. 5E). These effects were also observed when unstimulated ACC1-deficient

*i*NKT cells were treated with palmitate: *Pparg* expression rose (Fig. 5G) while glucose

uptake and cell death dropped (Fig. 5H). Thus, restoring the levels of palmitate, the

downstream product of ACC1 activity, restored PPARγ expression, which in turn

downregulated glycolysis and thereby protected the cells from cell death.

PPARγ activity can be regulated by a variety of ligands, including fatty acids, whose

transport to the nucleus is actively facilitated by fatty acid-binding proteins (FABPs) [39, 40, 41].

The unstimulated ACC1-deficient *i*NKT cells showed depressed expression of the genes

encoding FABP1, FABP3, and FABP5 (Fig. 5F). Thus, ACC1 deficiency downregulated the

expression of these genes. We then asked whether circumventing the ACC1 deficiency by

treating the unstimulated cells with palmitate could improve the expression of these FABPs,

as exogenous treatment of palmitate has been shown to induce FABPs in macrophages:

indeed, *Fabp1*, *Fabp3*, and especially *Fabp5* expression were upregulated in *i*NKT cells as

well (Fig. 5F).

We then determined whether the FABPs mediate the proglycolytic and pro-apoptotic

effects of ACC1 (and palmitate) deficiency in *i*NKT cells by culturing unstimulated

palmitate-treated ACC1-deficient *i*NKT cells with a FABP inhibitor, FABP-IN-1, which is

known to inhibit FABP 3, 5 and 7. Indeed, the inhibitor blocked the ability of palmitate to

restore PPARγ expression (Fig. 5G) and suppress glucose uptake and cell death (Fig. 5H).

These findings were also observed when CD3/CD28-stimulated ACC1-deficient *i*NKT cells

were treated with palmitate with and without FABP inhibitor: the FABP inhibitor reversed

the ability of palmitate to increase *Pparg* and gluconeogenesis gene expression and

downregulate glycolysis (Fig. 5E).

Since FABP-inhibitor treatment did not affect the PPARγ expression, glucose uptake,

and cell death in WT *i*NKT cells (Fig. 5E, G, H), it seems that the ACC1 deficiency in *i*NKT

cells mediated their metabolic reprogramming by abolishing palmitate production: this

downregulated their FABP expression, which in turn suppressed PPARγ expression, which

increased glycolysis and thereby promoted cell death (Fig. 5I).

On a side note, the WT and ACC1-deficient *i*NKT cells did not differ in terms of the

methylation of the promoter site of PPARγ (Fig. S3A), and treatment of the unstimulated

cells with histone deacetylase (HDAC) inhibitor did not alter their cell-death rates or

expression of *Pparg*, *Hk2*, *Bcl2*, or *Bak* (Fig. S3B, C). Thus, the regulation of PPARγ

expression in ACC1-deficient *i*NKT cells may not involve epigenetic changes such as

methylation and acetylation.

The ACC1-PPAR-γ axis in *i*NKT cells contributes to AHR and airway inflammation in allergic asthma

To confirm that ACC1-mediated metabolic programming in *i*NKT cells regulates the development of AHR, we pretreated WT or ACC1-deficient *i*NKT cells with palmitate or pioglitazone, adoptively transferred them into Jα18 KO mice, and then subjected the mice to OVA-induced asthma (Fig. 6) or HDM-induced asthma (Fig. S4). As shown above (Fig. 3), transfer of untreated WT *i*NKT cells was able restore AHR, airway inflammation scores, and high lymphocyte, neutrophil, eosinophil, and macrophage frequencies in the lungs and BALF. It also generated a CD4$^+$ T cell population in the lungs that expressed high levels of *Gata3*, *Il4*, *Il5*, and *Il13* and low levels of *Tbx21*, *Rorc*, *Foxp3*, *Ifng, Il17a,* and *Il10*. Moreover, the sorted *i*NKT cells cells in the lungs expressed high levels of *Il4* and *Il13* but low levels of *Ifng* and *Il17a.* By contrast, transfer of the untreated ACC1-deficient *i*NKT cells did not achieve any of these pathological changes or lung CD4$^+$ T cell or *i*NKT cell gene-expression profiles. However, when the ACC1-deficient *i*NKT cells were pretreated with palmitate or pioglitazone, all pathological and gene-expression changes were restored (Fig. 6A–G and Fig. S4).

As shown above (Fig. 3I), the transfer of untreated WT *i*NKT cells into mice that were then subjected to OVA-induced asthma led to a moderate *i*NKT cell population in the lungs that showed low apoptosis. By contrast, the transfer of untreated ACC1-deficient *i*NKT cells led to low lung *i*NKT cell numbers that frequently took up PI. However, pretreatment of the ACC1-deficient *i*NKT cells with palmitate or pioglitazone increased lung *i*NKT cell numbers and reduced their apoptosis (Fig. 6H, I). This is consistent with our *in vitro* experiments with the unstimulated and anti-CD3/CD28-stimulated *i*NKT cells above (Fig. 5). Notably, the sorted *i*NKT cell population in the lungs of Jα18 KO mice that received untreated ACC1-deficient *i*NKT cells also demonstrated less PPARγ expression and more glucose uptake and Glut1 expression compared to when untreated WT *i*NKT cells were transferred, and this was

reversed when the ACC1-deficient *i*NKT cells were pretreated with palmitate or pioglitazone

(Fig. 6J). Thus, treatment with palmitate or pioglitazone increased the survival of adoptively

transferred ACC1-deficient *i*NKT cells in Jα18 KO mice, which restored their ability to

induce AHR in allergic asthma. Meanwhile, chemokine receptor signaling is also implicated

in regulating homeostasis of *i*NKT cell in the periphery. In particular, Meyer et al. suggested

that *i*NKT cells require CCR4 to localize to the airways and to induce AHR[42]. Thus, we

examined the expression of several chemokine receptors, including CCR4. We found that WT

and ACC1-deficient *i*NKT cells did not differ in their chemokine receptor expressions,

suggesting that the chemokine signaling may not be critical for ACC1-mediated regulation in

AHR.

Altogether, these results show that ACC1-mediated metabolic programming in *i*NKT

cells maintains their PPARγ expression, which downregulates their glycolysis and thereby

increases their survival. This allows these cells to promote AHR and airway inflammation in

allergic asthma models.

The ACC1-PPARγ axis in *i*NKT cells may also contribute to allergic asthma in humans

We next asked whether patients with allergic asthma demonstrate altered expression of

fatty-acid synthesis-related metabolic enzymes in *i*NKT cells compared to healthy volunteers.

Thus, we sorted the *i*NKT, Treg, CD4$^+$ T, and CD8$^+$ T cells from the PBMCs of 10 healthy

volunteers, 10 patients with allergic asthma, and 10 patients with nonallergic asthma and

examined their mRNA expression of *ACC1* and *PPARG*. We also examined the expression of

FASN because it, like ACC1, is a key *de novo* fatty-acid synthesis enzyme and our

experiment with CD3/CD28-stimulated murine *i*NKT cells and conventional CD4$^+$ and CD8$^+$

T cells showed that the *i*NKT cells expressed much higher levels of *Fasn* as well as *Acc1*

than the other cells (Fig. 1C). Indeed, circulating *i*NKT cells from allergic asthma patients

expressed significantly higher levels of *ACC1, FASN*, and *PPARG* than *i*NKT cells from

healthy controls and nonallergic asthma patients (Fig. S5A). Similarly, the *i*NKT cells from

the allergic asthma patients expressed significantly higher *FBP1* levels than the control *i*NKT

cells, although this difference was not observed for *HK2* (Fig. S5B). Lastly, the expression

levels of *IL4* and *IL13* were significantly higher in *i*NKT cells from the allergic asthma

patients compared to those from healthy controls and nonallergic asthma patients (Fig. S5C).

These changes in *ACC1, FASN*, *PPARG*, and *FBP1* expression were much more pronounced

in the *i*NKT cells from allergic asthma patients than in the CD4$^+$ T cells, CD8$^+$ T cells, and

Tregs from the same patients (Fig. S5). Thus, *i*NKT cells from allergic asthma patients

express higher *ACC1, FASN* and *PPARG* levels and lower levels of a glycolysis which is

accompanied with higher levels of *IL4* and *IL13* than *i*NKT cells from healthy controls and

nonallergic asthma patients. These findings suggest that the ACC1-PPAR-γ axis in *i*NKT

cells may also be involved in human allergic asthma.

Discussion

The present study showed that *Cd4*-Cre*Acc1*^{fl/fl} mice exhibited attenuated AHR and

airway inflammation in OVA- and HDM-induced asthma models due to the enhanced

apoptosis and loss of *Gata3*, *Il4*, and *Il13* expression in lung *i*NKT cells. The mechanism

appeared to partly involve the low production in *i*NKT cells of the ACC1-downstream long-

chain fatty acid palmitate, which impaired FABP expression; this reduced PPARγ expression

and enhanced glycolysis in the *i*NKT cells, which promoted their apoptosis. Moreover, the

Cd4-Cre*Acc1*^{fl/fl} mice displayed significant defects in the thymic development and peripheral

homeostasis of *i*NKT cells in a cell-intrinsic manner; thus, homeostatic dysregulation of

*i*NKT cells may also contribute to the attenuation of AHR in *Cd4*-Cre*Acc1*^{fl/fl} mice.

The negative effect of ACC1 deficiency on the asthmogenic *i*NKT-cell functions is

consistent with several studies that suggest that ACC1-mediated *de novo* fatty-acid synthesis

can also shape the behavior of other T cells, including Th17 cell and Treg differentiation, IL-

5 production by Th2 cells in the lungs, the generation of memory CD4$^+$ T cells, and CD8$^+$ T

cell proliferation. Like us, these studies also show that blocking ACC1 function can improve

various diseases in mouse models [24, 29, 30, 43]. In this regard, Tregs may also play a major role

in asthma. However, the expression level of Foxp3 was comparable between WT and ACC1-

deficient Tregs. The level of Foxp3 to some extent, serves as a critical determinant of

suppressive function of Tregs[44, 45]. Thus, we speculate that they might not critically contribute

to the development of asthma, although we cannot completely rule out the contribution of

Tregs to our studies.

It should be noted that *Cd4*-Cre*Acc1*^{fl/fl} mice lack ACC1 expression in both conventional

CD4$^+$ T cells and *i*NKT cells. It should be noted that *Cd4*-Cre*Acc1*^{fl/fl} mice lack ACC1

expression in both conventional CD4$^+$ T cells and *i*NKT cells. While the use of *i*NKT cell-

specific Cre system would demonstrate critical role of ACC1 in *i*NKT cells regarding allergic

asthma, there is no *i*NKT cell-specific Cre system available yet. In addition, the study

conducted by Nakajima et al, which reported that the absence of ACC1 in CD4$^+$ T cells

resulted in reduced numbers and functional impairment of memory CD4$^+$ T cells, leading to

less airway inflammation further suggests possibility of involvement of conventional CD4$^+$ T

cells in regulation of allergic asthma.

However, based on our experimental results, we believe that *i*NKT cells more contribute

to the regulation of allergic asthma for the following reasons - (i) while the number of

*i*NKT cells were significantly reduced in *Cd4*-Cre*Acc1*$^{fl/fl}$ mice, the number of conventional

CD4$^+$ T cells were only slightly reduced, (ii) *Cd4*-Cre*Acc1*$^{fl/fl}$ mice were dramatically

decreased in their AHR in α-GalCer induced allergic asthma model, and (iii) Jα18 KO mice

that lack *i*NKT cells almost completely restore their AHR when adoptively transferred with

WT *i*NKT cells but not ACC1-deficient *i*NKT cells. These results indicate that ACC1-

mediated regulation of AHR is significantly dependent on *i*NKT cells, which might

contribute to AHR in the study conducted by Nakajima et al. as well. From these, we believe

that while ACC1 is a critical regulator of both conventional CD4$^+$ T cells and *i*NKT cells in

regulation of allergic asthma, *i*NKT cells may contribute more to regulation of allergic

asthma compared to CD4$^+$ T cells.

The possibility that ACC1 regulates survival of *i*NKT cells by programming their

metabolism was first revealed by our transcriptome analysis of treatment-naïve *i*NKT cells:

the ACC1-deficient *i*NKT cells exhibited downregulation of fatty-acid synthesis genes, as

could be expected, but also upregulation of glycolysis pathway-related genes. This impaired

the survival of the cells: compared to the WT *i*NKT cells, the ACC1-deficient *i*NKT cells

were prone to apoptosis. This was observed in ACC1-deficient *i*NKT cells regardless of

whether they were treatment-naïve, stimulated by TCR ligation *in vitro*, or activated during

OVA/HDM-mediated asthma. And the apoptotic tendency of these cells was completely

reversed by treatment with glycolysis inhibitor *in vitro*. These findings are consistent with

those of Kumar et al.[21], who reported that the production of lactate by *i*NKT cells when their

glycolysis is upregulated promotes their cell death.

Furthermore, the apoptotic tendency of the ACC1-deficient *i*NKT cells was

accompanied by their functional impairment. The ACC1-deficient *i*NKT cells exhibited

impaired viability and functionality. Treatment of glycolysis inhibitor in ACC1-deficient

*i*NKT cells not only restored cellular survival but also their functionalities. From these results,

we speculate that ACC1-mediated regulation of both cellular homeostasis and cytokine

production cooperatively contributed to the asthma phenotype.

Our transcriptome assays of naïve *i*NKT cells showed that of the multiple candidate

transcription factors that were examined, only *Pparg* was upregulated in WT *i*NKT cells but

not ACC1-deficient *i*NKT cells. This difference was augmented by TCR ligation. This

suggested that downregulation of PPARγ could mediate the enhanced glycolysis and

apoptosis in ACC1-deficient *i*NKT cells. This was supported by the fact that the PPARγ

agonist pioglitazone largely restored gluconeogenesis and downregulated glycolysis and

apoptosis in these cells. These findings are consistent with reports that PPARγ regulates the

glycolytic pathway, although heterogeneous effects have been noted [46, 47, 48, 49, 50, 51]. Our

study also supports Fu et al., who observed that lipid biosynthesis is increased in *i*NKT cells

after activation, and that this is mediated by PPARγ. Specifically, PPARγ acts synergistically

with the transcription factor PLZF (which drives the acquisition of T-helper effector

functions by innate and innate-like lymphocytes) to activate the *Srebf1* transcription factor [52],

which regulates lipid biosynthesis. This induces cholesterol synthesis, which is required for

the optimal production of IFNγ by the *i*NKT cells. This PPARγ/PLZ-*Srebf1*-cholesterol

pathway was found to contribute to anti-tumor immunity [53]. Previous studies also reported

that the PPARγ agonist ciglitazone promotes the TGFβ-dependent conversion of naïve

effector T cells into Tregs whereas PPARγ deficiency increases the infiltration of Th17 cells

into the central nervous system in experimental autoimmune encephalomyelitis. [54, 55]. Thus,

PPARγ can regulate the differentiation of naïve CD4$^+$ T cells into both Th17 or Treg cells,

which supports the role of PPARγ in *i*NKT-cell homeostasis and functions. However, further

research is needed to determine the mechanisms by which PPARγ regulates the metabolic

programming and biological changes in *i*NKT cells.

We also explored how PPARγ expression was downregulated in ACC1-deficient *i*NKT

cells. Several studies have reported that PPARγ expression can be regulated by endogenous

ligands such as FABPs, as well as by histone methylation and acetylation [56, 57], and that these

changes can be regulated by the intracellular lipid levels [58, 59]. Indeed, we observed that (i)

ACC1-deficient *i*NKT cells demonstrated low FABP expression, (ii) exogenous fatty acid not

only reversed the effects of ACC1 deficiency on glycolysis and apoptosis, it also restored the

expression of both FABPs and PPARγ, and (iii) treating ACC1-deficient *i*NKT cells with

exogenous palmitate or a PPARγ agonist and then transferring them into *i*NKT cell-deficient

mice restored AHR in OVA- or HDM-induced asthma models. Thus, the reduced *de novo*

fatty-acid synthesis in ACC1-deficient *i*NKT cells downregulated the expression of FABPs,

which decreased the expression and activity of PPARγ, which in turn promoted glycolysis

and apoptosis. This FABP-PPARγ axis was also observed in studies on hepatocytes and

adipocytes, which reported that FABPs interact with PPARγ, thereby inducing its

transactivation [39, 60]. However, we did not observe that PPARγ promoter methylation was

altered in the ACC1-deficient *i*NKT cells. Taken together, our findings indicate that the

ACC1-FABP-PPARγ axis drives the proglycolytic metabolic reprogramming in ACC1-

deficient *i*NKT cells, thereby downregulating *i*NKT-cell homeostasis, and functions,

including *i*NKT cell-mediated AHR in asthma models.

Finally, we showed that the ACC1-FABP-PPARγ axis in *i*NKT cells may also play

pathogenic roles in human allergic asthma: *i*NKT cells in the blood of patients with allergic

asthma expressed higher levels of *IL4*, *IL13*, *ACC1*, *FASN*, and *PPARG* and lower levels of a

glycolytic gene than the *i*NKT cells of nonallergic asthma patients and healthy controls.

Moreover, these genes were up/downregulated much more strongly in the *i*NKT cells from

allergic asthma patients than in the other T-cell subsets from the same patients. Notably, the

differences between allergic asthma patients and the control patients did not reflect lower

frequencies of *i*NKT cells in the PBMCs from the allergic asthma patients (data not shown).

This is consistent with speculation in the literature that *i*NKT cells participate in human

asthma by altering their phenotype rather than their frequencies: for example, although

asthma patients and controls do not differ in terms of *i*NKT-cell frequencies in the BALF, the

asthma *i*NKT cells express more IL-4 [11, 61]. Thus, the ACC1-FABP-PPARγ axis in *i*NKT

cells may be activated and thereby contribute to human allergic asthma pathology.

In conclusion, derangement of the ACC1-FABP-PPAR-γ axis induces glycolysis-related

impairment of cell viability in *i*NKT cells, which limits their ability to induce AHR and

airway inflammation in murine asthma models. Therefore, *de novo* lipid synthesis in *i*NKT

cells may play a critical regulatory role in the development of AHR.

Materials and methods

Mice

Female C57BL/6 (B6) mice were obtained from the Orient Company (Seoul, Korea). *Cd4-Cre* mice were purchased from the Jackson Laboratory (stock no. 017336; The Jackson Laboratory, Bar Harbor, ME, USA). *Acc1*$^{fl/fl}$ mice were originally generated and kindly provided by Dr. Salih J. Wakil of the Baylor College of Medicine (Houston, TX, USA). To generate *Cd4*-Cre*Acc1*$^{fl/fl}$ mice, the *Acc1*$^{fl/fl}$ mice were crossed with *Cd4*-Cre mice. The animals were bred and maintained under specific-pathogen-free conditions at the Biomedical Research Institute of Seoul National University Hospital (Seoul, Korea). All experiments were conducted on mice that were 8–10 weeks old. All animal experiments were approved by the Institutional Animal Care and Use Committee at Seoul National University Hospital (SNUH-IACUC). The animals were maintained in an AAALAC International (#001169)-accredited facility in accordance with the Guide for the Care and Use of Laboratory Animals 8th edition.

Generation of PBMCs from healthy subjects and asthmatic patients

In total, 30 human PBMC samples were obtained from healthy control subjects and patients with nonallergic and allergic asthma. The PBMCs were obtained from 30 mL peripheral blood by using Ficoll-Paque (GE Healthcare, Uppsala, Sweden). The *i*NKT, CD4$^+$ T, Foxp3$^+$ Tregs, and CD8$^+$ T cells in the PBMCs were sorted by FACSAria II (Becton Dickinson, San Jose, CA, USA; purity < 95%). All subjects provided informed written consent to participate in the study. The Institutional Review Board of Seoul National University Bundang Hospital approved the human studies (IRB #: B-1901/517–304).

Preparation and activation of murine T and *i*NKT cells

To determine the mRNA expression, transcriptome, metabolome, ECAR, and OCR of T and

*i*NKT cells, the cells were obtained from mice as follows. CD4$^+$ or CD8$^+$ T cells from the

thymi, spleens, lungs, and livers of *Acc1*$^{fl/fl}$ and *Cd4*-Cre*Accl1*$^{fl/fl}$ mice were enriched by

FACSAria II (Becton Dickinson). To obtain *i*NKT cells, the mononuclear cells of the livers

of *Acc1*$^{fl/fl}$ and *Cd4*-Cre*Accl1*$^{fl/fl}$ mice were isolated by Percoll gradient centrifugation.

Thereafter, the *i*NKT cells were isolated by using APC-conjugated α-GalCer/CD1d

tetramers-unloaded or loaded with PBS-57 from the tetramer facility of the National Institutes

of Health (Bethesda, MD, USA) followed by sorting using FACS (purity < 95%).

Alternatively, thymic and liver mononuclear cells were labeled with APC-conjugated ☐-

GalCer/CD1d tetramers, bound to anti-APC magnetic beads, and enriched on a MACS

separator (Miltenyi Biotec, Auburn, CA, USA; purity 89%). To analyze the development of

thymic *i*NKTs cells, we re-stained enriched cells with CD1d tetramer and gated out CD3 and

CD1d tetramer double positive cells via flow cytometry to identify thymic *i*NKT cells, which

were used for further analysis. For functional analysis, *i*NKT, CD4$^+$ T, and CD8$^+$ T cells were

stimulated with plate coated anti-CD3 and CD28 antibodies overnight.

Adoptive transfer of *i*NKT cells in allergic asthma models

*i*NKT cells were obtained from the lungs of at least 10 *Acc1*$^{fl/fl}$ or *Cd4*-Cre*Acc1*$^{fl/fl}$ mice.

Mouse lungs were finely chopped into small pieces using razor blades and were

enzymatically digested using type IV collagenase. *i*NKT cells from the lungs were sorted via

FACS using CD1d tetramers. Approximately, 6.0×10^5 of *i*NKT cells were obtained from at

least 10 mice and were adoptively transferred into individual recipient mouse via the

intratracheal route.

Antibodies, reagents, and preparation of fatty acids

The antibodies used in flow cytometry are summarized in Table S1. The PPAR□ agonist

pioglitazone (E6910), the histone deacetylase (HDAC) inhibitor MG-132 (474790), 2-deoxy-

D-glucose (2-DG, D6134), and palmitic acid (C16:0, P9767) were purchased from Sigma-

Aldrich (St. Louis, MO, USA). Briefly, palmitic acid was dissolved in sterile water using a

vortex and heated to 70°C for 10 min. Palmitic acid was conjugated to bovine serum albumin

(BSA, Biosesang, Seongnam, Korea) in serum-free RPMI containing 5% non-esterified fatty

acids (NEFA)-free BSA immediately after dissolving as described previously [62]. The

conjugated-palmitic acid was shaken at 140 rpm at 40°C for 1 h before being added to the

cells. Serum-free RPMI containing 5% NEFA-free BSA was used as the vehicle control.

Cells were treated with pioglitazone at 10 μM, 2-DG at 10 mM, MG-132 at 5 μM, or

palmitate at 100 μM.

Flow-cytometric analysis

Single-cell suspensions were preincubated with mouse anti-Fc receptor antibodies (BD

Biosciences, Franklin Lakes, NJ, USA), stained with antibodies for 30 min at 4□, washed

twice with $1 \times$ PBS, and analyzed using a BD LSRII flow cytometer (BD Biosciences). For

intracellular cytokine staining, cells were stained for surface markers, permeabilized, and

then stained for intracellular proteins. Cells were permeabilized and fixed with Intracellular

Fixation & Permeabilization Buffer (eBioscience, Waltham, MA, USA), according to the

manufacturer's protocols. Data were analyzed with FlowJo software (version 10; TreeStar,

Ashland, OR, USA).

Measurement of glucose uptake capacity using 2-NBDG assay

To assess the uptake of the fluorescent glucose analog 2-NBDG, cells were incubated for 30

min with 50 μM 2-NBDG (Cambridge Bioscience, Cambridge, UK)-supplemented glucose-

free RPMI 1640 or RPMI containing 10% (v/v) FBS supplemented with 12.5 nM

MitoTracker Deep Red (Thermo Fisher Scientific, Waltham, MA, USA) for 30 min at 37□,

5% (v/v) CO_2. The cells were washed with $1 \times$ PBS, fixed with 0.5% (w/w)

paraformaldehyde (PFA, Sigma), and then captured on a BD LSR Fortessa flow cytometer

(BD Biosciences). After treating 2-NBDG, the fluorescence intensity of cells were measured

using flow cytometry and represented the degree of glucose uptake in cells.

Generation of mixed BM chimeras

BM cells were prepared from the femurs and tibias of WT B6 (CD45.1[+] background) or *Cd4-*

Acc1[fl/fl] B6 (CD45.2[+] background) donor mice. Recipient mice (CD45.2[+] background) were

lethally irradiated (800 rads) and injected intravenously with a 1:1 mixture of BM cells from

WT and *Cd4*-Cre*Acc1*[fl/fl] mice (1×10^6 cells) in total. The chimeras were analyzed 8 weeks

after BM transplantation.

OVA- and HDM-induced asthma models

To induce the OVA asthma model, OVA antigen (Sigma) was dissolved in saline to a 1

mg/mL concentration. The mice were injected intraperitoneally with 4 mg aluminum

hydroxide (Thermo Fisher Scientific) mixed with 100 µL OVA in saline (1 mg/mL solution)

on days 0 and 7. On days 14, 15, 16, and 17, the isoflurane-anesthetized mice were

challenged with 50 µg OVA (1 mg/mL solution) intranasally. Twenty-four h after the last

OVA instillation, the mice were sacrificed to measure the AHR. To induce the HDM asthma

model, HDM (B82; Greer) ws dissolved in saline to a final concentration of 1 mg/mL.

Isoflurane-anesthetized mice were then treated intranasally on days 1–14 with 25 µg HDM

solution and, 96 h after the final intranasal instillation, the AHR and other inflammatory

variables were measured.

In vivo **intranasal inoculation of α-GalCer**

Mice were inoculated intranasally with 200 ng of α-GalCer dissolved in 50 µL PBS and, 24 h

after the last instillation, the mice were sacrificed to measure the AHR.

Measurement of AHR

AHR was measured by administering increasing doses of methacholine (0, 5, 10, 20, and 40

mg/mL) using the forced oscillation technique (FlexiVent System; SCIREQ, Montreal,

Quebec, Canada). A snapshot perturbation maneuver was imposed to measure the resistance

of the entire respiratory system.

Preparation of single-cell suspensions from lungs and BALF and differential counting

Mouse lungs were diced into small pieces using razor blades and incubated for 1 h in

digestion medium (RPMI-1640 with 1 mg/mL collagenase type 4, Sigma) at $37^{\circ}C$. The lung

cells were then filtered and treated with red blood cell lysis buffer. BALF cells were acquired

by administering 1 mL PBS supplemented with 2% FBS 3 times using a tracheal cannula.

Lavage fractions were pooled together for total and differential cell counts. Differential cell

counts were conducted with cytospin preparations after Giemsa staining and were validated

by flow cytometric analysis using a BD Fortessa (BD Biosciences).

Transcriptome analysis

*i*NKT cells from $Acc1^{\text{fl/fl}}$ and $Cd4$-Cre$Acc1^{\text{fl/fl}}$ mice were sorted and total RNA was extracted

using RNeasy with QIAshredders (Qiagen, Germantown, MD, USA). RNA quality and

quantity were determined using an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa

Clara, CA, USA) and Nanodrop (Nanodrop Technologies, Wilmington, DE, USA). RNA-seq

analyses were performed at the Theragen Bio Institute (Suwon, Korea). A Gene Set

Enrichment Assay (GSEA) was performed to identify significant expression differences

under the two biological conditions and to identify the gene classes that were overexpressed

or underexpressed in *i*NKT cells from *Cd4*-Cre*Acc1*[fl/fl] mice relative to those from WT mice.

The transcriptome data were from sorted *i*NKT cells from eight mice/group that were pooled

together. We also analyzed transcriptome data from our previous study to compare the *i*NKT

cells to conventional CD4[+]T cells (GSE103190) [32]. These analyses were performed using

GenePattern (https://genepattern.broadinstitute.org/).

Metabolome analysis

*i*NKT cells were sorted and pooled from eight WT mice, or eight *Cd4*-Cre*Accl1*[fl/fl] mice. The

metabolites were extracted from each cell group as previously described [63, 64], followed by

MeOX and MTBSTFA derivatization. Metabolic profiling was performed by GC-MS/MS

dynamic multiple reaction monitoring (dMRM) analyses using the Agilent 7890/7000 GC

triple quadrupole mass spectrometer system. Gas chromatography was conducted using an

Agilent J&W HP-5ms UI 15 m × 0.25 mm × 0.25 µm (P/N 19091S-431UI) capillary column.

Helium was used as the carrier gas with a constant column flow rate of 1.5 mL min^{-1}. The

initial oven temperature of 60°C was increased to 320°C at a rate of 10°C min^{-1}. The mass

spectrometer was operated in the electron ionization mode at 70 eV. The analyses were

performed in the dMRM mode. The area ratios of the analyte and the internal standard were

used for the evaluation. The split/splitless injector temperature was set to 280°C. The samples

were injected in the splitless mode with an autosampler Agilent 7683 A injector. Agilent

MassHunter Data Acquisition Software (ver. B.04.00) and MassHunter Workstation Software

for Quantitative Analysis (QQQ) were used for data acquisition and the quantitative analysis,

respectively. The results were normalized to the *i*NKT-cell numbers and/or total protein

concentrations.

Quantitative RT-PCR

To analyze gene-expression levels in the metabolic pathways, immune cells were sorted and

pooled from four-eight mice/group, and used in the analysis. Total RNA was isolated with

TRIzol reagent (Thermo Fisher Scientific) according to the manufacturer's protocol. RNA

was reverse-transcribed into cDNA using Maloney murine leukemia virus reverse

transcriptase Taq polymerase (Promega, Madison, WI, USA). For quantitative RT-PCR,

gene-specific PCR products were quantified by an Applied Biosystems 7500 Sequence

Detection System (Applied Biosystems, Foster City, CA, USA). The list of primers and their

sequences used are summarized in Table S2. Gene-expression levels were normalized to

those of *Actb* for mice and *ACTB* for humans.

Histology

Lung tissues were obtained from WT and J☐18 KO mice that did or did not undergo transfer

of *i*NKT cells from *Acc1*[fl/fl] and *Cd4*-Cre*Acc1*[fl/fl] mice during OVA- or HDM-induced asthma.

The tissues were fixed in 10% formalin, embedded in paraffin, and sectioned, followed by

staining with hematoxylin and eosin or periodic acid-Schiff (PAS). Lung inflammation and

goblet cell hyperplasia were graded using a semi-quantitative scoring system as previously

described [65, 66]. Briefly, inflammatory-cell numbers were graded as follows: 0: normal; 1: few

cells; 2: a ring of inflammatory cells that was 1 cell-layer deep; 3: a ring of inflammatory

cells that was 2–4 cell-layers deep; and 4: a ring of inflammatory cells that was >4 cell-layers

deep. The goblet-cell numbers in the airway were graded as follows: 0: < 0.5% PAS-positive

cells; 1: < 25%; 2: 25–50%; 3: 50–75%; and 4: > 75%. Two fields were counted for each

slide and mean score was calculated for each group. Two pulmonary pathologists reviewed

and analyzed representative images.

ECAR and OCR assays

*i*NKT cells were sorted from eight mice/group and pooled together. ECAR and OCR assays

were performed using a Seahorse Extracellular Flux Analyzer XF24e (Seahorse Bioscience

Inc., Santa Clara, CA, USA) according to the manufacturer's instructions. Briefly, the cells

were seeded in a Seahorse plate and cultured overnight to 70–80% confluence. The culture

medium was then replaced with cellular assay medium supplemented with 1 mM glutamine

for the ECAR assay and 1 mM pyruvate, 2 mM glutamine, and 10 mM glucose for the OCR

assay. The cells were incubated for 1 h in a CO_2-free incubator before measurement. The

assays were performed according to the Seahorse protocols with final concentrations of 10

mM glucose, 2 μM oligomycin, and 50 mM 2-DG for ECAR, and 2 μM oligomycin, 1 μM

carbonyl cyanide-4 (trifluoromethoxy) phenylhydrazone (FCCP), and 1 μM antimycin A for

OCR. These assays were performed in at least three independent experiments on separate

days.

Pyrosequencing assay

Genomic DNA was converted with bisulfite by using the EZ DNA Methylation Kit (Zymo

Research, Irvine, CA, USA) according to the manufacturer's protocol. PyroMarkTM software

was used to design the pyrosequencing primers. In brief, bisulfite-converted DNA was

subjected to PCR amplification in a total volume of 25 μL. DNA methylation was assessed

using a PyroMark Q48 Autoprep system (Qiagen) and PyroMark Q48 Advanced CpG

Reagents (Qiagen). The nucleotide dispensation order was generated by entering the

sequences into the PyroMark Q48 Autoprep software ver. 2.4.2 (Qiagen). The methylation at

each CpG site was determined using the PyroMark Q48 Autoprep software set in CpG mode.

The mean methylation of all CpG sites within the target region was determined by using the

methylation at the individual CpG sites.

Statistical analyses

GraphPad Prism 7 (GraphPad Software, La Jolla, CA, USA) was used for the statistical

analyses and graphical display of the data. Unpaired 2-tailed t-tests or Mann-Whitney U test

were performed to compare the groups. P values < 0.05 were considered statistically

significant.

References

1. Allende, M.L. *et al.* S1P1 receptor expression regulates emergence of NKT cells in peripheral tissues. *FASEB J* **22**, 307-315 (2008).

2. Brossay, L. *et al.* CD1d-mediated recognition of an alpha-galactosylceramide by natural killer T cells is highly conserved through mammalian evolution. *J Exp Med* **188**, 1521-1528 (1998).

3. Abdollahi-Roodsaz, S. *et al.* Stimulation of TLR2 and TLR4 differentially skews the balance of T cells in a mouse model of arthritis. *The Journal of clinical investigation* **118**, 205-216 (2008).

4. Osuchowski, M.F. *et al.* Minimum Quality Threshold in Pre-Clinical Sepsis Studies (MQTiPSS): An International Expert Consensus Initiative for Improvement of Animal Modeling in Sepsis. *Shock* **50**, 377-380 (2018).

5. Ma, C. *et al.* Gut microbiome-mediated bile acid metabolism regulates liver cancer via NKT cells. *Science* **360** (2018).

6. Terabe, M. & Berzofsky, J.A. Tissue-Specific Roles of NKT Cells in Tumor Immunity. *Front Immunol* **9**, 1838 (2018).

7. Kim, H.Y. *et al.* NKT cells promote antibody-induced joint inflammation by suppressing transforming growth factor beta1 production. *J Exp Med* **201**, 41-47 (2005).

8. Akbari, O. *et al.* Essential role of NKT cells producing IL-4 and IL-13 in the development of allergen-induced airway hyperreactivity. *Nat Med* **9**, 582-588 (2003).

9. Manson, M.L. *et al.* IL-13 and IL-4, but not IL-5 nor IL-17A, induce hyperresponsiveness in isolated human small airways. *Journal of Allergy and Clinical Immunology* **145**, 808-+ (2020).

10. Woo, Y.D., Koh, J., Kang, H.R., Kim, H.Y. & Chung, D.H. The invariant natural killer T cell-mediated chemokine X-C motif chemokine ligand 1-X-C motif chemokine receptor 1 axis promotes allergic airway hyperresponsiveness by recruiting CD103(+) dendritic cells. *J Allergy Clin Immunol* **142**, 1781-1792 e1712 (2018).

11. Akbari, O. *et al.* CD4+ invariant T-cell-receptor+ natural killer T cells in bronchial asthma. *N*

Engl J Med **354**, 1117-1129 (2006).

12.   Das, J. *et al.* Natural killer T cells and CD8+ T cells are dispensable for T cell-dependent
allergic airway inflammation. *Nat Med* **12**, 1345-1346; author reply 1347 (2006).

13.   Foell, D., Wittkowski, H., Vogl, T. & Roth, J. S100 proteins expressed in phagocytes: a novel
group of damage-associated molecular pattern molecules. *Journal of leukocyte biology* **81**,
28-37 (2007).

14.   Hamzaoui, A. *et al.* NKT cells in the induced sputum of severe asthmatics. *Mediators Inflamm*
2006, 71214 (2006).

15.   Pham-Thi, N. *et al.* Enhanced frequency of immunoregulatory invariant natural killer T cells in
the airways of children with asthma. *J Allergy Clin Immunol* **117**, 217-218 (2006).

16.   Thomas, S.Y., Lilly, C.M. & Luster, A.D. Invariant natural killer T cells in bronchial asthma. *N*
Engl J Med **354**, 2613-2616; author reply 2613-2616 (2006).

17.   Jung, J., Zeng, H. & Horng, T. Metabolism as a guiding force for immunity. *Nat Cell Biol* **21**,
85-93 (2019).

18.   Ganeshan, K. & Chawla, A. Metabolic Regulation of Immune Responses. *Annual Review of*
Immunology, Vol 32 **32**, 609-634 (2014).

19.   Yarosz, E.L., Chang, C.H. & Kumar, A. Metabolism in Invariant Natural Killer T Cells: An
Overview. *Immunometabolism* **3** (2021).

20.   Kim, Y.H., Kumar, A., Chang, C.H. & Pyaram, K. Reactive Oxygen Species Regulate the
Inflammatory Function of NKT Cells through Promyelocytic Leukemia Zinc Finger. *J Immunol*
199, 3478-3487 (2017).

21.   Kumar, A. *et al.* Enhanced oxidative phosphorylation in NKT cells is essential for their survival
and function. *P Natl Acad Sci USA* **116**, 7439-7448 (2019).

22.   Al-Khami, A.A. *et al.* Fuelling the mechanisms of asthma: Increased fatty acid oxidation in
inflammatory immune cells may represent a novel therapeutic target. *Clin Exp Allergy* **47**,

1170-1184 (2017).

23.   Tibbitt, C.A. *et al*. Single-Cell RNA Sequencing of the T Helper Cell Response to House Dust
830   Mites Defines a Distinct Gene Expression Signature in Airway Th2 Cells. *Immunity* **51**, 169-
831   184 (2019).

833   24.   Berod, L. *et al*. De novo fatty acid synthesis controls the fate between regulatory T and T
834   helper 17 cells. *Nat Med* **20**, 1327-1333 (2014).

836   25.   Soroosh, P. *et al*. Oxysterols are agonist ligands of RORgammat and drive Th17 cell
837   differentiation. *Proc Natl Acad Sci U S A* **111**, 12163-12168 (2014).

839   26.   Wang, C. *et al*. CD5L/AIM Regulates Lipid Biosynthesis and Restrains Th17 Cell
840   Pathogenicity. *Cell* **163**, 1413-1427 (2015).

842   27.   Wakil, S.J. & Abu-Elheiga, L.A. Fatty acid metabolism: target for metabolic syndrome. *Journal*
843   *of lipid research* **50 Suppl**, S138-143 (2009).

845   28.   Huang, S.C. *et al*. Cell-intrinsic lysosomal lipolysis is essential for alternative activation of
846   macrophages. *Nat Immunol* **15**, 846-855 (2014).

848   29.   Nakajima, T. *et al*. ACC1-expressing pathogenic T helper 2 cell populations facilitate lung and
849   skin inflammation in mice. *J Exp Med* **218** (2021).

851   30.   Endo, Y. *et al*. ACC1 determines memory potential of individual CD4(+) T cells by regulating
852   de novo fatty acid biosynthesis. *Nat Metab* **1**, 261-275 (2019).

854   31.   Lee, J. *et al*. Regulator of fatty acid metabolism, acetyl coenzyme a carboxylase 1, controls T
855   cell immunity. *J Immunol* **192**, 3190-3199 (2014).

857   32.   Oh, S.J., Ahn, J.Y. & Chung, D.H. Comparison of Invariant NKT Cells with Conventional T
858   Cells by Using Gene Set Enrichment Analysis (GSEA). *Immune network* **11**, 406-411 (2011).

860   33.   Meyer, E.H. *et al*. Glycolipid activation of invariant T cell receptor+ NK T cells is sufficient to
861   induce airway hyperreactivity independent of conventional CD4+ T cells. *Proc Natl Acad Sci*
862   *U S A* **103**, 2782-2787 (2006).

34.   Chandra, S. *et al.* A new mouse strain for the analysis of invariant NKT cell function. *Nat*
Immunol **16**, 799-800 (2015).

35.   Pajak, B. *et al.* 2-Deoxy-d-Glucose and Its Analogs: From Diagnostic to Therapeutic Agents.
Int J Mol Sci **21** (2019).

36.   Ahmadian, M. *et al.* PPARgamma signaling and metabolism: the good, the bad and the future.
Nat Med **19**, 557-566 (2013).

37.   Grygiel-Gorniak, B. Peroxisome proliferator-activated receptors and their ligands: nutritional
and clinical implications--a review. *Nutr J* **13**, 17 (2014).

38.   Moon, Y.A., Shah, N.A., Mohapatra, S., Warrington, J.A. & Horton, J.D. Identification of a
mammalian long chain fatty acyl elongase regulated by sterol regulatory element-binding
proteins. *J Biol Chem* **276**, 45358-45366 (2001).

39.   Wolfrum, C., Borrmann, C.M., Borchers, T. & Spener, F. Fatty acids and hypolipidemic drugs
regulate peroxisome proliferator-activated receptors alpha - and gamma-mediated gene
expression via liver fatty acid binding protein: a signaling path to the nucleus. *Proc Natl Acad*
Sci U S A **98**, 2323-2328 (2001).

40.   Patil, R. *et al.* A ligand-induced structural change in fatty acid-binding protein 1 is associated
with potentiation of peroxisome proliferator-activated receptor alpha agonists. *J Biol Chem*
294, 3720-3734 (2019).

41.   Hou, Y. *et al.* FABP5 controls macrophage alternative activation and allergic asthma by
selectively programming long-chain unsaturated fatty acid metabolism. *Cell Rep* **41**, 111668
(2022).

42.   Meyer, E.H. *et al.* iNKT cells require CCR4 to localize to the airways and to induce airway
hyperreactivity. *J Immunol* **179**, 4661-4671 (2007).

43.   Goto, Y. *et al.* Segmented filamentous bacteria antigens presented by intestinal dendritic cells
drive mucosal Th17 cell differentiation. *Immunity* **40**, 594-607 (2014).

44.   Fontenot, J.D., Gavin, M.A. & Rudensky, A.Y. Foxp3 programs the development and function

900 of CD4+CD25+ regulatory T cells. *Nat Immunol* **4**, 330-336 (2003).

902 45. Wing, J.B., Tanaka, A. & Sakaguchi, S. Human FOXP3(+) Regulatory T Cell Heterogeneity

903 and Function in Autoimmunity and Cancer. *Immunity* **50**, 302-316 (2019).

905 46. Lehmann, J.M. *et al.* An Antidiabetic Thiazolidinedione Is a High-Affinity Ligand for

906 Peroxisome Proliferator-Activated Receptor Gamma(Ppar-Gamma). *Journal of Biological*

907 *Chemistry* **270**, 12953-12956 (1995).

909 47. Masters, C.J., Reid, S. & Don, M. Glycolysis - New Concepts in an Old Pathway. *Molecular*

910 *and Cellular Biochemistry* **76**, 3-14 (1987).

912 48. Roberts, L.D. *et al.* The contrasting roles of PPARdelta and PPARgamma in regulating the

913 metabolic switch between oxidation and storage of fats in white adipose tissue. *Genome Biol*

914 **12**, R75 (2011).

916 49. Panasyuk, G. *et al.* PPARgamma contributes to PKM2 and HK2 expression in fatty liver. *Nat*

917 *Commun* **3**, 672 (2012).

919 50. Coman, C. *et al.* Simultaneous Metabolite, Protein, Lipid Extraction (SIMPLEX): A

920 Combinatorial Multimolecular Omics Approach for Systems Biology. *Mol Cell Proteomics* **15**,

921 1453-1466 (2016).

923 51. Calvier, L. *et al.* PPARgamma Links BMP2 and TGFbeta1 Pathways in Vascular Smooth

924 Muscle Cells, Regulating Cell Proliferation and Glucose Metabolism. *Cell Metab* **25**, 1118-

925 1134 e1117 (2017).

927 52. Mao, A.P. *et al.* Multiple layers of transcriptional regulation by PLZF in NKT-cell development.

928 *Proc Natl Acad Sci U S A* **113**, 7602-7607 (2016).

930 53. Fu, S. *et al.* Impaired lipid biosynthesis hinders anti-tumor efficacy of intratumoral iNKT cells.

931 *Nat Commun* **11**, 438 (2020).

933 54. Wohlfert, E.A., Nichols, F.C., Nevins, E. & Clark, R.B. Peroxisome proliferator-activated

934 receptor gamma (PPAR gamma) and immunoregulation: Enhancement of regulatory T cells

935 through PPAR gamma-dependent and -independent mechanisms. *Journal of Immunology*

936 **178**, 4129-4135 (2007).

55.   Klotz, L. *et al.* The nuclear receptor PPAR gamma selectively inhibits Th17 differentiation in a
T cell-intrinsic fashion and suppresses CNS autoimmunity. *Journal of Experimental Medicine*
206, 2079-2089 (2009).

56.   Huang, Q. *et al.* Mechanistic Insights Into the Interaction Between Transcription Factors and
Epigenetic Modifications and the Contribution to the Development of Obesity. *Front*
Endocrinol (Lausanne) **9**, 370 (2018).

57.   Furuhashi, M. & Hotamisligil, G.S. Fatty acid-binding proteins: role in metabolic diseases and
potential as drug targets. *Nat Rev Drug Discov* **7**, 489-503 (2008).

58.   Veerkamp, J.H. & van Moerkerk, H.T. Fatty acid-binding protein and its relation to fatty acid
oxidation. *Mol Cell Biochem* **123**, 101-106 (1993).

59.   Barak, Y. & Lee, C.H. The Molecular Basis of PPAR Function. *PPAR Res* **2010**, 510530
(2010).

60.   Tan, N.S. *et al.* Selective cooperation between fatty acid binding proteins and peroxisome
proliferator-activated receptors in regulating transcription. *Mol Cell Biol* **22**, 5114-5127 (2002).

61.   Shim, J.U. & Koh, Y.I. Increased Th2-like Invariant Natural Killer T cells in Peripheral Blood
From Patients With Asthma. *Allergy Asthma & Immunology Research* **6**, 444-448 (2014).

62.   Mayer, C.M. & Belsham, D.D. Palmitate attenuates insulin signaling and induces endoplasmic
reticulum stress and apoptosis in hypothalamic neurons: rescue of resistance and apoptosis
through adenosine 5' monophosphate-activated protein kinase activation. *Endocrinology* **151**,
576-585 (2010).

63.   Faubert, B. *et al.* AMPK Is a Negative Regulator of the Warburg Effect and Suppresses
Tumor Growth In Vivo. *Cell Metabolism* **17**, 113-124 (2013).

64.   Faubert, B. *et al.* Loss of the tumor suppressor LKB1 promotes metabolic reprogramming of
cancer cells via HIF-1 alpha. *Proceedings of the National Academy of Sciences of the United*
States of America **111**, 2554-2559 (2014).

65.   Kujur, W., Gurram, R.K., Haleem, N., Maurya, S.K. & Agrewala, J.N. Caerulomycin A inhibits

Th2 cell activity: a possible role in the management of asthma. *Sci Rep* **5**, 15396 (2015).

66.   Myou, S. *et al.* Blockade of inflammation and airway hyperresponsiveness in immune-
sensitized mice by dominant-negative phosphoinositide 3-kinase-TAT. *J Exp Med* **198**, 1573-
1582 (2003).

Acknowledgments

We would like to appreciate the NIH Tetramer Core Facility at the National Institute of
Health, USA for providing PBS-57 loaded CD1d tetramers.

Declaration of interests

The authors declare no competing interests.

Author contributions

JK, YDW, and YHJ conducted the experiments; KCJ provided mice; CJP, KSH and CYS
provided human samples and clinical information; KJH, KCJ, YKJ and HYK contributed to
interpretation of results; DHC, JK, and YDW designed the experiments, interpreted data, and
wrote the paper.

Funding Statement

This work was supported by the Basic Research Program through the National Research
Foundation of Korea (NRF) funded by the MSIT (NRF-2020R1A4A1017515) and the
National Research Foundation of Korea (NRF) grant funded by the Korea government
(MSIT) (NRF-2020R1A2C2008312).

Figure legends

Figure 1. The fatty-acid synthesis pathway is upregulated in *i*NKT cells during murine

allergic asthma.

(A–B) WT mice were subjected to OVA-induced asthma. (A) Schematic depiction of the

OVA-induced allergic asthma model (left). The airway resistance of naïve, OVA-sensitized,

and OVA-sensitized and OVA-challenged WT mice (right). (B) Gene expression of enzymes

related to glycolysis, the TCA cycle, β-oxidation, and fatty-acid synthesis in lung *i*NKT cells

that were sorted on d0, d7, and d17. The data shown in (A) and (B) are pooled data from four

independent experiments (n = 8/group). (C) *In vitro* gene expression of enzymes related to

glycolysis (*Hk2* and *Pkm2*), the TCA cycle (*Sdh2b* and *Idh1a*), and fatty-acid synthesis (*Acc1*

and *Fasn*) in CD8$^+$ T, CD4$^+$ T, and *i*NKT cells before and after CD3/CD28 stimulation.

Representative results from two independent experiments (n = 4/group) are shown. (D–I)

Cd4-Cre*Acc1*$^{fl/fl}$ mice and the *Acc1*$^{fl/fl}$ control mice were subjected to OVA-induced asthma

(D–F) or HDM-induced asthma (G–I). (D and G) Airway resistance. (E and H) Numbers and

survival status of *i*NKT cells that were sorted from the lung by flow cytometry. (F and I)

Cytokine expression in *i*NKT cells sorted from the lung. The data shown in (D–I) are pooled

data from four independent experiments (n = 8/group). All data are presented as mean ± SEM.

n.s, not significant. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, as determined by Unpaired 2-tailed

t-tests.

1020

Figure 2. ACC1 deficiency in murine *i*NKT cells associates with failure to develop α-

GalCer-induced AHR.

Cd4-Cre*Acc1*$^{fl/fl}$ and *Acc1*$^{fl/fl}$ mice were administered α-GalCer intranasally. (A) Airway

resistance. (B) Total immune cell numbers in the lungs and BALF. (C–D) Numbers of

immune-cell subsets in the BALF (C) and lungs (D). (E–F) Expression levels of transcription

factors (E) and cytokines (F) in the conventional CD4$^+$ T cell population that was sorted from

the lungs. (G) Numbers and survival status of *i*NKT cells sorted from the lungs. (H) Cytokine

expression in *i*NKT cells sorted from the lungs. All data are pooled from five independent

experiments (n = 10/group) and presented as mean ± SEM. n.s, not significant. *$p < 0.05$, **p

< 0.01, ***$p < 0.001$, as determined by Unpaired 2-tailed t-tests.

1031

Figure 3. Adoptive transfer of WT, but not ACC1-deficient, *i*NKT cells restores the

ability of NKT-knockout mice to develop OVA-induced asthma.

*i*NKT cells sorted from naïve *Cd4*-Cre*Acc1*$^{fl/fl}$ and *Acc1*$^{fl/fl}$ mice were adoptively transferred

into naïve WT and Jα18 KO mice, after which OVA-induced asthma was initiated. (A)

Airway resistance. (B) Total inflammatory cell numbers in the lungs and BALF. (C–D)

Numbers of immune-cell subsets in BALF (C) and lungs (D). (E–F) Expression levels of

transcription factors (E) and cytokines (F) in CD4^{+} T cells sorted from the lungs. (G)

Cytokine expression in *i*NKT cells sorted from the lungs. (H–I) Numbers (H) and survival

status (I) of *i*NKT cells from the lungs. (J) Representative histological lung section images

and histological scores. The data in (A–J) are pooled data from four independent experiments

(n = 8/group). All data are presented as mean ± SEM. n.s, not significant. *$p < 0.05$, **$p <$

0.01, ***$p < 0.001$, as determined by Unpaired 2-tailed t-tests.

1044

Figure 4. The high glycolytic capacity of ACC1-deficient *i*NKT cells associates with their

enhanced cell death.

*i*NKT cells were sorted from the liver of naïve *Cd4*-Cre*Acc1*$^{fl/fl}$ and *Acc1*$^{fl/fl}$ mice and used

directly in the experiments. The two *i*NKT cell preparations were compared. (A) Volcano

plot of the gene expression. (B) Signature gene expression, as determined by Gene Set

Enrichment Analysis (GSEA). (C) Heat map of the fatty-acid synthesis, glycolysis, cell death,

and cytokine genes. (D) Metabolome data map of the glycolysis, TCA cycle, and fatty-acid

synthesis metabolites. (E) Seahorse extracellular flux analysis of ECAR and OCR. (F)

Expression of glycolysis-related (*Hk2*, *Pkm2*) and gluconeogenesis-related (*Fbp1*, *G6pase*) genes. (G) 2-NBDG uptake. (H) GLUT1 expression. (I) Correlation between *i*NKT-cell death and maximal glycolytic activity. (J–K) The *i*NKT cells were treated with 2-DG to inhibit glycolysis. The effect of 2-DG on cell death (J) and intrinsic cell death-related gene expression (K) was then determined. In (A–D), the *i*NKT cells were from 8 mice/group. The data in (D) are presented as mean ± SEM. The data in (E–K) are pooled data from five independent experiments (n = 5/group) and are presented as mean ± SEM. n.s, not significant. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, as determined by Unpaired 2-tailed t-tests.

Figure 5. Low PPARγ expression in ACC1-deficient *i*NKT cells contributes to their enhanced cell death

*i*NKT cells sorted from the liver of naïve *Cd4*-Cre*Acc1*^{fl/fl} and *Acc1*^{fl/fl} mice were either used directly in the experiment (A) or cultured with and/or without anti-CD3/CD28 with and without the indicated additives (C–H). (A) Heatmap of the transcription factor genes and RT-PCR confirmation of *Pparg* expression in the uncultured *i*NKT-cell preparations. (B) CD3/CD28-stimulated and unstimulated *i*NKT cells were cultured with and without pioglitazone and *Pparg*, *Cepba*, *Fbp1*, *G6pase*, *Hk2*, and *Pkm2* expression was examined. (C–D) Unstimulated *i*NKT cells were cultured with and without pioglitazone and 2-NBDG uptake, GLUT1 expression (C), and cell death (D) were examined. (E) CD3/CD28-stimulated and unstimulated *i*NKT cells were cultured with and without palmitate and/or FABP inhibitor and *Pparg*, *Cepba*, *Fbp1*, *G6pase*, *Hk2*, and *Pkm2* expression was examined. (F) Unstimulated *i*NKT cells were cultured with and without palmitate and *Fabp1*, *Fabp3*, and *Fabp5* expression was examined. (G–H) Unstimulated *i*NKT cells were cultured with and without palmitate and/or FABP inhibitor and PPARγ expression (G), 2-NBDG uptake, and cell death (H) were examined. (I) Graphical abstract of ACC1-FABP-PPARγ in *i*NKT cells are shown. The data in (B–H) are pooled data from four independent experiments (n =

1079   4/group) and are presented as mean ± SEM. n.s, not significant. *$p < 0.05$, **$p < 0.01$, ***p

1080 < 0.001, as determined by Unpaired 2-tailed t-tests.

1081

Figure 6. Pretreatment of ACC1-deficient *i*NKT cells with PPARγ agonist or exogenous

fatty acid restores OVA-induced asthma.

*i*NKT cells sorted from naïve *Cd4*-Cre*Acc1*[fl/fl] and *Acc1*[fl/fl] mice were pretreated with and

without pioglitazone or palmitate and then adoptively transferred into naïve WT and Jα18

KO mice. OVA-induced asthma was then initiated. (A) Airway resistance. (B) Representative

histological images and histological scores of lung sections. (C) Total immune cell numbers

in the lungs and BALFs. (D–E) Immune-cell subsets in the BALFs (D) and lungs (E). (F–G)

Transcription factor (F) and cytokine (G) expression in the total CD4$^+$ T cell population from

the lungs. (H) Cell numbers and survival status of the lung *i*NKT cells. (I) Cytokine

expression in the lung *i*NKT cells. (J) PPARγ, 2-NBDG, and GLUT1 expression in the lung

*i*NKT cells. The data are pooled from four independent experiments (n = 8/group) and are

presented as mean ± SEM. n.s, not significant. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, as

determined by Unpaired 2-tailed t-tests.

1095

Legends for supplementary figures

Figure S1. *Cd4*-Cre*Acc1*[fl/fl−] mice fail to develop airway resistance in the HDM-induced

asthma model

(A) Gene set enrichment analysis (GSEA) in *i*NKT versus CD4$^+$ T cells. *Acc1*[fl/fl] and *Cd4*-

Cre*Acc1*[fl/fl] mice were intranasally challenged with OVA (B-F) or HDM (G-K), and various

parameters were measured. (B and G). Total cell counts from the lungs and BALs (C, D, H, I).

Estimation of immune cell subsets in the BALs (C and H) and lungs (D and I) via flow

cytometry (E, F, J, K). Expression levels of transcription factors (E and J) and cytokines (F

and K) in CD4$^+$ T (excluding *i*NKTs) cells sorted from lungs. Data were pooled from four

independent experiments with n = 8 per group and presented as means ± SEMs. n.s, not

significant. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ as determined by Unpaired 2-tailed t-tests

1107

Figure S2. Deficiency of ACC1 in *i*NKT cells perturbed their thymic development and

peripheral homeostasis in cell intrinsic manner

(A) The percentages of *i*NKT cells in the thymi, spleens, and livers from *Cd4*-Cre*Acc1*^{fl/fl} and

Acc1^{fl/fl} mice were measured via flow cytometry. (B) Analysis of CD4$^+$ SP thymocytes, CD8$^+$

SP thymocytes, CD4$^+$CD8$^+$ DP thymocytes, CD4$^-$CD8$^-$ DN thymocytes, and CD8$^+$T, CD4$^+$T

cells from the spleens of *Cd4*-Cre*Acc1*^{fl/fl} and *Acc1*^{fl/fl} mice. (C) Thymic *i*NKT cells from

thymus of *Cd4*-Cre*Acc1*^{fl/fl} and *Acc1*^{fl/fl} mice were enriched via magnetic-associated cell

sorting (MACS). Percentages of NKT1s, NKT2s, and NKT17s and developmental stages of

MACS purified CD3$^+$CD1d tetramer$^+$ *i*NKT cells were measured via flow cytometry. (A-C)

Data were pooled from four independent experiments with n = 4 per group. (D) Mixed bone

marrow chimeras were generated by adoptive transfer of a mixture of WT CD45.1 bone

marrow and CD45.2 *Cd4*-Cre*Acc1*^{fl/fl} bone marrow into irradiated WT mice. To investigate

the reconstitution, CD1d tetramer$^+$ *i*NKT cells were stained with CD45.1 and CD45.2

antibodies via flow cytometry. (E-F) The percentages of *i*NKT cells in the thymi, spleens,

and livers (E) and of CD4$^+$SP, CD8$^+$SP, CD4$^+$CD8$^+$DP, and CD4$^-$CD8$^-$DN thymocytes and

splenic CD4$^+$T, CD8$^+$T cells, CD3e$^+$T cells, B cells (F) were measured via flow cytometry.

(G-H) *i*NKT cells sorted from the livers of *Cd4*-Cre*Acc1*^{fl/fl} and *Acc1*^{fl/fl} mice were stimulated

with CD3/CD28 antibodies. (G) The levels of IL-4 and IFN-γ in the culture supernatants

were assessed by ELISA. (H) Expression of CD69 was estimated via flow cytometry. (D-H)

Data were pooled from three independent experiment with n = 30 per group and presented as

means ± SEMs. n.s, not significant. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ as determined by

Unpaired 2-tailed t-tests

1130

Figure S3. ACC1 deficiency does not affect epigenetic modification of *Pparg* in *i*NKT cells

(A) Schematic structure of the *Pparg* promoter and identification of the CpG-enriched regions. *Pparg* methylation was measured via pyrosequencing. (B and C) Expression levels of *Pparg*, *Hk2*, *Bcl2*, *Bak* (B), and cell death (C) of *i*NKT cells from *Cd4*-Cre*Acc1*$^{fl/fl}$ or *Acc1*$^{fl/fl}$ mice in the presence or absence of HDAC inhibitor treatment. Data were pooled from four independent experiments with n = 4 per group and presented as means ± SEMs. n.s, not significant. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ as determined by Unpaired 2-tailed t-tests

Figure S4. Adoptive transfer of Acc1-deficient *i*NKT cells pretreated with exogenous fatty acid or PPAR-γ agonist into *i*NKT cell-deficient mice (Jα18 KO) restores AHR in the HDM-induced asthma model

(A) Airway resistance was measured in WT, Jα18KO, and Jα18KO mice adoptively transferred with *i*NKT cells from *Acc1*$^{fl/fl}$ or *Cd4*-Cre*Acc1*$^{fl/fl}$ mice in the presence or absence of pioglitazone or palmitate treatment upon HDM challenge. (B) Representative histological images of lung sections from the indicated mouse groups challenged with HDM. (C-E) Total cell numbers in the lungs and BALs were counted (C), and subsets of immune cells in the BALs (D) and lungs (E) were estimated via flow cytometry. (F-G) Expression levels of transcription factors (F) and cytokines (G) were measured in CD4$^+$ T cells sorted from the lungs of mice with OVA-induced asthma. (H) Expression levels of cytokines were measured in *i*NKT cells sorted from the lungs of each mouse group. (I) Heat map of *Ccr4*, *Ccr6*, *Cxcr3*, and *Cxcr4* expression in WT and ACC1-deficient *i*NKT cells from RNA-Seq data and RT-PCR validation of the chemokine receptor expression levels in WT and ACC1-deficient *i*NKT cells. Data were pooled from four independent experiments with n = 8 per group and presented as means ± SEMs. n.s, not significant. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ as determined by Unpaired 2-tailed t-tests

1157

Figure S5. Genes for fatty acid metabolism are highly expressed in *i*NKT cells from

PBMCs of patients with allergic asthma

(A and B) The expression levels of *ACC1*, *FASN*, *PPARG* (A), *HK2*, and *FBP1* (B) in CD8$^+$

T, CD4$^+$ T, regulatory T cells, and *i*NKT cells from peripheral blood samples of healthy

donors and patients with non-allergic asthma or allergic asthma (n = 10 per group). (C) The

expression levels of *IL4*, *IL13, IFNG, IL10* were measured in *i*NKT cells from peripheral

bloods of healthy donors and patients with non-allergic asthma or allergic asthma. Data are

presented as means ± SEMs. n.s, not significant. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ as

determined by Mann-Whitney U test

1167

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2

Figure S3

A

PPAR gamma promoter site

19% 38% 22%

Acc1^{fl/fl}

CGA**TC**GA**TC**TACAGA**TC**AGA

16% 29% 22%

CD4-Cre*Acc1*^{fl/fl}

CGA**TC**GA**TC**TACAGA**TC**AGA

% DNA methylation

n.s

Acc1^{fl/fl}   *Cd4*-**Cre***Acc1*^{fl/fl}

B

● *Acc1*^{fl/fl}
● *Cd4*-Cre*Acc1*^{fl/fl}

Relative expression

n.s   n.s

inhibitor   inhibitor

Pparg

Relative expression

n.s   n.s

inhibitor   inhibitor

Hk2

Relative expression

n.s   n.s

inhibitor   inhibitor

Bcl2

Relative expression

n.s   n.s

inhibitor   inhibitor

Bak

C

DMSO   HDAC inhibitor

Acc1^{fl/fl}

CD4-Cre*Acc1*^{fl/fl}

PI

Apoptosis (%)

n.s   n.s

inhibitor   inhibitor

Figure S4

Figure S5

A

B

C