Water hardness variously influences tissue physiology of freshwater fish - *Cyprinus carpio*

***var koi*: A report on glucose, oxidative stress and antioxidant biomarkers**

Proteek Dasgupta*,

Department of Zoology,

Bangalore University, Jnana Bharathi Campus,

Bengaluru – 560056, Karnataka, India

Corresponding author - proteek.dasgupta@gmail.com

ORCID ID - 0000-0003-3962-0664

Abstract

Fishes endemic to freshwater habitat are strongly influenced by water hardness with

physiological consequences. The present study aimed to understand the effects of four-fold

sequential increase from soft to hard waters, on selected tissues of Koi carp, a popular

ornamental freshwater fish. Secondary stress markers - Glucose, Oxidative stress

(Malondialdehyde – Lipid Peroxidation damage) and Antioxidants (Catalase, Glutathione-S-

Transferase and Glutathione) were quantified in gills and white muscle after 14 days of exposure

to hardness of 75 (Soft - TS), 150 (Moderate - TM), 225 (Hard - TH) to 300 (Very Hard - TV)

mg $CaCO_3$/L. Both the examined tissues were distinctly affected by soft and moderate waters.

Glucose in gills ($p < 0.05$) was proportional to the rise in hardness concentration. Soft, moderate

and very hard waters (75, 150 and 300 mg $CaCO_3$/L) affected gills and muscle due to elevated

MDA concentrations ($p < 0.05$). CAT and GST provided antioxidative protection to the tissues.

The study results showed tissue-specific differential responses and more importantly,

concentrations below 225 mg $CaCO_3$/L elicited strong oxidative impairment in both gill and

muscle.

Keywords – Calcium; Carps; Gill; Muscle; Physiology; Water quality

Introduction

Hardness is perhaps the most significant physicochemical property of water because it directly or

indirectly influences physiology. Aquatic environments are constituted by varying concentrations

of hardness, majorly due to Calcium (Ca^{2+}) and Magnesium (Mg^{2+}) along with some trace

cations (Zn^{2+}, Mn^{2+}, etc) (Baldisserotto 2011; Romano et al. 2020). Based on the quantity of

major cations, water is classified into soft (< 75 mg $CaCO_3$/L) and hard (> 75 mg $CaCO_3$/L)

(Portz et al. 2006), both of which exert biological challenges. Soft water poses challenges to the

survival of fish causing efflux across ion channels and destabilising ionic balance while,

excessive hardness can cause hypercalcemia (Wendelaar Bonga et al. 1983) leading to bone

ossification (Blanksma et al. 2009). It is, therefore, quite clear that water hardness leads to

physiological changes. Without doubt, such changes can be studied through tissues which are

largely impacted (Gonzalez et al. 1998; Gundersen and Curtis 1995).

Gill and muscle have provided substantial information about adaptation to water hardness in

freshwater species such as Pinfish (*Lagodon rhomboides*) and Tilapia (Carrier and Evans 1976;

Flik and Verbost 1995; Wendelaar Bonga et al. 1983). While gills are the epicentre of Ca^{2+}

homeostasis and osmoregulation (Evans et al. 2005; Wendelaar Bonga et al. 1983), muscle is

susceptible to changes in constituent amino acids and ionic shifts due to water hardness

(Buentello and Gatlin 2002). Tendency of muscle mitochondria to take up Ca^{2+} from external

sources and compensate for ionic stress has been reported in *Tilapia mossambica* (Sulochana et

al. 1977). Both the tissues localise various redox reactions and such biomarker examinations

provide information about physiological adjustments due to hard waters (Lushchak 2011).

Secondary stress markers are instrumental in assessing the effects of hard and soft waters.

Reports by Copatti et al. (2019a) and Neves et al. (2017), highlight the usage of glucose to

evaluate the effects of water hardness. Malondialdehyde (MDA) is a marker of lipid peroxidation

(LPO), the consequences of which are prevented by enzymatic antioxidants (Catalase,

Glutathione-S-Transferase) or non-enzymatic (Glutathione) thereby, rectifying the

prooxidant/antioxidant ratio (Betteridge 2000; Lushchak 2016). Catalase is an important

antioxidant enzyme that protects cells and tissues from oxidative damage because it reduces

harmful hydrogen peroxide (H_2O_2) to water (H_2O) (Betteridge 2000). Activity of GST is specific

to detoxification of xenobiotics. It conjugates GSH to various electrophiles, thereby preventing

oxidative damage. Although, GSH can also independently scavenge free radicals to defend the

tissues from stress (Srikanth 2013). Despite broad insights offered by all the above, it is

noteworthy that investigations involving its usage have gained momentum recently (Copatti et al.

2019b; Michelotti et al. 2018) to understand the extent and efficiency of physiological

adaptations due to external hardness. Relevant information about the evaluation of secondary

markers can make way for further molecular insights.

Koi carp, an ornamental variety of common carp, is popular among aquarists for aesthetical

features. It is often spotted in various ponds, reservoirs, streams and lakes, which makes it an

ecologically relevant species to study hardness and physiology interactions. Further, this species

is physiologically sturdy under captive conditions as well. Therefore, the present study attempts

to evaluate the effects of water hardness (75, 150, 225 and 300 mg $CaCO_3$/L) on biomarkers

(glucose, oxidative stress and antioxidant profile) in gill and muscle of Koi carp.

Methods

Acclimation of fish

Juveniles (6.70 ± 0.15 g; 5.90 ± 0.12 cm) were purchased from the Ornamental Fish Research Centre, Bengaluru. Fish were randomly distributed in 50 L glass tanks filled with tap water. The tanks were maintained under natural photoperiod ($\approx$ 12 Light/12 Dark) with continuous aeration (Venus Aqua AP-608A, China). Fish were fed twice a day (09:00 and 18:00) at 2% body weight with commercial feed pellets (Taiyo Grow, India).

Experimental setup

The two-week study consisted of four levels of hardness: 75 (Soft - TS), 150 (Moderate - TM), 225 (Hard - TH), and 300 (Very Hard - TV) mg $CaCO_3$/L, selected based on its occurrence in natural systems (Stumm and Morgan 1996). Each treatment (hardness) was maintained in triplicate (6 fish per treatment). Hardness concentrations were prepared by the addition of Calcium carbonate ($CaCO_3$) to water and calibrated by complexometric EDTA titration (APHA 2005). Faecal matter accumulated in the tank was drained out by suction pipe with subsequent renewal of the specific hardness. The tanks were covered with a mesh net to prevent the fish from escaping. Water parameters were monitored for temperature (25.1 ± 1°C), pH (7.04 ± 0.1), dissolved oxygen (6.5 ± 0.08 mg/L) and alkalinity (213 ± 0.02 mg/L) (APHA 2005).

Sampling

Fish ($n = 6$ for each hardness level; total = 24 fish) were euthanized individually in clove oil solution (50 μL/L) and dissected for white muscle and gill. Tissues were washed with an ice-cold phosphate buffer solution (0.1M; pH 7.4) and homogenised in Potter-Elvehjem grinder. The homogenate (10% w/v) was then centrifuged at $5000 \times g$ following which supernatant (stored at

-20°C) was retained for all assays except Glutathione (homogenate precipitated with TCA before

centrifugation). The absorbance values were measured on a spectrophotometer (Systronics UV-

VIS 118, India).

Biochemical analyses

Glucose

Glucose was assayed according to Nelson (1944) and Somogyi (1952). The deproteinizing agent

$(Ba(OH)_2$ and $ZnSO_4)$ was added to the supernatant and centrifuged at $5000 \times g$ for 10 minutes.

Alkaline copper reagent (potassium-sodium tartrate; Na_2CO_3; $NaHCO_3$ and Na_2SO_4 in distilled

water) was added to the supernatant. The mixture was heated, followed by the addition of an

arseno-molybdate reagent. Optical density of the solution was recorded at 540 nm. A standard

curve was plotted to evaluate the glucose concentration.

Malondialdehyde (MDA)

Secondary product of Lipid peroxidation (LPO) - Malondialdehyde was estimated by using the

Niehaus and Samuelsson (1968) protocol. The supernatant was mixed with this TCA-TBA-HCl

reagent (15% Trichloroacetic acid, 0.38% Thiobarbituric acid and 0.25N Hydrochloric acid) in

the ratio of 1:2. This reaction mixture was heated in a boiling water bath for 15 minutes, cooled

and centrifuged at $1100 \times g$ for 5 minutes. The optical density of the solution was recorded at

535 nm. MDA concentration was calculated using extinction coefficient of $1.56 \times 10^5 \, M^{-1}$ cm.

Catalase (CAT)

Catalase activity was measured by using the Aebi (1984) protocol. The reaction was started by

adding the supernatant to an equimolar solution of H_2O_2 and phosphate buffer (50 mM; pH 7.1).

A decrease in absorbance was continuously recorded at 240 nm (UV) for 3 min.

Glutathione-S-Transferase (GST)

127   GST activity was estimated by using the Habig et al. (1974) protocol. The reaction mixture

128   contained the supernatant, phosphate buffer (0.1M; pH 6.5) and 2,4-Dinitrochlorobenzene

129   (30mM). The volume was adjusted with distilled water after which the reaction was initiated by

130   adding Glutathione (0.1 M). The optical density of the solution was recorded at 340 nm using a

131   molar extinction coefficient of $9.6 \times 10^3 M^{-1} cm^{-1}$.

Glutathione (GSH)

133   GSH was estimated by using the Moron et al. (1979) protocol. Homogenate was precipitated

134   with TCA (5%) and centrifuged at $3000 \times g$ for 10 minutes. The supernatant collected after

135   centrifugation was then added to the phosphate buffer (pH 6.5) and Ellman's reagent. The optical

136   density of the solution was recorded at 420 nm.

Total protein

138   Total protein content of the tissues was estimated according to Lowry et al. (1951) protocol.

139   Bovine serum albumin was used as standard. The optical density of the supernatant-reagents

140   mixture was recorded at 660 nm.

Statistical analysis

Data was summated as Mean ± SE. Normality and homoscedasticity were evaluated using the

Shapiro–Wilk and Levene tests, respectively. Inter-group comparisons were performed using

One-way ANOVA followed by a post-hoc test (Tukey). Significant differences were fixed at 95

% ($p < 0.05$). GraphPad Prism (Version 5.0, USA) and JASP (Version 0.16.2, Netherlands) were

used for statistical computation and visual presentations.

Results

Biomarkers in Gill

The glucose concentration increased progressively from TS to TV. Significant differences ($F =$

10.91; $p < 0.05$) were found between TV and the remaining treatments (Figure 1A). Soft water

TS elevated the concentration of MDA, followed by a spike in TV. Only TH differed

significantly ($F = 21.27$; $p < 0.001$) from the remaining treatments (Figure 2A). Highest

antioxidant Catalase activity was observed for TH, which differed significantly ($F = 50.26$; $p <$

0.001) from TS, TM and TV. Also, TV differed significantly from TM and TS ($F = 50.26$; $p <$

0.001) (Figure 3A). GST activity for TH and TV was comparatively higher than for TS and TM.

While no significant differences ($F = 26.45$; $p > 0.05$) were found between the low (TS and TM)

and high hardness groups (TH and TV), differences were observed between treatment pairs

(Figure 4A). The highest concentration of GSH was recorded for TV. Except TS that was not

significant ($F = 42.78$; $p > 0.05$) with TM and TH, remaining treatments recorded intergroup

differences (Figure 5A).

Biomarkers in White Muscle

The glucose concentration progressively increased from TS to TH but reduced at TV. Pairwise,

similar concentrations and no significant differences ($F = 22.98$; $p > 0.05$) were observed for TS

and TV and for TM and TH (Figure 1B). The highest level of MDA was observed for TM which

differed significantly ($F = 37.89$; $p < 0.001$) from TS, TH and TV (Figure 2B). The highest CAT

activity was observed in TM followed by a decrease for TH and TV. Catalase was the only

biomarker that varied significantly ($F = 49.38$; $p < 0.05$) among all treatments for white muscles

(Figure 3B). No significant differences ($F = 57.13$; $p > 0.05$) were recorded for GST between

TM and TH. The highest GST activity was recorded for TH, which differed significantly with TS

and TV ($F = 57.13$; $p < 0.001$) (Figure 4B). Glutathione (GSH) initially increased from TS to

TM, thereafter, reducing gradually for the remaining treatments. Only significant difference ($F =$

8.87; $p < 0.05$) was observed between TM and the remaining groups (Figure 5B).

Discussion

Glucose concentration

In the present study, glucose increased sequentially in gills indicating that it was more conserved

at higher levels of hardness. Progressive hardness led to increase in glucose, probably adding to

the energy reserves. Since glucose serves as a primary energy for metabolism, glucose estimation

can provide insights into the energy consumption for adaptations to hardness. Freshwater fish

take up Ca^{2+} through the gills, and this transcellular movement is dependent on the surrounding

Ca^{2+} concentrations, which affect the branchial permeability of the gills (Flik and Verbost 1995).

Generally, an environment with high hardness reduces gill permeability (through tightening of

cellular junctions) and subsequent loss of ions to water, ultimately conserving energy

(Golombieski et al. 2013). This is clear through the results of the present study. Contrarily,

muscle showed reductions in glucose at TV (300 mg $CaCO_3$/L), which was also observed in

juvenile Common Snook (*Centropomus undecimalis*) exposed to elevated hardness (Michelotti et

al. 2018). Probably, increased energy demands for energy lowered muscle glucose levels, and it

might have upregulated glycolysis.

Oxidative stress

Presence of enormous amounts of Polyunsaturated Fatty Acids (PUFAs), predisposes fishes to

lipid peroxidation, ultimately damaging cell membrane (Lushchak 2011) which is proportional to

MDA produced. In the present study, excluding 225 mg $CaCO_3$/L, MDA concentration for

remaining exposures was elevated in gills. Possibly, soft (75), moderate (150) and very hard

(300) waters, lead to damage of membrane associated with gills and white muscle. Subsequently,

this could have led to excess production of MDA through free radical generation. In

environments with low ionic concentration, certain membranes (such as apical membrane of gill)

mechanise uptake of divalent cations from hard waters through Ca^{2+} channels embedded in it to

meet the demand for necessary biological processes (Limbaugh et al. 2021). Elevated MDA is

indicative of membrane damage and cell injury.

Antioxidant response

Low catalase activity in gills exposed to soft (75) and moderate (30) waters indicated the failure

to prevent oxidative damage. Low ionic composition of freshwater environment might burden

the gills to reduce loss of ions to water for maintaining osmotic integrity. Therefore, efflux of

ions to the external environment with concentrations < 150 mg $CaCO_3$/L can ultimately lead to

oxidative stress. Contrarily, muscle showed sequential decrease in antioxidant activity exposures

above 150 mg $CaCO_3$/L, largely remaining unaffected at higher concentrations.

Increase in GST activity was noticed in gills exposed to hard (225) and very hard waters (300).

On the contrary, muscle showed elevated GST activity for all the exposures except soft waters

(75). Though, higher GST activity clearly indicated greater antioxidant capacity in both the

tissues, by far, antioxidative response was greater in muscle than gills. However, hardness < 75

mg $CaCO_3$/L is detrimental for both the tissues and might impact osmoregulatory functioning

(Evans 1987). Also, compared to Catalase, activity of GST was much higher in both the tissues

for exposures above 225 mg $CaCO_3$/L, indicating better antioxidant activity at higher levels of

water hardness.

Antioxidant GSH was relatively low for muscle in contrast to gills, indicating better antioxidant

capacity of muscle. In gills, a spike was observed for TH, which is conclusive that anything

above 300 mg $CaCO_3$/L is harmful. GSH can scavenge free radicals independently or in

conjunction with GST to provide antioxidant defence (Srikanth et al. 2013). Clearly, the study

results showed variance in GSH for both the tissues conforming to its specificity or tissue-

specific antioxidant response. This has previously been reported in other popular freshwater

species such as Nile Tilapia, Sharp Tooth Catfish (*Clarias lazera*) and Common carp (*Cyprinus*

carpio) (Hamed et al. 2004).

Conclusion

The study results reveal that soft and moderate (75 and 150 mg $CaCO_3$/L) hardness can alter

physiology of this freshwater species prompting antioxidative activity. Even after a prolonged

period of 14 days, fish showed considerable adaptability to all exposures. Further, glucose

concentration in gills was reserved for exposures to hardness > 225 mg $CaCO_3$/L, to cope up in

environments with low hardness. The results would benefit stakeholders and aquarist who
culture this valued commercial species.

List of abbreviations -

Malondialdehyde - MDA; Catalase - CAT; Glutathione-S-Transferase - GST; Glutathione GSH;
Glucose - GLU; Soft water - TS; Moderately hard water - TM; Hard water - TH; Very hard water
- TV

Ethics approval – Animal use within this study was approved by the welfare committee of
Bangalore University, India, in accordance with the national legislation.

Availability of data and material – All relevant data is within this article. Please contact
corresponding author for further queries

Conflict of interest – The author declares that there are no known competing financial interests
or personal relationships that could have appeared to influence the work reported in this paper.

Funding – This research did not receive any grant from funding agencies in the public,
commercial or not-for-profit sectors.

Acknowledgements – Author thanks Bangalore University for providing necessary
infrastructure.

References –

1. Aebi H (1984) Catalase in vitro. Methods in Enzymology 105:121–126. DOI:
https://doi.org/10.1016/s0076-6879(84)05016-3

2. APHA (2005) Standard methods for the examination of water and waste-water (21st ed)
Washington, DC: American Public Health Association.

3. Baldiserotto B (2011) Water pH and hardness affect growth of freshwater teleosts
Revista Brasileira de Zootecnia (40) 138-144

4. Betteridge DJ (2000) What Is Oxidative Stress? Metabolism 49(2):3-8. DOI:

10.1016/s0026-0495(00)80077-3.

5. Blanksma, C., Eguia, B., Lott, K., Lazorchak, J.M., Smith, M.E., Wratschko, M.,

Dawson, T.D., Elonen, C., Kahl, M., Schoenfuss, H.L., 2009. Effects of water hardness

on skeletal development and growth in juvenile fathead minnows. Aquaculture 286, 226–

232.

6. Buentello, J.A. and Gatlin, D.M., III (2002), Preliminary Observations on the Effects of

Water Hardness on Free Taurine and Other Amino Acids in Plasma and Muscle of

Channel Catfish. North American Journal of Aquaculture, 64: 95-102.

https://doi.org/10.1577/1548-8454(2002)064<0095:POOTEO>2.0.CO;2

7. Copatti CE, Baldisserotto B, Souza CF, Garcia LO (2019a) Protective effect of high

hardness in pacu juveniles (*Piaractus mesopotamicus*) under acidic or alkaline pH:

Biochemical and haematological variables. Aquaculture.; 502(1):250–57.

https://doiorg/10.1016/j.aquaculture.2018.12.028

8. Copatti CE, Baldisserotto B, de Freitas Souza C, Monserrat MJ, Garcia L (2019b) Water

pH and hardness alter ATPases and oxidative stress in the gills and kidney of pacu

(*Piaractus mesopotamicus*). Neotropical Ichthyology 17(4):e190032. DOI:

10.1590/1982-0224-20190032

9. Carrier JC, Evans DH. (1976) The role of environmental calcium in freshwater survival

of the marine teleost, *Lagodon rhomboides*. J Exp Biol. 1976;65(3):529-538.

10. Evans DH. The fish gill: site of action and model for toxic effects of environmental

pollutants. Environmental Health Perspectives. 1987; 71(1):54–58.

https://doi.org/10.1289/ehp.877147

273 11. Evans DH, Piermarini PM, Choe KP (2005) The multifunctional fish gill: Dominant site

274  of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous

275  waste. Physiol Rev. 85(1):97-177. https://doi.org/10.1152/ physrev.00050.2003

276 12. Flik, G., Verbost, P.M., 1995. Cellular mechanisms in calcium transport and homeostasis

277  in fishes. Biochemistry and Molecular Biology of Fishes 5, 252–263.

278 13. Golombieski J, Koakoski G, Becker A, Almeida A, Toni C, Finamor I, Pavanato M,

279  Almeida T, Baldisserotto B (2013) Nitrogenous and phosphorus excretions in juvenile

280  silver catfish (*Rhamdia quelen*) exposed to different water hardness, humic acid, and pH

281  levels. Fish Physiology and Biochemistry 39:837-849. DOI 10.1007/s10695-012-9744-8

282 14. Gonzalez RJ, Wood CM, Wilson RW, Patrick ML, Bergman HL, Narahara A, Val AL.

283  Effects of water pH and calcium concentration on ion balance in fish of the Rio Negro,

284  Amazon. Physiol Zool. 1998 Jan-Feb;71(1):15-22. doi: 10.1086/515893. PMID:

285  9472808.

286 15. Gundersen DT and Curtis LR (1995) Acclimation to hard or soft water at weakly alkaline

287  pH influences gill permeability and gill surface calcium binding in rainbow trout

288  (*Oncorhynchus mykiss*) Publication: Canadian Journal of Fisheries and Aquatic Sciences.

289  https://doi.org/10.1139/f95-848

290 16. Habig WH, Pabst MJ, Jakoby WB (1974) Glutathione S-transferases the first enzymatic

291  step in mercapturic acid formation. Journal of Biological Chemistry 249(22):7130–7139.

292 17. Hamed R, Maharem T, Guinidi R (2004) Glutathione and its related enzymes in the Nile

293  fish. Fish Physiol Biochem 30:189–199

294 18. Limbaugh N, Romano N, Egnew N, Shrivastava J, Bishop WM, Sinha AK (2021) Coping

295  strategies in response to different levels of elevated water hardness in channel catfish

(*Ictalurus punctatus*): Insight into ion-regulatory and histopathological modulations.

Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology

260:111040. doi: 10.1016/j.cbpa.2021.111040.

19. Lowry OH, Rosebrough NJ, Farr AL, Randal RJ (1951) Protein measurement with the

Folin-phenol reagent. The Journal of Biological Chemistry 193:265–275.

20. Lushchak IV (2011) Environmentally induced oxidative stress in aquatic animals.

Aquatic Toxicology 101:13–30. DOI: 10.1016/j.aquatox.2010.10.006

21. Lushchak IV (2016) Contaminant-induced oxidative stress in fish: a mechanistic

approach. Fish Physiol Biochem 42:711–747 DOI 10.1007/s10695-015-0171-5

22. Michelotti BT, Passini G, Carvalho C, Salbego J, Mori NC, Rodrigues RV, Baldisserotto

B, Cerqueira VR (2018) Growth and metabolic parameters of Common Snook juveniles

raised in freshwater with different water hardness. Aquaculture 482:31–35. DOI:

https://doi.org/10.1016/j.aquaculture.2017.08.029

23. Moron MS, Depierre JW, Mannervik M (1979) Levels of glutathione, glutathione

reductase and glutathione S-transferase activities in rat lung and liver. Biochimica et

Biophysica Acta (BBA) - General Subjects 582(1):67-78. DOI: 10.1016/0304-

4165(79)90289-7

24. Nelson N (1944) A Photometric Adaptation of the Somogyi Method for Determination of

Glucose. Journal of Biological Chemistry, 153:375-380.

25. Neves GC, Souza CF, Santos AC, Baldisserotto B, Lopes JM (2017) Waterborne calcium

and nitrite interaction: survival, growth, haematological and metabolic parameters in

silver catfish. Boletim do Instituto de Pesca, São Paulo 43 (3), 408–416.

https://doi.org/10.20950/1678-2305.2017v43n3p408.

26. Niehaus WG, Samuelsson B (1968) Formation of malonaldehyde from phospholipid
arachidonate during microsomal lipid peroxidation. European Journal of Biochemistry
6(1):126–130. DOI: 10.1111/j.1432-1033.1968.tb00428.x.

27. Portz DE, Woodley CM, Cech JJ (2006) Stress-associated impacts of short-term holding
on fishes. Rev Fish Biol Fisheries 16, 125–170 https://doi.org/10.1007/s11160-006-9012-
z

28. Romano N, Egnew N, Quintero H, Kelly A, Sinha AK, (2020) The effects of water
hardness on the growth, metabolic indicators and stress resistance of largemouth bass
Micropterus salmoides. Aquaculture 527:735469

29. Somogyi M (1952) Notes on sugar determination. Journal of Biological Chemistry
195(1):19-23.

30. Srikanth K, Pereira E, Duarte AC, Ahmad I. (2013) Glutathione and its dependent
enzymes' modulatory responses to toxic metals and metalloids in fish-a review. Environ
Sci Pollut Res Int. 20(4):2133 - 2149 doi:10.1007/s11356-012-1459-y

31. Stumm W, Morgan JJ (1996) Aquatic chemistry: chemical equilibria and rates in natural
waters. (3rd Ed.), pp 1022, John Wiley and Sons, New York.

32. Sulochana, H.R., Bashyam, I., Narayan, S. et al. Physiological correlates of calcium-
accumulating properties of mitochondria: Fish-muscle mitochondria. J Bioenerg
Biomembr 9, 337–348 (1977). https://doi.org/10.1007/BF00743149

33. Wendelaar Bonga SE, Lowik CJM, Van der Meij JCA (1983) Effects of external Mg^{2+}
and Ca^{2+} on branchial osmotic water permeability and prolactin secretion in the teleost
fish *Sarotherodon mossambicus*. General and Comparative Endocrinology 52:222-231

Figure Legends -

Figure 1 – Effect of hardness concentration of 75 (Soft - TS), 150 (Moderate - TM), 225 (Hard - TH) and 300 (Very Hard - TV) mg $CaCO_3$/L on glucose concentration in **(A)** gill and **(B)** white muscle of Koi carps ($n = 24$; 6 fish × 4 treatments). Superscripts among exposure groups indicate statistical significance ($p < 0.05$). (+) inside the box indicates the mean value of the exposure group.

Figure 2 – Effect of hardness concentration of 75 (Soft - TS), 150 (Moderate - TM), 225 (Hard - TH) and 300 (Very Hard - TV) mg $CaCO_3$/L on MDA concentration in **(A)** gill and **(B)** white muscle of Koi carps ($n = 24$; 6 fish × 4 treatments). Superscripts among exposure groups indicate statistical significance ($p < 0.05$). (+) inside the box indicates the mean value of the exposure group.

Figure 3 – Effect of hardness concentration of 75 (Soft - TS), 150 (Moderate - TM), 225 (Hard - TH) and 300 (Very Hard - TV) mg $CaCO_3$/L on Catalase activity in **(A)** gill and **(B)** white muscle of Koi carps ($n = 24$; 6 fish × 4 treatments). Superscripts among exposure groups indicate statistical significance ($p < 0.05$). (+) inside the box indicates the mean value of the exposure group.

Figure 4 – Effect of hardness concentration of 75 (Soft - TS), 150 (Moderate - TM), 225 (Hard - TH) and 300 (Very Hard - TV) mg $CaCO_3$/L on Glutathione-S-transferase (GST) activity in **(A)** gill and **(B)** white muscle of Koi carps ($n = 24$; 6 fish × 4 treatments). Superscripts among exposure groups indicate statistical significance ($p < 0.05$). (+) inside the box indicates the mean value of the exposure group.

Figure 5 – Effect of hardness concentration of 75 (Soft - TS), 150 (Moderate - TM), 225 (Hard - TH) and 300 (Very Hard - TV) mg $CaCO_3$/L on Glutathione (GSH) in **(A)** gill and **(B)** white muscle of Koi carps ($n = 24$; 6 fish × 4 treatments). Superscripts among exposure groups indicate

369   statistical significance ($p < 0.05$). (+) inside the box indicates the mean value of the exposure

370   group.

373   **Figure 1A & B**

374
375

383   **Figure 2A & B**

390 **Figure 3A & B**

397   **Figure 4A & B**

403   **Figure 5A & B**