Restructuring of olfactory representations in the fly brain around odor relationships in natural sources

Jie-Yoon Yang[1,2], Thomas F. O'Connell[1,2], Wei-Mien M. Hsu[3], Matthew S. Bauer[2], Kristina V. Dylla[2], Tatyana O. Sharpee[3], Elizabeth J. Hong [2,4]

[1] These authors contributed equally
[2] Division of Biology & Biological Engineering, California Institute of Technology, Pasadena, CA, USA
[3] Computational Neurobiology Laboratory, Salk Institute for Biological Studies, La Jolla, CA, USA; Department of Physics, University of California, San Diego, La Jolla, CA, USA
[4] Lead contact

ABSTRACT

A core challenge of olfactory neuroscience is to understand how neural representations of odor are generated and progressively transformed across different layers of the olfactory circuit into formats that support perception and behavior. The encoding of odor by odorant receptors in the input layer of the olfactory system reflects, at least in part, the chemical relationships between odor compounds. Neural representations of odor in higher order associative olfactory areas, generated by random feedforward networks, are expected to largely preserve these input odor relationships[1–3]. We evaluated these ideas by examining how odors are represented at different stages of processing in the olfactory circuit of the vinegar fly *D. melanogaster*. We found that representations of odor in the mushroom body (MB), a third-order associative olfactory area in the fly brain, are indeed structured and invariant across flies. However, the structure of MB representational space diverged significantly from what is expected in a randomly connected network. In addition, odor relationships encoded in the MB were better correlated with a metric of the similarity of their distribution across natural sources compared to their similarity with respect to chemical features, and the converse was true for odor relationships encoded in primary olfactory receptor neurons (ORNs). Comparison of odor coding at primary, secondary, and tertiary layers of the circuit revealed that odors were significantly regrouped with respect to their representational similarity across successive stages of olfactory processing, with the largest changes occurring in the MB. The non-linear reorganization of odor relationships in the MB indicates that unappreciated structure exists in the fly olfactory circuit, and this structure may facilitate the generalization of odors with respect to their co-occurence in natural sources.

INTRODUCTION

The search for organizing principles of olfaction has often focused on relating the chemical structure or physicochemical properties of odorants to their percept[4]. This approach is principled since odors are detected by their molecular interactions with large families of structurally diverse odorant receptor (OR) proteins expressed in ORNs[5]. Recently, significant inroads have been made in predicting a molecule's odor from its structure[6,7] but developing a generalized relationship between odorant structure and perception across the space of all possible odor stimuli remains challenging because of discontinuities in this relationship: small changes in

45 structure often result in dramatic changes in a molecule's odor[8,9]. This gap in understanding
46 motivates a search for additional organizational axes of odor space to complement structure-
47 based approaches towards gaining a better understanding of what determines a molecule's
48 smell.
49
Another important property of odorants is how they are organized relative to one another in
natural environments. Odors from natural sources are typically complex mixtures of dozens to
hundreds of monomolecular odorants, the composition of which is controlled by the conserved
biochemical and metabolic processes in the source[10,11]. The relative abundance or ratios of
volatiles in natural odor profiles can provide information about the value or state of the odor
source[12,13], for instance, if microbes that promote fermentation or spoilage are dominant. Thus,
the odor space of the natural world is highly structured, and this structure often contains
information about the identity or ethological value of the odor source.
58
We investigated how representations of odor at different stages of processing in the brain of the
vinegar fly *Drosophila melanogaster* relate to different odor properties, focusing on their
chemical properties or their relative abundances in behaviorally significant natural odor sources
like food. The fly has a compact olfactory system with a similar overall circuit architecture to its
vertebrate counterpart[14]. All ORNs that express the same OR project to a common synaptic
compartment, called a glomerulus, in the antennal lobe (AL), and the dendrites of uniglomerular
second-order projections neurons (PNs) extend into a single glomerulus[15]. Thus, each
glomerulus, corresponding to a specific OR, represents a fundamental unit of olfactory
processing. A major target of PN output from the AL is the mushroom body (MB), a cerebellum-
like associative center in the fly brain that encodes representations of odor identity[16]. Wiring of
PN inputs to Kenyon cells (KCs), the principal neurons of the MB, is probabilistic: each of the
~2000 KCs integrates input from a subset of PNs comprising ~10% of the ~50 olfactory
glomeruli in the system[17–19]. KCs have high spiking thresholds and act as coincidence detectors
that fire only when multiple input PNs are co-active[20,21], and local feedback inhibition between
KCs is provided from the arborizations of an unusual single, large GABAergic neuron called the
APL[22].
75
This circuit architecture recodes dense, distributed representations of odor in the ~50 glomeruli
of the PN layer into a sparse, high-dimensional representation in the MB layer[3,23–25] that
facilitates pattern separation and linear decoding by a smaller number of MB output neurons.
Theoretical studies of cerebellum-like circuits, characterized by expansion (PN input onto KCs)
and reconvergence (KC output onto MB output neurons), emphasize the role of random,
unstructured input for decorrelating activity patterns and maximizing the dimensionality of
representations[26,27]. Such features would promote efficient memory storage and reduced
synaptic inference during stimulus-specific associative learning and recall.
84
Feedforward random network models of the MB predict that odor relationships encoded in the
MB should be strongly decorrelated, invariant across individual brains, and should preserve
stimulus relationships encoded at the level of ORN input [1,28–30]. However, recent large-scale EM
reconstructions of MB synaptic connectivity demonstrated that some PN inputs are structured.

In particular, inputs from glomeruli tuned to odors common in food are more likely to converge
onto the same KC targets[19], though the functional impact on MB representations of odor
remains to be determined. Whereas random networks maximize coding capacity and promote
the separability of odor representations throughout odor space, structured networks can
correlate specific odor representations to promote generalization between odors sharing
particular ethological meaning. We investigated how odor coding in the fly olfactory circuit
balances these competing needs.
96
97
98   **RESULTS**
99
Population imaging of odor representations in the MB at cellular resolution
In the arc of sensorimotor transformation, representations of odor identity encoded in KC activity
patterns represent the output from the sensory arm of the pathway, which is flexibly coupled to
distinct downstream outputs and behaviors[31]. Thus, we began by investigating representations
of odor in the MB. The *Drosophila* olfactory circuit is the most comprehensively mapped
metazoan olfactory system, with the tuning of approximately half of the odorant receptors to a
large panel of 109 odors described by the Hallem dataset[32]. We selected 24 monomolecular
odors that spanned OR input space (Figure 1A) and investigated the extent to which those
relationships at the periphery are maintained in the MB.
109
We used in vivo volumetric two-photon microscopy to image odor-evoked calcium signals in the
MB of flies expressing nuclear-localized GCaMP6s selectively in KCs (directed by the *OK107-*
Gal4 driver[33]) (Figure 1B-C). In pilot experiments using cytoplasmic GCaMP6f as the calcium
reporter, the small size (~2-3 μm) and tight, regular packing of KC somata (Figure 1E)
presented challenges for motion correction, good cellular segmentation, and reliable pixel
assignment to individual KCs over the course of an imaging session, with poor trial-to-trial
reliability in odor panels larger than 8 odors. To expand the size of odor panels that could be
evaluated in a single brain, we turned to measuring nuclear calcium, which has slower response
dynamics, but strongly correlated response amplitudes compared to cytoplasmic calcium[34,35].
Localization of the calcium indicator to nuclei resulted in a several pixel gap between KCs that
facilitated reliable cellular segmentation (Figure 1E, H) and enabled recording of the
representations of between 8 to 17 odors in the same MB.
122
Flies were presented odors in pseudo-random sequences while rapidly z-scanning through the
KC cell body layer. Following volumetric motion correction, odor-evoked KC signals were
extracted using the Suite2P software package[36]. In brief, after correcting for motion in each
plane, regions of interests (ROIs) representing each KC were extracted (Figure 1H). Although
cell detection in Suite2P is usually neural activity-based, we extracted the spatial footprint of
each cell by performing anatomical segmentation on time-averaged images, resulting in
detection of between ~85-95% of the expected number of KCs. Since KC odor responses are
sparse and many KCs do not respond to any odor in even a relatively large panel, this
adjustment enabled reliable estimates of KC response rates. For a subset of experiments, odor
responses were registered to KCs across

Figure 1: Population representations of odor in the fly MB at cellular resolution.
A) t-SNE embedding of 109 odors based on the patterns of activity they elicit across 24 fly ORs in the Hallem dataset. A subset (open grey symbols) spanning the odor space was selected for measurement in the MB.

B) Odors were delivered to the antennae of immobilized flies expressing nuclear-localized GCaMP6s in all KCs, while imaging from KC somata with a two-photon microscope.

C) Imaging volumes comprising ~11 planes through the KC layer capture the activity of >85% KCs in an MB.

D) Configuration of volumetric imaging trial (3 Hz sweep rate).

E) Reconstruction of 3D ROIs corresponding to each KC from interleaved high-resolution anatomical imaging stacks. Scale bar, 10 μm.

F) Example odor-evoked calcium signals in an imaging block from a single KC.

G) Population representations in ~1800 KCs to 17 odors in the MB from a representative fly. Each row is a cell and each column is a trial. Cells are sorted by odor tuning.

H) Top: baseline fluorescence (left), weighted ROI masks of KCs (middle), and ROI lablels corresponding to individual KCs (right) from a single imaging plane in an MB. Bottom: odor-evoked patterns of KC activity in response to 1-pentanol, 1-hexanol, and ethyl propionate. Scale bar, 10 μm.

I) Matrix of pairwise correlation between KC population responses in individual odor trials, where trials are shown in presentation order (left) or grouped by odor (right). KC responses elicited by 1-hexanol were more similar to those elicited by 1-pentanol (green) than by ethyl propionate (orange).

Flies had genotype 20xUAS-nls-OpGCaMP6s-p10 (III)/+; OK107-Gal4 (IV)/+. Odors are pentyl acetate (pa), isoamyl acetate (iaa), methyl octanoate (mo), ethyl butyrate (eb), ethyl propionate (ep), 1-pentanol (1-5ol), 1-hexanol (1-6ol), 1-penten-3-ol (1-p3ol), 1-octanol (1-8ol), 2-heptanone (2-hep), 2-butanone (2-but), hexanal (6al), E2-hexenal (e2-h), benzaldehyde (bnz), methyl salicylate (msl), eugenol (eug), ethyl cinnamate (ec), linalool (lin), β-citronellol (b-cit), acetic acid (aac), propionic acid (pac), valeric acid (vac), γ-hexalactone (g-6lac), δ-decalactone (d-10lac).

133   different functional movies collected from the same MB by alignment of ROIs to 3D anatomical
134   models of KC somata constructed from high-resolution structural images through the MB (Figure
135   1E; Figure S1A).
136
Reproducible odor-specific response dynamics were observed in some cells (Figure 1F), but,
given the overall slow kinetics of nuclear-localized GCaMP6s, we focused our analysis in this
study on the peak amplitude of odor-evoked responses. KC population responses were stable,
odor-specific, and reliable across repeated trials of the same stimulus (Figure 1F-1G, Figure
S1A). The pairwise relationship between odor representations in MB coding space was
quantified using the correlation distance 1-r, where r is Pearson's correlation between the
vectors of KC responses to each pair of odor. Quantifying odor relationships using other metrics
such as cosine distance yielded similar results (e.g., Figure S3H). When we computed the
correlation distance between the KC response vector for every pairwise combination of trials in
an experiment (Figure 1I, left) and reordered the distance matrix to group together trials by odor,
we observed that KC responses to repeated presentations of the same odor were very strongly
correlated (on-diagonal blocks, Figure 1I, right). These results demonstrate the reliability of KC
odor responses across multiple presentations spanning the time course of an experiment. As
expected, the pairwise correlation distance between odors reflected the qualitative similarity of
their respective KC response patterns (Figure 1H-I), with visually similar activity patterns
corresponding to short odor distances.
153
154   The percentage of KCs responding to each odor was similar in each MB, ranging from ~5-13%
155   depending on the odor (Figure S1B). When compared against mean OR activity for each odor,
156   estimated by averaging the firing rates evoked by each odor across all ORs in the Hallem
157   dataset, KC response rate was not significantly correlated with mean ORN response strength
158   (Figure S1B, Spearman's rho=0.48, p=0.05). However, we note that the Hallem dataset

Figure 2: MB representational space is structured and invariant across individuals.
A) Correlation distance matrices for four different flies showing pairwise relationships between KC population responses in individual odor trials (upper triangles) or in trial-averaged responses for each odor (lower triangle).
B) Correlation distance between trial-averaged KC responses for every odor pair for the four flies in **A**. Odor pairs are in the same order in each row, arranged by the rank of their mean correlation distance across the four flies. Odors that evoke similar KC response patterns in one fly tend to also elicit similar response patterns in other flies.

C) Spearman's rank correlation between the rows of **B** (i.e., between flies), evaluating the similarity of the rank ordering of odor pairs between flies according to their representational distance in KCs.

D) Left: same as **C**, but for shuffled data in which the odor labels were randomly permuted for the responses of individual flies. The matrix shows the mean Spearman's correlation across 10,000 shuffles. Right: observed Spearman's correlation (black) and the mean and 95% CI of the Spearman's correlation across 10,000 shuffles (grey) for each fly pair. Red dotted line marks the mean observed Spearman's correlation (rho=0.76).

E) Correlation distance (mean and 95% CI) between KC responses for each odor pair, averaged across all flies in which the odor pair was sampled (n=3-22, see Supplemental Table 1). Each unique odor pair was assigned a reference ID (see Supplemental Table 1). A one-way ANOVA showed there was a significant difference in the odor-odor correlation distance between different odor pairs (F statistic=12.6, p=10^{-180}), consistent with odor-odor relationships being reliable across MBs in different flies. The correlation distance (mean and 95% CI across 100 model MBs) between predicted KC responses for odor pairs in the uniform, hemidraw, and hemibrain models are plotted for reference (see Figure 3).

F) Matrix of pairwise correlation distances between odor response profiles of every KC in the four flies in **A** that responded to at least one odor. The distance matrix was ordered by spectral clustering on the mean odor response vector of each cell. Each response cluster contained KCs from every fly.

G) Mean KC tuning profiles of each cluster, computed across KCs in each cluster in each fly. KCs with conserved odor tuning profiles are found in every MB.

159 underestimates ORN population responses to acids and amines since it does not include
160 odorant receptors from the ionotropic receptor (IR) family[37].
161
Overall, KCs were narrowly tuned, with most cells responding to two or fewer odors, and a
significant fraction (~34%) responding to no odor in a diverse 17-odor panel (Figure S2B).
However, compared against modeled KC responses (see below), observed odor responses in
KCs were more broadly tuned. This result is consistent with observations that existing MB
models poorly predict KC response rates to narrowly activating odors that selectively excite only
one or very few ORN classes (e.g., CO_2 or methyl salicylate, Figure S1D-E). These results
confirm that KC responses are sparse and selective. They also indicate that current
assumptions about MB circuit properties do not fully account for observed KC response rates to
all odors, particularly for narrowly activating odors.
171
Representations of odor in the MB are structured and invariant across individuals
In most circuits of the fly brain, neuronal connectivity is invariant across individuals, but the MB
is distinct in that the wiring of PN inputs is probabilistic: each of the ~2000 KCs, the principal
neurons of the MB, integrates input from a quasi-random subset of PNs comprising ~10% of the
~50 olfactory glomeruli in the system. As the number of possible glomerular combinations far
exceeds the number of KCs in any given MB, stereotyped KC connectivity, defined by a specific
set of synaptic inputs, does not exist across individual MBs. Indeed, a small set of genetically
defined KCs labeled by a sparse Gal4 driver did not exhibit stereotyped odor tuning[38]. However,
feedforward network models that assume random divergent connectivity between second- and
third-order olfactory layers predict that, while third-order olfactory responses will be
comparatively decorrelated, they will otherwise maintain relative pairwise odor relationships
present in the prior layer. Thus, these models predict that the geometry of third-order olfactory
representations will be invariant across different instantiations of the network (i.e., MBs), with
preserved odor relationships that are predictable from the stereotyped tuning of OR inputs.
186

To evaluate these ideas, we compared the correlational structure of odor representations
encoded in KC activity patterns in multiple flies. We found that the relationships between odors
in MB representational space were indeed structured and highly invariant across individual
brains (Figure 2A, S2A), consistent with prior work[3]. We evaluated the similarity of the
correlational structure across MBs by comparing the rank order of odor-odor correlation
distances in each MB (Figure 2B-C). The observed distribution of MB-MB correlations
(Spearman's *rho*=0.76 +/- 0.06) was significantly different from shuffled controls (Figure 2D, *p*
$<10^{-4}$). Also, the correlation distance between specific pairs of odors was consistent across
individual MBs: odors that evoked similar KC response patterns in one fly tended to also have
similar KC responses in other flies, and the same was observed for odors that evoked dissimilar
representations (Figure 2E, S2B; *p*=10^{-186}, one-way ANOVA). Finally, we combined all KCs that
respond to one or more odors from four different flies and clustered the KCs based on each
cell's odor response profile. This analysis identified reliable response types with specific odor
tuning profiles, and each response type was found in each of the four flies (Figure 2F). For
instance, KCs that are strongly co-tuned to 2-heptanone, isoamyl acetate, and pentyl acetate
were reliably observed in every MB (Figure 2G). These results demonstrate that the structure of
the representational space of odors is highly conserved across individual MBs in the fly.
204
Odor distances in MB coding space diverge from odor relationships predicted by a
random feedforward network model
To compare the observed structure of MB representational space to what is predicted by a
feedforward random network, we modeled KC population responses by adapting a previously
described, biologically plausible spiking model of the fly olfactory circuit[30] (Figure 3A). PN
responses, modeled from OR firing rates in the Hallem dataset[32], were used as input to a
population of 2000 spiking KCs. KCs were modeled as leaky integrate-and-fire units with a small
number of input sites, typically ~5-7 dendritic claws, that each receive input from a single PN
bouton. Under assumptions of random feedforward connectivity, the matrix of PN-KC
connections in each model instantiation was created by assigning each KC claw a single
glomerular input, with all glomeruli having an equal likelihood of being drawn ("uniform" model).
The inputs to the model were the firing rates of approximately half of fly ORs to 109 odors from
the Hallem dataset. KC activity was normalized by global feedback inhibition from the
GABAergic APL. KC spiking thresholds and APL-KC inhibitory weights were tuned to achieve a
mean KC response rate of ~10% across odors (see Methods), which matches experimental
observations.
221
The correlation between modeled KC responses for specific odor pairs was indeed consistent
across 100 simulated MBs with different, independently drawn random PN-KC connectivity
matrices (Figure 3B, error bars are 95% CI across simulations). However, the observed
structure of odor representations in KC coding space was only partially predicted by the
feedforward random model, with the observed relationship between odors deviating significantly
from model predictions for many odor pairs (Figure 2E, 3C, Di, S3H). In particular, observed
pairwise odor relationships were overall more decorrelated than those predicted from modeled
KC populations (Figure 3Di). Adjusting fit parameters to yield a lower mean KC response rate of
5% across odors did not appreciably

Figure 3: Odor relationships in MB representational space deviate from predicted relationships in a random feedforward network

A) Schematic of the fly olfactory network. Not depicted is a single GABAergic neuron (APL) in the MB that mediates feedback inhibition among KCs.

B) Comparison of relationships between predicted KC representations (uniform model) and observed OR tuning profiles for each odor pair (grey symbols). Error bars are 95% CI of KC correlation for each odor pair across 100 model MBs; each simulated MB has an independently generated PN-KC connectivity matrix drawn under assumptions of uniform input (each glomerulus drawn with equal probability).

C) Matrix of pairwise correlation distances between OR tuning profiles from the Hallem dataset **(Ci)**; predicted KC responses in the uniform model (**Cii**, left) or hemibrain model (**Cii**, right); or observed KC responses (**Ciii**) for 17 odors.

D) Comparison of correlation distances for each odor pair between observed KC responses for each odor pair and predicted KC responses in the uniform (**Di**) and hemibrain (**Dii**) models. Each symbol is an odor pair.

231   affect the systematic overprediction of the degree of correlation between odor representations,
232   nor did it affect the rank order of predicted odor-odor correlation distances (data not shown).
233
Recent analysis of the global structure of glomerular input sampling by third-order olfactory
neurons (Figure S3A) using large-scale EM-level reconstructions of part of the *Drosophila* brain
at synaptic resolution revealed that the wiring of PN inputs to KCs is not fully random[18,19], as

was previously believed. Indeed, analysis of the complete matrix of PN-KC connectivity from
two independently reconstructed fly brains confirmed that KCs sample particular combinations
of PN inputs at a higher rate than is expected based on their numerical frequency, and the
structure of this biased input onto KCs is similar between two MBs from different flies[19] (Figure
S3B-E). We also observed that the glomerular input structure to KCs bears significant similarity
to the structure of glomerular input to third-order olfactory neurons in the lateral horn (Figure
S3E), a brain region dedicated to innate odor processing in which neurons have stereotyped
connectivity and tuning[39,40].

245
To assess the possible impact of structured input on MB odor representations, we predicted KC
odor responses in the model under conditions in which the PN-KC connectivity matrix was
drawn according to the observed frequency of PN boutons corresponding to each glomerulus in
the hemibrain dataset ("hemidraw") or in which the experimentally reconstructed hemibrain PN-
KC connectivity matrix was directly implemented ("hemibrain") (see Methods). Neither
adjustment to PN-KC connectivity improved predictions of observed KC odor relationships
(Figure 3C, 3Dii, S3F-H); in fact, predicted responses for most odor pairs tended towards being
more correlated compared to the uniform model. These results indicate that observed biases in
PN-KC connectivity are unlikely to account for the differences between observed and predicted
KC odor responses; they suggest additional sources of structure are present in the olfactory
circuit that mediate the observed transformation of odor representations from the periphery to
the MB.

258
Reorganization of representations of odor in the MB around odor relationships in natural
sources
Comparisons of odor representations in the OR input layer and in the MB showed that, although
the representations of most pairs of odors were substantially decorrelated, as expected, some
odor pairs were comparatively less decorrelated between the OR and MB layers. This
transformation resulted in a significant regrouping of odor relationships in OR versus MB coding
space. For instance, the odors 2-heptanone, isoamyl acetate, and pentyl acetate emerged as a
cluster with similar KC representations, distinct from other odors in the panel, whereas the
representation of each of these odors is similar to many others at the level of its OR
representation (Figure 3Ci, iii).

269
To better understand the functional implications of this reorganization, we asked how odor
representations at different stages of olfactory processing relate to the properties of the odors,
focusing in particular on their chemical properties and on how they are correlated across natural
odor sources. For each odor, we computed molecular and physicochemical descriptors using
Mordred, an open-source molecular descriptor calculation software[41]. Since many descriptors
are highly correlated across odors, we identified a reduced set of 570 molecular descriptors that
captured the chemical relationships between odors equivalently to the full set of descriptors
(Supplemental Table S3).

278
279 *D. melanogaster* is an ecological generalist and human commensal[42]; as a starting point for
280 understanding the structure of natural odor space for *Drosophila*, we used a database of the

Figure 4: Reorganization of odor representations in the fly MB reflects odor relationships in natural sources.

A) Distribution of odors (monomolecular compounds, rows) in a literature-based database of the volatile headspace composition of many natural food odor sources (columns). The pie chart shows the distribution of odor sources in each food category.

B) Schematic summary of datasets for comparing the properties of odors (top) with their different neural representations (bottom).

C) Comparisons of odor-odor correlation distances in OR (left) or KC (right) coding space with their distances in chemical descriptor space.

D) Comparisons of odor-odor correlation distances in OR (left) or KC (right) coding space with their embedded distances in a hyperbolic model of natural source space.

E) Summary of correlation (lower triangle) and *p*-values (upper triangle) for comparisons between observed OR, observed KC, or modeled KC (uniform or hemibrain) odor relationships and odor-odor relationships in chemical or natural source space.

headspace volatile profiles for many food odor sources, compiled from 4407 published chemical
datasets primarily from the food and flavor science literature (Figure 4A). The odor sources are
biased towards fruits, plants, and vegetables, but also includes other common human foods
such as alcoholic beverages, meat, and dairy products. The database comprises thousands of
samples measured from 887 types of natural sources (apple, tomato, wine, etc.) and contains
over 8,000 monomolecular volatiles. Like other datasets profiling the chemical volatiles emitted
from natural sources[43], the majority of odorants occur sparsely in a small number of sources,
though a significant minority of odors are present broadly across many sources (Figure 4A).
289
Recently, we showed that the use of hyperbolic coordinates to embed the concentrations of
individual monomolecular odorants as they occur in natural odor sources better captures odor
relationships in the space of natural sources, compared to embeddings that use Euclidean
metrics[12,45]. Thus, to describe the similarity in distribution across natural odor sources for a
given pair of odors, we started with a normalized correlation coefficient computed between odor
abundances across different natural sources. Based on these distances, we performed
nonlinear dimensionality reduction using different curved and flat metrics[64]. The method
automatically adjusts the curvature of the embedding space and selects the best fitting
dimension based on the Bayesian Information Criterion[44]. We found that the natural odor
abundance data was best described by a three-dimensional hyperbolic space with a negative
curvature of -5.12. This observation mirrors previous results using a more limited dataset of
natural fruit or flower odor sources in which hyperbolic geometry also provided a significantly
better fit of the natural source space than the standard Euclidean geometry[12,45] . The intuitive
explanation for these results is that hyperbolic spaces provide good descriptions of the structure
of natural odor spaces because they arise as continuous approximations to tree-like hierarchical
networks. In the case of natural source data, the hierarchical relationships are hypothesized to
reflect dependencies produced by biochemical and metabolic pathways acting in plants and
other food sources, including in associated microbes like bacteria and fungi. In contrast,
nonlinear dimensionality reduction of odor distances computed from chemical descriptors
resulted in a hyperbolic embedding with higher dimension (dim=6) and much reduced curvature
(0.05) that did not provide a better fit of the dataset than a Euclidean model. Since odor
distances in the hyperbolic embedding better captured the correlational structure of abundances
of odors across natural sources compared to equivalent distances computed from a Euclidean
embedding, we used hyperbolic distances as a metric of odor relationships in natural source
space.
315
For each pair of odors in the dataset, we compared their representational distance in OR or KC
coding space with their relationship in chemical descriptor space or natural source space
(Figure 4B). We found that pairwise odor distances in OR coding space were better correlated
with their distances in chemical descriptor space (rho=0.27, p=2.4e-53) as compared with their
distances in natural source space (rho =0.087, p =4.8e-6) (Figure 4C-D). The converse was true
for odor distances in KC coding space: pairwise distances in KC coding space were better
correlated with their distances in natural source space (rho =0.23, p =7.1e-4) compared to their
distances in chemical descriptor space (rho =0.07, p=0.06) (Figure 4C-D). The correlation to
distances in KC coding space was observed only for natural source odor relationships quantified

by the hyperbolic embedding of natural source space, but not by Euclidean distance (Figure
4E), suggesting that capturing hierarchical relationships in the dataset is important and that the
low-dimensional embedding helps to de-noise the data. We also evaluated the relationship of
odor properties to odor distances in the modeled KC coding space under assumptions of
unstructured PN-KC connectivity. In contrast to the observed KC distances between odors,
modeled KC distances were better correlated with odor distances in chemical descriptor space
(*rho*=0.23, *p*=1.4e-38) compared to hyperbolic odor distances in natural source space
(*rho*=0.061, *p*=8.4e-3) (Figure 4E). This result is expected since, under assumptions of
unstructured connectivity in the olfactory circuit, the relationship of odors in KC coding space
stems from their relationships in OR input space. Similar results were observed for odor
relationships computed from KC responses predicted using the hemibrain circuit model,
indicating that the observed degree of glomerular input bias to KCs in the hemibrain does not
explain the reformatting of odor relationships in the MB. Overall, these data indicate that the
realignment of odor representation with natural source relationships that we observed in KCs
cannot be derived simply by random resampling of OR responses.
340
To determine if the relationships between odor representations and odor properties were driven
by only a small number of odor pairs in the dataset, we recomputed the correlations between
OR or KC representational distance and chemical descriptor or natural source distance using
100 subsamples comprising 75% of the odor pairs, dropping out a random 25% of the odor
pairs in each resampling. This analysis confirmed that odor distances in OR space were more
correlated with distances in chemical descriptor space as compared to in natural source space,
and vice versa for odor distances in KC space. These findings are consistent with a
reorganization of the fly olfactory code between the periphery and the MB from encoding the
chemical or structural relationships between odors to reflecting the odor relationships in
complex mixtures arising from natural sources.
351
Odor relationships are restructured across successive stages of olfactory processing
To better understand how representations of odor are reformatted from the periphery to the MB,
we used functional imaging to measure odor-evoked patterns of activity at each successive
stage of processing in the olfactory circuit. While delivering odors to the antennae of the fly, we
volumetrically imaged from all ORN axon terminals (labeled by the *pebbled-Gal4* driver) or from
~70% of PN dendrites (labeled by the *GH146-Gal4* driver) in the antennal lobe[46], where the
neurites from these cell populations are stereotypically organized into glomerular compartments
with characteristic size and position (Figure 5A-B). Population imaging from the terminals of all
ORN classes allowed odor relationships to be assessed in the complete OR input space to the
olfactory system, whereas the Hallem dataset measures the tuning of only approximately half of
all fly ORs (Supplemental Figure S4B). ROIs corresponding to individual glomeruli were
manually segmented from movies of odor responses in ORN axon or PN dendrites and their
odor response profiles extracted. To evaluate patterns of odor-evoked PN output, we
volumetrically imaged from the axonal boutons of ~70% of PNs (labeled by the *GH146-Gal4*
driver) in the calyx of the MB, where they synapse onto the sites of KC input (claws) (Figure 5A-
B). ROIs corresponding to PN boutons were automatically segmented using Suite2P; visual

Figure 5: Reformatting of odor representations across four stages of olfactory processing.
A) Anatomical schematic of fly olfactory circuit. ORN axon terminals are organized in a stereotyped glomerular map in the AL. Each uniglomerular PN has dendrites in a single AL glomerulus and projects to the calyx of the MB, where its axonal arbor terminates in multiple boutons that each synapse with KC input claws.
B) Representative baseline fluorescence and peak ΔF/F heatmaps of odor-evoked calcium signals in ORN axon terminals, PN dendrites, PN boutons, and KC somata to 1-pentanol, 1-hexanol, and ethyl propionate. KC data are reproduced from Figure 1. Scale bar, 10 μm. Genotypes: ORN axons are imaged in *pebbled-Gal4, UAS-GCaMP6f*;

+; *GH146-QF, QUAS-tdTomato*. PN dendrites and PN boutons are imaged in *GH146-Gal4/20xUAS-IVS-jGCaMP8m* (II).

C) Matrix of mean pairwise correlation distances (averaged across flies) of trial-averaged odor representations in ORN axons ($n = 4$ flies), PN dendrites ($n = 6$ flies), PN boutons ($n = 6$ flies), and KC somata ($n = 4$ flies) to a panel of 17 odors. Odors are displayed in the same order in each matrix according to the clustering of their representations in KC representational space.

D) Spearman's correlations measuring the similarity of rank ordering of odor response strengths at each stage of processing.

E) Hierarchical clustering of distance matrices in **C** showing best odor groupings in each representational space.

F) Pairwise adjusted mutual information (MI) score evaluating the similarity of clustering of odor representations in each representational space. The adjusted MI has a value of 1 when two clusterings are identical, and a value of 0 when the MI between two clustering is the value expected due to chance.

368   inspection confirmed that > 95% of ROIs reliably corresponded to single PN boutons. Odor
369   relationships were invariant across the brains of different individuals at each of these earlier
370   stages of olfactory processing (Figure S4A), consistent with the stereotyped connectivity of the
371   antennal lobe.
372
Comparing the pairwise correlation distances between odor representations in each layer of the
olfactory circuit, odor representations were progressively decorrelated at each feedforward
stage of processing (Figure 5C), with the mean and median odor-odor correlation becoming
systematically lower in each successive layer (Figure S4D). The largest decorrelation occurred
between PN boutons and KC somata, consistent with the divergent expansion of the circuit at
this synaptic step. However, the rank order of odor-odor distances was not maintained across
layers (ORN terminals versus KC somata, Spearman's rho=0.48), indicating that odor
relationships are transformed beyond a simple linear scaling between layers. For comparison, in
a model that assumes unstructured PN input to KCs and uniform inhibition in the AL and MB
("uniform"), the Spearman's correlation between odor relationships in the input OR layer and in
modeled KCs is rho=0.83. Furthermore, whereas the relative levels of odor-evoked activity in
PNs dendrites or boutons was well-predicted by levels of overall ORN input to the circuit, odor-
evoked KC response rates (and mean activity) were comparatively less well-predicted (Figure
5D, S4Ci-iv, and data not shown). These observations all point towards a non-linear
reformatting of odors in the MB representational space that alters the rank ordering of odor
relationships compared to earlier stages of coding.
389
We examined odor relationships at each step of processing using hierarchical clustering of odor
representations in each neural population, in order to better understand the origins of odor
relationships in MB coding space. For instance, one of the strongest odor clutsers at the level of
input representations – 2-heptanone, isoamyl acetate, and pentyl acetate – was maintained in
each subsequent stage of processing through to the KCs (Figure 5C, E). These odors share a
high degree of chemical similarity as well as co-occurrence in natural sources, which may
explain their robust grouping. However, other odor representations that were similarly well-
correlated and clustered in OR space (e.g., 1-pentanol and 2-butanone) were selectively
decorrelated in KC coding space, with each odor becoming reassigned to distinct, non-
overlapping groups of odors in KC coding space (Figure 5C, E). Additionally, the
representations for some odor pairs were observed to become reliably *more* similar at later
stages of olfactory processing. For instance, the pairwise correlation distances between 1-

pentanol and 1-hexanol was actually shorter in KC coding space as compared to in ORN axon,
PN dendrite, or PN bouton coding space.
404
Typically, these transformations in the representational geometry among odors occur
progressively at each successive stage of processing, with the largest change occurring
between PN boutons and KC somata (Figure 5C). However, we observed significant
reorganization of odor representations even between the inputs and outputs of the same cell
population, PNs; odor relationships encoded in PNs boutons better reflect odor relationships in
KC coding space compared to PN dendrites (Figure 5C, E). The nonlinear reformatting of odor
relationships across successive stages of olfactory coding indicates that prevailing models of
olfactory networks that assume global or uniform processing across neurons are incomplete,
and additional sources of structure exist in the fly olfactory circuit that mediate non-uniform
interactions between olfactory coding units (e.g., neurons or glomeruli).
415
416
417 **DISCUSSION**
418
We show that representations of odor are structured and invariant across MBs in individual flies,
and that the structure of MB odor coding space is only partially predicted by models that
assume random sampling of olfactory glomerular inputs by KCs. The latter is true even after
adjusting the MB model to account for the over- or under-representation of PN boutons from
specific glomeruli ("hemidraw" model) or the amount of over- and under-convergence of PN
inputs from specific sets of glomeruli onto KCs ("hemibrain" model). Thus, certain assumptions
of olfactory system architecture – for instance, of uniform strengths of unitary feedforward
synapses, uniform spiking thresholds, or nonselective, global inhibition in the AL or MB – are
likely oversimplifications, motivating a search for source(s) of structured interactions between
glomeruli or neurons in the olfactory circuit.
429
We demonstrate a significant transformation of the fly olfactory code between the periphery and
the MB, in which the encoding of odors by ORs better reflects the chemical relationships
between odors and the encoding of odors in the MB better reflects the distributions of odors
across behaviorally relevant natural sources. That representations at the olfactory periphery
better capture odor relationships in terms of their chemical or molecular properties is perhaps
unsurprising, since odor-OR interactions are governed by the structural features of the odor.
The reorganization of MB odor representations to better correlate with the relationships of these
odors in natural source space may reflect the progressive transformation of odor
representations in the olfactory network to encode latent variables that relate more directly to
behavioral value or perception.
440
Odor distances in the hyperbolic embedding of natural odor mixtures reflect correlations
between volatiles in the headspace profiles across natural sources that arise from conserved
metabolic pathways[45]. The transformation of odor representations in the MB is predicted to
facilitate the perceptual compression of odors that have shorter metabolic tree distances in the
hierarchical organization of natural odor mixtures relative to their chemical similarity. Consistent

with this idea, a recent preprint reports that distances between odors in a neural network
embedding trained on human olfactory perceptual labels are correlated with the metabolic
distance between odors in experimentally elucidated biochemical pathways[47].
449
The specific structure of odor relationships was largely invariant across individual brains at each
stage of olfactory processing (Figure 2), even in the MB where PN boutons connect
probabilistically with KCs. An important open question is the extent to which the invariance of
MB representational space may arise from genetically specified developmental processes
(sculpted by evolution) or from activity-dependent processes that reflect shared olfactory
experience[2]. Unlike their mammalian analogues in piriform cortex, KCs are not strongly
connected through recurrent excitation, although the degree to which PN-KC synapses or APL-
KC synapses may be regulated by experience is not well understood. We note that the structure
of PN overconvergence onto KCs bears significant similarity to the structure of PN input to third-
order olfactory neurons in the lateral horn (LH) (Figure S3E). LH neurons are stereotyped in
their anatomical connectivity and odor tuning and are believed to mediate innate olfactory
behaviors. The similarity in PN input structure between the MB and LH points to the likely
ethological significance of these over-represented glomerular combinations. It also raises the
possibility that the LH may have additional roles in shaping odor representations in the MB. For
instance, neurons downstream of the LH send centrifugal input to the MB calyx[48] which could
contribute to the remapping of odor representations in PN boutons. Other possible sources of
structure in the circuit are selective inhibition from the APL or MB-C1 neurons in the calyx, or
non-uniform PN-KC synaptic weights. Understanding the specific circuit mechanisms that shape
the reorganization of odor representations in the MB will be important for understanding how
odor relationships in natural sources become reflected in the olfactory code.
470
Theoretical studies of divergent expansive cerebellum-like circuits such as the MB emphasize
the computational benefits of random networks for maximizing coding capacity and promoting
the separability and discriminability of representations. However, structured networks can
correlate odor representations and promote generalization of odors along important directions of
odor space, for instance, related to the relationships of odorant molecules in behaviorally salient
natural odor sources. Our results suggest a revision of classic formulations of cerebellar-like
network architectures is warranted at least in some systems: the MB may trade-off the capacity
to maximally decorrelate activity patterns in parts of representational space for an increase in
coding capacity and robustness in a part of representational space of particular ethological
significance[49]. The recoding of odors in the MB to reflect relationships in natural sources would
predict greater generalization to odors with similar distributions in the environment, facilitating
the decoding of natural source identity in noisy or ambiguous situations.
483
484 **METHODS**
485
Experimental model
Drosophila melanogaster were raised on a 12:12 light:dark cycle at 25°C and 70% relative
humidity on cornmeal/molasses food containing: water (17.8 l), agar (136 g), cornmeal (1335.4
g), yeast (540 g), sucrose (320 g), molasses (1.64 l), CaCl2 (12.5 g), sodium tartrate (150 g),

tegosept (18.45 g), 95% ethanol (153.3 ml) and propionic acid (91.5 ml). All experiments were
performed in female flies aged 3-10 days post-eclosion. Unless otherwise noted, the
transgenes in this study were acquired from the Bloomington Drosophila Stock Center (BDSC)
and have been previously characterized as follows: *pebbled-Gal4* (X) directs expression in all
ORNs[50] (RRID:BDSC_80570); *GH146-Gal4* (II) directs expression in ~70% of PNs[51]
(RRID:BDSC_30026); *GH146-QF*, *QUAS-tdTomato* (III) expresses the red-fluorescent protein
tdTomato in ~70% of PNs[52] (RRID:BDSC_30037); *UAS-IVS-jGCaMP8m* (II) expresses the
calcium indicator jGCaMP8m[53] in a Gal4-dependent manner (RRID:BDSC_92591); *OK107-Gal4*
(IV) directs expression in all KCs[33] (RRID:BDSC_854); *UAS-OpGCaMP6f* (X) was from B. D.
Pfeiffer and D. J. Anderson (Caltech, Pasadena, CA) and expresses the calcium indicator
codon-optimized GCaMP6f[54] in a Gal4-dependent manner; and *UAS-nls-OpGCaMP6s-p10 (III)*
was from H. Chiu and D. J. Anderson (Caltech, Pasadena, CA) and expresses nuclear-
localized, codon-optimized OpGCaMP6s[35] in a Gal4-dependent manner.
503
Odor stimuli
Odors were delivered essentially as previously described[55]. A custom-built multi-channel
olfactometer delivered a constant 2 L/min stream of charcoal-filtered air. A 3-way solenoid
valve directed 200 mL/min of this flow either through a 20-ml glass vial containing 2-ml of odor
solution (valve open) or an equivalent vial containing 2-ml of the solvent. Air flow was
controlled using mass flow controllers (MC series, Alicat Scientific, Tucson, AZ). The 200
ml/min control or odor streams were carried by tubing of matched lengths and rejoined the
carrier stream at the same point along the carrier tube, approximately 10 cm from the fly. The
terminal end of the carrier tube had an inner diameter ~8mm and was ~1 cm away from the fly.
Odor concentrations refer to the v/v dilution factor of the odor solution in the vial. The
concentration of odor in the headspace is further diluted 10-fold in air prior to reaching the fly.
Unless otherwise indicated, all odors in this study were presented from a 10^{-3} (0.1%) dilution in
paraffin oil (J.T. Baker, VWR #JTS894), with the exception of movies collected from one fly, in
which odors were diluted to 10^{-2} (1%).
518
Two-photon calcium imaging
Volumetric, in vivo functional calcium imaging was performed essentially as previously
described[46,55]. After a brief period of cold anesthesia (<20 s), the fly was head-fixed, and the
cuticle, fat, and air sacs were removed to expose the brain region of interest. Two-photon
GCaMP6f or GCaMP8m fluorescence was excited with 925 nm light from a Mai Tai DeepSee
laser (Spectra-Physics, Santa Clara, CA). Images were acquired with an Olympus 20X/1.0
numerical aperture objective (XLUMPLFLN20XW) driven by a piezo motor that enabled fast z-
scanning, and the collection filter was centered at 525 nm with a 50 nm bandwidth.
Experiments were conducted at room temperature (~22°C). The brain was constantly perfused
by gravity flow with saline containing (in mM): 103 NaCl, 3 KCl, 5 N-Tris(hydroxymethyl)methyl-
2-aminoethane-sulfonic acid, 8 trehalose, 10 glucose, 26 NaHCO3, 1 NaH2PO4, 1.5 CaCl2,
and 4 MgCl2 (pH 7.3, osmolarity adjusted to 270– 275 mOsm). The saline was bubbled with
95% O_2/5% CO_2 and circulated in the bath at ~2-3 ml min^{-1}.
532

AL imaging. Flies were head-fixed with the dorsal surface of the head approximately parallel to
the imaging plane; the dorsal cuticle was removed, and the antennal lobes were exposed. The
antennae were snugly secured below the imaging chamber, keeping them dry and responsive
to odors. ORN axon terminals and PN dendrites were imaged with a galvo-galvo scanning
system (Thorlabs Bergamo). Flies were imaged from the dorsal side, with 5 planes spaced 12
μm apart. The depth of the first imaging plane was chosen to maximize the number of
glomeruli visible across the 5 planes. Images were acquired at a resolution of 192x192 pixels,
with typical fields-of-view of ~90-100μm^2 and volumetric sampling rates of ~1 Hz through the
antennal lobe. For ORN axon and PN dendrite experiments, odors were presented in a
pseudorandom order, with the three trials of each odor being presented contiguously. Trials
consisted of a 7 s baseline recording, the odor pulse (2s for ORN experiments and 3s for PN
experiments), and a 20 s post-stimulus period.
545
MB imaging. Flies were head fixed with the head tilted acutely downward, rotated ~70° from
its normal resting position. The posterior plate of the head was approximately parallel to the
imaging plane, and the antennae were dry underneath the imaging platform. The entire
perimeter of the head capsule was stabilized to the imaging platform with glue, and the
proboscis and legs were immobilized to minimize motion. KC somata and PN boutons were
imaged with a galvo/resonance scanning system (Thorlabs Bergamo), with the exception of
one fly in which KC somata were imaged on a galvo-galvo scanning system at a volumetric
sampling rate of 0.55 Hz. For PN bouton experiments, odors were presented in a
pseudorandom order, with the three trials of each odor being presented contiguously. Trials
consisted of a 7 s baseline recording, a 3s odor pulse, and a 20 s post-stimulus period, with
stimulus onset occurring every 30s. For KC somata experiments, stimulus order was
pseudorandomized such that all odors were presented before any were repeated; thus,
repetitions of a given odor usually did not occur on consecutive trials. If paraffin was included
as an odor stimulus, it was always presented first in each repetition block. Trials consisted of a
13 s baseline recording, a 2 s odor pulse, and a 45 s post-stimulus period, with stimulus onset
occurring every 60s.
For resonance imaging of KC somata, movies were collected at a frame rate of ~60 Hz
at 256 x 256 pixels in fast-z mode. The field-of-view was ~150 μm x 150 μm giving a pixel size
of ~0.6 μm/pixel. The bulk of the KC somata cluster was captured in 16 planes (and 4 flyback
frames) collected 2 μm apart, spanning a ~30 μm depth through the MB. This resulted in a
volumetric sampling rate of ~3 Hz. For a subset of experiments, KCs were sampled using fewer
imaging planes (eleven) with a z-step size of 3 μm. Since odor response rates and odor-odor
relationships were similar with these imaging parameters, KC somata datasets were combined.
For imaging of PN boutons in the MB calyx, movies were collected at a frame rate of ~60
Hz at a resolution of 256 x 256 pixels in fast-z mode. The field of view was ~75 μm x ~75
μm yielding a pixel size of ~0.3 μm/pixel. The full population of labeled PN boutons in the calyx
was captured using 8 planes (and 2 flyback frames) spaced ~3 μm apart, spanning a ~21 μm
depth in z, giving a volumetric sampling rate of ~6 Hz.
574
Image analysis
AL imaging

Image analysis was performed using custom Python scripts
(https://github.com/ejhonglab/al_analysis). Motion correction was performed separately in each
plane using the registration module in Suite2p. ROIs corresponding to individual glomeruli were
manually defined in each imaging plane in Fiji[56], using a combination of the resting
fluorescence and per-trial responses to visualize the glomerular boundaries. Responses (ΔF/F)
were calculated using a 6 s baseline (F) that ended 1 s prior to nominal odor onset in each trial.
The odor response was quantified as the mean ΔF/F during the 2 s after odor onset. Odor
tuning of identified glomeruli, assigned using a combination of anatomical position, size,
shape, and responses to a diagnostic panel of 10 narrowly activating odor stimuli, matched
previous descriptions[32,57,58].
587
MB imaging
Motion correction. For PN bouton and KC somata imaging, an initial round of 3D motion
correction was performed with the 'NormCorre' algorithm, using either the Matlab
implementation[59] (https://github.com/flatironinstitute/NoRMCorre) or Python implementation
from the calcium imaging analysis library 'CaImAn'[60]
(https://github.com/flatironinstitute/CaImAn). As necessary, planes were dropped following
motion correction due to field-of-view drift; typically the most superficial or deepest planes
were dropped if they were not consistently captured through the entire recording.
596
Source extraction. For movies of PN boutons and KC somata collected in the MB, source
extraction was carried out using either Suite2p[36] or CaImAn[60]. For all PN bouton, and for KC
somata data collected at a z-step of 2 μm, Suite2p was used to perform source extraction on a
plane-by-plane basis. ROIs corresponding to PN boutons and odor-evoked signals were
extracted in Suite2p's *functional* mode, using the activity-based algorithm. KC ROIs and
signals were extracted in Suite2p's *anatomical* mode, using the 'cellpose' model to perform
anatomical segmentation on the mean (time-averaged) image. Extracted fluorescence traces
were neuropil-corrected ($F_{corrected} = F - 0.7 *F_{neuropil}$) and normalized by z-scoring over time. The
'rastermap' function[61] was used to visualize $F_{corrected}$ for all extracted components, with cells
sorted to cluster those with similar patterns of activity. Stimulus-responsive clusters were
manually selected. For PN bouton movies, this step was used to filter out spurious ROIs – only
components belonging to stimulus-responsive clusters were included in subsequent analyses.
For KC movies, the relatively high level of baseline fluorescence in combination with the use of
a nuclear-targeted calcium indicator resulted in very high-quality cellular segmentation, with
very few (if any) spurious ROIs. Figure 1G shows the responses of all extracted cells from one
representative experiment, with cells (rows) corresponding to stimulus-responsive clusters
displayed at the top of the matrix, and non-responder cell clusters shown at the bottom.
CaImAn-MATLAB was used to perform 3D source extraction on KC somata data
collected at a z-step size of 3 μm. For each KC, a single ROI was extracted, which was roughly
spheroid and spanned multiple planes. The fluorescence baseline F_0 was computed at each
timepoint by taking a fixed percentile (ranging from 20-40%) of a rolling 30 s time window.
Extracted raw calcium signals were detrended and normalized to baseline fluorescence F0 to
compute ΔF/F responses. All components extracted by CaImAn were included in subsequent
analyses. Between ~1500-1900 KCs and ~500-800 PN bouton ROIs were extracted per movie.

Quantification of odor response. For Suite2p extracted signals, ROI response strength in a
single trial was computed by averaging the signal over an expected response peak time
window (0.25-2s post-stimulus onset for PN boutons, 2-8s post-stimulus for KCs), and
subtracting the mean pre-stimulus baseline (initiating 5s pre-stimulus for PN boutons, and 10s
pre-stimulus onset for KC soma). For CalmAn-extracted KC signals, the response strength for
each trial was calculated by subtracting the median 10s pre-stimulus baseline from the mean
15s post-stimulus ΔF/F signal. For all datasets, mean odor response strength was computed
by averaging across all trials in which the odor was presented.

KC response rate
KC response rates (Supplementary Figure S1) were computed from datasets in which source
extraction was performed using Suite2p, applying the 'cellpose' model to perform anatomical
segmentation. All extracted components were used in the analysis of response strength and
response breadth, including 'silent' cells that did not respond to any stimulus in the odor panel.
A cell was considered a 'responder' if its trial-averaged response to a given odor exceeded a
fixed threshold. The response threshold was determined separately for each dataset using
ethyl propionate (a consistently strongly activating odor) as a reference, in order to adjust for
small differences in responsiveness of different experimental preparations. The KC response
rate to ethyl propionate was computed over a range of thresholds, with a step size of 0.05
between threshold values. Threshold values resulting in a 12-15% ethyl propionate response
rate were selected. The median of these values was chosen as the final response threshold
and was used to compute response rates for the other odors in the dataset.

Analysis of the representational space of odors
Distances between population representations of odor. The correlation distance between the
population representation of two odors was computed as 1-*r*, where *r* is Pearson's correlation
between the population response vectors of the two odors. The population response vector
had length *l*, where *l* was the number of glomeruli, boutons, or cells in each experiment. Each
element of this vector was the trial-averaged response of each glomerulus/bouton/cell to the
stimulus. Representational dissimilarity matrices (RDM) show the pairwise correlation distances
for every pair of odors imaged and aligned in the same experiment (same MB) and were
computed using only the set of cells, glomeruli, or boutons in each experiment that responded
to one or more odor stimuli. For each stage of processing, we computed a mean RDM by
averaging the odor x odor RDMs across individual flies.

Comparison of representational space across individual MBs. For each MB, we generated an
odor-odor distance vector of the correlation distances for every odor pair measured in
common across the flies being compared. To evaluate the similarity of MB odor
representational space across flies, we computed the Spearman's rank correlation between
the odor-odor distance vectors for every pairwise combination of MBs. As a control, the cell x
odor response matrices for each MB were shuffled, and the same series of calculations
applied. For each iteration of the shuffle, odor responses were randomized by permuting the
columns of the cell x odor response matrix for each MB. The Spearman's correlation computed

from 10,000 iterations of this shuffle procedure was used to construct a distribution of
Spearman's correlations that would arise from chance for each pair of MBs, to be compared
against the observed correlation.

Cell clustering. KCs were grouped by spectral clustering carried out on KC odor tuning profiles
using 'sklearn.cluster.SpectralClustering'[62]. Trial-averaged odor response vectors for all KCs
that respond to one or more odors were pooled together from individual flies to create a grand
cell x odor response matrix. The number of rows in this matrix equaled the sum of the number
of responsive KCs across all flies and the number of columns equaled the number of odors
that were sampled in common across all flies. A KC x KC affinity matrix was computed by
taking the radial-basis transform of the KC x KC correlation distance matrix. Odor response
profiles for each cluster were calculated by averaging the responses of all cells assigned to
that cluster from each fly.

Modeling KC responses
We adapted a dynamic spiking model of the *Drosophila* olfactory network[30]
(https://github.com/annkennedy/mushroomBody) that implements the functional and
anatomical organization of the circuit. ORN input to the model was derived from a published
experimental dataset of the firing rate responses of 23 fly ORs to a panel of 109 odors[32]. The
model implements lateral inhibition in the AL with a divisive inhibition term that normalizes PN
firing rates. The model captures the response dynamics of PN and KC firing rates to an odor
pulse, where ORN input is estimated by simply convolving a step to the steady-state firing rate
with the cell's synaptic membrane filter. For our analyses in this study, we focused on the
mean firing rate of each cell (PN or KC) averaged over the odor pulse.
The model was implemented under several different assumptions of PN-KC
connectivity. In the 'uniform' model, each of 2000 KCs had a mean of six input glomeruli, with
each input independently assigned to a glomerulus at random. Each glomerulus had an equal
likelihood of being selected at every input. In the 'hemidraw' model, each KC in the model
(1748 cells) corresponded to a KC in the hemibrain connectome dataset[63] and was assigned its
observed number of claws (between 1 and 12, mean of 5.36 claws). Each claw was then
independently assigned to a glomerulus according to the frequency of PN boutons for that
glomerulus in the hemibrain MB; thus, glomeruli varied in their likelihood of being drawn as an
input to each claw. In the 'hemibrain' model, each KC in the model was directly assigned the
set of glomerular inputs of its corresponding cell in the hemibrain connectome. However, since
glomeruli in the model were limited to the 23 ORs available in the Hallem dataset, modeled
'hemibrain' KCs had only an average of ~4 inputs. To evaluate the impact of a reduced number
of inputs, versions of the 'uniform' and 'hemidraw' models were run with the distribution of the
number of claws per KC centered on 4. These models predicted very similar odor-odor
relationships to the earlier versions (data not shown).
The response rate of KCs to odor depends on both KC spiking threshold and the
strength of feedback inhibition to KCs from the anterior paired lateral (APL) GABAergic neuron.
In all versions of the model, KC spiking thresholds were assumed to be uniform across KCs
and were set to achieve an average KC response rate of 20% across odors. Global inhibition
by the APL was modeled as divisive inhibition at KC presynaptic terminals, and APL-KC

weights were adjusted to halve the mean response rate across odors to 10%. This procedure
was motivated by experimental observations that silencing APL output roughly doubles KC
response rates.[22] The APL was assumed to uniformly receive equal input from, and send equal
output to, all KCs. For modeling of KC responses to CO_2 (a stimulus not in the Hallem dataset),
an additional glomerulus (corresponding to glomerulus V) and odor (CO_2) was added into the
OR input matrix. To estimate an upper bound for KC response rates to CO_2, the firing rate of
glomerulus V to CO_2 was set to the maximum ORN firing rate (300 Hz) and set to zero for all
other odors[57].
717
Odor properties
Chemical descriptors were computed for each odor using the software Mordred[41]. Clustering
over features identified a reduced set of 570 descriptors that captured odor-odor relationships in
the full set of ~1800 descriptors; our analysis used this reduced set of descriptors
(Supplemental Table S3). Odor relationships in natural odor space were estimated from a large
database of headspace volatile profiles of natural food sources compiled from published
datasets from the food science literature (Volatile Compounds in Food, VCF16.9 database,
BeWiDo BV). The database contains 5564 observations from thousands of references. We
focused on 2190 observations for which volatile profiles were quantitatively described in
standardized units that could be compared between sources. For our analysis, we filtered
compounds to isolate those present in ten or more observations (775 odors) and used the log
scale of odor concentration.
730
Multidimensional scaling of odors in natural source space
We used hyperbolic non-metric multidimensional scaling (H-MDS)[64] to embed odors into a low-
dimensional hyperbolic space based on correlation distance, which reconstructs original
distances monotonically (with preserved rank-ordering). The minimum Bayesian information
criteria (BIC) determines the best dimension of the embedding. A hyperbolic metric was then
used to measure distances between odors within the low-dimensional embedding space. To
check that the results generalize across different subsets of natural odor sources, we repeated
the analysis for a separate, albeit smaller secondary dataset that was compiled from another set
of natural source literature references (see Supplemental Table S4). We find that odor pair
distances computed in the VCF and in the secondary natural source dataset were positively
correlated (rho=0.85, p=1.5e-8). The correlation between datasets increased with increasing the
minimum number of odor sources for which a monomolecular odorant was required to be
present in calculating distances between odorants. This result indicates increased stability of
distances between more common odorants, that are ubiquitously present in natural
environment. Furthermore, in both datasets, the hyperbolic space provided the best low-
dimensional description. The curvature and best fitting dimensionality (dim=3) were similar, with
curvature = -5.1 for the VCF dataset and curvature = -4.1 for the secondary natural source
dataset.
749
Hierarchical clustering of odor representations
Representational dissimilarity matrices (RDM) show the pairwise correlation distances between
every pairwise combination of odors imaged and aligned in the same experiment. For each

stage of processing, a mean RDM was computed by averaging the odor x odor RDMs across
individual flies. Dendrograms describing the relationships among odors at each stage of
processing were generated from the mean odor x odor RDM for each stage. Hierarchical
clustering of odors was implemented with 'sklearn.cluster.AgglomerativeClustering', using
Pearson's correlation as the distance metric and average linkage criterion (minimizes the
average of the distances between all observations of pairs of clusters). These odor x odor RDM
matrices were treated as feature matrices with dimensions (samples, features) rather than
distance matrices – each odor was a different sample/row, and that odor's features were its
distance to the odors.
762
Data inclusion criteria.
The number of flies (observations, *n*) in which each odor was measured is in Supplemental
Table S1. The number of observations of each odor-odor distance (odor pair measured in the
same MB) is in Supplemental Table S2. All flies analyzed in this study satisfied the following
criteria. First, any field-of-view drift and warping of structure could be fully corrected using
posthoc image registration, as evaluated by the 'crispness' of time-averaged movies (individual
nuclei distinct and separated). Second, stimulus-evoked responses were reliably observed over
the course of the entire recording in a 'bulk' fluorescence signal extracted in each frame from a
global ROI circumscribing the entire imaged structure. Third, trial-trial correlation distances for
repeated presentations of the same odor stimulus were clearly more similar to one another
than for presentations of different odor stimuli. Sample sizes were not predetermined using a
power analysis. We used sample sizes comparable to those used in similar types of studies[2,3].
775
776

ACKNOWLEDGEMENTS
We thank D. Anderson for sharing the *UAS-OpGCam6f* and *nls-OpGCaMP6s* fly lines. We
thank A. Kennedy for modeling code and assistance with adapting the olfactory network model
in this study. We thank V. Hauser for contributions to compiling natural odor source datasets.
We thank P. Kandimalla for advising on connectomic analyses. We thank members of the Hong
lab for careful readings of the manuscript. This work was supported by NIH grant
R01MH117825 to E.J.H., and by the NSF/CIHR/DFG/FRQ/UKRI-MRC Next Generation
Networks for Neuroscience Program (Award #2014217) to T.O.S. and E.J.H.

785
AUTHOR CONTRIBUTIONS
J.Y.Y., T. F.O., T.O.S., and E.J.H. conceived the project, analyzed data, and wrote the
manuscript. J.Y.Y. and K.V.D. performed imaging experiments in PN axons and KC somata.
T.F.O. performed imaging experiments in ORN axons and PN dendrites and supervised
curation of the natural odor source database. J.Y.Y., T.F.O., M.S.B., and W.M.H. analyzed data
and generated figures. E.J.H. and T.O.S. supervised the project and acquired funding.
792
793
794 **DECLARATION OF INTERESTS**
795
796 The authors declare no competing interests.

REFERENCES

1. Schaffer, E.S., Stettler, D.D., Kato, D., Choi, G.B., Axel, R., and Abbott, L.F. (2018). Odor Perception on the Two Sides of the Brain: Consistency Despite Randomness. Neuron *98*, 736-742.e3. 10.1016/j.neuron.2018.04.004.

2. Pashkovski, S.L., Iurilli, G., Brann, D., Chicharro, D., Drummey, K., Franks, K.M., Panzeri, S., and Datta, S.R. (2020). Structure and flexibility in cortical representations of odour space. Nature *583*, 253–258.

3. Endo, K., Tsuchimoto, Y., and Kazama, H. (2020). Synthesis of conserved odor object representations in a random, divergent-convergent network. Neuron *108*, 367–381.

4. Secundo, L., Snitz, K., and Sobel, N. (2014). The perceptual logic of smell. Current opinion in neurobiology *25*, 107–115.

5. Buck, L., and Axel, R. (1991). A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. Cell *65*, 175–187.

6. Khan, R.M., Luk, C.-H., Flinker, A., Aggarwal, A., Lapid, H., Haddad, R., and Sobel, N. (2007). Predicting odor pleasantness from odorant structure: pleasantness as a reflection of the physical world. Journal of Neuroscience *27*, 10015–10023.

7. Keller, A., Gerkin, R.C., Guan, Y., Dhurandhar, A., Turu, G., Szalai, B., Mainland, J.D., Ihara, Y., Yu, C.W., and Wolfinger, R. (2017). Predicting human olfactory perception from chemical features of odor molecules. Science *355*, 820–826.

8. Genva, M., Kenne Kemene, T., Deleu, M., Lins, L., and Fauconnier, M.-L. (2019). Is It Possible to Predict the Odor of a Molecule on the Basis of its Structure? International journal of molecular sciences *20*, 3018.

9. Sell, C.S. (2006). On the unpredictability of odor. Angewandte Chemie International Edition *45*, 6254–6261.

10. Dzialo, M.C., Park, R., Steensels, J., Lievens, B., and Verstrepen, K.J. (2017). Physiology, ecology and industrial applications of aroma formation in yeast. FEMS microbiology reviews *41*, S95–S128.

11. Junker, R.R., Kuppler, J., Amo, L., Blande, J.D., Borges, R.M., van Dam, N.M., Dicke, M., Dötterl, S., Ehlers, B.K., and Etl, F. (2018). Covariation and phenotypic integration in chemical communication displays: Biosynthetic constraints and eco-evolutionary implications. New Phytologist *220*, 739–749.

12. Ghaninia, M., Zhou, Y., Knauer, A.C., Schiestl, F.P., Sharpee, T.O., and Smith, B.H. (2022). Hyperbolic odorant mixtures as a basis for more efficient signaling between flowering plants and bees. PloS one *17*, e0270358.

13. Wright, G.A., and Schiestl, F.P. (2009). The evolution of floral scent: the influence of olfactory learning by insect pollinators on the honest signalling of floral rewards. Functional Ecology *23*, 841–851.

14. Wilson, R.I., and Mainen, Z.F. (2006). Early events in olfactory processing. Annual Review
of Neuroscience *29*, 163–201.

15. Vosshall, L.B., and Stocker, R.F. (2007). Molecular architecture of smell and taste in
Drosophila. Annu Rev Neurosci *30*, 505–533. 10.1146/annurev.neuro.30.051606.094306.

16. Modi, M.N., Shuai, Y., and Turner, G.C. (2020). The Drosophila mushroom body: from
architecture to algorithm in a learning circuit. Annual review of neuroscience *43*, 465–484.

17. Caron, S.J., Ruta, V., Abbott, L.F., and Axel, R. (2013). Random convergence of olfactory
inputs in the Drosophila mushroom body. Nature *497*, 113–117. 10.1038/nature12063.

18. Li, F., Lindsey, J.W., Marin, E.C., Otto, N., Dreher, M., Dempsey, G., Stark, I., Bates, A.S.,
Pleijzier, M.W., Schlegel, P., et al. (2020). The connectome of the adult Drosophila
mushroom body provides insights into function. eLife *9*, e62576. 10.7554/eLife.62576.

19. Zheng, Z., Li, F., Fisher, C., Ali, I.J., Sharifi, N., Calle-Schuler, S., Hsu, J., Masoodpanah,
N., Kmecova, L., and Kazimiers, T. (2022). Structured sampling of olfactory input by the fly
mushroom body. Current Biology *32*, 3334–3349.

20. Turner, G.C., Bazhenov, M., and Laurent, G. (2008). Olfactory representations by
Drosophila mushroom body neurons. Journal of neurophysiology *99*, 734–746.

21. Gruntman, E., and Turner, G.C. (2013). Integration of the olfactory code across dendritic
claws of single mushroom body neurons. Nature neuroscience *16*, 1821–1829.
10.1038/nn.3547.

22. Lin, A.C., Bygrave, A.M., De Calignon, A., Lee, T., and Miesenböck, G. (2014). Sparse,
decorrelated odor coding in the mushroom body enhances learned odor discrimination.
Nature neuroscience *17*, 559–568.

23. Honegger, K.S., Campbell, R.A., and Turner, G.C. (2011). Cellular-resolution population
imaging reveals robust sparse coding in the Drosophila mushroom body. The Journal of
neuroscience : the official journal of the Society for Neuroscience *31*, 11772–11785.
10.1523/JNEUROSCI.1099-11.2011.

24. Campbell, R.A., Honegger, K.S., Qin, H., Li, W., Demir, E., and Turner, G.C. (2013).
Imaging a population code for odor identity in the Drosophila mushroom body. The Journal
of neuroscience : the official journal of the Society for Neuroscience *33*, 10568–10581.
10.1523/JNEUROSCI.0682-12.2013.

25. Bhandawat, V., Olsen, S.R., Schlief, M.L., Gouwens, N.W., and Wilson, R.I. (2007). Sensory
processing in the Drosophila antennal lobe increases the reliability and separability of
ensemble odor representations. Nature Neuroscience *10*, 1474–1482.

26. Cayco-Gajic, N.A., and Silver, R.A. (2019). Re-evaluating circuit mechanisms underlying
pattern separation. Neuron *101*, 584–602.

27. Litwin-Kumar, A., Harris, K.D., Axel, R., Sompolinsky, H., and Abbott, L.F. (2017). Optimal
Degrees of Synaptic Connectivity. Neuron *93*, 1153-1164.e7.
10.1016/j.neuron.2017.01.030.

28. Babadi, B., and Sompolinsky, H. (2014). Sparseness and expansion in sensory
representations. Neuron *83*, 1213–1226.

29. Barak, O., Rigotti, M., and Fusi, S. (2013). The sparseness of mixed selectivity neurons
controls the generalization–discrimination trade-off. Journal of Neuroscience *33*, 3844–
3856.

30. Kennedy, A. (2019). Learning with naturalistic odor representations in a dynamic model of
the *Drosophila* olfactory system (Neuroscience) 10.1101/783191.

31. Owald, D., and Waddell, S. (2015). Olfactory learning skews mushroom body output
pathways to steer behavioral choice in Drosophila. Current opinion in neurobiology *35*, 178–
184.

32. Hallem, E.A., and Carlson, J.R. (2006). Coding of odors by a receptor repertoire. Cell *125*,
143–160.

33. Tanaka, N.K., Tanimoto, H., and Ito, K. (2008). Neuronal assemblies of the Drosophila
mushroom body. J Comp Neurol *508*, 711–755.

34. Eder, A., and Bading, H. (2007). Calcium signals can freely cross the nuclear envelope in
hippocampal neurons: somatic calcium increases generate nuclear calcium transients. BMC
neuroscience *8*, 1–11.

35. Jung, Y., Kennedy, A., Chiu, H., Mohammad, F., Claridge-Chang, A., and Anderson, D.J.
(2020). Neurons that function within an integrator to promote a persistent behavioral state in
Drosophila. Neuron *105*, 322–333.

36. Pachitariu, M., Stringer, C., Schröder, S., Dipoppa, M., Rossi, L.F., Carandini, M., and
Harris, K.D. (2016). Suite2p: beyond 10,000 neurons with standard two-photon microscopy.
BioRxiv, 061507.

37. Benton, R., Vannice, K.S., Gomez-Diaz, C., and Vosshall, L.B. (2009). Variant ionotropic
glutamate receptors as chemosensory receptors in Drosophila. Cell *136*, 149–162.
10.1016/j.cell.2008.12.001.

38. Murthy, M., Fiete, I., and Laurent, G. (2008). Testing odor response stereotypy in the
Drosophila mushroom body. Neuron *59*, 1009–1023.

39. Frechter, S., Bates, A.S., Tootoonian, S., Dolan, M.-J., Manton, J., Jamasb, A.R., Kohl, J.,
Bock, D., and Jefferis, G. (2019). Functional and anatomical specificity in a higher olfactory
centre. eLife *8*, e44590. 10.7554/eLife.44590.

40. Jefferis, G.S., Potter, C.J., Chan, A.M., Marin, E.C., Rohlfing, T., Maurer, C.R., Jr., and Luo,
L. (2007). Comprehensive maps of Drosophila higher olfactory centers: spatially segregated
fruit and pheromone representation. Cell *128*, 1187–1203.

41. Moriwaki, H., Tian, Y.-S., Kawashita, N., and Takagi, T. (2018). Mordred: a molecular
descriptor calculator. Journal of cheminformatics *10*, 1–14.

42. Markow, T.A. (2015). The secret lives of Drosophila flies. eLife *4*, e06793.
10.7554/eLife.06793.

43. Dunkel, A., Steinhaus, M., Kotthoff, M., Nowak, B., Krautwurst, D., Schieberle, P., and
Hofmann, T. (2014). Nature's chemical signatures in human olfaction: a foodborne
perspective for future biotechnology. Angewandte Chemie International Edition *53*, 7124–
7143.

44. Schwarz, G. (1978). Estimating the dimension of a model. The annals of statistics, 461–464.

45. Zhou, Y., Smith, B.H., and Sharpee, T.O. (2018). Hyperbolic geometry of the olfactory
space. Science advances *4*, eaaq1458.

46. Zocchi, D., Emily, S.Y., Hauser, V., O'Connell, T.F., and Hong, E.J. (2022). Parallel
encoding of CO_2 in attractive and aversive glomeruli by selective lateral signaling between
olfactory afferents. Current Biology.

47. Qian, W.W., Wei, J.N., Sanchez-Lengeling, B., Lee, B.K., Luo, Y., Vlot, M., Dechering, K.,
Peng, J., Gerkin, R.C., and Wiltschko, A. (2022). Metabolic activity organizes olfactory
representations. bioRxiv, 2022–07.

48. Dolan, M.-J., Frechter, S., Bates, A.S., Dan, C., Huoviala, P., Roberts, R.J., Schlegel, P.,
Dhawan, S., Tabano, R., and Dionne, H. (2019). Neurogenetic dissection of the Drosophila
lateral horn reveals major outputs, diverse behavioural functions, and interactions with the
mushroom body. Elife *8*, e43079.

49. Nguyen, T.M., Thomas, L.A., Rhoades, J.L., Ricchi, I., Yuan, X.C., Sheridan, A., Hildebrand,
D.G., Funke, J., Regehr, W.G., and Lee, W.-C.A. (2022). Structured cerebellar connectivity
supports resilient pattern separation. Nature, 1–7.

50. Sweeney, L.B., Couto, A., Chou, Y.H., Berdnik, D., Dickson, B.J., Luo, L., and Komiyama, T.
(2007). Temporal target restriction of olfactory receptor neurons by Semaphorin-1a/PlexinA-
mediated axon-axon interactions. Neuron *53*, 185–200. 10.1016/j.neuron.2006.12.022.

51. Stocker, R.F., Heimbeck, G., Gendre, N., and de Belle, J.S. (1997). Neuroblast ablation in
Drosophila P[GAL4] lines reveals origins of olfactory interneurons. Journal of Neurobiology
32, 443–456.

52. Potter, C.J., Tasic, B., Russler, E.V., Liang, L., and Luo, L. (2010). The Q system: a
repressible binary system for transgene expression, lineage tracing, and mosaic analysis.
Cell *141*, 536–548. 10.1016/j.cell.2010.02.025.

53. Zhang, Y., Rózsa, M., Bushey, D., Zheng, J., Reep, D., Broussard, G., Tsang, A., Tsegaye,
G., Patel, R., and Narayan, S. (2020). jGCaMP8 fast genetically encoded calcium indicators.
Janelia Research Campus *10*, 13148243.

54. Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter,
E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent
proteins for imaging neuronal activity. Nature *499*, 295–300. 10.1038/nature12354.

55. Hong, E.J., and Wilson, R.I. (2015). Simultaneous encoding of odors by channels with
diverse sensitivity to inhibition. Neuron *85*, 573–589. 10.1016/j.neuron.2014.12.040.

56. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., and Schmid, B. (2012). Fiji: an open-source platform
for biological-image analysis. Nature methods *9*, 676–682.

57. de Bruyne, M., Foster, K., and Carlson, J.R. (2001). Odor coding in the Drosophila antenna.
Neuron *30*, 537–552.

58. Silbering, A.F., Rytz, R., Grosjean, Y., Abuin, L., Ramdya, P., Jefferis, G.S., and Benton, R.
(2011). Complementary function and integrated wiring of the evolutionarily distinct
Drosophila olfactory subsystems. The Journal of neuroscience : the official journal of the
Society for Neuroscience *31*, 13357–13375. 10.1523/JNEUROSCI.2360-11.2011.

59. Pnevmatikakis, E.A., and Giovannucci, A. (2017). NoRMCorre: An online algorithm for
piecewise rigid motion correction of calcium imaging data. J Neurosci Methods *291*, 83–94.
10.1016/j.jneumeth.2017.07.031.

60. Giovannucci, A., Friedrich, J., Gunn, P., Kalfon, J., Brown, B.L., Koay, S.A., Taxidis, J.,
Najafi, F., Gauthier, J.L., and Zhou, P. (2019). CaImAn an open source tool for scalable
calcium imaging data analysis. elife *8*, e38173.

61. Stringer, C., and Pachitariu, M. (2019). Computational processing of neural recordings from
calcium imaging data. Current Opinion in Neurobiology *55*, 22–31.
10.1016/j.conb.2018.11.005.

62. Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., and Dubourg, V. (2011). Scikit-learn: Machine learning in
Python. the Journal of machine Learning research *12*, 2825–2830.

63. Scheffer, L.K., Xu, C.S., Januszewski, M., Lu, Z., Takemura, S., Hayworth, K.J., Huang,
G.B., Shinomiya, K., Maitlin-Shepard, J., Berg, S., et al. (2020). A connectome and analysis
of the adult Drosophila central brain. eLife *9*, e57443. 10.7554/eLife.57443.

64. Praturu, A., and Sharpee, T.O. (2022). A Bayesian Approach to Hyperbolic Multi-
Dimensional Scaling. bioRxiv, 2022–10.

65. Strona, G., Nappo, D., Boccacci, F., Fattorini, S., and San-Miguel-Ayanz, J. (2014). A fast
and unbiased procedure to randomize ecological binary matrices with fixed row and column
totals. Nature communications *5*, 4114.

66. Zheng, Z., Lauritzen, J.S., Perlman, E., Robinson, C.G., Nichols, M., Milkie, D., Torrens, O.,
Price, J., Fisher, C.B., Sharifi, N., et al. (2018). A Complete Electron Microscopy Volume of
the Brain of Adult Drosophila melanogaster. Cell *174*, 730-743.e22.
10.1016/j.cell.2018.06.019.

SUPPLEMENTAL MATERIALS

Supplementary Figure S1: KC response properties.
A) Odor-evoked responses in three example KCs from odor trials in different movies (imaging sessions with overlapping but distinct panels) collected in the same MB. Tracking and assignment of odor responses to KCs is reliable across imaging sessions.
B) Comparison of KC response rate (mean and 95% CI, n=4 flies) and mean evoked OR firing rate (across glomeruli) in the Hallem dataset to each odor.
C) Observed fraction of KCs (mean and 95% CI, n=4 flies) responding to the indicated number of odors for a panel containing 17 odors. The distribution of the fraction of modeled KCs responding to different numbers of odors in the uniform, hemidraw, and hemibrain models (see Figure 3) are plotted for comparison.

D) Predicted KC response rates to narrowly activating (CO_2, methyl salicylate) and broadly activating (ethyl butyrate) odors, under assumptions of uniform connectivity. For selective odors, the input firing rate for the cognate OR was set to saturating firing rates (300 Hz) to estimate the upper bound for KC response rate. For ethyl butyrate, the observed firing rates across ORs in the Hallem dataset was used as input to the model.
E) Observed KC response rates to varying concentrations of the odors in **D**.

Figure S2: KC population responses to another example odor panel, illustrating the invariance of MB representational space across individuals.

A) Matrices of correlation distances from three different flies showing pairwise relationships between KC population responses in individual odor trials (upper triangles) or in trial-averaged responses for each odor (lower triangle).

B) Correlation distances between KC response vectors from different trials of the same odor. For each odor, the trial-trial KC response correlation was computed for all pairs of trials of this odor in each fly and averaged. The plot shows the mean and 95% CI of the fly averages. Compare to Figure 2E. Weaker odors tend to be less reliable than stronger odors.

Figure S3: The structure of olfactory glomerular inputs to third-order olfactory neurons and the impact on MB representations of odor.

A) Example reconstruction from the hemibrain connectome[63] of a uniglomerular DM2 PN and two example postsynaptic partners, a Kenyon cell (KC) and a lateral horn neuron (LHN). Synaptic connections between identified neurons contained within the hemibrain volume are fully described.

B) For each third-order olfactory neuron, the set of glomeruli providing direct presynaptic input via PNs was extracted from connectome datasets (hemibrain and FAFB). For this analysis, synaptic connections were binarized (e.g., all glomerular inputs >5 synapses were treated equally regardless of synapse count).

C) The mapping of glomerular inputs to KCs was represented as a binary matrix, where a 1 in cell(i,j) indicates that neuron i receives input from glomerulus j. The 'Curveball' algorithm was used to generate random matrices (null model) that preserve the row and column totals of the original matrix[65].

D) For each pair of glomeruli, a distribution of the number of KCs receiving co-input from the two glomeruli in each shuffled matrix was generated. The observed number of KCs receiving co-input from each pair of glomeruli in the hemibrain was compared to this distribution to generate a z-score. An example distribution and z-score for the over-convergent glomerular pair DM1 and DM3 are given.

E) Glomerular input structure to KCs in the hemibrain MB[63] (left), KCs in the FAFB MB[66] (center), and lateral horn neurons in the hemibrain LH (right). The z-score for each pair of glomeruli measures the degree to which the glomeruli are over- or under-convergent in the observed population, compared to null models. A large positive value indicates strong over-convergence, and a large negative value indicates strong under-convergence. The ordering of glomeruli is the same in all matrices and was based on k-means clustering of the hemibrain MB matrix.

F) Matrix of pairwise correlation distances between predicted KC responses to 17 odors in the hemidraw model (left). The mean pairwise relationships for observed KC responses are reproduced from Figure **3Ciii** (right) for ease of comparison.

G) Comparison of odor-odor correlation distances between observed KC responses and predicted KC responses in the hemidraw model. Each symbol is an odor pair.

H) Matrix of pairwise cosine distances between predicted KC responses (uniform, hemidraw, or hemibrain models), or observed KC responses, for 17 odors.

987
988

Figure S4: Transformation of odor representations in the fly olfactory circuit.

A) Mean (top row) and standard deviation (bottom row) across flies of pairwise correlation distances of odor representations in ORN axons ($n = 4$ flies), PN dendrites ($n = 6$ flies), PN boutons ($n = 6$ flies), and KC somata ($n = 4$ flies) for a panel of 17 odors.

B) Pairwise correlation distances of OR tuning in the Hallem dataset (left) and ORN axon terminal responses measured in this study (right) for a panel of 17 odors.

C) Comparison of fractional KC response rates with odor-evoked **Ci)** firing rates across ORs in the Hallem dataset (grey); **Cii)** ORN terminal responses (grey); **Ciii)** PN dendrite responses (grey); and **Civ)** PN bouton responses (grey) for 17 odors. ORN terminal, PN dendrite, PN bouton, and KC responses are mean and 95% CI across flies of the ROI-averaged evoked response to each odor in each fly.

D) Distribution of odor-odor correlation distances for representations at each stage of olfactory processing.

990 **Supplemental Table S1:** Odor information and number of observations (flies) for each odor in
991 KC somata datasets.

Odor	Abbreviation	InChI	Sigma-Aldrich Cat. No.	CAS	No. flies sampled (n)
1-hexanol	1-6ol	InChI=1S/C6H14O/c1-2-3-4-5-6-7/h7H,2-6H2,1H3	471402	111-27-3	22
1-octanol	1-8ol	InChI=1S/C8H18O/c1-2-3-4-5-6-7-8-9/h9H,2-8H2,1H3	297887	111-87-5	16
1-pentanol	1-5ol	InChI=1S/C5H12O/c1-2-3-4-5-6/h6H,2-5H2,1H3	138975	71-41-0	10
1-penten-3-ol	1-p3ol	InChI=1S/C5H10O/c1-3-5(6)4-2/h3,5-6H,1,4H2,2H3	1984	616-25-1	7
2-butanone	2-but	InChI=1S/C4H8O/c1-3-4(2)5/h3H2,1-2H3	360473	78-93-3	21
2-heptanone	2-hep	InChI=1S/C7H14O/c1-3-4-5-6-7(2)8/h3-6H2,1-2H3	537683	110-43-0	14
acetic acid	aac	InChI=1S/C2H4O2/c1-2(3)4/h1H3,(H,3,4)	695092	64-19-7	5
B-citronellol	b-cit	InChI=1S/C10H20O/c1-9(2)5-4-6-10(3)7-8-11/h5,10-11H,4,6-8H2,1-3H3	C83201	106-22-9	12
benzaldehyde	bnz	InChI=1S/C7H6O/c8-6-7-4-2-1-3-5-7/h1-6H	418099	100-52-7	20
delta-decalactone	d-10lac	InChI=1S/C10H18O2/c1-2-3-4-6-9-7-5-8-10(11)12-9/h9H,2-8H2,1H3	74026	705-86-2	5
E2-hexenal	e2-h	InChI=1S/C6H10O/c1-2-3-4-5-6-7/h4-6H,2-3H2,1H3/b5-4+	158131000	6728-26-3	11
ethyl butyrate	eb	InChI=1S/C6H12O2/c1-3-5-6(7)8-4-2/h3-5H2,1-2H3	E15701	105-54-4	14
ethyl cinnamate	ec	InChI=1S/C11H12O2/c1-2-13-11(12)9-8-10-6-4-3-5-7-10/h3-9H,2H2,1H3/b9-8+	66761	103-36-6	5
ethyl propionate	ep	InChI=1S/C5H10O2/c1-3-5(6)7-4-2/h3-4H2,1-2H3	112305	105-37-3	16
eugenol	eug	InChI=1S/C10H12O2/c1-3-4-8-5-6-9(11)10(7-8)12-2/h3,5-7,11H,1,4H2,2H3	E51791	97-53-0	7
gamma-hexalactone	g-6lac	InChI=1S/C6H10O2/c1-2-5-3-4-6(7)8-5/h5H,2-4H2,1H3	68554	695-06-7	6
hexanal	6al	InChI=1S/C6H12O/c1-2-3-4-5-6-7/h6H,2-5H2,1H3	115606	66-25-1	15
isoamyl acetate	iaa	InChI=1S/C7H14O2/c1-6(2)4-5-9-7(3)8/h6H,4-5H2,1-3H3	W205532	123-92-2	12
linalool	lin	InChI=1S/C10H18O/c1-5-10(4,11)8-6-7-9(2)3/h5,7,11H,1,6,8H2,2-4H3	L2602	78-70-6	10
methyl octanoate	mo	InChI=1S/C9H18O2/c1-3-4-5-6-7-8-9(10)11-2/h3-8H2,1-2H3	21719	111-11-5	9
methyl salicylate	msl	InChI=1S/C8H8O3/c1-11-8(10)6-4-2-3-5-7(6)9/h2-5,9H,1H3	M6752	119-36-8	14
pentyl acetate	pa	InChI=1S/C7H14O2/c1-3-4-5-6-9-7(2)8/h3-6H2,1-2H3	109584	628-63-7	15
propyl acetate	pac	InChI=1S/C5H10O2/c1-3-4-7-5(2)6/h3-4H2,1-2H3	40858	109-60-4	7
valeric acid	va	InChI=1S/C5H10O2/c1-2-3-4-5(6)7/h2-4H2,1H3,(H,6,7)	240370	109-52-4	13

992
993

994 **Supplemental Table S2:** Mean, 95% CI, SEM, and number of observations (flies) of the
995 correlation distance for every unique odor pair in KC somata datasets.

996
997 **Supplemental Table S3:** Reduced list of Mordred molecular descriptors used in this study.

998
999 **Supplemental Table S4:** References contributing to a secondary natural odor source database.
1000