CRB3 navigates Rab11 trafficking vesicles to promote γTuRC assembly during

ciliogenesis

Bo Wang[1, 2, †], Zheyong Liang[1, 3, †], Tan tan[4], Miao Zhang[1, 2], Yina Jiang[5], Yangyang

Shang[1, 2], Xiaoqian Gao[1, 2], Shaoran Song[1, 2], Ruiqi Wang[1, 2], He Chen[1, 2], Jie Liu[1, 2],

Juan Li[1, 2], Yu Ren[6], and Peijun Liu[1, 2] *

[1]Center for Translational Medicine, the First Affiliated Hospital of Xi'an Jiaotong

University, Xi'an 710061, Shaanxi, P.R. China.

[2]Key Laboratory for Tumor Precision Medicine of Shaanxi Province, the First

Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, Shaanxi, P.R. China.

[3]Department of Cardiovascular Surgery, the First Affiliated Hospital of Xi'an Jiaotong

University, Xi'an 710061, Shaanxi, P.R. China.

[4]Center for Precision Medicine, Affiliated to The First People's Hospital of Chenzhou,

University of South China, Chenzhou 423000, Hunan, P.R. China.

[5]Department of Pathology, the First Affiliated Hospital of Xi'an Jiaotong University,

Xi'an 710061, Shaanxi, P.R. China.

[6]Department of Breast Surgery, the First Affiliated Hospital of Xi'an Jiaotong

University, Xi'an 710061, Shaanxi, P.R. China.

† These authors contributed equally to the work.

* Correspondence:

Peijun Liu

E-mail: liupeijun@xjtu.edu.cn

Running Title: CRB3 regulates ciliogenesis

Keywords: Primary cilium, Polarity protein, Breast cancer, CRB3, γ-TuRC assembly,

Rab11-positive endosome

Abstract

The primary cilium plays important roles in regulating cell differentiation, signal transduction, and tissue organization. Dysfunction of the primary cilium can lead to ciliopathies and cancer. The formation and organization of the primary cilium are highly associated with cell polarity proteins, such as the apical polarity protein CRB3. However, the molecular mechanisms by which CRB3 regulates ciliogenesis and the location of CRB3 remain unknown. Here, we show that CRB3, as a navigator, regulates vesicle trafficking in γ-TuRC assembly during ciliogenesis and cilium-related Hh and Wnt signaling pathways in tumorigenesis. *Crb3* knockout mice display severe defects of the primary cilium in the mammary ductal lumen and renal tubule, while mammary epithelial-specific *Crb3* knockout mice exhibit promotion of ductal epithelial hyperplasia and tumorigenesis. CRB3 is essential for lumen formation and ciliary assembly in the mammary epithelium. We demonstrate that CRB3 localizes to the basal body and that CRB3 trafficking is mediated by Rab11-positive endosomes. Significantly, CRB3 interacts with Rab11 to navigate GCP6/Rab11 trafficking vesicles to CEP290, resulting in intact γ-TuRC assembly. In addition, CRB3-depleted cells are unresponsive to the activation of the Hh signaling pathway, while CRB3 regulates the Wnt signaling pathway. Therefore, our studies reveal the molecular mechanisms by which CRB3 recognizes Rab11-positive endosomes to navigate apical vesicle trafficking in effective ciliogenesis, maintaining cellular homeostasis and tumorigenesis.

Introduction

Epithelial tissues are the most common and widely distributed tissues in the body, and they perform several particular functions. The integrity of the cellular architecture composed of epithelial cells is necessary for these functions. Establishing apical-basal cell polarity within epithelial cells is a crucial biological process for epithelial homeostasis. Unfortunately, the loss of apical-basal cell polarity is one of the most significant hallmarks of advanced malignant tumors[1], and several reports have shown that the loss of some polarity proteins causes disordered cell polarity within

epithelial cells, which cannot tightly control cellular growth and migration, ultimately

leading to tumor formation, progression, and metastasis[2, 3]. However, the

relationship between polarity proteins and tumorigenesis remains incomplete.

The Crumb complex, located apically, is vital in regulating and maintaining

apical-basal cell polarity within epithelial cells. In mammals, three Crumbs orthologs

are CRB1, CRB2, and CRB3, and only CRB3 is widely expressed in epithelial

cells[4]. Our previous studies have shown that CRB3 is expressed at low levels in

renal and breast cancers, and abnormal CRB3 expression leads to the disrupted

organization of MCF10A cells in three-dimensional (3D) cultures[5-7]. Inhibition of

CRB3 impairs contact inhibition and leads to migration, invasion and tumorigenesis

of cancer cells[6, 8]. Contact inhibition results in growth arrest and causes epithelial

cells to enter the quiescent phase, inducing primary cilium formation.

The primary cilium, an antenna-like microtubule-based organelle, is a sensorial

antenna that regulates cell differentiation, proliferation, polarity, and tissue

organization in most types of mammalian cells[9]. Dysfunction of the primary cilium

can lead to developmental and degenerative disorders called ciliopathies, such as

polycystic kidney disease, nephronophthisis, retinitis pigmentosa, and Meckel

syndrome[10]. Remarkably, many cancers, including melanoma and breast, renal,

pancreatic, and prostate cancer, exhibit loss of the primary cilium, most likely during

the early stages of tumorigenesis[11-15]. The primary cilium is a significant extension

on the apical surfaces of epithelial cells and is maintained by polarized vesicular

traffic[16]. Studies have shown that CRB3 localizes in the primary cilium and that

depletion of CRB3 leads to primary cilium loss in Madin-Darby canine kidney

(MDCK) cells[17, 18], indicating that CRB3 is necessary for the initiation of

ciliogenesis. However, how CRB3 promotes ciliogenesis on the apical surfaces of

epithelial cells remains unknown.

Here, we describe a novel conditional deletion of *Crb3* in mice and show that Crb3 is

required to assemble the primary cilium in the mammary gland, kidney and MEF cells

from *Crb3* knockout mice and that *Crb3* deletion promotes breast cancer progression

in vivo. Additionally, we found that CRB3 interacts with Rab11 and navigates

GCP6/Rab11 trafficking vesicles to the basal body of the primary cilium in mammary

epithelial cells. Furthermore, we identified that CRB3 regulates the ciliary Hedgehog

(Hh) and Wnt signaling pathways in breast cancer. Thus, there is unexpected diversity

in the polarity protein CRB3 navigating polarized Rab11-dependent traffic, leading to

a novel assembly mechanism of the γ-tubulin ring complex (γTuRC) in ciliogenesis.

Results

***Crb3* deletion mice exhibit smaller size and anophthalmia.**

Since *Crb3* knockout mice die shortly after birth[19], we generated a novel transgenic

mouse model with conditional deletion of *Crb3* using the Cre-loxP system. The loxP

sites flanked either side of exon 3 in the *Crb3* gene (Supplementary Fig. 1A). In

mammals, CRB3 has two isoforms, CRB3a and CRB3b, distinguished by alternative

splicing within the fourth exon of the *CRB3* gene, which in turn produces a protein

with 23 amino acid differences at the C terminus[17]. According to the construction

strategy, we are knocking out both isoforms *Crb3a* and *Crb3b* in the mice. The

positive embryonic stem (ES) cell clone was confirmed for the wild-type gene

(*Crb3*^{wt/wt}) and recombinant allele (*Crb3*^{wt/fl}) by Southern blotting (Supplementary Fig.

1B). The genotypes of the wild type, heterozygotes and homozygotes (*Crb3*^{fl/fl}) were

detected with specific primers (Supplementary Fig. 1C). We first intercrossed *Crb3*^{fl/fl}

with CMV enhancer/chicken β-actin promoter (CAG)-Cre mice and found that

Crb3^{fl/fl}, *Crb3*^{wt/fl}; CAG-Cre and *Crb3*^{fl/fl}; CAG-Cre pups were born normally and

well developed in size. However, most *Crb3*^{fl/fl}; CAG-*Cre* pups died within a few

days after birth. Few surviving *Crb3*^{fl/fl}; CAG-*Cre* mice were smaller and showed

anophthalmia than littermate *Crb3*^{fl/fl} mice at 4 weeks old (Fig. 1A-B). The eye is an

organ with perfect apical-basal cell polarity. However, *Crb3* knockout mice showed

ocular abnormalities. Together, these results indicate that Crb3 is necessary for eye

development.

Deletion of mammary epithelial-specific *Crb3* causes ductal epithelial

hyperplasia and tumorigenesis.

Based on our earlier studies, which showed that inhibiting CRB3 impairs the

organization of breast epithelium, we set out to directly explore the role of Crb3 in

mammary gland development. We first developed epithelial cell-specific deletion of

Crb3 in the mammary gland by intercrossing *Crb3*^fl/fl^ mice with mouse mammary

tumor virus long terminal repeat promoter (MMTV)-*Cre* mice. We confirmed the

expression of Crb3 in the mammary gland using immunoblotting (both isoforms

Crb3a and Crb3b), real-time quantitative PCR, and immunohistochemistry, and Crb3

expression was deficient in mammary epithelial cells with Crb3 knockout

(Supplementary Fig. 1D-F).

To assess the function of Crb3 in the development of mammary epithelial cells, we

analyzed whole mammary mounts from 8-week-old virgin *Crb3*^fl/fl^;MMTV-*Cre* and

littermate *Crb3*^fl/fl^ mice. *Crb3*^fl/fl^;MMTV-*Cre* mice displayed significantly increased

numbers of terminal end buds (TEBs) and bifurcated TEBs compared to *Crb3*^fl/fl^ mice

(Fig. 1 C-E). Histopathology results revealed that normal ductal epithelial cells were

arranged as a monolayer in *Crb3*^fl/fl^ mice, while *Crb3* knockout led to ductal epithelial

hyperplasia with increased ductal thickness in *Crb3*^fl/fl^;MMTV-*Cre* mice (Fig. 1 F).

To investigate the role of Crb3 in breast cancer tumorigenesis, we observed mammary

glands from 9-week-old virgin polyomavirus middle T antigen (PyMT)-cKO-Crb3

and PyMT-WT mice. The mammary glands of PyMT mice progressed to early

carcinomas and lost their normal epithelial architecture[20]. The PyMT mouse model

is widely used to study tumor initiation and development, with a histological

progression similar to that of human breast cancer. PyMT-WT mice exhibited loss of

epithelial features and developed breast cancer, while PyMT-cKO-Crb3 mice

developed larger tumors with faster tumor progression and a poorly differentiated

phenotype (Fig. 1G). Thus, these results suggest that Crb3 knockout leads to ductal

epithelial hyperplasia and promotes tumorigenesis.

CRB3 knockdown inhibits lumen formation in acini of mammary epithelial cells.

MCF10A cells are spontaneously immortalized human breast epithelial cells with the

characteristics of normal breast epithelium[21]. In 3D basement membrane culture

conditions, MCF-10A cells form acini similar to mammary epithelial acini. This

makes them a valuable system for studying the morphogenesis and oncogenesis of the

mammary epithelium[22]. To confirm the function of CRB3 in mammary epithelial

acini formation, we first knocked down the expression of both CRB3a and CRB3b

isoforms using lentiviral shRNA in MCF10A cells (Supplementary Fig. 2A and B).

Under the 3D culture system, we found that MCF10A cells infected with a

non-targeting shRNA lentivirus formed acini after 6 days (D6) and eventually formed

polarized acini with a hollow lumen at D14 (Fig. 2A). However, after CRB3

knockdown, MCF10A cells formed more and larger aberrant acini without a lumen

after D6 (Fig. 2A and B). Cell proliferation assays of MCF10A cells showed that

CRB3 knockdown increased the proliferation rate and accelerated G1 to S phase

progression (Supplementary Fig. 2C-E).

To investigate the reason for aberrant acini without lumens, we further studied

apoptosis and mitotic spindle orientation during lumen formation in a 3D culture

system. The immunofluorescence (IF) results showed that more internal

cleaved-caspase 3-positive acini were observed in the control groups than in the

CRB3 knockdown groups at D9 (Fig. 2C and D). Moreover, CRB3b overexpression

significantly promoted apoptosis in breast cancer cells (Supplementary Fig. 2F-H).

Similarly, the staining of α-tubulin in mitotic cells demonstrated more misorientation

of the mitotic spindle in CRB3 knockdown groups (Fig. 2E and F).

In addition to verifying the effect of *Crb3* deletion on promoting proliferation and

inhibiting apoptosis *in vivo*, we detected these markers in primary tumor tissue from

PyMT-WT and PyMT-cKO-Crb3 mice. *Crb3* deletion increased the numbers of

proliferative and mitotic cells, and decreased the numbers of apoptotic cells in

primary tumors (Fig 2G and H). These results demonstrate that CRB3 could promote

the proliferation of mammary epithelial cells, disrupt the mitotic spindle orientation,

and protect the internal cells from apoptosis during acini formation, ultimately leading

to irregular lumen formation and tumor progression.

CRB3 regulates primary cilium formation.

The primary cilium is an apical antenna-like extensions in epithelial cells that is

known to play an important role in controlling lumen formation[23]. Previous studies

have reported that CRB3 knockdown leads to primary cilium loss in MDCK cells[17,

18]. However, the molecular mechanism that regulates ciliogenesis on apical surfaces

is still unclear. To investigate the altered effect of CRB3 on ciliogenesis both *in vivo*

and *in vitro*, we examined the formation of primary cilia in control MCF10A cells and

CRB3 knockdown cells. We found that control MCF10A cells formed primary cilia

after becoming confluent, while CRB3 knockdown led to significant ciliogenesis

defects (Fig. 3A). The number of cells with primary cilium was significantly reduced

compared to the control group, but the length of the primary cilium that did exist was

not different (Fig. 3B). Conversely, CRB3b conditional overexpression by adding

doxycycline (+dox) resulted in restoring ciliary assembly in MCF7 cells (Fig. 3C,

Supplementary Fig. 3A). CRB3b overexpression increased the proportion of primary

cilium formation in breast cancer cells, and the length of the restored primary cilium

was increased (Fig. 3D). Importantly, the primary cilium could be directly visualized

using scanning electron microscopy (SEM) (Fig. 3E). After CRB3 knockdown, the

number of MCF10A cells with primary cilia was significantly decreased, while

CRB3b conditional overexpression did not increase these populations (Fig. 3F).

To further support these findings *in vivo*, we used breast and kidney tissue to assess

ciliogenesis in the *Crb3* knockout mouse model. Renal epithelial cells can form

prominent primary cilia, and the absence of primary cilia leads to ciliopathies. We

used immunofluorescence to detect ciliary formation in the mammary ductal lumen

and renal tubule. Compared to tissues of *Crb3*$^{\text{fl/fl}}$ mice, *Crb3* knockout led to the

absence of the primary cilium in the mammary ductal lumen and renal tubule from

Crb3$^{\text{fl/fl}}$; CAG-*Cre* mice (Fig. 3G, H). Specifically, the proportion of cells forming the

primary cilium plummeted (Fig. 3G, H). This same phenomenon was also observed in

mouse embryonic fibroblasts (MEFs) from *Crb3*$^{\text{fl/fl}}$; CAG-*Cre* mice (Supplementary

Fig. 3B, C). We conclude that CRB3 plays a central role in ciliogenesis in various

tissues and mammary cells.

CRB3 localizes to the basal body of the primary cilium.

CRB3, a polarity protein, is typically localized on the tight junctions of the apical

epithelium membrane[24]. Additionally, we noted that CRB3b is found to be closely

colocalized with centrosomes at prophase, metaphase, and anaphase[17]. To

investigate the relationship between CRB3 and ciliary formation, we first examined CRB3 localization in MCF10A cells expressing CRB3-GFP. In MCF10A cells, exogenous CRB3b was mainly localized at cell tight junctions and could colocalize with pericentrin (a centrosome marker) (Fig. 4A). And approximately 40% of cells showed this co-localization of exogenous CRB3b and pericentrin (Fig. 4B). We verified this phenomenon by observing the localization of endogenous CRB3 (both isoforms CRB3a and CRB3b) with another centrosome marker. Similarly, we detected that CRB3 accumulated on one side of the cytoplasm and had a CRB3 foci located at the basal body represented by γ-tubulin in MCF10A cells (Fig. 4C, D). Importantly, CRB3 knockdown disturbed the accumulation of γ-tubulin at the basal body in confluent quiescence cells (Fig. 4C, E). To demonstrate the relationship of this colocalization to the primary cilium, we employed acetylated tubulin to mark the primary cilium. Double immunostaining showed that this CRB3 foci was the basal body of the primary cilium, and CRB3 knockdown inhibited ciliary assembly in MCF10A cells (Fig. 4F, G). These results suggest that CRB3 is located at the basal body of the primary cilium and that CRB3 knockdown disturbs the accumulation of γ-tubulin in quiescent cells.

CRB3 trafficking is mediated by Rab11-positive endosomes.

To better investigate the molecular mechanisms of CRB3 in regulating primary cilium formation, we examined some important genes related to ciliogenesis. However, CRB3 knockdown did not affect the mRNA expression of these genes in MCF10A cells (Supplementary Fig. 4A). Then, we performed coimmunoprecipitation (co-IP) tandem mass spectrometry using tagged exogenous CRB3b as bait to identify its interacting proteins (Supplementary Fig. 4B). Pathway aggregation analysis of these proteins revealed that these interacting proteins of CRB3b were significantly involved in Golgi vesicle transport and vesicle organization (Fig. 5A). We focused on these pathways because smaller distal appendage vesicle (DAV) formation is critical for ciliogenesis initiation, and it requires the Rab GTPase Rab11-Rab8 cascade to function[25]. Interestingly, some Rab small GTPase family members identified as CRB3b-binding proteins were aggregated in these pathways, such as Rab10, Rab11A,

Rab11B, Rab14, Rab1A, Rab1B, Rab21, Rab2A, Rab32, Rab35, Rab38, Rab5B,

Rab5C, and Rab6A. Several centriolar proteins were also identified, including tubulin

gamma-1, tubulin gamma-2, CENP-E, CEP290, CEP192, CEP295, and CEP162 (Fig.

5B).

Rab11-positive vesicles bind to centrosomal Rabin8, leading to the formation of

ciliary membrane[26]. This finding suggested that the trafficking of Rab11-positive

vesicles and polarized vesicles is essential for early ciliary assembly. Next, we

examined the polarized vesicle traffic of CRB3 in MCF10A cells. Our studies found

that CRB3 could colocalize with EEA1-positive early endosomes, CD63-positive late

endosomes and Rab11-positive recycling endosomes (Fig. 5C). In addition, after

treatment with dynasore, an endocytosis inhibitor, CRB3 significantly accumulated at

the cell membrane after 2 hours, and the colocalization of CRB3 with EEA1-, CD63-,

and Rab11-positive endosomes was significantly decreased (Fig. 5C, D). We have

previously reported that CRB3 regulates cell contact inhibition, which is significantly

downregulated in confluent MCF10A cells[6]. Moreover, dynasore could rescue the

CRB3 downregulation in confluent MCF10A cells (Fig. 5E). The knockdown of

CRB3 did not affect Rab11 expression (Supplementary Fig. 4C). These results

suggest that CRB3 is highly dynamic among the endosome system, and in particular,

Rab11-positive endosomes are involved in the intracellular trafficking of CRB3.

CRB3 knockdown destabilizes γTuRC assembly during ciliogenesis.

According to our findings, we assumed that CRB3 might be involved in polarized

vesicle trafficking. The γ-Tubulin ring complex (γTuRC) is a central regulator of

microtubule nucleation and a nucleation site of α-Tubulin and β-Tubulin at the

microtubule organizing center in mitotic cells[27]. Additionally, the primary cilium,

a microtubule-based structure attached to the basal body, is transformed from the

mother centriole. As we found that CRB3 knockdown disturbed the accumulation of

γ-tubulin prompted us to analyze γTuRC assembly in ciliogenesis. The γTuRC is

composed of multiple copies of the γ tubulin small complex (γTuSC), GCP2 and

GCP3, plus GCP4, GCP5, and GCP6[27, 28] (Fig. 5F). Our findings indicate that

CRB3 downregulation did not affect the expression of γTuRC subunits (Fig. 5G).

However, we performed co-IPs directed against GCP6 in CRB3-depleted cells and
assessed the relative levels of both γTuSC-specific proteins compared with the control
shRNA-treated cells. Interestingly, CRB3 knockdown significantly reduced the
interaction of GCP6 with GCP3 (Fig. 5H). To verify this phenomenon, we observed
the localization of GCP3 with GCP6 foci in MCF10A cells. Correspondingly, CRB3
downregulation significantly disturbed the colocalization of GCP3 with GCP6 foci in
quiescence cells (Fig. 5I). To investigate whether CRB3 facilitate γTuRC assembly,
we fractionated MCF10A cytoplasmic proteins on sucrose gradients according to
published methods[29]. We noticed that γTuSC were sedimented mainly in fractions 3,
while γTuRC sedimented in fractions 6 (Fig. 5J). However, CRB3 downregulation
caused most of GCP6 and GCP5 to be precipitated independently of γTuRC in the
low-density fractions (Fig. 5J). Together, these results demonstrate that the depletion
of CRB3 disrupts molecular interactions between the γTuRC proteins, resulting in a
failure of microtubule formation in the primary cilium. However, this depletion may
not affect the expression of these molecules during γTuRC assembly. Based on these
findings, we hypothesize that CRB3 could facilitate the polarized vesicle trafficking
of some γTuRC-specific proteins through Rab11-positive endosomes.

CRB3 interacts with Rab11.
To fully prove this hypothesis, we first examined the interaction of CRB3 with
Rab11-positive endosomes and γTuRC-specific proteins in MCF10A cells. Using
CRB3 as bait, we identified that CRB3 could interact with Rab11, but not GCP6 and
GCP3 (Fig. 6A). In contrast, Rab11 could interact with CRB3 and GCP6 instead of
GCP3 by tagging Rab11 as bait (Fig. 6B). The co-IP results showed that CRB3
knockdown did not affect Rab11 interacting with GCP6, but the level of Rab11
binding to GCP6 tended to decrease in MCF10A cells with CRB3 knockdown (Fig.
6C). Consistent with other reports[30], Rab11 knockdown significantly inhibited
primary cilium formation in MCF10A cells (Supplementary Fig. 5A-C).
Based on these results, we further detected the region of CRB3b that interacts with
Rab11 by using coexpression and co-IP in HEK293 cells. CRB3b is composed of a
signal peptide (SP), extracellular domain, transmembrane (TM), FERM-binding

domain (FBD), and carboxy-terminal PDZ-binding domain (PBD). We constructed

CRB3-GFP fusion proteins by fusing GFP tags to the extracellular domain, as

previously reported[31] (Fig. 6D). According to these domains, we also constructed

the truncations of CRB3-GFP fusion proteins with serial C-terminal deletions (Fig. 6E)

and another Flag-Rab11a fusion protein. The coexpression and co-IP results showed

that Flag-Rab11a interacts with full-length CRB3 (1-120), CRB3 (1-116), CRB3

(1-83), and CRB3 (1-58) but not CRB3 (1-26) (Fig. 6F). Therefore, the region of

CRB3b that interacts with Rab11 is amino acids 27 to 58.

Rab11 is a small GTP-binding protein that plays an essential role in regulating the

dynamics of recycling endosomes, which must undergo GDP/GTP cycles. In

particular, Rab11a[S25N] is a GDP-locked form, while Rab11a[S20V] and

Rab11a[Q70L] are GTP-locked forms[30]. To further investigate the interaction

between the GTP-bound form of Rab11a and CRB3b, we constructed these mutant

variants of Rab11a. Next, we found that CRB3b interacts more strongly with

Rab11a[Q70L] and Rab11a[S20V], while weakly interacting with Rab11a[S25N],

compared to Rab11aWT (Fig. 6G). Together, these results indicate that the interaction

of CRB3b and Rab11a is dependent on the GTPase activity of Rab11a.

CRB3 navigates GCP6/Rab11 trafficking vesicles to CEP290 in the primary

cilium.

Since CRB3 knockdown does not affect the interaction with Rab11-positive

endosomes and γTuRC-specific proteins GCP6, our focus was to determine whether

CRB3 affects the localization of GCP6/Rab11 trafficking vesicles to the basal body of

the primary cilium. We reviewed the identification of CRB3 interacting proteins and

found some centriolar proteins. CEP290, for example, is located in the transition zone

of the primary cilium and is required for the formation of microtubule-membrane

linkers[32]. Then, we verified that exogenous CRB3b could bind to CEP290, Rab11,

GCP6 (Fig. 7A), and Rab11 also interacted with CEP290, CRB3-GFP, GCP6 (Fig.

7B). Similar to Fig. 6C, the amount of GCP6 bound to Rab11 was reduced within

CRB3 knockdown. Importantly, CRB3 knockdown made it difficult for Rab11 to bind

CEP290 (Fig. 7C). We corroborated these findings by checking the colocalization of

the basal body foci of GCP6 and γ-tubulin, γ-tubulin and Rab11 in MCF10A and MEF cells. In quiescence control cells, GCP6 foci and γ-tubulin formed a prominent focus at the basal body of the primary cilium, as observed by immunofluorescence. However, CRB3 downregulation eliminated this focus and significantly disturbed the basal body foci of GCP6 and γ-tubulin in MCF10A cells (Fig. 7D), and γ-tubulin and Rab11 in MEF cells (Fig. 7E). At the beginning of this experiment, we also attempted to show the interaction of CRB3, Rab11 and GCP6 by expressing fluorescence tagged GCP6 protein and using FRET assay. However, the centrosomal foci of fluorescent GCP6 was difficult to capture due to weak signal, and this may require re-overexpression of ninein-like protein (Nlp) to promote γTuRC enrichment[29, 33]. Although CRB3 knockdown had little effect on the formation of GCP6/Rab11 trafficking vesicles, it significantly inhibited γTuRC assembly during ciliogenesis. This, in turn, impacted the transport of GCP6/Rab11 trafficking vesicles to the basal body of the primary cilium. Since CRB3 affected the binding of Rab11 and CEP290, we speculate that CRB3 may act as a navigator, navigating GCP6/Rab11 trafficking vesicles to the basal body of the primary cilium.

CRB3 regulates the Hh and Wnt signaling pathways in tumorigenesis

To investigate the role of CRB3 in regulating ciliary assembly in breast cancer, we examined the relationship between CRB3 and the primary cilium in breast cancer tissues. We found that CRB3 was localized at the subapical surface of the mammary gland lumen in the paracarcinoma tissues, but no obvious expression or localization in breast cancer tissues (Fig. 8A). Immunofluorescence analysis showed that the intact primary cilium was observed in the adjacent para-carcinoma tissues, whereas the number of cells with primary cilia was much lower in breast cancer tissues (Fig. 8B, C). Consistent with previous findings, CRB3 was localized at the basal body of the primary cilium in adjacent para-carcinoma tissues (Fig. 8D). These results again suggest that a defect in CRB3 expression inhibits ciliary assembly, which could be involved in tumorigenesis.

To investigate the effects of primary cilium absence, we examined alterations in the Hedgehog (Hh) and Wnt signaling pathways in CRB3-depleted cells. Firstly, we

detected the translocation of SMO into the primary cilium and the expression of the

target gene GLI1. SAG, a potent Smoothened (SMO) receptor agonist, activates the

Hh signaling pathway. However, in MCF10A cells treated with SAG, CRB3

knockdown did not promote SMO translocation into the primary cilium (Fig 8E).

After SAG treatment, the mRNA expression of GLI1 was significantly decreased in

CRB3-depleted cells (Fig 8F). Crb3 knockout did not alter GLI1 expression and

subcellular localization in primary tumors of PyMT-cKO-Crb3 mice (Fig 8H). We

previously reported that CRB3 expression was high in immortalized mammary

epithelial cells, whereas loss of CRB3 expression was observed in breast cancer

cells[6]. On the other hand, western blotting showed that CRB3 knockdown resulted

in significant downregulation of GSK3-β in MCF10A cells. Additionally, β-catenin,

the major effector of the canonical Wnt signaling pathway, was upregulated. In breast

cancer cells, CRB3b overexpression led to upregulation of GSK3-β and

downregulation of β-catenin (Fig. 8G, Supplementary Fig. 6). Notably, β-catenin was

significantly upregulated and markedly localized in the nucleus of PyMT-cKO-Crb3

mice (Fig 8I). Overall, these data suggest that CRB3 knockdown cannot activate the

Hh signaling pathway but can activate the Wnt signaling pathway, which is ultimately

contributes to tumorigenesis.

Discussion

Apical-basal cell polarity is essential for cellular architecture, development and

homeostasis in epithelial tissues. However, most malignant cancers lack cell polarity

due to unconstrained cell cycle and cell migration with decreased adhesion. CRB3, an

important apical protein, is significantly reduced in various tumor cells and tissues,

and promotes metastasis and tumor formation in nude mice[5, 6, 8, 34-36]. However,

it has also been reported that CRB3 sequencing results are upregulated in breast

cancer samples[37]. Here, we generated a novel transgenic mouse model with

conditional deletion of Crb3 to investigate the role of polarity proteins in

tumorigenesis. *Crb3* knockout mice were reported to have cystic kidneys, improper

airway clearance in the lung, villus fusion, apical membrane blebs, and disrupted

microvilli in intestine airway clearances, which demonstrated that CRB3 is crucial for the development of the apical membrane and epithelial morphogenesis[19, 38]. Importantly, our studies in the *Crb3* knockout mouse model suggest that, in addition to death after birth, it is characterized by smaller size, ocular abnormality, bronchial smooth muscle layer defect, reduced pancreatic islets and hyperplasia mammary ductal and renal tubular epithelium (some data not shown). These important phenotypic changes are due to disruption of epithelial polarity homeostasis, which may ultimately lead to tumorigenesis.

In addition, we observed the absence of primary cilia in the mammary ductal lumen and renal tubule in the *Crb3* knockout mouse model. Consistent with other reports, CRB3 knockdown regulated ciliary assembly in MDCK cells[17, 18]. Furthermore, CRB3 was localized in the inner segments of photoreceptor cells and concentrated with the connecting cilium throughout the development of the mouse retina[39]. These ocular defect phenotypes closely resemble those of EHD1 knockout mice, which regulate ciliary vesicle formation in primary cilium assembly[25, 40, 41]. Hence, CRB3 can affect primary cilium formation in various epithelial tissues, such as the breast, kidney, and retina.

Previous literature has reported that CRB3 localizes in the primary cilium itself in differentiated MDCK cells[17, 18], but we mainly found that it is not located on the basal body of the primary cilium in mammary epithelial cells. The immunofluorescence results in these literatures showed that CRB3 was scattered on the primary cilia, with a strong foci at the basal body. Notably, in rat kidney collecting ducts, the localization of CRB3 on primary cilia was significantly reduced, while the obvious localization was at the basal body[18]. Another study also reported the co-localization of CRB3b and γ□tubulin in MDCK cells[17]. This finding is consistent with our conclusion, and we also verified its co-localization with the centrosome by overexpressing CRB3b in mammary epithelial cells, indicating that CRB3 mainly localizes to the basal body of the primary cilium. Since CRB3b also alters primary cilium formation in the mammary epithelium, it is important to investigate the molecular mechanisms by which CRB3 regulates primary cilium

formation.

Polarized vesicular traffic plays a crucial role in extending the primary cilium on
apical epithelial surfaces. However, the molecular mechanism and its relationship to
tumorigenesis remain unclear. Here, we reveal that Rab11-positive endosomes
mediate the intracellular trafficking of CRB3 and that CRB3 can navigate
GCP6/Rab11 trafficking vesicles to CEP290, promoting correct γTuRC assembly
during primary cilium formation. When cells form tight junctions, CRB3 is mainly
localized to the cell membrane, and subsequently the endosomes participate in the
intracellular degradation process of CRB3 on the cell membrane. The intracellular
CRB3 can bind to Rab11 through the extracellular domain, which in turn participates
in primary cilium assembly. Additionally, in the presence of CRB3, the epithelium can
form contact inhibition and an intact primary cilium in quiescence to achieve cellular
homeostasis. In contrast, the loss of contact inhibition and primary cilium formation
with CRB3 deletion can activate the Wnt signaling pathway, affect the Hh signaling
pathway, and disrupt the imbalance of this cellular homeostasis, resulting in
significant cell proliferation during tumorigenesis (Fig. 9).

The formation of the primary cilium has been divided into several distinct phases. It
begins with the maturation of the mother centriole, also known as the
centriole-to-basal-body transition[9, 42]. DAVs, which are small cytoplasmic vesicles
originating from the Golgi and the recycling endosome, accumulate in the vicinity of
the distal appendages of the mother centriole[25, 43, 44]. In growing RPE1 cells,
Rab8a is localized to cytoplasmic vesicles and the Golgi/trans-Golgi network, while
Rab8-positive vesicles are rapidly redistributed to the mother centriole by binding to
Rab11-positive recycling and endosomes under serum starvation[26, 45]. The EHD1
protein converts these DAVs to form larger ciliary vesicles, which elongate through
continuous fusion with Rab8-positive vesicles to produce the primary cilium
membrane[25, 30, 41]. Therefore, Rab11-positive recycling endosomes, as important
transporters, are necessary for primary cilium formation and centriole-basal-body
transition in the early phase. Our study found that CRB3 could bind to some members
of Rab small GTPase family, significantly participating in Golgi vesicle transport and

vesicle organization pathways through mass spectrometry identification. And coexpression and co-IP results showed that CRB3 interacted with Rab11 via the extracellular domain, which mediated the intracellular trafficking of CRB3 within Rab11-positive endosomes at different cell densities.

Unlike motile cilia, primary cilia have a 9+0 structure, consisting of only nine outer doublet microtubules without a central pair. Similar to microtubule assembly at microtubule□organizing centers (MTOCs) during mitosis, γ-tubulin, which is a part of γTuRC, is also concentrated in the basal body of the cilium in interphase[46]. γ□tubulin complexes have been shown to form microtubule templates and regulate microtubule nucleation through longitudinal contacts with α□tubulin and β□tubulin[27]. The γTuRC consists of multiple copies of γTuSC, which is composed of two copies of γ□tubulin and one each of GCP2 and GCP3. Then, GCP2 and GCP3 interact with GCP4, GCP5 and GCP6 to form γTuRC[47]. Our experiments showed that CRB3 knockdown did not affect the expression of the γTuRC-specific proteins GCP3, GCP6 and γ□tubulin, but inhibited the interaction between GCP6 and GCP3, leading to destabilization of γTuRC. Further analysis revealed that significant colocalization was observed between GCP6- and Rab11-positive vesicles at low cell density, while at high cell density, GCP6 accumulated at the centrosome with GCP3. CRB3 knockdown resulted in the disappearance of this accumulated GCP6. These results indicated that GCP6/Rab11 trafficking vesicles must be transported to the basal body for ciliogenesis. While it is still unclear how CRB3 interferes with γTuRC assembly during primary cilium formation, we hypothesize that Rab11-positive endosomes mediate the intracellular trafficking of CRB3 and that CRB3, as a navigator interacting with CEP290, can navigate GCP6/Rab11 trafficking vesicles to the transition zone for γTuRC assembly during primary cilium formation.

CEP290 is a centriolar or ciliary protein that localizes to the centrosomes in dividing cells, the distal mother centriole in quiescent cells, and the transition zone in the primary cilium[48]. Mutations in *CEP290* are associated with various diseases, such as the devastating blinding disease Leber's Congenital Amaurosis (LCA), nephronophthisis, Senior Løken syndrome (SLS), Joubert syndrome (JS),

Bardet-Biedl syndrome (BBS), and lethal Meckel-Gruber syndrome (MGS)[49]. In

particular, the phenotypes of CEP290-associated ciliopathies are very similar to those

in our *Crb3* knockout mouse model, with lethality and ocular and renal abnormalities.

CEP290 knockdown significantly decreased the number of cells with primary cilia,

and CEP290 knockout mice showed mislocalization of ciliary proteins in the

retinas[50, 51]. Additionally, depletion of CEP290 disrupts protein trafficking to

centrosomes and affects the recruitment of the BBSome and vesicular trafficking in

ciliary assembly[52, 53]. Although CEP290 serves as a hub to recruit many ciliary

proteins, it has not been reported to be involved in polarized vesicle trafficking or

localization. Our study verified that CEP290 could interact with the polarity protein

CRB3. CRB3 knockdown disrupts the interaction between CEP290 and

Rab11-positive endosomes. Thus, these results validate our hypothesis. Although

CRB3 navigates GCP6/Rab11 trafficking vesicles to CEP290, we further need to

detect the region of CRB3 interacting with CEP290 and what other cargos are

transported by CRB3/Rab11 trafficking vesicles in ciliary assembly.

In addition, our results showed that CRB3 knockdown failed to activate the Hh

signaling pathway with SAG in MCF10A cells. Under untreated conditions, PTCH1

located at the ciliary membrane inhibits SMO, and Suppressor of Fused (SuFu)

sequesters Gli transcription factors, leading to their degradation. SAG can activate the

Hh signaling pathway, resulting in dissociation into the nucleus to regulate the

expression of downstream target genes[54]. The Hh signaling pathway mainly

regulates cellular growth and differentiation and is abnormal in different types of

tumors. Some studies indicate that cilia are double-sided, promoting and preventing

cancer development through the Hh signaling pathway *in vivo*. It is thought that cilia

deletion could activate SMO to inhibit tumor growth, while promoting carcinogenesis

was induced by activated Gli2[55]. Although SAG cannot activate the Hh signaling

pathway in CRB3-depleted cells, further research is needed to determine the role of

CRB3 in the carcinogenic process through the ciliary Hh signaling pathway.

Additionally, another canonical Wnt signaling pathway was altered after CRB3

knockdown. The Wnt signaling pathway is an important cascade regulating

development and stemness, and aberrant Wnt signaling has been reported in many cancer entities, especially colorectal cancer[56]. In our study, CRB3 downregulated the expression of β-catenin. Thus, CRB3 can affect cilium-related signaling pathways in tumorigenesis.

In summary, our study provides novel evidence for the role of the apical polarity protein CRB3 in regulating viability, mammary and ocular development, and primary cilium formation in germline and conditional knockout mice. Deletion of CRB3 caused irregular lumen formation and ductal epithelial hyperplasia in mammary epithelial-specific knockout mice. Further study uncovered a new function of CRB3b in interacting with Rab11, navigating GCP6/Rab11 trafficking vesicles to CEP290 for γTuRC assembly during ciliogenesis. In addition, CRB3 also participates in regulating cilium-related Hh and Wnt signaling pathways in tumorigenesis. Understanding these polarity protein-mediated vesicle trafficking mechanisms will shed new light on luminal development and tumorigenesis.

Methods

Generation of floxed *Crb3* mice. All animal experiments were verified and approved by the Committee of Institutional Animal Care and Use of Xi'an Jiaotong University. Heterozygous $Crb3^{wt/fl}$ mice (C57BL/6J) were generated using Cas9/CRISPR-mediated genome editing (Cyagen Biosciences, Guangzhou, China). The gRNA to *Crb3* and Cas9 mRNA was coinjected into fertilized mouse eggs to generate targeted conditional knockout offspring. The *Crb3* gRNA sequences were gRNA1, 5'-GGCTGGGTCCACACCTACGGAGG-3'; gRNA2, 5'-ACCCACAAAGCCACGCCA GTGGG-3'. Mouse genotyping was screened using PCR with the primers F1, 5'-ACATAAGGCCTTCCGTTAAGCTG-3'; R1, 5'-GTGGATTCGGACCAGTCTGA-3'. $Crb3^{wt/fl}$ mice were intercrossed with MMTV-*Cre* mice or CAG-*Cre* mice to generate tissue-specific *Crb3* knockout mice. MMTV-*Cre* mice (FVB), CAG-*Cre* mice (C57BL/6N), and MMTV-PyMT mice (FVB) were purchased from Cyagen Biosciences. All mice were bred in the specific

pathogen-free (SPF) animal houses of the Laboratory Animal Center of Xi'an
Jiaotong University.

Mice were genotyped using PCR and DNA gel electrophoresis. Genomic DNA from
the mouse tail was extracted using the Mouse Direct PCR Kit (Bimake, TX, USA,
#B40015) according to the manufacturer's instructions. The genotyping primers were
F3, 5'-TTGAGAGTCTTAAGCAGTCAGGG-3'; R5, 5'- AACCTTTCCCAGGAGTA
TGTGAC-3'. PCR results showed that $Crb3^{wt/wt}$ was one band with 163 bp, $Crb3^{wt/fl}$
was two bands with 228 bp and 163 bp, and $Crb3^{fl/fl}$ was one band with 228 bp. The
primers for identifying *Cre* alleles were F,
5'-TTGAGAGTCTTAAGCAGTCAGGG-3'; R,
5'-TTACCACTCCCAGCAAGACAC-3', and amplification with one 247 bp band.
The reaction conditions of *Crb3* and *Cre* PCR were as follows: 94°C for 5 min, 94°C
for 20 s, 60°C for 30 s, and 72°C for 20 s for 35 cycles.

Mouse mammary gland whole mount analysis. After euthanizing the mice, the
inguinal mammary gland tissues were removed intact and fully spread onto slides.
The tissues were rapidly fixed in Carnoy's fixative for 2 h at room temperature.
Subsequently, the tissues were washed in 70% ethanol solution for 15 min at room
temperature, and in 50% and 30% ethanol and distilled water for 5 min each. Staining
was performed in carmine alum solution at 4°C overnight. After washing with a 70%
ethanol solution, the tissues were washed again with 70%, 95%, and 100% ethanol for
15 min each. Tissues were cleared in xylene overnight at room temperature and then
mounted with neutral balsam. Images were obtained using a microscope (Leica DMi8,
Wetzlar, Germany).

Cell culture and transfection. The MCF10A, MCF7, T47D, MDA-MB-231,
HCC1806, T47D, and MDA-MB-453 cell lines were purchased from Shanghai
Institute of Biochemistry and Cell Biology (Chinese Academy of Sciences, Shanghai,
China). MCF10A cells were routinely grown in DMEM/F12 (1:1) media (HyClone,
UT, USA) supplemented with 5% horse serum, 20 ng/ml human EGF, 10 μg/ml
insulin, and 0.5 μg/ml hydrocortisone. MCF7, MDA-MB-231 and MDA-MB-453
cells were cultured in high glucose DMEM media (HyClone, UT, USA) supplemented

with 10% FBS (HyClone, UT, USA, #SH30084.03). T47D and HCC1806 cells were grown in RPMI-1640 media (HyClone, UT, USA) supplemented with 10% FBS. All cell lines were incubated in 5% CO_2 at 37°C.

siRNAs were purchased from GenePharma company (Shanghai, China). Cultured cells were transfected with siRNAs or plasmids using Lipofectamine 2000 (Invitrogen, CA, USA, #11668-019) following the manufacturer's protocol. The negative control shRNA (NC) and shRNA against CRB3 (both isoforms CRB3a and CRB3b) were packaged into lentivirus as previously described[5, 6].

DNA constructs and stable cell lines. CRB3b was PCR amplified from RNA extracted from MCF10A cells and cloned into the pCMV-Blank vector (Beyotime, Shanghai, China, D2602) using T4 ligase (NEB, MA, USA). Then, the GFP sequence was inserted between the extracellular and transmembrane CRB3 domains to generate the pCMV-CRB3-GFP vector, as described previously[31]. The full-length CRB3 (1-120), CRB3 (1-116), CRB3 (1-83), CRB3 (1-58), and CRB3 (1-26) were amplified by PCR from the pCMV-CRB3-GFP vector using specific primers to create EcoR Ⅰ and Xho Ⅰ restriction sites, and then cloned into the pCMV-Blank vector. The full-length was also cloned into the lentiviral vector pLVX-TetOne-Puro (Clontech, Takara, #631849). The plasmid pECMV-3×FLAG-RAB11A was purchased from SinoBiological (Beijing, China). The plasmids of pECMV-3×FLAG-RAB11A[S20V]/[S25N]/[Q70L] mutants were constructed using the *Fast* Mutagenesis System (Transgen, Beijing, China, FM111-01).

Lentiviruses were produced in HEK293T cells. CRB3-GFP in MCF10A and MCF7 cells was generated using a lentivirus expression system. Stable cell lines were screened using puromycin after lentiviral infection for 72 h.

3D morphogenesis. To initiate 3D morphogenesis, growth factor-reduced Matrigel (Corning, USA, #354230) was added to a four-well chamber slide system (Corning, USA, #177437) and then incubated at 37°C for solidification. MCF10A cells were plated into this chamber slide and cultured in DMEM/F12 (1:1) media (HyClone, UT, USA) supplemented with 2% horse serum, 5 ng/ml human EGF, 10 μg/ml insulin, 0.5 μg/ml hydrocortisone, and 2.5% Matrigel. 3D morphogenesis was photographed using

a microscope (Leica DMi8, Wetzlar, Germany) at different time points. On day 14 after cell culture, 3D MCF10A cells were fixed and stained.

Immunohistochemistry. The paraffin-embedded sections of samples were baked, deparaffinized in xylene, and sequentially rehydrated in gradient concentrations of ethanol, sequentially. Then, antigen retrieval was performed in Tris-EDTA buffer (pH 9.0) heated to 95°C for 20 min. Endogenous peroxidase activity was reduced in 3% hydrogen peroxide for 10 min at room temperature. Then, 5% goat plasma was used for blocking at 37°C for 30 min, and the sections were incubated with specific primary antibodies overnight at 4°C. Primary antibodies used were Ki67 (Abcam, UK, ab279653), phospho-histone H3 (Abcam, UK, ab267372), cleaved caspase 3 (CST, USA, #9661), GLI1 (CST, USA, #3538), and β-catenin (CST, USA, #8480). The following steps used biotin-streptavidin HRP detection kits (ZSGB-BIO, China, SP-9001, SP-9002) according to the manufacturer's instructions. The sections were stained with DAB (ZSGB-BIO, China, ZLI-9017), counterstained with hematoxylin, dehydrated in gradient concentrations of ethanol, cleared in xylene, and mounted with neutral balsam, sequentially. Images were obtained using a slide scanner (Leica SCN400, Wetzlar, Germany).

Immunofluorescence. Cells were plated on coverslips and fixed in 4% paraformaldehyde for 30 min. Then, the membranes were permeabilized using 0.2% Triton X-100. The cells were blocked with 5% BSA solution for 1 h at room temperature. Cells were incubated with specific primary antibodies overnight at 4°C. Caspase 3 (CST, USA, #9662S), α-tubulin (CST, USA, #3873), acetylated tubulin (Proteintech, USA, #662001-IG), γ-tubulin (Proteintech, USA, #15176-1-AP), pericentrin (Abcam, UK, ab4448), CRB3 (Sigma, USA, HPA013835), EEA1 (BD, USA, #610456), Rab11 (BD, USA, #610823), and GCP6 (Abcam, UK, ab95172) were used as primary antibodies. The secondary antibodies were Alexa Fluor 488-labeled or 594-labeled (Invitrogen, USA, A32731, A32744). DPIA (5 μg/ml) was used for DNA staining, and images were captured using a confocal microscope (Leica SP5 □, Wetzlar, Germany).

MEFs isolation and maintenance. The embryos were harvested from

Crb3^{wt/fl};CAG-*Cre* mice crosses at E13.5, and the head, tail, and internal organs were

removed. The tail was used for genotyping, and DNA was extracted from it. Then, the

body was minced and digested with trypsin in a 37°C incubator for 30 min. The

resulting cells were resuspended in MEF media (DME, 10% FBS, 1%

penicillin/streptomycin) and plated onto 10 cm cell culture dishes with one embryo in

each. They were subsequently passaged 1:3 with MEF media to expand the culture.

Real-time PCR assay. Total RNA from cell lines was isolated using TRIzol reagent

(Invitrogen, USA, #15596026), and 5 μg RNA was converted to cDNA with the

RevertAid first strand cDNA synthesis kit (Thermo, USA, K1622) according to the

manufacturer's instructions. Real-time PCR was prepared with SYBR qPCR Premix

(Takara, Japan, RR420L), and then performed with a real-time PCR detection

instrument (Bio–Rad CFX96, USA). The primers used for real-time PCR were

purchased from Tsingke Biotech (Beijing, China), and the sequences are listed in

Table 1. The mRNA expression was normalized to *GAPDH*, and fold changes were

calculated using the ΔΔCt method.

Immunoprecipitation and immunoblotting. Cultured cells were lysed in RIPA

buffer supplemented with protease inhibitors (Roche, NJ, USA). Then, the cell lysates

were subjected to SDS–PAGE separation and were transferred to PVDF membranes.

The membranes were subjected to immunoblot assays using antibodies against CRB3

(Santa Cruz, USA, sc-292449, both isoforms CRB3a and CRB3b can be detected),

GCP2 (Santa Cruz, USA, sc-377117), GCP3 (Santa Cruz, USA, sc-373758), GCP4

(Santa Cruz, USA, sc-271876), GCP5 (Santa Cruz, USA, sc-365837), GCP6 (Abcam,

UK, ab95172), γ-tubulin (Proteintech, USA, #66320-1-Ig), Rab11 (Abcam, UK,

ab128913), GFP (Roche, USA, #11814460001), Flag (Sigma, USA, F7425), CEP290

(Santa Cruz, USA, sc-390637), GSK3-β (CST, USA, #12456), β-catenin (CST, USA,

#8480), GAPDH (Proteintech, USA, #HRP-6004), and β-actin (Proteintech, USA,

#HRP-60008). HRP-conjugated secondary antibodies (CST, USA, #7074, #7076)

were then incubated at room temperature for 1 h in the dark. Final detection was

achieved by using ECL Plus (Millipore, Germany, WBULS0500). Proteins for the

immunoprecipitation assay were lysed with NP40 buffer. The cell lysates were

detected by a Dynabeads protein G immunoprecipitation kit (Invitrogen, USA,

#10007D) following the manufacturer's instructions. The elution protein complexes

were subjected to an immunoblot assay.

Protein identification and bioinformatics analyses. LC-MS/MS analysis was

conducted by PTM Bio (Zhejiang, China). Q Exactive^{TM} Plus (Thermo, MA, USA)

was used for tandem mass spectrometry data analysis. The Blast2GO program against

the UniProt database was used to perform functional annotation and classification of

the identified proteins. Subsequently, DAVID tools were used to perform pathway

enrichment analysis.

Scanning electron microscope (SEM) observations. MCF10A cells with

CRB3-GFP and CRB3 knockdown were plated on culture slides, and doxycycline

(Dox) was added to induce the expression of CRB3-GFP. After the cells were fully

confluent, the slides were fixed with 4% paraformaldehyde at 4°C overnight, followed

by 1% osmium tetroxide at 4°C for 1 h. Dehydration was performed at room

temperature, and the samples were dried using the critical point drying method.

Images were obtained by using a scanning electron microscope (Hitachi TM1000,

Japan).

Clinical data. Patient data, breast cancer tissues, and adjacent para-cancerous tissues

were collected from the First Affiliated Hospital of Xi'an Jiaotong University

(Shaanxi, China). All patients signed informed consent forms before surgery. This

research was authorized by the Ethics Committee of the First Affiliated Hospital of

Xi'an Jiaotong University and conducted in conformity with the Declaration of

Helsinki.

Statistics. Statistical analysis was performed using SPSS statistics 23.0 for Windows

(IBM, Armonk, USA). All experiments were repeated at least three times. The values

are expressed as the mean ± standard deviation (SD). Unpaired Student's t test was

used to compare the differences between two groups. One-way ANOVA followed by

Dunnett's multiple comparisons test was used for multiple comparisons. The χ^2 test

was used to assess the significance of the observed frequencies. $P<0.05$ was

considered an indicator of statistical significance.

Acknowledgments

This work was supported by grants from the National Natural Science Foundation of

China (No. 81872272 and 82173023), Innovation Capability Support Program of

Shaanxi (No. 2020TD-046) and Clinical Research Award of the First Affiliated

Hospital of Xi'an Jiaotong University (No. XJTU1AF-CRF-2017-007). We are

grateful to Prof. Ceshi Chen (Kunming Institute of Zoology, China) and Prof.

Yongping Shao (Xi'an Jiaotong University, China) for their guidance in this research.

We also thank Qi Tian, Lizhe Zhu and Yan Zhou for their help with the experiments.

Conflict of interest

The authors declare no conflicts of interest.

Author contributions

P.L. designed the research; B.W., Z.L., T.T, Y.J., Y.S., R.W., H.C., J.L. and P.L.

performed the experimental work, and M.Z., X.G., J.L. and S.S. analyzed the data;

B.W., Z.L., Y.R. and P.L. wrote the original manuscript. P.L. conceived the study and

critically revised the manuscript. All authors read and approved the final manuscript.

Data availability

The datasets used and/or analyzed during the current study are available from the

corresponding author on reasonable request.

References
715

1. Wodarz A, Näthke I. Cell polarity in development and cancer. Nature cell biology.

2007;9(9):1016-24. Epub 2007/09/01. doi: 10.1038/ncb433. PubMed PMID: 17762893.

2. McCaffrey LM, Macara IG. Epithelial organization, cell polarity and tumorigenesis. Trends

in cell biology. 2011;21(12):727-35. Epub 2011/07/26. doi: 10.1016/j.tcb.2011.06.005.

PubMed PMID: 21782440.

3. Etienne-Manneville S. Polarity proteins in migration and invasion. Oncogene.

2008;27(55):6970-80. Epub 2008/11/26. doi: 10.1038/onc.2008.347. PubMed PMID:

19029938.

4. Elsum I, Yates L, Humbert PO, Richardson HE. The Scribble-Dlg-Lgl polarity module in

development and cancer: from flies to man. Essays in biochemistry. 2012;53:141-68. Epub

2012/08/30. doi: 10.1042/bse0530141. PubMed PMID: 22928514.

5. Li P, Wang Y, Mao X, Jiang Y, Liu J, Li J, et al. CRB3 downregulation confers breast

cancer stem cell traits through TAZ/β-catenin. Oncogenesis. 2017;6(4):e322. Epub

2017/04/25. doi: 10.1038/oncsis.2017.24. PubMed PMID: 28436991; PubMed Central PMCID:

PMCPMC5520500.

6. Mao X, Li P, Wang Y, Liang Z, Liu J, Li J, et al. CRB3 regulates contact inhibition by

activating the Hippo pathway in mammary epithelial cells. Cell death & disease.

2017;8(1):e2546. Epub 2017/01/13. doi: 10.1038/cddis.2016.478. PubMed PMID: 28079891;

PubMed Central PMCID: PMCPMC5386381.

7. Mao X, Li P, Ren Y, Li J, Zhou C, Yang J, et al. Cell polarity protein CRB3 is an

independent favorable prognostic factor for clear cell renal cell carcinoma. International journal

of   oncology.   2015;46(2):657-66.   doi: 10.3892/ijo.2014.2763.   PubMed   PMID:

WOS:000347671200025.

8.   Karp CM, Tan TT, Mathew R, Nelson D, Mukherjee C, Degenhardt K, et al. Role of the

polarity determinant crumbs in suppressing mammalian epithelial tumor progression. Cancer

research. 2008;68(11):4105-15. Epub 2008/06/04. doi: 10.1158/0008-5472.Can-07-6814.

PubMed PMID: 18519669; PubMed Central PMCID: PMCPMC2696887.

9.   Sánchez I, Dynlacht BD. Cilium assembly and disassembly. Nature cell biology.

2016;18(7):711-7. Epub 2016/06/29. doi: 10.1038/ncb3370. PubMed PMID: 27350441;

PubMed Central PMCID: PMCPMC5079433.

10.   Reiter JF, Leroux MR. Genes and molecular pathways underpinning ciliopathies. Nature

reviews   Molecular   cell   biology.   2017;18(9):533-47.   Epub   2017/07/13.   doi:

10.1038/nrm.2017.60.   PubMed   PMID: 28698599;   PubMed   Central   PMCID:

PMCPMC5851292.

11.   Kim J, Dabiri S, Seeley ES. Primary cilium depletion typifies cutaneous melanoma in situ

and   malignant   melanoma.   PloS   one.   2011;6(11):e27410.   Epub   2011/11/19.   doi:

10.1371/journal.pone.0027410.   PubMed   PMID: 22096570;   PubMed   Central   PMCID:

PMCPMC3214062.

12.   Menzl I, Lebeau L, Pandey R, Hassounah NB, Li FW, Nagle R, et al. Loss of primary cilia

occurs   early   in   breast   cancer   development.   Cilia.   2014;3:7.   Epub   2014/07/06.   doi:

10.1186/2046-2530-3-7.   PubMed   PMID: 24987519;   PubMed   Central   PMCID:

PMCPMC4076761.

13.   Basten SG, Willekers S, Vermaat JS, Slaats GG, Voest EE, van Diest PJ, et al. Reduced

cilia frequencies in human renal cell carcinomas versus neighboring parenchymal tissue. Cilia.

2013;2(1):2. Epub 2013/02/02. doi: 10.1186/2046-2530-2-2. PubMed PMID: 23369289;

PubMed Central PMCID: PMCPMC3564780.

14. Seeley ES, Carrière C, Goetze T, Longnecker DS, Korc M. Pancreatic cancer and

precursor pancreatic intraepithelial neoplasia lesions are devoid of primary cilia. Cancer

research. 2009;69(2):422-30. Epub 2009/01/17. doi: 10.1158/0008-5472.Can-08-1290.

PubMed PMID: 19147554; PubMed Central PMCID: PMCPMC2629528.

15. Hassounah NB, Nagle R, Saboda K, Roe DJ, Dalkin BL, McDermott KM. Primary cilia are

lost in preinvasive and invasive prostate cancer. PloS one. 2013;8(7):e68521. Epub

2013/07/12. doi: 10.1371/journal.pone.0068521. PubMed PMID: 23844214; PubMed Central

PMCID: PMCPMC3699526.

16. Bazellières E, Aksenova V, Barthélémy-Requin M, Massey-Harroche D, Le Bivic A. Role

of the Crumbs proteins in ciliogenesis, cell migration and actin organization. Seminars in cell &

developmental biology. 2018;81:13-20. Epub 2017/10/24. doi: 10.1016/j.semcdb.2017.10.018.

PubMed PMID: 29056580.

17. Fan S, Fogg V, Wang Q, Chen XW, Liu CJ, Margolis B. A novel Crumbs3 isoform

regulates cell division and ciliogenesis via importin beta interactions. The Journal of cell

biology. 2007;178(3):387-98. Epub 2007/07/25. doi: 10.1083/jcb.200609096. PubMed PMID:

17646395; PubMed Central PMCID: PMCPMC2064851.

18. Fan S, Hurd TW, Liu CJ, Straight SW, Weimbs T, Hurd EA, et al. Polarity proteins control

ciliogenesis via kinesin motor interactions. Current biology : CB. 2004;14(16):1451-61. Epub

2004/08/25. doi: 10.1016/j.cub.2004.08.025. PubMed PMID: 15324661.

19.   Whiteman EL, Fan S, Harder JL, Walton KD, Liu CJ, Soofi A, et al. Crumbs3 is essential

for proper epithelial development and viability. Molecular and cellular biology.

2014;34(1):43-56. Epub 2013/10/30. doi: 10.1128/mcb.00999-13. PubMed PMID: 24164893;

PubMed Central PMCID: PMCPMC3911272.

20.   Ye X, Tam WL, Shibue T, Kaygusuz Y, Reinhardt F, Ng Eaton E, et al. Distinct EMT

programs control normal mammary stem cells and tumour-initiating cells. Nature.

2015;525(7568):256-60. Epub 2015/09/04. doi: 10.1038/nature14897. PubMed PMID:

26331542; PubMed Central PMCID: PMCPMC4764075.

21.   Soule HD, Maloney TM, Wolman SR, Peterson WD, Jr., Brenz R, McGrath CM, et al.

Isolation and characterization of a spontaneously immortalized human breast epithelial cell

line, MCF-10. Cancer research. 1990;50(18):6075-86. Epub 1990/09/25. PubMed PMID:

1975513.

22.   Halsne R, Tandberg JI, Lobert VH, Østby GC, Thoen E, Ropstad E, et al. Effects of

perfluorinated alkyl acids on cellular responses of MCF-10A mammary epithelial cells in

monolayers and on acini formation in vitro. Toxicol Lett. 2016;259:95-107. Epub 2016/08/12.

doi: 10.1016/j.toxlet.2016.08.004. PubMed PMID: 27511595.

23.   Jonassen JA, San Agustin J, Follit JA, Pazour GJ. Deletion of IFT20 in the mouse kidney

causes misorientation of the mitotic spindle and cystic kidney disease. The Journal of cell

biology. 2008;183(3):377-84. Epub 2008/11/05. doi: 10.1083/jcb.200808137. PubMed PMID:

18981227; PubMed Central PMCID: PMCPMC2575779.

24.   Martin-Belmonte F, Perez-Moreno M. Epithelial cell polarity, stem cells and cancer.

Nature reviews Cancer. 2011;12(1):23-38. Epub 2011/12/16. doi: 10.1038/nrc3169. PubMed

PMID: 22169974.

25. Lu Q, Insinna C, Ott C, Stauffer J, Pintado PA, Rahajeng J, et al. Early steps in primary

cilium assembly require EHD1/EHD3-dependent ciliary vesicle formation. Nature cell biology.

2015;17(3):228-40. Epub 2015/02/17. doi: 10.1038/ncb3109. PubMed PMID: 25686250;

PubMed Central PMCID: PMCPMC4344897.

26. Westlake CJ, Baye LM, Nachury MV, Wright KJ, Ervin KE, Phu L, et al. Primary cilia

membrane assembly is initiated by Rab11 and transport protein particle II (TRAPPII)

complex-dependent trafficking of Rabin8 to the centrosome. Proceedings of the National

Academy of Sciences of the United States of America. 2011;108(7):2759-64. Epub 2011/01/29.

doi: 10.1073/pnas.1018823108. PubMed PMID: 21273506; PubMed Central PMCID:

PMCPMC3041065 R.H.S., and P.K.J. are employees of Genentech, Inc.

27. Kollman JM, Merdes A, Mourey L, Agard DA. Microtubule nucleation by γ-tubulin

complexes. Nature reviews Molecular cell biology. 2011;12(11):709-21. Epub 2011/10/14. doi:

10.1038/nrm3209. PubMed PMID: 21993292; PubMed Central PMCID: PMCPMC7183383.

28. Liu P, Zupa E, Neuner A, Böhler A, Loerke J, Flemming D, et al. Insights into the

assembly and activation of the microtubule nucleator γ-TuRC. Nature. 2020;578(7795):467-71.

Epub 2019/12/20. doi: 10.1038/s41586-019-1896-6. PubMed PMID: 31856152.

29. Farache D, Jauneau A, Chemin C, Chartrain M, Rémy MH, Merdes A, et al. Functional

Analysis of γ-Tubulin Complex Proteins Indicates Specific Lateral Association via Their

N-terminal Domains. The Journal of biological chemistry. 2016;291(44):23112-25. Epub

2016/10/30. doi: 10.1074/jbc.M116.744862. PubMed PMID: 27660388; PubMed Central

PMCID: PMCPMC5087730.

30. Knödler A, Feng S, Zhang J, Zhang X, Das A, Peränen J, et al. Coordination of Rab8 and

Rab11 in primary ciliogenesis. Proceedings of the National Academy of Sciences of the United

States of America. 2010;107(14):6346-51. Epub 2010/03/24. doi: 10.1073/pnas.1002401107.

PubMed PMID: 20308558; PubMed Central PMCID: PMCPMC2851980.

31. Djuric I, Siebrasse JP, Schulze U, Granado D, Schlüter MA, Kubitscheck U, et al. The

C-terminal domain controls the mobility of Crumbs 3 isoforms. Biochimica et biophysica acta.

2016;1863(6 Pt A):1208-17. Epub 2016/03/16. doi: 10.1016/j.bbamcr.2016.03.008. PubMed

PMID: 26975581.

32. Craige B, Tsao CC, Diener DR, Hou Y, Lechtreck KF, Rosenbaum JL, et al. CEP290

tethers flagellar transition zone microtubules to the membrane and regulates flagellar protein

content. The Journal of cell biology. 2010;190(5):927-40. Epub 2010/09/08. doi:

10.1083/jcb.201006105. PubMed PMID: 20819941; PubMed Central PMCID:

PMCPMC2935561.

33. Casenghi M, Meraldi P, Weinhart U, Duncan PI, Körner R, Nigg EA. Polo-like kinase 1

regulates Nlp, a centrosome protein involved in microtubule nucleation. Developmental cell.

2003;5(1):113-25. Epub 2003/07/11. doi: 10.1016/s1534-5807(03)00193-x. PubMed PMID:

12852856.

34. Li P, Feng C, Chen H, Jiang Y, Cao F, Liu J, et al. Elevated CRB3 expression suppresses

breast cancer stemness by inhibiting β-catenin signalling to restore tamoxifen sensitivity.

Journal of cellular and molecular medicine. 2018;22(7):3423-33. Epub 2018/03/31. doi:

10.1111/jcmm.13619. PubMed PMID: 29602199; PubMed Central PMCID:

PMCPMC6010813.

35. Thiery JP, Acloque H, Huang RY, Nieto MA. Epithelial-mesenchymal transitions in

development and disease. Cell. 2009;139(5):871-90. Epub 2009/12/01. doi:

10.1016/j.cell.2009.11.007. PubMed PMID: 19945376.

36. Varelas X, Samavarchi-Tehrani P, Narimatsu M, Weiss A, Cockburn K, Larsen BG, et al.

The Crumbs complex couples cell density sensing to Hippo-dependent control of the

TGF-β-SMAD pathway. Developmental cell. 2010;19(6):831-44. Epub 2010/12/15. doi:

10.1016/j.devcel.2010.11.012. PubMed PMID: 21145499.

37. Lin W-H, Asmann YW, Anastasiadis PZ. Expression of polarity genes in human cancer.

Cancer informatics. 2015;14:CIN. S18964.

38. Charrier LE, Loie E, Laprise P. Mouse Crumbs3 sustains epithelial tissue morphogenesis

in vivo. Scientific reports. 2015;5:17699. Epub 2015/12/04. doi: 10.1038/srep17699. PubMed

PMID: 26631503; PubMed Central PMCID: PMCPMC4668553.

39. Herranz-Martín S, Jimeno D, Paniagua AE, Velasco A, Lara JM, Aijón J, et al.

Immunocytochemical evidence of the localization of the Crumbs homologue 3 protein (CRB3)

in the developing and mature mouse retina. PloS one. 2012;7(11):e50511. Epub 2012/12/12.

doi: 10.1371/journal.pone.0050511. PubMed PMID: 23226298; PubMed Central PMCID:

PMCPMC3511585.

40. Arya P, Rainey MA, Bhattacharyya S, Mohapatra BC, George M, Kuracha MR, et al. The

endocytic recycling regulatory protein EHD1 Is required for ocular lens development.

Developmental biology. 2015;408(1):41-55. Epub 2015/10/13. doi:

10.1016/j.ydbio.2015.10.005. PubMed PMID: 26455409; PubMed Central PMCID:

PMCPMC4688215.

41. Rainey MA, George M, Ying G, Akakura R, Burgess DJ, Siefker E, et al. The endocytic

recycling regulator EHD1 is essential for spermatogenesis and male fertility in mice. BMC Dev

Biol. 2010;10:37. Epub 2010/04/03. doi: 10.1186/1471-213x-10-37. PubMed PMID: 20359371;

PubMed Central PMCID: PMCPMC2856533.

42. Kobayashi T, Dynlacht BD. Regulating the transition from centriole to basal body. The

Journal of cell biology. 2011;193(3):435-44. Epub 2011/05/04. doi: 10.1083/jcb.201101005.

PubMed PMID: 21536747; PubMed Central PMCID: PMCPMC3087006.

43. Kobayashi T, Kim S, Lin YC, Inoue T, Dynlacht BD. The CP110-interacting proteins

Talpid3 and Cep290 play overlapping and distinct roles in cilia assembly. The Journal of cell

biology. 2014;204(2):215-29. Epub 2014/01/15. doi: 10.1083/jcb.201304153. PubMed PMID:

24421332; PubMed Central PMCID: PMCPMC3897186.

44. Schmidt KN, Kuhns S, Neuner A, Hub B, Zentgraf H, Pereira G. Cep164 mediates

vesicular docking to the mother centriole during early steps of ciliogenesis. The Journal of cell

biology. 2012;199(7):1083-101. Epub 2012/12/21. doi: 10.1083/jcb.201202126. PubMed

PMID: 23253480; PubMed Central PMCID: PMCPMC3529528.

45. Das A, Guo W. Rabs and the exocyst in ciliogenesis, tubulogenesis and beyond. Trends

in cell biology. 2011;21(7):383-6. Epub 2011/05/10. doi: 10.1016/j.tcb.2011.03.006. PubMed

PMID: 21550243; PubMed Central PMCID: PMCPMC3128673.

46. Guichard P, Laporte MH, Hamel V. The centriolar tubulin code. Seminars in cell &

developmental biology. 2021. Epub 2021/12/14. doi: 10.1016/j.semcdb.2021.12.001. PubMed

PMID: 34896019.

47. Liu P, Würtz M, Zupa E, Pfeffer S, Schiebel E. Microtubule nucleation: The waltz between

γ-tubulin ring complex and associated proteins. Current opinion in cell biology. 2021;68:124-31.

Epub 2020/11/16. doi: 10.1016/j.ceb.2020.10.004. PubMed PMID: 33190097.

48.   Chen HY, Kelley RA, Li T, Swaroop A. Primary cilia biogenesis and associated retinal

ciliopathies. Seminars in cell & developmental biology. 2021;110:70-88. Epub 2020/08/05. doi:

10.1016/j.semcdb.2020.07.013.   PubMed   PMID: 32747192;   PubMed   Central   PMCID:

PMCPMC7855621.

49.   Drivas TG, Bennett J. CEP290 and the primary cilium. Advances in experimental

medicine   and   biology.   2014;801:519-25.   Epub   2014/03/26.   doi:

10.1007/978-1-4614-3209-8_66. PubMed PMID: 24664739.

50.   Tsang WY, Bossard C, Khanna H, Peränen J, Swaroop A, Malhotra V, et al. CP110

suppresses primary cilia formation through its interaction with CEP290, a protein deficient in

human   ciliary   disease.   Developmental   cell.   2008;15(2):187-97.   Epub   2008/08/13.   doi:

10.1016/j.devcel.2008.07.004.   PubMed   PMID: 18694559;   PubMed   Central   PMCID:

PMCPMC3987787.

51.   Chang B, Khanna H, Hawes N, Jimeno D, He S, Lillo C, et al. In-frame deletion in a novel

centrosomal/ciliary protein CEP290/NPHP6 perturbs its interaction with RPGR and results in

early-onset   retinal   degeneration   in   the   rd16   mouse.   Human   molecular   genetics.

2006;15(11):1847-57. Epub 2006/04/25. doi: 10.1093/hmg/ddl107. PubMed PMID: 16632484;

PubMed Central PMCID: PMCPMC1592550.

52.   Kim J, Krishnaswami SR, Gleeson JG. CEP290 interacts with the centriolar satellite

component PCM-1 and is required for Rab8 localization to the primary cilium. Human

molecular   genetics.   2008;17(23):3796-805.   Epub   2008/09/06.   doi: 10.1093/hmg/ddn277.

PubMed PMID: 18772192; PubMed Central PMCID: PMCPMC2722899.

53. Stowe TR, Wilkinson CJ, Iqbal A, Stearns T. The centriolar satellite proteins Cep72 and

Cep290 interact and are required for recruitment of BBS proteins to the cilium. Molecular

biology of the cell. 2012;23(17):3322-35. Epub 2012/07/07. doi: 10.1091/mbc.E12-02-0134.

PubMed PMID: 22767577; PubMed Central PMCID: PMCPMC3431927.

54. Wheway G, Nazlamova L, Hancock JT. Signaling through the Primary Cilium. Frontiers in

cell and developmental biology. 2018;6:8. Epub 2018/02/24. doi: 10.3389/fcell.2018.00008.

PubMed PMID: 29473038; PubMed Central PMCID: PMCPMC5809511.

55. Wong SY, Seol AD, So PL, Ermilov AN, Bichakjian CK, Epstein EH, Jr., et al. Primary cilia

can both mediate and suppress Hedgehog pathway-dependent tumorigenesis. Nature

medicine. 2009;15(9):1055-61. Epub 2009/08/25. doi: 10.1038/nm.2011. PubMed PMID:

19701205; PubMed Central PMCID: PMCPMC2895420.

56. Zhan T, Rindtorff N, Boutros M. Wnt signaling in cancer. Oncogene. 2017;36(11):1461-73.

Epub 2016/09/13. doi: 10.1038/onc.2016.304. PubMed PMID: 27617575; PubMed Central

PMCID: PMCPMC5357762.

928
929

Figure 1. *Crb3* knockout mice exhibit smaller sizes and ocular abnormalities, and mammary epithelial cell-specific *Crb3* knockout leads to ductal epithelial hyperplasia and promotes tumorigenesis.

A, B. Representative whole bodies (A) and eyes (B) from littermate *Crb3*$^{fl/fl}$, *Crb3*$^{wt/fl}$;CAG-*Cre* and *Crb3*$^{fl/fl}$;CAG-*Cre* mice at 4 weeks old. C. Representative mammary whole mounts from littermate *Crb3*$^{fl/fl}$ and *Crb3*$^{fl/fl}$;MMTV-*Cre* mice at 8 weeks old with Carmine-alum staining. (scale bars, 200 μm) D, E. Quantification of the average number of TEBs and bifurcated TEBs in littermate *Crb3*$^{fl/fl}$ (n=10) and *Crb3*$^{fl/fl}$;MMTV-*Cre* (n=10) mice at 8 weeks old. F. Representative images of mammary glands in littermate *Crb3*$^{fl/fl}$ and *Crb3*$^{fl/fl}$;MMTV-*Cre* mice stained with H&E. (scale bars, 50 μm) G. Representative images of primary tumors stained with H&E in PyMT-WT and PyMT-cKO-*Crb3* mice at 9 weeks old. (scale bars, left 500 μm, right 50 μm) The magnified areas of boxed sections are shown in the right panels. Bars represent the means ± SD; Unpaired Student's *t* test, ***$P<0.001$.

Figure 2. CRB3 knockdown inhibits acinar formation of mammary epithelial cells in a 3D culture system.

A. Representative effect of CRB3 on acinar formation in the 3D culture system at days 3, 6, 9 and 14. (The magnified areas of the marked arrows are shown in the lower right corner) B. Quantification of the average number, diameter and aberration of acini (At least 3-4 random micrographs were analyzed for each condition in each of three independent experiments, n=10). C. Immunofluorescence showing apoptosis during lumen formation. Caspase 3 (red), α-tubulin (green), and DNA (blue). D. Quantification of the proportion of caspase 3 positive acini (At least 3-4 random micrographs were analyzed for each condition in each of three independent experiments). E. Immunofluorescence showing the mitotic spindle orientation during lumen formation. α-tubulin (green) and DNA (blue). F. Quantification of division angle (At least 3-4 acini were randomly examined for each condition in each of three independent experiments). G. Immunohistochemical analyses of Ki67, phospho-histone H3 and cleaved caspase 3 in primary tumors from PyMT-WT (n=3)

and PyMT-cKO-*Crb3* (n=3) mice at 9 weeks old. (Positive cells are marked by arrows; scale bars, 25 μm) H. Quantification of the number of Ki67, phospho-histone H3 and cleaved caspase 3 positive cells from IHC staining images. Bars represent means ± SD; Unpaired Student's *t* test, * *P*<0.5, ** *P*<0.01, ***P*<0.001.

Figure 3. CRB3 alters primary cilium formation in mammary cells, mammary ductal lumen and renal tubule from *Crb3*^fl/fl^;CAG-*Cre* mice.

A. C. Representative images of immunofluorescent staining of primary cilium formation with CRB3 knockdown in MCF10A cells and CRB3b conditional overexpression upon dox induction in MCF7 cells. Acetylated tubulin (red), γ-tubulin (green), and DNA (blue). (The primary cilium is marked by arrows; scale bars, 10 μm) B. D. Quantification of the proportion of cells with primary cilium formation and the length of the primary cilium (At least 3-4 random micrographs were analyzed for each condition in each of three independent experiments, n=10). E. Representative scanning electron microscope images of primary cilium formation with CRB3 knockdown and CRB3b conditional overexpression in MCF10A cells. (The primary cilium is marked by arrows; scale bars, 50 μm) F. Quantification of the proportion of cells with primary cilium formation (At least 3-4 random micrographs were analyzed for each condition in each of three independent experiments, n=10). G. H. Representative immunofluorescent staining of primary cilium formation in the mammary ductal lumen and renal tubule from *Crb3*^fl/fl^ (n=10) and *Crb3*^fl/fl^;CAG-*Cre* (n=10) mice, respectively. Acetylated tubulin (red), γ-tubulin (green), and DNA (blue). (scale bars, 25 μm) Bars represent the means ± SD; Unpaired Student's *t* test, *P*<0.05, ***P*<0.001.

Figure 4. CRB3 localizes to the basal body of the primary cilium.

A. Immunofluorescence showing the colocalization of exogenous CRB3b with centrosomes in MCF10A cells. Pericentrin, a marker of centrosome (red), CRB3-GFP (green), and DNA (blue). (Colocalization is marked by arrows; scale bars, 10 μm) B. Quantification of the proportion of cells with pericentrin and exogenous CRB3b colocalization (At least 3-4 random micrographs were analyzed for each condition in

each of three independent experiments, n=10). C. Another colocalization of endogenous CRB3 with the basal body in MCF10A cells. γ-tubulin is a marker of the centrosome and basal body of the primary cilium (green), CRB3 (red), and DNA (blue). (Colocalization is marked by arrows; scale bars, 10 μm) D. Corresponding fluorescence intensity profile across a section of the array, as indicated by the dashed white line in (C). E. Quantification of the number of γ-tubulin foci in MCF10A cells from IF staining images (At least 3-4 random micrographs were analyzed for each condition in each of three independent experiments, n=10). F. Double immunostaining displaying the colocalization of CRB3 with the primary cilium in MCF10A cells. Acetylated tubulin (green), CRB3 (red), and DNA (blue). (Colocalization is marked by arrows; scale bars, 10 μm) G. Fluorescence 3D reconstruction of CRB3 and primary cilium colocalization. Acetylated tubulin (red), CRB3 (green), and DNA (blue). Bars represent the means ± SD; Unpaired Student's *t* test, ***$P<0.001$.

Figure 5. CRB3 trafficking is mediated by Rab11-positive endosomes, and CRB3 knockdown destabilizes γTuRC assembly during ciliogenesis.

A. Pathway aggregation analysis of CRB3b binding proteins identified by mass spectrometry in MCF10A cells. B. Table of some Rab small GTPase family members and centriolar proteins identified as CRB3b binding proteins. C. Immunofluorescence showing the colocalization of CRB3 with EEA1-, CD63-, and Rab11-positive endosomes in MCF10A cells. EEA1, CD63, Rab11 (green), CRB3 (red), and DNA (blue). (scale bars, 10 μm) D. Quantification of the proportion of cells with these markers and CRB3 colocalization (n=10). E. Western blotting showing the levels of CRB3 in MCF10A cells treated with dynasore at different cell densities. F. The structure diagram of γTuRC (γ-tubulin ring complex). G. Immunoblot analysis of the effect of CRB3 on γTuRC molecules in MCF10A cells. H. Coimmunoprecipitation showing the interacting proteins with GCP6 in MCF10A cells with CRB3 knockdown. I. Representative images of immunofluorescent staining of GCP3 and GCP6 colocalization in MCF10A cells with the corresponding fluorescence intensity profile. GCP3 (red), GCP6 (green), and DNA (blue). (At least 3-4 random micrographs were

1021 analyzed for each condition in each of three independent experiments, scale bars, 10

1022 µm) J. The comparison of cytoplasmic extracts from MCF10A cells and cells with

1023 CRB3 knockdown after fractionation in sucrose gradients. The γTuSC sedimentation

1024 was mainly in fractions 3, and γTuRC sedimentation was mainly in fractions 6. Bars

1025 represent the means ± SD; Unpaired Student's *t* test, ***$P<0.001$.

1026

Figure 6. CRB3 interacts with Rab11.

A. Coimmunoprecipitation of CRB3 with Rab11, GCP6 and GCP3 in MCF10A cells.

B. Coimmunoprecipitation of Rab11 with CRB3, GCP6 and GCP3 in MCF10A cells.

C. Coimmunoprecipitation of Rab11 with GCP6 in control and CRB3 knockdown

MCF10A cells. D. Schematic diagram of CRB3b domains. E. Diagram truncations of

CRB3b-GFP fusion proteins with serial C-terminal deletions. F. Domain mapping of

CRB3b-GFP for Flag-Rab11a binding. Flag antibody co-IP of the full-length

CRB3b-GFP and truncations of CRB3b-GFP with Flag-Rab11a were cotransfected

into HEK293 cells for 48 h. Immunoblot analysis was performed using GFP and Flag

antibodies. G. Coimmunoprecipitation of Rab11 mutant variants with full-length

CRB3b-GFP. Flag antibody co-IP of the full-length CRB3b-GFP with

Flag-Rab11aWT, Flag-Rab11a[Q70L], Flag-Rab11a[S20V] and Flag-Rab11a[S25N]

were cotransfected into HEK293 cells for 48 h. Immunoblot analysis was performed

using GFP and Flag antibodies. Bars represent the means ± SD, and the experiments

were performed in triplicate; Unpaired Student's *t* test, *$P<0.05$, ***$P<0.001$.

1042

Figure 7. CRB3 navigates GCP6/Rab11 trafficking vesicles to the basal body of the

primary cilium.

A. Coimmunoprecipitation of exogenous CRB3b with Rab11, GCP6, GCP3 and

CEP290 in MCF10A cells. B. Coimmunoprecipitation of Rab11 with exogenous

CRB3b, GCP6, GCP3 and CEP290 in MCF10A cells. C. Coimmunoprecipitation of

Rab11 with GCP6 and CEP290 in control and CRB3 knockdown MCF10A cells. D.

Representative immunofluorescent images of GCP6 and γ-tubulin colocalization of

the basal body foci in MCF10A cells with CRB3 knockdown. γ-tubulin (green), GCP6

(red), and DNA (blue). (Colocalization is marked by arrows; scale bars, 25 μm) E.

Representative immunofluorescent images of Rab11 and γ-tubulin colocalization of

the basal body foci in MEF cells from $Crb3^{fl/fl}$ and $Crb3^{fl/fl}$;CAG-Cre mice. γ-tubulin

(green), Rab11 (red), and DNA (blue). Foci are marked by arrows; scale bars, 25 μm.

Bars represent the means ± SD; Unpaired Student's t test, ***P<0.001.

1056

Figure 8. Defects in CRB3 expression inhibit ciliary assembly in breast cancer tissues,

and activate the Wnt signaling pathway in mammary cells and PyMT mouse model.

A. Representative immunofluorescent images of CRB3 in breast cancer tissues (n=50).

CRB3 (green) and DNA (blue). (scale bars, 25 μm) B. Representative

immunofluorescent images of the primary cilium in breast cancer tissues (n=50).

Acetylated tubulin (red), γ-tubulin (green), and DNA (blue). (scale bars, 10 μm) C.

Quantification of the proportion of cells with primary cilium formation in breast

cancer tissues (n=50). D. Representative immunofluorescent images of CRB3 and

γ-tubulin colocalization in adjacent paracarcinoma tissues. CRB3 (green), γ-tubulin

(red), and DNA (blue). (scale bars, 25 μm) E. Quantification of the proportion of

MFC10A cells with SMO translocation after CRB3 knockdown (n=6). F. Real-time

quantitative PCR showing the relative mRNA expression of GLI1 upon SAG

treatment in CRB3-depleted MFC10A cells (n=6). G. Immunoblot analyses of the

effect of CRB3 on the molecules of the Wnt signaling pathway in mammary cells, and

the experiments were performed in triplicate. H. I. Immunohistochemical analyses of

GLI1 and β-catenin in primary tumors from PyMT-WT and PyMT-cKO-$Crb3$ mice at

9 weeks old, respectively. (scale bars, 25 μm) Bars represent the means ± SD;

Unpaired Student's t test, ***P<0.001.

1075

Figure 9. Schematic model of CRB3 regulating ciliary assembly.

Graphic summary of prominent phenotypes observed after CRB3 deletion. CRB3 is

localized on apical epithelial surfaces and participates in tight junction formation to

maintain contact inhibition and cell homeostasis in quiescence. Inside the cell,

Rab11-positive endosomes mediate the intracellular trafficking of CRB3, and CRB3

navigates GCP6/Rab11 trafficking vesicles to CEP290. Then, GCP6 is involved in

normal γTuRC assembly in ciliogenesis. In CRB3 deletion cells, the primary cilium

fails to assemble properly, and the Wnt signaling pathway is activated through

β-catenin upregulation and nuclear localization, but the Hh signaling pathway fails to

be activated. This cellular imbalance is disrupted, leading to tumorigenesis.

Abbreviations: γTuRC, γ-tubulin ring complex; TJ, tight junction; EE, early

endosome; LE, late endosome.

1088

Supplementary Figure 1. Generation of *Crb3* knockout mice using the Cre-loxP

system.

A. Schematic representation of the strategy for loxP sequence insertion and PCR

genotyping. The loxP sites (gray arrow) flanked either side of exon 3 (orange) in the

Crb3 gene. The F3&R5 and F4&R6 primer pair were used to differentiate the alleles.

B. Southern blot analysis of a positive ES cell clone confirmed homologous

recombination. The genomic DNA from ES cell clones and wild-type cells was

digested with Avr Ⅱ or Sac Ⅰ and hybridized with a 5' external probe or a 3' internal

probe. C. PCR identification of mouse genotyping showing $Crb3^{wt/wt}$, $Crb3^{wt/fl}$ and

$Crb3^{fl/fl}$ alleles. D. The expression of Crb3 was detected by immunoblotting in

mammary gland tissues in $Crb3^{fl/fl}$ and $Crb3^{fl/fl}$;MMTV-*Cre* mice at 8 weeks old. E.

Real-time quantitative PCR showing the relative mRNA expression of *Crb3* in

mammary gland tissues (n=10 mice). F. Crb3 expression was assessed using

immunohistochemistry in mammary gland tissues. The mammary epithelial cells were

marked by areas of white dotted lines. (scale bars, 50 μm) Bars represent the means ±

SD; Unpaired Student's *t* test, ***P<0.001.

1105

Supplementary Figure 2. CRB3 knockdown promotes proliferation of mammary

epithelial cells, while overexpression promotes apoptosis of breast cancer.

A. B. Immunoblot and real-time quantitative PCR analysis of CRB3 knockout

efficiency in MCF10A cells. C. MCF10A proliferation assay over six successive days.

D. Cell cycle showing the distribution of MCF10A cells. E. Quantification of cell

cycle distribution. F.G.H. Apoptosis of MCF7, T47D, and MDA-MB-453 breast

cancer cells with CRB3b overexpression and quantification of apoptotic cells. The

data are three independent experiments performed in triplicate. Bars represent the

means ± SD; Unpaired Student's *t* test, * *P*<0.05, **P<0.01.

1115

1116   **Supplementary Figure 3.** CRB3 alters primary cilium formation in MEFs from

1117 $Crb3^{fl/fl}$;CAG-*Cre* mice.

1118   A. Immunoblot analysis of CRB3b conditional overexpression upon dox induction

efficiency in MCF7 cells. B. Representative immunofluorescent staining of primary cilium formation in MEFs from *Crb3*^{fl/fl} and *Crb3*^{fl/fl};CAG-*Cre* mice. Acetylated tubulin (red), γ-tubulin (green), and DNA (blue). (scale bars, 10 μm) C. Quantification of the proportion of cells with primary cilium formation (At least 3-4 random micrographs were analyzed for each condition in each of three independent experiments, n=10). Bars represent the means ± SD; Unpaired Student's *t* test, ***$P<0.001$.

Supplementary Figure 4. The effect of CRB3 on the expression of some ciliogenesis-related genes and Rab11.

A. Quantification of mRNA expression of ciliogenesis-related genes in MCF10A cells (normalized to GAPDH, data are three independent experiments performed in triplicate). B. Coomassie blue staining showing the gel electrophoresis of exogenous CRB3b immunoprecipitated precipitates. C. Immunoblot analysis of the effect of CRB3 on Rab11 in MCF10A cells. Bars represent the means ± SD; Unpaired Student's *t* test.

Supplementary Figure 5. Rab11 knockdown caused defects in primary cilium.

A. Representative images of immunofluorescent staining of primary cilium formation with Rab11 knockdown in MCF10A cells. Acetylated tubulin (red), γ-tubulin (green), and DNA (blue). (primary cilium is marked by arrows; scale bars, 20 μm) B. Quantification of the proportion of cells with primary cilium formation in MCF10A cells (At least 3-4 random micrographs were analyzed for each condition in each of three independent experiments, n=10). C. Immunoblot analysis of Rab11 knockout efficiency in MCF10A cells. Bars represent the means ± SD; Unpaired Student's *t* test, ***$P<0.001$.

Supplementary Figure 6. Western blotting quantification of the molecules in the Wnt signaling pathway.

A. The relative protein levels of CRB3, GSK3-β, and β-catenin in CRB3-depleted MFC10A cells (normalized to β-actin). B. The relative protein levels of CRB3,

GSK3-β, and β-catenin in MDA-MB-231 and HCC 1806 breast cancer cells with CRB3b overexpression (normalized to β-actin). The data are three independent experiments performed in triplicate. Bars represent the means ± SD; Unpaired Student's *t* test, **$P<0.01$, ***$P<0.001$.

A

Crb3^{fl/fl}   Crb3^{wt/fl};CAG-Cre   Crb3^{fl/fl};CAG-Cre

B

Crb3^{fl/fl}   Crb3^{wt/fl};CAG-Cre   Crb3^{fl/fl};CAG-Cre

C

Crb3^{fl/fl}

Crb3^{fl/fl};MMTV-Cre

D

TEBs/gland (N)
***
Crb3^{fl/fl}   Crb3^{fl/fl};MMTV-Cre

E

Bifurcated TEBs (N)
***
Crb3^{fl/fl}   Crb3^{fl/fl};MMTV-Cre

F

Crb3^{fl/fl}   Crb3^{fl/fl};MMTV-Cre

G

PyMT-WT   PyMT-cKO-Crb3

A

	Day 3	Day 6	Day 9	Day 14
NC				
shCRB3				

B

Number of acini
NC shCRB3
**

Diameter of acini (μm)
NC shCRB3
***

Aberrant acini (%)
NC shCRB3
**

C

	Caspase 3	α-tubulin	DAPI	Merge
NC				
shCRB3				

D

Caspase 3 positive acini (%)
NC shCRB3
***

E

	α-tubulin	DAPI	Merge
NC			
shCRB3			

F

NC   shCRB3
90°   90°   Division angle
45°   45°
1 2   1 2 Cell#

Division angle (°)
NC shCRB3
*

G

	Ki67	Phospho-histone H3	Cleaved caspase 3
PyMT-WT			
PyMT-cKO-Crb3			

H

Ki67 positive cells (%)
NC shCRB3
*

No. of pHH3 positive cells per visual field
NC shCRB3
0.06

No. of cleaved caspase 3 positive cells per visual field
NC shCRB3
**

A

MCF10A

Pericentrin CRB3-GFP DAPI Merge

B

Cells with pericentrin and
CRB3 colocalization (%)

MCF10A

E

No. of γ-tubulin foci
per visual field

***

NC shCRB3

C

γ-tubulin CRB3 DAPI Merge

NC

shCRB3

D

Intensity

Distance (pixel)

F

Ace-tubulin CRB3 DAPI Merge

NC

shCRB3

G

Ace-tubulin CRB3 DAPI

5 μm 5 μm 1 μm

1 μm

A

neutrophil mediated immunity
neutrophil degranulation
neutrophil activation involved in immune response
neutrophil activation
Golgi vesicle transport
vesicle organization
antigen processing and presentation
Fc receptor signaling pathway
vesicle budding from membrane
formation of cytoplasmic translation initiation complex

Count
● 10
● 20
● 30
● 40

qvalue
1e-07
2e-07
3e-07

B

Rab small GTPases family
Rab-10, 11A, 11B, 14, 1A, 1B, 21, Rab-2A, 32, 35, 38, 5B, 5C, 6A

Centriolar proteins
Tubulin gamma-1,Tubulin gamma-2 CENP-E, CEP290, CEP192, CEP295, CEP162

C

EEA1 CRB3 DAPI
DMSO
Dynasore

CD63 CRB3 DAPI
DMSO
Dynasore

Rab11 CRB3 DAPI
DMSO
Dynasore

D

Cells with markers and CRB3 colocalization (%)

DMSO
Dynasore

EEA1 CD63 Rab11

*** *** ***

E

Cell density
MDSO   Dynasore
L   H   L   H   kDa

CRB3   28

GAPDH   36

F

γ-tubulin
GCP2, 3
GCP4, 5, 6

Primary cilium

γ-TuRC
γ-tubulin ring complex

G

NC shCRB3 kDa

CRB3   28
GCP6   200
GCP5   118
GCP3   104
GCP2   102
GCP4   76
γ-tubulin   48
β-actin   43

H

10% Input   Ig G   IP: GCP6

NC shCRB3 NC shCRB3 NC shCRB3

GCP3
GCP6
IgG

I

GCP3 GCP6 DAPI
NC
shCRB3

Intensity
200
100
0
0 20 40 60 80 100
Distance (pixel)

Intensity
200
100
0
0 100 200 300
Distance (pixel)

J

NC
γTuSC   γTuRC
5% - - - - - - - - - - - - - - - 40%

shCRB3
5% - - - - - - - - - - - - - - - - 40% kDa

GCP6   200
GCP5   118
GCP3   104
GCP2   102
GCP4   76
γ-tubulin   48

Fraction 1 2 3 4 5 6 7 8 9   1 2 3 4 5 6 7 8 9