CRB3 navigates Rab11 trafficking vesicles to promote γTuRC assembly during ciliogenesis

Bo Wang[1, 2, †], Zheyong Liang[1, 3, †], Tan tan[4], Miao Zhang[1, 2], Yina Jiang[5], Yangyang Shang[1, 2], Xiaoqian Gao[1, 2], Shaoran Song[1, 2], Ruiqi Wang[1, 2], He Chen[1, 2], Jie Liu[1, 2], Juan Li[1, 2], Yu Ren[6], and Peijun Liu[1, 2] *

[1]Center for Translational Medicine, the First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, Shaanxi, P.R. China.

[2]Key Laboratory for Tumor Precision Medicine of Shaanxi Province, the First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, Shaanxi, P.R. China.

[3]Department of Cardiovascular Surgery, the First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, Shaanxi, P.R. China.

[4]Center for Precision Medicine, Affiliated to The First People's Hospital of Chenzhou, University of South China, Chenzhou 423000, Hunan, P.R. China.

[5]Department of Pathology, the First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, Shaanxi, P.R. China.

[6]Department of Breast Surgery, the First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, Shaanxi, P.R. China.

† These authors contributed equally to the work.
* Correspondence:
Peijun Liu
E-mail: liupeijun@xjtu.edu.cn

Running Title: CRB3 regulates ciliogenesis

Keywords: Primary cilium, Polarity protein, Breast cancer, CRB3, γ-TuRC assembly, Rab11-positive endosome

Abstract

The primary cilium plays important roles in regulating cell differentiation, signal transduction, and tissue organization. Dysfunction of the primary cilium can lead to ciliopathies and cancer. The formation and organization of the primary cilium are highly associated with cell polarity proteins, such as the apical polarity protein CRB3. However, the molecular mechanisms by which CRB3 regulates ciliogenesis and CRB3 location remain unknown. Here, we show that CRB3, as a navigator, regulates vesicle trafficking in γ-TuRC assembly during ciliogenesis and cilium-related Hh and Wnt signaling pathways in tumorigenesis. *Crb3* knockout mice display severe defects of the primary cilium in the mammary ductal lumen and renal tubule. CRB3 is essential for lumen formation and ciliary assembly in the mammary epithelium. We demonstrate that CRB3 localizes to the basal body and that CRB3 trafficking is mediated by Rab11-positive endosomes. Significantly, CRB3 directly interacts with Rab11 to navigate GCP6/Rab11 trafficking vesicles to CEP290, resulting in intact γ-TuRC assembly. In addition, CRB3-depleted cells cannot respond to the activation of the Hh signaling pathway, while CRB3 regulates the Wnt signaling pathway. Therefore, our studies reveal the molecular mechanisms by which CRB3 recognizes Rab11-positive endosomes to navigate apical vesicle trafficking in effective ciliogenesis, maintaining cellular homeostasis and tumorigenesis.

Introduction

Epithelial tissues are the most common, widely distributed, and perform several particular functions. The integrity of the specific cellular architecture composed of epithelial cells is necessary for these functions of epithelial tissues. Establishing apical-basal cell polarity within epithelial cells is a crucial biological process for epithelial homeostasis. Most human cancers originate from epithelial tissues, and loss of apical-basal cell polarity is one of the most significant hallmarks of advanced malignant tumors[1]. Several reports have shown that loss of some polarity proteins causes disordered cell polarity within epithelial cells, which cannot tightly control cellular growth and migration, ultimately leading to tumor formation, progression,

and metastasis[2, 3]. However, the relationship between polarity proteins and tumorigenesis remains incomplete.

The crumb complex, located apically, is vital in regulating and maintaining apical-basal cell polarity within epithelial cells. In mammals, three Crumbs orthologs are CRB1, CRB2, and CRB3, and only CRB3 is widely expressed in epithelial cells[4]. Our previous studies have shown that CRB3 is expressed at low levels in renal and breast cancers, and abnormal CRB3 expression leads to the disrupted organization of MCF10A cells in three-dimensional (3D) cultures[5-7]. Inhibition of CRB3 impairs contact inhibition and leads to migration, invasion and tumorigenesis of cancer cells[6, 8]. The contact inhibition results in growth arrest and causes epithelial cells to enter the quiescent phase, inducing primary cilium formation.

The primary cilium, an antenna-like microtubule-based organelle in most types of mammalian cells, is a sensorial antenna that regulates cell differentiation, proliferation, polarity, and tissue organization[9]. Dysfunction of the primary cilium can lead to developmental and degenerative disorders called ciliopathies, such as polycystic kidney disease, nephronophthisis, retinitis pigmentosa, and Meckel syndrome[10]. Remarkably, it has been reported that many cancers, including melanoma and breast, renal, pancreatic, and prostate cancer, exhibit loss of the primary cilium, most likely during the early stages of tumorigenesis[11-15]. The primary cilium is a significant extension on the apical surfaces of epithelial cells and is maintained by polarized vesicular traffic[16]. Some studies have shown that CRB3 localizes in the primary cilium and that depletion of CRB3 leads to primary cilium loss in Madin-Darby canine kidney (MDCK) cells[17, 18], indicating that CRB3 is necessary for the initiation of ciliogenesis. However, how CRB3 promotes ciliogenesis on the apical surfaces of epithelial cells remains unknown.

Here, we describe a novel conditional deletion of *Crb3* in mice and show that Crb3 is required to assemble the primary cilium in the gland, kidney and MEF cells from *Crb3* knockout mice and that *Crb3* deletion promotes breast cancer progression *in vivo*. We found that CRB3, which directly interacts with Rab11, navigates GCP6/Rab11 trafficking vesicles to the basal body of the primary cilium in mammary

epithelial cells. Additionally, we identified that CRB3 regulates the ciliary Hedgehog

(Hh) and Wnt signaling pathways in breast cancer. Thus, we found unexpected

diversity in the polarity protein CRB3 navigating polarized Rab11-dependent traffic,

leading to a novel assembly mechanism of the γ-tubulin ring complex (γTuRC) in

ciliogenesis.

Results

Crb3 **deletion mice exhibit smaller and anophthalmia.**

Since *Crb3* knockout mice die shortly after birth[19], we generated a novel transgenic

mouse model with conditional deletion of *Crb3* using the Cre-loxP system. The loxP

sites flanked either side of exon 3 in the *Crb3* gene (Supplementary Fig. 1A). The

positive embryonic stem (ES) cell clone was confirmed for the wild-type gene

($Crb3^{wt/wt}$) and recombinant allele ($Crb3^{wt/fl}$) by Southern blotting (Supplementary Fig.

1B). The genotypes of the wild type, heterozygotes and homozygotes ($Crb3^{fl/fl}$) were

detected with specific primers (Supplementary Fig. 1C). We first intercrossed $Crb3^{fl/fl}$

with CMV enhancer/chicken β-actin promoter (CAG)-Cre mice and found that

$Crb3^{fl/fl}$, $Crb3^{wt/fl}$; CAG-Cre and $Crb3^{fl/fl}$; CAG-Cre pups were born normally and

looked well developed in size. However, most $Crb3^{fl/fl}$; CAG-*Cre* pups died within a

few days after birth. Few surviving $Crb3^{fl/fl}$; CAG-*Cre* mice were smaller and showed

anophthalmia than littermate $Crb3^{fl/fl}$ mice at 4 weeks old (Fig. 1A-B). The eye is an

organ with perfect apical-basal cell polarity. However, *Crb3* knockout mice showed

ocular abnormalities. Together, these results indicated that Crb3 is necessary for eye

development.

Mammary epithelial-specific *Crb3* deletion causes ductal epithelial hyperplasia

and tumorigenesis.

Based on our earlier studies showing that inhibition of CRB3 impairs the organization

of breast epithelium, we set out to directly explore the role of Crb3 in mammary gland

development. We first developed epithelial cell-specific deletion of *Crb3* in the

mammary gland by intercrossing $Crb3^{fl/fl}$ mice with mouse mammary tumor virus

long terminal repeat promoter (MMTV)-*Cre* mice. We demonstrated the expression of

Crb3 in the mammary gland using immunoblotting, real-time quantitative PCR, and

immunohistochemistry, and the level of Crb3 expression was deficient in mammary

epithelial cells with Crb3 knockout (Supplementary Fig. 1D-F).

To assess the function of Crb3 in the development of mammary epithelial cells, we

analyzed whole mammary mounts from 8-week-old virgin *Crb3*^{fl/fl};MMTV-*Cre* and

littermate *Crb3*^{fl/fl} mice. *Crb3*^{fl/fl};MMTV-*Cre* mice displayed significantly increased

numbers of terminal end buds (TEBs) and bifurcated TEBs than *Crb3*^{fl/fl} mice (Fig. 1

C-E). The histopathology results showed that normal ductal epithelial cells were

arranged as a monolayer in *Crb3*^{fl/fl} mice, while *Crb3* knockout led to ductal epithelial

hyperplasia with increased ductal thickness in *Crb3*^{fl/fl};MMTV-*Cre* mice (Fig. 1 F).

To observe the role of Crb3 in the tumorigenesis of breast cancer, we observed

mammary glands from 9-week-old virgin polyomavirus middle T antigen

(PyMT)-cKO-Crb3 and PyMT-WT mice whose mammary glands progressed to early

carcinomas and lost their normal epithelial architecture[20]. The PyMT mouse model

has been widely used in the study of tumor initiation and development with similar

histological progression to human breast cancer. PyMT-WT mice exhibited loss of

epithelial features and development of breast cancer, while PyMT-cKO-Crb3 mice

grew larger tumors with faster tumor progression and a poorly differentiated

phenotype (Fig. 1G). Thus, Crb3 knockout leads to ductal epithelial hyperplasia and

promotes tumorigenesis, and it is essential for branching morphogenesis in mammary

gland development and tumor progression.

CRB3 knockdown inhibits lumen formation in acini of mammary epithelial cells.

MCF10A cells are spontaneously immortalized human breast epithelial cells with the

characteristics of normal breast epithelium[21]. MCF-10A cells could form acini

similar to mammary epithelial acini under 3D basement membrane culture conditions.

It has become a valuable system to study the morphogenesis and oncogenesis of the

mammary epithelium[22]. To confirm the function of CRB3 in mammary epithelial

acini formation, we first knocked down the expression of CRB3 by lentiviral shRNA

in MCF10A cells (Supplementary Fig. 2A and B). Under the 3D culture system, we

found that MCF10A cells infected with no target shRNA lentivirus could form acini

after 6 days (D6) and finally formed polarized acini with a hollow lumen at D14 (Fig. 2A). However, MCF10A cells, after CRB3 knockdown, formed more number, larger diameter and aberrant acini after D6, and these acini had no lumen (Fig. 2A-D). Cell proliferation assays of MCF10A cells showed that CRB3 knockdown increased the proliferation rate and accelerated G1 to S phase progression (Supplementary Fig. 2C-E).

Apoptosis and mitotic spindle orientation during lumen formation were further investigated in a 3D culture system. The immunofluorescence (IF) results showed that more internal cleaved-caspase 3-positive acini were observed in the control groups than in the CRB3 knockdown groups at D9 (Fig. 2E). CRB3 overexpression significantly promoted apoptosis in breast cancer cells (Supplementary Fig. 2F-H). Similarly, the staining of α-tubulin in mitotic cells exhibited more misorientation of the mitotic spindle in CRB3 knockdown groups (Fig. 2F-H).

In addition to verifying the effect of *Crb3* deletion on promoting proliferation and inhibiting apoptosis *in vivo*, these markers were detected in primary tumor tissue from PyMT-WT and PyMT-cKO-Crb3 mice. *Crb3* deletion increased the numbers of proliferative and mitotic cells and decreased the numbers of apoptotic cells in primary tumors (Fig 2I). These results demonstrate that CRB3 could promote the proliferation of mammary epithelial cells, disturb the mitotic spindle orientation, and protect the internal cells from apoptosis during acini formation, finally leading to irregular lumen formation and tumor progression.

CRB3 regulates primary cilium formation.

The primary cilium, one of the apical antenna-like extensions in epithelial cells, has been reported to play an important role in controlling lumen formation[23]. CRB3 knockdown leads to primary cilium loss in MDCK cells[17, 18], but the molecular mechanism regulating ciliogenesis on apical surfaces is still unclear. Investigating the altered effect of CRB3 on ciliogenesis *in vivo* and *in vitro*, we found that control MCF10A cells formed primary cilia after becoming confluent, while CRB3 knockdown led to significant ciliogenesis defects (Fig. 3A). Compared with the control, the number of cells with primary cilium was significantly increased, and the

length of the existing primary cilium was not different (Fig. 3B). Conversely, CRB3

conditional overexpression, adding doxycycline (+dox), resulted in restoring ciliary

assembly in MCF7 cells (Fig. 3C). CRB3 overexpression increased the proportion of

primary cilium formation in breast cancer cells, and the length of the restored primary

cilium was increased (Fig. 3D). Importantly, the primary cilium could be directly

visualized using scanning electron microscopy (SEM) (Fig. 3E). MCF10A cells with

primary cilia were significantly decreased after CRB3 knockdown, while CRB3

conditional overexpression did not increase these populations (Fig. 3F).

To further corroborate these findings *in vivo*, we used breast and kidney tissue to

assess ciliogenesis in the *Crb3* knockout mouse model. Renal epithelial cells can form

prominent primary cilia, and the absence of primary cilia leads to ciliopathies. We

used immunofluorescence to detect ciliary formation in the mammary ductal lumen

and renal tubule. Compared with tissues of $Crb3^{fl/fl}$ mice, *Crb3* knockout led to the

absence of primary cilium in the mammary ductal lumen and renal tubule from

$Crb3^{fl/fl}$; CAG-*Cre* mice (Fig. 3G, H). In particular, the proportion of cells forming the

primary cilium plummeted (Fig. 3G, H). This same phenomenon was also observed in

mouse embryonic fibroblasts (MEFs) from $Crb3^{fl/fl}$; CAG-*Cre* mice (Supplementary

Fig. 3A, B). We conclude that CRB3 plays a central role in ciliogenesis in various

tissues and mammary cells.

CRB3 localizes to the basal body of the primary cilium.

CRB3, a polarity protein, was mostly reported to be localized on the tight junctions of

the apical epithelium membrane[24]. Additionally, we noted that CRB3 could be

closely colocalized with centrosomes at prophase, metaphase, and anaphase[17]. To

investigate the relationship between CRB3 and ciliary formation, we first examined

CRB3 localization in MCF10A cells that expressed CRB3-GFP. In MCF10A cells,

exogenous CRB3 was mainly localized at cell tight junctions and could colocalize

with pericentrin (a centrosome marker) (Fig. 4A). And approximately 40% of cells

had this co-localization of exogenous CRB3 and pericentrin (Fig. 4B). We observed

the localization of endogenous CRB3 with another centrosome marker to verify this

phenomenon. Similarly, we detected that CRB3 accumulated on one side of the

cytoplasm and had a CRB3 focous located at the basal body represented by γ-tubulin in MCF10A cells (Fig. 4C, D). Importantly, CRB3 knockdown disturbed the accumulation of γ-tubulin at the basal body in the confluent quiescence cells (Fig. 4C). To demonstrate the relationship of this colocalization to the primary cilium, we employed acetylated tubulin to mark the primary cilium. Double immunostaining showed that this CRB3 focous was the basal body of the primary cilium, and CRB3 knockdown inhibited ciliary assembly in MCF10A cells (Fig. 4E, F). These results suggest that CRB3 is located at the basal body of the primary cilium and that CRB3 knockdown disturbs the accumulation of γ-tubulin in quiescent cells.

CRB3 trafficking is mediated by Rab11-positive endosomes.

To better investigate the molecular mechanisms of CRB3 in regulating primary cilium formation, we detected a series of genes related to ciliogenesis, including markers of centriole or primary cilium, basal body (BB) components, intraflagellar transport (IFT)-A and IFT-B anterograde transport. However, CRB3 knockdown did not alter the mRNA expression of these genes (Supplementary Fig. 4A). Then, we carried out coimmunoprecipitation (co-IP) tandem mass spectrometry using tagged exogenous CRB3 as bait to identify its interacting proteins (Supplementary Fig. 4B). Pathway aggregation analysis of these proteins revealed that these interacting proteins of CRB3 were significantly involved in Golgi vesicle transport and vesicle organization (Fig. 5A). We focused on these pathways because smaller distal appendage vesicle (DAV) formation is critical for ciliogenesis initiation, and it requires the Rab GTPase Rab11-Rab8 cascade to function[25]. Interestingly, some Rab small GTPase family members identified as CRB3-binding proteins were aggregated in these pathways, such as Rab10, Rab11A, Rab11B, Rab14, Rab1A, Rab1B, Rab21, Rab2A, Rab32, Rab35, Rab38, Rab5B, Rab5C, and Rab6A. Several centriolar proteins were also identified, including tubulin gamma-1, tubulin gamma-2, CENP-E, CEP290, CEP192, CEP295, and CEP162 (Fig. 5B).

Rab11-positive vesicles bind to centrosomal Rabin8, leading to ciliary membrane formation[26]. This report suggested that the trafficking of Rab11-positive vesicles and polarized vesicles is an essential process for early ciliary assembly. Next, we

examined the polarized vesicle traffic of CRB3 in MCF10A cells. Our studies found

that CRB3 could colocalize with EEA1-positive early endosomes, CD63-positive late

endosomes and Rab11-positive recycling endosomes (Fig. 5C). In addition, after 2

hours of treatment with dynasore, an endocytosis inhibitor, CRB3 significantly

accumulated at the cytomembrane, and the colocalization of CRB3 with EEA1-,

CD63-, and Rab11-positive endosomes was significantly decreased (Fig. 5C, D). We

have reported that CRB3 expression correlates with the contact-dependent change,

which is CRB3 expression reduced considerably in confluent MCF10A cells[6]. Then,

dynasore could also rescue the CRB3 downregulation induced by the increasing cell

density of MCF10A cells (Fig. 5E). CRB3 knockdown did not affect Rab11

expression (Supplementary Fig. 4C). These results suggest that the intracellular

trafficking of CRB3 is mediated by Rab11-positive endosomes at different cell

densities.

CRB3 knockdown destabilizes γTuRC assembly during ciliogenesis.

According to our findings, we assumed that CRB3 might play an important role in

polarized vesicle trafficking. The γ-Tubulin ring complex (γTuRC) is a central

regulator of microtubule nucleation and is a nucleation site of α-Tubulin and

β-Tubulin at the microtubule-organizing center in mitotic cells[27]. The primary

cilium is a microtubule-based structure attached to the basal body transformed from

the mother centriole. As we found that CRB3 knockdown disturbed the accumulation

of γ-tubulin, we analyzed γTuRC assembly in ciliogenesis. The γTuRC is composed

of multiple copies of the γ-tubulin small complex (γTuSC), GCP2 and GCP3, plus

GCP4, GCP5, and GCP6[27, 28] (Fig. 5F). CRB3 downregulation did not affect the

expression of γTuRC subunits (Fig. 5G). Then, we performed co-IPs directed against

GCP6 in CRB3-depleted cells and assessed the relative levels of both γTuSC-specific

proteins compared with the control shRNA-treated cells. Interestingly, CRB3

knockdown significantly reduced the interaction of GCP6 with GCP3 (Fig. 5H). To

verify this phenomenon, we observed the localization of GCP3 with GCP6 foci in

MCF10A cells. Correspondingly, CRB3 downregulation significantly disturbed the

colocalization of GCP3 with GCP6 foci in quiescence cells (Fig. 5I). To investigate

whether CRB3 facilitate γTuRC assembly, we fractionated MCF10A cytoplasmic

proteins on sucrose gradients according to the method published[29]. We noticed that

γTuSC were regimented mainly in fractions 3 and γTuRC sediment in fractions 6 (Fig.

5J). However, CRB3 downregulation caused most of GCP6 and GCP5 to be

precipitated independently of γTuRC in the low-density fractions (Fig. 5J). Together,

these results have established that the depletion of CRB3 interferes with the molecular

interactions between the γTuRC proteins, failing microtubule formation in the primary

cilium. However, it may not affect the expression of these molecules during γTuRC

assembly. According to these results, we hypothesized that CRB3 could mediate the

polarized vesicle trafficking of some γTuRC-specific proteins during Rab11-positive

endosomes.

CRB3 directly interacts with Rab11.

To fully prove this hypothesis, we first examined CRB3 interaction with

Rab11-positive endosome and γTuRC-specific proteins in MCF10A cells. Using

CRB3 as bait, we identified that CRB3 could interact with Rab11 instead of GCP6

and GCP3 (Fig. 6A). In contrast, Rab11 could interact with CRB3 and GCP6 instead

of GCP3 by tagging Rab11 as bait (Fig. 6B). Thus, does CRB3 knockdown affect the

trafficking of γTuRC-specific proteins during Rab11-positive vesicles? The co-IP

results showed that CRB3 knockdown did not affect Rab11 interacting with GCP6,

and the level of Rab11 binding to GCP6 had a tendency to decrease in MCF10A cells

with CRB3 knockdown (Fig. 6C). The colocalization rate of Rab11 and GCP6 was

significantly decreased in MCF10A cells with CRB3 knockdown (Supplementary Fig.

5A-C). Consistent with other reports[30], Rab11 knockdown significantly inhibited

primary cilium formation in MCF10A cells (Supplementary Fig. 5D-F).

Based on these results, we further detected the region of CRB3 interacting with Rab11

by using coexpression and co-IP in HEK293 cells. CRB3 consists of a signal peptide

(SP), extracellular domain, transmembrane (TM), FERM-binding domain (FBD), and

carboxy-terminal PDZ-binding domain (PBD). Due to the specificity of the C- and

N-terminal domains, we constructed CRB3-GFP fusion proteins where GFP tags were

fused to the extracellular domain as previously reported[31] (Fig. 6D). According to

these domains, we also constructed the truncations of CRB3-GFP fusion proteins with

serial C-terminal deletions (Fig. 6E) and another Flag-Rab11a fusion protein. The

coexpression and co-IP results showed that Flag-Rab11a interacted with full-length

CRB3 (1-120), CRB3 (1-116), CRB3 (1-83), and CRB3 (1-58) but not CRB3 (1-26)

(Fig. 6F). Therefore, the region of CRB3 interacting with Rab11 is amino acids 27 to

58.

Rab11, a small GTP-binding protein, is an essential regulator of the dynamics in

recycling endosomes, which are required to undergo GDP/GTP cycles. In particular,

Rab11a[S25N] is a GDP-locked form, Rab11a[S20V], and Rab11a[Q70L] is a

GTP-locked form[30]. To further investigate the GTP-bound form of Rab11a

interacting with CRB3, we constructed these mutant variants of Rab11a. Next, we

found that the weaker interaction of CRB3 and these mutant variants, Rab11a[Q70L],

Rab11a[S20V] and Rab11a[S25N], compared with Rab11aWT, was observed using

co-IP analysis (Fig. 6G). Together, these results indicate that CRB3 directly interacts

with Rab11a, depending on the GTPase activity of Rab11a.

CRB3 navigates GCP6/Rab11 trafficking vesicles to CEP290 in the primary

cilium.

Since CRB3 knockdown does not affect the interaction with Rab11-positive

endosome and γTuRC-specific proteins GCP6, we wanted to know whether CRB3

affected the localization of GCP6/Rab11 trafficking vesicles to the basal body of the

primary cilium. Reviewing the identification of CRB3 interacting proteins, there are

some centriolar proteins. CEP290 is located in the transition zone of the primary

cilium and is required for the formation of microtubule-membrane linkers[32]. Then,

we verified that exogenous CRB3 could bind to CEP290, Rab11, GCP6 (Fig. 7A), and

Rab11 also interacted with CEP290, CRB3-GFP, GCP6 (Fig. 7B). Similar to Fig. 6C,

the amount of GCP6 bonded to Rab11 was reduced within CRB3 knockdown.

Importantly, CRB3 knockdown showed that Rab11 hardly bonded to CEP290 (Fig.

7C). We checked the colocalization of the basal body foci of GCP6 and γ-tubulin,

γ-tubulin and Rab11 in MCF10A and MEF cells to corroborate this result. In

quiescence control cells, GCP6 foci and γ-tubulin formed a prominent focus at the

basal body of the primary cilium, as observed by immunofluorescence. However, CRB3 downregulation eliminated this focus and significantly disturbed the basal body foci of GCP6 and γ-tubulin in MCF10A cells (Fig. 7D), and γ-tubulin and Rab11 in MEF cells (Fig. 7E). At the beginning of this experiment, we also tried to show the interaction of CRB3, Rab11 and GCP6 with expressing fluorescence tagged GCP6 protein and FRET assay. However, the centrosomal foci of fluorescent GCP6 was difficult to capture because of too weak signal, and this may require re-overexpression of ninein-like protein (Nlp) to promote γTuRC enrichment[29, 33].

Although CRB3 knockdown had little effect on the formation of GCP6/Rab11 trafficking vesicles, it could significantly inhibit the γTuRC assembly during ciliogenesis, thus affecting the transport of GCP6/Rab11 trafficking vesicles to the basal body of the primary cilium. Since CRB3 affected the combination of Rab11 and CEP290, we speculate that CRB3 could be a navigator, navigating GCP6/Rab11 trafficking vesicles to the basal body of the primary cilium.

CRB3 regulates the Hh and Wnt signaling pathways in tumorigenesis

To examine the role of CRB3 in regulating ciliary assembly in breast cancer, we explored the relationship between CRB3 and the primary cilium in breast cancer tissues. We found that CRB3 was localized at the subapical surface of the mammary gland lumen in the paracarcinoma tissues. CRB3 had no obvious expression or localization in breast cancer tissues (Fig. 8A). The immunofluorescence analysis results showed that the intact primary cilium could be observed in the adjacent para-carcinoma tissues, whereas the number of cells with primary cilia was much lower in breast cancer tissues (Fig. 8B, C). Consistent with previous findings, CRB3 was localized at the basal body of the primary cilium in adjacent para-carcinoma tissues (Fig. 8D). These results again suggest that a defect in CRB3 expression inhibits ciliary assembly, which could be involved in tumorigenesis.

To detect some functional consequences of the absence of the primary cilium, we measured alterations in the Hedgehog (Hh) and Wnt signaling pathways in CRB3-depleted cells. We first detected the translocation of SMO into the primary cilium and the expression of the target gene GLI1. SAG is a potent Smoothened

(SMO) receptor agonist that activates the Hh signaling pathway. CRB3 knockdown

did not promote the translocation of SMO into the primary cilium in MCF10A cells

treated with SAG (Fig 8E). After SAG treatment, the mRNA expression of GLI1 was

significantly decreased in CRB3-depleted cells (Fig 8F). Crb3 knockout did not alter

the expression or cellular localization in primary tumors from PyMT-cKO-Crb3 mice

(Fig 8H). We have reported that CRB3 expression was high in immortalized

mammary epithelial cells while loss of CRB3 expression in breast cancer cells[6]. On

the other hand, western blotting results showed that GSK3-β was significantly

downregulated after CRB3 knockdown in MCF10A cells. In addition, β-catenin, the

major effector of the canonical Wnt signaling pathway, was upregulated. CRB3

overexpression resulted in upregulation of GSK3-β and downregulation of β-catenin

in breast cancer cells (Fig. 8G). Visibly, β-catenin was significantly upregulated and

markedly localized in the nucleus from PyMT-cKO-Crb3 mice (Fig 8I). Together,

these data suggest that CRB3 knockdown cannot activate the Hh signaling pathway

but can activate the Wnt signaling pathway, which is ultimately involved in

tumorigenesis.

Discussion

Apical-basal cell polarity is essential for cellular architecture, development and

homeostasis in epithelial tissues. Most malignant cancers lack cell polarity because of

the unconstrained cell cycle and cell migration with decreased adhesion. CRB3, an

important apical protein, is significantly reduced in various tumor cells and tissues

and promotes metastasis and tumor formation in nude mice[5, 6, 8, 34-36]. Here, we

generate a novel transgenic mouse model with conditional deletion of Crb3 to study

the role of polarity proteins in tumorigenesis. *Crb3* knockout mice were reported to

have cystic kidneys, improper airway clearance in the lung, villus fusion, apical

membrane blebs, and disrupted microvilli in intestine airway clearances, which

demonstrated that CRB3 is crucial for the development of the apical membrane and

epithelial morphogenesis[19, 37]. Importantly, our studies in the *Crb3* knockout

mouse model suggest that in addition to death after birth, it is characterized by smaller

size, ocular abnormality, bronchial smooth muscle layer defect, reduced pancreatic islets and hyperplasia mammary ductal and renal tubular epithelium (some data not shown). These important phenotypic changes are due to disruption of epithelial polarity homeostasis, which may ultimately lead to tumorigenesis.

In addition, we also found the absence of primary cilia in the mammary ductal lumen and renal tubule in the *Crb3* knockout mouse model. Consistent with other reports, CRB3 knockdown regulated ciliary assembly in MDCK cells[17, 18]. Furthermore, CRB3 was localized in the inner segments of photoreceptor cells and concentrated with the connecting cilium during the whole development of the mouse retina[38]. These ocular defect phenotypes remarkably resemble EHD1 knockout mice, which regulate ciliary vesicle formation in primary cilium assembly[25, 39, 40]. Hence, CRB3 can affect primary cilium formation in various epithelial tissues, such as the breast, kidney, and retina.

Previous literature has reported that CRB3 localizes in the primary cilium in differentiated MDCK cells[17, 18], but we mainly found that it is not located on the basal body of the primary cilium in mammary epithelial cells. The immunofluorescence results in these literatures showed that CRB3 was scattered on the primary cilia, and had a strong foci at the basal body. Especially in rat kidney collecting ducts, the localization of CRB3 on primary cilia was significantly reduced, while the obvious localization was basal body[18]. Another literature also reported the co-localization of CRB3 and γ-tubulin in MDCK cells[17]. This result is consistent with our conclusion, and we also verified its co-localization with centrosome by overexpressing CRB3 in mammary epithelial cells, indicating that CRB3 mainly localized to the basal body of the primary cilium. Given that CRB3 also alters primary cilium formation in the mammary epithelium, it is important to investigate the molecular mechanisms by which CRB3 regulates primary cilium formation.

Polarized vesicular traffic plays an important role in extending the primary cilium on apical epithelial surfaces, and the molecular mechanism and relationship to tumorigenesis remain unclear. Here, we reveal that Rab11-positive endosomes mediate the intracellular trafficking of CRB3 and that CRB3 can navigate

GCP6/Rab11 trafficking vesicles to CEP290 for correct γTuRC assembly during primary cilium formation. Moreover, in the presence of CRB3, the epithelium can form contact inhibition and an intact primary cilium in quiescence to achieve cellular homeostasis. In contrast, loss of contact inhibition and primary cilium formation with CRB3 deletion can activate the Wnt signaling pathway, affect the Hh signaling pathway, and disrupt the imbalance of this cellular homeostasis, resulting in significant cell proliferation during tumorigenesis (Fig. 9).

The formation of the primary cilium has been divided into several distinct phases. It begins with the maturation of the mother centriole, also known as a centriole-to-basal-body transition[9, 41]. DAVs, which are small cytoplasmic vesicles originating from the Golgi and the recycling endosome, accumulate in the vicinity of distal appendages of the mother centriole[25, 42, 43]. Rab8a is localized to cytoplasmic vesicles and the Golgi/trans-Golgi network in growing RPE1 cells, while Rab8-positive vesicles are rapidly redistributed to the mother centriole with binding to Rab11-positive recycling and are endosomes under serum starvation[26, 44]. The EHD1 protein converts these DAVs to form larger ciliary vesicles, which elongate through continuous fusion with Rab8-positive vesicles to produce the primary cilium membrane[25, 30, 40]. Thus, Rab11-positive recycling endosomes, as important transporters, are necessary for primary cilium formation and a centriole-basal-body transition in the early phase. Our study found that CRB3 could bind to some Rab small GTPase family members to significantly participate in Golgi vesicle transport and vesicle organization pathways through mass spectrometry identification. And coexpression and co-IP results showed that CRB3 directly interacted with Rab11 via extracellular domain, which mediated the intracellular trafficking of CRB3 within Rab11-positive endosomes at different cell densities.

Unlike motile cilia, primary cilia have only nine outer doublet microtubules without a central pair of microtubules (9+0 structure). Similar to microtubule assembly at microtubule-organizing centers (MTOCs) during mitosis, γ-tubulin, which is a part of γTuRC, is also concentrated in the basal body of the cilium in interphase[45]. γ-Tubulin complexes have been shown to form microtubule templates and regulate

microtubule nucleation through longitudinal contacts with α-tubulin and β-tubulin[27].

The γTuRC consists of multiple copies of γTuSC, which is composed of two copies of

γ-tubulin and one each of GCP2 and GCP3. Then, GCP2 and GCP3 interact with

GCP4, GCP5 and GCP6 to form γTuRC[46]. Our experiments showed that CRB3

knockdown does not affect the expression of the γTuRC-specific proteins GCP3,

GCP6 and γ-tubulin but inhibits the interaction between GCP6 and GCP3, resulting in

destabilization of γTuRC. Further analysis revealed significant colocalization between

GCP6- and Rab11-positive vesicles at low cell density, while at high cell density,

GCP6 accumulated at the centrosome together with GCP3. CRB3 knockdown results

in the disappearance of this accumulated GCP6. These results indicated that

GCP6/Rab11 trafficking vesicles must be transported to the basal body for

ciliogenesis. While it is still unclear how CRB3 interferes with γTuRC assembly

during primary cilium formation, we hypothesize that Rab11-positive endosomes

mediate the intracellular trafficking of CRB3 and that CRB3, as a navigator

interacting with CEP290, can navigate GCP6/Rab11 trafficking vesicles to the

transition zone for γTuRC assembly in primary cilium formation.

CEP290 is considered a centriolar or ciliary protein localized to the centrosomes in

dividing cells, the distal mother centriole in quiescent cells, and the transition zone in

the primary cilium[47]. *CEP290* mutations are associated with various diseases, such

as the devastating blinding disease Leber's Congenital Amaurosis (LCA),

nephronophthisis, Senior Løken syndrome (SLS), Joubert syndrome (JS),

Bardet-Biedl syndrome (BBS), and lethal Meckel-Gruber syndrome (MGS)[48]. In

particular, the phenotypes of CEP290-associated ciliopathies are very similar to those

in our *Crb3* knockout mouse model, with lethality and ocular and renal abnormalities.

CEP290 knockdown significantly decreased the number of cells with primary cilia,

and CEP290 knockout mice showed mislocalization of ciliary proteins in the

retinas[49, 50]. Additionally, the depletion of CEP290 disrupts protein trafficking to

centrosomes and affects the recruitment of the BBSome and vesicular trafficking in

ciliary assembly[51, 52]. Although CEP290 serves as a hub to recruit many ciliary

proteins, it has not been reported to be involved in polarized vesicle trafficking or

localization. Our study verified that CEP290 could interact with the polarity protein

CRB3. CRB3 knockdown disrupts the interaction between CEP290 and

Rab11-positive endosomes. Thus, these results validate our hypothesis. Although

CRB3 navigates GCP6/Rab11 trafficking vesicles to CEP290, we further need to

detect the region of CRB3 interacting with CEP290 and what other cargos are

transported by CRB3/Rab11 trafficking vesicles in ciliary assembly.

In addition, our results showed that CRB3 knockdown could not activate the Hh

signaling pathway with SAG in MCF10A cells. Under untreated conditions, PTCH1

located at the ciliary membrane inhibits SMO, and Suppressor of Fused (SuFu)

sequesters Gli transcription factors, leading to its degradation. SAG can activate the

Hh signaling pathway, resulting in dissociation into the nucleus to regulate the

expression of downstream target genes[53]. The Hh signaling pathway mainly

regulates cellular growth and differentiation and is abnormal in different types of

tumors. Some studies indicate that cilia are double-sided, promoting and preventing

cancer development through the Hh signaling pathway *in vivo*. It is thought that cilia

deletion could activate SMO to inhibit tumor growth, while promoting carcinogenesis

was induced by activated Gli2[54]. Although SAG cannot activate the Hh signaling

pathway in CRB3-depleted cells, CRB3 regulation of the carcinogenic process

through the ciliary Hh signaling pathway still needs further research. In addition,

another canonical Wnt signaling pathway was altered after CRB3 knockdown. The

Wnt signaling pathway is an important cascade regulating development and stemness,

and aberrant Wnt signaling has been reported in many more cancer entities, especially

colorectal cancer[55]. CRB3 downregulated the expression of β-catenin in our study.

Thus, CRB3 can affect cilium-related signaling pathways in tumorigenesis.

In summary, our study provides novel evidence for the apical polarity protein CRB3

in the regulation of viability, mammary and ocular development, and primary cilium

formation in germline and conditional knockout mice. CRB3 deletion causes irregular

lumen formation and ductal epithelial hyperplasia in mammary epithelial-specific

knockout mice. Further study uncovered a new function of CRB3 in directly

interacting with Rab11, navigating GCP6/Rab11 trafficking vesicles to CEP290 for

γTuRC assembly during ciliogenesis. In addition, CRB3 also participates in regulating

cilium-related Hh and Wnt signaling pathways in tumorigenesis. Understanding these

polarity protein-mediated vesicle trafficking mechanisms will shed new light on

luminal development and tumorigenesis.

Methods

Generation of floxed *Crb3* mice. All animal experiments were verified and approved

by the Committee of Institutional Animal Care and Use of Xi'an Jiaotong University.

Heterozygous $Crb3^{wt/fl}$ mice (C57BL/6J) were generated by Cas9/CRISPR-mediated

genome editing (Cyagen Biosciences, Guangzhou, China). The gRNA to *Crb3* and

Cas9 mRNA was coinjected into fertilized mouse eggs to generate targeted

conditional knockout offspring. The sequences of the *Crb3* gRNA were gRNA1,

5'-GGCTGGGTCCACACCTACGGAGG-3';   gRNA2,

5'-ACCCACAAAGCCACGCCA GTGGG-3'. Mouse genotyping was screened by

PCR with the primers F1, 5'-ACATAAGGCCTTCCGTTAAGCTG-3'; R1, 5'-

GTGGATTCGGACCAGTCTGA-3'. $Crb3^{wt/fl}$ mice were intercrossed with

MMTV-*Cre* mice or CAG-*Cre* mice to generate tissue-specific *Crb3* knockout mice.

MMTV-*Cre* mice (FVB), CAG-*Cre* mice (C57BL/6N) and MMTV-PyMT mice (FVB)

were purchased from Cyagen Biosciences. All mice were bred in the specific

pathogen-free (SPF) animal houses of the Laboratory Animal Center of Xi'an

Jiaotong University.

Mice were genotyped by using PCR and DNA gel electrophoresis. Genomic DNA

from the mouse tail was extracted by using the Mouse Direct PCR Kit (Bimake, TX,

USA, #B40015) according to the manufacturer's instructions. The genotyping primers

were   F3,   5'-TTGAGAGTCTTAAGCAGTCAGGG-3';   R5,   5'-

AACCTTTCCCAGGAGTA TGTGAC-3'. PCR results showed that $Crb3^{wt/wt}$ was one

band with 163 bp, $Crb3^{wt/fl}$ was two bands with 228 bp and 163 bp, and $Crb3^{fl/fl}$ was

one band with 228 bp. The primers for identifying *Cre* alleles were F,

5'-TTGAGAGTCTTAAGCAGTCAGGG-3';   R,

5'-TTACCACTCCCAGCAAGACAC-3', and amplification with one 247 bp band.

The reaction conditions of *Crb3* and *Cre* PCR were as follows: 94°C for 5 min, 94°C

for 20 s, 60°C for 30 s, and 72°C for 20 s for 35 cycles.

Mouse mammary gland whole mount analysis. After mice were euthanized, the

inguinal mammary gland tissues were removed intact and fully spread onto slides.

Tissues were rapidly fixed in Carnoy's fixative for 2 h at room temperature. Then, the

tissues were washed in 70% ethanol solution for 15 min at room temperature and in

50 and 30% ethanol and distilled water for 5 min each. Staining was carried out in

carmine alum solution at 4°C overnight. After washing with 70% ethanol solution,

tissues were washed again in 70, 95 and 100% ethanol for 15 min each. Tissues were

cleared in xylene overnight at room temperature and then mounted with neutral

balsam. Images were obtained by using a microscope (Leica DMi8, Wetzlar,

Germany).

Cell culture and transfection. MCF10A, MCF7, T47D, MDA-MB-231, HCC1806,

T47D, MDA-MB-453 cell lines were purchased from Shanghai Institute of

Biochemistry and Cell Biology (Chinese Academy of Sciences, Shanghai, China).

MCF10A cells were routinely grown in DMEM/F12 (1:1) media (HyClone, UT, USA)

supplemented with 5% horse serum, 20 ng/ml human EGF, 10 μg/ml insulin, and 0.5

μg/ml hydrocortisone. MCF7, MDA-MB-231 and MDA-MB-453 cells were cultured

in high glucose DMEM media (HyClone, UT, USA) supplemented with 10% FBS

(HyClone, UT, USA, #SH30084.03). T47D and HCC1806 cells were grown in

RPMI-1640 media (HyClone, UT, USA) supplemented with 10% FBS. All cell lines

were incubated in 5% CO_2 at 37°C.

siRNAs were purchased from GenePharma company (Shanghai, China). Cultured

cells were transfected with siRNAs or plasmids by using Lipofectamine 2000

(Invitrogen, CA, USA, #11668-019) following the manufacturer's protocol. The

negative control shRNA (NC) and shRNA against CRB3 were packaged into

lentivirus as previously described[5, 6].

DNA constructs and stable cell lines. CRB3 was PCR amplified from RNA

extracted from MCF10A cells and cloned into the pCMV-Blank vector (Beyotime,

Shanghai, China, D2602) using T4 ligase (NEB, MA, USA). Then, the GFP sequence

was inserted between the extracellular and transmembrane CRB3 domains to generate

the pCMV-CRB3-GFP vector as described previously[31]. The full-length CRB3

(1-120), CRB3 (1-116), CRB3 (1-83), CRB3 (1-58) and CRB3 (1-26) were amplified

by PCR from the pCMV-CRB3-GFP vector using specific primers to create EcoR I

and Xho I restriction sites and cloned into the pCMV-Blank vector. Full-length was

cloned into the lentiviral vector pLVX-TetOne-Puro (Clontech, Takara, #631849).

pECMV-3×FLAG-RAB11A was purchased from SinoBiological (Beijing, China).

The plasmids of pECMV-3×FLAG-RAB11A[S20V]/[S25N]/[Q70 L] mutants were

constructed by using the *Fast* Mutagenesis System (Transgen, Beijing, China,

FM111-01).

Lentiviruses were produced in HEK293T cells. CRB3-GFP in MCF10A and MCF7

cells was generated by using a lentivirus expression system. Screening of stable cell

lines used puromycin after lentiviral infection for 72 h.

3D morphogenesis. The growth factor-reduced Matrigel (Corning, USA, #354230)

was added to the four-well chamber slide system (Corning, USA, #177437) and then

incubated at 37°C for solidification. MCF10A cells were plated into this chamber

slide and cultured in DMEM/F12 (1:1) media (HyClone, UT, USA) supplemented

with 2% horse serum, 5 ng/ml human EGF, 10 μg/ml insulin, 0.5 μg/ml

hydrocortisone, and 2.5% Matrigel. 3D morphogenesis was photographed using a

microscope (Leica DMi8, Wetzlar, Germany) at different time points. 3D MCF10A

cells were fixed and stained on day 14 after cell culture.

Immunohistochemistry. The paraffin-embedded sections of samples were baked,

deparaffinized in xylene, and rehydrated sequentially in gradient concentrations of

ethanol, sequentially. Then, antigen retrieval was performed in Tris-EDTA buffer (pH

9.0) heated to 95°C for 20 min. Endogenous peroxidase activity was reduced in 3%

hydrogen peroxide for 10 min at room temperature. Then, 5% goat plasma was used

for blocking at 37°C for 30 min, and the sections were incubated with specific

primary antibodies overnight at 4°C. Ki67 (Abcam, UK, ab279653), phospho-histone

H3 (Abcam, UK, ab267372), cleaved caspase 3 (CST, USA, #9661), GLI1 (CST,

USA, #3538), and β-catenin (CST, USA, #8480) were used as primary antibodies. The following steps used biotin-streptavidin HRP detection kits (ZSGB-BIO, China, SP-9001, SP-9002) according to the manufacturer's instructions. The sections were stained with DAB (ZSGB-BIO, China, ZLI-9017), counterstained with hematoxylin, dehydrated in gradient concentrations of ethanol, cleared in xylene, and mounted with neutral balsam, sequentially. Images were obtained by using a slide scanner (Leica SCN400, Wetzlar, Germany).

Immunofluorescence. Cells were plated on coverslips and fixed in 4% paraformaldehyde for 30 min. Then, the membranes were permeabilized by using 0.2% Triton X-100. The cells were blocked with 5% BSA solution for 1 h at room temperature. Cells were incubated with specific primary antibodies overnight at 4°C. Caspase 3 (CST, USA, #9662S), α-tubulin (CST, USA, #3873), acetylated tubulin (Proteintech, USA, #662001-IG), γ-tubulin (Proteintech, USA, #15176-1-AP), pericentrin (Abcam, UK, ab4448), CRB3 (Sigma, USA, HPA013835), EEA1 (BD, USA, #610456), Rab11 (BD, USA, #610823), and GCP6 (Abcam, UK, ab95172) were used as primary antibodies. The secondary antibodies were Alexa Fluor 488-labeled or 594-labeled (Invitrogen, USA, A32731, A32744). DPIA (5 μg/ml) was used for DNA staining, and images were taken by using a confocal microscope (Leica SP5 II, Wetzlar, Germany).

MEFs isolation and maintenance. The embryos were harvested from $Crb3^{wt/fl}$;CAG-Cre mice crosses at E13.5. Head, tail and all innards were removed from these embryos. DNA could be extracted from the tails for genotyping. Then, the body was minced and digested with trypsin in a 37°C incubator for 30 min. Cells were resuspended in MEF media (DME, 10% FBS, 1% penicillin/streptomycin) and plated on 10 cm cell culture dishes with one embryo. They were passaged 1:3 with MEF media for further expansion.

Real-time PCR assay. Total RNA from cell lines was isolated by using TRIzol reagent (Invitrogen, USA, #15596026), and 5 μg RNA was converted to cDNA with the RevertAid first strand cDNA synthesis kit (Thermo, USA, K1622) according to the manufacturer's instructions. Real-time PCR was prepared with SYBR qPCR

Premix (Takara, Japan, RR420L) and then performed with a real-time PCR detection

instrument (Bio–Rad CFX96, USA). The primers used for real-time PCR were

purchased from Tsingke Biotech (Beijing, China), and the sequences are listed in

Table 1. The mRNA expression was normalized to *GAPDH*, and fold changes were

calculated by using the ΔΔCt method.

Immunoprecipitation and immunoblotting. Cultured cells were lysed in RIPA

buffer supplemented with protease inhibitors (Roche, NJ, USA). Then, the cell lysates

were subjected to SDS–PAGE separation and transferred to PVDF membranes. The

membranes were subjected to immunoblot assays using antibodies against CRB3

(Santa Cruz, USA, sc-292449), GCP2 (Santa Cruz, USA, sc-377117), GCP3 (Santa

Cruz, USA, sc-373758), GCP4 (Santa Cruz, USA, sc-271876), GCP5 (Santa Cruz,

USA, sc-365837), GCP6 (Abcam, UK, ab95172), γ-tubulin (Proteintech, USA,

#66320-1-Ig), Rab11 (Abcam, UK, ab128913), GFP (Roche, USA, #11814460001),

Flag (Sigma, USA, F7425), CEP290 (Santa Cruz, USA, sc-390637), GSK3-β (CST,

USA, #12456), β-catenin (CST, USA, #8480), GAPDH (Proteintech, USA,

#HRP-6004), and β-actin (Proteintech, USA, #HRP-60008). HRP-conjugated

secondary antibodies (CST, USA, #7074, #7076) were incubated at room temperature

for 1 h in the dark. Final detection was detected by ECL Plus (Millipore, Germany,

WBULS0500). Proteins for the immunoprecipitation assay were lysed with NP40

buffer. The cell lysates were detected by a Dynabeads protein G immunoprecipitation

kit (Invitrogen, USA, #10007D) according to the manufacturer's instructions. Elution

protein complexes were subjected to immunoblot assay.

Protein identification and bioinformatics analyses. LC−MS/MS analysis was

conducted by PTM Bio (Zhejiang, China). Q Exactive^{TM} Plus (Thermo, MA, USA)

was used for tandem mass spectrometry data analysis. The Blast2GO program against

the UniProt database was used to analyze the functional annotation and classification

of the identified proteins. Then, pathway enrichment analysis was enriched by using

DAVID tools.

Scanning electron microscope (SEM) observations. CRB3-GFP and CRB3

knockdown MCF10A cells were plated on culture slides, and doxycycline (Dox) was added to induce the expression of CRB3-GFP. After the cells were fully confluent, the slides were fixed with 4% paraformaldehyde at 4°C overnight and then with 1% osmium tetroxide at 4°C for 1 h. Dehydration was performed at room temperature, and the samples were dried using the critical point drying method. Images were obtained by using a scanning electron microscope (Hitachi TM1000, Japan).

Clinical data. Patient data, breast cancer tissues and adjacent para-cancerous tissues were collected from the First Affiliated Hospital of Xi'an Jiaotong University (Shaanxi, China). All patients signed informed consent forms before surgery. This research was authorized by the Ethics Committee of the First Affiliated Hospital of Xi'an Jiaotong University and conducted in conformity with the Declaration of Helsinki.

Statistics. Statistical analysis was performed using SPSS statistics 23.0 for Windows (IBM, Armonk, USA). All experiments were repeated at least three times. The values are expressed as the mean ± standard deviation (SD). Unpaired Student's t test was used to compare the differences between two groups. One-way ANOVA followed by Dunnett's multiple comparisons test was used for multiple comparisons. The χ^2 test was used to assess the significance of the observed frequencies. $P<0.05$ was considered an indicator of statistical significance.

Acknowledgments

This work was supported by grants from the National Natural Science Foundation of China (No. 81872272 and 82173023), Innovation Capability Support Program of Shaanxi (No. 2020TD-046) and Clinical Research Award of the First Affiliated Hospital of Xi'an Jiaotong University (No. XJTU1AF-CRF-2017-007). We are grateful to Prof. Ceshi Chen (Kunming Institute of Zoology, China) and Prof. Yongping Shao (Xi'an Jiaotong University, China) for help in guiding the research. We thank Qi Tian, Lizhe Zhu and Yan Zhou for help with the experiments.

Conflict of interest

The authors declare no conflicts of interest.

Author contributions

P.L. designed the research; B.W., Z.L., T.T, Y.J., Y.S., R.W., H.C., J.L. and P.L.

performed the experimental work, and M.Z., X.G., J.L. and S.S. analyzed the data;

B.W., Z.L., Y.R. and P.L. wrote the original manuscript. P.L. conceived the study and

critically revised the manuscript. All authors read and approved the final manuscript.

Data availability

The datasets used and/or analyzed during the current study are available from the

corresponding author on reasonable request.

References

1.   Wodarz A, Näthke I. Cell polarity in development and cancer. Nature cell biology.

2007;9(9):1016-24. Epub 2007/09/01. doi: 10.1038/ncb433. PubMed PMID: 17762893.

2.   McCaffrey LM, Macara IG. Epithelial organization, cell polarity and tumorigenesis. Trends

in cell biology. 2011;21(12):727-35. Epub 2011/07/26. doi: 10.1016/j.tcb.2011.06.005.

PubMed PMID: 21782440.

3.   Etienne-Manneville S. Polarity proteins in migration and invasion. Oncogene.

2008;27(55):6970-80. Epub 2008/11/26. doi: 10.1038/onc.2008.347. PubMed PMID:

19029938.

4.   Elsum I, Yates L, Humbert PO, Richardson HE. The Scribble-Dlg-Lgl polarity module in

development and cancer: from flies to man. Essays in biochemistry. 2012;53:141-68. Epub

2012/08/30. doi: 10.1042/bse0530141. PubMed PMID: 22928514.

5.   Li P, Wang Y, Mao X, Jiang Y, Liu J, Li J, et al. CRB3 downregulation confers breast

cancer stem cell traits through TAZ/β-catenin. Oncogenesis. 2017;6(4):e322. Epub

2017/04/25. doi: 10.1038/oncsis.2017.24. PubMed PMID: 28436991; PubMed Central PMCID:

PMCPMC5520500.

6.   Mao X, Li P, Wang Y, Liang Z, Liu J, Li J, et al. CRB3 regulates contact inhibition by

activating the Hippo pathway in mammary epithelial cells. Cell death & disease.

2017;8(1):e2546. Epub 2017/01/13. doi: 10.1038/cddis.2016.478. PubMed PMID: 28079891;

PubMed Central PMCID: PMCPMC5386381.

7.   Mao X, Li P, Ren Y, Li J, Zhou C, Yang J, et al. Cell polarity protein CRB3 is an

independent favorable prognostic factor for clear cell renal cell carcinoma. International journal

of oncology. 2015;46(2):657-66. doi: 10.3892/ijo.2014.2763. PubMed PMID:

WOS:000347671200025.

8. Karp CM, Tan TT, Mathew R, Nelson D, Mukherjee C, Degenhardt K, et al. Role of the

polarity determinant crumbs in suppressing mammalian epithelial tumor progression. Cancer

research. 2008;68(11):4105-15. Epub 2008/06/04. doi: 10.1158/0008-5472.Can-07-6814.

PubMed PMID: 18519669; PubMed Central PMCID: PMCPMC2696887.

9. Sánchez I, Dynlacht BD. Cilium assembly and disassembly. Nature cell biology.

2016;18(7):711-7. Epub 2016/06/29. doi: 10.1038/ncb3370. PubMed PMID: 27350441;

PubMed Central PMCID: PMCPMC5079433.

10. Reiter JF, Leroux MR. Genes and molecular pathways underpinning ciliopathies. Nature

reviews Molecular cell biology. 2017;18(9):533-47. Epub 2017/07/13. doi:

10.1038/nrm.2017.60. PubMed PMID: 28698599; PubMed Central PMCID:

PMCPMC5851292.

11. Kim J, Dabiri S, Seeley ES. Primary cilium depletion typifies cutaneous melanoma in situ

and malignant melanoma. PloS one. 2011;6(11):e27410. Epub 2011/11/19. doi:

10.1371/journal.pone.0027410. PubMed PMID: 22096570; PubMed Central PMCID:

PMCPMC3214062.

12. Menzl I, Lebeau L, Pandey R, Hassounah NB, Li FW, Nagle R, et al. Loss of primary cilia

occurs early in breast cancer development. Cilia. 2014;3:7. Epub 2014/07/06. doi:

10.1186/2046-2530-3-7. PubMed PMID: 24987519; PubMed Central PMCID:

PMCPMC4076761.

13. Basten SG, Willekers S, Vermaat JS, Slaats GG, Voest EE, van Diest PJ, et al. Reduced

747 cilia frequencies in human renal cell carcinomas versus neighboring parenchymal tissue. Cilia.

748 2013;2(1):2. Epub 2013/02/02. doi: 10.1186/2046-2530-2-2. PubMed PMID: 23369289;

749 PubMed Central PMCID: PMCPMC3564780.

750 14. Seeley ES, Carrière C, Goetze T, Longnecker DS, Korc M. Pancreatic cancer and

751 precursor pancreatic intraepithelial neoplasia lesions are devoid of primary cilia. Cancer

752 research. 2009;69(2):422-30. Epub 2009/01/17. doi: 10.1158/0008-5472.Can-08-1290.

753 PubMed PMID: 19147554; PubMed Central PMCID: PMCPMC2629528.

754 15. Hassounah NB, Nagle R, Saboda K, Roe DJ, Dalkin BL, McDermott KM. Primary cilia are

755 lost in preinvasive and invasive prostate cancer. PloS one. 2013;8(7):e68521. Epub

756 2013/07/12. doi: 10.1371/journal.pone.0068521. PubMed PMID: 23844214; PubMed Central

757 PMCID: PMCPMC3699526.

758 16. Bazellières E, Aksenova V, Barthélémy-Requin M, Massey-Harroche D, Le Bivic A. Role

759 of the Crumbs proteins in ciliogenesis, cell migration and actin organization. Seminars in cell &

760 developmental biology. 2018;81:13-20. Epub 2017/10/24. doi: 10.1016/j.semcdb.2017.10.018.

761 PubMed PMID: 29056580.

762 17. Fan S, Fogg V, Wang Q, Chen XW, Liu CJ, Margolis B. A novel Crumbs3 isoform

763 regulates cell division and ciliogenesis via importin beta interactions. The Journal of cell

764 biology. 2007;178(3):387-98. Epub 2007/07/25. doi: 10.1083/jcb.200609096. PubMed PMID:

765 17646395; PubMed Central PMCID: PMCPMC2064851.

766 18. Fan S, Hurd TW, Liu CJ, Straight SW, Weimbs T, Hurd EA, et al. Polarity proteins control

767 ciliogenesis via kinesin motor interactions. Current biology : CB. 2004;14(16):1451-61. Epub

768 2004/08/25. doi: 10.1016/j.cub.2004.08.025. PubMed PMID: 15324661.

19.   Whiteman EL, Fan S, Harder JL, Walton KD, Liu CJ, Soofi A, et al. Crumbs3 is essential

for proper epithelial development and viability. Molecular and cellular biology.

2014;34(1):43-56. Epub 2013/10/30. doi: 10.1128/mcb.00999-13. PubMed PMID: 24164893;

PubMed Central PMCID: PMCPMC3911272.

20.   Ye X, Tam WL, Shibue T, Kaygusuz Y, Reinhardt F, Ng Eaton E, et al. Distinct EMT

programs control normal mammary stem cells and tumour-initiating cells. Nature.

2015;525(7568):256-60. Epub 2015/09/04. doi: 10.1038/nature14897. PubMed PMID:

26331542; PubMed Central PMCID: PMCPMC4764075.

21.   Soule HD, Maloney TM, Wolman SR, Peterson WD, Jr., Brenz R, McGrath CM, et al.

Isolation and characterization of a spontaneously immortalized human breast epithelial cell

line, MCF-10. Cancer research. 1990;50(18):6075-86. Epub 1990/09/25. PubMed PMID:

1975513.

22.   Halsne R, Tandberg JI, Lobert VH, Østby GC, Thoen E, Ropstad E, et al. Effects of

perfluorinated alkyl acids on cellular responses of MCF-10A mammary epithelial cells in

monolayers and on acini formation in vitro. Toxicol Lett. 2016;259:95-107. Epub 2016/08/12.

doi: 10.1016/j.toxlet.2016.08.004. PubMed PMID: 27511595.

23.   Jonassen JA, San Agustin J, Follit JA, Pazour GJ. Deletion of IFT20 in the mouse kidney

causes misorientation of the mitotic spindle and cystic kidney disease. The Journal of cell

biology. 2008;183(3):377-84. Epub 2008/11/05. doi: 10.1083/jcb.200808137. PubMed PMID:

18981227; PubMed Central PMCID: PMCPMC2575779.

24.   Martin-Belmonte F, Perez-Moreno M. Epithelial cell polarity, stem cells and cancer.

Nature reviews Cancer. 2011;12(1):23-38. Epub 2011/12/16. doi: 10.1038/nrc3169. PubMed

PMID: 22169974.

25. Lu Q, Insinna C, Ott C, Stauffer J, Pintado PA, Rahajeng J, et al. Early steps in primary

cilium assembly require EHD1/EHD3-dependent ciliary vesicle formation. Nature cell biology.

2015;17(3):228-40. Epub 2015/02/17. doi: 10.1038/ncb3109. PubMed PMID: 25686250;

PubMed Central PMCID: PMCPMC4344897.

26. Westlake CJ, Baye LM, Nachury MV, Wright KJ, Ervin KE, Phu L, et al. Primary cilia

membrane assembly is initiated by Rab11 and transport protein particle II (TRAPPII)

complex-dependent trafficking of Rabin8 to the centrosome. Proceedings of the National

Academy of Sciences of the United States of America. 2011;108(7):2759-64. Epub 2011/01/29.

doi: 10.1073/pnas.1018823108. PubMed PMID: 21273506; PubMed Central PMCID:

PMCPMC3041065 R.H.S., and P.K.J. are employees of Genentech, Inc.

27. Kollman JM, Merdes A, Mourey L, Agard DA. Microtubule nucleation by γ-tubulin

complexes. Nature reviews Molecular cell biology. 2011;12(11):709-21. Epub 2011/10/14. doi:

10.1038/nrm3209. PubMed PMID: 21993292; PubMed Central PMCID: PMCPMC7183383.

28. Liu P, Zupa E, Neuner A, Böhler A, Loerke J, Flemming D, et al. Insights into the

assembly and activation of the microtubule nucleator γ-TuRC. Nature. 2020;578(7795):467-71.

Epub 2019/12/20. doi: 10.1038/s41586-019-1896-6. PubMed PMID: 31856152.

29. Farache D, Jauneau A, Chemin C, Chartrain M, Rémy MH, Merdes A, et al. Functional

Analysis of γ-Tubulin Complex Proteins Indicates Specific Lateral Association via Their

N-terminal Domains. The Journal of biological chemistry. 2016;291(44):23112-25. Epub

2016/10/30. doi: 10.1074/jbc.M116.744862. PubMed PMID: 27660388; PubMed Central

PMCID: PMCPMC5087730.

30. Knödler A, Feng S, Zhang J, Zhang X, Das A, Peränen J, et al. Coordination of Rab8 and

Rab11 in primary ciliogenesis. Proceedings of the National Academy of Sciences of the United

States of America. 2010;107(14):6346-51. Epub 2010/03/24. doi: 10.1073/pnas.1002401107.

PubMed PMID: 20308558; PubMed Central PMCID: PMCPMC2851980.

31. Djuric I, Siebrasse JP, Schulze U, Granado D, Schlüter MA, Kubitscheck U, et al. The

C-terminal domain controls the mobility of Crumbs 3 isoforms. Biochimica et biophysica acta.

2016;1863(6 Pt A):1208-17. Epub 2016/03/16. doi: 10.1016/j.bbamcr.2016.03.008. PubMed

PMID: 26975581.

32. Craige B, Tsao CC, Diener DR, Hou Y, Lechtreck KF, Rosenbaum JL, et al. CEP290

tethers flagellar transition zone microtubules to the membrane and regulates flagellar protein

content. The Journal of cell biology. 2010;190(5):927-40. Epub 2010/09/08. doi:

10.1083/jcb.201006105. PubMed PMID: 20819941; PubMed Central PMCID:

PMCPMC2935561.

33. Casenghi M, Meraldi P, Weinhart U, Duncan PI, Körner R, Nigg EA. Polo-like kinase 1

regulates Nlp, a centrosome protein involved in microtubule nucleation. Developmental cell.

2003;5(1):113-25. Epub 2003/07/11. doi: 10.1016/s1534-5807(03)00193-x. PubMed PMID:

12852856.

34. Li P, Feng C, Chen H, Jiang Y, Cao F, Liu J, et al. Elevated CRB3 expression suppresses

breast cancer stemness by inhibiting β-catenin signalling to restore tamoxifen sensitivity.

Journal of cellular and molecular medicine. 2018;22(7):3423-33. Epub 2018/03/31. doi:

10.1111/jcmm.13619. PubMed PMID: 29602199; PubMed Central PMCID:

PMCPMC6010813.

35. Thiery JP, Acloque H, Huang RY, Nieto MA. Epithelial-mesenchymal transitions in

development and disease. Cell. 2009;139(5):871-90. Epub 2009/12/01. doi:

10.1016/j.cell.2009.11.007. PubMed PMID: 19945376.

36. Varelas X, Samavarchi-Tehrani P, Narimatsu M, Weiss A, Cockburn K, Larsen BG, et al.

The Crumbs complex couples cell density sensing to Hippo-dependent control of the

TGF-β-SMAD pathway. Developmental cell. 2010;19(6):831-44. Epub 2010/12/15. doi:

10.1016/j.devcel.2010.11.012. PubMed PMID: 21145499.

37. Charrier LE, Loie E, Laprise P. Mouse Crumbs3 sustains epithelial tissue morphogenesis

in vivo. Scientific reports. 2015;5:17699. Epub 2015/12/04. doi: 10.1038/srep17699. PubMed

PMID: 26631503; PubMed Central PMCID: PMCPMC4668553.

38. Herranz-Martín S, Jimeno D, Paniagua AE, Velasco A, Lara JM, Aijón J, et al.

Immunocytochemical evidence of the localization of the Crumbs homologue 3 protein (CRB3)

in the developing and mature mouse retina. PloS one. 2012;7(11):e50511. Epub 2012/12/12.

doi: 10.1371/journal.pone.0050511. PubMed PMID: 23226298; PubMed Central PMCID:

PMCPMC3511585.

39. Arya P, Rainey MA, Bhattacharyya S, Mohapatra BC, George M, Kuracha MR, et al. The

endocytic recycling regulatory protein EHD1 Is required for ocular lens development.

Developmental biology. 2015;408(1):41-55. Epub 2015/10/13. doi:

10.1016/j.ydbio.2015.10.005. PubMed PMID: 26455409; PubMed Central PMCID:

PMCPMC4688215.

40. Rainey MA, George M, Ying G, Akakura R, Burgess DJ, Siefker E, et al. The endocytic

recycling regulator EHD1 is essential for spermatogenesis and male fertility in mice. BMC Dev

Biol. 2010;10:37. Epub 2010/04/03. doi: 10.1186/1471-213x-10-37. PubMed PMID: 20359371;

PubMed Central PMCID: PMCPMC2856533.

41. Kobayashi T, Dynlacht BD. Regulating the transition from centriole to basal body. The

Journal of cell biology. 2011;193(3):435-44. Epub 2011/05/04. doi: 10.1083/jcb.201101005.

PubMed PMID: 21536747; PubMed Central PMCID: PMCPMC3087006.

42. Kobayashi T, Kim S, Lin YC, Inoue T, Dynlacht BD. The CP110-interacting proteins

Talpid3 and Cep290 play overlapping and distinct roles in cilia assembly. The Journal of cell

biology. 2014;204(2):215-29. Epub 2014/01/15. doi: 10.1083/jcb.201304153. PubMed PMID:

24421332; PubMed Central PMCID: PMCPMC3897186.

43. Schmidt KN, Kuhns S, Neuner A, Hub B, Zentgraf H, Pereira G. Cep164 mediates

vesicular docking to the mother centriole during early steps of ciliogenesis. The Journal of cell

biology. 2012;199(7):1083-101. Epub 2012/12/21. doi: 10.1083/jcb.201202126. PubMed

PMID: 23253480; PubMed Central PMCID: PMCPMC3529528.

44. Das A, Guo W. Rabs and the exocyst in ciliogenesis, tubulogenesis and beyond. Trends

in cell biology. 2011;21(7):383-6. Epub 2011/05/10. doi: 10.1016/j.tcb.2011.03.006. PubMed

PMID: 21550243; PubMed Central PMCID: PMCPMC3128673.

45. Guichard P, Laporte MH, Hamel V. The centriolar tubulin code. Seminars in cell &

developmental biology. 2021. Epub 2021/12/14. doi: 10.1016/j.semcdb.2021.12.001. PubMed

PMID: 34896019.

46. Liu P, Würtz M, Zupa E, Pfeffer S, Schiebel E. Microtubule nucleation: The waltz between

γ-tubulin ring complex and associated proteins. Current opinion in cell biology. 2021;68:124-31.

Epub 2020/11/16. doi: 10.1016/j.ceb.2020.10.004. PubMed PMID: 33190097.

47. Chen HY, Kelley RA, Li T, Swaroop A. Primary cilia biogenesis and associated retinal

ciliopathies. Seminars in cell & developmental biology. 2021;110:70-88. Epub 2020/08/05. doi:

10.1016/j.semcdb.2020.07.013. PubMed PMID: 32747192; PubMed Central PMCID:

PMCPMC7855621.

48. Drivas TG, Bennett J. CEP290 and the primary cilium. Advances in experimental

medicine and biology. 2014;801:519-25. Epub 2014/03/26. doi:

10.1007/978-1-4614-3209-8_66. PubMed PMID: 24664739.

49. Tsang WY, Bossard C, Khanna H, Peränen J, Swaroop A, Malhotra V, et al. CP110

suppresses primary cilia formation through its interaction with CEP290, a protein deficient in

human ciliary disease. Developmental cell. 2008;15(2):187-97. Epub 2008/08/13. doi:

10.1016/j.devcel.2008.07.004. PubMed PMID: 18694559; PubMed Central PMCID:

PMCPMC3987787.

50. Chang B, Khanna H, Hawes N, Jimeno D, He S, Lillo C, et al. In-frame deletion in a novel

centrosomal/ciliary protein CEP290/NPHP6 perturbs its interaction with RPGR and results in

early-onset retinal degeneration in the rd16 mouse. Human molecular genetics.

2006;15(11):1847-57. Epub 2006/04/25. doi: 10.1093/hmg/ddl107. PubMed PMID: 16632484;

PubMed Central PMCID: PMCPMC1592550.

51. Kim J, Krishnaswami SR, Gleeson JG. CEP290 interacts with the centriolar satellite

component PCM-1 and is required for Rab8 localization to the primary cilium. Human

molecular genetics. 2008;17(23):3796-805. Epub 2008/09/06. doi: 10.1093/hmg/ddn277.

PubMed PMID: 18772192; PubMed Central PMCID: PMCPMC2722899.

52. Stowe TR, Wilkinson CJ, Iqbal A, Stearns T. The centriolar satellite proteins Cep72 and

Cep290 interact and are required for recruitment of BBS proteins to the cilium. Molecular

biology of the cell. 2012;23(17):3322-35. Epub 2012/07/07. doi: 10.1091/mbc.E12-02-0134.

PubMed PMID: 22767577; PubMed Central PMCID: PMCPMC3431927.

53. Wheway G, Nazlamova L, Hancock JT. Signaling through the Primary Cilium. Frontiers in

cell and developmental biology. 2018;6:8. Epub 2018/02/24. doi: 10.3389/fcell.2018.00008.

PubMed PMID: 29473038; PubMed Central PMCID: PMCPMC5809511.

54. Wong SY, Seol AD, So PL, Ermilov AN, Bichakjian CK, Epstein EH, Jr., et al. Primary cilia

can both mediate and suppress Hedgehog pathway-dependent tumorigenesis. Nature

medicine. 2009;15(9):1055-61. Epub 2009/08/25. doi: 10.1038/nm.2011. PubMed PMID:

19701205; PubMed Central PMCID: PMCPMC2895420.

55. Zhan T, Rindtorff N, Boutros M. Wnt signaling in cancer. Oncogene. 2017;36(11):1461-73.

Epub 2016/09/13. doi: 10.1038/onc.2016.304. PubMed PMID: 27617575; PubMed Central

PMCID: PMCPMC5357762.

914
915

Figure 1. *Crb3* knockout mice exhibit smaller sizes and ocular abnormalities, and

mammary epithelial cell-specific *Crb3* knockout leads to ductal epithelial hyperplasia

and promotes tumorigenesis.

A, B. Representative whole bodies (A) and eyes (B) from littermate *Crb3*$^{fl/fl}$,

Crb3$^{wt/fl}$;CAG-*Cre* and *Crb3*$^{fl/fl}$;CAG-*Cre* mice at 4 weeks old. C. Representative

mammary whole mounts from littermate *Crb3*$^{fl/fl}$ and *Crb3*$^{fl/fl}$;MMTV-*Cre* mice at 8

weeks old with Carmine-alum staining. (scale bars, 200 μm) D, E. Quantification of

the average number of TEBs (n=10) and bifurcated TEBs (n=10) in littermate *Crb3*$^{fl/fl}$

and *Crb3*$^{fl/fl}$;MMTV-*Cre* mice at 8 weeks old. F. Representative images of mammary

glands in littermate *Crb3*$^{fl/fl}$ and *Crb3*$^{fl/fl}$;MMTV-*Cre* mice stained with H&E. (scale

bars, 50 μm) G. Representative images of primary tumors stained with H&E in

PyMT-WT and PyMT-cKO-*Crb3* mice at 9 weeks old. (scale bars, left 500 μm, right

50 μm) Magnified areas of boxed sections are shown in the right panels. Bars

represent the means ± SD; Unpaired Student's *t* test, ***$P<0.001$.

Figure 2. CRB3 knockdown inhibits acinar formation of mammary epithelial cells in

a 3D culture system.

A. Representative effect of CRB3 on acinar formation in the 3D culture system at

days 3, 6, 9 and 14. (Magnified areas of marked arrows are shown in the lower right

corner) B-D. Quantification of the average number, diameter and aberration of acini

(n=10). E. Immunofluorescence showing apoptosis during lumen formation. Caspase

3 (red), α-tubulin (green), and DNA (blue). F. Immunofluorescence showing the

mitotic spindle orientation during lumen formation. α-Tubulin (green) and DNA

(blue). G, H. Quantification of division angle. I. Immunohistochemical analyses of

Ki67, phospho-histone H3 and cleaved caspase 3 in primary tumors from PyMT-WT

and PyMT-cKO-*Crb3* mice at 9 weeks old. (positive cells marked by arrows) Scale

bars, 25 μm, bars represent means ± SD; Unpaired Student's *t* test, * $P<0.5$, **

$P<0.01$, ***$P<0.001$.

Figure 3. CRB3 alters primary cilium formation in mammary cells, mammary ductal

lumen and renal tubule from *Crb3*^{fl/fl};CAG-*Cre* mice.

A. C. Representative images of immunofluorescent staining of primary cilium

formation with CRB3 knockdown in MCF10A cells and CRB3 conditional

overexpression upon dox induction in MCF7 cells. Acetylated tubulin (red), γ-tubulin

(green), and DNA (blue). (primary cilium marked by arrows; scale bars, 10 μm) B. D.

Quantification of the proportion of cells with primary cilium formation and the length

of the primary cilium (n=10). E. Representative scanning electron microscope images

of primary cilium formation with CRB3 knockdown and conditional overexpression

in MCF10A cells. (primary cilium marked by arrows; scale bars, 50 μm) F.

Quantification of the proportion of cells with primary cilium formation (n=10). G. H.

Representative immunofluorescent staining of primary cilium formation in the

mammary ductal lumen and renal tubule from *Crb3*^{fl/fl} and *Crb3*^{fl/fl};CAG-*Cre* mice,

respectively. Acetylated tubulin (red), γ-tubulin (green), and DNA (blue). (n=10; scale

bars, 25 μm) Bars represent the means ± SD; Unpaired Student's *t* test, *$P<0.05$,

***$P<0.001$.

Figure 4. CRB3 localizes to the basal body of the primary cilium.

A. Immunofluorescence showing the colocalization of exogenous CRB3 with

centrosomes in MCF10A cells. Pericentrin, a marker of centrosome (red), CRB3-GFP

(green), and DNA (blue). (Colocalization marked by arrows; scale bars, 10 μm) B.

Quantification of the proportion of cells with pericentrin and exogenous CRB3

colocalization (n=10). (bars represent the means ± SD) C. Another colocalization of

endogenous CRB3 with the basal body in MCF10A cells. γ-Tubulin is a marker of the

centrosome and basal body of the primary cilium (green), CRB3 (red), and DNA

(blue). (Colocalization marked by arrows; scale bars, 10 μm) D. Corresponding

fluorescence intensity profile across a section of the array, as indicated by the dashed

white line in (C). E. Double immunostaining displaying the colocalization of CRB3

with the primary cilium in MCF10A cells. Acetylated tubulin (green), CRB3 (red),

and DNA (blue). (Colocalization marked by arrows; scale bars, 10 μm) F.

Fluorescence 3D reconstruction of CRB3 and primary cilium colocalization.

Acetylated tubulin (red), CRB3 (green), and DNA (blue).

Figure 5. CRB3 trafficking is mediated by Rab11-positive endosomes, and CRB3

knockdown destabilizes γTuRC assembly during ciliogenesis.

A. Pathway aggregation analysis of CRB3 binding proteins identified by mass

spectrometry in MCF10A cells. B. Table of some Rab small GTPase family members

and centriolar proteins identified as CRB3 binding proteins. C. Immunofluorescence

showing the colocalization of CRB3 with EEA1-, CD63-, and Rab11-positive

endosomes in MCF10A cells. EEA1, CD63, Rab11 (green), CRB3 (red), and DNA

(blue). (scale bars, 10 μm) D. Quantification of the proportion of cells with these

markers and CRB3 colocalization (n=10). E. Western blots showing the levels of

CRB3 in MCF10A cells treated with dynasore at different cell densities. F. Structure

diagram of γTuRC (γ-tubulin ring complex). G. Immunoblot analysis of the effect of

CRB3 on γTuRC molecules in MCF10A cells. H. Coimmunoprecipitation showing

the interacting proteins with GCP6 in MCF10A cells with CRB3 knockdown. I.

Representative images of immunofluorescent staining of GCP3 and GCP6

colocalization in MCF10A cells with the corresponding fluorescence intensity profile.

GCP3 (red), GCP6 (green), and DNA (blue). (scale bars, 10 μm) J. Comparison of

cytoplasmic extracts from MCF10A cells and cells with CRB3 knockdown after

fractionation in sucrose gradients. The γTuSC sedimentation was mainly in fractions 3,

and γTuRC sedimentation was mainly in fractions 6. Bars represent the means ± SD;

Unpaired Student's *t* test, ***P<0.001.

Figure 6. CRB3 directly interacts with Rab11.

A. Coimmunoprecipitation of CRB3 with Rab11, GCP6 and GCP3 in MCF10A cells.

B. Coimmunoprecipitation of Rab11 with CRB3, GCP6 and GCP3 in MCF10A cells.

C. Coimmunoprecipitation of Rab11 with GCP6 in control and CRB3 knockdown

MCF10A cells. D. Schematic diagram of CRB3 domains. E. Diagram truncations of

CRB3-GFP fusion proteins with serial C-terminal deletions. F. Domain mapping of

CRB3-GFP for Flag-Rab11a binding. Flag antibody co-IP of the full-length

CRB3-GFP and truncations of CRB3-GFP with Flag-Rab11a were cotransfected into

HEK293 cells for 48 h. Immunoblot analysis was performed using GFP and Flag

antibodies. G. Coimmunoprecipitation of Rab11 mutant variants with full-length

CRB3-GFP. Flag antibody co-IP of the full-length CRB3-GFP with Flag-Rab11aWT,

Flag-Rab11a[Q70L], Flag-Rab11a[S20V] and Flag-Rab11a[S25N] were cotransfected

into HEK293 cells for 48 h. Immunoblot analysis was performed using GFP and Flag

antibodies. Bars represent the means ± SD; Unpaired Student's t test, *P<0.05.

1013

Figure 7. CRB3 navigates GCP6/Rab11 trafficking vesicles to the basal body of the

primary cilium.

A. Coimmunoprecipitation of exogenous CRB3 with Rab11, GCP6, GCP3 and

CEP290 in MCF10A cells. B. Coimmunoprecipitation of Rab11 with exogenous

CRB3, GCP6, GCP3 and CEP290 in MCF10A cells. C. Coimmunoprecipitation of

Rab11 with GCP6 and CEP290 in control and CRB3 knockdown MCF10A cells. D.

Representative immunofluorescent images of GCP6 and γ-tubulin colocalization of

the basal body foci in MCF10A cells with CRB3 knockdown. γ-Tubulin (green),

GCP6 (red), and DNA (blue). (Colocalization marked by arrows; scale bars, 25 μm) E.

Representative immunofluorescent images of Rab11 and γ-tubulin colocalization of

the basal body foci in MEF cells from $Crb3^{fl/fl}$ and $Crb3^{fl/fl}$;CAG-Cre mice. γ-tubulin

(green), Rab11 (red), and DNA (blue). Foci marked by arrows; scale bars, 25 μm.

Bars represent the means ± SD; Unpaired Student's t test, ***P<0.001.

1027

Figure 8. Defects in CRB3 expression inhibit ciliary assembly in breast cancer tissues,

and activate the Wnt signaling pathway in mammary cells and PyMT mouse model.

A. Representative immunofluorescent images of CRB3 in breast cancer tissues (n=50).

CRB3 (green) and DNA (blue). (scale bars, 25 μm) B. Representative

immunofluorescent images of the primary cilium in breast cancer tissues (n=50).

Acetylated tubulin (red), γ-tubulin (green), and DNA (blue). (scale bars, 10 μm) C.

Quantification of the proportion of cells with primary cilium formation in breast

cancer tissues (n=50). D. Representative immunofluorescent images of CRB3 and

γ-tubulin colocalization in adjacent paracarcinoma tissues. CRB3 (green), γ-tubulin (red), and DNA (blue). (scale bars, 25 μm) E. Quantification of the proportion of MFC10A cells with SMO translocation after CRB3 knockdown (n=6). F. Real-time quantitative PCR showing the relative mRNA expression of GLI1 upon SAG treatment in CRB3-depleted MFC10A cells (n=6). G. Immunoblot analyses of the effect of CRB3 on the molecules of the Wnt signaling pathway in mammary cells. H. I. Immunohistochemical analyses of GLI1 and β-catenin in primary tumors from PyMT-WT and PyMT-cKO-*Crb3* mice at 9 weeks old, respectively. (scale bars, 25 μm) Bars represent the means ± SD; Unpaired Student's *t* test, ***$P<0.001$.

Figure 9. Schematic model of CRB3 regulating ciliary assembly.

Graphic summary of prominent phenotypes observed after CRB3 deletion. CRB3 is localized on apical epithelial surfaces and participates in tight junction formation to maintain contact inhibition and cell homeostasis in quiescence. Inside the cell, Rab11-positive endosomes mediate the intracellular trafficking of CRB3, and CRB3 navigates GCP6/Rab11 trafficking vesicles to CEP290. Then, GCP6 is involved in normal γTuRC assembly in ciliogenesis. In CRB3 deletion cells, the primary cilium does not assemble properly, and the Wnt signaling pathway is activated through β-catenin upregulation and nuclear localization, but the Hh signaling pathway fails to be activated. This cellular imbalance is disrupted, leading to tumorigenesis. γTuRC, γ-tubulin ring complex; TJ, tight junction; EE, early endosome; LE, late endosome.

Supplementary Figure 1. Generation of *Crb3* knockout mice by using the Cre-loxP

system.

A. Schematic representation of the strategy for loxP sequence insertion and PCR

genotyping. The loxP sites (gray arrow) flanked either side of exon 3 (orange) in the

Crb3 gene. The primer pair F3&R5 and F4&R6 were used to distinguish the alleles. B.

Southern blot analysis of a positive ES cell clone confirmed homologous

recombination. The genomic DNA from ES cell clones and wild-type cells was

digested with Avr II or Sac I and hybridized with a 5' external probe or 3' internal

probe. C. PCR identification of mouse genotyping showing $Crb3^{wt/wt}$, $Crb3^{wt/fl}$ and

$Crb3^{fl/fl}$ alleles. D. Crb3 expression was detected by immunoblotting in mammary

gland tissues in $Crb3^{fl/fl}$ and $Crb3^{fl/fl}$;MMTV-*Cre* mice at 8 weeks old. E. Real-time

quantitative PCR showing the relative mRNA expression of *Crb3* in mammary gland

tissues (n=10). F. Crb3 expression was assessed by using immunohistochemistry in

mammary gland tissues. The mammary epithelial cells were marked by areas of white

dotted lines. (scale bars, 50 μm) Bars represent the means ± SD; Unpaired Student's *t*

test, ***P<0.001.

1074

Supplementary Figure 2. CRB3 knockdown promotes proliferation of mammary

epithelial cells, and overexpression promotes apoptosis of breast cancer.

A. B. Immunoblot and real-time quantitative PCR analysis of CRB3 knockout

efficiency in MCF10A cells. C. MCF10A proliferation assay for six successive days.

D. Cell cycle showing the distribution of MCF10A cells. E. Quantification of the cell

cycle distribution. F.G.H. Apoptosis of MCF7, T47D, and MDA-MB-453 breast

cancer cells with CRB3 overexpression and quantification of apoptotic cells. Bars

represent the means ± SD; Unpaired Student's *t* test, * *P*<0.05, **P<0.01.

1083

1084 **Supplementary Figure 3.** CRB3 alters primary cilium formation in MEFs from

1085 $Crb3^{fl/fl}$;CAG-*Cre* mice.

1086 A. Representative immunofluorescent staining of primary cilium formation in MEFs

1087 from $Crb3^{fl/fl}$ and $Crb3^{fl/fl}$;CAG-*Cre* mice. Acetylated tubulin (red), γ-tubulin (green),

and DNA (blue). (scale bars, 10 μm) B. Quantification of the proportion of cells with primary cilium formation (n=10). Bars represent the means ± SD; Unpaired Student's *t* test, ***P*<0.001.

Supplementary Figure 4. The effect of CRB3 on a series of ciliogenesis-related genes and Rab11 expression.

A. Quantification of the mRNA expression of genes encoding markers of centriole or primary cilium, BB components, IFT-A and IFT-B anterograde transport (normalized to GAPDH). B. Coomassie blue staining showing the gel electrophoresis of exogenous CRB3 immunoprecipitated precipitates. C. Immunoblot analysis of the effect of CRB3 on Rab11 in MCF10A cells. Bars represent the means ± SD; Unpaired Student's *t* test

Supplementary Figure 5. CRB3 knockdown disturbed the colocalization of GCP6 and Rab11, and Rab11 knockdown caused primary cilium defects.

A. Representative immunofluorescent staining of the colocalization between Rab11 and GCP6 in MCF10A cells. GCP6 (red), Rab11 (green), and DNA (blue). (scale bars, 20 μm) B. Fluorescence colocalization analysis in (A). C. Quantification of the proportion of cells with Rab11 and GCP6 colocalization. D. Representative images of immunofluorescent staining of primary cilium formation with Rab11 knockdown in MCF10A cells. Acetylated tubulin (red), γ-tubulin (green), and DNA (blue). (primary cilium marked by arrows; scale bars, 20 μm) E. Quantification of the proportion of cells with primary cilium formation. F. Immunoblot analysis of Rab11 knockout efficiency in MCF10A cells. Bars represent the means ± SD; Unpaired Student's *t* test, ***P*<0.01, ***P*<0.001.

A

Crb3$^{fl/fl}$ Crb3$^{wt/fl}$;CAG-Cre   Crb3$^{fl/fl}$;CAG-Cre

B

Crb3$^{fl/fl}$ Crb3$^{wt/fl}$;CAG-Cre   Crb3$^{fl/fl}$;CAG-Cre

C

Crb3$^{fl/fl}$

Crb3$^{fl/fl}$;MMTV-Cre

D

TEBs/gland (N)

***

Crb3$^{fl/fl}$
Crb3$^{fl/fl}$;MMTV-Cre

E

Bifurcated TEBs (N)

***

Crb3$^{fl/fl}$
Crb3$^{fl/fl}$;MMTV-Cre

F

Crb3$^{fl/fl}$ Crb3$^{fl/fl}$;MMTV-Cre

G

PyMT-WT   PyMT-cKO-Crb3

A

GFP — 40
Rab11 — 24
GCP6 { short time / long time — 200
GCP3 — 104
CEP290 — 290
IgG — 55

B

Rab11 — 24
GFP — 40
GCP6 — 200
GCP3 — 104
CEP290 — 290
IgG — 55

C

GCP6 — 200
CEP290 — 290
IgG — 55

D

MCF10A

NC   shCRB3

γ-tubulin/GCP6/DAPI

E

MEFs

CRB3^fl/fl   CRB3^fl/fl;CAG-Cre

γ-tubulin/Rab11/DAPI