Title: Selective loss of specific subsets of hair cell and spiral ganglion cell types in an outbred mouse model of age-related hearing loss

Authors: Ely Cheikh Boussaty[1*], Neil Tedeschi[2*], Mark Novotny[2], Yuzuru Ninoyu[1], Eric Du[1], Clara Draf[1], Yun Zhang[2], Richard H. Scheuermann[2,3†], Rick Friedman[1†]

Affiliations:
[1]Department of Otolaryngology, University of California, San Diego, CA
[2]J. Craig Venter Institute, La Jolla, CA
[3]Department of Pathology, University of California, San Diego, CA

*Contributed equally
[†]Contributed equally; to whom correspondence should be addressed: rscheuermann@jcvi.org, rafriedman@health.ucsd.edu

Key words: single nucleus RNA sequencing, cell ontology, marker genes, age-related hearing loss, genetic determinants of hearing loss

Abstract: Age☐related hearing loss (ARHL) is the most common cause of hearing loss and is one of the most prevalent conditions affecting the elderly globally. Despite evidence from our lab and others about its polygenic nature, little is known about the genes, cell types and pathways involved, preventing the development of therapeutic intervention. In this manuscript we describe, for the first time, the complete cell-type specific cochlear transcriptome of the aging mouse cochlea in outbred mice, using snRNA-seq, in relation to auditory threshold variation. We annotated cochlear cell types identified using unsupervised clustering via a three-tiered approach. First, we annotated clusters by linking to expression of known marker genes. Second, we used the NS-Forest algorithm to select minimum cluster-specific marker genes, which were first used to reduce dimensional feature space for statistical comparison of our clusters with existing publicly available data sets on the gEAR website (https://umgear.org/). Finally, we validated and refined our annotation using Multiplexed Error Robust Fluorescence In Situ Hybridization (MERFISH) and the cluster-specific marker genes as probes. We report on 60 unique cell-types (clusters) expanding the number of defined cochlear cell types by more than two times. Additionally, we show cluster variation with hearing thresholds demonstrating specific cell type increases and decrease in response to hearing loss implicating hair cell (HC) subtypes, ganglion cell subtypes, and cell subtypes withing the stria vascularis in this model of ARHL. These results provide a view into the cellular and molecular mechanisms responsible for age-related hearing loss and pathways for therapeutic targeting.

Introduction:

Age-related hearing loss (ARHL), also known as presbycusis, is the most common cause of hearing loss and one of the most prevalent conditions affecting the global population. Progressive, bilateral sensorineural hearing impairment affects approximately 25% of people aged 65-74 and 50% aged 75 and older. In the United States, ARHL occurs in two thirds of people over 70 years of age and is a risk factor for depression and cognitive decline[2, 6-7]. Aside from the detrimental impact on quality of life, presbycusis also carries an increasing economic burden as the cost of ARHL medical expenditures is expected to reach $60 billion in 2030. It is projected that 2.45 billion people will have hearing loss by 2050, a 56.1% increase from 2019, despite stable age-standardized prevalence[8]. Notably, a substantial fraction of patients with ARHL have no identifiable mutation in any known hearing loss gene among the 100+ that have been identified, suggesting that a significant fraction of hearing loss is due to unidentified monogenic or polygenic causes[9].

As part of a broad approach to studying the genetic landscape of ARHL, we have begun a large-scale effort to phenotypically characterize the auditory function of young and aged Crl:CFW(SW)-US_P08 (hereafter CFW) outbred mice[10]. Genome-wide association analyses and genome-wide exome studies have provided some insights into the genetics of ARHL in both mice and humans[11-13]. Another powerful approach to the identification of candidate genes and pathways has been the use of single cell/nucleus transcriptomics. This approach has provided insights into the molecular mechanisms underlying cochlear development[14] and response to damaging noise[15-16]. Recently, Li et al. looked at the transcriptome of inner and outer hair cells (OHCs) and the stria vascularis in aging CBA/J mice, implicating changes in several processes including gene transcription, DNA damage, autophagy and oxidative stress[17].

Developing therapies for ARHL requires an understanding of the genes and pathways involved. The cellular complexity of the inner ear, including the sensory and supporting cells of the organ of Corti, the lateral wall (stria vascularis) and the auditory neurons, necessitates a single cell approach to gain an understanding of the pathway changes associated with ARHL. Due to this complexity, a few laboratories have used single-cell RNA sequencing (scRNA-seq) to characterize the inner ear transcriptome during development, aging in a single mouse strain, and after acoustic trauma[18-24].

Single cell analyses of the numerous cell types of the adult mouse inner ear, particularly the organ of Corti, have been hampered by the difficulty of dissociating the cells due to their actin cytoskeleton and tight junctions[25]. In this study we used single nuclear RNA-seq (snRNA-seq) across 48 genetically diverse outbred mice at 10 months of age to provide for a more unbiased representation of cell types and were able to identify gene expression signatures for various cell types withing the cochlea, including novel markers for inner and outer HCs. Furthermore, to our knowledge this is the first snRNA-seq study to examine differential gene expression across all cochlear cell types in genetically diverse outbred mice with varying degrees of hearing loss.

Results:

Carworth Farms White (CFW) mice are a genetically diverse outbred mouse population distinguished by degrading linkage disequilibrium (LD) between nearby alleles and shorter LD ranges compared to other commercially available inbred mouse strains[26-27], which makes them an ideal model to capture high-resolution mapping in genome-wide association studies (GWAS). Our ongoing efforts to phenotype the hearing function in CFW mice have categorized

them into eight distinct patterns of hearing, ranging from normal hearing (Figure 1-A); to moderate mid- (Figure 1-B) or high-frequency (Figure 1-C) hearing loss; to severe high-frequencies (Figure 1-D, E) hearing loss; to a moderate (Figure 1-F) , or severe (Figure 1-G), or profound (Figure 1-H) all-frequencies hearing loss, with each of those hearing patterns worsening with age.

A group of 48 mice were selected to represent those hearing patterns (Figure 1-I), based on their proportion in the general CFW cohort, to explore the molecular and cellular correlates of hearing loss using single nucleus RNA sequencing (snRNA-seq) of dissected cochlea. During the dissection of inner ear tissue, special care was given to quickly process the samples while avoiding any shear force. The vestibular parts of the inner ear, including utricle, saccule, and semicircular canal ampulla, were removed before collecting cochlear tissue. snRNA-seq processing was chosen over scRNA-seq to avoid the stress responses induced during the single cell processing procedure and to provide for a more unbiased representation of cell types from solid tissues[28] .

Iterative unsupervised clustering was used to group snRNA-seq transcriptional profiles into transcriptome clusters. The initial Leiden community detection yielded 26 clusters, a subset of which (13) were processed through a second round of sub-clustering based on Silhouette score and manual inspection to yield a final collection of 60 transcriptome clusters (Figure 2A). The similarity relationships between the clusters were determined using hierarchical clustering (Figure 2B) and the relative abundance of each putative cell type determined (Figure 2C). The hierarchical clustering results confirmed the transcriptional relationships between the initial clusters and the common origin of sub-clusters from the second round of sub-clustering.

As an initial approach toward determining which cell types correspond to these transcriptomic clusters, we examined the expression levels of several genes known to be expressed in specific cochlear cell types (Figure 2B). The organ of Corti supporting cell gene Epyc[29]   is uniquely expressed in the two related Clusters 10.1 and 10.2. The inner ear progenitor gene Lgr5[30]   is uniquely expressed in Cluster 6. The spindle-root gene Slc26a4 is uniquely expressed in the two related Clusters 19.1 and 19.2[31]   . The Kcnq1 marginal cell gene and the Slc26a7 Reissner's membrane gene are preferentially expressed in Cluster 5 and Cluster 4, respectively. The hematopoietic stem cell-related gene Runx1[32]   is expressed in twelve related clusters. The intermediate cell genes Kcnj10[33]   and Met[34]   are uniquely and preferentially expressed in Cluster 9. The glial and Schwann's cell genes Mpz, Mbp, and Plp1 are expressed in the related Clusters 25, 17.1, 17.2, and 18. The HC genes Xirp2[35]   and Cabp2[36]   are expressed in related Clusters 23.1, 23.2, 23.3, and 23.4. The spiral ganglion neuron (SGN) genes Slc4a4[37] and Grm8[38]   are expressed in related Clusters 12.1 and 12.2. The basal cell gene Cldn11[39] is expressed in related Clusters 2.1 and 2.2. Finally, the supporting fibrocyte genes Slco1c1[40] and Cald1[41]   are expressed in seventeen related clusters.

While the expression patterns of known marker genes was useful in providing an initial annotation and validation of the snRNA-seq analysis, there were many examples where finer grained distinctions were identified from the unsupervised clustering results. To further extend this cell type classification, the NS-Forest algorithm was used to identify the minimum sets of marker genes for each cluster. NS-Forest uses random forest machine learning and a binary expression scoring method to identify necessary and sufficient marker genes, optimally capturing the essence of cell type identity[42-43]   . NS-Forest analysis of the 60 cell type clusters yielded 117 marker genes with high cell type specificity illustrated by a diagonal expression pattern of markers across the clustered dataset (Figure 3A) and relatively high F-beta values of

classification accuracy (Supplementary Table 1). For example, HC Clusters 23.2 and 23.4 showed F-beta values of 0.93 and 0.88 using the single cell type markers Slc26A5 (Prestin)[44] and Ripor3, respectively. For the other HC clusters, combinatorial expression of two marker genes gave optimal classification accuracy of F-beta = 0.85 with Gm1113 and Ofcc1 markers for Clusters 23.1 and markers Ush2a and C230072F16Rik for Cluster 23.3. NS-Forest marker genes for the SGN populations included Ntng1, Mdga1, Cdh9, and Meg3, which have been shown to be among the highest differentially expressed genes during the diversification process of SGN (Petitpré et al., 2022).

NS-Forest also produces an extended set of marker genes that show specific binary expression patterns across the cell clusters. In the case of the four HC clusters (Clusters 23.1, 23.3, 23.2 and 23.4), the genes Slc26a5, Ocm, and Cacnald encoding for Prestin[44], Oncomodulin[45], and Calcium voltage dependent channel (Cav1.3)[46] respectively, are highly expressed in Cluster 23.2 suggesting this group is an OHC population (Figure 3B). Cluster 23.4 shows specific expression of Slc17a8[47] and Nrxn1[48], known IHCs markers encoding for Vesicular Glutamate Transporter-3 and a presynaptic adhesion protein, respectively. Cluster 23.3 shows specific expression of Myo15b, while Cluster 23.1 shows high expression of Kcnab1, which encodes a potassium voltage-gated channel, a HC marker conserved across species[49] and shown to be more IHC specific in mature mouse cochlea[50-51]. Genes recently validated as HCs gene markers in newborn cochlea[19] were not identified in our dataset, suggesting that those markers may be more specific for early postnatal stages.

To extend the granular cell type annotations, the transcriptomics clusters were compared against five recently released snRNA and scRNA postnatal dataset (Supplementary Table 2) hosted by the gEAR database (https://umgear.org) using FR-Match, a cluster-to-cluster cell type matching algorithm that incorporates shared information among cells to determine the relationship between two transcriptome clusters[52] using the cluster-specific marker genes identified by NS-Forest to provide a reduced dimensional space and support their statistical comparison. A match was considered high confidence if a match was found in both directions (query to reference and reference to query) and/or if a match was found to the same cell type with two or more datasets; a match was considered low confidence if there was a match in only one dataset in only one direction. The FR-Match comparisons show that 29 clusters were matched to gEAR cell types with high confidence (Supplementary Figure 1 and Table 1). The analysis results confirmed the classification of strial cells (marginal, intermediate, and basal) and spindle-root populations with high confidence (up to 8 matches). The clusters corresponding to cochlear OHCs, glial/Schwann cells, immune cells, and a subset of fibrocyte populations were also confirmed with high confidence, albeit with lower numbers of matches. The cell populations characterized as Hensen's cells, Dieters cells, and pillar cells and the remaining fibrocytes populations were matched with lower confidence. The results of FR-Match analysis were also consistent with the cluster hierarchy, highlighting the advantage of using the FR-Match to identify corresponding cell types.

To validate the specificity of NS-Forest cell-type-specific marker expression and to characterize the spatial relationships between the cell types, the recently developed multiplexed error-robust fluorescent in situ hybridization technique (MERFISH), a single-molecule imaging approach that allows identification and spatial localizations of RNA transcripts for several different cell types[77-78], was used in cross-sections from P5 cochleae. The NS-Forest cell-type specific marker genes (Figure 3A) were utilized as MERFISH probes to confirm the consistency between the cell-type annotations and transcripts localizations. For example, marker genes for HC clusters

localized to IHCs (Cluster 23.1, Gm1113+/Ofcc1+) (Fig. 4A), OHCs (Cluster 23.2, Slc26A5+) (Fig. 4B), both IHC and OHC (Cluster 23.4, Ripor3+) (Fig. 4C). Moreover, the MERFISH imaging confirmed the annotation of some subcluster categories with the lower confidence FR-Match results, such as border cells for Cluster 20.1 (Dgkb+/Klhl14+) (Fig. 4D), inner and outer pillar cells for Cluster 20.2 (Mdm1+) (Fig. 4E), Deiter's cell and inner phalangeal cell for Cluster 20.3 (Dgkb+/Nckap5+/Slc1a3+) (Fig. 4F), where the localization pattern and clustering matched conventional anatomical localization nomenclature in the cochlea. Interestingly, some marker genes localized to the same anatomical cell-types but indicated different cell populations among the subcategories, such as Cluster 12 for spiral ganglion cells (Fig. 4H and I) and Cluster 17 for Schwann cells (Fig. 4J). These data suggested that the NS-Forest algorithm successfully identified unique marker genes for the transcriptomic clusters and the spatial transcriptomic analysis verified the FR-match annotations based on anatomical and physiological properties of cochlear cells.

For further evaluation of the proportion of cell types, we investigated the spatial organization of the scala media containing the organ of Corti, which is the receptor organ for hearing, by MERFISH (Supplementary Fig.2A and 2B). In addition to HC and supporting cell clusters (Fig.4), we successfully identified various cell clusters in the cochlear duct, such as epithelial cells of Reisner membrane (Cluster4, Soc26a7+/Tmem72+) (Supplementary Fig.2C), marginal cells (Cluster 5, Kcnq1+/Stac+) and intermediate cells (Cluster9, Dct+) in the stria vascularis (Supplementary Fig.2D), spindle cells (Cluster 19.2, Anxa1+/Dpp10+)(Supplementary Fig.2E), root cells in the spiral prominence (Cluster 6, Lgr5+/Slc26a4+)(Supplementary Fig.2F), inner and outer sulcus cells (Cluster 10, Epyc+/Gata3+)(Supplementary Fig.2G), fibrocytes in the tympanic covering layer of basilar membrane (Cluster 11, Mybpc1+/Nav3)(Supplementary Fig.2H), interdental cells in the spiral limbus (Cluster16, Otoa+)(Supplementary Fig.2I). Notably, the clustering successfully distinguished transcriptional profiles even among rare cell types in the spiral prominence of the cochlea, including spindle cells (Cluster19) and root cells (Cluster6). These data demonstrate that the snRNA-seq approach robustly profiled cell type-specific transcriptomes in the cochlear tissue.

By combining the information derived from manual marker gene evaluation, gEAR reference matching, and MERFISH localization, cell type names and definitions for the 60 transcriptomics clusters were determined (Table 1). In order to provide for more descriptive cell type names for each of the transcriptomic clusters, we adopted a convention established by the ontology development community for defining and naming cell types derived from single cell transcriptomics experiments for incorporation into the Provisional Cell Ontology[28,53], incorporating information about anatomic location, parent cell class, and specific marker gene combinations to provide for unique cell type names and experimentally useful definitions (Table 1 and Supplementary Table 1) to serve as future references.

In order to determine if loss of hearing acuity is associated with the loss of specific cochlear cell types, the auditory brainstem response (ABR) threshold was compared with the cell type proportions in the 48 outbred CFW mice with three patterns observed (Figure 5A). For example, the proportion of cells in Clusters 23.2, 23.4, 12.1, and 12.2 showed an inverse correlation with sound intensity (Figure 5A, red; Figure 5B, C, D), whereas the proportion of cells in Clusters 2.1 and 2.2 showed a positive correlation (Figure 5A, blue), while the vast majority of clusters showed no correlation (Figure 5A, black and Supplementary Figure 3). In support of this observation, whole-mount immunohistochemistry showed that OHC and IHC structures are normal in mice with good hearing (Figure 5E) compared to aged mice with poor hearing (Figure

5F), demonstrating that HC loss is a primary driver of ARHL in CFW and an ideal model to replicate the phenotype seen in humans[54].

In comparing the cell type proportions with each other (Supplementary Figure 4), all cell types that show proportional loss with hearing loss are positively correlated with each other (e.g., Cluster 23.2 with Clusters 23.4, 12.1, and 12.2 (top row, left)), with the correlation strongest for cell types that are closely related to each other (e.g., Cluster 12.1 and 12.2, third row); all cell types that show proportionally increases with hearing loss are positive correlated with each other (e.g., Cluster 2.1 with Clusters 2.2, 19.1, and 19.2 (fifth row, middle)); all of the cell types that show proportional increases with hearing loss are negatively correlated with cell types that show proportional loss with hearing (e.g., Cluster 2.1 with Clusters 23.2, 23.4, 12.1, and 12.2 (fifth row, left)); cell types that show no correlation with hearing loss show no correlation with cell types that show some correlation with hearing loss (e.g., Cluster 23.3 with Clusters 23.2, 23.4, 12.1, 12.2, 2.1, 2.2, 19.1, and 19.2 (bottom row, left and middle)). Interestingly, although the proportions of HC Clusters 23.1 and 23.3 do not correlate with HC clusters 23.2 and 23.4, they show strong correlation with each other. These results suggest that hearing loss is associated with specific concerted changes in cell populations in this genetic model of ARHL.

Discussion:

In this manuscript we have described the first snRNA-seq study of the aged adult outbred mouse cochlea correlated with auditory physiological data. In contrast to a recent study, using scRNA-seq, which identified 27 cell types within the cochlea of a single mouse strain, C57BL/6J (Sun G et al.), we report the identification of 60 distinct cell types in this outbred mouse model using snRNA-seq to avoid the stress responses induced during the single cell processing procedure and to provide for a more unbiased representation of cell types from solid tissues[28].

Presbycusis, ARHL, is the most common sensory disorder in man and is characterized by reduced hearing sensitivities and speech understanding, particularly in noisy environments. Classically described, there are four subtypes, sensory, neural, strial and mixed, each associated with a corresponding cellular senescence[55]. The findings reported here confirm the complex nature of ARHL regarding the survival/senescence of specific cell types. It is well known that SGN numbers decrease with age and are responsible for mixed and sensory forms of hearing loss including their ultimate decline after synaptic ribbon loss[56]. In this study, two of the four SGN clusters decreased with increasing hearing loss, and yet, two subtypes of SGN cells identified in this analysis did not. It is also well known that HC loss is a common finding in most acquired hearing loss, yet, of the four subtypes of HCs detected in our in-depth analysis, only two decline with hearing loss. We also observe that pillar and inner border cell populations decline with hearing loss resulting in a loss of cytoarchitecture in the organ of Corti. Elevated hearing thresholds were associated with loss of marginal cell clusters in the stria vascularis, the portion of the ear responsible for maintenance of the endocochlear potential. The increased number of basal cells and spindle cells positively correlating with higher hearing thresholds suggest a compensatory role for these cells in maintaining the endocochlear potential with aging[57]. The increased numbers of macrophages positively correlated with hearing loss likely represents the inflammatory response to the age-related cochlear stressors[58-60]. Finally, our analysis identified seven separate clusters of fibrocytes with two out of the seven declining in numbers with hearing loss.

The use of snRNA-seq to identify and quantify cell types in the cochlea of elderly mice with varying levels of age-related hearing loss (ARHL) identified ten cell types whose proportions

inversely correlated with hearing acuity, including two HC types (cochlear OHC Slc26a5 and HC Ripor3), two spiral ganglion neuron types (cochlear spiral ganglion neuron C130073E24Rik Ntng1 and cochlear spiral ganglion neuron Mdga1 Tmem108), one Schwan cell (cochlear Schwan cell Pde1c Lama1), four border cell types (cochlear inner / outer sulcus Prss36 Hmcn1, cochlear border cell Dgkb Klhl14, cochlear pillar cell Mdm1, and cochlear phalageal cell Dgkb Nckap5), and one marginal cell type (cochlear marginal Stac Kcnq1) (Supplementary Table 3). Each of these distinct cell types is characterized by the expression of specific combinations of marker genes. Since these marker genes show very specific expression in the cell type that they mark, they may play a critical role in the function of that cell type. For example, the two marker genes for cochlear marginal Stac Kcnq1 cells are both involved in ion channel function that regulate plasma membrane potential[61-62] and are likely to be essential for the proper functioning of the cell. Since these cell types are critical for maintaining hearing with age, it is possible that their marker genes would also be essential for hearing. Indeed three of the 19 marker genes for these ARHL-associated cell types show genetic associations with hearing deficits (Supplementary Table 3). Kvlqt1 (aka Kcnq1) -/- mice are completely deaf due to defects in inner ear development[63]. Mutations in KVLQT1 in humans cause Jervell and Lange-Nielsen (JLN) syndrome, an inherited autosomal recessive disease characterized by a congenital bilateral deafness associated with QT prolongation and ventricular arrhythmias[64]. Mutations in PDE1C are associated with an autosomal dominant form of nonsyndromic postlingual progressive deafness in human[65]. Mutations in SLC26A5 (aka Prestin) are associated with familial nonsyndromic hearing loss in humans[66]. Homozygous Slc26a5 mutant mice showed a loss of outer HC electromotility in vitro and a loss of cochlear sensitivity in vivo without disruption of mechanoelectrical transduction in outer HCs[67]. The finding that some of these cell type specific marker genes show genetic association with hearing deficits supports the observation that loss of these specific cell types would lead to ARHL.

Interestingly, marker genes specific for other cell types that did not correlate with ARHL in our mouse model also show genetic associations with certain types of hearing deficits (Supplementary Table 4). For example, although the proportion of cochlear HC Ush2a C230072F16Rik did not diminish with ARHL, mutations in its marker gene USH2A causes Usher syndrome type IIa, an autosomal recessive disorder characterized by moderate to severe sensorineural hearing loss and retinitis pigmentosa[68]. These findings suggest that five other cochlear cell types identified in our study are involved in other types of hearing loss syndromes.

In conclusion, by combining information from manual marker gene annotation, gEAR reference data matching and MERFISH spatial transcriptomics analysis, 60 unique transcriptomic clusters were identified and annotated with cell type identities and specific marker gene characterization. Several of these specific cell types showed preferential loss in the aging cochlea that also correlated with quantitative measures of hearing loss. The genes specifically expressed in these cells could serve as candidate targets for novel therapeutics in the future.

Materials and Methods:

Animals

Animals originated from the same stock of outbred mice Crl:CFW(SW)-US_P08 (CFW), maintained by Charles River Laboratories (CRL) in Portage, Michigan, were used in this study. All procedures were performed in accordance with guidelines from NIH and with approval by the Institutional Care and Use Committee (IACUC) at University of California San Diego (IACUC S17178).

Auditory phenotyping/ Hearing Patterns determination

All CFW mice were subsequently aged until a final age of ten months. Within this period, phenotypic audiometric assessment using Auditory Brainstem Response (ABR) was completed at three time points: young adults (5-8 weeks), 6 months, and 10 months. Auditory phenotyping and determination of hearing patterns were previously detailed[69]. Briefly, upon aging to the stated time points, anesthetized mice were presented to auditory signals as tone pips ranging from 20 dB to 100 dB SPL at the frequencies 4 kHz, 8 kHz, 12 kHz, 16 kHz, 24 kHz, and 32 kHz.

The detection distinctive ABR waveform at each frequency was used to determine the hearing thresholds and characterize ARHL within the CFW mice into one of eight distinct hearing patterns: Normal hearing; isolated mid-frequency (12 kHz or 16 kHz) loss; Moderate high-frequency (24 kHz or 32 kHz) loss; Severe high-frequency (only 32 kHz) loss; Severe high-frequency (24 kHz + 32 kHz) loss; Moderate all-frequency loss; Severe all-frequency loss; and Profound all-frequency loss.

Isolation of cochlear tissue

A group of 48 mice were sacrificed and their inner ears were utilized to generate single nucleus RNA-seq transcriptomes. Mice were selected to represent the hearing pattern group described above based on the proportion of such pattern in the general CFW cohort. Special care was taken to complete the microdissection of the 2 cochleae from each mouse in less than 7 minutes while ensuring a minimum physical stretching on the organ of Corti. Samples were collected at the same time of day across individual and batches.

Following the ABR testing at 10 months age final time point, the anesthetized mice were decapitated, and their inner ears were quickly transferred into ice-cold DPBS buffer (Thermo Fisher Scientific) for microdissection. Tissue from utricule and saccule were carefully removed to avoid including unwanted vestibular cell types before removing the bony wall of the cochlea. The microdissected tissue from each mouse was pooled in a tube containing 1 ml of Iscove's Modified Dulbecco's Medium (IMDM) (Thermo Fisher Scientific).

Single nucleus isolation and RNA sequencing

Immediately after isolation, bulk cochlear tissues in cell culture media were processed for snRNA-seq serially in sets of four specimens and kept on ice for the entire procedure. Dounce homogenization was performed to isolate individual nuclei from the bulk cochlear tissue followed by Fluorescent Automated Cell Sorting (FACS) to provide purified, intact nuclei for RNA-seq as detailed in [Krishnaswami 2016] with the following modifications. Briefly, a total of 5 mLs of Homogenization Buffer with Propidium Iodide (PI) (1.5 µM, Thermo Fisher cat. #P3566) and Calcien, AM (1.0 µM) (Thermo Fisher cat. #C1430) was used to stain intact nuclei and exclude whole cells from being sorted. The cochlear tissue in cell culture media was briefly

centrifuged to remove the supernatant to approximately 100 µL volume and the remaining tissue transferred using a 1 mL wide bore pipette tip into 1 mL of chilled Homogenization Buffer in a Dounce Homogenizer (Wheaton cat. #357538) stored on ice. A total of 10 strokes with the loose piston followed by 14 strokes with the tight piston was followed by filtration through a series of two cell strainers (Becton Dickenson Falcon cat. #352235) and loading onto a Beckton Dickenson FACS Aria II cell sorter with a 70 mm nozzle. A chilled 96-well plate was used with 4 µL of Homogenization Buffer placed in wells A1 to A4 as a destination for sorted nuclei. FACS gating of nuclei were accomplished with forward and side scatter doublet nuclei gating exclusion, green channel (Calcien, AM) exclusion, and triggered on the red (PI) channel with approximately 1-7% of the total event population consisting of intact nuclei. A total of 36,000 sorted nuclei from each of the 4 samples were sorted into wells A1 to A4 with a final volume of approximately 60 µL after completion of sorting.

Isolated nuclei in the 4 wells were kept on ice and processed for RNA-seq using the Chromium Next GEM Chip G Single Cell Kit (10X Genomics cat. #1000120) according to manufactures instructions with the following modifications. A total of 43.2 µL of sorted volume containing approximately 26,000 nuclei was added to 31.8 µL of the Single Cell Master mix for a total of 75 µL, followed by loading of 70 µL (approx. 24,267 nuclei) onto the Next GEM chip. cDNAs were amplified for 13 cycles. Recovered cDNAs were quality controlled by loading 1 µL of cDNA onto an Agilent Bioanalyzer using a High Sensitivity DNA chip kit (cat. #5067-4626), followed by library prep according to manufacturer's instructions for dual index barcoded libraries.

The nuclei isolation, sorting, and Next Gem chip cDNA synthesis procedures were repeated 12 more times for a total of 48 cDNA library samples. A total of 48 single nuclei barcoded RNA-seq libraries were subsequently generated from the cDNAs using the Single Index Kit T Set A (10X Genomics cat. #1000213) and 16 cycles of library amplification followed by purification according to manufactures instructions. A 1 µL loading of a 10-fold dilution of each library was performed for quality control and quantification on an Agilent High Sensitivity chip. Equimolar amounts (6 nM) of a subset of 16 libraries were pooled and diluted to create a final loading of a 300 pM library pool onto a dual-lane NovaSeq 6000 (Illumina, Inc.) using the S2 Reagent Kit v1.5 Paired End 2x50 base (100 cycle) sequencing kit (Illumina cat. #20028316). Sequencing parameters were set for R1 at 28 cycles, I7 Index at 8 cycles, I5 index at 0 cycles, and Read 2 at 91 cycles for a calculated R2 read count of 25,600 per cell. Two more NovaSeq 6000 S2 runs of 16 library pools were subsequently loaded for a total of 3 runs to complete the RNA-seq of all 48 samples. A range of X to Y reads per cell were generated.

Read alignment and nucleus detection

The cellranger count command with the –include-intron option from the 10x Cell Ranger 5.0.1 package[76] was used to align reads and count barcodes and UMIs. Reads were aligned to the Cell Ranger reference package refdata-gex-mm10-2020-A.

Quality control

Data quality was assessed using the number of nuclei per sample, the number of UMIs (library size), the number of genes detected and the number of mitochondrial genes per nuclei. Nuclei were removed if they had fewer than 1000 UMIs and/or fewer than 500 detected genes. Nuclei that had the number of detected mitochondrial genes tagged as outliers by the isOutlier function from the scuttle package were also removed[74].

Normalization, feature selection and dimensionality reduction
UMI counts were normalized using logNormCounts from the scuttle package. Highly variable genes were selected using modelGeneVar and getTopHVGs functions from the scran package[73]. PCA was performed using the runPCA function from the scater package. Nineteen principal components were used for clustering, which accounted for 50% of the variance.

Clustering and doublet detection
To minimize the inclusion of droplets that have more than one nucleus, putative doublets were identified using the scDblFinder package[71]. Since scDblFinder uses cluster information to predict doublets, nuclei were first clustered using the Leiden community detection algorithm as implemented in the leidenalg python package[75]. First, a shared nearest-neighbor graph was constructed with buildSNNGraph from the scran package[74]. Then the leidenalg.find_partition function with partition_type set to ModularityVertexPartition was called. Finally, sclDblFinder was called using the resultant cluster labels. About 6% of the droplets were predicted to be doublets and removed from downstream analysis (Supplementary Figure 5). The remaining nuclei were re-processes repeating quality control, normalization, feature selection, dimensionality reduction and clustering.

Visualization
The clustered data set was visualized by running runUMAP and runTSNE from the scater package[74]. For runUMAP, n_neighbors was set to 100; for runTSNE, perplexity was set to 100.

Sub-clustering
The initial clustering using Leiden community detection yielded 26 clusters. To determine if any of these clusters should undergo another round of clustering, two methods were used: the silhouette width, which compares the average distance of a nucleus to all other nuclei within the same cluster to the average distance to nuclei in the nearest neighboring cluster[70], and manual inspection. Supplementary Figure 6 shows an example of this analysis where Clusters 1, 10, and 17 contain separate areas of nuclei with negative silhouette widths colored in red as candidates for sub-clustering. Other candidate clusters were detected by manual inspection of the tSNE embeddings (Figure 2A), for example, Clusters 20 and 23, which appear to have two distinct "islands" suggesting the need for sub-clustering. Supplementary Figure 7 shows examples of the resulting subclusters. By applying those criteria on the initial clusters, 13 were retained and 13 went through a second round of sub-clustering, which brought the total number of transcriptomic clusters identified to 60. The final clustered cell-by-gene expression matrix can be found at the gEAR resource - https://umgear.org/p?l=c2acc279 .

Cluster markers
NS-Forest was used to identify the minimum number of necessary and sufficient markers for each cluster[43]. NS-Forest was performed with the number of trees set to 1000 and the number of genes to test set to 6.

Multiplexed error-robust fluorescence in situ hybridization (MERFISH) on the cochlear tissue

MERFISH samples were prepared in accordance with company instructions (Vizgen, Cambridge, MA, USA). Briefly, C57Bl/6J mice cochleae of postnatal day 5 were harvested and fixed with 4% paraformaldehyde (PFA) in 0.1 M phosphate-buffered saline (PBS; pH 7.4) at 4 °C overnight. Samples were dehydrated in graded-sucrose series (10% and 20% for 30 min, and 30% for overnight at 4 °C) with RNase inhibitor [New England Biolabs (NEB), M0314L, Ipswich, MA, USA]. Samples were placed in a cryomold and embedded with O.C.T. compound (Sakura Finetek, Torrance, CA, USA), then frozen with dry ice/ethanol bath. The embedded tissue was sectioned into 10μm thick slices using a Leica CM1860 (Leica Biosystems, Nussloch, Germany) cryostat, and 3 to 5 cochlear mid-modiolar sections were mounted onto a center of MERSCOPE slide glass (Vizgen, #20400001, USA). The mounted sections were washed with 0.1 M PBS, then permeabilized in 70% ethanol at 4 °C for 24 h. The cell boundary staining was performed by using a primary antibody mix (Vizgen, #20300010, USA), followed by a secondary antibody mix (Vizgen, #20300011, USA) for 1 h at 23 □, respectively. Stained samples were incubated with an encoding probe [MERSCOPE 140 Gene Panel Mix (Vizgen, #20300006, USA)] for 36 hours at humidified 37 □ cell culture incubator. After post-encoding hybridization wash with formamide wash buffer (Vizgen, #20300002, USA), samples were embedded with a gel embedding solution [gel embedding premix (Vizgen, #20300004, USA), 5 ml; 10% ammonium persulfate solution (Millipore-Sigma, 09913-100G, Burlington, MA, USA), 25 µl; N,N,N',N'-tetramethylethylenediamine (Millipore-Sigma, T7024-25ML, USA), 2.5 µl]. For tissue clearing, samples were incubated in digestion premix (Vizgen, #20300005, USA) with RNase inhibitor (NEB, USA) for 1 h at 23 °C, followed by clearing premix [clearing premix (Vizgen, #20300003, USA), 5ml; proteinase K (NEB, P8107S, USA), 50□l] for 48 h at humidified 37 □ cell culture incubator. After the tissue became transparent, samples were washed with the wash buffer (Vizgen, #20300001, USA) and incubated with 4′,6-diamidino-2-phenylindole (DAPI) and polythymine (polyT) staining reagent (Vizgen, #20300021, USA) for 15 min with agitation. Images were taken by using MERSCOPE (Vizgen, USA). DAPI / polyT and cell-boundary staining 2 was utilized for the cell segmentation parameter respectively, then image processing analysis was done on the MERSCOPE. The images were visualized and analyzed on the MERSCOPE Visualizer (Vizgen, USA).

Figure Legends:
Figure 1 - Phenotyping CFW outbred mice shows eight distinct hearing patterns. Auditory Brain Response (ABR) thresholds measure the health of hearing function and correspond to the response to sound stimulus with known intensity at a specific frequency. The mouse is considered to have mild-moderate hearing impairment if ABR thresholds were between 41 - 60 dB, severe impairment if between 61 – 80 dB, and profound impairment if greater than 80 dB. Symbols indicate ABR levels in individual outbred mice. Dotted lines indicate median thresholds for each of the eight groups. A) Normal hearing; B) moderate mid-frequency (12 kHz or 16 kHz) loss; C) moderate high-frequency (24 kHz or 32 kHz) loss; D) severe high-frequency (only 32 kHz) loss; E) severe high-frequency (24 kHz + 32 kHz) loss; F) moderate all-frequency loss; G) severe all-frequency loss; H) profound all-frequency loss. I) Individual ABR plots for mice selected to represent the eight patterns for cochlear tissue collection and snRNA-seq analysis.

Figure 2 – Sixty snRNA-seq transcriptomic clusters from mouse cochlea. A) Iterative unsupervised clustering of the processed cell-by-gene snRNA-seq expression matrix performed

using Leiden community detection identified sixty distinct transcriptomic clusters (numbers). tSNE embedding of the clustering results is shown. Cell type annotations of each transcriptomic cluster based on subsequent analysis (Table 1) is shown. The data can be explored at https://umgear.org/index.html?multigene_plots=0&layout_id=c2acc279&gene_symbol_exact_match=1&gene_symbol=slc26a5 . B) Heatmap of marker genes known to be expressed in specific ear cell types arranged according to hierarchical clustering of transcriptomic clusters, which captures the transcriptional relationships between the discrete clusters. The horizontal banding patterns of these known marker genes validates the structure of the hierarchical clustering taxonomy. C) Box plots of the proportion (log10 transformed) of each cell type across the 48 outbred mouse cochlear samples, with median proportion ranging from ~10^{-4} for Cluster 22.1 to ~10^{-1} for Cluster 4.

Figure 3 – Cell type-specific marker genes. NS-Forest was used to identify the minimum set of necessary and specific marker genes for each cell type, with gene expression (rows) across each transcriptomic cluster (columns) illustrated in violin plots. A) Minimum NS-Forest marker genes identified from the complete cell-by-gene expression matrix, with diagonal expression patterns and little off-diagonal expression showing marker gene specificity. In addition to the minimum site of marker genes, NS-Forest also produces an extended set of ten genes selected based on binary expression (on-off) patterns in the target cluster. B) Top ten binary genes were identified within the four Cluster 23 iterative subclusters using the pooled Cluster 23 subcluster cell-by-gene expression matrix as input, showing strong and distinct binary expression patterns, especially for Clusters 23.3 and 23.4.

Figure 4 – Spatial organization of cell types in the mouse cochlea. MERFISH images with selected marker genes for Cluster 23 (cluster of HC) (A-C), Cluster 20 (cluster of supporting cell) (D-F), Cluster 12 (cluster of spiral ganglion cell) (I-L) and Cluster 17 (cluster of Schwann cell) on the cochlear mid-modiolar cryosections at P4. Each pseudo-color dots represent transcripts of different marker genes and are merged with cell-boundary staining in dark grey. Cell boundaries are depicted by white or white-dashed line. Myo7a was used for pan-HC marker gene (yellow) (A-F). A) Cluster 23.1 for IHC with Gm113(magenta, arrow), Ofcc1(blue, arrowhead). B) Cluster 23.2 for OHC with Slc26a4 (red, arrow). C) Cluster 23.4 for unknown hair-cell subtype with Ripor3 (green, arrow). D) Cluster 20.1 for inner border cell with Dgkb (green, arrowhead) and Klhl14 (magenta, arrow). E) Cluster 20.2 for pillar cell with Mdm1 (red, arrow). F) Cluster 20.3 for inner phalangeal cell (white line) and Deiter's cell (white-dashed line) with Dgkb (green, arrowhead), Nckap5 (blue) and Slc1a3(orange, arrowhead). I) Localization of marker genes for cluster 12 in cochlear spiral ganglion cells (arrow) with Ntng1 (yellow), Mdga1 (light blue) and Meg3 (dark blue). H. I. Binary color images of cluster 12.1 with Ntng1(arrow) and C130073E24Rik (thick arrow), of cluster 12.2 with Cdh9 (double-head arrow), Tmem108 (arrowhead), and Mdga1, and of cluster 12.3 with Meg3, and merged image with pseudo colors (I). Representative cell boundaries for each subcluster are depicted by white-dot (cluster 12.1) and white-dashed (cluster 12.2), and asterisk indicates a cell of cluster 23.3. Mpz (brown) is a marker gene for Schwann cell. M. Images for cluster 17.1 with Pde1c (red, arrowhead) and Lama1(green, thick arrow), and for cluster 17.2 with Ntng1 (light blue, double-head arrow) and Cdh19 (yellow, arrow). Meg3 (dark blue) and Mpz (brown) indicate spiral ganglion cells and Schwann cells, respectively. Sale bars; 15μm (A-F), 250μm (G), and 25μm (H-J).

Figure 5 – Loss of specific cochlear cell types with age-related hearing loss. The proportion of each cell type in the 48 outbred mouse cochlea was compared against the ABR sound intensity score at different sound frequencies. A) Correlation between selected cell type proportions (Y-axis) and ABR sound intensity thresholds at 32 kHz (x-axis). HC Clusters 23.2 and 23.4 and ganglion neuron Clusters 12.1 and 12.2 show significant inverse correlations with intensity threshold (red). Basal cell Clusters 2.1 and 2.2 show significant positive correlation with intensity threshold (blue). In contrast to the inverse correlation observed with Clusters 23.2 and 23.4, HC Clusters 23.1 and 23.3 show no correlation with intensity threshold. Spearman's rho correlation and p-value are show at the top of each plot. B) Sound intensity threshold of individual mice ranked in order of hearing loss. C) Proportion of Cluster 23.2 cells in individual mice. D) Proportion of Cluster 23.4 cells in individual mice. Whole mount projections of organ of Corti corresponding to areas encoding 16 kHz frequency show that mice with profound hearing loss (F) exhibit IHC and OHC loss that worsens through the higher frequencies while the organ of corti from mice with normal hearing (E) show intact sturctures with no HC loss.

ACKNOWLEDGEMENTS

Research reported in this publication was supported by NIDCD of the National Institutes of Health under award number: R01DC018566.

REFERENCES

1. Fransen E., Bonneux S., Corneveaux J. J., Schrauwen I., Di Berardino F., White C. H., et al.. (2015). Genome-wide association analysis demonstrates the highly polygenic character of age-related hearing impairment. *Eur. J. Hum. Genet.* 23, 110–115. 10.1038/ejhg.2014.

2. Bainbridge KE, Wallhagen MI. Hearing loss in an aging American population: extent, impact, and management. *Annu Rev Public Health*. 2014;35:139-152. doi:10.1146/annurev-publhealth-032013-182510

3. Deal, Jennifer A., et al. "Hearing impairment and incident dementia and cognitive decline in older adults: the health ABC study." *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences* 72.5 (2017): 703-709.

4. Deal, J. A., Goman, A. M., Albert, M. S., Arnold, M. L., Burgard, S., Chisolm, T., ... & ACHIEVE Investigators. (2018). Hearing treatment for reducing cognitive decline: Design and methods of the Aging and Cognitive Health Evaluation in Elders randomized controlled trial. Alzheimer's & Dementia: Translational Research & Clinical Interventions, 4, 499-507.

5. Lin, F. R., Niparko, J. K., & Ferrucci, L. (2011). Hearing loss prevalence in the United States. *Archives of internal medicine*, *171*(20), 1851-1853.

6. Gurgel, R. K., Ward, P. D., Schwartz, S., Norton, M. C., Foster, N. L., & Tschanz, J. T. (2014). Relationship of hearing loss and dementia: a prospective, population-based study. *Otology & neurotology: official publication of the American Otological Society,*

American Neurotology Society [and] European Academy of Otology and Neurotology, *35*(5), 775.

7. Jayakody, D. M., Friedland, P. L., Martins, R. N., & Sohrabi, H. R. (2018). Impact of aging on the auditory system and related cognitive functions: a narrative review. *Frontiers in neuroscience*, 125.

8. Vos, T., Lim, S. S., Abbafati, C., Abbas, K. M., Abbasi, M., Abbasifard, M., ... & Bhutta, Z. A. (2020). Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet*, *396*(10258), 1204-1222.

9. Bowl, M. R., & Dawson, S. J. (2019). Age-related hearing loss. *Cold Spring Harbor perspectives in medicine*, *9*(8), a033217.

10. Parker, C. C., Gopalakrishnan, S., Carbonetto, P., Gonzales, N. M., Leung, E., Park, Y. J., ... & Palmer, A. A. (2016). Genome-wide association study of behavioral, physiological and gene expression traits in outbred CFW mice. *Nature genetics*, *48*(8), 919-926.

11. Lewis, M. A., Nolan, L. S., Cadge, B. A., Matthews, L. J., Schulte, B. A., Dubno, J. R., ... & Dawson, S. J. (2018). Whole exome sequencing in adult-onset hearing loss reveals a high load of predicted pathogenic variants in known deafness-associated genes and identifies new candidate genes. *BMC medical genomics*, *11*(1), 1-12.

12. Di Stazio, M., Morgan, A., Brumat, M., Bassani, S., Dell'Orco, D., Marino, V., ... & Girotto, G. (2020). New age-related hearing loss candidate genes in humans: An ongoing challenge. *Gene*, *742*, 144561.

13. Ingham, N. J., Rook, V., Di Domenico, F., James, E., Lewis, M. A., Girotto, G., ... & Steel, K. P. (2020). Functional analysis of candidate genes from genome-wide association studies of hearing. *Hearing research*, *387*, 107879.

14. Brown, S. D., Hardisty-Hughes, R. E., & Mburu, P. (2008). Quiet as a mouse: dissecting the molecular and genetic basis of hearing. *Nature Reviews Genetics*, *9*(4), 277-290.

15. Lavinsky, J., Crow, A. L., Pan, C., Wang, J., Aaron, K. A., Ho, M. K., ... & Friedman, R. A. (2015). Genome-wide association study identifies nox3 as a critical gene for susceptibility to noise-induced hearing loss. *PLoS genetics*, *11*(4), e1005094.

16. Liu, W., Johansson, Å., Rask-Andersen, H., & Rask-Andersen, M. (2021). A combined genome-wide association and molecular study of age-related hearing loss in H. sapiens. *BMC medicine*, *19*(1), 1-23.

17. Li, Y., Liu, H., Giffen, K. P., Chen, L., Beisel, K. W., & He, D. Z. (2018). Transcriptomes of cochlear inner and outer hair cells from adult mice. *Scientific data*, *5*(1), 1-12.

18. Hoa, M., Olszewski, R., Li, X., Taukulis, I., Gu, S., DeTorres, A., ... & Kelley, M. W. (2020). Characterizing adult cochlear supporting cell transcriptional diversity using Single-Cell RNA-Seq: validation in the adult mouse and translational implications for the adult human cochlea. *Frontiers in molecular neuroscience*, *13*, 13.

19. Kolla, L., Kelly, M. C., Mann, Z. F., Anaya-Rocha, A., Ellis, K., Lemons, A., ... & Kelley, M. W. (2020). Characterization of the development of the mouse cochlear epithelium at the single cell level. *Nature communications*, *11*(1), 1-16.

20. Korrapati, S., Taukulis, I., Olszewski, R., Pyle, M., Gu, S., Singh, R., ... & Hoa, M. (2019). Single cell and single nucleus RNA-Seq reveal cellular heterogeneity and homeostatic regulatory networks in adult mouse stria vascularis. *Frontiers in molecular neuroscience*, *12*, 316.

21. Petitpré, C., Wu, H., Sharma, A., Tokarska, A., Fontanet, P., Wang, Y., ... & Lallemend, F. (2018). Neuronal heterogeneity and stereotyped connectivity in the auditory afferent system. *Nature communications*, *9*(1), 1-13.

22. Shrestha, B. R., Chia, C., Wu, L., Kujawa, S. G., Liberman, M. C., & Goodrich, L. V. (2018). Sensory neuron diversity in the inner ear is shaped by activity. *Cell*, *174*(5), 1229-1246.

23. Sun, S., Babola, T., Pregernig, G., So, K. S., Nguyen, M., Su, S. S. M., ... & Müller, U. (2018). Hair cell mechanotransduction regulates spontaneous activity and spiral ganglion subtype specification in the auditory system. *Cell*, *174*(5), 1247-1263.

24. Petitpré, C., Faure, L., Uhl, P., Fontanet, P., Filova, I., Pavlinkova, G., ... & Lallemend, F. (2022). Single-cell RNA-sequencing analysis of the developing mouse inner ear identifies molecular logic of auditory neuron diversification. *Nature communications*, *13*(1), 1-15.

25. Burns, J. C., Kelly, M. C., Hoa, M., Morell, R. J., & Kelley, M. W. (2015). Single-cell RNA-Seq resolves cellular complexity in sensory organs from the neonatal inner ear. *Nature communications*, *6*(1), 1-16.

26. Rice MC, O'Brien SJ. Genetic variance of laboratory outbred Swiss mice. Nature. 1980;283:157–161.

27. Chia R, Achilli F, Festing MFW, Fisher EMC. The origins and uses of mouse outbred stocks. Nat Genet. 2005;37:1181–1186.

28. Bakken, T. E., Hodge, R. D., Miller, J. A., Yao, Z., Nguyen, T. N., Aevermann, B., Barkan, E., Bertagnolli, D., Casper, T., Dee, N., Garren, E., Goldy, J., Graybuck, L. T., Kroll, M., Lasken, R. S., Lathia, K., Parry, S., Rimorin, C., Scheuermann, R. H., Schork, N. J., … Tasic, B. (2018). Single-nucleus and single-cell transcriptomes compared in matched cortical cell types. *PloS one*, *13*(12), e0209648. https://doi.org/10.1371/journal.pone.0209648

29. Hanada, Y., Nakamura, Y., Ishida, Y., Takimoto, Y., Taniguchi, M., Ozono, Y., Koyama, Y., Morihana, T., Imai, T., Ota, Y., Sato, T., Inohara, H., & Shimada, S. (2017). Epiphycan is specifically expressed in cochlear supporting cells and is necessary for normal hearing. *Biochemical and biophysical research communications*, *492*(3), 379–385. https://doi.org/10.1016/j.bbrc.2017.08.092

30. Chen, Y., Qiang, R., Zhang, Y., Cao, W., Wu, L., Jiang, P., Ai, J., Ma, X., Dong, Y., Gao, X., Li, H., Lu, L., Zhang, S., & Chai, R. (2021). The Expression and Roles of the Super Elongation Complex in Mouse Cochlear Lgr5+ Progenitor Cells. *Frontiers in cellular neuroscience*, *15*, 735723. https://doi.org/10.3389/fncel.2021.735723

31. Nishio, A., Ito, T., Cheng, H., Fitzgerald, T. S., Wangemann, P., & Griffith, A. J. (2016). Slc26a4 expression prevents fluctuation of hearing in a mouse model of large vestibular aqueduct syndrome. *Neuroscience*, *329*, 74–82. https://doi.org/10.1016/j.neuroscience.2016.04.042

32. de Bruijn, M., & Dzierzak, E. (2017). Runx transcription factors in the development and function of the definitive hematopoietic system. *Blood*, *129*(15), 2061–2069. https://doi.org/10.1182/blood-2016-12-689109

33. Marcus, D. C., Wu, T., Wangemann, P., & Kofuji, P. (2002). KCNJ10 (Kir4.1) potassium channel knockout abolishes endocochlear potential. *American journal of physiology. Cell physiology*, *282*(2), C403–C407. https://doi.org/10.1152/ajpcell.00312.2001

34. Shibata, S., Miwa, T., Wu, H. H., Levitt, P., & Ohyama, T. (2016). Hepatocyte Growth Factor-c-MET Signaling Mediates the Development of Nonsensory Structures of the Mammalian Cochlea and Hearing. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *36*(31), 8200–8209. https://doi.org/10.1523/JNEUROSCI.4410-15.2016

35. Scheffer, D. I., Zhang, D. S., Shen, J., Indzhykulian, A., Karavitaki, K. D., Xu, Y. J., Wang, Q., Lin, J. J., Chen, Z. Y., & Corey, D. P. (2015). XIRP2, an actin-binding protein essential for inner ear hair-cell stereocilia. *Cell reports*, *10*(11), 1811–1818. https://doi.org/10.1016/j.celrep.2015.02.042

36. Yang, T., Scholl, E. S., Pan, N., Fritzsch, B., Haeseleer, F., & Lee, A. (2016). Expression and Localization of CaBP Ca2+ Binding Proteins in the Mouse Cochlea. *PloS one*, *11*(1), e0147495. https://doi.org/10.1371/journal.pone.0147495

37. Grandi, F. C., De Tomasi, L., & Mustapha, M. (2020). Single-Cell RNA Analysis of Type I Spiral Ganglion Neurons Reveals a *Lmx1a* Population in the Cochlea. *Frontiers in molecular neuroscience*, *13*, 83. https://doi.org/10.3389/fnmol.2020.00083

38. Sun, S., Siebald, C., & Müller, U. (2021). Subtype maturation of spiral ganglion neurons. *Current opinion in otolaryngology & head and neck surgery*, *29*(5), 391–399. https://doi.org/10.1097/MOO.0000000000000748

39. Gow, A., Davies, C., Southwood, C. M., Frolenkov, G., Chrustowski, M., Ng, L., Yamauchi, D., Marcus, D. C., & Kachar, B. (2004). Deafness in Claudin 11-null mice reveals the critical contribution of basal cell tight junctions to stria vascularis function. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *24*(32), 7051–7062. https://doi.org/10.1523/JNEUROSCI.1640-04.2004

40. Ng, L., Liu, Y., Liu, H., & Forrest, D. (2021). Cochlear Fibrocyte and Osteoblast Lineages Expressing Type 2 Deiodinase Identified with a Dio2CreERt2 Allele. *Endocrinology*, *162*(12), bqab179. https://doi.org/10.1210/endocr/bqab179

41. Scheffer, D. I., Shen, J., Corey, D. P., & Chen, Z. Y. (2015). Gene Expression by Mouse Inner Ear Hair Cells during Development. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *35*(16), 6366–6380. https://doi.org/10.1523/JNEUROSCI.5126-14.2015

42. Aevermann, B. D., Novotny, M., Bakken, T., Miller, J. A., Diehl, A. D., Osumi-Sutherland, D., Lasken, R. S., Lein, E. S., & Scheuermann, R. H. (2018). Cell type discovery using single-cell transcriptomics: implications for ontological representation. *Human molecular genetics*, *27*(R1), R40–R47. https://doi.org/10.1093/hmg/ddy100

43. Aevermann, B., Zhang, Y., Novotny, M., Keshk, M., Bakken, T., Miller, J., Hodge, R., Lelieveldt, B., Lein, E., & Scheuermann, R. H. (2021). A machine learning method for the discovery of minimum marker gene combinations for cell type identification from single-cell RNA sequencing. *Genome research*, *31*(10), 1767–1780. https://doi.org/10.1101/gr.275569.121

44. Yamashita, T., Hakizimana, P., Wu, S., Hassan, A., Jacob, S., Temirov, J., Fang, J., Mellado-Lagarde, M., Gursky, R., Horner, L., Leibiger, B., Leijon, S., Centonze, V. E., Berggren, P. O., Frase, S., Auer, M., Brownell, W. E., Fridberger, A., & Zuo, J. (2015). Outer Hair Cell Lateral Wall Structure Constrains the Mobility of Plasma Membrane Proteins. *PLoS genetics*, *11*(9), e1005500. https://doi.org/10.1371/journal.pgen.1005500

45. Sakaguchi, N., Henzl, M. T., Thalmann, I., Thalmann, R., & Schulte, B. A. (1998). Oncomodulin is expressed exclusively by outer hair cells in the organ of Corti. *The journal of histochemistry and cytochemistry : official journal of the Histochemistry Society*, *46*(1), 29–40. https://doi.org/10.1177/002215549804600105

46. Chen, J., Chu, H., Xiong, H., Chen, Q., Zhou, L., Bing, D., Liu, Y., Gao, Y., Wang, S., Huang, X., & Cui, Y. (2012). Expression patterns of Ca(V)1.3 channels in the rat cochlea. *Acta biochimica et biophysica Sinica*, *44*(6), 513–518. https://doi.org/10.1093/abbs/gms024

47. Ruel, J., Emery, S., Nouvian, R., Bersot, T., Amilhon, B., Van Rybroek, J. M., Rebillard, G., Lenoir, M., Eybalin, M., Delprat, B., Sivakumaran, T. A., Giros, B., El Mestikawy, S., Moser, T., Smith, R. J., Lesperance, M. M., & Puel, J. L. (2008). Impairment of SLC17A8 encoding vesicular glutamate transporter-3, VGLUT3, underlies nonsyndromic deafness DFNA25 and inner hair cell dysfunction in null mice. *American journal of human genetics*, *83*(2), 278–292. https://doi.org/10.1016/j.ajhg.2008.07.008

48. Mozhui, K., Wang, X., Chen, J., Mulligan, M. K., Li, Z., Ingles, J., Chen, X., Lu, L., & Williams, R. W. (2011). Genetic regulation of Nrxn1 [corrected] expression: an

integrative cross-species analysis of schizophrenia candidate genes. *Translational psychiatry*, *1*(7), e25. https://doi.org/10.1038/tp.2011.24

49. Janesick, A., Scheibinger, M., Benkafadar, N., Kirti, S., Ellwanger, D. C., & Heller, S. (2021). Cell-type identity of the avian cochlea. *Cell reports*, *34*(12), 108900. https://doi.org/10.1016/j.celrep.2021.108900

50. Liu, H., Pecka, J. L., Zhang, Q., Soukup, G. A., Beisel, K. W., & He, D. Z. (2014). Characterization of transcriptomes of cochlear inner and outer hair cells. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *34*(33), 11085–11095. https://doi.org/10.1523/JNEUROSCI.1690-14.2014

51. Zhenghong Bi, Xiang Li, Minhui Ren, Yunpeng Gu, Tong Zhu, Shuting Li, Guangqin Wang, Suhong Sun, Yuwei Sun, Zhiyong Liu, Development and transdifferentiation into inner hair cells require Tbx2, *National Science Review*, 2022;, nwac156, https://doi.org/10.1093/nsr/nwac156

52. Zhang, Y., Aevermann, B. D., Bakken, T. E., Miller, J. A., Hodge, R. D., Lein, E. S., & Scheuermann, R. H. (2021). FR-Match: robust matching of cell type clusters from single cell RNA sequencing data using the Friedman-Rafsky non-parametric test. *Briefings in bioinformatics*, *22*(4), bbaa339. https://doi.org/10.1093/bib/bbaa339

53. Tan, S. Z. K., Kir, H., Aevermann, B., Gillespie, T., Hawrylycz, M., Lein, E., ... & Osumi-Sutherland, D. (2021). Brain Data Standards Ontology: A data-driven ontology of transcriptomically defined cell types in the primary motor cortex. *bioRxiv*.

54. Wu, P. Z., O'Malley, J. T., de Gruttola, V., & Liberman, M. C. (2020). Age-Related Hearing Loss Is Dominated by Damage to Inner Ear Sensory Cells, Not the Cellular Battery That Powers Them. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *40*(33), 6357–6366. https://doi.org/10.1523/JNEUROSCI.0937-20.2020

55. Schuknecht, H. F., & Gacek, M. R. (1993). Cochlear pathology in presbycusis. *The Annals of otology, rhinology, and laryngology*, *102*(1 Pt 2), 1–16. https://doi.org/10.1177/00034894931020S101

56. Fernandez, K. A., Jeffers, P. W., Lall, K., Liberman, M. C., & Kujawa, S. G. (2015). Aging after noise exposure: acceleration of cochlear synaptopathy in "recovered"

ears. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, *35*(19), 7509–7520. https://doi.org/10.1523/JNEUROSCI.5138-14.2015

57. Hibino, H., & Kurachi, Y. (2006). Molecular and physiological bases of the K+ circulation in the mammalian inner ear. *Physiology (Bethesda, Md.)*, *21*, 336–345. https://doi.org/10.1152/physiol.00023.2006

58. Hough, K., Verschuur, C. A., Cunningham, C., & Newman, T. A. (2022). Macrophages in the cochlea; an immunological link between risk factors and progressive hearing loss. *Glia*, *70*(2), 219–238. https://doi.org/10.1002/glia.24095

59. Rai, V., Wood, M. B., Feng, H., Schabla, N. M., Tu, S., & Zuo, J. (2020). The immune response after noise damage in the cochlea is characterized by a heterogeneous mix of adaptive and innate immune cells. *Scientific reports*, *10*(1), 15167. https://doi.org/10.1038/s41598-020-72181-6

60. Noble, K., Brown, L., Elvis, P., & Lang, H. (2022). Cochlear Immune Response in Presbyacusis: a Focus on Dysregulation of Macrophage Activity. *Journal of the Association for Research in Otolaryngology : JARO*, *23*(1), 1–16. https://doi.org/10.1007/s10162-021-00819-x

61. Wong King Yuen, S. M., Campiglio, M., Tung, C. C., Flucher, B. E., & Van Petegem, F. (2017). Structural insights into binding of STAC proteins to voltage-gated calcium channels. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(45), E9520–E9528. https://doi.org/10.1073/pnas.1708852114

62. Schroeder, B. C., Waldegger, S., Fehr, S., Bleich, M., Warth, R., Greger, R., & Jentsch, T. J. (2000). A constitutively open potassium channel formed by KCNQ1 and KCNE3. *Nature*, *403*(6766), 196–199. https://doi.org/10.1038/35003200

63. Lee, M. P., Ravenel, J. D., Hu, R. J., Lustig, L. R., Tomaselli, G., Berger, R. D., Brandenburg, S. A., Litzi, T. J., Bunton, T. E., Limb, C., Francis, H., Gorelikow, M., Gu, H., Washington, K., Argani, P., Goldenring, J. R., Coffey, R. J., & Feinberg, A. P. (2000). Targeted disruption of the Kvlqt1 gene causes deafness and gastric hyperplasia in mice. *The Journal of clinical investigation*, *106*(12), 1447–1455. https://doi.org/10.1172/JCI10897

64. Neyroud, N., Tesson, F., Denjoy, I., Leibovici, M., Donger, C., Barhanin, J., Fauré, S., Gary, F., Coumel, P., Petit, C., Schwartz, K., & Guicheney, P. (1997). A novel mutation in the potassium channel gene KVLQT1 causes the Jervell and Lange-Nielsen cardioauditory syndrome. *Nature genetics*, *15*(2), 186–189. https://doi.org/10.1038/ng0297-186

65. Wang, L., Feng, Y., Yan, D., Qin, L., Grati, M., Mittal, R., Li, T., Sundhari, A. K., Liu, Y., Chapagain, P., Blanton, S. H., Liao, S., & Liu, X. (2018). A dominant variant in the PDE1C gene is associated with nonsyndromic hearing loss. *Human genetics*, *137*(6-7), 437–446. https://doi.org/10.1007/s00439-018-1895-y

66. Liu, X. Z., Ouyang, X. M., Xia, X. J., Zheng, J., Pandya, A., Li, F., Du, L. L., Welch, K. O., Petit, C., Smith, R. J., Webb, B. T., Yan, D., Arnos, K. S., Corey, D., Dallos, P., Nance, W. E., & Chen, Z. Y. (2003). Prestin, a cochlear motor protein, is defective in non-syndromic hearing loss. *Human molecular genetics*, *12*(10), 1155–1162. https://doi.org/10.1093/hmg/ddg127

67. Liberman, M. C., Gao, J., He, D. Z., Wu, X., Jia, S., & Zuo, J. (2002). Prestin is required for electromotility of the outer hair cell and for the cochlear amplifier. *Nature*, *419*(6904), 300–304. https://doi.org/10.1038/nature01059

68. Eudy, J. D., Weston, M. D., Yao, S., Hoover, D. M., Rehm, H. L., Ma-Edmonds, M., Yan, D., Ahmad, I., Cheng, J. J., Ayuso, C., Cremers, C., Davenport, S., Moller, C., Talmadge, C. B., Beisel, K. W., Tamayo, M., Morton, C. C., Swaroop, A., Kimberling, W. J., & Sumegi, J. (1998). Mutation of a gene encoding a protein with extracellular matrix motifs in Usher syndrome type IIa. *Science (New York, N.Y.)*, *280*(5370), 1753–1757. https://doi.org/10.1126/science.280.5370.1753

69. Du, E. Y., Boussaty, E. C., La Monte, O. A., Dixon, P. R., Zhou, T. Y., & Friedman, R. A. (2022). Large-scale phenotyping and characterization of age-related hearing loss in outbred CFW mice. *Hearing research*, *424*, 108605. https://doi.org/10.1016/j.heares.2022.108605

70. Amezquita R, Lun A, Hicks S and Gottardo R (2021). Advanced Single-Cell Analysis with Bioconductor. Version 1.2.1. http://bioconductor.org/books/3.14/OSCA.advanced/index.html

71. Germain P, Lun A, Macnair W and Robinson M (2021). Doublet identification in single-cell sequencing data using scDblFinder. *f1000research*. doi: 10.12688/f1000research.73600.1.

72. Krishnaswami SR, Grindberg RV, Novotny M, Venepally P, Lacar B, Bhutani K, Linker SB, Pham S, Erwin JA, Miller JA, Hodge R, McCarthy JK, Kelder M, McCorrison J, Aevermann BD, Fuertes FD, Scheuermann RH, Lee J, Lein ES, Schork N, McConnell MJ, Gage FH, Lasken RS. Using single nuclei for RNA-seq to capture the transcriptome of postmortem neurons. *Nat Protoc*. 2016 Mar;11(3):499-524. doi: 10.1038/nprot.2016.015.

73. Lun ATL, McCarthy DJ and Marioni JC (2016). A step-by-step workflow for low-level analysis of single-cell RNA-seq data with Bioconductor. *F1000Res*. 5, 2122. doi: 10.12688/f1000research.9501.2

74. McCarthy DJ, Campbell KR, Lun ATL and Willis QF (2017). Scater: pre-processing, quality control, normalisation and visualisation of single-cell RNA-seq data in R. *Bioinformatics*, 33, 1179-1186. doi: 10.1093/bioinformatics/btw777.

75. Traag VA, Waltman L, Van Eck NJ (2018). From Louvain to Leiden: guaranteeing well-connected communities. *arXiv*:1810.08473

76. Zheng, Grace XY, Terry JM, et al. (2017). Massively parallel digital transcriptional profiling of single cells. *Nature Communications*. 8: 1-12, doi:10.1038/ncomms14049

77. Moffitt, J. R., & Zhuang, X. (2016). RNA imaging with multiplexed error-robust fluorescence in situ hybridization (MERFISH). In *Methods in enzymology* (Vol. 572, pp. 1-49). Academic Press.

78. Zhuang, X. (2021). Spatially resolved single-cell genomics and transcriptomics by imaging. *Nature methods*, *18*(1), 18-22.

Fig.1: Phenotyping CFW outbred mice shows eight distinct patterns of hearing. Auditory Brain Response (ABR) thresholds measure the health of hearing function and correspond to the response to sound stimulus with known intensity at a specific frequency. The mouse is considered to have mild-moderate hearing impairment if ABR thresholds were between 41 - 60 dB, severe if between 61 – 80 dB, and profound if greater than 80 db. **A** Normal hearing. **B** Moderate mid-frequency (12 kHz or 16 kHz) loss. **C** Moderate high-frequency (24 kHz or 32 kHz) loss. **D** Severe high-frequency (only 32 kHz) loss. **E** Severe high-frequency (24 kHz + 32 kHz) loss. **F** Moderate all-frequency loss. **G** Severe all-frequency loss. **H** Profound all-frequency loss. **I** individual ABR plots for mice selected to collect cochlear tissue for snRNA seq processing.

Figure 2 – Sixty snRNA-seq transcriptomic clusters from mouse cochlea. A) Iterative unsupervised clustering of the processed cell-by-gene snRNA-seq expression matrix performed using Leiden community detection identified sixty distinct transcriptomic clusters (numbers). tSNE embedding of the clustering results is shown. Cell type annotations of each transcriptomic cluster based on subsequent analysis (Table 1) is shown. The data can be explored   at https://umgear.org/index.html?multigene_plots=0&layout_id=c2acc279&gene_symbol_exact_match=1&gene_symbol=slc26a5 . B) Heatmap of marker genes known to be expressed in specific ear cell types arranged according to hierarchical clustering of transcriptomic clusters, which captures the transcriptional relationships between the discrete clusters. The horizontal banding patterns of these known marker genes validates the structure of the hierarchical clustering taxonomy. C) Box plots of the proportion (log10 transformed) of each cell type across the 48 outbred mouse cochlear samples, with median proportion ranging from $\sim 10^{-4}$ for Cluster 22.1 to $\sim 10^{-1}$ for Cluster 4.

Figure 3 – Cell type-specific marker genes. NS-Forest was used to identify the minimum set of necessary and specific marker genes for each cell type, with gene expression (rows) across each transcriptomic cluster (columns) illustrated in violin plots. A) Minimum NS-Forest marker genes identified from the complete cell-by-gene expression matrix, with diagonal expression patterns and little off-diagonal expression showing marker gene specificity. In addition to the minimum site of marker genes, NS-Forest also produces an extended set of ten genes selected based on binary expression (on-off) patterns in the target cluster. B) Top ten binary genes were identified within the four Cluster 23 iterative subclusters using the pooled Cluster 23 subcluster cell-by-gene expression matrix as input, showing strong and distinct binary expression patterns, especially for Clusters 23.3 and 23.4.

Figure 4 – Spatial organization of cell types in the mouse cochlea. MERFISH images with selected marker genes for Cluster 23 (cluster of HC) (A-C), Cluster 20 (cluster of supporting cell) (D-F), Cluster 12 (cluster of spiral ganglion cell) (I-L) and Cluster 17 (cluster of Schwann cell) on the cochlear mid-modiolar cryosections at P4. Each pseudo-color dots represent transcripts of different marker genes and are merged with cell-boundary staining in dark grey. Cell boundaries are depicted by white or white-dashed line. Myo7a was used for pan-HC marker gene (yellow) (A-F). A) Cluster 23.1 for IHC with Gm113(magenta, arrow), Ofcc1(blue, arrowhead). B) Cluster 23.2 for OHC with Slc26a4 (red, arrow). C) Cluster 23.4 for unknown hair-cell subtype with Ripor3 (green, arrow). D) Cluster 20.1 for inner border cell with Dgkb (green, arrowhead) and Klhl14 (magenta, arrow). E) Cluster 20.2 for pillar cell with Mdm1 (red, arrow). F) Cluster 20.3 for inner phalangeal cell (white line) and Deiter's cell (white-dashed line) with Dgkb (green, arrowhead), Nckap5 (blue) and Slc1a3(orange, arrowhead). I) Localization of marker genes for cluster 12 in cochlear spiral ganglion cells (arrow) with Ntng1 (yellow), Mdga1 (light blue) and Meg3 (dark blue).

H. I. Binary color images of cluster 12.1 with Ntng1(arrow) and C130073E24Rik (thick arrow), of cluster 12.2 with Cdh9 (double-head arrow), Tmem108 (arrowhead), and Mdga1, and of cluster 12.3 with Meg3, and merged image with pseudo colors (I). Representative cell boundaries for each subcluster are depicted by white-dot (cluster 12.1) and white-dashed (cluster 12.2), and asterisk indicates a cell of cluster 23.3. Mpz (brown) is a marker gene for Schwann cell. M. Images for cluster 17.1 with Pde1c (red, arrowhead) and Lama1(green, thick arrow), and for cluster 17.2 with Ntng1 (light blue, double-head arrow) and Cdh19 (yellow, arrow). Meg3 (dark blue) and Mpz (brown) indicate spiral ganglion cells and Schwann cells, respectively. Sale bars; 15μm (A-F), 250μm (G), and 25μm (H-J).

Figure 5 – Loss of specific cochlear cell types with age-related hearing loss. The proportion of each cell type in the 48 outbred mouse cochlea was compared against the ABR sound intensity score at different sound frequencies. A) Correlation between selected cell type proportions (Y-axis) and ABR sound intensity thresholds at 32 kHz (x-axis). HC Clusters 23.2 and 23.4 and ganglion neuron Clusters 12.1 and 12.2 show significant inverse correlations with intensity threshold (red). Basal cell Clusters 2.1 and 2.2 show significant positive correlation with intensity threshold (blue). In contrast to the inverse correlation observed with Clusters 23.2 and 23.4, HC Clusters 23.1 and 23.3 show no correlation with intensity threshold. Spearman's rho correlation and p-value are show at the top of each plot. B) Sound intensity threshold of individual mice ranked in order of hearing loss. C) Proportion of Cluster 23.2 cells in individual mice. D) Proportion of Cluster 23.4 cells in individual mice. Whole mount projections of organ of Corti corresponding to areas encoding 16 kHz frequency show that mice with profound hearing loss (F)

exhibit IHC and OHC loss that worsens through the higher frequencies while the organ of corti from mice with normal hearing (E) show intact sturctures with no HC loss.

Hierarchical cluster order	Cluster ID	Cluster name	Cell type name	FR-Match high confidence	FR-Match low confidence	MERFISH cell type | anatomic structure	enrichR
1	20.2	cochlear pillar cell	cochlear pillar cell Mdm1		root	inner / outer pillar cell | epithelium of cochlear duct	mixed
2	20.1	cochlear border cell	cochlear border cell Dgkb Klhl14		root	border cell of cochlea | epithelium of cochlear duct	mixed
3	20.3	cochlear phalageal cell	cochlear phalageal cell Dgkb Nckap5		root	inner phalangeal cell / Deiters' cell | epithelium of cochlear duct	
4	16	cochlear interdental cell	cochlear interdental cell Otoa Ceacam16		root/Deiter	interdental cell of cochlea | tympanic lip of limbus of osseous spiral lamina	lymphoid
5	21	cochlear supporting cell	cochlear supporting cell Gpc5 Otogl		root/Deiter	supporting cell of cochlea | epithelium of cochlear duct	
6	10.1	cochlear sulcus cell 1	cochlear inner / outer sulcus cell Prss36 Hmcn1	root	root	border cell / Claudius cell | spiral sulcus	
7	10.2	cochlear sulcus cell 2	cochlear inner / outer sulcus cell Adamtsl1 Col11a1	root	root	border cell / Claudius cell | spiral sulcus	
8	6	cochlear root cell	cochlear root cell Lgr5 Slc26a4	root	root	root cell | outer spiral sulcus	
9	10.3	cochlear sulcus cell 3	cochlear inner / outer sulcus cell Rasgef1b Lmo3	root	root	border cell / Claudius cell | spiral sulcus	
10	5	cochlear marginal	cochlear marginal Stac Kcnq1	marginal	marginal	strial marginal cell | stria vascularis of cochlear duct	epithelial
11	4	cochlear Reissner's membrane	cochlear Reissner's membrane Gm48447	Reissner's	Reissner's	epithelial cell | Membrane of Reissner	retinal/sensory
12	19.1	cochlear spindle 1	cochlear spindle Gm43154 Agbl1	spindle	spindle	spindle cell | spiral prominence of cochlear duct	
13	19.2	cochlear spindle 2	cochlear spindle Dpp10 Anxa1	spindle	spindle	spindle cell | spiral prominence of cochlear duct	
14	22.1	cochlear neutrophil 1	cochlear neutrophil S100a8 Hba-a1		neutrophil	neutrophil | cochlea	neutrophils
15	22.2	cochlear neutrophil 2	cochlear neutrophil S100a8 Retnlg	neutrophil	neutrophil	neutrophil | cochlea	
16	13.2	unknown 1	unknown Dnah12 Rgs22			?? | ??	
17	13.6	unknown 2	unknown Bpifa1			?? | cochlear modiolus	mixed
18	24	cochlear erythroblast	cochlear erythroblast Kel	erythroblast	erythroblast	erythroblast | cochlea	erythroblast
19	1.6	cochlear NK/T cell	cochlear NK/T cell Gm2682 Skap1			mature NK T cell | cochlear nerve	NK/T cell
20	1.8	cochlear CD8+ T cells	cochlear CD8+ T cells Runx3 Grap2			CD8-positive, alpha-beta T cell | cochlea	CD8+ T cells
21	1.3	cochlear leukocyte	cochlear leukocyte Rnf220 Bcl11a			leukocyte | cochlea	
22	1.5	cochlear B cell	cochlear B cell Chst3 Pax5	B cell	B cell	B cell | cochlear nerve	B cell
23	1.1	cochlear macrophage 1	cochlear macrophage 1	Macrophage	macrophage	tissue-resident macrophage | cochlea	various immune cell types
24	1.9	cochlear macrophage 2	cochlear macrophage 2 Spic			tissue-resident macrophage | cochlea	macrophage
25	14	cochlear neutrophil 3	cochlear neutrophil Abca13 Adpgk	neutrophil	neutrophil	neutrophil | cochlea	myeloid/neutrophil
26	1.2	cochlear neutrophil 4	cochlear neutrophil F13a1 Lyn			neutrophil | cochlea	

#	ID	Cell type	Markers			Annotation	Category
27	1.4	cochlear leukocyte	cochlear leukocyte Rab44 Cdk6			leukocyte	cochlea	myeloid/neutrophil
28	1.7	cochlear myeloid/neutrophil	cochlear myeloid/neutrophil Gm20528 Tspoap1			myeloid leukocyte	cochlea	melanocytes
29	9	cochlear Intermediate	cochlear Intermediate Dct	Intermediate	intermediate	strial intermediate cell	stria vascularis of cochlear duct	oligodendrocyte
30	25	cochlear oligodendrocyte	cochlear oligodendrocyte Prr5l Mog		oligodendrocyte	oligodendrocyte	cochlear nerve	stromal
31	17.2	cochlear Schwann cell 1	cochlear Schwann cell Ntng1 Cdh19			Schwann cell	cochlear ganglion / cochlear nerve	neuron
32	17.1	cochlear Schwann cell 2	cochlear Schwann cell Pde1c Lama1			Schwann cell	cochlear ganglion / cochlear nerve	Schwan cell
33	18	cochlear Schwann cell 3	cochlear Schwann cell Mpz Gm12068	glia	glia/Schwann cell	Schwann cell	cochlear ganglion / cochlear nerve	myocytes
34	23.1	cochlear inner hair cell	cochlear inner hair cell Gm1113 Ofcc1			cochlea inner hair cell	epithelium of cochlear duct	neuron/ganglion
35	23.3	cochlear hair cell 1	cochlear hair cell Ush2a C230072F16Rik			cochlea auditory hair cell	epithelium of cochlear duct	mixed
36	23.2	cochlear outer hair cell	cochlear outer hair cell Slc26a5	outer hair cell	outer hair cell	cochlear outer hair cell	epithelium of cochlear duct	mixed
37	23.4	cochlear hair cell 2	cochlear hair cell Ripor3			cochlea auditory hair cell	epithelium of cochlear duct	astrocyte
38	26	cochlear astrocyte	cochlear astrocyte Gabrb1 Slc39a12		Hensens	astrocyte	cochlear nerve	
39	12.3	cochlear spiral ganglion neuron 1	cochlear spiral ganglion neuron 3 Meg3			spiral ganglion neuron	cochlear ganglion	neuron/ganglial cell
40	12.1	cochlear spiral ganglion neuron 2	cochlear spiral ganglion neuron 1 C130073E24Rik Ntng1			spiral ganglion neuron	cochlear ganglion	neuron/ganglial cell
41	12.2	cochlear spiral ganglion neuron 3	cochlear spiral ganglion neuron 2 Mdga1 Tmem108			spiral ganglion neuron	cochlear ganglion	
42	2.1	cochlear basal 1	cochlear basal Sorcs3 Spaar	basal	basal	basal cell	spiral ligament	
43	2.2	cochlear basal 2	cochlear basal Dync1i1 Gm12153	basal	basal	basal cell	cochlear ganglion	
44	7.3	cochlear fibrocyte 1	cochlear fibrocyte Kcnk2	fibrocyte	fibrocyte	fibrocyte	spiral ligament	
45	7.1	cochlear fibrocyte 2	cochlear fibrocyte Slc8a3 Ucma	fibrocyte	fibrocyte	fibrocyte	spiral ligament	
46	7.2	cochlear fibrocyte 3	cochlear fibrocyte Slc4a10 Slc4a11	fibrocyte	fibrocyte	fibrocyte	spiral ligament	
47	13.5	cochlear endothelial	cochlear endothelial Ptprb	fibrocyte	fibrocyte	endothelial cell	spiral modiolar artery	ciliated epithelial/endothelial
48	13.3	cochlear ciliated epithelial 1	cochlear ciliated epithelial Lgr6 Fam78b	fibrocyte	fibrocyte	ciliated epithelial cell	??	ciliated epithelial/osteocytes
49	13.1	cochlear ciliated epithelial 2	cochlear ciliated epithelial Satb2 Fap	fibrocyte	fibrocyte	ciliated epithelial cell	cochlear modiolus	ciliated epithelial
50	13.8	cochlear ciliated epithelial 3	cochlear ciliated epithelial Cxcl12 Svep1	fibrocyte	fibrocyte	ciliated epithelial cell	cochlear modiolus	ciliated epithelial/fibroblast
51	13.4	cochlear ciliated epithelial 4	cochlear ciliated epithelial Mecom	fibrocyte	fibrocyte	ciliated epithelial cell	??	ciliated epithelial
52	13.7	cochlear smooth muscle cell	cochlear smooth muscle cell Mrvi1 Trpc3	fibrocyte	fibrocyte	smooth muscle cell	spiral modiolar artery	ciliated epithelial/smooth muscle
53	15	cochlear glial	cochlear glial Slc6a13	fibrocyte	fibrocyte	glial cell	cochlear nerve	glial
54	3.2	cochlear basal 3	cochlear basal Bnc2 Fn1	fibrocyte	basal	basal cell	cochlea	
55	3.1	cochlear basal 4	cochlear basal Slc47a1 Slc4a10	fibrocyte	basal	basal cell	cochlea	

56	3.3	cochlear basal 5	cochlear basal Clic4 Gm26740	fibrocyte	basal	basal cell	cochlea	
57	8.3	cochlear fibrocyte 4	cochlear fibrocyte Themis Itga8	fibrocyte	fibrocyte	fibrocyte	cochlear ganglion	
58	11	cochlear fibrocyte 5	cochlear fibrocyte Mybpc1 Nav3	fibrocyte	fibrocyte	fibrocyte	basilar membrane of cochlea	various stromal cell types
59	8.1	cochlear fibrocyte 6	cochlear fibrocyte Sorcs3 Lama2	fibrocyte	fibrocyte	fibrocyte	cochlear ganglion	
60	8.2	cochlear fibrocyte 7	cochlear fibrocyte Itga8 Slit2	fibrocyte	fibrocyte	fibrocyte	spiral ligament / cochlear ganglion	