Barley shows reduced Fusarium Head Blight under drought and modular expression of differential expressed genes under combined stress

Felix Hoheneder[1]*, Christina E. Steidele[1]*, Maxim Messerer[2], Klaus Mayer[2], Nikolai Köhler[1,3], Christine Wurmser[4], Michael Heß[1], Michael Gigl[5], Corinna Dawid[5], Remco Stam[1,6], Ralph Hückelhoven[1]§

* these authors contributed equally and are listed alphabetically

§ correspondence: hueckelhoven@tum.de

[1] Chair of Phytopathology, TUM School of Life Sciences, HEF World Agricultural Systems Center, Technical University of Munich, Emil-Ramann Str. 2, 85354 Freising-Weihenstephan, Germany
[2] Plant Genome and Systems Biology, Helmholtz Center Munich, German Research Center for Environmental Health, Ingolstädter Landstraße 1, 85764 Neuherberg, Germany
[3] LipiTUM, Chair of Experimental Bioinformatics, TUM School of Life Sciences, Technical University of Munich, Maximus-von-Imhof Forum 3, 85354 Freising-Weihenstephan, Germany
[4] Chair of Animal Physiology and Immunology, TUM School of Life Sciences, Technical University of Munich, Weihenstephaner Berg 3/I, 85354 Freising-Weihenstephan, Germany
[5] Chair of Food Chemistry and Molecular Sensory Science, TUM School of Life Sciences, Technical University of Munich, Lise-Meitner-Straße 34, 85354 Freising-Weihenstephan, Germany
[6] Institute of Phytopathology, Christian Albrecht University of Kiel, Hermann-Rodewald-Straße 9, 24118 Kiel, Germany

Running title: Barley Fusarium Head Blight under drought stress

#tables: 0
#figures: 4
#supplemental tables/data: 6
#supplemental figures: 3

word count: 5664

date of submission: 15.02.2023

Current email addresses:

Felix Hoheneder:	felix.hoheneder@tum.de
Christina E. Steidele:	christina.steidele@tum.de
Maxim Messerer:	maxim.messerer@helmholtz-muenchen.de
Klaus Meyer:	k.mayer@helmholtz-muenchen.de
Nikolai Köhler:	nikolai.koehler@tum.de
Christine Wurmser:	christine.wurmser@tum.de
Michael Heß:	michael.hess@uba.de
Michael Gigl:	michael.gigl@tum.de
Corinna Dawid:	corinna.dawid@tum.de
Remco Stam:	remco.stam@phytomed.uni-kiel.de
Ralph Hückelhoven:	hueckelhoven@tum.de

ORCID:
Felix Hoheneder (0000-0001-9541-983X)
C. Steidele (0000-0003-1284-6604)
Maxim Messerer (0000-0003-0554-5045)
Klaus Mayer (0000-0001-6484-1077)
Nikolai Köhler (0000-0002-5103-883X)
Christine Wurmser
Michael Heß (0000-0003-1202-5224)
Michael Gigl (0000-0002-5617-9236)
Corinna Dawid (0000-0001-5342-2600)
Remco Stam (0000-0002-3444-6954)
R. Hückelhoven (0000-0001-5632-5451)

Highlight

Co-expression network analysis reveals association of physiological stress markers and gene
expression modules under biotic and abiotic stress

Abstract

Plants often face simultaneous abiotic and biotic stress conditions. However, physiological and
transcriptional responses of plants under combined stress situations are little understood.
Spring barley is susceptible to Fusarium Head Blight (FHB), which is strongly affected by
weather conditions. We therefore studied the potential influence of drought on FHB severity
and responses in three differently susceptible spring barley varieties and found strongly
reduced FHB severity in susceptible varieties under drought. Quantity of differentially
expressed genes (DEGs) and strength of transcriptomic regulation reflected the concentration
of physiological stress markers such as abscisic acid or fungal DNA contents. Infection-related
gene expression associated rather with susceptibility than resistance. Weighted gene
correlation network analysis uncovered 18 modules of co-expressed genes, which reflect the
pathogen or drought response in the used varieties. A generally infection-related module
contained co-expressed genes for defence, programmed cell death and mycotoxin-
detoxification indicating that diverse genotypes use a similar defence strategy towards FHB
albeit with different success. Further DEGs showed co-expression in drought or genotype-
associated modules correlating with measured phytohormones or the osmolyte proline. The
combination of drought stress with infection lead to highest numbers of DEGs and provoked a
modular composition of single stress responses rather than a specific transcriptional readout.

Keywords and Abbreviations

barley, drought stress, Fusarium head blight, 3'-RNAseq, transcriptome, weighted gene co-
expression network analysis

Abbreviations:

28	ABA	= abscisic acid
29	Ba	= Barke
30	DFc	= drought stress, infected with *Fusarium culmorum*
31	DEG	= differently expressed gene
32	DM	= drought stress, mock-inoculated
33	DPA	= dihydrophaseic acid
34	*F.*	= *Fusarium*
35	Fc	= *Fusarium culmorum*
36	FHB	= Fusarium Head Blight
37	IAA	= indol 3-acetic acid
38	Mo	= Morex
39	PA	= phaseic acid
40	Pb	= Palmella Blue
41	SA	= salicylic acid
42	SOTA	= self-organizing tree algorithm
43	WFc	= watered, infected with *Fusarium culmorum*
44	WGCNA	= weighted gene correlation network analysis
45	WM	= watered, mock-inoculated
46	WRKY	= WRKY transcription factor

Introduction

Under natural conditions, plants are facing multiple abiotic and biotic stress factors affecting growth, crop yield and pathogen defence. The simultaneous occurrence of two or more stress factors triggers complex regulatory responses resulting in synergistic, antagonistic or neutral effects on expression and balancing of stress responses (Choi *et al.* 2013; Zhang and Sonnewald 2017; Pandey *et al.* 2015; Ramegowda and Senthil-Kumar 2015; Pandey and Senthil-Kumar 2019). Many stress responses share common factors of plant signalling and metabolism, and co-occurrence of stressors is often fatal for crop productivity (Farooq *et al.* 2009; Suzuki *et al.* 2014; Zandalinas *et al.* 2021). Transcriptional analyses in *Arabidopsis thaliana* indicate that plants respond to combined abiotic and biotic stress with tailored regulatory responses differing from responses towards single stressors (Gupta *et al.* 2016). Stress-stress interactions are often complex and not well understood (Atkinson and Urwin 2012). One of the most limiting factors of plant growth, performance and crop productivity is water availability (Kang *et al.* 2009). In conjunction with global warming, strong and frequent drought periods are predicted to increase in near future and represent serious threats to food security and supply (Meza *et al.* 2020) and especially spring barley will strongly be affected by droughts (Olesen *et al.* 2011; Xie *et al.* 2018). A recent study further predicts that on a global scale proportions of fungal soil borne pathogens will increase with global warming (Delgado-Baquerizo *et al.* 2020). Hence, vulnerability of agriculture may rise with a simultaneously predicted increase in global food demands leading to increasing gaps in food security. In this context, understanding plant physiology under complex stress combinations could unfold regulatory pathways and marker genes important for advanced genotype selection and breeding of disease resistant and stress tolerant crops (Pandey and Senthil-Kumar 2019).

Fusarium head blight (FHB) is one of the most destructive diseases of small grain cereal crops worldwide and its epidemiology is strongly associated with weather conditions. The disease is caused by a pathogen complex of several *Fusarium* species such as *F. graminearum* and *F. culmorum*. Infections with *Fusarium* pathogens can cause yield loss but also quality losses and contamination with hazardous mycotoxins like deoxynivalenol (DON) or zearalenone (Wegulo *et al.* 2015). The occurrence and strength of the disease is dependent on multiple environmental factors especially temperature, air humidity and rainfalls around anthesis, which collectively influence disease incidence and severity in the field (Birr *et al.* 2020; Hoheneder *et al.* 2022). FHB resistance of wheat (Buerstmayr *et al.* 2021; Mesterhazy 2020; Buerstmayr and Lemmens 2015) and barley (Buerstmayr and Lemmens 2015; Ogrodowicz *et al.* 2020) is a quantitative trait expressed from various QTLs and hence difficult to exploit. Furthermore, effective chemical control of FHB is limited to a short time window at around anthesis, when primary head infection usually occurs. Additionally, control measurements and pathogenicity of the fungus strongly interfere with weather conditions, both increasing complexity of effective FHB management (Wegulo *et al.* 2015).

In order to obtain physiological markers for FHB responses under complex environmental conditions, understanding gene regulation networks in response to single and combined abiotic and biotic stressors is crucial. The transcriptional response of wheat and barley to FHB is increasingly well understood (Hameed *et al.* 2022; Kazan and Gardiner 2018) but little information is available for complex stress situations. Hormones regulate various stress responses and physiological and developmental processes in plants. Under drought, plant hormones act as notable endogenous plant growth regulators and mediate abiotic stress tolerance (Farooq *et al.* 2009) and pathogen defence (Bari and Jones 2009). Predominantly, abscisic acid (ABA) governs growth, root-shoot ratio and regulates stomatal conductance to reduce transpiration and water loss under drought over relatively long distances from the root to the shoot. However, while ABA improves abiotic stress tolerance during vegetative growth, the downstream effects of ABA during reproductive stage can be contrary (Dolferus *et al.*

2011). There is evidence that reproductive tissue is most sensitive to water deficit in general (Blum 2009) and drought stress strongly delays flowering or even leads to abortion of spikelets in cereal crops (Barnabás *et al.* 2008). When it comes to pathogen infection, the genetic regulatory mechanisms to balance hormonal regulations associated with specific stress responses are not sufficiently investigated, especially when invading microbes are able to manipulate regulatory networks in the host by biosynthesis of phytohormones (Bari and Jones 2009), which was previously described for *Fusarium* species (Dörffling and Petersen 1984; Jaroszuk-Ściseł *et al.* 2014; Luo *et al.* 2016; Qi *et al.* 2016) Salicylic acid (SA), jasmonic acid (JA) and ethylene are mainly involved in diverse pathogen defence responses against various pathogens and expression of pathogenesis related (PR) genes (Bari and Jones 2009). ABA induces several transcription factors, which contribute to specific abiotic stress but also pathogen defence regulations (Yao *et al.* 2021). SA is further associated as signal for systemic acquired resistance mechanisms. However, the function of hormonal defence signalling network is dependent on the pathogen´s trophic lifestyles (Adie *et al.* 2007), which is often not clearly classifiable and changes during pathogenesis of the hemibiotrophic *Fusarium* species from biotrophic to necrotrophic (Brown *et al.* 2010). However, the antagonistically acting SA and JA/ethylene associated pathways are both involved in responses to *Fusarium* infections (Wang *et al.* 2018). In particular, when it comes to combined stress, functions of either synergistically or antagonistically interacting plant hormones provoke complex stress responses. Hence, the host plant may prioritize particular stress responses, which can lead to either resistance or susceptibility (Bari and Jones 2009; Wang *et al.* 2018; Gupta *et al.* 2020). Although, previous studies revealed associations of ethylene (Chen *et al.* 2009; Xiao *et al.* 2013) auxin (Brauer *et al.* 2019) or ABA (Buhrow *et al.* 2021; Qi *et al.* 2016) with susceptibility or SA (Makandar *et al.* 2012), JA (Sun *et al.* 2016) and gibberellic acid (Buhrow *et al.* 2021) with resistance against *Fusarium* spp., roles of phytohormones remain complex and ambiguous. It was further shown, that their role during infection with *F. graminearum* strongly depends on distinct stages of infection demonstrating strong interactions of the pathogen with plant physiology (Ding *et al.* 2011; Ameye *et al.* 2015; Makandar *et al.* 2010; Makandar *et al.* 2012). Although, several biological and physiological mechanisms of *Fusarium* spp.-barley pathosystem were described, understanding of FHB susceptibility and resistance under complex environmental stress conditions poses new challenges but could support future breeding and selection of pathogen resistant and stress tolerant genotypes. For this purpose, the present study aims to investigate genotype-dependent quantitative resistance and regulatory networks of barley under infection with *Fusarium culmorum* in combination with drought stress. This revealed genotype-independent stress response markers, infection success-related gene expression clusters and a modular gene expression network under combined stress.

Material and methods

Description of greenhouse experiments
Three spring barley genotypes (Barke, Morex and Palmella Blue) were grown under controlled conditions in glass house cabins with air conditioning for temperature control. The genotypes were preselected according to different resistance to FHB as assessed in inoculation trials in the field (Hoheneder *et al.* 2022) and preliminary greenhouse.
Six barley grains were sown in each pot containing 3 L peat substrate (Einheitserde C700, Stender, Germany). 12 pots were prepared per barley genotype and stress treatment. We applied daily automatic watering and additional lightning for 16 h per day. Plants were randomized twice a week. Temperature was set to 18 °C (day) and 16 °C (night) with a relative air humidity of 60%.
Drought conditions were set for half of the pots from seven days before expected flowering (growth stage 65) on by a stop of automatic watering on separate flooding tables. To prevent

plants from early and premature plant death due to fast desiccation of the substrate, each pot
received little amount of water (50 mL) in the first three days after stopping daily irrigation. All
plants without irrigation showed reduced turgor pressure and partial loss of green leaf area
indicating strong drought stress of the plants. Continuously watered plants showed normal
growth and phenology.
Seven days after the beginning of drought conditions, spikes of irrigated or drought stressed
plants were sprayed with spray flasks till run-off either with *F. culmorum* spore solution or mock
solution to obtain four different treatment contrasts (WM: watered-mock, WFc: watered-
infected, DM: drought-mock, DFc: drought-infected). To maintain optimal moist conditions for
infection (99% relative air humidity), spikes were covered and sealed with transparent
polythene bags for two days. Respective mock sprayed plants were similarly treated. Two and
four days after inoculation, individual spikes were cut from each pot and flash frozen in liquid
nitrogen for further DNA, RNA and metabolite extraction.

162
Preparation of Fusarium culmorum inoculum
Fugal inoculum was cultured and propagated according to (Linkmeyer *et al.* 2013). Therefore,
three isolates of *F. culmorum* (Fc002, Fc03, Fc06 – culture collection, Chair of Phytopathology,
Technical University of Munich) known to strongly infect barley spikes and to produce DON
(Linkmeyer *et al.* 2013; Hofer *et al.* 2016; Hoheneder *et al.* 2022) were combined in equal
amounts. Spore solution was adjusted to 50,000 conidia per mL tap water and contained
1 mL/L Tween 80 to improve wetting of the spike tissue. The respective mock solution
contained the same amount of Tween 80.

171
Sample preparation, DNA and RNA extraction from immature spike tissue
Each individual sample of each biological triplicate per treatment variation was divided into two
pieces using one part for RNA extraction. The remaining spike tissues was put together to a
pooled sample of three spikes for DNA extraction. After separation, the spike samples were
immediately ground in liquid nitrogen and stored at minus 70 °C. DNA extraction was carried
out according to the protocol of (Fraaije *et al.* 1999) with minor modifications as described by
(Hofer *et al.* 2016). DNA concentration was adjusted to 20 ng total DNA μL^{-1} with nuclease free
water. RNA extraction was performed with Direct-zol RNA Miniprep Plus Kit (ZymoResearch,
USA) according to manufacturer's protocol.

181
Quantification of fungal DNA
Fungal and barley DNA was determined with qPCR according to (Nicolaisen *et al.* 2009) using
species specific primers and 10-fold dilution series of pure target DNA as standards. Non
template controls only contained water. Quantitative PCR reactions were carried out using
Takyon Low ROX SYBR 2X MasterMix blue dTTP (Eurogentec, Belgium) with a AriaMx real-
time PCR system (Agilent Technologies, USA). DNA contents of each sample was determined
in duplicates. Finally, fungal DNA was normalized with barley DNA in pg *F. culmorum* DNA
ng^{-1} barley^{-1} DNA^{-1}.

190
Library preparation for Illumina HiSeq2500 sequencing
Preparation of libraries for 3'-RNA sequencing was carried out with Lexogen QuantSeq 3'-
RNA-Seq Library Prep Kit (FWD) (Lexogen, Austria) for Illumina sequencing according to
manufacturer's protocol. Quantification of input RNA was conducted with Qubit Fluorometer
2.0 (Invitrogen, USA) and Qubit RNA BR (broad range) Assay Kit (Invitrogen, USA) according
to manufacturer's protocol in a range of 10 to 1200 ng total RNA. Quantification of library size
was performed with Qubit Fluorometer 2.0 und Qubit DNA High Sensitivity Assay Kit
(Invitrogen, USA). A check for distribution of mRNA fragment size of each library was
performed with am Agilent 2100 Electrophoresis Bioanalyzer (Agilent Technologies, USA).
Final quantification of each library was determined according to Illumina qPCR guide with a
KAPA SYBR Fast Mastermix Low ROX (Peqlab, Germany) in a QuantStudio 5 real-time PCR
system (Applied Biosystems, USA). Final libraries were normalized with elution buffer (Qiagen,
Germany) to a final concentration to 2 nM and pooled with an equal amount of each sample

library. Denaturation and dilution of libraries for HiSeq Clustering was performed according to
protocol A of user guide (Illumina) with a concentration for clustering of 10 pM.
206
Illumina HiSeq2500 sequencing
Sequencing of libraries was carried out on a HiSeq2500 sequencing platform using HiSeq
Rapid SR Cluster Kit v2 (Illumina, USA) and HiSeq Rapid SBS Kit v2 (50 Cycle) with run
parameters set for multiplexed single-reads (read 1: 100 cycles) and single-indexed reads of
7 cycles. HiSeq Control Software 2.2.70 was used for sequencing. Image analysis and base
calling was carried out with Real-Time Analysis (RTA) 1.18.66.4. Fastq-files were generated
with CASAVA BCL2FASTQ Conversion Software v2.20. Raw data have been stored on NCBI-
Gene Expression Omnibus with the accession number GSE223521.
215
3'-RNAseq trimming, mapping and read count
The quality of the raw 3'-RNAseq data was analysed with FastQC
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc). The trimming step was done with
Trimmomatic (Bolger *et al.* 2014) using the parameters ILLUMINACLIP:Illumina-
SE.fasta:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:40. We used
Hisat2 (Kim *et al.* 2015) with parameters -U --rna-strandness FR to map the trimmed data to
the Morex v2 assembly (Monat *et al.* 2019). In order to add an artificial 3'UTR because of the
3'-RNAseq reads, we used a custom script to elongate the last exon either by 3 kb or until the
next gene in the gff file of Morex v2. The counts of the generated bam files were assigned to
genes using featureCounts (Liao *et al.* 2014) with the parameters -t gene -s 1 -M -O.
226
227 *CPM calculation*
228 CPM (counts per million) were calculated using the *edgeR*-package (Robinson *et al.* 2010).
229 The CPMs were calculated from the "DGEList" object with log = FALSE, normalized.lib.sizes
230 = TRUE.
231
Differential gene expression and cluster analysis
The differential gene expression analyses were carried out using the *edgeR*-package
(Robinson *et al.* 2010). We kept all genes with a CPM equal or bigger than 15 in at least 10%
of all samples. To increase the signal-to-noise ratio for the analyses, we pooled the two time
points (48h and 96h post infection) for each sample and calculated the differential expressed
genes with the following contrast formula: "((variety_condition_48h + variety_condition_96h)/2)
- ((variety_watered-mock_48h + variety_watered-mock_96h)/2)" with variety := Barke, Morex,
Palmella Blue and condition := watered-infected, drought-mock, drought-infected.
The cluster-analyses of the DEGs was done with the Multiple Expression Viewer (MeV, (Howe
et al. 2011) using the implemented self-organizing tree algorithm (SOTA) with the default
parameters. As input we used all differentially expressed genes with an FDR < 0.05 in at least
one variety for one certain stress condition. For each analyses the log_2 fold-change values over
all samples for the respective gene sets were used.
245
GO enrichment analysis
GO-enrichment analyses were performed on the differentially expressed genes using the
topGO-R-package (Alexa and Rahnenfuehrer 2022). The GO-terms for each barley gene were
downloaded from https://biit.cs.ut.ee/gprofiler/gost. As gene universe we have used all genes
found in our RNAseq analyses that passed the threshold of having a CPM equal or bigger than
15 in at least 10% of all samples (35657 genes). The enrichment analyses for "molecular
function" and "biological process" was done using the following settings: algorithm="weight01",
statistic="fisher". Shown are the graphs for the 5 most significant nodes.
254
Weighted gene correlation network analysis (WGCNA)
Co-expression analyses were performed using the R-package WGCNA (Langfelder and
Horvath 2008, 2012). First we calculated the mean of the CPM values for the three replicates
for each time point, variety, treatment and stress condition. We then selected the 12818 genes
which showed in the differential gene expression analysis a sig. regulation with an FDR < 0.05

in at least one variety. The mean CPM values of those genes were normalized using variance
stabilizing transformation from the *vsn* R-package (Huber *et al.* 2002). For the WGCNA
analysis we have used the default settings with a soft power of 6 for the calculation of network
adjacency of gene counts and the topological overlap matrix, clustering the genes into 18
modules plus 42 unassigned genes collected in the grey module.
For all the external traits (phytohormones, plant metabolites and fungal DNA) that we have
correlated with our modules, we previously calculated the mean of the three replicates for each
time point, variety, treatment and stress condition. The mean value was than subjected into
the network analysis.
For the network analysis, the 25 edges between any two genes with the highest edge weight
in each module have been selected and the corresponding network of those selected edges
was drawn using the *igraph* R-package (Csardi and Nepusz 2006). For better visualization the
networks were manually adjusted.
273
Measurement of phytohormones and plant metabolites
Contents of phytohormones and secondary plant metabolites (abscisic acid, abscisic acid
glucoside, phaseic acid, dihydrophaseic acid, auxin, salicylic acid, salicylic acid glucoside and
proline) was carried out with mass spectrometry according to methods described in
(Chaudhary *et al.* 2020) and (Abramov *et al.* 2021). For this purpose, 150 mg of grinded barley
spike tissue as used for RNA extraction was used to determine contents of phytohormones
and plant metabolites in relation to fresh weight.
281

Results

*Global transcription analysis reflects variety-dependent susceptibility to FHB and
enhanced resistance under drought*

To study variety-dependent and -independent stress responses in barley and the effect of
drought stress on FHB, we grew three diverse spring barley varieties in pots under controlled
conditions in the greenhouse. The varieties Barke, Morex and Palmella Blue were selected
based on differences in their resistance against FHB in preliminary greenhouse experiments
and in inoculation trials in the field (Hoheneder *et al.* 2022) with Barke showing the most
resistant phenotype and Palmella Blue being the most susceptible variety. Morex showed an
intermediate resistance phenotype and represents the variety with the sequenced reference
genome for the later analysis. Further genotype characteristics and agronomic traits for all
three varieties can be found in the supplemental table T1.

Plants were either continuously irrigated (watered samples) or irrigation was stopped (drought
samples) seven days before flowering (GS 57-59), a time period in which weather conditions
are of pivotal importance for FHB pathogenesis in the field (Hoheneder *et al*, 2022). Inoculation
or mock-treatment of the spikes was carried out with *Fusarium culmorum* (*Fc*) spore
suspension on watered samples (watered-Fc, WFc or watered-mock, WM) or drought samples
(drought-Fc, DFc or drought-mock, DM) at mid of flowering (GS 65) and samples were taken
48 and 96 hours post-infection (hpi) (Fig. 1A).

In order to better understand the underlying differences in response to infection, drought stress
or the combination of drought with infection, we analysed the gene expression of barley spikes
using 3'-RNA-sequencing. Mapping of 3'-RNA reads on the reference genome, identified
transcripts from 41746 (Barke), 42653 (Morex) or 41363 (Palmella Blue) barley genes in our
samples. For 34969 genes, we obtained reliable reads from all three varieties. We compared
for each variety all treatments against the watered, not-infected (watered-mock) samples and
counted all significant (false discovery rate (FDR) corrected $p < 0.05$) differentially expressed
genes (DEGs) (Fig. 1B). There were large variety-dependent differences in the amount of

DEGs and in the distribution of up- and downregulated genes, ranging for example from 229
DEGs in Palmella Blue under drought stress over 953 genes in Morex up to 7447 genes in
Barke. Under infection, by contrast, susceptible Palmella Blue showed most DEGs. In total and
over all genotypes, we found 2949 DEGs after *F. culmorum* infection, 7879 DEGs under
drought, and 10188 DEGs under drought plus infection. Because the gene sets overlap, this
corresponds to a sum of 12818 DEGs (Supplemental data D1.1).

We used quantification of fungal DNA to assess the severity of infection in the same
experiment. Under irrigation we observed strong differences in the amount of fungal DNA from
Barke (Ba, 6 pg *Fc* DNA ng^{-1} barley^{-1} DNA^{-1}) over Morex (Mo, 45 pg *Fc* DNA ng^{-1} barley^{-1}
DNA $^{-1}$) to Palmella Blue (Pb, 70 pg *Fc* DNA ng^{-1} barley^{-1} DNA^{-1}; averages of 48 and 96 h post
inoculation) (Fig. 1C), corresponding well to the previously observed differences in basal
resistance. When compared to this, all three varieties showed a reduced amount of fungal
DNA when the plants were exposed to drought before infection (Ba 5 pg *Fc* DNA ng^{-1} barley^{-1}
DNA^{-1}; Mo 3 pg *Fc* DNA ng^{-1} barley^{-1} DNA^{-1}; Pb 13 pg *Fc* DNA ng^{-1} barley^{-1} DNA^{-1}).

We measured the accumulation of abscisic acid in all spike samples as one of the major
responses to drought stress. We observed low values of abscisic acid in most of the irrigated
samples, whereas the samples from drought-stressed barley showed clearly elevated levels
of ABA, with strong genotype-dependent differences (Ba 4260 ng ABA g^{-1} FW^{-1}; Mo 2057 ng
ABA g^{-1} FW^{-1}; Pb 1578 ng ABA g^{-1} FW^{-1}; averages of 48 and 96 h post inoculation, 9 or 11
days after stop of irrigation) (Fig. 1C).

Despite strong differences in susceptibility to *F. culmorum* infection and in drought responses
as measured by accumulation of ABA and the stress-associated amino acid proline
(Supplemental data D2), our data also revealed DEGs that were regulated in all three
genotypes under either infection-related stress (146 DEGs), drought stress (64 DEGs) or the
combination of both (167 DEGs) (Supplemental data D1.2 - D1.4). Because our genotypes are
diverse in geographic origin and pedigree (Supplemental table T1), those genes may serve as
general variety-independent markers for the core response of barley to the respective stresses.
In support of this, several of the DEGs in those lists are identical or homologous to previously
reported DEGs in other barley or wheat genotypes infected by *Fusarium graminearum* or
suffering from drought. Examples for such generally *Fusarium*-responsive genes are Fusarium
resistant-orphan protein, tryptophan decarboxylases, anthranilate synthase, laccases,
HvWRKY23 and DMR6-like 2-oxoglutarate and Fe(II)-dependent oxygenase genes
(Buerstmayr *et al.* 2021; Low *et al.* 2020; Soni *et al.* 2020; Boddu *et al.* 2006; Boddu *et al.*
2007; Karre *et al.* 2019; Perochon *et al.* 2019; Tucker *et al.* 2021). Additionally, many enzyme
genes that are potentially involved in detoxifying DON are among the commonly infection-
regulated DEGs such as glycosyltransferases and Glutathione-S-transferase genes and a
cysteine synthase (compare (Gardiner *et al.* 2010)). Prominent examples in that respect are
HvUGT13248 (HORVU.MOREX.r2.5HG0384710) and HvUGT6
(HORVU.MOREX.r2.5HG0430540) (He *et al.* 2020; Michlmayr *et al.* 2018; Schweiger *et al.*
2010). In the list of genotype-independent drought-associated DEGs, we find previously
reported dehydrins and late-embryogenesis-abundant protein genes, potential ABA-receptor
complex PP2c protein phosphatases and a downregulated SAUR auxin response protein
gene. 46 of the combined stress-associated variety-independent DEGs are also variety-
independently regulated in one of the single stress situations. Those genes hence reliably
showed the infection-related stress response even under drought and the drought-related
response in additionally infected situations over all barley varieties (Supplemental data D1.5).

Interestingly, there are only two DEGs (HORVU.MOREX.r2.5HG0401150;
HORVU.MOREX.r2.5HG0372030) that show significant regulation in all genotypes under

combined stress but not after one of the single stresses in at least one genotype. Hence, barley
seems to express no genotype-independent specific response to the applied combined stress.

The questions arose whether we find genotype-dependent stress markers in our data sets. We
found many Barke-specific DEGs under drought and many Palmella Blue-specific DEGs after
infection, because those genotypes showed the strongest gene expression responses to the
respective stresses. We therefore more specifically asked whether there are Barke-specific
DEGs under *Fusarium* stress that may explain higher quantitative resistance. This generated
a list of 20 upregulated and 47 downregulated genes, which are neither differentially expressed
in Morex nor in Palmella Blue (Supplemental data D1.6). Vice versa, we found 148 genes that
were upregulated in both susceptible Morex and Palmella Blue but not in Barke (Supplemental
data D1.7). Those lists possibly contain new factors for quantitative resistance or susceptibility
to FHB.

With a self-organizing tree algorithm (SOTA) analysis, DEGs clustered according to their
differential expression pattern over all varieties and treatments (Fig. 1C). Under infection with
F. culmorum, the expression strength of infection-related cluster I11 (2283 genes; 77.4% of all
infection-related DEGs) reflects the amount of fungal DNA (Fig. 1C). Cluster I9 and I10 (223
and 243 genes) show downregulation only in the most susceptible variety Palmella Blue. Taken
together, 93.2% of all infection-related DEGs are in those clusters (I9-I11) and their expression
strength rather reflects disease progression than resistance to FHB.

Under drought stress, the quantity of DEGs and the strength of expression in most of the
clusters reflected the amount of ABA that we measured in individual varieties with Barke
showing the strongest ABA accumulation (all drought-related clusters except D3, Fig. 1B, C).

The combination of drought stress with infection lead to the highest number of DEGs (10188
genes). Most of the clusters (DI1, 2, 4, 5, 9, 10, 11; 6394 genes, 62.8%) mirror very well the
expression pattern under drought stress alone, indicating that those genes are mainly
responsive to the drought stress. Cluster DI3 contains rather *Fusarium*-responsive genes, as
they are not upregulated under drought stress alone, but show a strong infection-strength
dependent regulation, similar to the before mentioned cluster I11. In particular, Palmella Blue,
which is still strongly infected under drought, shows upregulation of genes in cluster DI3.
Additionally, there are 3 clusters with variety-dependent expression that is similar in all stress
situations (DI6 for Palmella Blue, DI7 for Morex and DI8 for Barke).

Taken together, we observed variety-dependent differences both in infection severity and
strength of drought responses and all plants were less infected when plants were exposed to
drought before infection compared to their respective controls. Each single stress provokes a
mostly stress-specific gene expression response and the combination of drought stress and
infection was mainly dominated by the firstly applied drought stress response in our
experiment. However, in highly susceptible Palmella Blue many of combined stress-regulated
DEGs are also infection-related DEGs. In many cases, number of DEGs and strength of gene
regulation reflect an increase of a stress marker such as content of ABA or fungal DNA in the
same samples.

Expression network analysis supports global similarities in the FHB-responses, but
also differences in drought and combined stress responses
400
For further analysis of the DEGs we performed a weighted gene co-expression network
analysis (Zhang and Horvath 2005) to find DEGs that cluster into modules according to their
expression pattern over all samples, time points and treatments (Fig. 2A), resulting in 19

modules of DEGs. 42 DEGs did not match any co-expression pattern (grey module) (Fig. 2A).
The rest of the DEGs clustered into one of the remaining 18 modules, range in size from 63
genes up to 5287 genes in the turquoise module.

We checked for each barley variety and stress the relative contribution of modules to the
overall stress response and displayed the five modules with the highest contribution to the sum
of DEGs (Fig. 2B, supplementary Fig. S1). The blue module contains more than 60% of the
infection-related largely upregulated DEGs for all three varieties. This may indicate that all
three varieties show a similar response to infection with *F. culmorum* despite the fact that p-
value based numbers of DEGs strongly differed. Gene ontology analyses ranked protein
phosphorylation and response to biotic stress as most significantly enriched biological process
(supplementary data D3 and D4A) and heme binding, DNA-binding transcription factor activity
and protein serine/threonine kinase activity as most significantly enriched molecular functions
in the blue module (supplementary data D3 and D4) possibly reflecting a general pathogen
response in the blue module.

The drought stress response in Barke and Morex is dominated by the turquoise co-expression
module but also the green module contributes to the drought response in all genotypes. In
Palmella Blue the salmon module contains most DEGs but the turquoise module is also
represented. Gene ontology analyses identified photosynthesis related functions as most
significantly enriched in the turquoise module, and most of the genes are downregulated under
drought (supplementary data D1.3, D3 and D4). The green module contains many similarly
up-regulated genes and is most significantly enriched in gene ontology terms photorespiration,
response to stress ending in defence response and embryo development ending in seed
dormancy, and the salmon module with protein dephosphorylation and response to water
(supplementary data D3 and D4).

After the combination of drought stress and infection most of the DEGs in Palmella Blue cluster
in the blue module as seen for infection alone. In Barke and Morex the turquoise module is still
the biggest one. Interestingly, none of the modules showed a significant association with the
trait "combined stress" (see "DFc against all" in supplemental Figure S2A).

Additionally, we can also see genotype-related modules such as the yellow module being
regulated in Barke under all three stress conditions but contributing less to the sum of DEGs
in Morex and Palmella Blue DEGs (Fig. 2B, supplemental Figure S1).

The contribution pattern of individual modules to the overall stress response supports the
previous observation that the combination of drought stress with infection is mainly dominated
by the drought stress in Barke, because the distribution of the biggest modules is almost not
altered in Barke under drought stress alone and under drought plus infection. In comparison,
the distribution pattern of DEGs under drought plus infection matches better to infection alone
in Palmella Blue (Fig. 2B, supplemental Figure S1). Morex shows a more complex response,
but the drought stress pattern is largely recovered under combined stress.

Co-expression modules correlate with FHB-severity, phytohormones or stress marker
444
For plant stress responses, little information exists on association of gene co-expression
clusters with quantitative physiological traits of stress responses. We analysed module-trait
associations by correlating WGCNA-module-sample eigengenes with determined contents of
phytohormones and their derivates, abundance of the stress marker proline and the amount
of fungal DNA to identify significant associations (Fig. 3).

The blue module is the only module which has a significant positive correlation with the amount
of fungal DNA and the phytohormone auxin (IAA). The turquoise module shows the strongest
negative correlation with ABA; most genes in this module are downregulated under drought.
The green module has the highest positive correlation with the content of ABA. Similarly, to
the turquoise module, the tan module shows a significant negative correlation with ABA
contents. The salmon module, containing most of the DEGs in Palmella Blue under drought,
shows a significant positive correlation with proline, ABA and the ABA derivates phaseic acid,
dihydrophaseic acid and ABA glucoside. Hence, the module indicates a specific set of DEGs
for stress regulation under drought. Additionally, many modules show by-trend contrasting
correlations with salicylic acid or auxin versus ABA or derivates of ABA. The purple module is
even significantly positively associated with phaseic acid and dihydrophaseic acid but
negatively with salicylic acid. Hence, module-trait associations added functional value in
showing significant linkages between stress physiology and complex gene expression patterns
(Fig. 3).

Co-expression levels reveal potential hub genes in selected modules

For a better understanding of the inter-connection and functionality-related linkage between
the most co-regulated genes within selected modules we took the 25 edges with the highest
edge weight from each of those modules. The edge weight represents the connection strength
between two nodes, corresponding to the co-expression of those two genes.

The blue module is the only one with a significant positive correlation with the amount of fungal
DNA within barley spikes (Fig. 3) and most of the heavy edges are directly connected with the
DON-detoxifying   Glycosyltransferase   HvUGT13248   (HORVU.MOREX.r2.5HG0384710)
(Mandalà *et al.* 2019; Schweiger *et al.* 2010) (Fig. 4A, B, supplemental table D5). All of the
selected genes are infection-specifically expressed and the expression strength partially
reflects the amount of fungal DNA detected in each variety, with Palmella Blue having the
highest amount of fungal DNA detected by qPCR (see Fig. 1C). Most of the genes show a
positive fold change after infection of watered and drought stressed plants, but no significant
regulation under drought stress alone (Fig. 4B). Manual gene reannotations suggested a
possible function of genes in DON-toxin response (HvUGT13248 and a GSTU6-like
Glutathione S-transferase, HORVU.MOREX.r2.7HG0532730), plant defence and regulation of
cell death (e.g. heat shock transcription factor similar to rice lesion mimic gene *spl7*,
HORVU.MOREX.r2.1HG0067240; a NAC transcription factor, similar to OsNAC4 involved in
cell death responses, HORVU.MOREX.r2.3HG0247070; disease resistance protein a, which
represents   a   Toll-interleukin   receptor   1   domain   only   protein,
HORVU.MOREX.r2.2HG0110590) (Kaneda *et al.* 2009; Yamanouchi *et al.* 2002) and of plant
immunity (homologs of Arabidopsis PUB21 and PUB23 plant U-box domain-containing family
proteins, HORVU.MOREX.r2.6HG0521570, HORVU.MOREX.r2.7HG0555030) (Trujillo *et al.*
2008).

The salmon module shows positive correlation with drought stress associated ABA and ABA-
derivates and the drought stress marker proline but no association with fungal infection
success (Fig. 3). The genes corresponding to the 25 highest co-expression values are in all
three varieties upregulated under drought stress, both alone and in combination with infection
(Fig. 4D, supplemental table D5). Typical drought and dehydration stress-associated genes
are   among   the   strongly   co-expressed   genes.   These   are   for   instance   dehydrins
(HORVU.MOREX.r2.6HG0516710, HORVU.MOREX.r2.6HG0516720) or a seed maturation
protein (HORVU.MOREX.r2.7HG0529950). Additionally, we find a phytoene synthase
potentially   involved   in   ABA   biosynthesis   (HORVU.MOREX.r2.5HG0419050)   and
hexosyltransferase similar to galactinol synthase 2, a drought stress tolerance protein that

catalyses raffinose synthesis for osmoregulation (HORVU.MOREX.r2.2HG0090050) (Gu *et al.*
2016; Li *et al.* 2020; Selvaraj *et al.* 2017) and an ABA-related Arabidopsis AtHB-7-like
homeobox transcription factor (HORVU.MOREX.r2.6HG0496860) (Söderman *et al.* 1996).

The green module is significantly correlated with ABA and proline but not fungal infection
success (Fig. 3). The genes corresponding to the 25 highest co-expression values are, similar
to what we found for the salmon module, genes associated with responses to dehydration
stress, like late-embryogenesis abundant proteins (HORVU.MOREX.r2.1HG0049080,
HORVU.MOREX.r2.1HG0049090,   HORVU.MOREX.r2.3HG0213440,
HORVU.MOREX.r2.5HG0352360), potential osmoregulatory sodium/hydrogen_exchanger
(HORVU.MOREX.r2.7HG0560260) (Brini *et al.* 2007) or desiccation-induced 1VOC proteins
(HORVU.MOREX.r2.4HG0322480) (Mulako *et al.* 2008) (Fig. 4E, supplemental table D5). A
lot of the genes show a stronger regulation in Barke compared to the two others varieties
potentially reflecting higher ABA contents in Barke under drought (Fig. 4F).

The turquoise module is by far the biggest module (5287 genes) (Fig. 2A) and is negatively
correlated with ABA (Fig.3). Most of the strongest co-expressed genes are either associated
with photosynthesis or are associated with translation (Suppl. Fig. S3A+B, supplemental table
D5), two process being already known to be downregulated under stress.

The midnightblue module is not associated with any measured trait (Fig. 3), but is among the
modules dominating the drought stress response of especially Palmella Blue (Fig. 2B).
Interestingly, 8 out of the highly connected 11 genes with strongest co-expression are
annotated as heat-shock proteins (Suppl. Fig. S3C+D, supplemental table D5).

Taken together the correlation of the modules and phytohormones or stress-markers is further
strongly supported by the predicted functions of most strongly co-expressed genes in each
module. Corresponding gene annotations support the hypothesis that the blue module may
partly link to DON-responses and cell death during differentially progressing pathogenesis in
the different varieties. The salmon and green module showed a significant correlation with
several typical drought stress markers, like ABA or proline, and also the genes showing the
strongest co-expression in these modules strongly support a key function in the response to
drought stress and ABA.

Discussion

Comparably little is known about the response of plants to a combination of different stresses
and there is a high demand for understanding complex stress responses in crop plants (Rivero
et al. 2022). Severity, frequency and combination of stresses collectively define the likelihood
that stress results in tissue damage or complete organ or plant death (Buchanan 2000).
Literature provides examples for plant responses to diverse simultaneous stresses that act
additive or mutually inhibitory or synergistic (Ben Alaya *et al.* 2021; Loo *et al.* 2022; Rivero *et*
al. 2022). We wanted to better understand how drought, which increasingly often occurs before
flowering and seed set in mid European barley growing areas, may influence FHB infection.
We used diverse spring barley genotypes that have different degrees of quantitative resistance
to FHB under field and glasshouse conditions (Hoheneder *et al.* 2022) to survey a broad
spectrum of possible responses in barley. We observed strong differences in the amount of
the drought stress hormone ABA (Fig. 1C; Supplemental data D2) whereas the drought stress
metabolite and osmolyte proline accumulated more uniformly among the three genotypes
(Supplemental data D2). This might indicate that the genotypes experienced a similar change
of water potential but Barke particularly strongly responded to this by accumulation of ABA and
the strongest drought-related gene expression response. In contrast, Barke reacted least to

F. culmorum infection, although it showed the highest degree of quantitative resistance as
assessed by fungal DNA in infected ears (Fig. 1). By contrast, highly susceptible Palmella Blue
showed most DEGs and strongest upregulation under infection. Morex showed an
intermediate response in both aspects. Therefore, the global gene expression rather reflects a
transcriptional response associated with successful FHB pathogenesis. However, many genes
that are significantly regulated in susceptible barley are also infection-responsive in more
resistant Barke albeit at a lower level and not significantly different from controls (Fig. 1;
Supplemental data D1.1). We speculate that in less susceptible Barke less tissue gets in direct
contact to the developing fungus and our low numbers of DEGs in most resistant Barke may
be biased due to a dilution effect from healthy parts of the tissue. Nevertheless, high numbers
and strong regulation of DEGs in susceptible barley do not reflect efficient defence, an
observation that has been made before for FHB in wheat (Biselli *et al.* 2018; Buerstmayr *et al.*
2021; Wang *et al.* 2018).

Application of drought stress before FHB infection resulted in less successful pathogenesis.
This effect was strong in Morex and moderate in Palmella Blue. In little susceptible Barke,
drought did not have a major impact on FHB development. Accordingly, global gene
expression responses to the fungus were reduced in susceptible barley under drought.
However, in Palmella Blue, the amount of DEGs that appear to be primarily infection-
responsive under combined stress was still high, and genes showed strong upregulation.
Again, strong gene expression responses to fungal infection associate with fungal success,
even if the success is limited under drought.

SOTA analysis allowed for a quick overview about global gene expression patterns. Generally,
SOTA-resolved gene expression suggested that most combined stress regulated clusters
showed a modular gene expression trend. This means that they often reflect gene expression
that similarly appeared under drought or infection alone. We did not find clear indications for a
specific response of barley to the combined stress but data rather support a combination of
the drought and the infection response. If we consider the 2949 infection-related DEGs, we
see that most clusters show a lower responsiveness under combined stress, when compared
to infection stress alone. However, data do not resolve whether this reflects an inhibitory effect
of drought stress on the pathogen response or rather less fungal infection success, although
we maintained optimal infection conditions by covering barley spikes with polythene bags for
two days after inoculation. Morex profited most strongly from drought when considering
reduced fungal infection success. Interestingly, quite some clusters from the combined stress
situation, show also mild pathogen- and drought-responsiveness in Morex (Fig. 1C, clusters
DI5, 7, 9-11). It is tempting to speculate, that in this case the drought response positively added
to the pathogen response and supported effective pathogen defence.

We used the full complexity of our data set and all DEGs to calculate gene co-expression
networks by WGCNA. This uncovered modules of co-expressed genes, which largely reflect
the pathogen response (blue module) or drought response (turquoise, green, salmon modules)
in the used varieties. Similar information was extracted from the SOTA and as such not a
surprise, e.g. infection related clusters of the SOTA analysis were rediscovered in the blue
module of the WGCNA (supplementary dataset D1.1). Added value of the WGCNA derived
from physiological trait association and diving into most strongly co-expressed gene sets
(arbitrarily extracted 25 heaviest edges of big modules). Association of modules with
physiological data of fungal development and of plant stress markers supported that the blue
module is positively correlated with fungal DNA and auxin contents (Fig. 3). Previous reports
from wheat FHB had already reported a connection between FHB susceptibility and auxin
contents or responses (Brauer *et al.* 2019; Luo *et al.* 2016; Wang *et al.* 2018). This could hence
reflect a general association in FHB of *Triticeae*. The blue module subnetwork of 25 most
strongly co-regulated genes centered around the DON-detoxifying glycosyltransferase

HvUGT13248 and more genes with predicted functions in toxin-response, dampening of
immune-responses and programmed cell death regulation (Kaneda *et al.* 2009; Mandalà *et al.*
2019; Buerstmayr *et al.* 2021) (Fig. 4A). In this context it is interesting, that overexpression of
the barley cell death suppressor BAX Inhibitor-1 enhances resistance to *F. graminearum* ear
and crown infection in barley (Babaeizad *et al.* 2009). Since gene selection was based solely
on edge weight in the network, and several discovered genes have partially described function
in *Fusarium* spp. or general immune response, outcome suggests that our pipeline was
suitable for finding potentially pivotal gene functions in big co-expression modules. This is
supported by the fact that also the green and salmon module subnetworks of heavy edges
reflect gene annotations known from drought responses and seed maturation (Fig. 4C, E).
Self-pollinating barley is fertilized already before open flowering and seed set is initiated. We
applied drought stress before open flowering and our data suggest that seed maturation was
accelerated under drought, storage proteins and dehydrins are upregulated and also sugar
osmolytes may enrich. This change in plant resource allocation could have also influenced
fungal infection success because quick ripening might have limited the hemibiotrophic fungus
in access to plant metabolites.

The interplay of biotic and abiotic stress responses is complex and difficult to study. Literature
provides examples of mutually inhibitory or supportive stress responses, such that combined
or consecutive stresses may result in enhanced or reduced stress resistance. Often plant
hormone-triggered pathways act antagonistically in complex interactions with the environment,
which may have evolved as a measure of checks and balances to fine tune responses for
optimal fitness. Our data also show multiple tendencies of opposite correlation of gene
expression in WGCNA modules and salicylic acid versus ABA or auxin versus ABA and related
metabolites (compare grey-blue versus green squares in vertical columns of Fig. 3). Perhaps,
ABA responses and early ripening limited pathogenesis-associated auxin responses and
thereby reduced susceptibility to fungal infection.

The question arises, whether we can generalize the observations we made in terms of drought
associated resistance to fungal infections in barley. Indeed, a recent study found many drought
stress related genes to be differentially expressed during Fusarium crown rot development in
barley and wheat (Su *et al.* 2021). In a similar context, a NAC transcription factor (HvSNAC1),
involved in drought tolerance, was found to mediate resistance towards Ramularia leaf spot
disease of barley (McGrann *et al.* 2015). We observed earlier, that Barke and fourteen other
spring barley genotypes showed enhanced resistance to Ramularia leaf spots when artificially
exposed to long lasting drought stress in the field. However, in open fields, severe Ramularia
leaf spots occurred even in hot and dry summers if single short rain events provided sufficient
leaf wetness (Hoheneder *et al.* 2021). It is hence difficult to generalize that drought would
mitigate severity of fungal infections. The timing of drought stress application may further be
pivotal for its influence on fungal infection success. In preliminary experiments, we found that
drought can also enhance development of FHB disease, when applied simultaneously instead
of before infection. It remains speculative, whether and how altered gene regulation is
depending on the order of occurring abiotic and biotic stress situations. We found several NAC
and WRKY transcription factors regulated under combined stress and clustering in the blue
module (Supplemental data D1.1), which reveals importance of single and versatile acting
genes on multiple stress responses. This and future studies may provide an increasing amount
of data that allows for a deep understanding of how stress responses interact depending on
plant genotype and phenology and timing and combination of stresses.

The complexity of gene expression data is a boon and a bane of combined stress response
analyses. In front of thousands of stress-regulated genes in our study, using diverse genotypes
allowed for identification of genotype-independent stress markers that are reliable both under
single and combined stresses (Supplemental data D1). Computational association of gene

expression modules with physiological or agronomic traits appears helpful for interpretation of
complex data. Enhancing the complexity of experimental setups and data points also supports
co-expression analyses and identification of subnetworks that may function in stress resistance
or susceptibility and can identify trait-associated candidate genes for functional analysis.

Supplementary data

Table T1. Detailed information about the three used barley varieties

Figure S1. Full list of co-expression modules in each variety for each stress treatment

Figure S2. Relationships of consensus module eigengenes with binary comparisons

Figure S3. Co-expression levels reveal potential hub genes in selected modules

Supplemental data D1. List of DEGs

Supplemental data D2. Data table with measured phytohormones, stress markers and fungal
DNA

Supplemental data D3. Statistics of GO term enrichment for molecular function and biological
process in selected modules

Supplemental data D4. Hierarchical trees of GO term enrichment for molecular function and
biological process in selected modules

Supplemental data D5. Gene numbers, abbreviations and CPM values for genes with the
highest co-expression values in selected module

Acknowledgements

669 The authors thank Lena Forster and Carolin Hutter for technical assistance in the laboratory
670 and greenhouse. We further thank the gardener and technician team of the Plant Technology
671 Center at Technical University of Munich for maintenance of greenhouse experiments.

Author contributions

FH, CES, MH and RH: conceptualization.
FH, CES and RH wrote the manuscript
FH and conducted the greenhouse experiments.
CW performed the RNA sequencing.
MG and CD measured the phytohormones and stress markers.
MM and KM mapped the RNAseq reads to Barley genome.
CES, NK, RS and RH analysed the obtained sequencing results.
681

Conflict of interest

683 No conflict of interest declared

Funding

This project was financially supported by the Bavarian State Ministry of the Environment and Consumer Protection in frame of the Project network BayKlimaFit I and II (www.bayklimafit.de); subproject 10: TGC01GCUFuE69781 and subproject 6: TEW01002P-77746 to R.H. and by the German Research Foundation in the collaborative research center SFB924 to C.D..

References

Abramov, A; Hoffmann, T; Stark, T D *et al.* (2021): Engineering of benzoxazinoid biosynthesis in *Arabidopsis thaliana*: Metabolic and physiological challenges. In: *Phytochemistry* 192, p.112947. DOI: 10.1016/j.phytochem.2021.112947.

Adie, B A T; Godoy, M; Pérez-Pérez, J; Pérez-Pérez, M M; Sánchez-Serrano, J-J; Schmelz, E A; Solano, R (2007): ABA is an essential signal for plant resistance to pathogens affecting JA biosynthesis and the activation of defenses in *Arabidopsis*. In: *The Plant cell* 19 (5), p.1665–1681. DOI: 10.1105/tpc.106.048041.

Alexa, A; Rahnenfuehrer, J (2022): topGO. enrichment analysis for gene ontology. Version R package version 2.50.0.

Ameye, M; Audenaert, K; Haesaert, G; Smagghe, G; Steppe, K; van Meulebroek, L; Vanhaecke, L; Vleesschauwer, D de; Zutter, N de (2015): Priming of wheat with the green leaf volatile Z-3-hexenyl acetate enhances defense against *Fusarium graminearum* but boosts Deoxynivalenol production. In: *Plant physiology* 167 (4), p.1671–1684. DOI: 10.1104/pp.15.00107.

Atkinson, N J; Urwin, P E (2012): The interaction of plant biotic and abiotic stresses: from genes to the field. In: *Journal of experimental botany* 63 (10), p.3523–3543. DOI: 10.1093/jxb/ers100.

Babaeizad, V; Imani, J; Kogel, K-H; Eichmann, R; Hückelhoven, R (2009): Over-expression of the cell death regulator BAX inhibitor-1 in barley confers reduced or enhanced susceptibility to distinct fungal pathogens. In: *Theoretical and Applied Genetics* 118 (3), p.455–463. DOI: 10.1007/s00122-008-0912-2.

Bari, R; Jones, J D G (2009): Role of plant hormones in plant defence responses. In: *Plant molecular biology* 69 (4), p.473–488. DOI: 10.1007/s11103-008-9435-0.

Barnabás, B; Jäger, K; Fehér, A (2008): The effect of drought and heat stress on reproductive processes in cereals. In: *Plant, Cell & Environment* 31 (1), p.11–38. DOI: 10.1111/j.1365-3040.2007.01727.x.

Ben Alaya, A; Rabhi, F; Hessini, K; Djébali, N (2021): *Pyrenophora teres* growth and severity of net blotch on barley under salt stress. In: *European Journal of Plant Pathology* 161 (3), p.709–722. DOI: 10.1007/s10658-021-02355-z.

Birr, T; Hasler, M; Verreet, J-A; Klink, H (2020): Composition and Predominance of Fusarium Species Causing Fusarium Head Blight in Winter Wheat Grain Depending on Cultivar Susceptibility and Meteorological Factors. In: *Microorganisms* 8 (4), p.617. DOI: 10.3390/microorganisms8040617.

Biselli, C; Bagnaresi, P; Faccioli, P; Hu, X; Balcerzak, M; Mattera, M G; Yan, Z; Ouellet, T; Cattivelli, L; Valè, G (2018): Comparative Transcriptome Profiles of Near-Isogenic Hexaploid

Wheat Lines Differing for Effective Alleles at the 2DL FHB Resistance QTL. In: *Frontiers in plant science* 9, p.37. DOI: 10.3389/fpls.2018.00037.

Blum, A (2009): Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. In: *Field Crops Research* 112 (2-3), p.119–123. DOI: 10.1016/j.fcr.2009.03.009.

Boddu, J; Cho, S; Kruger, W M; Muehlbauer, G J (2006): Transcriptome Analysis of the Barley-*Fusarium graminearum* Interaction. In: *Molecular plant-microbe interactions* 19 (4), p.407–417. DOI: 10.1094/MPMI-19-0407.

Boddu, J; Cho, S; Muehlbauer, G J (2007): Transcriptome Analysis of Trichothecene-Induced Gene Expression in Barley. In: *Molecular plant-microbe interactions* 20 (7), p.1364–1375. DOI: 10.1094/MPMI-20-11-1364.

Bolger, A M; Lohse, M; Usadel, B (2014): Trimmomatic: a flexible trimmer for Illumina sequence data. In: *Bioinformatics* 30 (15), p.2114–2120. DOI: 10.1093/bioinformatics/btu170.

Brauer, E K; Rocheleau, H; Balcerzak, M; Pan, Y; Fauteux, F; Liu, Z; Wang, L; Zheng, W; Ouellet, T (2019): Transcriptional and hormonal profiling of *Fusarium graminearum*-infected wheat reveals an association between auxin and susceptibility. In: *Physiological and Molecular Plant Pathology* 107, p.33–39. DOI: 10.1016/j.pmpp.2019.04.006.

Brini, F; Hanin, M; Mezghani, I; Berkowitz, G A; Masmoudi, K (2007): Overexpression of wheat Na+/H+ antiporter TNHX1 and H+-pyrophosphatase TVP1 improve salt- and drought-stress tolerance in *Arabidopsis thaliana* plants. In: *Journal of experimental botany* 58 (2), p.301–308. DOI: 10.1093/jxb/erl251.

Brown, N A; Urban, M; van de Meene, A M L; Hammond-Kosack, K E (2010): The infection biology of *Fusarium graminearum*: defining the pathways of spikelet to spikelet colonisation in wheat ears. In: *Fungal biology* 114 (7), p.555–571. DOI: 10.1016/j.funbio.2010.04.006.

Buchanan, K L (2000): Stress and the evolution of condition-dependent signals. In: *Trends in Ecology & Evolution* 15 (4), p.156–160. DOI: 10.1016/s0169-5347(99)01812-1.

Buerstmayr, H; Lemmens, M (2015): Breeding healthy cereals: genetic improvement of Fusarium resistance and consequences for mycotoxins. In: *World Mycotoxin Journal* 8 (5), p.591–602. DOI: 10.3920/WMJ2015.1889.

Buerstmayr, M; Wagner, C; Nosenko, T; Omony, J; Steiner, B; Nussbaumer, T; Mayer, K F X; Buerstmayr, H (2021): Fusarium head blight resistance in European winter wheat: insights from genome-wide transcriptome analysis. In: *BMC Genomics* 22 (1), p.1–17. DOI: 10.1186/s12864-021-07800-1.

Buhrow, L M; Liu, Z; Cram, D; Sharma, T; Foroud, N A; Pan, Y; Loewen, M C (2021): Wheat transcriptome profiling reveals abscisic and gibberellic acid treatments regulate early-stage phytohormone defense signaling, cell wall fortification, and metabolic switches following *Fusarium graminearum*-challenge. In: *BMC Genomics* 22 (1), p.1–21. DOI: 10.1186/s12864-021-08069-0.

Chaudhary, A; Chen, X; Gao, J; Leśniewska, B; Hammerl, R; Dawid, C; Schneitz, K (2020): The Arabidopsis receptor kinase STRUBBELIG regulates the response to cellulose deficiency. In: *PLoS genetics* 16 (1), e1008433. DOI: 10.1371/journal.pgen.1008433.

Chen, X; Steed, A; Travella, S; Keller, B; Nicholson, P (2009): *Fusarium graminearum* exploits ethylene signalling to colonize dicotyledonous and monocotyledonous plants. In: *The New phytologist* 182 (4), p.975–983. DOI: 10.1111/j.1469-8137.2009.02821.x.

Choi, H-K; Iandolino, A; da Silva, F G; Cook, D R (2013): Water deficit modulates the response of Vitis vinifera to the Pierce's disease pathogen *Xylella fastidiosa*. In: *Molecular plant-microbe interactions* 26 (6), p.643–657. DOI: 10.1094/MPMI-09-12-0217-R.

Csardi, G; Nepusz, T (2006): The Igraph Software Package for Complex Network Research: InterJournal, complex systems, (Complex Systems).

Delgado-Baquerizo, M; Guerra, C A; Cano-Díaz, C; Egidi, E; Wang, J-T; Eisenhauer, N; Singh, B K; Maestre, F T (2020): The proportion of soil-borne pathogens increases with warming at the global scale. In: *Nature Climate Change* 10 (6), p.550–554. DOI: 10.1038/s41558-020-0759-3.

Ding, L; Xu, H; Yi, H; Yang, L; Kong, Z; Zhang, L; Xue, S; Jia, H; Ma, Z (2011): Resistance to Hemi-Biotrophic *F. graminearum* Infection Is Associated with Coordinated and Ordered Expression of Diverse Defense Signaling Pathways. In: *PLoS ONE* 6 (4), e19008. DOI: 10.1371/journal.pone.0019008.

Dolferus, R; Ji, X; Richards, R A (2011): Abiotic stress and control of grain number in cereals. In: *Plant Science* 181 (4), p.331–341. DOI: 10.1016/j.plantsci.2011.05.015.

Dörffling, K; Petersen, W (1984): ABA in phytopathogenic fungi of the genera *Botrytis*, *Ceratocystis*, *Fusarium* and *Rhizoctonia*. In: *Zeitschrift für Naturforschung* 39 (6), p.683–684.

Farooq, M; Wahid, A; Kobayashi, N; Fujita, D; Basra, S M A (2009): Plant Drought Stress: Effects, Mechanisms and Management. In: E. Lichtfouse, M. Navarrete, P. Debaeke, S. Véronique andC. Alberola (eds.): *Sustainable Agriculture*, p.153–188.

Fraaije, B A; Lovell, D J; Rohel, E A; Hollomon, D W (1999): Rapid detection and diagnosis of Septoria tritici epidemics in wheat using a polymerase chain reaction/PicoGreen assay. In: *Journal of Applied Microbiology* 86 (4), p.701–708. DOI: 10.1046/j.1365-2672.1999.00716.x.

Gardiner, S A; Boddu, J; Berthiller, F; Hametner., C; Stupar, R M; Adam, G; Muehlbauer, G J (2010): Transcriptome analysis of the Barley–Deoxynivalenol interaction: evidence for a role of glutathione in Deoxynivalenol detoxification. In: *Molecular plant-microbe interactions* 23 (7), p.962–976. DOI: 10.1094/MPMI-23-7-0962.

Gu, L; Zhang, Y; Zhang, M; Li, T; Dirk, L M A; Downie, B; Zhao, T (2016): ZmGOLS2, a target of transcription factor ZmDREB2A, offers similar protection against abiotic stress as ZmDREB2A. In: *Plant molecular biology* 90 (1), p.157–170. DOI: 10.1007/s11103-015-0403-1.

Gupta, A; Sarkar, A K; Senthil-Kumar, M (2016): Global Transcriptional Analysis Reveals Unique and Shared Responses in *Arabidopsis thaliana* Exposed to Combined Drought and Pathogen Stress. In: *Frontiers in plant science* 7, p.686. DOI: 10.3389/fpls.2016.00686.

Gupta, A; Sinha, R; Fernandes, J L; Abdelrahman, M; Burritt, D J; Tran, L-S P (2020): Phytohormones regulate convergent and divergent responses between individual and combined drought and pathogen infection. In: *Critical reviews in biotechnology* 40 (3), p.320–340. DOI: 10.1080/07388551.2019.1710459.

Hameed, A; Poznanski, P; Noman, M; Ahmed, T; Iqbal, A; Nadolska-Orczyk, A; Orczyk, W (2022): Barley Resistance to *Fusarium graminearum* Infections: From Transcriptomics to Field with Food Safety Concerns. In: *Journal of agricultural and food chemistry* 70 (46), p.14571–14587. DOI: 10.1021/acs.jafc.2c05488.

He, Y; Wu, L; Liu, X; Jiang, P; Yu, L; Qiu, J; Wang, G; Zhang, X; Ma, H (2020): TaUGT6, a Novel UDP-Glycosyltransferase Gene Enhances the Resistance to FHB and DON Accumulation in Wheat. In: *Frontiers in plant science* 11. DOI: 10.3389/fpls.2020.574775.

Hofer, K; Barmeier, G; Schmidhalter, U; Habler, K; Rychlik, M; Hückelhoven, R; Hess, M (2016): Effect of nitrogen fertilization on Fusarium head blight in spring barley. In: *Crop Protection* 88, p.18–27. DOI: 10.1016/j.cropro.2016.05.007.

Hoheneder, F; Biehl, E M; Hofer, K; Petermeier, J; Groth, J; Herz, M; Rychlik, M; Heß, M; Hückelhoven, R (2022): Host Genotype and Weather Effects on Fusarium Head Blight Severity and Mycotoxin Load in Spring Barley. In: *Toxins* 14 (2), p.125. DOI: 10.3390/toxins14020125.

Hoheneder, F; Hofer, K; Groth, J; Herz, M; Heß, M; Hückelhoven, R (2021): Ramularia leaf spot disease of barley is highly host genotype-dependent and suppressed by continuous drought stress in the field. In: *Journal of Plant Diseases and Protection* 128 (3), p.749–767. DOI: 10.1007/s41348-020-00420-z.

Howe, E A; Sinha, R; Schlauch, D; Quackenbush, J (2011): RNA-Seq analysis in MeV. In: *Bioinformatics* 27 (22), p.3209–3210. DOI: 10.1093/bioinformatics/btr490.

Huber, W; von Heydebreck, A; Sültmann, H; Poustka, A; Vingron, M (2002): Variance stabilization applied to microarray data calibration and to the quantification of differential expression. In: *Bioinformatics* 18 (Suppl. 1), S96-S104. DOI: 10.1093/bioinformatics/18.suppl_1.S96.

Jaroszuk-Ściseł, J; Kurek, E; Trytek, M (2014): Efficiency of indoleacetic acid, gibberellic acid and ethylene synthesized in vitro by *Fusarium culmorum* strains with different effects on cereal growth. In: *Biologia* 69 (3), p.281–292. DOI: 10.2478/s11756-013-0328-6.

Kaneda, T; Taga, Y; Takai, R; Iwano, M; Matsui, H; Takayama, S; Isogai, A; Che, F-S (2009): The transcription factor OsNAC4 is a key positive regulator of plant hypersensitive cell death. In: *The EMBO Journal* 28 (7), p.926–936. DOI: 10.1038/emboj.2009.39.

Kang, Y; Khan, S; Ma, X (2009): Climate change impacts on crop yield, crop water productivity and food security – A review. In: *Progress in Natural Science* 19 (12), p.1665–1674. DOI: 10.1016/j.pnsc.2009.08.001.

Karre, S; Kumar, A; Yogendra, K; Kage, U; Kushalappa, A; Charron, J-B (2019): HvWRKY23 regulates flavonoid glycoside and hydroxycinnamic acid amide biosynthetic genes in barley to combat Fusarium head blight. In: *Plant molecular biology* 100 (6), p.591–605. DOI: 10.1007/s11103-019-00882-2.

Kazan, K; Gardiner, D M (2018): Transcriptomics of cereal-*Fusarium graminearum* interactions: what we have learned so far. In: *Molecular plant pathology* 19 (3), p.764–778. DOI: 10.1111/mpp.12561.

Kim, D; Langmead, B; Salzberg, S L (2015): HISAT: a fast spliced aligner with low memory requirements. In: *Nature Methods* 12 (4), p.357–360. DOI: 10.1038/nmeth.3317.

Langfelder, P; Horvath, S (2008): WGCNA: an R package for weighted correlation network analysis. In: *BMC bioinformatics* 9, p.559. DOI: 10.1186/1471-2105-9-559.

Langfelder, P; Horvath, S (2012): Fast R Functions for Robust Correlations and Hierarchical Clustering. In: *Journal of statistical software* 46 (11).

Li, S; Chen, N; Li, F; Mei, F; Wang, Z; Cheng, X; Kang, Z; Mao, H (2020): Characterization of wheat homeodomain-leucine zipper family genes and functional analysis of TaHDZ5-6A in drought tolerance in transgenic *Arabidopsis*. In: *BMC plant biology* 20 (1), p.1–23. DOI: 10.1186/s12870-020-2252-6.

Liao, Y; Smyth, G K; Shi, W (2014): featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. In: *Bioinformatics* 30 (7), p.923–930. DOI: 10.1093/bioinformatics/btt656.

Linkmeyer, A; Götz, M; Hu, L; Asam, S; Rychlik, M; Hausladen, H; Hess, M; Hückelhoven, R (2013): Assessment and Introduction of Quantitative Resistance to Fusarium Head Blight in Elite Spring Barley. In: *Phytopathology* 103 (12), p.1252–1259. DOI: 10.1094/PHYTO-02-13-0056-R.

Loo, E P-I; Tajima, Y; Yamada, K *et al.* (2022): Recognition of Microbe- and Damage-Associated Molecular Patterns by Leucine-Rich Repeat Pattern Recognition Receptor Kinases Confers Salt Tolerance in Plants. In: *Molecular plant-microbe interactions* 35 (7), p.554–566. DOI: 10.1094/MPMI-07-21-0185-FI.

Low, Y C; Lawton, M A; Di, R (2020): Validation of barley 2OGO gene as a functional orthologue of Arabidopsis DMR6 gene in Fusarium head blight susceptibility. In: *Scientific Reports* 10 (1), p.1–13. DOI: 10.1038/s41598-020-67006-5.

Luo, K; Rocheleau, H; Qi, P-F; Zheng, Y-L; Zhao, H-Y; Ouellet, T (2016): Indole-3-acetic acid in *Fusarium graminearum*: Identification of biosynthetic pathways and characterization of physiological effects. In: *Fungal biology* 120 (9), p.1135–1145. DOI: 10.1016/j.funbio.2016.06.002.

Makandar, R; Nalam, V; Chaturvedi, R; Jeannotte, R; Sparks, A A; Shah, J (2010): Involvement of Salicylate and Jasmonate Signaling Pathways in *Arabidopsis* Interaction with *Fusarium graminearum*. In: *Molecular plant-microbe interactions* 23 (7), p.861–870. DOI: 10.1094/MPMI-23-7-0861.

Makandar, R; Nalam, V J; Lee, H; Trick, H N; Dong, Y; Shah, J (2012): Salicylic Acid Regulates Basal Resistance to Fusarium Head Blight in Wheat. In: *Molecular plant-microbe interactions* 25 (3), p.431–439. DOI: 10.1094/MPMI-09-11-0232.

Mandalà, G; Tundo, S; Francesconi, S; Gevi, F; Zolla, L; Ceoloni, C; D'Ovidio, R (2019): Deoxynivalenol Detoxification in Transgenic Wheat Confers Resistance to Fusarium Head Blight and Crown Rot Diseases. In: *Molecular plant-microbe interactions* 32 (5), p.583–592. DOI: 10.1094/MPMI-06-18-0155-R.

McGrann, G R D; Steed, A; Burt, C; Goddard, R; Lachaux, C; Bansal, A; Corbitt, M; Gorniak, K; Nicholson, P; Brown, J K M (2015): Contribution of the drought tolerance-related Stress-responsive NAC1 transcription factor to resistance of barley to Ramularia leaf spot. In: *Molecular plant pathology* 16 (2), p.201–209. DOI: 10.1111/mpp.12173.

Mesterhazy, A (2020): Updating the Breeding Philosophy of Wheat to Fusarium Head Blight (FHB): Resistance Components, QTL Identification, and Phenotyping—A Review. In: *Plants* 9 (12), p.1702. DOI: 10.3390/plants9121702.

Meza, I; Siebert, S; Döll, P *et al.* (2020): Global-scale drought risk assessment for agricultural systems. In: *Natural Hazards and Earth System Sciences* 20 (2), p.695–712. DOI: 10.5194/nhess-20-695-2020.

Michlmayr, H; Varga, E; Malachová, A *et al.* (2018): UDP-Glucosyltransferases from Rice, *Brachypodium*, and Barley: Substrate Specificities and Synthesis of Type A and B Trichothecene-3-O-β-d-glucosides. In: *Toxins* 10 (3), p.111. DOI: 10.3390/toxins10030111.

Monat, C; Padmarasu, S; Lux, T *et al.* (2019): TRITEX: chromosome-scale sequence assembly of Triticeae genomes with open-source tools. In: *Genome Biology* 20 (1), p.1–18. DOI: 10.1186/s13059-019-1899-5.

Mulako, I; Farrant, J M; Collett, H; Illing, N (2008): Expression of Xhdsi-1VOC, a novel member of the vicinal oxygen chelate (VOC) metalloenzyme superfamily, is up-regulated in leaves and roots during desiccation in the resurrection plant *Xerophyta humilis* (Bak) Dur and Schinz. In: *Journal of experimental botany* 59 (14), p.3885–3901. DOI: 10.1093/jxb/ern226.

Nicolaisen, M; Supronienė, S; Nielsen, L K; Lazzaro, I; Spliid, N H; Justesen, A F (2009): Real-time PCR for quantification of eleven individual *Fusarium* species in cereals. In: *Journal of Microbiological Methods* 76 (3), p.234–240. DOI: 10.1016/j.mimet.2008.10.016.

Ogrodowicz, P; Kuczyńska, A; Mikołajczak, K; Adamski, T; Surma, M; Krajewski, P; Ćwiek-Kupczyńska, H; Kempa, M; Rokicki, M; Jasińska, D (2020): Mapping of quantitative trait loci for traits linked to fusarium head blight in barley. In: *PLoS ONE* 15 (2), e0222375. DOI: 10.1371/journal.pone.0222375.

Olesen, J E; Trnka, M; Kersebaum, K C; Skjelvåg, A O; Seguin, B; Peltonen-Sainio, P; Rossi, F; Kozyra, J; Micale, F (2011): Impacts and adaptation of European crop production systems to climate change. In: *European Journal of Agronomy* 34 (2), p.96–112. DOI: 10.1016/j.eja.2010.11.003.

Pandey, P; Ramegowda, V; Senthil-Kumar, M (2015): Shared and unique responses of plants to multiple individual stresses and stress combinations: physiological and molecular mechanisms. In: *Frontiers in plant science* 6, p.723. DOI: 10.3389/fpls.2015.00723.

Pandey, P; Senthil-Kumar, M (2019): Plant-pathogen interaction in the presence of abiotic stress: What do we know about plant responses? In: *Plant Physiology Reports* 24 (4), p.541–549. DOI: 10.1007/s40502-019-00483-7.

Perochon, A; Kahla, A; Vranić, M; Jia, J; Malla, K B; Craze, M; Wallington, E; Doohan, F M (2019): A wheat NAC interacts with an orphan protein and enhances resistance to Fusarium head blight disease. In: *Plant biotechnology journal* 17 (10), p.1892–1904. DOI: 10.1111/pbi.13105.

Qi, P-F; Balcerzak, M; Rocheleau, H; Leung, W; Wei, Y-M; Zheng, Y-L; Ouellet, T (2016): Jasmonic acid and abscisic acid play important roles in host–pathogen interaction between *Fusarium graminearum* and wheat during the early stages of fusarium head blight. In: *Physiological and Molecular Plant Pathology* 93, p.39–48. DOI: 10.1016/j.pmpp.2015.12.004.

Ramegowda, V; Senthil-Kumar, M (2015): The interactive effects of simultaneous biotic and abiotic stresses on plants: Mechanistic understanding from drought and pathogen combination. In: *Journal of Plant Physiology* 176, p.47–54. DOI: 10.1016/j.jplph.2014.11.008.

Rivero, R M; Mittler, R; Blumwald, E; Zandalinas, S I (2022): Developing climate-resilient crops: improving plant tolerance to stress combination. In: *The Plant Journal* 109 (2), p.373–389. DOI: 10.1111/tpj.15483.

Robinson, M D; McCarthy, D J; Smyth, G K (2010): edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. In: *Bioinformatics (Oxford, England)* 26 (1), p.139–140. DOI: 10.1093/bioinformatics/btp616.

Schweiger, W; Boddu, J; Shin, S; Poppenberger, B; Berthiller, F; Lemmens, M; Muehlbauer, G J; Adam, G (2010): Validation of a Candidate Deoxynivalenol-Inactivating UDP-Glucosyltransferase from Barley by Heterologous Expression in Yeast. In: *Molecular plant-microbe interactions* 23 (7), p.977–986. DOI: 10.1094/MPMI-23-7-0977.

Selvaraj, M G; Ishizaki, T; Valencia, M *et al.* (2017): Overexpression of an Arabidopsis thaliana galactinol synthase gene improves drought tolerance in transgenic rice and increased grain yield in the field. In: *Plant biotechnology journal* 15 (11), p.1465–1477. DOI: 10.1111/pbi.12731.

Söderman, E; Mattsson, J; Engström, P (1996): The *Arabidopsis* homeobox gene ATHB-7 is induced by water deficit and by abscisic acid. In: *The Plant Journal* 10 (2), p.375–381. DOI: 10.1046/j.1365-313X.1996.10020375.x.

Soni, N; Hegde, N; Dhariwal, A; Kushalappa, A C (2020): Role of laccase gene in wheat NILs differing at QTL-Fhb1 for resistance against Fusarium head blight. In: *Plant Science* 298, p.110574. DOI: 10.1016/j.plantsci.2020.110574.

Su, Z Y; Powell, J J; Gao, S; Zhou, M; Liu, C (2021): Comparing transcriptional responses to Fusarium crown rot in wheat and barley identified an important relationship between disease resistance and drought tolerance. In: *BMC plant biology* 21 (1), p.73. DOI: 10.1186/s12870-020-02818-1.

Sun, Y; Xiao, J; Jia, X; Ke, P; He, L; Cao, A; Wang, H; Wu, Y; Gao, X; Wang, X (2016): The role of wheat jasmonic acid and ethylene pathways in response to Fusarium graminearum infection. In: *Plant Growth Regulation* 80 (1), p.69–77. DOI: 10.1007/s10725-016-0147-1.

Suzuki, N; Rivero, R M; Shulaev, V; Blumwald, E; Mittler, R (2014): Abiotic and biotic stress combinations. In: *New Phytologist* 203 (1), p.32–43. DOI: 10.1111/nph.12797.

Trujillo, M; Ichimura, K; Casais, C; Shirasu, K (2008): Negative Regulation of PAMP-Triggered Immunity by an E3 Ubiquitin Ligase Triplet in *Arabidopsis*. In: *Current Biology* 18 (18), p.1396–1401. DOI: 10.1016/j.cub.2008.07.085.

Tucker, J R; Legge, W G; Maiti, S; Hiebert, C W; Simsek, S; Yao, Z; Xu, W; Badea, A; Fernando, W G D (2021): Transcriptome Alterations of an in vitro-Selected, Moderately Resistant, Two-Row Malting Barley in Response to 3ADON, 15ADON, and NIV Chemotypes of *Fusarium graminearum*. In: *Frontiers in plant science* 12, p.1567. DOI: 10.3389/fpls.2021.701969.

Wang, L; Li, Q; Liu, Z; Surendra, A; Pan, Y; Li, Y; Zaharia, I L; Ouellet, T; Fobert, P R (2018): Integrated transcriptome and hormone profiling highlight the role of multiple phytohormone pathways in wheat resistance against fusarium head blight. In: *PLoS ONE* 13 (11), e0207036. DOI: 10.1371/journal.pone.0207036.

Wegulo, S N; Baenziger, P S; Hernandez Nopsa, J; Bockus, W W; Hallen-Adams, H (2015): Management of Fusarium head blight of wheat and barley. In: *Crop Protection* 73, p.100–107. DOI: 10.1016/j.cropro.2015.02.025.

Xiao, J; Jin, X; Jia, X *et al.* (2013): Transcriptome-based discovery of pathways and genes related to resistance against Fusarium head blight in wheat landrace Wangshuibai. In: *BMC Genomics* 14 (1), p.1–19. DOI: 10.1186/1471-2164-14-197.

Xie, W; Xiong, W; Pan, J; Ali, T; Cui, Q; Guan, D; Meng, J; Mueller, N D; Lin, E; Davis, S J (2018): Decreases in global beer supply due to extreme drought and heat. In: *Nature Plants* 4 (11), p.964–973. DOI: 10.1038/s41477-018-0263-1.

Yamanouchi, U; Yano, M; Lin, H; Ashikari, M; Yamada, K (2002): A rice spotted leaf gene, Spl7, encodes a heat stress transcription factor protein. In: *Proceedings of the National Academy of Sciences* 99 (11), p.7530–7535. DOI: 10.1073/pnas.112209199.

Yao, T; Zhang, J; Xie, M; Yuan, G; Tschaplinski, T J; Muchero, W; Chen, J-G (2021): Transcriptional Regulation of Drought Response in *Arabidopsis* and Woody Plants. In: *Frontiers in plant science* 11, p.2044. DOI: 10.3389/fpls.2020.572137.

Zandalinas, S I; Sengupta, S; Fritschi, F B; Azad, R K; Nechushtai, R; Mittler, R (2021): The impact of multifactorial stress combination on plant growth and survival. In: *The New phytologist* 230 (3), p.1034–1048. DOI: 10.1111/nph.17232.

Zhang, B; Horvath, S (2005): A general framework for weighted gene co-expression network analysis. In: *Statistical applications in genetics and molecular biology* 4. DOI: 10.2202/1544-6115.1128.

Zhang, H; Sonnewald, U (2017): Differences and commonalities of plant responses to single and combined stresses. In: *The Plant Journal* 90 (5), p.839–855. DOI: 10.1111/tpj.13557.

Figure legends

Fig. 1. Global transcription analysis of variety-dependent, reduced susceptibility against FHB under drought stress in the greenhouse. (A) Experimental setup: Three different barley varieties (Barke, Ba; Morex, Mo; and Palmella Blue, Pb) were grown in pots under controlled conditions in the greenhouse. Drought stress was induced by stopped irrigation 7 days before anthesis (growth stage 61-69; dark grey arrow). Irrigated and drought-stressed plants were inoculated with conidia of *Fusarium culmorum* or mock (dark grey arrow) and samples were harvested 2 and 4 days post-inoculation (light grey arrows). (B) Number of DEGs. For each variety all treatments were compared against the watered, mock-infected samples. Differentially expressed genes were counted (FDR $p<0.05$) and number of DEGs were plotted. Upwards directed columns show the numbers of upregulated genes and downwards directed columns show downregulated genes. (C) Heat maps of key experimental outcomes and expression $\log_2$ fold changes of clusters of genes that behave similarly according to self-organizing tree algorithm. Amount of *Fusarium culmorum* DNA per ng barley DNA was determined by qPCR. Values were colour-coded with grey (0), yellow ($\log_2$ 1 pg *Fc* DNA ng^{-1} barley^{-1} DNA^{-1}) to orange ($\log_2$ 6 pg *Fc* DNA ng^{-1} barley^{-1} DNA^{-1}). The average DNA content from 2 and 4 dpi was calculated and the $\log_2$ used for data representation. Abscisic acid content [in pg ng^{-1} fresh^{-1} weight^{-1}] in all spike samples was quantified using mass spectrometry. Values were colour-coded with light-green (0 ng ABA ng^{-1} fresh^{-1} weight^{-1}) to dark-green (4000 ng ABA ng^{-1} fresh^{-1} weight^{-1}). For each variety DEGs (relative to watered, not infected samples) with a FDR-corrected p-value < 0.05 were counted under each stress treatment and the quantity was colour-coded with light-blue (0) to dark-blue (7500). All genes which were significantly regulated ($p < 0.05$) in at least one variety in one particular stress treatment (infection: 2949 genes; drought: 7879 genes; drought + infection: 10188 genes) were subjected to a cluster analysis using a self-organizing tree algorithm (SOTA) using the expression pattern over all varieties and stress treatments. The respective stress treatment for which the DEGs were selected is highlighted by a black frame. All DEGs were split into 11 clusters, labelled with numbers 1-11 and a capital letter for the stress (I=infection, D=drought, DI=drought + infection) next to it and the corresponding number of genes in each cluster. The mean $\log_2$ fold change for each cluster was colour-coded ranging from blue (> -4) over white (0) to red (< 4).

Fig. 2. Modules of DEGs after weighted gene co-expression network analysis. (A) Co-expression modules. Based on the expression pattern over all time points, varieties and

treatments all DEGs clustered into 18 modules (color-coded) with 63 to 5287 genes after WGCNA-analysis. 42 genes didn't follow any co-expression pattern and were clustered in the grey module. The numbers reflect the amount of DEGs in each module. (B) Top 5 biggest co-expression modules in each variety for each stress treatment. We counted the DEGs in each module for each variety under each stress treatment. We show the 5 modules with most DEGs in each stress scenario in percentage relative to the total number of DEGs per variety (Barke, Ba; Morex, Mo; and Palmella Blue, Pb).

Fig. 3. Relationships of consensus module eigengenes with FHB-severity, different phytohormones or stress markers. Each column in the table corresponds to a module and each row to one of the physiological traits: FHB-severity, a phytohormone, its derivate or proline as a stress marker. Module names are shown on top. Square colours in the figure represent the correlations of corresponding module eigengenes and measured stress parameters. The FDR-corrected p-values are coded by size (the bigger the square, the more significant). Highly significant correlations ($p < 0.01$) are highlighted with bold frames. DNA, *F. culmorum* DNA relative to plant DNA; ABA, abscisic acid; PA, phaseic acid; DPA, dihydrophaseic acid; IAA, indol 3-acetic acid; SA, salicylic acid.

Fig. 4. Co-expression levels reveal potential hub genes in selected modules. For three co-expression modules (blue, salmon and green), the 25 edges with the highest edge weight have been selected and their corresponding network was schematically drawn (A blue module, C salmon module, E green module). Each node represents one gene, and in the heat maps on the left the log fold changes of the gene expression values are shown with the respective colour codes and individual scales for each module (B blue module, D salmon module, F green module). The size of the square represents the false discovery rate-corrected p-values (FDR), with bigger squares indicating higher confidence levels. The bold squares indicate a significant FDR p-value smaller than 0.05. For gene identifiers, see supplemental dataset D5.

Fig. 1. Global transcription analysis of variety-dependent, reduced susceptibility against FHB under drought stress in the greenhouse.

Fig. 2. Modules of DEGs after weighted gene co-expression network analysis.

Fig. 3. Relationships of consensus module eigengenes with FHB-severity, different phytohormones or stress markers.

Fig. 4. Co-expression levels reveal potential hub genes in selected modules.