1

Impaired inflammation resolution in murine bone marrow failure is rescued by Resolvin E1 treatment

Rachel Grazda[1]*, Allison N. Seyfried[1,4]*, Krishna Rao Maddipatti[2], Gabrielle Fredman[3], Katherine C. MacNamara[1#]

*co-first authors

[1]Department of Immunology and Microbiology, Albany Medical College, Albany, New York, USA;

[2]Department of Pathology, Lipidomics Core Facility, Wayne State University, Detroit, Michigan, USA;

[3]Department of Molecular and Cellular Physiology, Albany Medical College, Albany, New York, USA;

[4]Current address: Institute for Clinical Pharmacodynamics, Schenectady, NY, USA

[#] Corresponding author:
Katherine C. MacNamara
Department of Immunology and Microbial Disease
Albany Medical College, MC-151
47 New Scotland Avenue
Albany, NY 12208
Phone: 518-262-0921
Email: macnamk@mail.amc.edu

33 Running Title: **Resolvin E1 improves SAA**

Key Points:

- Deficiencies in efferocytosis during SAA are associated with expansion of MerTK^{hi} SIRPα^{hi} monocytes and increased CD47 expression.

- Imbalanced pro-inflammatory/pro-resolving lipid mediators correlated with SAA progression; RvE1 treatment improves efferocytosis and disease.

Abstract

Severe aplastic anemia (SAA) is an acquired form of bone marrow failure characterized by a profound loss of hematopoietic stem cells and failure to produce blood and immune cells. T cells are well-known drivers of disease, though underlying mechanisms causing persistent inflammation remain unknown. Typically, inflammation resolves via an active process mediated by specialized pro-resolving mediators (SPMs), such as resolvins. SPMs dampen inflammation without compromising host immunity, in part, through their ability to induce efficient clearance of apoptotic cells by phagocytes (i.e., efferocytosis). In a murine model of SAA, we identify a novel population of marrow monocytes that co-express the efferocytic receptor MerTK and the inhibitory receptor SIRPα. At the same time, the marrow contained a significant increase in apoptotic and dead cells that expressed the 'don't eat me' marker CD47. Myeloid cells from the marrow of SAA-induced mice exhibited reduced phagocytosis and efferocytosis, relative to controls. Consistent with impaired resolution responses, lipidomic analysis of the marrow revealed an imbalance of pro-inflammatory to pro-resolving lipid mediators. Specifically, we observed significantly reduced 18-hydroxyeicosapentaenoic acid (18-HEPE), a precursor to the E-series resolvins, followed by markedly increased prostaglandins and thromboxane in SAA. Treatment with exogenous Resolvin E1 (RvE1) increased BM cellularity, platelet count, and survival in SAA-induced mice, correlating with improved efferocytotic function. Our findings suggest that defective resolution pathways contribute to disease progression in SAA and demonstrate that SPMs, such as RvE1, may offer new strategies to treat disease that do not rely on immune suppression.

Word count: 244

Introduction

Idiopathic severe aplastic anemia (SAA) is a rare form of bone marrow failure (BMF) in which T cell-derived IFNγ and macrophage-derived tumor necrosis factor (TNF) drive the loss of hematopoietic stem and progenitor cells (HSPCs), resulting in pancytopenia.[1,2] Patients diagnosed with SAA have significantly increased interferon gamma (IFNγ) and TNF levels in the blood and BM.[3,4] Current therapies for SAA remain limited to immunosuppressive therapy (IST; anti-thymocyte globulin and cyclosporine) and BM transplantation.[5-7] Although the mechanism was unknown in 1981, treatment of patients with anti-thymocyte globulin in conjunction with BM transplantation increased survival odds immensely.[5]

Using IFNγ adenylate-uridylate–rich element (ARE)–deleted (del) mice, which exhibit increased IFNγ levels, Lin et al demonstrated that autoreactive T cells did not directly mediate destruction of HSCs during SAA, rather, IFNγ acted on the HSPC compartment to inhibit progenitor cell maturation and differentiation.[1] It has also been shown in a murine model of SAA induced by lymphocyte transfer, that TNF produced by host macrophages drives IFNγ production and infiltration of T cells into the BM during SAA.[2] Additionally, macrophages were shown to drive disease in an IFNγR- and CCR5-dependent manner.[8,9] Therefore, although T cells are an important target of current IST, additional cell types likely contribute to disease pathogenesis.

In 1995, researchers noted that patients with aplastic anemia exhibited an increased frequency of apoptotic CD34+ progenitor cells in the BM, and the frequency of apoptotic cells correlated with severity of disease.[10] These data suggested that increased apoptosis of progenitors contributed to disease progression. In addition,

rampant cell death, driven by the Fas/FasL pathway for apoptosis, leads to the

destruction of the hematopoietic stem and progenitor pool[11], as well as other cells[12].

Macrophages are critical for SAA progression, in part through producing and

augmenting inflammatory mediators[2,8,9]. Macrophages are also important phagocytes

within the BM, among other tissues, where they remove apoptotic cells[13,14], and

promote tissue regeneration[15].

Efficient clearance of apoptotic and dead cells by phagocytes, termed

efferocytosis, is crucial for inflammation-resolution and tissue function at homeostasis.[16]

Apoptotic cells provide both "find me" and "eat me" signals to phagocytes[17], and

phosphatidylserine (PS) is a well-known "eat me" signal recognized by tissue-resident

phagocytes and efferocytic monocyte-derived macrophages.[18,19] Apoptotic signaling

promotes rearrangement of the phospholipid bilayer to expose PS on the outer leaflet.

Externalized PS is either recognized directly, or via the linker molecules Gas6 or Protein

S, by TAM receptors on phagocytes such as MerTK. Signaling through MerTK induces

cytoskeleton rearrangements that initiates engulfment of the apoptotic cell.[17,20]

Efferocytic function is regulated, however, to prevent clearance of healthy cells. Indeed,

healthy, viable cells express "don't eat me" signals, such as CD47, that interact with

inhibitory receptors expressed by phagocytes, such as SIRPα, resulting in the inhibition

of phagocytosis.[21,22]

Resolution of inflammation is an active process mediated by a class of lipid

mediators termed specialized pro-resolving mediators (SPMs).[23] SPMs, such as the D-

and E-series resolvins, derived from omega-3 polyunsaturated fatty acids, possess

potent pro-resolving actions and are essential for resolution.[23] Resolvin E1 (RvE1) has

been shown to improve efferocytosis in various inflammatory disease models such as

periodontitis[24], colitis[25], lung cancer[26], acute lung injury and bacterial pneumonia[27],

asthma[28], and atherosclerosis[29], to name a few. Using an established, preclinical

murine model of SAA, we demonstrate dysfunctional inflammation resolution programs,

as determined by impaired efferocytosis and an imbalance in pro-inflammatory and pro-

resolving lipid mediators. We identify a unique population of MerTK^{hi} SIRPα^{hi}

monocytes, and an accumulation of CD47^{+} apoptotic cells within the BM during SAA,

that correlate with impaired cell clearance. RvE1 treatment, given to mice with

established SAA, provided significant protection and improved efferocytosis. Our

findings provide new insight to inflammation resolution programs and SAA

pathogenesis. Moreover, our work offers key rationale to pursue novel therapeutic

strategies to treat BMF that improve inflammation resolution without limiting host

defense.

Methods

Mice: All animal protocols were approved by the Institutional Animals Care and Use Committee at Albany Medical College. C57BL/6 (H$^{b/b}$) and BALB/cAnN (H-2$^{d/d}$) mice were purchased from Taconic (Albany, NY, https://www.taconic.com/). Macrophage insensitive to IFNγ (MIIG) mice were generously gifted by Dr. Michael Jordan. Hybrid B6 F1 (H-2$^{b/d}$) mice were generated by crossing C57BL/6 or MIIG mice on a C57BL/6 background with BALB/c mice. All mice were bred and housed in the Animal Resource Facility under microisolator conditions at Albany Medical College.

Bone Marrow Failure Induction: A splenocyte-infusion method was used to induce bone marrow failure (BMF) in Hybrid F1 mice aged 6-9 weeks. Hybrid F1 mice were subjected to sub-lethal radiation (300 RADs) using a ^{137}Cs source. Irradiated mice received adoptive transfer of 6.5 x 10^6 C57BL/6 splenocytes from age- and gender-matched donors via intraperitoneal injection. Mice were euthanized via CO_2 inhalation followed by cervical dislocation.

Blood Collection and CBCs: Blood was collected from euthanized mice into EDTA-coated tubes via cardiac puncture and analyzed with a Heska Element HT5 for complete blood count (CBC).

Cell Preparation and Flow Cytometry: Whole bone marrow was flushed from femurs and tibias. After RBC lysis, cell suspensions were plated and stained. See supplemental

157   Table 1 for antibody details. Data were collected using a FACSymphony A3 (BD

158   Biosciences) with FACSDiva software or Cytek Northern Light (Cytek Bioscienes) with

159   SpectroFlo software and analyzed using FlowJo software (TreeStar, Ashland, OR).

160
161
Phagocytosis Assay: 200μL fluorescent Dil (DilC$_{18}$(3))-labeled liposomes (Liposoma)

were administered to mice via retro-orbital I.V. injection on day 9 post induction. BM and

blood was harvested 16 hours post injection.

Efferocytosis Assay: Whole BM was flushed and incubated with phosphatidylserine

(PS)-coated lipid microparticles (Echelon Biosciences) for 1 hour at 37°C. Cells were

then stained for flow cytometry.

SIRPα Neutralization: Anti-SIRPα (clone P84; 200 μg; BioXCell) antibody was diluted

in PBS and administered to mice via intraperitoneal injection on days 7 and 9 post-

induction for day 10 harvest, or days 7, 9, 11, and 13 for day 14 harvest.

173
174
RvE1 Treatment: Resolvin E1 (250ng; Cayman Chemical) was diluted in PBS and

administered to mice via intraperitoneal injection on days 7, 9, 11 for day 12 harvest, or

days 7, 9, 11, and 13 for day 14 harvest.

Gene Expression: Whole BM was flushed and pooled from hind limbs of three mice

per group. RBCs were lysed and cells were stained to sort purified monocytes (SSC^{low},

CD11b$^+$, Ly6C^{hi}, Ly6G$^-$). Monocytes were collected in RLT lysis buffer (Qiagen). mRNA

was isolated (Qiagen RNeasy Mini Kit) and quantitative-RT-PCR was performed

(Eppendorf *realplex[2]* Mastercycler).

Protein analysis: Whole BM was flushed from the pelvis. BM cell lysates were

homogenized manually with a pestle in a buffer containing IGEPAL CA-630 and

protease inhibitors. Soluble Mer and Gas6 were quantified using ELISA kits from Ray

Biotech (accession #Q60805 and Q61592, respectively) and normalized to whole

protein using the Pierce BCA protein assay kit (Thermo Fisher, catalog #23255).

Lipidomic analysis: Whole BM was flushed from mouse hind limbs (femurs and tibias)

with ice cold PBS. Samples were flash frozen on dry ice before shipment to Wayne

State University Lipidomics Core facility for analysis. Fatty acyl lipidomic analysis was

performed as per published procedures.[30,31] Briefly, the frozen samples were thawed on

ice at the time of analysis, and homogenized using Zirconium beads (Precellys,

Biotage). The homogenate was extracted for fatty acyl lipids using StrataX columns

(Phenomenex) following supplementation with internal standards. The extracts were

analyzed by LC-MS/MS using Multiple Reaction Monitoring (MRM) method on a

QTRAP5500 mass analyzer (Sciex). Identities of the individual lipid mediators were

confirmed from the retention times and spectra recorded for each detected peak and

were quantified relative to the internal standards. The data were normalized against

protein content of the samples (ng/mg protein).

Survival Studies: PBS or 250 ng RvE1 was administered via intraperitoneal injection days 7, 9, 11, and 13 post induction. Mice were monitored twice daily for 28 or 37 days and euthanized when moribund. Euthanasia criteria was based on signs of dehydration, response to physical stimuli, and mobility — as previously described.[9]

Statistical Analysis: Data were analyzed with GraphPad Prism software (version 8.0, La Jolla, CA). Statistical analysis was performed using two-tailed Student's T-test or Two-Way ANOVA with Tukey's post-hoc analysis, and details are provided in each figure legend. Lipidomic data was analyzed using GraphPad Prism software (version 9.0, La Jolla, CA). Principal component analysis was performed using 22 lipid mediators detected across all samples from mice 3 days post-radiation for healthy (no radiation), radiation controls, and SAA-induced mice. Data were standardized and principal components were selected using Kaiser-Guttman's rule. Heat map analysis was performed on LC-MS/MS data from radiation controls and SAA-induced mice, normalized to healthy controls.

All methods for flow cytometric analysis and lipidomic analysis are described in detail in the supplemental materials (available on the Blood website), and data will be made available upon request.

Results

Monocytosis, cytopenias, and increased marrow cell death during SAA

Thrombocytopenia and decreased BM cellularity are well-known characteristics of severe aplastic anemia (SAA).[1,8,9] Using a murine model of SAA induced by adoptive transfer of splenocytes to sub-lethally irradiated F1 mice[8,32,33], we observed a significant decrease in red blood cells (**Figure 1A**) and platelets (**Figure 1B**) by 10 days post splenocyte transfer (dpst). We also observed a dramatic loss of blood lymphocytes correlating with an expansion of the circulating blood monocyte population (**Figure 1C**). Mice induced to develop SAA exhibited significant hypocellularity by 10 dpst (**Figure 1D**), though BM monocytes (CD11b$^+$Ly6C^{hi}Ly6G$^-$) were increased by both frequency and number, whereas neutrophils (CD11b$^+$ Ly6C^{low} Ly6G^{hi}) were decreased (**Figure 1E-F;** supplemental Fig. 1A-B). Therefore, disease progression is associated with a hematopoietic program that favors monopoiesis.

To investigate cell death during disease progression we performed Annexin V and 7-AAD staining (**Figure 1G**). Similar frequencies of apoptotic (AnnV$^+$7-AAD$^-$) and dead cells were observed between radiation controls and SAA-induced mice at 8 dpst (**Figure 1H**). However, by 10 dpst the SAA-induced mice had a striking increase in dead (7-AAD$^+$) cells, compared to radiation controls. IFNγ can induce apoptosis[34] and is a known driver of disease. We previously demonstrated that transgenic mice containing macrophages insensitive to IFNγ (referred to as MIIG mice[8]) exhibit protection from SAA. Here we observed that SAA-induced MIIG mice exhibited a decrease in the frequency of dead cells, relative to littermate control (LC) counterparts (supplemental

Fig. 2). These data suggest that the accumulation of dead cells and increased

monocytes, during the early stages of SAA, correlate with disease progression.

Increased surface expression of MerTK and SIRPα on monocytes during SAA

Under healthy conditions, apoptotic cells are rapidly cleared, and the accumulation

of dead cells during SAA suggested that efferocytosis may be impaired. Monocytes and

macrophages (MΦs) are the main efferocytes within the BM[35], and MerTK is a critical

receptor for efferocytosis that, when bound to PS on apoptotic cells, initiates

cytoskeletal rearrangement and the subsequent uptake of dead or dying cells.[36] We

hypothesized that reduced MerTK expression or cleavage via ADAM17, may contribute

to impaired clearance of dead cells in the marrow during SAA. To address this, we

performed flow cytometry to evaluate MerTK expression on myeloid cells 10 days post

SAA induction. In contrast to our prediction, the frequency and absolute number of

MerTK expressing monocytes expanded dramatically in the BM of SAA mice (**Figure**

2A-C). MerTK expression on macrophages also increased during SAA, though these

changes were not statistically significant (supplemental Fig. 3A-C). Analysis of

monocyte gene expression indicated an increase in *Mertk*, as well as *Ifnγ* and

Tnf (known drivers of disease)[1,2], in SAA mice, relative to radiation controls

(supplemental Fig. 4 A-C).

Under inflammatory conditions, MerTK can be cleaved by ADAM17[37], releasing

soluble Mer (sMer) into the extracellular milieu to act as a decoy for Gas6 and Protein

S.[38] Under SAA conditions, we observed increased Mer in BM homogenates (BM-H),

though this did not reach statistical significance (supplemental Fig. 4D). To complement

this analysis, we also measured Gas6 in the marrow and found a significant decrease in

SAA mice, relative to radiation control (supplemental Fig. 4E). These data suggest that,

during SAA, increased Mer cleavage without sufficient production and secretion of Gas6

may contribute to impaired efferocytosis, and the subsequent accumulation of dead and

dying cells.

Healthy cells can actively suppress their engulfment via "don't eat-me" signaling,

such as the CD47-SIRPα axis. We found that the population of monocytes with high

MerTK expression during SAA also exhibited a significant increase in

SIRPα expression, resulting in a striking increase in the absolute number of MerTK^{hi}

SIRPa^{hi} monocytes **(Figure 2D-F**). We also observed an increase in SIRPα on MerTK^{hi}

macrophages (supplemental Fig. 3D-F). Together, these data suggest that, under SAA

conditions, monocytes are primed for efferocytosis but expression of SIRPα may impair

this function. Monocytes from MIIG mice expressed significantly reduced MerTK and

SIRPα, correlating with resistance of MIIG mice to SAA progression (supplemental Fig.

5). Taken together, our data are consistent with a role of aberrantly high MerTK

expression and the unique population of MerTK^{hi} SIRPα^{hi} monocytes in disease

pathogenesis.

Increased CD47 expression on dead cells

CD47 is an important marker of "self" for healthy viable cells, and apoptotic cells

lose expression of CD47 enabling phagocytic clearance.[39,40] During SAA, however, we

observed increased CD47 expression on both live and apoptotic BM cells (**Figure 2G-I**).

These data suggest that increased CD47 on apoptotic cells, combined with increased

SIRPα^{hi} monocytes, may prevent efficient clearance in the marrow during SAA.

To better understand whether the SIRPα–CD47 axis contributes to disease, SAA-

induced mice were treated with anti-SIRPα neutralizing antibody days 7, 9, and 11 post

induction (supplemental Fig. 6A). Although we did observe a significant decrease in the

frequency of dead cells and CD47^{+} apoptotic cells (supplemental Fig. 6C-D), anti-SIRPα

treatment exacerbated anemia during SAA by 14 dpst (supplemental Fig. 6F-I).

Together, these data support the idea that the SIRPα–CD47 axis contributes to the

accumulation of dead cells during SAA, but as this axis is necessary for RBC

circulation[41], we reasoned it would not be an effective therapeutic target in SAA.

Decreased phagocytosis and efferocytosis in SAA

To examine whether phagocytes were functionally impaired during SAA, F1 mice

were induced to develop disease and administered fluorescently-labeled (Dil-labeled)

liposomes day 9 post-induction to evaluate phagocytotic capacity. BM was harvested

day 10 to evaluate the uptake of liposomes via flow cytometry (supplemental Fig. 7A).

SAA mice exhibited a decrease in Dil liposome-positive F4/80^{+} and CD11b^{+} cells,

suggesting impairments in phagocytosis (supplemental Fig. 7B-E).

To better address whether efferocytosis is defective during SAA, we utilized

phosphatidylserine (PS)-coated microparticles to mimic apoptotic cells. BM from SAA-

induced mice and radiation controls was incubated with fluorescently tagged PS-coated

lipid microparticles (PS-MPs), and uptake was determined via flow cytometry (**Figure**

3A-B). We observed a significant reduction in PS-MP^{+} monocytes and F4/80^{+} cells in

marrow isolated from SAA-induced mice, relative to controls, at 8 dpst (**Figure 3C-F**).

Consistent with the role of impaired efferocytosis in disease progression, SAA-induced

MIIG mice did not exhibit a significant decrease in PS-MPs, as compared to LC

counterparts (supplemental Fig. 8). Together, these data demonstrate that during SAA

progression, myeloid cells exhibit reduced phagocytosis and efferocytosis.

Imbalance of lipid mediators that regulate inflammation resolution in SAA

Based on our observation that efferocytosis is impaired during SAA progression,

we next addressed how key resolution programs may be impacted in disease. To

examine the lipid mediators associated with disease initiation and progression we

performed LC-MS/MS to analyze the polyunsaturated fatty acid (PUFA) metabolome in

BM of healthy, radiation control, and SAA-induced mice. Principal component analysis

(PCA) revealed that a majority of the variation (PC1; 43%) at day 3 post-radiation was

due to differences in mono-hydroxy eicosapentaenoic acids, including a decrease in 18-

HEPE in SAA mice, relative to radiation controls (**Figure 4A,** supplemental Fig. 9A).

Indeed, limited 18-HEPE, the precursor to the E-series resolvins, may contribute to

impaired generation of these SPMs in the marrow during SAA (**Figure 4B**). By day 8,

SAA-induced mice had significantly elevated PGE2, PGD2, and TXB2 relative to

radiation controls (**Figure 4C-E,** supplemental Fig. 9B, supplemental Table 2). Our data

suggest that in the context of SAA, omega-3 derived lipid precursors are decreased in

the marrow. Together, these data demonstrate that SAA pathogenesis correlates with

unregulated prostanoid and thromboxane synthesis supporting our conclusion that

resolution kinetics are defective.

Resolvin E1 provides therapeutic benefit in SAA

To address whether SPMs may be viable therapeutic targets in mitigating the pathogenesis of SAA, we first examined expression of key SPM receptors on BM monocytes. We determined by flow cytometry that ChemR23, the receptor for RvE1[42], was significantly upregulated on BM monocytes and neutrophils, but not on macrophages or LK cells (supplemental Fig. 10), which provided rationale for treating mice with exogenous RvE1.

To directly test the therapeutic efficacy of exogenous RvE1, we administered RvE1 or vehicle every other day, beginning at day 7 post-induction, a time point when marrow is hypocellular (**Figure 5A**). RvE1 treatment did not improve lymphopenia or anemia, or limit SAA-induced monocytosis (**Figure 5B**-**C**), however, a significant increase in circulating white blood cells and platelets was noted by 14 dpst (**Figure 5D-E**). Although not significant, we also observed increased overall BM cellularity by day 14 (**Figure 5F**). The number of BM monocytes remained unchanged; however, neutrophil numbers, total LK cells, and HSC progenitor populations were increased with RvE1 treatment (**Figure 5G-H;** supplemental Fig. 11). Moreover, we observed that RvE1 significantly improved survival when mice were administered therapeutic doses of RvE1, on days 7, 9, 11, and 13 post-SAA induction (**Figure 5I**).

Consistent with a role for RvE1 in improving resolution of inflammation, we observed that RvE1 treatment decreased expression of the inhibitory receptor SIRPα, (**Figure 6A**), which correlated with increased PS-MP$^+$ monocytes and F4/80$^+$ cells by 12

360   dpst (**Figure 6B-E**). Together, our data suggest that RvE1 therapy improves efferocytic

361   function and bone marrow cellularity, which rescued disease. Moreover, these studies

362   define a novel approach to treat autoimmune marrow failure.

Discussion

Efferocytosis is a crucial component of resolution pathways and maintains tissue

homeostasis by limiting inflammation. By engaging in specific ligand-receptor signaling,

such as PS and MerTK, efferocytes promote the resolution of inflammation by clearing

dying cells, releasing anti-inflammatory mediators like IL-10 and TGF-β, and driving

SPM biosynthesis.[20,37] Apoptotic cells that are not properly cleared enter secondary

necrosis, characterized by the breakdown of the cellular membrane and the leakage of

inflammatory DAMPs that may exacerbate inflammation.[20] Although impairments in

resolution are known to underly several chronic inflammatory diseases[43-46], whether

resolution responses are impaired during SAA or in other BMF diseases has yet to be

investigated.

Here we find an increase in the frequency of dead and dying cells within the BM

of SAA mice. While it is known that SAA is associated with increased apoptosis, the

accumulation of necrotic cells also suggested deficiencies in cell clearance. The

increase in dead and dying cells correlated with the expansion of a unique population of

SIRPα^{hi} MerTk^{hi} monocytes, not observed in radiation controls. Indeed, IFNγ is known to

drive monopoiesis[47], and earlier studies demonstrated that increased monocytes

preceded the accumulation of macrophages in the bone marrow during SAA

pathogenesis.[8] Consistent with a potential role of SIRPα^{hi} MerTk^{hi} monocytes in driving

disease, we noted that this population was significantly reduced in the MIIG mouse

model, a strain that is protected from developing SAA.[8] We reasoned that although the

monocytes highly expressed the efferocytic receptor MerTK, they were likely impaired in

efferocytic function due to co-expression of SIRPα, an inhibitory receptor.

Why MerTK expression is increased so strikingly requires further investigation,

though our data suggests that MerTK expression is driven by IFNγ during SAA. The

pathways necessary to initiate inflammation are also thought to program resolution[48,49],

and induction of *Mertk* expression may be one mechanism whereby IFNγ, typically

associated with acute inflammation, also participates in resolution. It is also known that

under inflammatory conditions MerTK is susceptible to cleavage by ADAM17,

generating soluble Mer (sMer).[50] The cleavage of MerTK is known to limit SPM

biosynthesis and recent studies in atherosclerosis demonstrated that increased MerTK

cleavage resulted in defective efferocytosis and disease progression.[37,51] Notably, sMer

can antagonize full-length Mer by binding to and sequestering the ligand Gas6 and

inhibiting Gas-mediated MerTK activation.[38] Our observation that SAA mice exhibited a

slight increase in sMer and a decrease in Gas6 in the marrow compared to radiation

controls, suggests that sMer may be contributing to improper MerTK activation and

deficiencies in resolution responses.

Healthy viable cells can expose PS under normal physiological conditions. To

prevent improper engulfment by phagocytes, healthy cells utilize "don't eat me"

markers, such as CD47. Dying cells typically lose or alter the expression of CD47 to

enable cell clearance, as in the case of aging RBCs, where CD47 is expressed on

normal RBCs, preventing their elimination by phagocytes.[41] Therefore, it is of no

surprise that anti-SIRPα therapy worsened anemia in SAA-induced mice. High CD47

expression on dying cells has also been shown to promote inefficient clearance via the

"nibbling" of cells, rather than whole-cell engulfment by phagocytes.[43] TNF and IFNγ,

which are increased during SAA[2], have been shown to upregulate CD47

expression.[52,53] As defective efferocytosis is known to elicit pro-inflammatory cytokines,

it is possible that CD47, TNF, and IFNγ play a critical role in the feedback loop that

inhibits cellular clearance and inflammation-resolution, thus potentiating inflammation.

413   Specialized pro-resolving mediators (SPMs) actively drive the resolution of

inflammation and a return to homeostasis.[23] Class switching of lipid mediators, from pro-

inflammatory to pro-resolving, is essential for resolution to occur.[54] Lipid mediator

analysis of the BM from SAA-induced mice demonstrated an imbalance of pro-

inflammatory to pro-resolving mediators with exuberant prostanoid synthesis. SPM

biosynthesis occurs via transcellular mechanisms that require close contact of cells

expressing distinct enzymes, and it is notable that during SAA the bone marrow

becomes increasingly hypocellular, which may prevent SPM generation. A loss of

proper SPM biosynthesis or signaling has been linked with the exacerbation of

inflammation in various disease settings.[55-57] While the cause is still unclear, a lack of

SPM production and an increase in prostanoid synthesis, combined with dysregulated

unalamation[58], appears to be contributing to persistent inflammation observed during

SAA. A recent report uncovered a prostanoid storm limited efferocytosis in vitro and in

atherosclerotic plaques[59], thus supporting the notion that sustained elevated

prostanoids, without a compensatory increase in SPMs, may contribute to defective

efferocytosis in the bone marrow during SAA.

429   Treatment with exogenous SPMs has been utilized to improve resolution in

multiple experimental settings.[25,45,60] SPMs have a unique safety profile since they,

unlike immunosuppressive therapies, do not compromise the host's natural immune

response.[27,61] During SAA, 18-HEPE, the precursor for the E-series resolvins is

significantly reduced, relative to radiation controls. Meanwhile, the receptor for RvE1,

ChemR23, was increased on monocytes and neutrophils in SAA. Together, these data

suggested that RvE1 could be an ideal therapeutic for treating SAA. Indeed, RvE1

therapy, starting at a timepoint where the BM is already hypocellular, improved platelet

count, BM cellularity, survival, and efferocytosis (**Figure 7**).

Current treatments for SAA rely on HSC transplantation, when possible, and IST,

though IST has a high refractory rate and hinders the patient's natural immune

response. Recently approved Etrombopag, an agonist of the thrombopoietin receptor,

has shown improved response times in combination with IST, although, overall

responses were not improved in older patients or those that began treatment with

severe disease.[62] Recent preclinical studies using the Janus kinase (Jak1/2) inhibitor

Ruxolitinib (Rux) demonstrated effectiveness in a murine model of SAA with markedly

reduced T lymphocytes in the marrow.[63] However, toxicity associated with Rux use has

been noted in humans and mouse models.[64,65] Therefore, improved therapeutic

strategies are still needed for durable treatment of SAA. Our findings that efferocytosis

was impaired in SAA, and that SPM therapy was efficacious, provide strong rationale for

targeting inflammation-resolution during BM failure. Therapies aimed at improving

resolution, perhaps in conjunction with lower dosing of IST, may provide a more

effective and safe treatment for SAA that does not compromise the patient's natural

immune response.

Word count: 3941

Acknowledgements

The authors would like to thank Jesse Bonin for technical assistance. This work was

supported by CDMRP Bone Marrow Failure Research Program-IDA (BM190079) to

K.C.M. This study was also supported in part by National Center for Research

Resources, National Institutes of Health Grant S10RR027926 (K.R.M.)

Authorship

R.G. and A.N.S. performed *in vivo* experiments, analyzed data, and prepared figures.

R.G. wrote the manuscript. K.R.M. provided oversight and guidance for LC-MS/MS

studies and analyzed data. G.F. contributed to experimental design and data analysis

for the project. K.C.M. conceived of the project, analyzed data, and wrote the

manuscript. The order of the first authors was determined on the basis of the effort and

contributions to writing of the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

References

1. Lin FC, Karwan M, Saleh B, et al. IFN-gamma causes aplastic anemia by altering hematopoietic stem/progenitor cell composition and disrupting lineage differentiation. *Blood*. Dec 11 2014;124(25):3699-708. doi:10.1182/blood-2014-01-549527
2. Sun W, Wu Z, Lin Z, et al. Macrophage TNF-alpha licenses donor T cells in murine bone marrow failure and can be implicated in human aplastic anemia. *Blood*. Dec 27 2018;132(26):2730-2743. doi:10.1182/blood-2018-05-844928
3. Zoumbos NC, Gascon P, Djeu JY, Young NS. Interferon is a mediator of hematopoietic suppression in aplastic anemia in vitro and possibly in vivo. *Proc Natl Acad Sci U S A*. Jan 1985;82(1):188-92. doi:10.1073/pnas.82.1.188
4. Merli P, Quintarelli C, Strocchio L, Locatelli F. The role of interferon-gamma and its signaling pathway in pediatric hematological disorders. *Pediatr Blood Cancer*. Apr 2021;68(4):e28900. doi:10.1002/pbc.28900
5. Gratwohl A, Osterwalder B, Nissen C, et al. [Treatment of severe aplastic anemia]. *Schweiz Med Wochenschr*. Oct 10 1981;111(41):1520-2. Behandlung der schweren aplastischen Anamie.
6. Xu L, Fu B, Wang W, et al. Haploidentical hematopoietic cell transplantation for severe acquired aplastic anemia: a case-control study of post-transplant cyclophosphamide included regimen vs. anti-thymocyte globulin & colony-stimulating factor-based regimen. *Sci China Life Sci*. Jun 2020;63(6):940-942. doi:10.1007/s11427-019-9585-x
7. Rosenfeld S, Follmann D, Nunez O, Young NS. Antithymocyte globulin and cyclosporine for severe aplastic anemia: association between hematologic response and long-term outcome. *JAMA*. Mar 5 2003;289(9):1130-5. doi:10.1001/jama.289.9.1130
8. McCabe A, Smith JNP, Costello A, Maloney J, Katikaneni D, MacNamara KC. Hematopoietic stem cell loss and hematopoietic failure in severe aplastic anemia is driven by macrophages and aberrant podoplanin expression. *Haematologica*. Sep 2018;103(9):1451-1461. doi:10.3324/haematol.2018.189449
9. Seyfried AN, McCabe A, Smith JNP, Calvi LM, MacNamara KC. CCR5 maintains macrophages in the bone marrow and drives hematopoietic failure in a mouse model of severe aplastic anemia. *Leukemia*. Nov 2021;35(11):3139-3151. doi:10.1038/s41375-021-01219-z
10. Philpott NJ, Scopes J, Marsh JC, Gordon-Smith EC, Gibson FM. Increased apoptosis in aplastic anemia bone marrow progenitor cells: possible pathophysiologic significance. *Exp Hematol*. Dec 1995;23(14):1642-8.
11. Liu CY, Fu R, Wang HQ, et al. Fas/FasL in the immune pathogenesis of severe aplastic anemia. *Genet Mol Res*. May 30 2014;13(2):4083-8. doi:10.4238/2014.May.30.3
12. Gupta P, Niehans GA, LeRoy SC, et al. Fas ligand expression in the bone marrow in myelodysplastic syndromes correlates with FAB subtype and anemia, and predicts survival. *Leukemia*. Jan 1999;13(1):44-53. doi:10.1038/sj.leu.2401233

13.   Michalski MN, Koh AJ, Weidner S, Roca H, McCauley LK. Modulation of
Osteoblastic Cell Efferocytosis by Bone Marrow Macrophages. *J Cell Biochem*. Dec
2016;117(12):2697-2706. doi:10.1002/jcb.25567
14.   McCauley LK, Dalli J, Koh AJ, Chiang N, Serhan CN. Cutting edge: Parathyroid
hormone facilitates macrophage efferocytosis in bone marrow via proresolving
mediators resolvin D1 and resolvin D2. *J Immunol*. Jul 1 2014;193(1):26-9.
doi:10.4049/jimmunol.1301945
15.   Wynn TA, Vannella KM. Macrophages in Tissue Repair, Regeneration, and
Fibrosis. *Immunity*. Mar 15 2016;44(3):450-462. doi:10.1016/j.immuni.2016.02.015
16.   Doran AC, Yurdagul A, Jr., Tabas I. Efferocytosis in health and disease. *Nat Rev
Immunol*. Apr 2020;20(4):254-267. doi:10.1038/s41577-019-0240-6
17.   Elliott MR, Koster KM, Murphy PS. Efferocytosis Signaling in the Regulation of
Macrophage Inflammatory Responses. *J Immunol*. Feb 15 2017;198(4):1387-1394.
doi:10.4049/jimmunol.1601520
18.   Shiratsuchi A, Osada S, Kanazawa S, Nakanishi Y. Essential role of
phosphatidylserine externalization in apoptosing cell phagocytosis by macrophages.
Biochem Biophys Res Commun. May 19 1998;246(2):549-55.
doi:10.1006/bbrc.1998.8663
19.   Fadok VA, Voelker DR, Campbell PA, Cohen JJ, Bratton DL, Henson PM.
Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers
specific recognition and removal by macrophages. *J Immunol*. Apr 1 1992;148(7):2207-
16.
20.   Korns D, Frasch SC, Fernandez-Boyanapalli R, Henson PM, Bratton DL.
Modulation of macrophage efferocytosis in inflammation. *Front Immunol*. 2011;2:57.
doi:10.3389/fimmu.2011.00057
21.   Gardai SJ, McPhillips KA, Frasch SC, et al. Cell-surface calreticulin initiates
clearance of viable or apoptotic cells through trans-activation of LRP on the phagocyte.
Cell. Oct 21 2005;123(2):321-34. doi:10.1016/j.cell.2005.08.032
22.   Okazawa H, Motegi S, Ohyama N, et al. Negative regulation of phagocytosis in
macrophages by the CD47-SHPS-1 system. *J Immunol*. Feb 15 2005;174(4):2004-11.
doi:10.4049/jimmunol.174.4.2004
23.   Serhan CN, Levy BD. Resolvins in inflammation: emergence of the pro-resolving
superfamily of mediators. *J Clin Invest*. Jul 2 2018;128(7):2657-2669.
doi:10.1172/JCI97943
24.   Hasturk H, Kantarci A, Ohira T, et al. RvE1 protects from local inflammation and
osteoclast- mediated bone destruction in periodontitis. *FASEB J*. Feb 2006;20(2):401-3.
doi:10.1096/fj.05-4724fje
25.   Ishida T, Yoshida M, Arita M, et al. Resolvin E1, an endogenous lipid mediator
derived from eicosapentaenoic acid, prevents dextran sulfate sodium-induced colitis.
Inflamm Bowel Dis. Jan 2010;16(1):87-95. doi:10.1002/ibd.21029
26.   Kantarci A, Kansal S, Hasturk H, Stephens D, Van Dyke TE. Resolvin E1
Reduces Tumor Growth in a Xenograft Model of Lung Cancer. *Am J Pathol*. Oct
2022;192(10):1470-1484. doi:10.1016/j.ajpath.2022.07.004
27.   Seki H, Fukunaga K, Arita M, et al. The anti-inflammatory and proresolving
mediator resolvin E1 protects mice from bacterial pneumonia and acute lung injury. *J
Immunol*. Jan 15 2010;184(2):836-43. doi:10.4049/jimmunol.0901809

28.   Haworth O, Cernadas M, Yang R, Serhan CN, Levy BD. Resolvin E1 regulates
interleukin 23, interferon-gamma and lipoxin A4 to promote the resolution of allergic
airway inflammation. *Nat Immunol*. Aug 2008;9(8):873-9. doi:10.1038/ni.1627
29.   Hasturk H, Abdallah R, Kantarci A, et al. Resolvin E1 (RvE1) Attenuates
Atherosclerotic Plaque Formation in Diet and Inflammation-Induced Atherogenesis.
Arterioscler Thromb Vasc Biol. May 2015;35(5):1123-33.
doi:10.1161/ATVBAHA.115.305324
30.   Norris PC, Skulas-Ray AC, Riley I, et al. Identification of specialized pro-
resolving mediator clusters from healthy adults after intravenous low-dose endotoxin
and omega-3 supplementation: a methodological validation. *Sci Rep*. Dec 21
2018;8(1):18050. doi:10.1038/s41598-018-36679-4
31.   Maddipati KR, Romero R, Chaiworapongsa T, et al. Eicosanomic profiling
reveals dominance of the epoxygenase pathway in human amniotic fluid at term in
spontaneous labor. *FASEB J*. Nov 2014;28(11):4835-46. doi:10.1096/fj.14-254383
32.   Chen J, Lipovsky K, Ellison FM, Calado RT, Young NS. Bystander destruction of
hematopoietic progenitor and stem cells in a mouse model of infusion-induced bone
marrow failure. *Blood*. Sep 15 2004;104(6):1671-8. doi:10.1182/blood-2004-03-1115
33.   Bloom ML, Wolk AG, Simon-Stoos KL, Bard JS, Chen J, Young NS. A mouse
model of lymphocyte infusion-induced bone marrow failure. *Exp Hematol*. Dec
2004;32(12):1163-72. doi:10.1016/j.exphem.2004.08.006
34.   Ossina NK, Cannas A, Powers VC, et al. Interferon-gamma modulates a p53-
independent apoptotic pathway and apoptosis-related gene expression. *J Biol Chem*.
Jun 27 1997;272(26):16351-7. doi:10.1074/jbc.272.26.16351
35.   Poon IK, Lucas CD, Rossi AG, Ravichandran KS. Apoptotic cell clearance: basic
biology and therapeutic potential. *Nat Rev Immunol*. Mar 2014;14(3):166-80.
doi:10.1038/nri3607
36.   Scott RS, McMahon EJ, Pop SM, et al. Phagocytosis and clearance of apoptotic
cells is mediated by MER. *Nature*. May 10 2001;411(6834):207-11.
doi:10.1038/35075603
37.   Cai B, Thorp EB, Doran AC, et al. MerTK cleavage limits proresolving mediator
biosynthesis and exacerbates tissue inflammation. *Proc Natl Acad Sci U S A*. Jun 7
2016;113(23):6526-31. doi:10.1073/pnas.1524292113
38.   Sather S, Kenyon KD, Lefkowitz JB, et al. A soluble form of the Mer receptor
tyrosine kinase inhibits macrophage clearance of apoptotic cells and platelet
aggregation. *Blood*. Feb 1 2007;109(3):1026-33. doi:10.1182/blood-2006-05-021634
39.   Lv Z, Bian Z, Shi L, et al. Loss of Cell Surface CD47 Clustering Formation and
Binding Avidity to SIRPalpha Facilitate Apoptotic Cell Clearance by Macrophages. *J
Immunol*. Jul 15 2015;195(2):661-71. doi:10.4049/jimmunol.1401719
40.   Burger P, Hilarius-Stokman P, de Korte D, van den Berg TK, van Bruggen R.
CD47 functions as a molecular switch for erythrocyte phagocytosis. *Blood*. Jun 7
2012;119(23):5512-21. doi:10.1182/blood-2011-10-386805
41.   Oldenborg PA, Zhelezynak A, Fang YF, Lagenaur CF, Gresham HD, Lindberg
FP. Role of CD47 as a marker of self on red blood cells. *Science*. Jun 16
2000;288(5473):2051-4. doi:10.1126/science.288.5473.2051

42.   Ohira T, Arita M, Omori K, Recchiuti A, Van Dyke TE, Serhan CN. Resolvin E1
receptor activation signals phosphorylation and phagocytosis. *J Biol Chem*. Jan 29
2010;285(5):3451-61. doi:10.1074/jbc.M109.044131
43.   Gerlach BD, Marinello M, Heinz J, et al. Resolvin D1 promotes the targeting and
clearance of necroptotic cells. *Cell Death Differ*. Feb 2020;27(2):525-539.
doi:10.1038/s41418-019-0370-1
44.   Rymut N, Heinz J, Sadhu S, et al. Resolvin D1 promotes efferocytosis in aging
by limiting senescent cell-induced MerTK cleavage. *FASEB J*. Jan 2020;34(1):597-609.
doi:10.1096/fj.201902126R
45.   El Kebir D, Gjorstrup P, Filep JG. Resolvin E1 promotes phagocytosis-induced
neutrophil apoptosis and accelerates resolution of pulmonary inflammation. *Proc Natl
Acad Sci U S A*. Sep 11 2012;109(37):14983-8. doi:10.1073/pnas.1206641109
46.   Martin-Rodriguez O, Gauthier T, Bonnefoy F, et al. Pro-Resolving Factors
Released by Macrophages After Efferocytosis Promote Mucosal Wound Healing in
Inflammatory Bowel Disease. *Front Immunol*. 2021;12:754475.
doi:10.3389/fimmu.2021.754475
47.   de Bruin AM, Libregts SF, Valkhof M, Boon L, Touw IP, Nolte MA. IFNgamma
induces monopoiesis and inhibits neutrophil development during inflammation. *Blood*.
Feb 9 2012;119(6):1543-54. doi:10.1182/blood-2011-07-367706
48.   Serhan CN. Pro-resolving lipid mediators are leads for resolution physiology.
Nature. Jun 5 2014;510(7503):92-101. doi:10.1038/nature13479
49.   Serhan CN, Savill J. Resolution of inflammation: the beginning programs the end.
Nat Immunol. Dec 2005;6(12):1191-7. doi:10.1038/ni1276
50.   Thorp E, Vaisar T, Subramanian M, Mautner L, Blobel C, Tabas I. Shedding of
the Mer tyrosine kinase receptor is mediated by ADAM17 protein through a pathway
involving reactive oxygen species, protein kinase Cdelta, and p38 mitogen-activated
protein kinase (MAPK). *J Biol Chem*. Sep 23 2011;286(38):33335-44.
doi:10.1074/jbc.M111.263020
51.   Cai B, Thorp EB, Doran AC, et al. MerTK receptor cleavage promotes plaque
necrosis and defective resolution in atherosclerosis. *J Clin Invest*. Feb 1
2017;127(2):564-568. doi:10.1172/JCI90520
52.   Kojima Y, Volkmer JP, McKenna K, et al. CD47-blocking antibodies restore
phagocytosis and prevent atherosclerosis. *Nature*. Aug 4 2016;536(7614):86-90.
doi:10.1038/nature18935
53.   Ye ZH, Jiang XM, Huang MY, et al. Regulation of CD47 expression by interferon-
gamma in cancer cells. *Transl Oncol*. Sep 2021;14(9):101162.
doi:10.1016/j.tranon.2021.101162
54.   Levy BD, Clish CB, Schmidt B, Gronert K, Serhan CN. Lipid mediator class
switching during acute inflammation: signals in resolution. *Nat Immunol*. Jul
2001;2(7):612-9. doi:10.1038/89759
55.   Chiang N, Dalli J, Colas RA, Serhan CN. Identification of resolvin D2 receptor
mediating resolution of infections and organ protection. *J Exp Med*. Jul 27
2015;212(8):1203-17. doi:10.1084/jem.20150225
56.   Kain V, Jadapalli JK, Tourki B, Halade GV. Inhibition of FPR2 impaired
leukocytes recruitment and elicited non-resolving inflammation in acute heart failure.
Pharmacol Res. Aug 2019;146:104295. doi:10.1016/j.phrs.2019.104295

57. Tourki B, Kain V, Pullen AB, et al. Lack of resolution sensor drives age-related cardiometabolic and cardiorenal defects and impedes inflammation-resolution in heart failure. *Mol Metab*. Jan 2020;31:138-149. doi:10.1016/j.molmet.2019.10.008

58. Maddipati KR. Non-inflammatory Physiology of "Inflammatory" Mediators - Unalamation, a New Paradigm. *Front Immunol*. 2020;11:580117. doi:10.3389/fimmu.2020.580117

59. Hosseini Z, Marinello M, Decker C, et al. Resolvin D1 Enhances Necroptotic Cell Clearance Through Promoting Macrophage Fatty Acid Oxidation and Oxidative Phosphorylation. *Arterioscler Thromb Vasc Biol*. Mar 2021;41(3):1062-1075. doi:10.1161/ATVBAHA.120.315758

60. Sulciner ML, Serhan CN, Gilligan MM, et al. Resolvins suppress tumor growth and enhance cancer therapy. *J Exp Med*. Jan 2 2018;215(1):115-140. doi:10.1084/jem.20170681

61. Spite M, Norling LV, Summers L, et al. Resolvin D2 is a potent regulator of leukocytes and controls microbial sepsis. *Nature*. Oct 29 2009;461(7268):1287-91. doi:10.1038/nature08541

62. Groarke EM, Patel BA, Gutierrez-Rodrigues F, et al. Eltrombopag added to immunosuppression for children with treatment-naive severe aplastic anaemia. *Br J Haematol*. Feb 2021;192(3):605-614. doi:10.1111/bjh.17232

63. Groarke EM, Feng X, Aggarwal N, et al. Efficacy of JAK1/2 inhibition in murine immune bone marrow failure. *Blood*. Jan 5 2023;141(1):72-89. doi:10.1182/blood.2022015898

64. Zeiser R, Polverelli N, Ram R, et al. Ruxolitinib for Glucocorticoid-Refractory Chronic Graft-versus-Host Disease. *N Engl J Med*. Jul 15 2021;385(3):228-238. doi:10.1056/NEJMoa2033122

65. Verstovsek S, Mesa RA, Gotlib J, et al. A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *N Engl J Med*. Mar 1 2012;366(9):799-807. doi:10.1056/NEJMoa1110557

Figure Legends

Figure 1. Cytopenias and BM hypocellularity are associated with increased cell
death in SAA. F1 hybrid mice were induced to develop severe aplastic anemia via the
radiation followed by splenocyte transfer model. Mice were euthanized 10 days post
splenocyte transfer (dpst) and complete blood count data are shown for total WBCs and
RBCs (**A**) and platelets (**B**). (**C**) The breakdown of all WBCs is shown. (**D**) Overall BM
cellularity in radiation controls and SAA mice. (**E**) Frequency of BM monocytes and
neutrophils. (**F**) Absolute number of BM monocytes (CD11b$^+$ Ly6C^{hi} Ly6G$^-$) and
neutrophils (CD11b$^+$ Ly6C^{low} Ly6G$^+$). Data shows two pooled independent experiments
showing mean ± SEM, n=7-10 per group, significance was determined using a
Student's t-test . ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$. (**G**) Gating strategy for AnnexinV
and 7-AAD staining to differentiate live, apoptotic, and dead BM cells, plots
representative for radiation controls and SAA mice. Numbers reflect the percent of cells
within the gated region. (**H**) Frequency of BM cells that are live (open), apoptotic (grey),
or dead (black) for both radiation controls and SAA mice days 8 and 10 post splenocyte
transfer. Data shows two pooled independent experiments per time point showing mean
± SEM, n=5-10 per group, significance was determined using a Student's t-test . *
$p<0.05$, *** $p<0.001$

Figure 2. The expansion of MerTK^{hi} SIRPα^{hi} monocytes and CD47 on apoptotic
cells in SAA mice. Mice were induced to develop SAA and BM harvested 10 days post
splenocyte transfer (dpst). (**A**) Representative gating for MerTK^{hi} monocytes. Numbers
on plots reflect the percent of cells within the gated region. The frequency and absolute
number of MerTK^{hi} monocytes is shown in **B** and **C** respectively. (**D**) SIRPα^{hi} staining on
MerTK^{hi} monocytes. The frequency (**E**) and absolute number (**F**) of SIRPα^{hi} MerTK^{hi}
monocytes. Data represents two pooled independent experiments showing mean ±
SEM, n=7-10 per group, significance using a Student's t-test. **** $p<0.0001$. (**G**) Gating
strategy for CD47 on live and apoptotic BM cells. Numbers on plots reflect the percent
of cells within the gated region. Frequency (**H**) and absolute number (**I**) of CD47 for live

and apoptotic cells. Data representative of one experiment showing mean ± SD, n=3-5

per group. Significance was determined using a Two-way ANOVA with Tukey's multiple

comparison test. **** $p<0.0001$.

Figure 3. Enhanced MerTK, SIRPα, and CD47 expression is associated with

impaired efferocytosis. (**A**) F1 hybrid mice were induced with SAA and BM harvested

8 dpst. Whole BM was flushed and incubated with PS lipid microparticles for 1 hour at

37°C. (**B**) The lipid microparticles contain attached PS with the phospholipid headgroup

exposed and trace amounts of a fluorescent lipid PE-NBD embedded in the particles.

(**C**) Representative staining of monocytes and frequency of cells positive for PS-coated

microparticles (PS-MPs). Numbers on plots reflect the percent of cells within the gated

region. (**D**) Average frequencies of PS-MPs-positive cells among total monocytes. (**E**)

Representative staining of macrophages and frequency of cells positive for PS-MPs.

Numbers on plots are the percent of cells within the gated region. (**F**) Average

frequencies of PS-MPs-positive cells among total F4/80$^+$ BM cells. Significance was

determined using a Student's t-test. ** $p<0.01$, *** $p<0.001$.

Figure 4. Imbalanced pro-Inflammatory- pro-resolving lipid mediators. Bone

marrow was collected and analyzed by LC-MS/MS. (**A**) Principal component analysis of

samples taken from mice without exposure to radiation (healthy), radiation only (RC), or

radiation plus splenocytes to induce SAA (SAA) day 3 post-radiation. Each dot

represents data from a single mouse and the size of the dot reflects concentration of 18-

HEPE. (**B**) Schematic of the biosynthesis of resolvins from ω-3 polyunsaturated fatty

acids. Fold change in concentrations of PGD2 (**C**) PGE2 (**D**), or TXB2 (**E**) in the bone

marrow of SAA mice, relative to radiation controls (RC), 8 dpst. Data representative of

two independent experiments that included 2-3 mice each. Significance was determined

using a Student's t-test. * $p<0.05$, ** $p<0.01$.

Figure 5. Exogenous RvE1 improves platelets, BM cellularity and survival. (**A**) SAA-induced F1 hybrid mice were treated with 250ng of RvE1 days 7, 9, 11, and 13 post induction. BM and blood was evaluated day 12 or 14. The breakdown of all WBCs (**B**) and total RBCs (**C**), WBCs (**D**), and platelets (**E**) are shown. (**F**) Total BM cellularity in F1 mice treated with vehicle or RvE1. Absolute number of BM monocytes and neutrophils (**G**), as well as the frequency of LK cells (**H**). Data representative of one experiment per dpst showing mean ± SD, n=5-8 per group. Significance was determined using a Student's t-test. * $p < 0.05$. (**I**) Percent survival 30 days post-induction; SAA mice treated with vehicle (solid black line) $n = 18$; SAA mice treated with RvE1 (black dashed line) $n = 19$. Data pooled from three independent experiments.

Figure 6. Exogenous RvE1 improves aberrant monocyte phenotype and efferocytosis. Mice were induced to develop SAA and treated with 250ng RvE1 days 7, 9, 11, and 13 post induction. BM was harvest day 12 or 14 dpst. Representative staining (**A**) and the frequency (**B**) of SIRPα on BM monocytes 14 dpst. Staining (**B** and **D**) and frequency (**C** and **E**) of PS-MPs on monocytes and F4/80$^+$ cells 12 dpst, respectively. Numbers on flow plots are the frequency of cells within the gated region. Data from one experiment with mean ± SD, n=5-8 per group. Significance was determined using a Student's t-test. * $p < 0.05$

Figure 7. RvE1 therapy reduces SIRPα^{hi} monocytes, increases efferocytosis, and improves resolution. Schematic model illustrating the role of impaired efferocytosis in dysfunctional inflammation resolution and the development of SAA. An increase in SIRPα^{hi} monocytes and CD47$^+$ apoptotic cells prevent clearance of apoptotic and dead cells, which is associated with disease progression (left panel). Exogenous RvE1 treatment decreases SIRPα^{hi} monocytes and increases efferocytosis, correlating with improved survival (right panel).

Figure 1

Figure 2

A Day 0: SAA Induction | **Day 8** | incubate with PtdSer-coated microparticles | examine uptake by flow cytometry

300 Rads + 6.5x10⁷ B6 (H-2^b/d) splenocytes

F1 Hybrid (H-2^b/d)

harvest bone marrow

B PE-NBD

PtdSer

3 µm

C Rad. Control   SAA

Ly6C

PS-Microparticles

8.73   3.12

D Freq PS-MPs (%Mono)

**

Rad   SAA

E Rad. Control   SAA

F4/80

PS-Microparticles

12.6   5.76

F Freq PS-MPs (%F4/80⁺)

***

Rad   SAA

Figure3

A PC scores

B

C

D

E

Figure 4

Figure 5

Figure 7