Title: Highly sensitive *in vivo* detection of dynamic changes in enkephalins following acute stress

Authors: Marwa O. Mikati[1,2,3], Petra Erdmann-Gilmore[4], Rose Connors[4], Sineadh M. Conway[1,2,3], Jim Malone[4], Justin Woods[1,2,3], Robert W. Sprung[4], R. Reid Townsend[4], Ream Al-Hasani [1,2,3].

[1]Department of Anesthesiology, [2]Washington University Pain Center, Washington University in St. Louis, MO, USA; [3]Center for Clinical Pharmacology, University of Health Science and Pharmacy in St. Louis, MO, USA, [4]Department of Medicine, Washington University School of Medicine, Washington University in St. Louis, St. Louis, MO, USA

Correspondence: Ream Al-Hasani, PhD al-hasanir@wustl.edu.

Abstract: Enkephalins are opioid peptides that modulate analgesia, reward, and stress. *In vivo* detection of enkephalins remains difficult due to transient and low endogenous concentrations and inherent sequence similarity. We present an analytical method for Met- and Leu-Enkephalin detection in the mouse Nucleus Accumbens shell after acute stress. This approach increases spatiotemporal resolution, optimizes the detection of Met-Enkephalin through methionine oxidation, and provides insight into the relationship between Met- and Leu-Enkephalin following stress.

Main

Endogenous opioid peptides are released at significantly lower concentrations than small molecule neurotransmitters and undergo post-translational processing that yields ~30 unique peptides from 4 precursor molecules. Together, these properties have made sensitive peptide detection difficult to achieve with existing technologies [1]. Here, we describe an analytical method that enables quantifying dynamic *in vivo* changes in both Leu- and Met-enkephalin (Leu-Enk and Met-Enk) peptides in rodents. We show unique release profiles between Leu- and Met-Enk in response to two different stressors and gain new insight into peptide release dynamics; plus our method offers improved spatiotemporal resolution. More broadly, this approach will enable understanding of how neuropeptide levels change in response to acute or chronic behavioral, pharmacological, and neurophysiological manipulations.

Enkephalins can act on both the delta and/or mu opioid receptors, but their activity often depends on receptor and peptide expression patterns, which differ across the brain [2–5]. Confirming an opioid peptide's identity based on the activity of the receptor alone poses difficulties. For example, in a single brain region, the mu opioid receptor may be activated by an enkephalin or endorphin peptide [3]. It has been particularly challenging to measure Met- and Leu-Enk selectively and independently *in vivo*; in fact, both peptides are typically referred to more generally as 'enkephalins', which oversimplifies their distinct properties. Though antibody-based techniques [6–8] or liquid chromatography/mass spectrometry (LC-MS) [9–12] have improved Met- and Leu-Enk detection somewhat, limitations such as spatiotemporal restrictions and poor selectivity between

enkephalins remain. Our group previously improved real time *in vivo* detection of evoked opioid peptide release using microdialysis/nanoLC-MS (nLC-MS) after photostimulation [13]. From this strong foundation, we have now optimized a novel opioid peptide detection technique. For the first time, we are able to selectively measure real-time *in vivo* levels of basal peptide release as well as during behavioral manipulations. Our primary goals in establishing this analytical workflow to measure opioid peptide dynamics *in vivo* were to 1) increase temporal resolution allowing for the detection of dynamic changes to short external stimuli and acute conditions, 2) lower the detection threshold to allow for accurate measurements from small brain regions and using less volume of sample, 3) measure endogenous release of peptides in the native state without the need for photo- or chemogenetic stimulation. To meet these goals and measure both Leu- and Met-Enk we used an improved and specialized microdialysis technique coupled with nLC-MS.

Our workflow (**Fig 1a**) starts with stereotaxic surgery to implant the microdialysis probe in the mouse nucleus accumbens shell (NAcSh). Along with others, we have shown that the opioid receptor system in the NAcSh mediates both reward and aversion through photostimulation of peptide-expressing neurons across the dorsal/ventral axis [14–20]. Our aim was to measure NAcSh peptide release in response to aversive, stressful stimuli. We use custom-made microdialysis probes that are similar in size to commonly used fiber photometry probes to avoid extensive tissue damage caused by traditional microdialysis probes (**Suppl. Fig. 1a**). After recovery, we begin microdialysis collection every 13 minutes, circulating artificial cerebrospinal fluid (aCSF) and collecting interstitial fluid (ISF) at a flow rate of 0.8 μL/min. After each sample collection, we add a consistent known concentration of isotopically labeled internal standard of Met-Enk and Leu-Enk to the collected ISF for the accurate identification and quantification of endogenous peptide. These internal standards have a different mass/charge (m/z) ratio than endogenous Met- and Leu-Enk. Thus, we can now assess the quality of the signal through confirming: the presence of three transitions, the shape of the peak, retention time (**Suppl Fig. 2**). Importantly, we have resolved one of the main barriers preventing the two peptides from being distinguished: the instability of the methionine residue of Met-Enk during sample processing. We doubly oxidize the samples overnight prior to nLC-MS analysis. Critically, this double oxidation allows us to reliably and sensitively detect Met-Enk, clearly differentiating it from Leu-Enk [21]. Beyond opioid peptides, this approach represents a key advance and can be applied to any peptide containing a methionine residue. As seen in **Suppl. Fig. 2e**, before

the methionine oxidation reaction, Met-Enk can be observed in unoxidized, and singly- and doubly-oxidized forms with varying intensities. However, after the reaction is complete, most of the Met-Enk is observed as doubly oxidized with much higher intensity enabling accurate quantification (**Suppl. Fig. 2f)**. To increase the signal to noise ratio, we also perform solid-phase extraction after the oxidation reaction to remove any competing species in the collected samples. Using forward and reverse curves using varying concentrations of Met- and Leu-Enk standards (**Fig. 1a**) we determined that the lower limit of quantification (LLOQ), for both Met-Enk and Leu-Enk is 40 amol/sample (**Figure 1b-c**). The improved quantification limit means increased temporal resolution. We now need less volume, (10 μL or less), and shorter sampling times: 10 minutes or less, rather than 15-20 minutes as compared to previous studies [13,22]. To test the utility of the method in measuring real-time changes in peptide levels *in vivo*, we evoked the release of the peptides using high K^+ ringer's solution, which depolarizes the cells and causes the release of dense core vesicles containing neuropeptides [23]. This depolarization state significantly increased the release of Leu- and Met-Enk in the NAcSh **Figure 1d**. We then quantified the ratio of the Met- to Leu-Enk after applying high K+ solution and showed that this ratio is 3:1 in the NacSh (**Fig. 1e-f**). The release of Met-Enk is postulated both in vitro and ex vivo to be between 3 to 4 times that of Leu-Enk [24–26]. We are the first to not only corroborate the predicted ratio *in vivo,* but specifically in the NAcSh. Interestingly, this 3:1 ratio is almost identical to that previously reported in homogenized human tissue from the sympathoadrenal systems using radioimmunoassays [27]. Importantly, this conserved ratio highlights the usefulness of rodent models for investigating the highly conserved opioid system and and demonstrates its potential in translation to humans.

To highlight the applicability and value of this improved approach, we investigated the changes in the levels of enkephalins following acute stress. It has been hypothesized that enkephalins act as anti-stress signals based on studies that have shown increased preproenkephalin transcript levels following acute stress [28–31]. Moreover, decreased enkephalin expression in the NAc was correlated with an increase in the susceptibility to social defeat [32]. Our group has also recently demonstrated that the photostimulation of peptidergic neurons in the ventral NAcSh drives aversion [14], which further suggests a role for neurons in the NAcSh during aversive stimuli such as stress. Despite the wealth of data implicating these peptides in the stress response, it has not been possible to measure the dynamics of real-time peptide release in response to acute stressful stimuli. We

sought to test the hypothesis that Met- and Leu-Enk are released in the NAcSh following acute stress. Using the nLC-MS detection method, we successfully monitored dynamic changes of Met- and Leu-Enk during two different stressors. The first stressor was defined as experimenter handling, which involves the experimenter scruffing the mouse and connecting the inlet tubing and outlet tubing of the microdialysis setup to the microdialysis probe implanted in the mouse brain (**Fig. 1a**). The second stressor was exposure to predator odor (i.e., fox urine). We show that both Met- and Leu-Enk are released following experimenter handling, and that the levels decrease within minutes after exposure to the stress (**Fig. 2a**). We also analyzed the data with sex as a main effect and did not find any differences between males and females after correcting for multiple comparisons (**suppl. Fig. 3a**). We corroborate our finding from **Fig. 1e-f** showing that the ratio of 3:1 Met-Enk to Leu-Enk holds following experimenter handling (**Fig. 2c**). Another key finding is that peptide release appears to be precisely controlled. We show that the amol/sample of Met-Enk released at the beginning of the experiment is predictive of later release. For example, a higher level released (in the fmol/sample range) during the first collections results in lower released (in the amol/sample range) during later collections and vice versa (**Fig. 2d**). We show a negative correlation (~ -0.3) between the first two samples (~26 minutes) and the later timepoints, samples 3-10 (~104 minutes). Importantly, this relationship is not due to Met-Enk depletion as the circulation of high K+ aCSF is sufficient to drive the release of Met-Enk at the end of an experiment and reverse the negative correlation (**Fig. 2e**).

To test whether our findings regarding peptide release correlated with neuronal activity following the same stressor, we also monitored enkephalinergic neuron activity using fiber photometry following experimenter handling. To do this, we stereotaxically injected a Cre-dependendent adeno-associated virus containing the calcium sensor GCaMP6s in the NAcSh of preproEnkephalin-Cre (pEnk-Cre) mice and implanted fiber photometry probes (**Fig. 2f**). During handling, we observed increased calcium activity in enkephalinergic neurons, even after it is repeated twice within a few minutes (**Fig. 2g**). Individual response to experimenter handling varied appreciably, as shown by the heatmaps and bar plots (**Fig. 2h-i**); such variation could be due to differences in probe placement (**Suppl. Fig 2c-d**) as peptidergic neurons in the dNAcSh encode reward while neurons in the vNAcSh encode aversion [14,33]. Furthermore, the enkephalin precursor is expressed across the dorsal-ventral axis as shown in the *in situ* hybridization atlas (Allen Brain Institute). The fiber photometry data

and peptide release data are concordant, as enkephalinergic neurons are activated following experimenter handling, which drives Met- and Leu-Enk somatodendritic release. Critically, the photometry and peptide release data differ in that repeated neuronal activation may not necessarily lead to repeated peptide release, as suggested by the decrease in peptide release over time in **Fig. 2d**. Additionally, fiber photometry cannot provide information about Met- and Leu-Enk independently, as it relies on the precursor gene, *penk*. Taken together, our data suggest that fiber photometry results must be coupled with in vivo measurements of neuropeptides to determine release dynamics.

Prior reports have shown that fox urine leads to an increase in the expression of the neuronal activation-associated transcription factor fos in enkephalin-positive neurons in the NAcSh [34]. Therefore, we sought to determine whether enkephalins are released in the NAcSh following the exposure to fox urine. We first measured calcium activity of enkephalinergic neurons using fiber photometry after the exposure to fox urine (**Fig. 3a**). We show that upon first exposure, the activity of enkephalinergic neurons increases markedly (**Fig. 3b-d**). However, this enkephalinergic activity is significantly attenuated during second and third exposures that are separated by a few minutes suggesting habituation to the stimulus (**Fig. 3b-d**). We also wondered whether the response to fox urine was also due to the weigh boat used to introduce the fox urine, as it represents a novel object in the home cage, which could be stressful. Using fiber photometry, we measured enkephalinergic calcium activity after introducing a weigh boat containing water to the home cage (**suppl. Fig. 4a**). Interestingly, our data suggest that enkephalinergic neurons also respond to the introduction of a novel object, and the response to the water-containing weigh boat is attenuated upon repeated exposure, which is what we observed for fox urine (**Suppl. Fig. 4b-d**). We compared the enkephalinergic neuron activation in response to both water and fox urine for each individual mouse that experienced both. We see similar calcium transient responses to both the introduction of a novel object and the odor of a predator, suggesting the same subpopulation of enkephalinergic neurons may have been activated (**Suppl Fig. 4e**). Interestingly, only recently have researchers shown that a higher level of spiking in the striatum is not always reflected in greater photometry transients. In the future, it will be important to determine whether the fox urine response indeed shows higher action potential spiking than the weigh boat with water [35]. Although the calcium responses to both the fox-urine and water-containing weigh boats were similar in magnitude, it may be that different subpopulations of enkephalinergic neurons are activated in the

NacSh. Further exploration of the specific neurons that respond to the introduction of a novel object or the odor of a predator is warranted. We concluded that enkephalinergic neurons respond to predator odor and novel object introduction; therefore, we tested whether enkephalinergic neuron activation leads to Met- and Leu-Enk release in the NAcSh following predator odor exposure (**Fig. 3e**). In this separate group of animals, we show that there is a significant increase in Met-Enk release (**Fig. 3f**). The response to predator odor varies appreciably: although most animals show increased Met-Enk release, some show continued release with no elevation in Met-Enk levels, and a minority show no release (**Fig. 3g**). Enkephalins are thought to mediate resilience to stress, so the individual differences in responses to predator odor may indicate differences in stress adaptability [31,36–38]. Moreover, no sex differences in the responses to fox urine were observed, similar to responses to experimenter handling (**Suppl. Fig. 3b**). It is also worth noting that we only observe Leu-Enk release in 3 of 16 experiments. This is likely due to Leu-Enk not being released following predator odor or because it is released at a level below the LLOQ, as suggested by the observed 3:1 ratio of Met-Enk to Leu-Enk (**Fig. 3g**).

This work allows for multiple unique advances in studying endogenous opioids and neuropeptides in general. First, we have developed a method of methionine stabilization that is transferrable to other neuropeptides and have decreased the LLOQ and temporal resolution. Second, we quantified the relationship between Leu- and Met-Enk and offer a definitive *in vivo* ratio that has long been in question; this ratio corroborates human data and suggests that mice are a useful model system for the study of enkephalinergic release dynamics. Third, we identify a role for the NAcSh in the acute stress response, which has been suggested by studies showing changes in transcript levels of the precursor molecule of enkephalins [32,39]. Fourth, when a limited amount of peptide is released over time, we show that peptide release is biologically constrained and can be experimentally driven using high K+ buffer or photo or chemogenetic stimulation [13]. Finally, we show that Met- and Leu-Enk are released during acute stress, offering evidence for their role in gating the stress response [31]. We also show that data from fiber photometry corroborate peptide release data; however, fiber photometry does not offer the necessary specificity to correlate activation with release, or the distinction between different peptides that arise from the same precursor molecule. Importantly, this method can immediately be applied to Met- and Leu-Enk samples from other species – enabling the detection of these endogenous opioids in human CSF samples.

Figures and Legends:

Figure 1: An optimized approach for *in vivo* Met- and Leu-Enk measurement. **a.** Timeline of *in vivo* sample collection on day 1 and methionine oxidation reaction overnight, sample processing on day 2, and data acquisition on the LC-MS, days 3-5. The microdialysis probe is implanted via stereotaxic surgery in the ventral Nucleus Accumbens Shell. The mouse is allowed to recover before being connected to the microdialysis lines. Samples are then collected at a rate of 0.8 μL/min for 13 minutes each. After collection is completed, the samples are oxidized overnight. On day 2 the samples undergo solid phase extraction (SPE) and are then dehydrated and reconstituted using formic acid (HCOOH) before being acquired on the LC-MS. **b**. Forward curves of Leu-Enk and Met-Enk showing the peak area ratios as the light standard levels are varied. **c**. Reverse curve of Leu-Enk and Met-enk showing the relationship between the heavy standard concentration applied and the measured concentration based on the instrument. **d.** Violin plots showing that high K+ ringer's solution increases the release of both Leu-Enk and Met-Enk compared to baseline levels in artificial cerebrospinal fluid (aCSF) (Leu-Enk n=9, Met-Enk n=18). The dashed line indicates the median. **e.** The evoked concentrations of Met-Enk to Leu-Enk in the same samples show that Met-Enk is consistently released at a higher level than Leu-Enk (n=9). **f.** Met-Enk is released at a factor of 2.97 that of Leu-Enk as shown by linear regression analysis of the data in f, suggesting a linear relationship between the two peptides. Data in **d**, **e**, and

7 **f** are transformed to a log scale. In **d**, 2-way ANOVA analysis shows a main effect of peptide, solution, and
8 interaction. The *p*-values reported were calculated using a Sidak's multiple comparisons test.

Figure 2: Experimenter handling evokes the release of Met- and Leu-Enk in the NAcSh. a. Experimenter handling during microdialysis causes a significant increase in the release of Leu- and Met-Enk in comparison to baseline. **b.** During experimenter handling, Met-Enk is consistently released at a higher rate than Leu-Enk in the same samples. **c.** Linear regression analysis of the data in **b** shows that Met-enk is released at a rate of 3.12 times the rate of Leu-Enk during experimenter handling suggesting a linear relationship between the two peptides. **d.** A negative correlation (-0.3141) shows that if a high concentration of Met-Enk is released in the first two samples, the concentration released in later samples is affected; such influence suggests that there is regulation of the maximum amount of peptide to be released in NAcSh. **e.** The negative correlation in panel d is reversed by using high K+ buffer to evoke Met-Enk release, suggesting that the limited release observed in **d** is due to modulation of peptide release rather than depletion of reserves. Data in **a-e** are transformed to a log scale and n=24 animals. In panel **a**, 2-way ANOVA analysis shows a main effect of peptide, treatment, and interaction. The *p*-values reported were calculated using a Sidak's multiple comparisons test. **f.** Schematic describing the viral strategy and probe placement for the fiber photometry experiment in pEnk-Cre mice injected with the calcium sensor GCaMP6s in the NAcSh. **g.** Average z-score trace following the first, second, and third events involving experimenter handling. **h**. Heatmaps showing individual mouse z-scored fiber photometry responses before and after experimenter handling. **i**. Bar graphs showing the averaged z-score responses before experimenter handling and after experimenter handling (n=10 for **g-i**).

Figure 3. Fox odor exposure activates enkephalinergic neurons and drives the release of Met-Enk in the NAcSh. a. Viral strategy and probe placement for the fiber photometry experiment in pEnk-Cre mice injected with the calcium sensor GCaMP6s in the NAcSh. **b.** Averaged z-score traces of the first, second, and third exposure to fox odor. **c.** Bar graphs showing the averaged z-scored fiber photometry responses before and after exposure to fox odor **d**. Heatmaps showing individual mouse z-scored fiber photometry responses before and after exposure to fox odor. (n=9, for **a-d**) **e**. Experimental timeline describing the fox odor microdialysis experiment. **f**. There is a significant increase in Met-Enk concentration during fox odor exposure (sample 4), in comparison to sample, 3 before odor exposure. Data is transformed to a log scale and the *p*-value was calculated using a paired t-test (n=11). **g.** Heatmaps showing the variation in Met-Enk and Leu-Enk

release profiles in response to exposure to fox odor, suggesting that such exposure selectively causes the release of Met-Enk but not Leu-Enk in most experiments.

Materials and Methods:

List of Abbreviatons:

FA, formic acid
MeCN, acetonitrile
MeOH, methanol
DIA, Data-Independent Acquisition
AGC, Automatic Gain Control
nano-LC-MS, capillary liquid chromatography interfaced to a mass spectrometer
MS1, mass spectra of peptide precursors
MS2, fragmentation mass spectrum of peptide from precursor ion

Reagents

Acetonitrile (MeCN; J.T. Baker, cat. no. 9829-03)

Water (J.T. Baker, cat. no. 4218-03)

Formic Acid (FA; Sigma-Aldrich, cat. no. 56302)

Hydrogen Peroxide (Fisher Chemical, cat. no. H325-100)

Methanol (MeOH; Fluka, cat. no. 34966)

Equipment

Stage tips

Laboratory pipetting needles with 90° blunt ends, 16-gauge, 2-inch length; Cadence Science, cat. no. 7938)

C18 extraction disks, diam. = 47 mm, 20 pack; (Empore, cat. no. 66883-U)

Adapter for stage tipping (Glygen, cat. no. CEN.24)

1.5-ml tubes (Axygen, Cat. No. MCT-175-C)
Eppendorf centrifuge (Eppendorf, model 5424R)

Vortex (Labnet VX100)

Autosampler vials (Sun-Sri, Cat. No. 200046)

Autosampler vial caps (Sun-Sri, cat. no. 501 382)

UPLC system for LC-MS of analysis of TMT-labeled peptides (EASY nL-LC 1000, Thermo Scientific)

Mass spectrometer for analysis of TMT-labeled peptides (Q-Exactive™ Plus Hybrid Quadrupole Orbitrap™, Thermo Scientific)

EASY-Spray™ column, 75 µm x 50 cm PepMap (Thermo Scientific™, cat. no. ES903)

Animals

Adult male and female C57BL/6 mice at 8-15 weeks of age were used for microdialysis procedures. For fiber photometry, adult preproEnkephalin-Cre (pEnk-Cre) male and female mice at ages between 15-24 weeks of age were used to measure enkephalinergic cell activity. Mice were group housed and kept at a 12-hour

light/dark cycle with *ad libitum* access to water and food. The holding facility was both temperature and

humidity controlled. All surgical and behavioral procedures were approved by the Washington University

School of Medicine animal care guidelines in accordance with federal animal use regulations.

Chemicals

Met and Leu Enkephalin standards including isotopically labeled forms were synthesized at New England

Peptide and can be ordered by the research community. The heavy isotope labeled Leu Enkephalin is

synthesized with a fully labeled L-Leucine ($^{13}C_6$, ^{15}N) at the C terminus of the peptide (YGGF**L**). For Met

Enkephalin, a fully labeled L-Phenylalanine ($^{13}C_9$, ^{15}N) was added (YGG**F**M). The resulting mass shift between

the endogenous (light) and heavy isotope-labeled peptide are 7Da and 10Da, respectively. Heavy isotope

amino acids are guaranteed >99% pure. Peptide purity specifications are set at ≥ 95% pure by area on a

regular HPLC gradient with UV detection. Artificial cerebrospinal fluid (aCSF) contained 124 mM NaCl, 2.5 mM

KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$, 1.25 mM NaH_2PO_4, 24 mM $NaHCO_3$, 5 mM HEPES, 12.5 mM Glucose

adjusted to pH 7.4 with NaOH. High K^+ ringer solution contained 48 mM NaCl, 100 mM KCl, 2.4 mM $CaCl_2$,

0.85 mM $MgCl_2$ adjusted to pH 7.4 with NaOH as described in (Al-Hasani, 2018). All salt compounds were

purchased from Sigma (St. Louis, MO).

***In vivo* microdialysis**

Custom microdialysis probes (BASI Inc. part MD-2206) of 5 mm shaft length and 1 mm Polyacrylonitrile

membrane with a 30 kDa molecular weight cutoff were used for the *in vivo* experiments. The probes were

flushed prior to surgical implantation for 15 minutes at 10 μL/min with artificial cerebrospinal fluid as directed by

the manufacturer using a BASI syringe pump (MD 1001). For the surgical procedure, mice were briefly

anaesthetized in an induction chamber at 3% isofluorane before being placed in a stereotaxic frame (Kopf)

secured with ear and bite bars. During the surgery, the isofluorane level was kept at 1.5-2%. The probe was

then inserted into the ventral Nucleus Accumbens shell (stereotaxic coordinates from bregma: +1.3 [AP],±0.5

[ML], −5.0 mm [DV]). Implanted probes were secured with dental adhesive (C&B metabond kit) followed by

cyanoacrylate (Loctite superglue gel control).

After the completion of the surgery, each mouse was allowed to recover for 30 minutes. The mouse was then

connected to an inlet perfusion line to circulate aCSF or high K+ ringer solution and an outlet perfusion line to

collect the interstitial fluid from the NAcSh. The outlet line directly collected sample in a tube that is kept at 4

°C throughout the experiment. The flow of aCSF was then then turned on and sample collection time started as sample began to collect in the outlet tube. The flow rate of aCSF was set to 0.8 μL/min during sample collection. Fractions were collected every 13 minutes resulting in a 10 μL sample. After each collection, the samples were spiked with 2 μL of 12.5 fM isotopically labeled Met-Enkephalin and Leu-Enkephalin. The microdialysis rig (BASI Raturn MD 1404) was equipped with bedding, regular mouse chow and water ad libitum. While connected to the perfusion lines, the mice were allowed to move freely. For the experiments measuring evoked release, the high K+ ringer solution was perfused through the lines instead of aCSF at 0.8 μL/min and fractions were collected every 13 minutes as described previously. The samples were similarly spiked with 2 μL of 12.5 fM isotopically labeled Met-Enkephalin and Leu-Enkephalin.

For the experiments where the animals were exposed to the fox urine odor, 3 mL of fox urine was absorbed into a kim wipe contained in a chemical weigh boat. The weigh boats were prepared and remained in a fume hood the experiment to prevent the odor permeating the experimental space before the planned collection. Then, the experimenter placed the weigh boat in the microdialysis rig and allowed the mouse to explore it. The exposure was standardized to span 3 fractions (39 minutes) during each of the experiments while aCSF was being perfused through the lines.

In vivo sample processing

Methionine Oxidation Reaction

For met-enkephalin detection and stabilization, the animal samples underwent a methionine oxidation reaction. To each sample vial, 28 μl of 0.1% (vol/vol) formic acid (FA) was added and then the samples were transferred to 0.5 mL tubes from the collection tubes. Afterwards, 40 μl of a mixture of 6% FA (vol/vol) and 6% hydrogen peroxide (vol/vol) was added to each of the sample tubes and mixed and gently spun. The samples were then incubated overnight at room temperature.

Solid Phase Extraction with C18 stage tips

After the methionine oxidation reaction, a solid phase extraction process was performed using C18 stage tips. Stage tips were conditioned before sample loading. To each of the stage tips, 100 μl of methanol is added and then spun for 2 minutes at 2,000 rpm in microcentrifuge tubes. The stage tips were then rinsed with 100 μl of a mixture of 50% acetonitrile (MeCN; (vol/vol)) and 0.1% FA (vol/vol) followed by a 2-minute spin at 2,000 rpm. This was followed by a rinse in 0.1% FA (vol/vol) and a 2-minute spin at 3,000 rpm. The final rinse was done

by adding 100 µl of 1% FA (vol/vol) to each of the stage tips. Following the final rinse, the samples were added

to the C18 stage tips and spun for 10 minutes at 0.8 rpm. After ensuring that the samples passed through the

stage tips, the samples were desalted with 0.1% FA (vol/vol) and spun for 1.5 minutes at 3,000 rpm twice. After

the spins were done, the stage tips were directly placed into autosampler vials to elute the analyte with 60 µl of

50% MeCN, 0.1% FA (vol/vol) using a syringe. Eluates in the autosampler vial were dried in a speedvac,

reconstituted in 3 µl 1% FA (vol/vol) and set up on the autosampler for injection on the mass spectrometer.

***nano*-LC-MS/MS.** The samples in formic acid (1%) were loaded (2.5 µl) onto a 75 µm i.d. × 50 cm Acclaim®

PepMap 100 C18 RSLC column (Thermo-Fisher Scientific) on an EASY *nano*LC (Thermo Fisher Scientific) at

a constant pressure of 700 bar at 100% A (1%FA). Prior to sample loading the column was equilibrated to

100%A for a total of 11 µL at 700 bar pressure. Peptide chromatography was initiated with mobile phase A (1%

FA) containing 5%B (100%MeCN, 1%FA) for 1 min, then increased to 20% B over 19 min, to 32% B over 10

min, to 95% B over 1 min and held at 95% B for 19 min, with a flow rate of 250 nL/min. The data was acquired

in data-independent acquisition (DIA) mode. The full-scan mass spectra were acquired with the Orbitrap mass

analyzer with a scan range of m/z = 150 to 1500 and a mass resolving power set to 70,000. Data-independent

(DIA) high-energy collisional dissociations were performed according to the inclusion list with a mass resolving

power set to 17,500, an isolation width of 2 Da, and a normalized collision energy setting of 27. The maximum

injection time was 60 ms for parent-ion analysis and product-ion analysis. The automatic gain control (AGC)

was set at a target value of 3e6 ions for full MS scans and 2e5 ions for MS2. The inclusion list contained 4

peptide entries. The loop count was set at 30. Retention time windows were set at 20 min.

Stable-isotope dilution assay for measurement of enkephalins

Peptides were synthesized as high purity natural abundance and stable isotope-labeled pairs. Assays were

developed as Tier 2 assays according to CPTAC guidelines (Mol. Cell. Proteomics. 2014;13:907–917). Co-

eluting peak boundaries for the internal standard and endogenous peptides were manually refined using the

Skyline software package (Bioinformatics. 2010 Apr 1;26(7):966-8. doi: 10.1093/bioinformatics/btq054. Epub

2010 Feb 9. PMID: 20147306). Integrated peak areas for the three most intense fragment ions were then

exported for determination of peak area ratios of endogenous to internal standard signal, which were used as a

basis for determining abundance and quantifying changes in enkephalin levels in mouse studies.

In vivo fiber photometry

Stereotaxic surgery

For the surgical procedure, mice were briefly anaesthetized in an induction chamber at 3% isofluorane before being placed in a stereotaxic frame (Kopf) secured with ear and bite bars. During the surgery, the isofluorane level was kept at 1.5-2%. Then we performed a craniotomy and unilaterally injected using a Nanoject III (Drummond) 200-250 nL of AAV5.Syn.Flex.GCaMP6s.WPRE.SV40 (Addgene: 100845-AAV5) at the following coordinates :+1.3 [AP],±0.5 [ML], −5.0 mm [DV]. To allow for viral expression, the fiber photometry probes (5 mm shaft length, doric lenses) were implanted 3 weeks later at +1.3 [AP], ±0.5 [ML], −4.75 mm [DV]. Implanted probes were secured with dental adhesive (C&B metabond kit) followed by cyanoacrylate (Loctite superglue gel control).

Fiber photometry recording procedure

Before recording, the implanted probes were tethered to an optic fiber using a ferrule sleeve (Doric, catalog no. ZR_2.5). Two light-emitting diodes (LEDs) were used to excite GCaMP6s. A 331 Hz sinusoidal LED light (Doric) of wavelength 470 nm 470 ± 20 nm was used for the Ca^{2+} -dependent signal. A 210 Hz sinusoidal LED light of wavelength 405 ± 10 nm (Doric, LED driver) was used to evoke the Ca^{2+} -independent signal and serve as the isosbestic control. GCaMP6s fluorescence traveled through the same optic fiber before being bandpass filtered (525 ± 25 nm, Doric, catalog no. FMC4), transduced by a femtowatt photoreceiver (New Focus Model 2151) and recorded by a real-time processor (TDT, RZ5P). The signals were extracted by the Synapse software (TDT). During the recording, two behavioral manipulations were introduced. The first was after a 10-minute recording in a home cage environment, an experimenter handled the mouse while the recording was ongoing. The mouse was then let go and the handling was repeated another two times in a single session. The second behavioral manipulation was introduced to the same mice a week after the first recording session. After a 10-minute recording in the home cage environment, a weight boat containing a kim wipe that had absorbed 3 mL of fox urine was placed in the cage as the recording proceeded. The weigh boat was removed after a couple of minutes of exploration then re-introduced into the cage for another two times during the session.

Fiber Photometry Data Analysis

Data analysis was performed through the python toolbox GuPPy [40]. Briefly, the isosbestic control signal was used to correct for motion and photobleaching artifacts. The control channel is fitted to the signal channel using a least squares polynomial fit of degree 1. Transient detection occurred after using a 15s moving window for

thresholding transients. Then, high-amplitude events were identified and their timestamp of occurrence during the session was reported. To determine the effect of handling or fox urine exposure on enkephalinergic cell activity, the event timestamp was used to observe changes in activity before and after the behavioral manipulation. The average of all trials is then calculated for repeated exposures and for exposures in multiple animals. The fiber photometry data was represented as z-scores.

Immunohistochemistry and verification of probe placement

After the completion of the experiment, mice were anaesthetized with a mixture of ketamine, xylazine and acepromazine and transcardially perfused with ice-cold 4% paraformaldehyde in phosphate buffer. After the perfusion, brains were dissected and fixed for 24 hours in 4% paraformaldehyde at 4°C and then placed in 30% sucrose solution for cryoprotection for a minimum of 48 hours. The brains were then sliced into 30 um sections, washed three times in PBS and blocked in PBS containing 0.5% Triton X-100 and 5% goat serum for one hour. Following the blocking step, the slices were stained for the glial fibrillary acidic protein (GFAP), an astrocyte marker to determine glial scarring around the probe site (guinea pig anti-GFAP, synaptic systems) at 1:500 dilution and incubated at 4°C for 16 hours. The next day, the sections were washed three times with PBS and stained with the secondary antibody, mouse anti-guinea pig, Alexa Fluor 594 (Invitrogen) at 1:500 dilution and incubated for two hours at room temperature. After the incubation, the slices were washed three times in PBS, followed by two washes with PB. Finally, the slices were mounted on glass slides with Hard set Vectashield with DAPI staining (Vector Labs) and imaged using an epifluorescence microscope (Leica DM6 B). Correct probe placement in the NAcSh is represented in **supplementary figure 1b**.

For the verification of the injection and probe placement for fiber photometry, the same procedure described above was performed for tissue preparation. After slicing, the slices were washed in a 0.1% Triton for one hour in order to permeabilize the cells followed by two 10-minute PBS washes. The slices were then stained with Neurotrace (Nissl) red (Invitrogen) to mark neurons by adding 5 μL to each well for 2 hours at room temperature followed by 3 10-minute PBS washes and 3 10-minute PB washes. Finally, the slices were mounted using Hard set Vectashield (Vector Labs). GCaMP6s expression was determined by observing green fluorescent protein expression colocalized with Neurotrace staining. Correct probe placement in the NAcSh is represented in **supplementary figure 1d**.

Funding

This work was supported by National Institutes on Drug Abuse R00 DA038725 (RA) and R21DA048650 (RA). The NARSAD Young Investigator Grant from the Brain and Behavior Research Foundation, grant no. 28243 (RA). The WU-PSR is supported in part by the WU Institute of Clinical and Translational Sciences (NCATS UL1 TR000448), the Mass Spectrometry Research Resource (NIGMS P41 GM103422; R24GM136766) and the Siteman Comprehensive Cancer Center Support Grant (NCI P30 CA091842). Cognitive, Computational, and Systems Neuroscience Fellowship (MOM).

Author Contributions

- Marwa Mikati conducted the *in vivo* experiments, analyzed the data, and wrote the manuscript with the assistance and supervision of Robert Sprung PhD, Reid Townsend MD, Ph.D., and Ream Al-Hasani Ph.D.

- Petra Erdmann-Gilmore and Rose Connors processed and analyzed the microdialysis samples for nLC-MS.

- Sineadh Conway, PhD assisted with *in vivo* microdialysis and fiber photometry experiments.

- Jim Malone assisted with proteomic data analysis and visualization.

- Justin Woods assisted with the immunohistochemistry experiments.

- The proteomic experiments were performed at the Washington University Proteomics Shared Resource (WU-PSR), under the supervision and guidance of R Reid Townsend M.D., Ph.D., Director and Robert Sprung, PhD., Co-Director.

- Ream Al-Hasani, PhD conceptualized the study and supervised the experiments.

Acknowledgments

The figure schematics were created with biorender.com. We would like to thank Justin W. Baldwin for his guidance on appropriate data visualization tools and for his support in proofreading the manuscript. We would also like to thank Dr. Jordan McCall for his advice on data analysis and visualization as well as proofreading the manuscript. Thank you to Emily Wheeler for proofreading the manuscript and providing feedback on sentence structure and flow. We would also like to thank the entirety of the proteomics core team including Dr. Yiling Mi for her assistance. Thank you to the entirety of the Al-Hasani and McCall labs for their feedback on

experimental design and analysis. Finally, thank you to Dr. Tim Holy, Dr. Robyn Klein, and Dr. Alexxai Kravitz

for their feedback.

References

[1]   L.D. Fricker, E.B. Margolis, I. Gomes, L.A. Devi, Five Decades of Research on Opioid Peptides: Current Knowledge and Unanswered Questions, Mol Pharmacol. 98 (2020) 96–108. https://doi.org/10.1124/MOL.120.119388.

[2]   M.R. Banghart, X.J. He, B.L. Sabatini, A Caged Enkephalin Optimized for Simultaneously Probing Mu and Delta Opioid Receptors, ACS Chem Neurosci. 9 (2018) 684–690. https://doi.org/10.1021/ACSCHEMNEURO.7B00485/ASSET/IMAGES/LARGE/CN-2017-00485A_0003.JPEG.

[3]   I. Gomes, S. Sierra, L. Lueptow, A. Gupta, S. Gouty, E.B. Margolis, B.M. Cox, L.A. Devi, Biased signaling by endogenous opioid peptides, Proceedings of the National Academy of Sciences. 117 (2020) 11820–11828. https://doi.org/10.1073/pnas.2000712117.

[4]   A. Mansour, M.T. Hoversten, L.P. Taylor, S.J. Watson, H. Akil, The cloned μ, δ and κ receptors and their endogenous ligands: Evidence for two opioid peptide recognition cores, Brain Res. 700 (1995) 89–98. https://doi.org/10.1016/0006-8993(95)00928-J.

[5]   M.R. Banghart, S.Q. Neufeld, N.C. Wong, B.L. Sabatini, Enkephalin Disinhibits Mu Opioid Receptor-Rich Striatal Patches via Delta Opioid Receptors, Neuron. 88 (2015) 1227–1239. https://doi.org/10.1016/J.NEURON.2015.11.010.

[6]   N.T. Maidment, D.R. Brumbaugh, V.D. Rudolph, E. Erdelyi, C.J. Evans, Microdialysis of extracellular endogenous opioid peptides from rat brain in vivo, Neuroscience. 33 (1989) 549–557. https://doi.org/10.1016/0306-4522(89)90407-7.

[7]   P.W. Marinelli, L. Bai, R. Quirion, C. Gianoulakis, A Microdialysis Profile of Met-Enkephalin Release in the Rat Nucleus Accumbens Following Alcohol Administration, Alcohol Clin Exp Res. 29 (2005) 1821–1828. https://doi.org/10.1097/01.alc.0000183008.62955.2e.

[8]   L. Hemandez, B.G. Hoebel, Food reward and cocaine increase extracellular dopamine in the nucleus accumbens as measured by microdialysis, 1988.

[9]   M.M. Nieto, J. Wilson, A. Cupo, B.P. Roques, F. Noble, Chronic Morphine Treatment Modulates the Extracellular Levels of Endogenous Enkephalins in Rat Brain Structures Involved in Opiate Dependence: A Microdialysis Study, The Journal of Neuroscience. 22 (2002) 1034–1041. https://doi.org/10.1523/JNEUROSCI.22-03-01034.2002.

[10]   O.S. Mabrouk, Q. Li, P. Song, R.T. Kennedy, Microdialysis and Mass Spectrometric Monitoring of Dopamine and Enkephalins in the Globus Pallidus Reveal Reciprocal Interactions that Regulate Movement, J Neurochem. 118 (2011) 24. https://doi.org/10.1111/J.1471-4159.2011.07293.X.

[11]   Q. Li, Jon-Kar Zubieta, R.T. Kennedy, Practical Aspects of in Vivo Detection of Neuropeptides by Microdialysis Coupled Off-Line to Capillary LC with Multi-stage MS, Anal Chem. 81 (2009) 2242. https://doi.org/10.1021/AC802391B.

[12]   H.M. Baseski, C.J. Watson, N.A. Cellar, J.G. Shackman, R.T. Kennedy, Capillary liquid chromatography with MS3 for the determination of enkephalins in microdialysis samples from the striatum of anesthetized and freely-moving rats, Journal of Mass Spectrometry. 40 (2005) 146–153. https://doi.org/10.1002/jms.733.

[13]   R. Al-Hasani, J.-M.T. Wong, O.S. Mabrouk, J.G. McCall, G.P. Schmitz, K.A. Porter-Stransky, B.J. Aragona, R.T. Kennedy, M.R. Bruchas, In vivo detection of optically-evoked opioid peptide release, Elife. 7 (2018). https://doi.org/10.7554/eLife.36520.

[14]   R. Al-Hasani, J.G. McCall, G. Shin, A.M. Gomez, G.P. Schmitz, J.M. Bernardi, C.-O. Pyo, S. il Park, C.M. Marcinkiewcz, N.A. Crowley, M.J. Krashes, B.B. Lowell, T.L. Kash, J.A. Rogers, M.R. Bruchas, Distinct Subpopulations of Nucleus Accumbens Dynorphin Neurons Drive Aversion and Reward., Neuron. 87 (2015) 1063–77. https://doi.org/10.1016/j.neuron.2015.08.019.

[15]   B.E. Pirino, M.B. Spodnick, A.T. Gargiulo, G.R. Curtis, J.R. Barson, A.N. Karkhanis, Kappa-opioid receptor-dependent changes in dopamine and anxiety-like or approach-avoidance

behavior occur differentially across the nucleus accumbens shell rostro-caudal axis, Neuropharmacology. 181 (2020). https://doi.org/10.1016/J.NEUROPHARM.2020.108341.

[16] B.H. Trieu, B.C. Remmers, C. Toddes, D.D. Brandner, E.M. Lefevre, A. Kocharian, C.L. Retzlaff, R.M. Dick, M.A. Mashal, E.A. Gauthier, W. Xie, Y. Zhang, S.S. More, P.E. Rothwell, Angiotensin-converting enzyme gates brain circuit–specific plasticity via an endogenous opioid, Science (1979). 375 (2022) 1177–1182. https://doi.org/10.1126/SCIENCE.ABL5130/SUPPL_FILE/SCIENCE.ABL5130_DATA_S1.ZIP.

[17] S.M. Reynolds, K.C. Berridge, Positive and negative motivation in nucleus accumbens shell: bivalent rostrocaudal gradients for GABA-elicited eating, taste "liking"/"disliking" reactions, place preference/avoidance, and fear, J Neurosci. 22 (2002) 7308–7320. https://doi.org/10.1523/JNEUROSCI.22-16-07308.2002.

[18] D.C. Castro, M.R. Bruchas, A Motivational and Neuropeptidergic Hub: Anatomical and Functional Diversity within the Nucleus Accumbens Shell, Neuron. 102 (2019) 529–552. https://doi.org/10.1016/J.NEURON.2019.03.003.

[19] A.G. Schindler, D.I. Messinger, J.S. Smith, H. Shankar, R.M. Gustin, S.S. Schattauer, J.C. Lemos, N.W. Chavkin, C.E. Hagan, J.F. Neumaier, C. Chavkin, Stress produces aversion and potentiates cocaine reward by releasing endogenous dynorphins in the ventral striatum to locally stimulate serotonin reuptake, J Neurosci. 32 (2012) 17582–17596. https://doi.org/10.1523/JNEUROSCI.3220-12.2012.

[20] D.C. Castro, K.C. Berridge, Opioid Hedonic Hotspot in Nucleus Accumbens Shell: Mu, Delta, and Kappa Maps for Enhancement of Sweetness "Liking" and "Wanting," Journal of Neuroscience. 34 (2014) 4239–4250. https://doi.org/10.1523/JNEUROSCI.4458-13.2014.

[21] J.J. Pesavento, B.A. Garcia, J.A. Streeky, N.L. Kelleher, C.A. Mizzen, Mild performic acid oxidation enhances chromatographic and top down mass spectrometric analyses of histones, Mol Cell Proteomics. 6 (2007) 1510–1526. https://doi.org/10.1074/MCP.M600404-MCP200.

[22] A.G. Difeliceantonio, O.S. Mabrouk, R.T. Kennedy, K.C. Berridge, Enkephalin surges in dorsal neostriatum as a signal to eat, Current Biology. 22 (2012) 1918–1924. https://doi.org/10.1016/J.CUB.2012.08.014.

[23] T.L. Ripley, J. Jaworski, P.K. Randall, R.A. Gonzales, Repeated perfusion with elevated potassium in in vivo microdialysis - A method for detecting small changes in extracellular dopamine, J Neurosci Methods. 78 (1997) 7–14. https://doi.org/10.1016/S0165-0270(97)00129-5.

[24] M. Comb, P.H. Seeburg, J. Adelman, L. Eiden, E. Herbert, Primary structure of the human Met- And Leu-enkephalin precursor and its mRNA, Nature. 295 (1982) 663–666. https://doi.org/10.1038/295663a0.

[25] J. HUGHES, H.W. KOSTERLITZ, T.W. SMITH, THE DISTRIBUTION OF METHIONINE-ENKEPHALIN AND LEUCINE-ENKEPHALIN IN THE BRAIN AND PERIPHERAL TISSUES, Br J Pharmacol. 120 (1997) 428–436. https://doi.org/10.1111/j.1476-5381.1997.tb06829.x.

[26] G. Henderson, J. Hughes, H.W. Kosterlitz, In vitro release of Leu- and Met-enkephalin from the corpus striatum, Nature 1978 271:5646. 271 (1978) 677–679. https://doi.org/10.1038/271677a0.

[27] T. Yoshimasa, K. Nakao, H. Ohtsuki, S. Li, H. Imura, Methionine-enkephalin and leucine-enkephalin in human sympathoadrenal system and pheochromocytoma, Journal of Clinical Investigation. 69 (1982) 643–650. https://doi.org/10.1172/JCI110491.

[28] J.A. Mansi, S. Laforest, G. Drolet, Effect of stress exposure on the activation pattern of enkephalin-containing perikarya in the rat ventral medulla, J Neurochem. 74 (2000) 2568–2575. https://doi.org/10.1046/J.1471-4159.2000.0742568.X.

[29] E.C. Dumont, R. Kinkead, J.F. Trottier, I. Gosselin, G. Drolet, Effect of chronic psychogenic stress exposure on enkephalin neuronal activity and expression in the rat hypothalamic paraventricular nucleus, J Neurochem. 75 (2000) 2200–2211. https://doi.org/10.1046/J.1471-4159.2000.0752200.X.

[30] S. Ceccatelli, C. Orazzo, Effect of different types of stressors on peptide messenger ribonucleic acids in the hypothalamic paraventricular nucleus, Acta Endocrinol (Copenh). 128 (1993) 485–492. https://doi.org/10.1530/ACTA.0.1280485.

[31] R.J. Valentino, E. van Bockstaele, Endogenous opioids: The downside of opposing stress, Neurobiol Stress. 1 (2015) 23–32. https://doi.org/10.1016/J.YNSTR.2014.09.006.

[32] H. Nam, R. Chandra, T.C. Francis, C. Dias, J.F. Cheer, M.K. Lobo, Reduced nucleus accumbens enkephalins underlie vulnerability to social defeat stress, Neuropsychopharmacology. 44 (2019) 1876–1885. https://doi.org/10.1038/s41386-019-0422-8.

[33] J.W. de Jong, S.A. Afjei, I. Pollak Dorocic, J.R. Peck, C. Liu, C.K. Kim, L. Tian, K. Deisseroth, S. Lammel, A Neural Circuit Mechanism for Encoding Aversive Stimuli in the Mesolimbic Dopamine System, Neuron. 101 (2019) 133-151.e7. https://doi.org/10.1016/J.NEURON.2018.11.005.

[34] A.L.O. Hebb, R.M. Zacharko, M. Gauthier, F. Trudel, S. Laforest, G. Drolet, Brief exposure to predator odor and resultant anxiety enhances mesocorticolimbic activity and enkephalin expression in CD-1 mice, European Journal of Neuroscience. 20 (2004) 2415–2429. https://doi.org/10.1111/j.1460-9568.2004.03704.x.

[35] A.A. Legaria, B.A. Matikainen-Ankney, B. Yang, B. Ahanonu, J.A. Licholai, J.G. Parker, A. v. Kravitz, Fiber photometry in striatum reflects primarily nonsomatic changes in calcium, Nature Neuroscience 2022 25:9. 25 (2022) 1124–1128. https://doi.org/10.1038/s41593-022-01152-z.

[36] P. Bérubé, J.F. Poulin, S. Laforest, G. Drolet, Enkephalin Knockdown in the Basolateral Amygdala Reproduces Vulnerable Anxiety-Like Responses to Chronic Unpredictable Stress, Neuropsychopharmacology 2014 39:5. 39 (2013) 1159–1168. https://doi.org/10.1038/npp.2013.316.

[37] P. Bérubé, S. Laforest, S. Bhatnagar, G. Drolet, Enkephalin and dynorphin mRNA expression are associated with resilience or vulnerability to chronic social defeat stress, Physiol Behav. 122 (2013) 237–245. https://doi.org/10.1016/J.PHYSBEH.2013.04.009.

[38] I. Melo, E. Drews, A. Zimmer, A. Bilkei-Gorzo, Enkephalin knockout male mice are resistant to chronic mild stress, Genes Brain Behav. 13 (2014) 550–558. https://doi.org/10.1111/GBB.12139.

[39] A.L.O. Hebb, R.M. Zacharko, M. Gauthier, F. Trudel, S. Laforest, G. Drolet, Brief exposure to predator odor and resultant anxiety enhances mesocorticolimbic activity and enkephalin expression in CD-1 mice, European Journal of Neuroscience. 20 (2004) 2415–2429. https://doi.org/10.1111/j.1460-9568.2004.03704.x.

[40] V.N. Sherathiya, M.D. Schaid, J.L. Seiler, G.C. Lopez, T.N. Lerner, GuPPy, a Python toolbox for the analysis of fiber photometry data, Scientific Reports 2021 11:1. 11 (2021) 1–9. https://doi.org/10.1038/s41598-021-03626-9.