Title: Growth enhances colors, shapes and biological functions of Turing color patterns of Eukaryotes

Author: Pierre Galipot[1,2]*

Affiliations:

[1]Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles; CP50, 75005 Paris, France.

[2]Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Tokyo, Japan

*Corresponding author. Email: pierregalipotpro@gmail.com

Abstract:

Evidenced in zebrafishes and *Mimulus*, Turing-like mechanisms are probably responsible for many periodic color patterns of Eukaryotes. Nevertheless, patterns such as leopard iconic rosettes required additional ingredients to explain their formation. Growth being the main candidate, we extensively explore its multiple facets, at the Eukaryotes scale. We show that putative-growth Turing color patterns are present in many diverse lineages. Using models, we show the many ways growth can induce new shapes and colors, and that putative-growth pattern locations correlates with tissue hot spots of growth, suggesting the latter as the underlying mechanism. By reverse reasoning, we show that growth effects could reveal crucial information about pattern formation. Finally, we show how putative growth patterns can contribute to influence organisms visibility, thereby improving camouflage or aposematism.

One-Sentence Summary: Growth-putative Turing color patterns are diverse, adorn thousands of Eukaryotic species and enhance their visual effects.

Repeated color patterns are known to allow countless ecological roles such as be more visible to congeners, for partner recognition (*1*) or communication (*2*), or to other species, in an attractively manner for pollination (*3*) or repellingly for aposematism (*4*). Repeated color patterns may also avoid detection or recognition, in camouflage (*5*).

Turing-like systems are composed of a couple of activator and inhibitor in their simplest molecular form (*6*). After years of mathematical formalization (*7*) and modeling (*8*) they have been described *in vivo* (*9*) and are responsible for many biological repeated structures called Turing patterns. Characterized by their periodicity and the range of repeated motifs shapes (spots, stripes, mazes and not much more (*10*), they form for example the vertebrate digits (*11*), and fingerprints (*12*) or the xylem openwork patterns of plants (*13*). For color patterns, their cellular and molecular bases have been unveiled in zebrafishes (*14*) and recently in plants of the

genus *Mimulus* (*15*), while currently being explored in more and more other eukaryotic taxa, like the Felidae (*16*).

The intriguing and iconic leopard (*Panthera pardus*) pattern, characterized by brown rosettes with an orange heart on a light-yellow background, is not under the scope of a classic Turing pattern (*10*). Consequently, the leopard color pattern has been the focus of attempts to explain its formation *(14, 15)*, with recent models impressive in both their realism and their biological plausibility (*19*). Those models perfectly reproduced leopard rosettes by adding the most probable missing ingredient: tissue growth during pattern formation. But what about beyond leopard famous color pattern? Is there a biodiversity of putative-growth Turing color patterns (hereafter called PGTCPs), and what are their qualitative and quantitative characteristics?

To answer this question, we sought to measure in the broadest possible way the importance that growth could have in the formation and the diversity of the periodic color patterns in the Eukaryotes, beyond the biological curiosity of this feline coat.

This study is based on the observation of color and shapes of color patterns, which we then relate to processes. Thanks to the specificities of patterns produced by Turing-like systems (periodicity and motif diversity: spots, stripes, mazes), we make the assumption that we can predict the nature of the underlying mechanism from the characteristics of the pattern. For a reason of accuracy, we therefore refer to *putative classic Turing-like color patterns,* sometimes simplified as *'classic'* and *putative growth Turing-like color patterns*, abbreviated as PGTCPs. 'Turing-like' refers to the fact that Turing systems can be of many biological scales and natures (*10*).

Five research themes were addressed in this study: i) What are the specific geometric and color diversities of PGTCPs and what is their biodiversity in Eukaryotes? ii) By which ways growth can influence the final shapes and colors of Turing patterns and how can it be modeled? iii) Since no functional study has so far directly proven the involvement of growth in PGTCPs, what clues are pointing to it? iv) Can PGTCPs phenotypes be geometrically interpreted and give important information on the Turing system itself and/or on the processes underwent by the tissues during morphogenesis? Finally, v) could PGTCPs improve or provide new biological functions compared to putative classic Turing-like color patterns?

We then discussed the factors controlling the potential rarity of PGTCPs observed in some taxa, whether it is real (linked to ecological, evolutionary or mechanical reasons) or artificial (linked to observation bias).

Results

Classic and putative growth Turing-like color patterns (PGTCPs) are common and diversified in Angiosperms and Animalia but seem absent in Fungi and other lineages of Eukaryotes

While the leopard has been for a long time the only species studied for its PGTCP, some other animal species have been proposed as candidates during the last years, such as the whale shark (*20*), the ground squirrel *Ictidomys tridecemlineatus*, or some species of *Dendrobates* frogs (*19*). In order to provide a most comprehensive picture possible of the presence of these motifs, we have extended our research to all Eukaryotes. A first large-scale screening showed that PGTCPs seem absent from certain lineages of multicellular Eukaryotes, in particular in Plantae other than Angiosperms (probably due to the absence of biotic pollination), as well as in Fungi (possibly due to the presence of a pre-existing mechanism of repeated pattern formation by physical cracking (*21*)) and to unicellular lineages of Eukaryotes (where visual signals might be less relevant at the microscopic scale). We then restricted our study to Metazoans and Angiosperms, trying to find from multiple sources at least one example of a species bearing PGTCPs for each of the 620 orders of animals and 422 families (or suitable taxonomic rank when family did not exist) of Angiosperms. The PGTCPs were classified in 4 categories, which could be combined: line-and-dot alternation, mixed colors, rosettes (and mazes of mazes) and intermediate bands (Fig. 1.A). The complete results are available in a database accessible at the following URL (www.eukolorpatterns@cnrs.fr).

Figures 1.B and 1.E show the mapping of the different types of patterns (classic or PGTCPs) onto the phylogenetic trees of animals and Angiosperms. In the former, PGTCPs were found in 62 orders (10% of the total), while in 268 orders (43%) only classic patterns with or without suspicion of PGTCPs were found. The remaining 290 orders (47%) did not seem to show any Turing-like pattern, whether classic or PGTCPs.

For Angiosperms, where to our knowledge no species had been explicitly described with PGTCPs, we found 38 families with at least one species displaying PGTCPs, 22 families with potential PGTCPs, plus 5 families with only classic Turing patterns, 33 families with potential classic Turing patterns and finally 339 families for which no species displaying any putative Turing color pattern was found.

In both Angiosperms and animals, PGTCPs have been found in widely separated orders and families, suggesting several independent evolutions. Some subgroups such as the Lamiales or Actinopterygii appear to harbor a high proportion of PGTCPs; by contrast, some taxa with high species diversity and high coloration diversity seem to harbor fewer PGTCPs than expected, such as insects or birds. We proposed possible ecological, evolutionary and/or biological constraints explanations for these unexpected scarcities in the Discussion part and in Table 1.

Finally, we carried out a quasi-exhaustive species-scale survey of color patterns in two groups that are iconic in terms of color patterns and where species are well described in the literature: mammals (6649 species, Fig. 1.C) and orchids (29573 species, Fig. 1.D). Among the former, 10% of the species were found to clearly display putative Turing-like patterns (541 species) or PGTCPs (73 reliable species, 79 potential). Groups with relatively few species such as Feliformia (to which the leopard belongs) have a large proportion of species bearing PGTCPs,

while others with a high specific richness exhibit few or no PGTCPs, such as the bats (Chiroptera).

In orchids, where only PGTCPs have been searched for, 525 species (1.8%) clearly display PGTCPs, 528 others are potential candidates. Again, while many genera possess at least one species with a PGTCP, some contribute more both in proportion and in absolute value, such as the genus *Stanhopea*, of which 33 of the 81 species (41%) display conspicuous PGTCPs.

This first overview of PGTCPs in Eukaryotes suggests a very high diversity, which deserves to be the subject of a greater attention in the future and raises the question of the selective forces that may have driven the evolution of PGTCPs.

Disruption induced by tissue growth can generate new shapes like rosettes, lines-and-dots alternation or intermediate bands, together with new colors by mixing and memory mechanisms in Turing patterns

The iconic pattern of the leopard has been the focus of much attention over the last twenty years, where several mechanisms of rosette formation have been proposed *(15, 16)*. The current consensus is that a rosette is an old dot reorganized into smaller dots in response to the increase in tissue size (Fig. 2.A and 2.B). Because Turing systems are sensitive to scale, a motif extension drive by tissue growth will be out of its equilibrium and cause its rearrangement. PGTCPs are then the more or less transient states produced during a return toward the equilibrium. The second particularity of leopard rosette, *i.e.*, the presence of orange coloration in its center, intermediate between dark brown and light yellow background, has been recently reproduced by de Gomensoro Malheiros et al (*19*) by growth combined to a memory mechanism where each particular location of the final pattern is the average of the all the previous states throughout the pattern formation. Their simulations perfectly reproduce natural leopard patterns. For the leopard rosettes or also the emu intermediary bands, we postulate that melanocytes are regularly produced in the skin tissues, becoming black (eumelanin) or yellow (pheomelanin) melanocytes depending on the current local state of Turing system, and at the end, tissues regions where Turing state knew the two states through time are composed of a mosaic of black and yellow melanocytes. This leads to intermediary color at the macroscopic scale (by our eye interpretation), in this case the orangish hair of leopard and light-brown feathers of the emu.

As we saw, leopard pattern combines both novel shapes and colors, two possible effects of growth. Nevertheless, some species can exhibit rosettes without intermediary color like the orchid *P. hieroglyphica,* which on the other hand has stretched rosettes. We reproduced them by simulated anisotropic growth without memory (Fig. 2.A, up and 2.B, right).

To better classify growth effects and PGTCPs, we suggest that rosettes belong to a bigger family of disrupted patterns. Considering the three classic Turing motifs shapes (dots, stripes and mazes), rosettes are then disrupted spots, intermediate bands (like the emu ones) and stripes-and-spots alternations are disrupted stripes, and nested patterns of some pufferfishes like *Tetraodon mbu* are mazes disrupted into mazes of mazes (*22*).

Despite the efficiency of this memory model, it is *a priori* not compatible with all biological situations including some systems now well described, such as the zebrafish (*Danio rerio*) stripes

formation. In fact, although we observe intermediate colors in a large number of Actinopterygii fish species, the cellular Turing mechanisms described in zebrafishes are not compatible with the conception of additive coloration through memory of earlier stages proposed for the leopard, as these fish colored cells constantly rearrange *(9, 19)*. We therefore propose a new type of mechanism that could fit the current biological knowledge of zebrafishes stripe formation, without superimposition of several color states during tissue growth, which we called "non-equilibrium intermediate coloration".

We considered that the intermediate colors could be produced by intermediate values of the Turing system parameters (typically activator or inhibitor medium-level concentrations). When tissue growth disrupts an already formed pattern, for example stripes, new ones are generated from the middle of the pre-existing ones. At the equilibrium, new sharp frontiers will be generated, but during their formation, intermediate values may be reached over a sufficiently long period and over a large width to be observed, (Figure 2.C, 1^{st} row). This implies that if colors are produced in real time, species will display intermediate coloration during their growth phase and intermediate color will be transiently visible (e.g., *Pomacanthus imperator* juvenile pattern, Fig 2.D). Furthermore, if the pattern dynamics are stopped before reaching its equilibrium, intermediary colors can become definitive by being snap-shot. Indeed, many natural PGTCPs could be 'out of equilibrium' patterns frozen in time by the biological readout of the Turing system state at a given time.

This doublet of mechanisms may also explain why intermediate colors are rare in Angiosperms, even though rosettes are very common, which show that growth can effectively disrupt many plant color patterns. According to the pioneer study by Ding *et al* (*15*) on the flowers of *Mimulus*, their color mechanism is based on a reaction-diffusion involving two proteins. The preservation of memory seems more complicated (although not impossible) to imagine this at the molecular level (*23*), given the short half-life of molecules. Nevertheless, some flowers like those of the Lamiaceae *Scutellaria rubropunctata* exhibit striking leopard-like purple rosettes, with intermediary colors on their middle, showing that is it not totally impossible to produce them in Angiosperms (*24*).

The last category, lines-and-dots alternation is a bit different in terms of mechanics.

Less common than other PGTCPs, it is harbored by very distantly-related species like *Stapelia gigantea* on its flowers, the squirrel *Ictidomys tridecemlineatus* on its fur (Fig. 1.D) or the whale shark *Rhincodon typus* on its skin.These species generate dots or sometimes mazes in the middle of bands. Several models have been proposed to reproduce them, some related to biological phenomena such as neural crest migration (*19*). Our proposition is based on both internal and external factors, respectively random noise of the system and accelerating growth of the tissue.

A stripe as initial condition (1^{st} generation stripe, Fig. 1C, 2^{nd} row, t=0 and t=1000) generates a multi-striped pattern (2^{nd} generation stripes, Fig. 1C, 2^{nd} row, t=5000 and t=22000) in a small tissue. If a higher growth rate follows, eventually helped by a highly noisy Turing system, this could disrupt the formation of 3^{rd} generation stripes that will break into aligned dots (Figure 1.C, 2^{nd} row, t=22700), leading to alternations of stripes and dots. We can note that like all PGTCPs, these patterns are only meso-stable, if the dynamics of the system continues to be active, the fate of all stripes is to break into dots (Fig. 1.C, 2^{nd} row, t=10000, biologically observe in whale

sharks) unless a forcing mechanism maintains them or snapshot the pattern. Lines-and-dots patterns are a family containing variations, if growth is very high, more complex patterns than aligned dots can form (*19*).

In summary, all the types of PGTCPs (rosettes, intermediate colorations, line-and-dots alternation, intermediate bands and all their variations) could be view as the result of a triad: i) tissue growth, ii) scale sensibility of Turing-like systems and iii) system memory, as Turing systems dynamically rearrange new patterns from the previous ones as new initial conditions.

While a definitive functional proof of the implication of growth is still lacking, we then gather a cluster of clues suggesting that it is the main candidate for the great diversity of PGTCPs in Eukaryotes.

The distribution of PGTCPs occurrences suggests a universal mechanism, and morphometric analyses show that their presence correlates with the most-growing areas of the tissues

The parameter(s) conditioning the presence or absence of PGTCPs are undoubtedly common and universal, as evidenced by the examples of presence/absence found at all scales of life starting with the individual scale, in a spatial and temporal manner (Fig. 3.A). Indeed, in most species with PGTCPs on their external tissues, we noticed that not all the tissue is covered by PGTCPs. At the temporal scale, the fish *Pomacanthus imperator* has only intermediate bands during the transition from juvenile to adult morphs (Figure 3.A). At the intraspecific scale, we documented individuals belonging to the same species, some of them displaying these patterns and others not, for instance in zebras (*Equus quagga*, Fig. 3.A). Finally, the large distribution in the phylogenetic tree of these motifs presented in Fig. 1.B and 1.E shows that if a common mechanism governs all these motifs, it must be shared by a large amount of very diverse species. All these examples suggest the existence of a widespread mechanism underlying the diversity of patterns, and growth fits this requirement.

To strongly suggest its role in PGTCPs formation, we estimated through morphometric studies the tissue areas undergoing the greatest cumulative expansion during the development of organisms whose developmental sequence was available (Fig. 3.B and 3.C, left), and compared them to the map of presence or absence of growth patterns (Fig. 3.C, middle and right). In the Burchell's zebra (*Equus quagga burchellii),* despite the approximation limitations of this method, areas where growth is more intense predict rather satisfactorily the areas of presence of intermediate bands, *i.e.,* on the trunk and on the proximal part of the limbs, with an absence of patterns on the head, neck and distal part of the limbs (Figure 3.C). It is also supported by the fact that in most mammals, PGTCPs are preferentially located on the trunk and absent from the head, which is in agreement with the fact that the head undergoes much less growth than the body during the embryonic development (*25*).

Effects of growth on Turing patterns can be used as reverse tools to extract information on pattern timing and system characteristics such as wavelength

The effects of growth on Turing patterns reported in this article are reminiscent of earlier stages: rosettes are ancient dots, mixed colors are produced in ancient dark or light zone, and so on.

This led us to ask if this could be exploited to obtain information on earlier stages, in particular the crucial timing of pattern formation, which is invisible to the experimenter in many species, because very often production of colors occurs much later (*26*). Such data could increase both the raw knowledge of Turing pattern systems together with allowing comparative analyses, but also it could facilitate the implementation of functional experiments by allowing to focus on a reduced temporal window.

Periodicity is the crucial characteristic of Turing systems for the reverse method we have developed (presented in Fig. 4.A). Indeed, by assuming that for a given organism i) the whole pattern is set up at the same time in all points of the tissue and that ii) the parameters of the system are homogeneous in space and time (meaning that periodicity is respected in all points of the tissue), we could deduce that at the time of the formation of the pattern, all the motifs must be regularly spaced on the tissue, and have the same size. However, since for a large part of the species this is not what is observed on the final pattern, the cause of this discrepancy is the differences in the relative growth between the tissue parts during development, from the stage of the end of pattern production to the moment of observation.

By extension, concerning growth patterns such as rosettes, if we consider either a rosette as a single motif, or several motifs (by counting how many sub-motifs it is divided into), it could theoretically be possible to calculate both the timing of the initial dots pattern, and the pattern of rosette formation.

To test this hypothesis, we use the cat (*Felis silvestris*) and leopard models, the first having a well described embryogenesis for which morphometric measurements are applicable, the second for its PGTCPs, while being morphologically close to the first at least until its juvenile stages (Fig. 4.B). For leopard dots, predicted timings are rather scattered, but all estimations point to late timings, post stage 22. This dispersion may have one or several non-mutually exclusive causes such as: approximation of landmark distances and positions, approximation in the motif counting, differences in external morphology between cat and leopard, non-trivial relationship between number of motifs and tissue size or finally, different timing of pattern formation depending on the tissue location. If we focus on the two distances that undergo the greatest increase in relative size with respect to the head (trunk and tail), which *a priori* are the two most reliable values for estimation, they suggest stage 22 as the stage of ending of phase 1 of patterning (formation of the 1st generation of dots, which will give the rosettes). Furthermore, when we look at the estimates of the 2nd generation patterns, it points to much more recent stages (Figure 4.B), juvenile or even adult, which is coherent with the observations of Liu et al. (*18*), which show that the pattern continues to be set up after the juvenile stages.

To improve predictions, we simulated the formation of rosette patterns at different growth rates (Fig. 4.C, solid lines), and compared to the theoretical number of motifs at each step (Fig. 4.C, dashed lines), i.e., at an equivalent size, but without growth.

For all simulations, whatever the growth rate, we observe a discrepancy between the number of patterns expected and the number of patterns observed, which can be initially attributed to the inertia of the system (the time taken to respond to the perturbation). Nevertheless, if we let the system run long enough after the end of growth, we observe very clearly that the final number of patterns is lower than the expected one, by about 15% for all growth rates in our example. This means that when the Turing system undergoes a first pattern generation, then responds to growth, it imposes constraints which prevent the system from reaching its maximum number of motifs at the equilibrium.

This shows that this relationship between growth rate and motif number is not totally trivial and must be used with great caution in our predictions. However, several things could counterbalance this. First, it should be noted that in cases where the growth rate is too high, the system does not have time to produce growth patterns. This means that if growth patterns are observed, the growth rate is necessarily reduced, and therefore the shift with the expected number of patterns should be lower.

In our models, the "delta" was between 16 and 20%, for growth rates low enough for growth patterns production. In order to determine if these values were verified for other conditions, we carried out other simulations with higher final size values. We then obtained 'delta' values all between 10 and 20% (not shown). Without knowing what the relationship is between tissue growth rate and response rate of biological systems, we therefore propose to use a correction of +15%, +/-5% to the number of motifs counted in presence of rosettes, which should bring the estimated value closer to the theoretical value and improve the accuracy of our reverse method.

Finally, we develop a method to better follow and estimate tissue growth thanks to artificial landmarks. Some plant tissues like sepal outer epidermis are accessible during their development and therefore available for marking. We chose to simply apply spot landmarks with waterproof ink. This relatively non-invasive technique allows to estimate tissue deformations and so local growth through developmental time. By tracking the relative position of ink points using 2D photography, or *in vivo* photogrammetry (Fig. 4D, lower panel) in the case of 3-dimensional folded tissues, geometric morphometry techniques can be applied.

Although imperfect, these methods should be able to estimate the timing of production of classical Turing and growth patterns on many species and allow large-scale comparative studies (Fig. 4.D). Moreover, as precise patterning timings will be revealed by functional experiments, the validity of the prediction of this method could be verified and improved. Finally, while the proxy used here is the number of motifs, a study assisted by deep learning could help find finer estimators (*27*), which would therefore improve the prediction of the timing, and its corollary: the natural wavelength of the pattern.

PGTCPs could have evolved under selection for enhancing or reducing pattern visibility

Resulting from the interaction between a Turing system and tissue growth contingent to the development, PGTCPs could appear as only by-products of growth and not under the scope of an efficient selection. However, we have previously documented numerous examples of

presence/absence of growth patterns, within and among species, suggesting instead possible ways of control and selection.

Like color patterns in general, classic Turing-like color patterns are perceived by con- and hetero-specifics, and could therefore be under selection for signaling or camouflage (*28*). We therefore wondered whether the geometric and color features of the PGTCPs could improve signaling or camouflage compared to classical Turing patterns.

Concerning camouflage, the mixed colorations caused by growth could allow the species displaying them to retain the camouflage capacities of Turing patterns repetition, while reducing the segregation of colors that is intrinsically linked to them. Indeed, a classic Turing system produces a segregation of colors, not very compatible with the production of intermediate colors, which are likely to be useful in camouflage. For example, the shades of brown resulting from the mixture of pigment cells in the eumelanin (black) / pheomelanin (yellow/orange) system are the basis of many camouflages, especially in mammals and birds (22, 23). If dark and light melanocytes are separated by the Turing system, the resulting patterns are often very contrasting, as they lose the potential for color shades. This is illustrated by the many examples of mammalian species with black and light Turing patterns from a genus where their closely related species have intermediate-colored coats (Fig. 5.A). In some specific cases, the effects of growth could then be beneficial in terms of camouflage and be positively selected during evolution. For example, the geometrical particularities of some growth patterns such as leopard rosettes could further improve the camouflage efficiency compared to a classical spotted pattern, by adding highest motif variability, novel shapes and a more fractal pattern (Fig. 5.B, (24, 25)). We observe that all the Ratites juveniles possess orange and brownish colors but for each of these four species, the origin and nature of these intermediate colors seem different: Rhea and emu (*Dromaius novaehollandiae*) exhibit orange intermediate bands, i.e., PGTCPs (Fig. 5.B, left and right), the juvenile ostrich (*Struthio camelus*) has orange colors on its head and neck, probably due to a particular type of pheomelanin (*33*) as for the cassowary *(Cassuarius sp.)*, and also has very large feathers (for a down coat) intermingled on its back, which have the effect of averaging and mixing the colors produced by the melanocytes in the skin. These convergent traits could be a sign that intermediate colors are under selection and have a specific biological function. In particular, we can hypothesize that they enhance their cryptic pattern, making it more difficult to detect by potential predators (Fig. 5.B, right) (*34*). Growth would thus be a mean to produce intermediate colors.

On the other side of the spectrum, some characteristics of growth patterns could enhance their visibility and their attractiveness or repulsion to conspecifics or other species, as in the case of aposematism. Indeed, the periodicity of Turing patterns makes them easily detectable by a large number of visual neural networks, and their attractiveness, including in the human species, is directly related to their low processing time for the brain (Fig. 5.C, (*35*)). We hypothesize that growth could be a non-negligible source of striking geometries, positively selected during evolution. Typically, the aposematic ring patterns of the greater blue-ringed octopus (*Hapalochlaena lunulata* (*36*)) and Dendrobatid frogs (*32*, *33*) line-and-dots patterns have all the characteristics to have been produced by growth (Fig. 5.D, up) and it can be hypothesized that these motifs improve pattern detection, compared to classical dotted patterns (Fig. 5.D). Contrary to periodic stripes or dots (*39*), ring patterns and line-and-dot alternations are at our knowledge

very rare or absent in abiotic environments, which might confer them a higher detectability by
vision systems.

More than mere by-products of developing tissue growth, we showed that growth patterns could
potentially enhance camouflage capabilities, and conversely improve pattern visibility in
situations of aposematism, interspecific attraction or sexual selection. Behavioral and
evolutionary studies would then provide an even more complete picture of the functional
possibilities of these particular patterns.

Discussion

Factors controlling the taxa-specific absence or sparsity of putative-growth Turing color
patterns (PGTCPs)

Although PGTCPs are widely distributed in the phylogeny tree of Eukaryotes, there are hotspots
(e.g., Chondrichthyes and Actinopterygii for Animals or Lamiids for Angiosperms) and quasi-
empty areas (e.g., Rosids for Angiosperms or Fungi (tree not shown)). As PGTCPs are the
combination of growth and an active Turing color system, the absence of the latter is a first
trivial reason of the absence of PGTCPs, and it can be spot in the phylogeny in the grey zones
(neither blue nor yellow in Figure 1). In contrast, there are some taxa with some or many
examples of species with classic periodic color patterns, but in which it was impossible or very
difficult to find examples of PGTCPs, typically birds where to our present knowledge, only one
species exhibits a PGTCP, the emu *Dromaius novaehollandiae*. As the explanations could be
very different from one taxon to another, we tried to propose several hypotheses, which are
summarized in Table 1. For plants, many families seem not to be adorned with putative-Turing-
like patterns (grey tips), but when they are, in most of the time PGTCPs are also found. There is
no clear clade regrouping numerous families with only classic Turing patterns and no PGTCPs.

We then detail here one example, the case of insects, perhaps the most informative in many ways
for the factors controlling the presence/absence of PGTCPs. Indeed, insects are the metazoan
class containing the largest number of species (estimated at more than 5 million species (*40*)) and
are extremely diverse in terms of color patterns, both for colors and shapes. We would expect to
find many examples of classical Turing patterns, and also of PGTCPs, which is not the case for
either. More surprisingly, many closely related taxa belonging like Insects to the Pancrustaceae,
exhibit numerous Turing color patterns and PGTCPs, Decapoda being one example. Last but not
least, Turing systems have been found in Insects, but producing other types of structures than
color patterns (such as Turing nanopatterns of coat corneae (*41*) or tarsal attachment patterns
(*42*)), showing the compatibility between the external tissues of insects and Turing systems.

So, to explain this apparently rarity of Turing color patterns and PGTCPs in insects, we invoked
three types of explanations, not mutually exclusive. First, many parts of insect bodies are
segmented and compartmented, such as the abdomen or wings. This causes a division of space
and could reduce the size of the tissue where a hypothetical Turing system must express. The
boundaries between compartments may be too disruptive for Turing system actors and prevent
them from forming a large-scale proper pattern. If Turing systems could theoretically be
produced at any scale, as evidenced by the nanopatterns mentioned above, this
compartmentalization may prevent large motifs to occur, and therefore causing an impossibility
to ensure a proper biological function (*e.g.,* too small and tight motifs may not be visible to inter

or intra-specific individuals). The fact that insects are on average small organisms goes in the same direction, as larger animals would be favored for the formation of proper Turing color patterns. This is supported by the fact that the few examples of PGTCPs that we found in insects were in relatively large body parts and relatively large species, typically wings of *Pseudopanthera macularia* (Lepidoptera), *Eupteryx decemnotata* (Hemiptera) *Xanthippus corallipes* (Orthoptera) and *Dynastes tityus* (Coleoptera). Second, because insect body is already segmented, it may favor other types of color pattern mechanisms and thus disfavor Turing patterning during evolution (for example, black and white stripes in some silkworm strains are simply generated by the expression of Toll ligand Spätzle3 gene in the anterior part of every segment (*43*), generating a bicolor pattern, but *de facto* leads to a multi-repeated pattern thanks to the presence of multiple body segments). Finally, the growth conditions of insect outer tissues may also be a hindrance in the formation of PGTCPs, as most outer tissues are renewed by molting (*44*). This discontinuous growth may favor i) equilibrium patterns that disfavor PGTCPs, ii) the resetting of the Turing system at each molt and the *de novo* production of a color pattern, with a little increase in tissue absolute size because its step-by-step nature, thus again disfavoring PGTCPs. To summarize, the optimal conditions for observing PGTCPs are a tissue which a large increase in its absolute size and a continuous growth, which is more or less the opposite of the insect growth modalities. We believe that this reasoning can be extended to other taxa and be the subject of further and more precise studies.

Hidden PGTCPs: spatial decoupling, blurring, or pixelating

From the Turing-like system to the final coloration phenotype many mechanisms may act as filters or transformers, also called "secondary mechanisms" (*21*) taking a rough molecular state (in case of classic Turing based on morphogens) to a complex color pattern, the one which is observed in the end. Information and complexity can be added through the process, or removed, or even translated into another nature. For this reason, if growth influence the Turing-like system, and produced hidden PGCTPs, these might not be kept until the end of the process. We propose here three types of mechanisms aborting PGTCPs (Fig. 6).

First, is the spatial decoupling between the structure initially producing the structure (e.g., the melanocytes in the epidermis or dermis of bird or mammal skin) and the structure bearing the colors (e.g., feather or hair). A completely disordered fur or down will erase the underlying pattern, and so potential Turing-like patterns and PGTCPs. Juvenile ostriches harbor a rather disheveled down on their dorsum, compared to their head and neck. As a result, while the plucked skin shows a repeated pattern (consisting of stripes and dots), it is only observable on the head and neck (Fig. 6B). But this is not the only mechanism that appears to be at work in the juvenile ostrich back, which also appears to involve what we call "pixelating".

This second mechanism is when the final-colored structure is discrete (like a feather) and characterized by a lower "resolution" than the structures producing the initial pattern (Fig 6C). It acts as a filter by mixing several contiguous system values, therefore total information is reduced, and possibly most of the pattern characteristics are lose. In fact, Juvenile ostriches present nearly submillimeter initial skin pattern resolution, but nevertheless it possesses very low density of feathers on the back, which are very large compared to the down of the neck and head.

In other words, they have a very low resolution compared to the putative people underlying melanocyte skin pattern. This could lead to the loss of the geometric signal at the plumage scale.

The third type of mechanism leading to pattern loss is called blurring (Fig. 6D), and we assume that it could be quite frequent in Angiosperms PGTCPs. In this case, PGTCPs could be set up, but several mechanisms like a high sensitivity to Turing morphogens leading to a high level of color production with a maximum plateau could erase the difference between "medium" zones (typically the center of a rosette) and "high" zones (the crown a rosette), resulting in a more classical spot pattern. On-Off Filters with sigmoidal kinetics like allosteric enzymes could also make invisible the effects of growth and PGTCPs.

Experimental bias that need to be addressed to obtain a more accurate picture of PGTCPs of Eukaryotes

By definition, PGTCPs are produced by growth and individuals undergo the most growing phases during embryonic and post-embryonic development. However, most of the biodiversity textbooks and atlas focus on adult external morphology. This lack of embryonic and juvenile data clearly causes an underestimation of PGTCPs occurrences. In this study we also have noticed several times that when male/female dimorphism existed, only male individuals were represented in some textbooks. This was typically the case of *Cervus elaphus* where spotted Turing pattern of some females and juveniles was nearly missed if there had been no subsequent check.

Last but not least, inter-individual color pattern variability is very high in many species, like the Lamiaceae *Scutellaria rubropunctata,* which exhibit on its petal a range of color patterns from uniform blue or white to classic putative Turing spot patterns and striking PGTCPs like classic or open rosettes depending on the strain (*24*). A most comprehensive view of PGTCPs is therefore necessarily through the examination of many individuals.

References and Notes

1. W. Himstedt, The Significance of Color Signals in Partner Recognition of the Newt Triturus alpestris. *Copeia*. **1979**, 40–43 (1979).

2. R. Futahashi, R. Kurita, H. Mano, T. Fukatsu, Redox alters yellow dragonflies into red. *Proc Natl Acad Sci U S A*. **109**, 12626–12631 (2012).

3. A. S. Leonard, D. R. Papaj, 'X' marks the spot: The possible benefits of nectar guides to bees and plants. *Functional Ecology*. **25**, 1293–1301 (2011).

4. M. Elias, Evolution of aposematism and mimicry in butterflies: Causes, consequences and paradoxes. *Comptes Rendus Biologies*. **342**, 256–258 (2019).

5. M. Q. R. Pembury Smith, G. D. Ruxton, Camouflage in predators. *Biological Reviews*. **95**, 1325–1340 (2020).

6. S. Kondo, The present and future of Turing models in developmental biology. *Development*. **149**, dev200974 (2022).

7. A. M. Turing, The chemical basis of morphogenesis. *Bulletin of Mathematical Biology*. **52**, 153–197 (1990).

8. J. B. L. Bard, A model for generating aspects of zebra and other mammalian coat patterns. *Journal of Theoretical Biology*. **93**, 363–385 (1981).

9. S. Kondo, R. Asal, A reaction-diffusion wave on the skin of the marine angelfish Pomacanthus. *Nature*. **376**, 765–768 (1995).

10. S. Kondo, An updated kernel-based Turing model for studying the mechanisms of biological pattern formation. *Journal of Theoretical Biology*. **414**, 120–127 (2017).

11. K. L. Cooper, Self-organization in the limb: a Turing mechanism for digit development. *Curr Opin Genet Dev*. **32**, 92–97 (2015).

12. J. D. Glover, Z. R. Sudderick, B. B.-J. Shih, C. Batho-Samblas, L. Charlton, A. L. Krause, C. Anderson, J. Riddell, A. Balic, J. Li, V. Klika, T. E. Woolley, E. A. Gaffney, A. Corsinotti, R. A. Anderson, L. J. Johnston, S. J. Brown, S. Wang, Y. Chen, M. L. Crichton, D. J. Headon, The developmental basis of fingerprint pattern formation and variation. *Cell*. **186**, 940-956.e20 (2023).

13. Y. Nagashima, S. Tsugawa, A. Mochizuki, T. Sasaki, H. Fukuda, Y. Oda, A Rho-based reaction-diffusion system governs cell wall patterning in metaxylem vessels. *Sci Rep*. **8**, 11542 (2018).

14. A. Nakamasu, G. Takahashi, A. Kanbe, S. Kondo, Interactions between zebrafish pigment cells responsible for the generation of Turing patterns. *Proceedings of the National Academy of Sciences*. **106**, 8429–8434 (2009).

15. B. Ding, E. L. Patterson, S. V. Holalu, J. Li, G. A. Johnson, L. E. Stanley, A. B. Greenlee, F. Peng, H. D. Bradshaw, M. L. Blinov, B. K. Blackman, Y.-W. Yuan, Two MYB Proteins in a Self-Organizing Activator-Inhibitor System Produce Spotted Pigmentation Patterns. *Current Biology*. **30**, 802-814.e8 (2020).

16. C. B. Kaelin, K. A. McGowan, G. S. Barsh, Developmental genetics of color pattern establishment in cats. *Nat Commun*. **12**, 5127 (2021).

17. J. D. Murray, How the Leopard Gets Its Spots. *Scientific American*. **258**, 80–87 (1988).

18. R. T. Liu, S. S. Liaw, P. K. Maini, Two-stage Turing model for generating pigment patterns on the leopard and the jaguar. *Phys. Rev. E*. **74**, 011914 (2006).

19. M. De Gomensoro Malheiros, H. Fensterseifer, M. Walter, *ACM Trans. Graph.*, in press, doi:10.1145/3386569.3392478.

20. S. Kondo, H. Shirota, Theoretical analysis of mechanisms that generate the pigmentation pattern of animals. *Seminars in Cell & Developmental Biology*. **20**, 82–89 (2009).

21. P. Galipot, C. Damerval, F. Jabbour, The seven ways eukaryotes produce repeated colour motifs on external tissues. *Biological Reviews*. **96**, 1676–1693 (2021).

22. H. Shoji, Y. Iwasa, Labyrinthine versus straight-striped patterns generated by two-dimensional Turing systems. *Journal of Theoretical Biology*. **237**, 104–116 (2005).

23. V. A. Grieneisen, B. Scheres, P. Hogeweg, A. F. M Marée, Morphogengineering roots: comparing mechanisms of morphogen gradient formation. *BMC Syst Biol*. **6**, 37 (2012).

24. H. Yoshimura, S. Arakaki, M. Hamagawa, Y. Kitamura, M. Yokota, T. Denda, Differentiation of germination characteristics in Scutellaria rubropunctata (Lamiaceae) associated with adaptation to rheophytic habitats in the subtropical Ryukyu Islands of Japan. *J Plant Res*. **132**, 359–368 (2019).

25. C. F. Stevens, Darwin and Huxley revisited: the origin of allometry. *Journal of Biology*. **8**, 14 (2009).

26. N. Haupaix, C. Curantz, R. Bailleul, S. Beck, A. Robic, M. Manceau, The periodic coloration in birds forms through a prepattern of somite origin. *Science*. **361**, eaar4777 (2018).

27. S. Lee, "EXTRACTING INSIGHTS OF CLASSIFICATION FOR TURING PATTERN WITH FEATURE ENGINEERING" in (2020; https://www.semanticscholar.org/paper/EXTRACTING-INSIGHTS-OF-CLASSIFICATION-FOR-TURING-Lee/b7aee75fb7a5544290076fbd2cb3502df8358113).

28. M. J. How, J. M. Zanker, Motion camouflage induced by zebra stripes. *Zoology*. **117**, 163–170 (2014).

29. T.-L. Gluckman, G. C. Cardoso, The dual function of barred plumage in birds: camouflage and communication. *Journal of Evolutionary Biology*. **23**, 2501–2506 (2010).

30. C. Di Bernardi, A.-M. Thierry, N. E. Eide, D. E. Bowler, L. Rød-Eriksen, S. Blumentrath, L. Tietgen, B. K. Sandercock, Ø. Flagstad, A. Landa, Fitness and fur colouration: Testing the camouflage and thermoregulation hypotheses in an Arctic mammal. *Journal of Animal Ecology*. **90**, 1328–1340 (2021).

31. W. L. Allen, I. C. Cuthill, N. E. Scott-Samuel, R. Baddeley, Why the leopard got its spots: relating pattern development to ecology in felids. *Proceedings of the Royal Society B: Biological Sciences*. **278**, 1373–1380 (2010).

32. A. J. Bies, D. R. Blanc-Goldhammer, C. R. Boydston, R. P. Taylor, M. E. Sereno, Aesthetic Responses to Exact Fractals Driven by Physical Complexity. *Frontiers in Human Neuroscience*. **10** (2016) (available at https://www.frontiersin.org/articles/10.3389/fnhum.2016.00210).

33. D. M. Zaghloul, S. El-Gendy, Anatomical, light and scanning electron microscopical study of ostrich (Struthio camelus) integument. *Journal of Veterinary Anatomy*. **7**, 33–54 (2014).

34. R. R. Baker, G. A. Parker, G. Salt, The evolution of bird coloration. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*. **287**, 63–130 (1997).

35. B. Spehar, S. Wong, S. van de Klundert, J. Lui, C. Clifford, R. Taylor, Beauty and the beholder: the role of visual sensitivity in visual preference. *Frontiers in Human Neuroscience*. **9** (2015) (available at https://www.frontiersin.org/articles/10.3389/fnhum.2015.00514).

36. L. M. Mäthger, G. R. R. Bell, A. M. Kuzirian, J. J. Allen, R. T. Hanlon, How does the blue-ringed octopus (Hapalochlaena lunulata) flash its blue rings? *Journal of Experimental Biology*. **215**, 3752–3757 (2012).

37. R. A. Saporito, R. Zuercher, M. Roberts, K. G. Gerow, M. A. Donnelly, Experimental Evidence for Aposematism in the Dendrobatid Poison Frog Oophaga pumilio. *Copeia*. **2007**, 1006–1011 (2007).

38. A. M. M. Stuckert, K. Summers, Investigating signal modalities of aposematism in a poison frog. *Journal of Evolutionary Biology*. **n/a**, doi:10.1111/jeb.14111.

39. B. T. Werner, B. Hallet, Numerical simulation of selforganized stone stripes. *Nature*. **361**, 142–145 (1993).

40. N. E. Stork, How Many Species of Insects and Other Terrestrial Arthropods Are There on Earth. *Annual Review of Entomology* (2018), doi:10.1146/annurev-ento-020117-043348.

41. A. Blagodatski, A. Sergeev, M. Kryuchkov, Y. Lopatina, V. L. Katanaev, Diverse set of Turing nanopatterns coat corneae across insect lineages. *Proceedings of the National Academy of Sciences*. **112**, 10750–10755 (2015).

42. T. H. Büscher, M. Kryuchkov, V. L. Katanaev, S. N. Gorb, Versatility of Turing patterns potentiates rapid evolution in tarsal attachment microstructures of stick and leaf insects (Phasmatodea). *Journal of The Royal Society Interface*. **15**, 20180281 (2018).

43. Y. KonDo, S. Yoda, T. Mizoguchi, T. Ando, J. Yamaguchi, K. Yamamoto, Y. Banno, H. Fujiwara, Toll ligand Spätzle3 controls melanization in the stripe pattern formation in caterpillars. *Proc Natl Acad Sci U S A*. **114**, 8336–8341 (2017).

44. B. S. Heming, *Insect Development and Evolution* (Cornell University Press, 2018).

45. M. Leménager, J. Burkiewicz, D. J. Schoen, S. Joly, Studying flowers in 3D using photogrammetry. *New Phytologist*. **237**, 1922–1933 (2023).

46. F. J. Rohlf, The tps series of software. *Hystrix It. J. Mamm.* **26**, 9–12 (2015).

47. I. L. Dryden, K. V. Mardia, *Statistical Shape Analysis: With Applications in R* (John Wiley & Sons, 2016).

48. I. T. Jolliffe, "Generalizations and Adaptations of Principal Component Analysis" in *Principal Component Analysis*, I. T. Jolliffe, Ed. (Springer, New York, NY, 1986; https://doi.org/10.1007/978-1-4757-1904-8_12), *Springer Series in Statistics*, pp. 223–234.

49. R. da S. N. Barreto, M. N. Rodrigues, R. C. Carvalho, F. M. De Oliveira E. Silva, N. N. Rigoglio, J. C. F. Jacob, E. L. Gastal, M. A. Miglino, Organogenesis of the Musculoskeletal System in Horse Embryos and Early Fetuses. *The Anatomical Record*. **299**, 722–729 (2016).

50. F. Kuhar, L. Terzzoli, E. Nouhra, G. Robledo, M. Mercker, Pattern formation features might explain homoplasy: fertile surfaces in higher fungi as an example. *Theory Biosci.* **141**, 1–11 (2022).

Acknowledgments:

I deeply thank Dr. Marianne Elias and Dr. Sylvain Gerber for their corrections, remarks and advice about all phases of the project.

I deeply thank Prof. Marcelo de Gomensoro Malheiros from Centro de Ciências Computacionais, Brasil for his valuable remarks and suggestions concerning the numerical simulations.

I deeply thank Dr. Bart Buyck (ISYEB, MNHN, France), Dr. Tsuyoshi Hosoya, Dr. Jean-Philippe Rioult (UniCAEN, France) and Pr. Marc-Andre Selosse (ISYEB, MNHN, France) for the valuable mycological discussions concerning the links between Fungi, colors and Turing-like patterns.

I deeply thank Dr. Violaine Llaurens (ISYEB, MNHN, France) for her precious advice on the organization of the project and her remarks on the notions of Evolution mentioned in the article.

I deeply thank Dr. Visotheary Ung (ISYEB, MNHN, France) for having helped to set up the website hosting the database and for her precious advice on the data formatting.

I deeply thank Marion Leménager and Haruhiko Fujiwara for their help concerning the photogrammetry experiments on Phalaenopsis hieroglyphica.

I deeply thank Émilie Dupré for her work on classic Turing and PGTCPs pattern recognition by deep learning, which will help complete the database and make it as exhaustive as possible in the future.

I deeply thank Florian Jabbour and Catherine Damerval for their advice and corrections on the ultimate version of the project.

Funding: JSPS (Japan Society For the Promotion of Science), Short-Term Fellowship.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: Survey Data of Fig.1 is available at www.eukolorpatterns.cnrs.fr

Materials and Methods

PGTCPs survey among Eukaryotes

Putative Growth Turing-like Color Patterns (PGTCPs) were classified into 4 categories, which could be combined within the same species: rosettes (and mazes of mazes), intermediate stripes, mixed colors, and line-and-dot alternation.

We checked both for classic Turing Patterns and for PGTCPs, and in these two categories four types of labels have been given be for each group: Y (Yes, if at least one species with a clear pattern has been found), M (Maybe Yes, if at least one species with a potential but not clear pattern has been found), N? (Maybe No, if no species with pattern has been found, but if the search was non exhaustive) and N (Probably No, if no species with pattern has been found, if the search was almost exhaustive).

The complete results are available in a database accessible at the following URL (www.eukolorpatterns@cnrs.fr) open to contributions. They comprised the complete list of taxa explored, the names of the candidate species and a link to a representative picture of patterns. For the almost exhaustive surveys of mammals and orchids, lists of species were respectively extracted from ASM Mammal Diversity database (https://www.mammaldiversity.org/, January 2022), and COL, Catalogue of Life (https://www.catalogueoflife.org/).

We use multiple sources for photographic or textual proofs attesting a classic Turing pattern or a PGTCP, Handbook of the Mammals of the World (HMW), Volume 1 to 9, Lynx Edicions for mammals and http://www.orchidspecies.com/ for orchids, with additional sources from the scientific literature when a photograph was not available. When possible, in mammals, juvenile patterns were also looked at, although the vast majority of sources are limited to the adult, and most often the male. In some species, sexual dimorphism resulted in the presence of a Turing pattern only in the female (e.g., *Cervus elaphus*), and in others, only the juveniles showed a PGTCP (e.g., the wild boar *Sus scrofa*). Many examples of PGTCPs may have been missed due to poor sources on juvenile phenotypes.

The lists of orders for animals and families for Angiosperms (or higher or lower taxonomic level if order/family did not exist) were extracted from COL, Catalogue of Life (https://www.catalogueoflife.org/). On a case-by-case basis but not exhaustively, corrections were made if scientific literature no longer recognized the taxon. Again, the sources of photography were multiple, scientific literature was preferred, then online encyclopedias Wikipedia (https://en.wikipedia.org/) and Wikimedia Commons (https://commons.wikimedia.org), eventually alternative sources (through search engines) with keywords such as "pattern", "spotted", "stripes", when the first three were not conclusive.

PGTCPs mapping on phylogenetic trees

Phylogenetic relationships between species were built with phyloT (V2, https://phylot.biobyte.de/) based on NCBI taxonomy. Unrecognized taxa were resolved by hand. Phylogenetic trees were plotted and annotated in iTOL (Interactive Tree Of Life, version 6.5.8, https://itol.embl.de/). Tree leaves were name after the found species name if at least one example was found, or by the name of the order/family if not.

PGTCPs modeling

A custom Turing model was built in Processing 4.0 software language (https://processing.org/),
based on De Gomensoro Malheiros et al equations (*19*).

An example of code can be found at
(https://github.com/PierreGalipot/growthpattern/blob/9b3fa4db636e7ca7410c7c346d2032829bcd
6172/leopardgrowth)

Fixed parameters for all simulations presented in this study:

s = 6

r = 30

dt = 0.002

minA = 0

maxA = 4.4

minB = 3.5

maxB = 100

Growth was indirectly implemented by its effect on diffusion (via s parameter).

PGTCPs occurrences morphometrics

For Burchell's zebra (*Equus quagga burchellii*), we hypothesized that its external embryonic
development was close to that of the horse (*Equus ferus caballus*, whose embryogenesis is well
documented. We selected a panel of landmarks spaced throughout the body delineating 22
segments whose lengths were measured at each developmental stage. For each segment, the
relative increase in size was calculated (growth factor), allowing us to associate with each point
of the tissue an estimated value of local growth depending on its distance to each of the 22
segments and on each segment growth factor.

$$growth\_factor(pixel) = \frac{\sum_{i=1}^{22} \frac{growth\_factor(segment(i))}{distance(pixel, segment(i))^2}}{\sum_{i=1}^{22} \frac{1}{distance(pixel, segment)^2}}$$

A color code was then attributed for each pixel and plotted in Processing 4.0 software language
(https://processing.org/).

Code can be found at this URL:
https://github.com/PierreGalipot/growthpattern/blob/a9909be6f70d25cfc8d835d02ea5492548aa0
a56/occurences_NCTPs_VS_Growthmaps.

This growth map was then compared to a zebra map of PGTCPs occurrence build by hand by
reporting tissue zones where PGTCPs are observed.

Reverse method morphometrics

Two data were needed to estimate the timing of the pattern implementation: i) tissue
morphometrics during development, for this purpose distances between landmarks were used as
a proxy of the shape and ii) the theoretical shape of the tissue at the formation of the pattern,
obtained by using adult pattern and applying corrections from the number of patterns present on
each distance. For example, if the neck-tail distance of an adult leopard is x times longer than the
head, but there are y times fewer motifs along it, this means that at the time of pattern formation
the relative size of the neck-tail distance was $1/(x*y)$ that of the head. By repeating this for
independent distances on the tissue, if the estimations tend to converge to a specific
developmental stage, we could make the reasonable assumption that this is the timing of the end
of Turing pattern production.

We used the cat (*Felis silvestris*) and leopard models, the first having a well described
embryogenesis for which morphometric measurements are applicable, the second for its non-
canonical patterns, while being morphologically close to the first at least until its juvenile stages.
We selected 11 distances representative of external morphology (head, trunk, tail, neck, limbs),
and measurable from early embryonic stages to adults. All these distances were plotted relative
to head length (Fig. 4.B) during development. The head was chosen because its dynamics is far
from most other distances, which allows for maximum contrast in relative distances. Motifs
counts of the adult cat and adult leopard were used to calculate theoretical values for each
distance, and by graphical reading, the most likely approximate theoretical stage to which the
distance refers was estimated and represented by ellipses, whose size was representative of the
relative distance range covered during development (in this case, tail and trunk have the most
contrasted dynamics compared to the head). This technique was also applied to PGTCPs,
considering either that a rosette (or equivalent) was a single motif (to calculate the timing of
production of the first generation of dots), or by considering its constituent sub-patterns (to date
the second generation of motifs).

From each timing prediction, the wavelength of the pattern at its formation can be calculated,
since the number of patterns along its distance is known and the distances are also known in an
absolute way by morphometry. In the same way, if these predictions converge towards a value,
we can assume that this value is close to the real value of the wavelength of the initial pattern.

Reverse method modeling

To test the prediction validity, and how well patterns can be used as predictors of morphometry,
we performed simulations of PGTCPs formation and counted the number of motifs obtained over
time and compared to the theoretical ones. To calculate the theoretical value of the motifs
number, simulations were run without growth and at different tissue sizes. As expected, the final
number of motifs is proportional to the tissue size, for a given set of Turing system parameters.
Different growth rates were simulated (0.01, 0.02, 0.05, 0.1, 0.2) starting from a tissue size = 1
with different maximum sizes (1/0.4, 1/0.25, 1/0.185). The number of iterations was between
t=75000 and t=100000 and a screenshot was taken every t=250. The number of motifs was
counted in an automated way with a custom macro on Fiji software (version 2.3.0) based on the

"Find maxima" process, after having verified the validity of the pattern detection algorithm in
several representative situations.

Although we let the simulations continuing after growth had stopped, in order to observe the
behavior of the system, the most important timing is the moment when the tissue reached its final
size. The difference between the number of motifs observed at this timing and the expected
number of motifs was calculated and compared between different values of growth rate and final
size.
757

Artificial landmarks and in vivo monitoring and estimation of tissue local growth

We adapted a photogrammetry method (*45*) to an *in vivo* flower bud of *Phalaenopsis*
hieroglyphica. Points were disposed by hand with a laboratory waterproof ink marker pen.
Photographs have been taken with an Olympus OMD E-M10 14-42mm camera. For preliminary
test of geometric morphometrics, landmark coordinates were manually recorded using tpsDig2
(*46*) and processed following standard procedures with tpsRelw (*46*). Landmark configurations
were optimally aligned by generalized Procrustes analysis (*47*), which extract shape data by
filtering out differences in scale, location, and orientation among raw landmark configurations.
The patterns of shape variation were explored using the principal component analysis (PCA)
(*48*).
768

Fig. 1. Diversity of growth-putative
Turing color patterns (PGTCPs) of
Eukaryotes. (**A**) Details of PGTCPs in
Animalia (left) and Angiosperms (right),
divided into four categories: line-and-dots
alternation, mixed color, rosettes,
intermediate bands. (**B**) Occurrences of
classic Turing patterns and PGTCPs in all
Animalia orders mapped on Animalia
phylogenetic tree. Grey leaf: no Turing
pattern found in this order, blue leaf: only
classic Turing pattern found in this order,
yellow leaf: PGTCP found in at least one
species. (**C**) Proportions of classic Turing
Patterns and PGTCPs for mammals
exhaustive focus. (**D**) Proportions of
classic Turing Patterns and PGTCPs for
orchids exhaustive focus. (**E**) Occurrences
of classic Turing patterns and PGTCPs in
all Angiosperms families mapped on
Angiosperms phylogenetic tree. Grey leaf:
no Turing pattern found in this order, blue
leaf: only classic Turing pattern found in
this order, yellow leaf: PGTCP found in at
least on species.

Fig. 2. Modeling of tissue growth reproduce all types of PGTCPs via diverse mechanisms.
(**A**) First row: simulation of *Phalaenopsis hieroglyphica* rosette patterns (with higher lateral growth). Second row: simulation of leopard (*Panthera pardus*) rosette patterns (with uniform growth). (**B**) Leopard-like pattern is obtained through superimposition of early and late patterns via a memory mechanism, absent in *P. hieroglyphica* (**C**) First row: simulation of *Pomacanthus imperator* intermediate band pattern. Second row: simulation of line-and-dots patterns, with the example of *Ictidomys tridecemlineatus*. (**D**) Overview and details of P. imperator, *I. tridecemlineatus* and *Stapelia gigantea* PGTCPs.

Fig. 3. PGTCPs correlate with most-expensing regions of the tissue, suggesting the responsibility of growth in their formation. (**A**) Examples of presence/absence of PGTCPs are found at every biological scale, suggesting a universal mechanism of control, growth being the main candidate. (**B**) Approximation of zebra (*Equus quagga*) external morphology dynamics via measurements of absolute distances between landmarks at several developmental stages (in horse embryo, pictures taken from (*49*)) and in zebra adult. (**C**) Comparison between most-expensing regions map, left, obtained by assigning to every tissue location a growth value depending to its distance to every distance and its respective growth factor, and PGTCPs occurrences map, right, obtained from adult zebra skin photographs, middle, with a satisfying correlation between the two maps.

843
Fig. 4. PGTCPs can be a source of information thanks to a reverse method. (**A**) Left: examples of
distortions in classic Turing pattern (*Mimulus sp.*) and a PGTCP (*Panthera pardus*) Right: reverse method
principles and assumptions. (**B**) Dynamics of relative size of various distances to head size during
development of the cat (*Felix sylvestris*, stages 14 to 23) and leopard (*Panthera pardus,* stage 24 and 25)
and predicted classic and PGTCPs pattern formation. (**C**) Simulation of dynamics of motifs number
during pattern formation under tissue growth. Y-axis: number of motifs simulated at different growth
rates (solid lines), compared to the theoretical number of motifs (dashed lines). Up left and down right: 4
details of pattern phenotype through growth for g = 0.05, at different times (1: early, 2: pattern at the start
of growth, 3: pattern just before growth stops, 4: late pattern at the equilibrium). (**D**) Tissue growth
monitoring with hand-written landmarks on the tissue, example of *Phalaenopsis hieroglyphica* sepal outer
epidermis. Upper panel: Sequence of landmarked flower bud. Lower panel: Left, four views extracted
from the photogrammetry model of *P. Hieroglyphica* flower bud (at Day 6). Right, preliminary 2D
geometric morphometrics landmarks assessment and comparison between young stage (Day 6, 1st and 3rd
pictures) and final stage (Day 41, 2nd picture). Fourth picture: grid deformation between young and late
stage after Procrustes analysis.

859
Fig. 5. Potential biological functions enhanced by PGTCPs. (**A**) Turing systems are a source of color
segregation, leading to contrasted patterns with more extreme and less intermediate colorations. Species
or individual with intermediary color patterns (up) compared to relative ones with Turing patterns (down).
(**B**) Intermediate colors and novel shapes induced by growth could improve camouflage of species. Up:
leopard rosettes have characteristics that can potentially improve the effectiveness of camouflage. Down:
Different ways of producing intermediate colors, improving camouflage in juvenile Ratites: via pigments
giving intermediate colors like orange pheomelanin (cassowary, ostrich and maybe *Rhea*), via feathers
whose large size and arrangement mix the colors produced by melanocytes (ostrich), or via PGTCPs (emu
and maybe *Rhea*) (**C**) Positive correlation between pattern preference and pattern sensitivity (time needed
for the brain to process the signal). Adapted from (*35*) (**D**) *Dendrobates leucomelas* line-and-dots

alternations and *Hapalochlaena lunulata* ringed rosettes have characteristics that can potentially improve
the effectiveness of their aposematic pattern, like unusual motifs, more fractal and periodic patterns.

Fig. 6. Three ways to lose PGTCPs information from prepattern to final phenotype. A) no loss. B)
Decoupling C) Pixelating D) Blurring.

Taxon	Classic Turing-like Color patterns estimated sparsity (+, ++, +++, absence)	PGTCPs estimated sparsity (+, ++, +++)	Potential causes for Turing-like patterns (*in italics*) and PGTCPs (**in bold**) sparsity, other than chance.
Insecta	++	+++	• *Segmented and compartmented body (too little space to produce proper motifs)* • *Segmented and compartmented body (the pre-existing repeated structures of the body may favor other types of color pattern mechanisms, as the ability of Turing systems to produce repeated patterns may be de facto obsolete)* • **Growth modalities of insect body parts often involves a discontinuous growth and a small absolute increase in tissue surface at each iteration, making it potentially more difficult to form PGTCPs.**
Aves	Juveniles (+), adults (+++)	Juveniles (+++), adults (+++)	• *Many juveniles have no down or colored skin. Disordered down or a too low feather density could hide potential Turing-patterns or PGTCPs (by decoupling or pixelation).* • *The feathers of adults make it difficult or impossible for a skin-level 2D Turing pattern to keep its visible geometry at the scale of the plumage (decoupling).* • **Many species harbor repeated patterns on their feathers, but the underlying mechanism is *a priori* not compatible with the 2D growth necessary for PGTCPs formation.**
Fungi	Absent?	Absent?	• *Physical cracking patterning (such as the very famous* Amanita muscaria *red and white pattern or black and white desquamation pattern of* Coprinopsis picacea*) could have been preferentially selected during Evolution at the expense of potential Turing mechanism).* • *Ecologically, there might by a reduced evolutionary pressure to produce repeated color patterns on Fungi caps (even if Amanitaceae produce some striking ones) and other ways to attract/repel other species might have been favorized like poisons or attractive flavors. Many Fungi taxa seem to exhibit a poor relationship between colors and attractiveness/repellency.* • *Turing patterns not involved in coloration might be responsible for the great hymenium shape diversity (50). This putative ability of Fungi to produced Turing patterns, together with their absence for color patterning strengthens the previous point.*
Angiosperms	++	+	• *Families whose species are not based on zoogamy seem to be spared from repeated color patterns, which would then explain the absence of classic Turing patterns and PGTCPs.* • **Some species could produce prepatterns of PGTCPs but may be hidden when the colors are produced, by blurring. We have observed many orchid species that keep PGTCPs stigmas (like spots with tortuous edges) but many of the typical characters are absent (rosettes, intermediate colors).**

882   Table 1. Potential causes for Turing-like patterns and PGTCPs sparsity in specific taxa