Extensive profiling of histidine-containing dipeptides reveals species-specific distribution and

metabolism in mice, rats and humans

Short title: Profiling of histidine-containing dipeptides

Thibaux Van der Stede[1,2,†], Jan Spaas[1,3,4,†], Sarah de Jager[1,†], Jana De Brandt[4,5], Camilla Hansen[2], Bjarne

Vercammen[1], Siegrid De Baere[6], Siska Croubels[6], Ruud Van Thienen[1], Kenneth Verboven[4,5], Dominique

Hansen[4,5,7], Thierry Bové[8], Bruno Lapauw[9], Charles Van Praet[10,11], Karel Decaestecker[10,11], Bart

Vanaudenaerde[12], Bert O Eijnde[3,13,14], Lasse Gliemann[2], Ylva Hellsten[2], Wim Derave[1,*]

[1] Department of Movement and Sports Sciences, Ghent University, Belgium.

[2] Department of Nutrition, Exercise and Sports, Copenhagen University, Denmark.

[3] University MS Center (UMSC) Hasselt – Pelt, Belgium.

[4] BIOMED Biomedical Research Institute, Hasselt University, Belgium.

[5] REVAL Rehabilitation Research Center, Hasselt University, Belgium.

[6] Department of Pathobiology, Pharmacology and Zoological Medicine, Ghent University, Belgium.

[7] Heart Center Hasselt, Jessa Hospital Hasselt, Belgium.

[8] Department of Cardiac Surgery, Ghent University Hospital, Belgium.

[9] Department of Endocrinology, Ghent University Hospital, Belgium.

[10] Department of Urology, Ghent University Hospital, Belgium.

[11] Department of Human Structure and Repair, Ghent University, Belgium.

[12] Department of Chronic Diseases and Metabolism, KU Leuven, Belgium.

[13] SMRC Sports Medical Research Center, BIOMED Biomedical Research Institute, Hasselt University,

Belgium.

[14] Division of Sport Science, Stellenbosch University, South Africa.

[†] These authors have contributed equally

* Correspondence: Wim.Derave@UGent.be

Abstract

Histidine-containing dipeptides (HCDs) are pleiotropic homeostatic molecules linked to inflammatory, metabolic and neurological diseases, as well as exercise performance. Using a sensitive UHPLC-MS/MS approach and an optimized quantification method, we performed a systematic and extensive profiling of HCDs in the mouse, rat, and human body (in n=26, n=25, n=19 tissues, respectively). Our data show that tissue HCD levels are uniquely regulated by carnosine synthase, an enzyme preferentially expressed by fast-twitch skeletal muscle fibers and brain oligodendrocytes. Cardiac HCD levels are remarkably low. The low abundant HCD N-acetylcarnosine is enriched in human skeletal muscles. Here, N-acetylcarnosine is continuously secreted into the circulation as the most stable plasma HCD, which is further induced by acute exercise in a myokine-like fashion. Carnosine is preferentially transported within red blood cells in humans but not rodents. We provide a novel basis to unravel tissue-specific, paracrine, and endocrine roles of HCDs in human health and disease.

Teaser: Human muscle releases N-acetylcarnosine at rest and especially during exercise, potentially initiating tissue crosstalk.

Introduction

Carnosine synthase (CARNS1) is presumably the only enzyme in animals capable of synthesizing an abundant class of endogenous dipeptides. The enzyme links L-histidine to either β-alanine or γ-aminobutyric acid (GABA), respectively rendering carnosine or homocarnosine. These parent dipeptides and their methylated (anserine and balenine) and acetylated (N-acetylcarnosine) analogs are collectively called the histidine-containing dipeptides (HCDs).

Since the initial discovery in 1900 by Vladimir Gulevich (*1*), carnosine and the other HCDs have been linked to various physiological functions, mostly serving to preserve redox status and cellular homeostasis (for a full overview, see (*2*)). The most relevant biochemical properties for their functions relate to proton buffering, metal chelation and antioxidant capacity, which further translates to protection against advanced glycation and lipoxidation end products (*3, 4*). The physiological importance of tissue HCD content is underscored by an extensive body of research ranging from enhancement of exercise performance (*5*) to treatment of cardiometabolic (*6, 7*) or neurological diseases (*8*) in rodents. Major differences between animal and human HCD metabolism may be present however, given that high carnosinase (CN1) activity in human, but not rodent, plasma results in rapid degradation of carnosine (*9, 10*).

Nevertheless, even more than 120 years after the initial discovery of carnosine and 10 years after the molecular identification of CARNS1 (*11*), there remains a lack of basic understanding of HCD synthesis, distribution, and metabolism throughout the animal and especially the human body. It is thought that HCDs are primarily expressed in excitable tissues such as skeletal and cardiac muscle and the central nervous system (CNS), but current literature mostly consists of scattered observations focussing on a limited number of tissues or species. Information on cardiac levels is sparse, although HCDs could play an important role in cardiomyocyte homeostasis (*12*). Furthermore, there is unclarity regarding the synthesis and physiological role of HCDs in kidney, lung, liver, and other non-excitable tissues. A first profiling of HCDs in rat tissues from Aldini *et al.* (*13*) did not detect HCDs in non-excitable tissues, although this and other previous endeavours were potentially limited from lower detection sensitivity compared to the currently available technology. For example, the low abundant HCDs anserine, balenine and N-acetylcarnosine have never been extensively characterized in animal or human tissues.

Here, we have performed the first systematic profiling of the five main HCDs, combined with determination of CARNS1 expression levels, in the mouse, rat and human body. Various human tissue samples were collected from live donors, except for post-mortem collected brain regions. We uncovered profound differences in HCD distribution and metabolism between tissues and species. For

72   instance, we demonstrate that humans have a unique way of circulating HCDs and releasing it from

73   carnosine-synthesizing tissues such as skeletal muscle.

Results

<u>CARNS1 is the unique and rate-limiting enzyme for HCD synthesis</u>

Using whole-body *Carns1*-knockout (KO) mice, we aimed to investigate whether CARNS1 deficiency

results in a complete lack of endogenous (homo)carnosine and their derivatives in a variety of tissues,

which would imply that CARNS1 is the unique and rate-limiting enzyme for HCD synthesis. CARNS1 was

successfully knocked out at the gene (**Fig 1A**) and protein (100 kDa, **Fig 1B**) level, thereby also validating

our Western blot antibody for specifically detecting the CARNS1 protein. As described previously,

Carns1-KO mice displayed normal growth and survival (*14*). The deletion of *Carns1* led to an absence

of carnosine and homocarnosine in all investigated tissues (**Fig 1C**). Similarly, these mice were devoid

of the carnosine-derived analogs anserine, balenine and N-acetylcarnosine (of which only anserine is

consistently present in mouse tissues, cfr. infra, **Fig 1C**).

In addition, tissue from *Carns1*-KO mice was used to optimise our quantitative UHPLC-MS/MS-based

detection of HCDs. Concentration levels in muscle and brain were compared using matrix-matched

standard calibration curve preparation in *Carns1*-KO tissue matrix (i.e. muscle or brain homogenates)

and water (as current standard practice in HCD research). In water, HCD levels were overestimated by

~2 to 3-fold (**Fig 1D-E**), indicating the importance of utilizing a corresponding blank tissue matrix for

HCD quantification. This approach was used for all further analyses in this paper (except human

cerebrospinal fluid), rendering the HCD quantifications more accurate than previously reported.

<u>HCDs are not excitable tissue-specific metabolites</u>

Using our sensitive UHPLC-MS/MS method, we performed a systematic profiling of HCDs in various

tissues of the mouse, rat, and human body (**Fig 1F**, **Table S1**). Carnosine was the only HCD present in

all studied tissues across the three species. Skeletal muscles contained the largest amounts of HCDs,

followed by the CNS, up to the millimolar range. However, not all excitable tissue contained large

amounts of HCDs, since unexpectedly low HCD levels were observed in the heart of all three species

(50-100 times lower than skeletal muscles). These low HCD levels in cardiac tissue better reflect those

measured in non-excitable tissues. Besides skeletal muscle, CNS, heart, kidney, adipose, lung and liver

tissue (presented in **Fig 1F**), we also found low levels of HCDs in the mouse and rat stomach wall, gallbladder, pancreas, small intestine, colon, thymus, spleen, and eye (**Table S1**).

Figure 1. Species- and tissue-specific distribution of histidine-containing dipeptides. (A) PCR gel and **(B)** Western blot showing the successful knockout of the *Carns1* gene and the absence of CARNS1 protein in mice. **(C)** HCD measurements by UHPLC-MS/MS showing HCDs are absent from various tissues of *Carns1*-KO compared to WT mice. **(D)** Carnosine and **(E)** homocarnosine measured by UHPLC-MS/MS, followed by quantification based on a standard calibration curve prepared in water *vs. Carns1*-KO tissue matrix. **(F)** HCD measurements by UHPLC-MS/MS in skeletal muscle, CNS, heart, kidney, adipose tissue, lung, and liver tissue from mice (M), rats (R) and humans (H). Values were averaged if more than 1 type of the respective tissue was present (e.g. soleus and EDL for rodent muscle). ANS, anserine; BAL, balenine; CAR, carnosine; Cereb., cerebellum; CNS, central nervous system; EDL, extensor digitorum longus; Fr. ctx., frontal cortex; HCAR, homocarnosine; Hippoc., hippocampus; HZ, heterozygous; KO, knockout; NAC, N-acetylcarnosine; OB, olfactory bulb; Sp. cord, spinal cord; WT, wild type.

CARNS1 content drives fiber type-related differences in HCD content in skeletal muscle

We next aimed to profile the HCD content in skeletal muscle in more detail, with a focus on potential fiber type-specific differences. In mice and rats, we determined the HCD content in soleus (more oxidative, slow-twitch) and extensor digitorum longus (EDL; more glycolytic, fast-twitch) muscles. Our

results confirmed previous reports (*15-17*) that carnosine and anserine content is higher in EDL muscle (**Fig 2A**). Also the homocarnosine content was ~2.5-fold higher in EDL compared to soleus in both mice and rats (**Fig 2B**). To get more insights if CARNS1 content (i.e. HCD production) is the main driver for the fiber type-related differences, we first explored a publicly available human muscle fiber type-specific RNAseq dataset (*18*). These data show a ~2-fold higher *CARNS1* mRNA content in type IIa *vs.* type I fibers (**Fig 2C**). We next determined CARNS1 protein levels in the soleus and EDL muscles from the mice and rats (**Fig 2D-E**). CARNS1 levels were indeed higher in EDL muscles, with a larger difference between soleus and EDL muscles in mice (6.6-fold) than in rats (1.4-fold). To translate mRNA differences in human muscle to the protein level, we used Western blotting on pools of pre-classified type I or type IIa fibers (**Fig S1**). This revealed a ~2-fold higher CARNS1 content in type IIa fibers (**Fig 2D-E**), consistent with findings at the mRNA level. These results suggest that CARNS1 expression is the main driver regulating the clear fiber type-related differences in HCD content across the three species.

Figure 2. Muscle and muscle fiber type-specific differences in histidine-containing dipeptides and CARNS1. (A) Carnosine and anserine and (**B**) measurements by UHPLC-MS/MS in soleus and EDL muscles from mice and rats. **(C)** Human muscle fiber type-specific calculation of *CARNS1* mRNA based on a previously published dataset. **(D)** Protein levels of CARNS1 determined by Western blot in soleus and EDL muscle from mice and rats, and human type I and type II fiber pools. **(E)** Representative Western blot and loading controls. ANS, anserine; CAR, carnosine; EDL, extensor digitorum longus.

<u>Homocarnosine is the dominant HCD in the CNS, except for the olfactory bulb of rodents</u>

Apart from skeletal muscle, HCDs are also highly present in the CNS. By analyzing seven different

regions from the mouse, rat and human CNS, we could confirm that the highest levels of carnosine are

found in the olfactory bulb in rodents, reaching concentrations of ~1200 μmol/kg tissue, which is

similar to or even higher than skeletal muscle carnosine levels (**Fig 3A, Table S1).** In contrast, the

human olfactory bulb contained approximately 15 times less carnosine (~80 μmol/kg). In mice and

rats, the olfactory bulb was the only CNS region containing more carnosine than homocarnosine, whilst

in humans all regions contained more homocarnosine than carnosine. Similar to our findings in human

CNS tissue, homocarnosine was also abundantly present in human cerebrospinal fluid (**Fig 3B**).

CARNS1 protein levels also showed considerable variability between CNS regions (**Fig 3C-D**). Mice and

rats exhibited high expression in olfactory bulb, as well as the spinal cord and medulla oblongata, but

not the rest of the CNS. In human tissues, we found lower CARNS1 levels in the olfactory bulb, but

instead observed greater amounts in the white matter, thalamus, and frontal cortex.

Immunofluorescence was used to further study the localisation of CARNS1 in the CNS. We chose the

region exhibiting the greatest CARNS1 protein levels among rodents (mouse olfactory bulb) and

humans (white matter). Double-labeling of CARNS1 and OLIG2, an oligodendrocyte lineage marker,

revealed that CARNS1 resides in oligodendrocytes of human white matter (**Fig 3E**). Cell markers for

microglia (CD68, **Fig S2A**), astrocytes (GFAP, **Fig S2B**) and neurons/axons (NF-H, **Fig S2C**) did not co-

localise with CARNS1. In the mouse olfactory bulb, CARNS1 appeared in spherical structures near the

surface of the olfactory bulb, i.e. the glomeruli, where olfactory nerve terminals form synapses with

dendrites from projection neurons that carry signals into the brain (**Fig 3F**).

<u>CARNS1 expression scales with tissue HCD levels on a whole-body level</u>

To investigate if tissue CARNS1 expression closely relates to tissue HCD levels, CARNS1 protein levels

(Western blot) were plotted against HCD concentrations (UHPLC-MS/MS). If the tissue CARNS1 level is

the main determinant of tissue HCD levels, a linear relation between both variables is expected. In all

non-excitable tissues, no CARNS1 could be detected by Western blot (**Fig 4A-C**). HCD levels in these

tissues probably reflect transmembrane HCD uptake. On a whole-body level, CARNS1 scaled with HCD

levels in mice (**Fig 4A**), rats (**Fig 4B)** and humans (**Fig 4C**). However, when comparing within organs,

more CARNS1 was not always directly linked to a higher HCD concentration. In the human CNS, for

example, CARNS1 was 13-fold higher in white matter than in the cerebellum, but both tissues had

similar HCD levels. This indicates that although CARNS1 is the only enzyme responsible for HCD

synthesis, other factors, such as exchange of HCDs between organs or a high HCD turnover rate (synthesis/degradation), could also influence intracellular HCD levels.

Figure 3. Region-specific levels of histidine-containing dipeptides and CARNS1 in the mouse, rat and human central nervous system. (A) HCD measurements by UHPLC-MS/MS in seven different central nervous system regions from mice (M), rats (R) and humans (H). **(B)** HCD measurements by UHPLC-MS/MS in human cerebrospinal fluid. **(C)** Protein levels of CARNS1 determined by Western blot in seven different central nervous system regions from mice, rats and humans. Data are mean ± SD. **(D)** Representative Western blot and loading controls. **(E)** Immunohistochemical detection of CARNS1 and OLIG2 in human white matter. **(F)** Immunohistochemical detection of CARNS1 in mouse olfactory bulb. In panels (A), (C) and (D), spinal cord tissue is from the cervical region, mouse white matter is from the corpus callosum,

and human frontal cortex is from the superior frontal gyrus. ANS, anserine; AU, arbitrary units; BAL, balenine; CAR, carnosine; Cereb., cerebellum; CNS, central nervous system; Fr. ctx., frontal cortex; HCAR, homocarnosine; MO, medulla oblongata; NAC, N-acetylcarnosine; OB, olfactory bulb; Sp. cord, spinal cord; Thal., thalamus; W. Ma., white matter. Scale bars are 50 μm (E), 100 μm (F).

Figure 4. Association between CARNS1 and tissue histidine-containing dipeptide levels. Plotted relationship between CARNS1 levels (Western blot) and HCD measurements (UHPLC-MS/MS) in a variety of **(A)** mouse, **(B)** rat, and **(C)** human tissues. The central nervous system (CNS) regions that are shown, besides the olfactory bulb (OB), are frontal cortex (FrCtx), cerebellum (Cereb), spinal cord (SpC), thalamus (Thal), white matter (WMa) and medulla oblongata (MO). Mouse muscles are soleus (Sol) and extensor digitorum longus (EDL). Human muscles are m. vastus lateralis (VL) and m. pectoralis (Pect). Human adipose tissue is subcutaneous and visceral fat (Adip). Human kidney is medulla and cortex. Data are mean ± SEM. AU, arbitrary units.

N-acetylcarnosine and balenine are mainly found in human skeletal muscle

Whilst the parent HCDs (carnosine and homocarnosine) were ubiquitously expressed, most of the examined tissues also contained at least one methylated (anserine or balenine) or acetylated (N-acetylcarnosine) carnosine analog (**Fig 1F**). Rodent skeletal muscles contained by far the highest anserine levels (up to ~1500 μmol/kg in the rat EDL, **Fig 5A**). Besides anserine (~40 μmol/kg, **Fig 5A**), human skeletal muscle also contained N-acetylcarnosine (~25 μmol/kg, **Fig 5B**) and balenine (~5 μmol/kg, **Fig 5C**). In fact, human skeletal muscle was the tissue where we observed the highest N-acetylcarnosine and balenine levels. **Fig 5D-F** display the proportion of methylated or acetylated carnosine variants in different species and tissues.

Figure 5. Anserine, N-acetylcarnosine and balenine levels in mouse, rat and human tissues. (A) Anserine, **(B)** N-
acetylcarnosine, and **(C)** balenine measurements by UHPLC-MS/MS in mouse (M), rat (R), and human (H) tissues. The
figures display the 8 tissues with the highest concentration of each dipeptide. Data are mean ± SD. **(D-F)** Relative
proportion of anserine, N-acetylcarnosine, and balenine in **(D)** mouse, **(E)** rat, and **(F)** human tissues. Adip., adipose;
ANS, anserine; BAL, balenine; CNS, central nervous system; NAC, N-acetylcarnosine; n.d., not detectable; Olf. bulb,
olfactory bulb; Sk., skeletal.

Acetylation of carnosine provides stability in the human circulation, which is not required in red blood cells or rodents

Since rodents and humans differ substantially in presence and activity of the hydrolyzing enzyme CN1 in the circulation (*2, 10*), we attempted to map the circulating content of the five main HCDs. As expected, levels of plasma carnosine and anserine were very high in mice (~1500 nM) and rats (~3500 nM), but in the low nanomolar range in humans (**Fig 6A**). Low levels of homocarnosine could be detected in all 3 species, while balenine was only present in human plasma (**Fig 6A**). Interestingly, N-acetylcarnosine was the most abundant HCD in the human circulation, accounting for ~45% of the total HCDs (**Fig 6B**). Although HCD levels in red blood cells (RBCs) were in the nanomolar range in all 3

species, striking differences were observed compared to plasma (**Fig 6A** and **6C**). For both rodent species, HCD levels were drastically lower in RBCs than plasma (**Fig 6D** and **S3**). On the contrary, human carnosine and anserine levels were higher in every RBC sample compared to plasma, whilst N-acetylcarnosine levels were lower in RBCs than plasma (**Fig 6E**). These data suggest that carnosine in the human circulation is rendered more stable via acetylation to N-acetylcarnosine (which is resistant to hydrolysis by CN1) or via transport inside RBCs.

Figure 6. Species differences in histidine-containing dipeptides in plasma and red blood cells. (A) HCD measurements by UHPLC-MS/MS in mouse, rat and human plasma and red blood cells (data are mean ± SD). **(B)** Relative proportion of each HCD in mouse (M), rat (R) and human (H) plasma. **(C)** Relative proportion of each HCD in mouse, rat and human red blood cells. **(D)** Direct comparison of HCDs in plasma and red blood cells collected from the same mice. **(E)** Direct comparison of HCDs in plasma and red blood cells collected from the same human individuals. ANS, anserine; BAL, balenine; CAR, carnosine; HCAR, homocarnosine; NAC, N-acetylcarnosine; RBC, red blood cells.

<u>Oral β-alanine supplementation affects HCDs in skeletal muscle and circulating N-acetylcarnosine levels in humans</u>

We next investigated the effects of chronic supplementation of the rate-limiting precursor β-alanine on the content of all five HCDs in human skeletal muscle. As expected, β-alanine supplementation increased muscle carnosine content (+89%, **Fig 7A** and **S4A**). In addition, we observed increases in muscle anserine (+38%, **Fig 7A** and **S4B),** N-acetylcarnosine (+97%, **Fig 7A** and **S4C**) and balenine (+199%, **Fig 7A** and **S4D)**, and these increases were proportional to the carnosine increase (**Fig 7B**). However, homocarnosine content significantly decreased (-50%) after 12 weeks of β-alanine supplementation (**Fig 7A** and **S4E**). Plasma N-acetylcarnosine increased by 48% after the supplementation period (**Fig 7A** and **S4E**), which correlated at the individual level with muscle

carnosine content (**Fig 7C**), suggesting a possible link between intramuscular and circulating HCD levels.

In summary, oral β-alanine supplementation is a potent stimulus affecting all HCDs in skeletal muscle,

and plasma N-acetylcarnosine may reflect muscle HCD levels under baseline conditions and during β-

alanine supplementation.

<u>N-acetylcarnosine is released from human skeletal muscle during exercise</u>

Given this likely relationship between intra- and extracellular HCDs, and given that skeletal muscle is

the main active organ during exercise, we explored HCD dynamics during exercise. First, we collected

muscle interstitial fluid at rest and during exercise in humans. During exercise, interstitial levels for

every HCD increased (**Fig 7D**). This increase could however be primarily caused by sarcolemmal

damage following insertion of the microdialysis probe (*19*). To check this, we performed two follow-

up experiments. Firstly, *in vitro* human primary muscle cells were electrically stimulated for 2-3 h to

simulate muscle contraction. This did not result in secretion of carnosine into the culture medium

immediately after the electrical stimulation or following 30 min recovery (**Fig 7E**). Other HCDs could

not be detected in the cell culture medium. Secondly, we collected interstitial fluid from mouse skeletal

muscle, with a previously published method in which the muscle is not mechanically affected (*20, 21*).

Interstitial levels of carnosine and anserine were not higher in exercised mice compared to control

mice (**Fig S5A**). Next, we collected samples of the femoral artery and vein at rest and during exercise

(from a group of postmenopausal women). Our results clearly indicate a release of N-acetylcarnosine

from muscle tissue at rest of ~13 nmol/min (from one leg), which further increased ~2-fold during

exercise (**Fig 7F**). No release at rest or during exercise was observed for any of the other HCDs (**Fig 7F**).

These results were confirmed using a similar experimental setup in healthy young men, and with more

sampling time points during rest, passive movement, exercise and recovery. Here, we again showed a

release of N-acetylcarnosine at rest (~14 nmol/min from one leg), which increased 3.4-fold during

exercise and quickly returned back to baseline during recovery (**Fig 7G**). Exercise did not induce

carnosine release or uptake within RBCs (**Fig S5B**). Also in mice, no changes in carnosine or anserine

levels within plasma were observed following 60 min exercise (**Fig S5C**). Taken together, these data

indicate that N-acetylcarnosine in humans is likely the major, or most stable, HCD released during

exercise in a myokine-like fashion.

Figure 7. β-alanine supplementation and N-acetylcarnosine release during exercise from human muscle. (A) Changes in HCD levels, measured by UHPLC-MS/MS, in human skeletal muscle (m. vastus lateralis) after β-alanine supplementation. **(B)** Correlation between supplementation-induced changes in skeletal muscle carnosine and N-acetylcarnosine, measured by UHPLC-MS/MS. **(C)** Correlation between skeletal muscle carnosine and plasma N-acetylcarnosine, measured by UHPLC-MS/MS. **(D)** HCD measurements by UHPLC-MS/MS in human skeletal muscle interstitial fluid at rest and following exercise (Ex.). **(E)** Carnosine measurements by UHPLC-MS/MS in culture medium from primary human muscle cells in control condition, after 2-3 h of electrical stimulation (Stim), and after 2-3 h of electrical stimulation followed by 30 min recovery (Stim+Rec). **(F)** HCD measurements by UHPLC-MS/MS in human arterial and venous plasma samples at rest and following exercise (Ex.). Positive values indicate a net release, negative

Discussion

This is the first study to systematically and extensively profile the organ distribution of HCDs and their differences between mice, rats and humans. Although present in all investigated non-excitable tissues in minute amounts, mainly excitable tissues contained high HCD levels in all species. Yet, our data show surprisingly low values in cardiac tissue across species, and a different distribution of HCDs in CNS regions between species. The enzyme CARNS1 is the unique enzyme responsible for endogenous carnosine and homocarnosine synthesis, and is a major determinant for tissue HCD levels. In human CNS white matter, CARNS1 appears restricted to cells from the oligodendrocyte lineage. We also uncover that N-acetylcarnosine is the primary circulating HCD in human plasma and is continuously secreted from skeletal muscle into the circulation, which is further increased by physical exercise. An overview of the main findings is visualized in **Fig 8**.

Figure 8. Graphical summary. Visual representation of the main findings, based on our extensive profiling of histidine-containing dipeptides (HCDs) in mice, rats and humans, and related follow-up experiments. Arrow up indicates high abundance, arrow down indicates low abundance. CNS, central nervous system; RBC, red blood cell.

Previous endeavors to profile HCDs mostly focused on rodent tissues, and resulted in fragmented and sometimes contradictory literature (*13, 22-25*). Our systematic approach and very sensitive state-of-the-art UHPLC-MS/MS methodology facilitate direct comparison between tissues and species. Our data contradicted some of the previous findings, e.g. that human muscle contains only carnosine and no other HCDs (*2*), that rat kidney, lung, plasma and liver lack HCDs (*13*), or that homocarnosine is

exclusively found in the CNS (*26*), with no presence of carnosine in the human brain or cerebrospinal

fluid (*27*). We confirmed that anserine is predominantly found in rodent skeletal muscles (*2, 16*), but

add that N-acetylcarnosine and balenine were primarily enriched in human skeletal muscles, although

still lower than anserine and (homo)carnosine. From our systematic approach, we calculate that 99.1%

of the total amount of HCDs in the human body is found in skeletal muscle tissue, which confirms

previous estimations (*28*). Moreover, HCD concentrations in the present study are often different than

those previously reported. More specifically, we report lower HCD concentrations in human skeletal

muscle than previous studies (*29-31*). This is most likely explained by the use of tissue-specific *Carns1*-

KO tissue matrix for our quantification method, which is known to be important for MS-based

quantification (*32*). Furthermore, Peters *et al.* reported very high carnosine (1800 µmol/kg) and

anserine (4000 µmol/kg) concentrations in human kidney (*23*), which are approximately 1000-fold

higher than the concentrations in our dataset (~2 µmol/kg). Besides the use of a different detection

technique and quantification method, it is unclear where such differences may have originated from.

CARNS1 and HCDs, especially carnosine, have long been recognised as enriched compounds within the

olfactory tract of rodents (*33, 34*). In human olfactory bulbs, we found remarkably low levels of

carnosine compared to homocarnosine. In addition, this is the first study to unequivocally ascribe

CARNS1 expression to a specific cell type within the CNS; outside of the olfactory bulb. Our discovery

of CARNS1 localisation within cells of the oligodendrocyte lineage (human white matter) confirms

suggestions from recent RNA sequencing databases of the mouse and human CNS that reported

Carns1/*CARNS1* as an oligodendrocyte-enriched gene (*35, 36*).

Traditionally, the highest HCD levels are assigned to the excitable tissues. Though this holds true with

respect to skeletal muscle and CNS tissue, it was quite compelling that we found extremely low

amounts of HCDs in cardiac muscle tissue. In contrast to previous suggestions that the rat heart

contains ~10 mM HCDs (*37*), we report 100-fold lower levels. This was consistently found in all three

species we investigated, and is also 50- to 100-fold lower compared to the concentrations in skeletal

muscle. This can potentially be attributed to more accurate and sensitive quantification compared to

the older technology. Nevertheless, HCDs are thought to play a crucial role in cardiac function and

recovery from injury (*12, 38*). For instance, isolated cardiac myocytes from *Carns1*-transgenic hearts

were protected against hypoxia reoxygenation injury (*39*), whilst *Carns1*-KO rats have impaired cardiac

contractility accompanied by reduced Ca^{2+} peaks and slowed Ca^{2+} removal (*40*). This underscores the

physiological importance of HCDs and indicates that even low HCD levels can contribute significantly

to cell/organ function and health. It remains to be determined, however, which biochemical properties

and physiological roles of these pleiotropic molecules are the most important in different tissues and

under different conditions. With respect to cardiomyocytes, it has been proposed that carnosine

functions as a Ca^{2+}/H^+ exchanger to shuttle calcium towards and protons away from the sarcomere site

(*41, 42*).

The parent HCDs carnosine and homocarnosine share the same synthesizing enzyme CARNS1. This also

underlies our observation that oral β-alanine intake leads to increased carnosine but reduced

homocarnosine levels in human muscle, implying substrate inhibition between GABA and β-alanine for

CARNS1. Additionally, high expression of CARNS1 can lead to high tissue content of either carnosine

or homocarnosine, probably dependent of the local precursor availability (GABA *vs.* β-alanine). *Carns1*-

KO mice did not produce HCDs, whereas in WT mice, rats and humans, there appears to be a

relationship between CARNS1 expression and HCD content on a whole-body level. Moreover, the

differences in HCD content between slow- and fast-twitch muscle fibers were paralleled by similar

differences in CARNS1 expression. However, the correlation between CARNS1 expression and HCD

content was not perfect, suggesting that there might be inter-organ exchange or that there is higher

HCD turnover in tissues that have an important role for carnosine consumption/recycling, for example

through oxidative stress and reactions with toxic metabolites in pathological conditions (*8*).

Alternatively, the fact that CARNS1 shows a preference for β-alanine compared to GABA as a substrate

may skew this relationship considering that some tissues contain more carnosine than homocarnosine

and vice versa (*11*).

Tissues that have no or minimal CARNS1 expression likely rely on transmembrane uptake of HCDs

derived from exogenous/dietary sources or from production in CARNS1-expressing organs. It has

remained unclear, however, if and how HCDs are transported between tissues. The detection of HCDs

in various rodent and human tissues likely illustrates that HCDs can be exchanged between organs with

and without synthesizing capacity. Indeed, mice that lack the carnosine transporter PEPT2 have altered

(mostly reduced) HCD levels in various organs but increased levels in skeletal muscle tissue, which is

capable of synthesizing carnosine itself (*22*). This issue remains largely unclear in humans, in which

high activity of the CN1 enzyme quickly degrades circulating carnosine (*2*). It has long been suggested

that circulating HCDs are extremely low or absent in human plasma (*43-45*), although more recent

reports already detected carnosine (*46, 47*). We now demonstrate that N-acetylcarnosine is the most

stable carnosine analog in plasma, indicating that acetylation of the β-alanine residue protects against

the hydrolyzing activity of CN1. Thus, N-acetylcarnosine may be the primary HCD that is exchanged

between tissues in humans. Interestingly, plasma N-acetylcarnosine levels were correlated to muscle

carnosine (and N-acetylcarnosine) levels, possibly indicating that plasma N-acetylcarnosine can serve

as a surrogate marker for intramuscular HCD levels. Plasma N-acetylcarnosine levels increased

following β-alanine intake, showing that circulating N-acetylcarnosine levels may also be a marker for

muscle carnosine/HCD loading. In addition, our data indicate that transport of carnosine in RBCs is an

alternative strategy to protect against CN1 in human plasma, as recently suggested (*48*). This was not

true for rodents, who had lower HCD levels in RBCs than humans, despite more than 25 times higher

plasma HCD levels.

It has been proposed that carnosine is released from muscles during periods of contractile activity,

potentially serving as a health-promoting myokine. This hypothesis is primarily based on a study in

rats, where plasma carnosine levels increased during the dark/active phase when animals were

provided with a running wheel (*49*). We report that in humans, N-acetylcarnosine is the only HCD that

is consistently released from muscle tissue into plasma at rest, which is further increased during

periods of muscular activity (exercise). This opens various new research avenues on N-acetylcarnosine

as an exercise-induced myokine (*50, 51*). Future experiments should determine its relevance for

exercise training adaptations and cell/organ crosstalk (*52*). The average N-acetylcarnosine release of

14.3 nmol/min from non-contracting leg muscles at rest is striking. Extrapolation of this release to the

whole body, assuming that all muscles have the same N-acetylcarnosine secretion, suggests that in

theory the blood N-acetylcarnosine concentration should increase by 23.9 µM every 24 hours. Despite

this continuous and substantial release of N-acetylcarnosine into the circulation, resting plasma N-

acetylcarnosine levels only reach 50-350 nM in most subjects, indicating that there is a large

uptake/utilization of N-acetylcarnosine in other organs, or urinary excretion. Based on the amount of

N-acetylcarnosine release measured in arterio-venous samples from the leg, we also estimated that a

daily turnover of 25% of the total muscle N-acetylcarnosine pool is needed to maintain stable muscle

N-acetylcarnosine levels, suggesting a rather dynamic HCD homeostasis in human muscle. Specific

description of these calculations and used assumptions can be found in the **Supplementary Text**. In

muscle interstitial fluid samples, all HCDs appear to increase during exercise. However, it is hard to

distinguish physiological exercise-induced release from sarcolemmal rupture caused by insertion of

the microdialysis probes (*19*), as also supported by (I) the absence of a carnosine release during

electrical stimulation of primary human muscle cells or interstitial fluid sampled from mice post-

exercise, and (II) the lack of other HCD release (besides N-acetylcarnosine) in the venous effluent of

contracting muscles.

Despite being the first study to systematically and extensively study the distribution of HCDs in three

species, we acknowledge that the mouse and rat data cannot be fully extrapolated to all mouse and

rat strains, since these can differ somewhat (*53*). We also decided to focus on the two parent HCDs

(carnosine and homocarnosine) and carnosine's methylated and acetylated analogs (anserine,

balenine, N-acetylcarnosine). Other HCD conjugates do exist, but these are mostly very low abundant

products from reactions with other (toxic) compounds (e.g. carnosine-propanol or 2-oxo-carnosine

(*54-56*)).

In conclusion, we extensively profiled the organ distribution of the five main HCDs and discovered new
physiological routes of HCD metabolism. Our results can be used to generate various new research
hypotheses and highlight that findings derived from animal research on HCDs can not always be
translated to humans. In particular, the apparent inter-cell and inter-tissue para- and endocrine
regulation of HCDs, as well as its relevance to human health, disease and exercise
performance/adaptation, deserve further investigation.

Materials and methods

<u>HCD profiling - Rodent tissue collection</u>

All mouse tissues were obtained from an in-house breeding of *Carns1*-KO and WT mice with a C57/BL6
background, kindly provided by Prof. M. Eckhardt (*14, 57*). Genotypes of the offspring from
heterozygous parents were determined in toe samples using a previously published protocol (*14*).
Wistar rats were supplied by Envigo (The Netherlands). Mice and rats were housed under standard
room conditions (12h:12h light:dark cycle, 20-24°C, relative humidity 30-60%) and had *ad libitum*
access to drinking water and food pellets. For tissue collection, female and male animals were
sacrificed at an age of 6-10 w (mice) or 7-8 w (rats) old. Following overdose injection of Dolethal
(200 mg/kg, i.p.), blood was collected from the right ventricle, kept in Multivette® 600 K3 EDTA vials
on ice, centrifuged (5 min, 3500 rpm), and plasma was stored at -80°C. Before tissue dissection, mice
and rats were perfused with 0.9% NaCl solution containing heparin (25 UI/mL) via a left ventricular
puncture. For determination of HCD levels by UHPLC-MS/MS and CARNS1 expression by Western blot,
tissues were immediately frozen in liquid nitrogen, before being stored at -80°C. For
immunohistochemistry, whole mouse brains were carefully placed on a metal plate cooled by dry ice
in a foam box for several minutes, wrapped in aluminum foil, and stored at -80°C. Mouse exercise
experiments were performed on a treadmill (6 m/min, speed increased every 2 min by 2 m/min until
16 m/min, total duration 60 min), and were preceded by a 1-week adaptation period (3 running
sessions, gradually increasing exercise intensity and duration). Sedentary mice were placed on a
stationary treadmill for 60 min. Immediately after exercise, plasma was obtained and mice were
perfused as described above. To collect interstitial fluid, gastrocnemius muscles were placed on 20 μM
nylon net filters (Millipore, cat# NY2004700) and centrifuged (10 min, 800 × g) (*20, 21*). All animal
procedures were approved by the Ethical Committee on Animal Experiments at Hasselt University
(202074B, 202127 and 202145).

 HCD profiling - Human tissue collection

All human samples were obtained after written informed consent.

Human vastus lateralis muscle: Muscle biopsies were collected from the m. vastus lateralis of healthy,
young volunteers using the Bergström needle biopsy technique with suction, as described previously
(*58*). One part of the samples was immediately snap-frozen in liquid nitrogen and stored at -80°C until
UHPLC-MS/MS analysis. The other part was submerged in 1-1.5 mL of RNA*later* (Thermo Fisher
Scientific), stored at 4°C for maximum 48 h and subsequently stored at -80°C for later individual fiber
dissection.

Human heart and pectoralis muscle: Heart and pectoralis muscle samples were collected from
patients undergoing open heart surgery under general anaesthesia. Heart samples consisted of the
right atrial appendage, harvested at the time of venous drainage cannulation for cardiopulmonary
bypass. Samples were immediately snap-frozen in liquid nitrogen and stored at -80°C. For **Fig 1F**, HCD
concentrations of vastus lateralis and pectoralis muscle were averaged.

Human kidney: Kidney samples were collected from patients undergoing radical nephrectomy. In case
of kidney cancer, tissue was sampled as far away from the site of the tumor to obtain the healthiest
part of the kidney, immediately snap-frozen in liquid nitrogen and stored at -80°C. Medulla and cortex
were sampled separately (**Table S1**), but data was later averaged for analysis and visualization.

Human adipose tissue: Visceral and subcutaneous adipose tissue were sampled from lean and obese
individuals during abdominal surgery following an overnight fast (*59*). Tissue pieces were rinsed, freed
from visible blood and connective tissue, and snap-frozen in liquid nitrogen. The two subtypes (**Table
S1**) were later averaged for analysis and visualization.

Human lung: Lung tissue was obtained from unused healthy donor lungs that were not used for
transplantation from the BREATH KULeuven biobank (S51577).

Human liver: Liver samples were surgically removed from fasted subjects with obesity during gastric
bypass surgery. Exclusion criteria were the presence of malignancies, drinking more than two units
(women) or three units (men) of alcohol per day, having known liver pathologies other than non-
alcoholic fatty liver disease. For the current analysis, five samples with NAS score 0 or 1 were selected
(*60*).

Human plasma and RBC: Blood samples were always collected in pre-cooled EDTA tubes, after a 1-2-
day lacto-ovo vegetarian diet to ensure no influence of dietary HCD intake. For plasma, samples were
immediately centrifuged (10 min, 3000 × g, 4°C), followed by immediate deproteinization (110 µL of

35% 5-sulfosalisylic acid per 1 mL of plasma) and second centrifugation (5 min, 15000 × g, 4°C). Plasma

samples were then stored at -80°C. For RBC isolation (*61*), blood tubes were centrifuged for 15 min at

120 × g at room temperature. Plasma was carefully removed and 200 µL RBCs were collected in 1.8 mL

of ice-cold methanol (55% v/v). Samples were then stored at -80°C.

Human CNS: Seven different human CNS regions were obtained from The Netherlands Brain Bank

(NBB), Netherlands Institute for Neuroscience, Amsterdam (open access www. brainbank.nl). All

donors were 'non-demented controls', indicating the absence of neurological and psychiatric disease.

From 2 subjects, all 7 regions were available. For immunohistochemical analyses, white matter tissue

from healthy controls and unaffected white matter tissue of multiple sclerosis patients from a previous

study was used.

Human cerebrospinal fluid: Cerebrospinal fluid from subjects without neurological disease at the time

of sampling was obtained via the University Biobank Limburg (UbiLim), with approval from the Medical

Ethics Committee at Hasselt University (CME2021-004). Lumbar cerebrospinal fluid was collected into

PPS tubes, kept at 4°C, centrifuged to remove cells (10 min, 500 × g, 4°C), and supernatant was stored

at -80°C.

<u>β-alanine supplementation study in humans</u>

Vastus lateralis muscle biopsies obtained via the Bergström needle biopsy technique with suction

(n=11 β-alanine, n=11 placebo) and plasma samples (n=19 β-alanine, n=17 placebo) were collected

before and after 12 weeks of β-alanine supplementation in patients with COPD (sustained-release

CarnoSyn®, NAI). Methodological details have been described previously (*62*). Snap-frozen biopsies

were freeze-dried for 48 h, followed by manual removal of non-muscle material (fat, connective tissue,

blood) under a light microscope. Effects of β-alanine were analyzed using a two-way repeated

measures ANOVA (group *vs*. time) for each HCD separately, followed by Sidak's multiple comparisons

tests. Correlations were performed using Pearson correlation (Δ muscle carnosine *vs.* Δ N-

acetylcarnosine) or Spearman rank correlation (muscle carnosine *vs.* plasma N-acetylcarnosine).

<u>Human HCD release experiments</u>

Microdialysis experiment: Detailed methodology has been described previously (*63*). In short,

interstitial samples from m. vastus lateralis were collected using the microdialysis technique at rest

and after 30 min of one-legged knee extensor exercise (15 min 6 W, 15 min 50% peak power). Subjects

consisted of a group of young (n=7) and old (n=13) healthy men (results are pooled together since no differences between groups could be observed). Concentrations were corrected for probe recovery as determined by relative loss of [2–3H]-labeled adenosine in the dialysate. Interstitial levels during exercise were compared to resting values using a multiple Wilcoxon matched-pairs signed rank test with Holm-Sidak multiple comparison test.

Arterio-venous balance experiment 1: Arterial and venous samples were collected from the femoral artery/vein at rest and after 30min one-legged knee extensor exercise (6W) in a group of healthy postmenopausal women (n=19). Methodological details can be found in (*64*). Samples were deproteinized with 35% 5-sulfosalisylic acid, as described above, on the day of the UHPLC-MS/MS analysis. Exercise *vs.* resting HCD levels were compared using a multiple Wilcoxon matched-pairs signed rank test with Holm-Sidak multiple comparison test.

Arterio-venous balance experiment 2: Seven healthy, young men (28 ± 4 years old, BMI of 24 ± 2, VO_{2max} of 49 ± 7 mL/min/kg) participated in this experiment. After passive transport to the lab, catheters were inserted in the femoral artery and vein. Next, arterial and venous samples were collected every 30 min during a 90 min supine resting period. After this, 20 min of passive leg movement was performed, followed by 60 min of active one-legged knee extensor exercise (20 min 10 W, 20 min 20 W, 20 min 10 W). Samples were collected at the end of each exercise bout. Finally, arterio-venous samples were collected after 30 min and 2 h of recovery. Plasma and RBC samples were collected as described above. Exercise-induced effects were analyzed using a mixed-effect model with repeated measurements over time, with post-hoc comparison of every time point *vs.* baseline (Holm-Sidak test).

Primary cell culture experiments

Biopsy samples (~150 mg) were obtained from m. vastus lateralis from young, healthy men. Primary skeletal muscle cells were isolated with homemade antibody-coated magnetic beads and cultured as previously described (*65, 66*). Cultured skeletal muscle cells were used for analysis on day 5 or 6 after the onset of differentiation. At this time, most of the myocytes have differentiated into multinucleated myotubes and can easily be identified as muscle cells. Myotubes were starved with media containing 0.1% Bovine Serum Albumin (DMEM without phenol, D-glucose and L-glutamine) for 16 hours before experiments. The skeletal muscle cells were electro-stimulated as described previously (*67*), with the minor addition of 5μM (S)-nitro-Blebbistatin (Cayman Chemical, CAS. 856925-75-2) to the stimulation buffer to inhibit the spontaneous contraction of the myotubes (*68*). The cells were stimulated for 2-3 h (50 Hz, 0.6s/0.4 s trains, 1 ms pulse width, 10 V, homemade electrical stimulator connected to a

Digitimer MultiStim SYSTEM-D330). The extracellular medium was collected immediately or 30 min
after the end of stimulation, and medium from non-stimulated control cells was harvested
simultaneously. Changes over time were analyzed using a mixed-effect model with repeated
measurements over time, with post-hoc comparison of stimulated and stimulated+recovery *vs.* control
cells (Holm-Sidak test).

<u>HCD determination by UHPLC-MS/MS</u>

Details of the validation of the in-house developed UHPLC-MS/MS analysis has been described
previously (*69*), with the exception that all experiments were performed on a Xevo® TQ-S MS/MS
system with 2.5 µL injection volume. The limit of detection was determined to be 5-10 nM (in plasma),
corresponding to 0.38-0.76 µmol/kg tissue for our homogenization protocol. Pure carnosine and
anserine were kindly provided by Flamma S.p.a. (Chignolo d'Isola, Bergamo, Italy), and pure balenine
by NNB Nutrition (Frisco, Texas, USA). Homocarnosine (#33695) and N-acetyl-L-carnosine (#18817)
were bought from Cayman Chemical (Ann Arbor, Michigan, USA). UHPLC-MS/MS data extraction and
analysis was performed using Masslynx software 4.2 (Waters, Milford, USA).

Tissues: All tissues were prepared similarly, based on the method described previously (*8*). Frozen
tissues were quickly weighed and immediately homogenized in extraction solution (ultrapure water
with 10 mM HCl and internal standard carnosine-d4) in a ratio of 95 µL extraction solution per 5 mg
tissue in a QIAGEN TissueLyser II (1 min, 30 Hz). The concentration of carnosine-d4 varied according to
expected HCD concentrations in the tissues: muscle (20 µM), CNS (5 µM), liver/lung/spleen (0.5 µM)
and all other tissues (1 µM). Then, homogenates were centrifuged (20 min, 3000 × g, 4°C).
Supernatants were immediately diluted in a 3:1 ratio with ice-cold acetonitrile (-20°C), vortexed and
kept on ice for 15min. After a second centrifugation step (20 min, 3000 × g, 4°C), samples were stored
at -80°C until the day of the UHPLC-MS/MS analysis. Samples were combined with 75:25
acetonitrile:water in a 4:1 ratio before injection in the UHPLC-MS/MS device. For skeletal muscle and
CNS, samples were also injected after an extra initial dilution (1:25 for muscle, 1:10 for CNS) for the
determination of carnosine (skeletal muscle) and homocarnosine (CNS) content. Standard calibration
curves were prepared for each individual run and for each tissue separately in the respective tissue of
the *Carns1*-KO mice to account for possible tissue matrix effects (for all three species). Differences
between soleus and EDL HCD levels in mice and rats were analyzed using paired t-tests or Wilcoxon
signed-rank tests, depending on normality of the data.

Plasma: Deproteinized plasma (150 µL) was combined with acetonitrile containing 1% formic acid (215
µL), 1 µM carnosine-d4 as internal standard (10 µL) and ultrapure water (25 µL). For mouse plasma

analysis, volumes were scaled down to available plasma (75 μL). After thoroughly vortexing, samples were centrifuged (15 min, 15000 × g, 4°C). The supernatant was collected and injected in the UHPLC-MS/MS. A standard calibration curve was prepared in a pool of human deproteinized plasma that was collected after a 2-day lacto-ovo vegetarian diet to minimize circulating HCDs (for all three species).

RBC: First, 190 μL of the RBC samples was combined with 10 μL of internal standard carnosine-d4 (2μM) and 10 μL of 75:25 acetonitrile:water (with 1% formic acid). This mixture was vortexed and then centrifuged in a 10 kDa filter (Nanosep® Centrifugal Device with Omega Membrane™, Pall Corporation). The supernatant was subsequently evaporated at 40°C and the droplet resuspended in 40 μL of 75:25 acetonitrile:water (with 1% formic acid) before injection in the UHPLC-MS/MS device. A standard calibration curve was prepared in a pool of human RBC samples that were collected after a 2-day lacto-ovo vegetarian diet to minimize circulating HCDs (for all three species).

Cerebrospinal fluid: Human cerebrospinal fluid was treated identically as plasma, but the standard calibration curve was prepared in ultrapure water since no *Carns1*-KO tissue matrix was available, possibly resulting in overestimation of the absolute concentrations.

Interstitial fluid: Interstitial fluid (15 μL) was combined with 30 μL acetonitrile containing 1% formic acid and 1 μM carnosine-d4 and 10 μL ultrapure water. This mixture was vortexed, centrifuged (15 min, 15000 × g, 4°C) and the supernatant was used to inject in the UHPLC-MS/MS. For detection of carnosine in humans, samples were first diluted 1:30. For detection of carnosine and anserine in mice, samples were first diluted 1:500. A standard calibration curve was prepared in Ringer-Acetate buffer as this was used to perfuse the microdialysis probes.

Cell culture medium: Extracellular medium (150 μL) was mixed with acetonitrile containing 1% formic acid (240 μL) and internal standard carnosine-d4 (2 μM, 10 μL), vortexed and injected in the UHPLC-MS/MS. A standard calibration curve was prepared in DMEM culture medium.

CARNS1 protein levels by Western blot

Tissues were diluted in RIPA buffer (300 μL per 10 mg tissue; 50 mM Tris pH 8.0, 150 mM NaCl, 0.5% sodium deoxycholate, 0.1% SDS, 1% Triton-X100, and freshly added protease/phosphatase inhibitors [Roche]), and homogenized using stainless steel beads and a QIAGEN TissueLyser II (shaking 1 min, 30 Hz). Following centrifugation (15 min, 12000 × g, 4°C), supernatants were stored at -80°C. Pierce™ BCA Protein Assay Kit (Thermo Fisher) was used according to manufacturer's instructions to determine protein concentrations (read at 570 nm wavelength). For detection of CARNS1 protein levels, 1 μg of protein was diluted in loading buffer solution (63 mM Tris Base pH 6.8, 2% SDS, 10% glycerol, 0.004%

Bromophenol Blue, 0.1 M DTT), heated for 4 min at 95°C, and separated in polyacrylamide gels (4-15%,

Mini-PROTEAN TGX, Bio-Rad) at 100-140 V on ice. Next, stain-free gels were imaged following UV

exposure to visualise total protein content (ChemiDoc MP Imaging System, Bio-Rad). Proteins were

transferred from the gel to an ethanol-immersed PVDF membrane in transfer buffer (30 min, 25 V, 1.0

A, Trans-Blot Turbo Transfer System, Bio-Rad). Membranes were briefly washed in Tris-buffered saline

with 0.1% Tween20 (TBS-T), and blocked for 30 min using 3% milk powder in TBS-T. Following overnight

incubation at 4°C with primary antibodies against CARNS1 (rabbit polyclonal, 1:1000 in 3% milk/TBS-T,

HPA038569, Sigma), membranes were washed (3 × 5 min), incubated with secondary HRP-conjugated

goat anti-rabbit antibodies (1:5000 in 3% milk/TBS-T) for 60 min at room temperature, washed again

(3 × 5 min), and chemiluminescent images were developed in a ChemiDoc MP Imaging System (Bio-

Rad) using Clarity Western ECL substrate (Bio-Rad). CARNS1 protein bands were quantified with Image

Lab 6.1 software (Bio-Rad), and normalized to total protein content from the stain-free image. Finally,

bands were expressed relative to total CARNS1 expression (sum of all bands) of a particular mouse,

rat, or human. For muscle, band densities were expressed as fold changes relative to the average

density from soleus muscles per blot. Linearity of the signal was determined for every tissue.

Differences between soleus and extensor digitorum longus CARNS1 content were analyzed using a

paired t-test (rat) or Wilcoxon signed-rank test (mouse), depending on normality of the data.

<u>CARNS1 protein level in human single muscle fibers.</u>

CARNS1 protein levels in type I *vs*. type II muscle fibers were determined based on a previously

published method (*70*). Muscle samples in RNA*later* (Thermo Fisher Scientific) were thawed and

subsequently transferred to a petri dish filled with fresh RNA*later* (Thermo Fisher Scientific) solution.

Individual muscle fibers were manually dissected under a light microscope and immediately

submerged in a new 0.5 mL tube with 5 μL ice-cold Laemmli buffer (125 mM Tris-HCl (pH 6.8), 10%

glycerol, 125 mM SDS, 200 mM DTT, 0.004% bromophenol blue). Tubes were then incubated for 15

min at 4°C, followed by 10 min at 70°C and then stored at -80°C until the next step of the analysis. A

total of 40-72 fibers were isolated from 9 biopsies. Next, muscle fiber type (based on myosin heavy

chain expression) was determined using dot blotting techniques. For this, 0.5 μL of the muscle fiber

lysate was spotted onto two activated and equilibrated PVDF membranes, one for MHC I and one for

MHC IIa. After air-drying the PVDF membrane for 30 min, it was re-activated in 96% ethanol and

equilibrated in transfer buffer (8 mM Tris-base, 39 mM glycine, 0.015% SDS, 20% ethanol). The next

steps are similar to standard Western blotting as described above, with primary antibodies for MHC I

(A4.840, 1:1000 in 3% milk in TBS-T, DSHB) or MHC IIa (A4.74, 1:1000 in 3% milk in TBS-T, DSHB). Fiber

lysates were classified as type I or IIa fibers based on a positive stain for only the MHC I or IIa antibody (**Fig S1**). Next, fibers of the same type from the same subject were pooled (n=10-26 for type I and n=9-22 for type IIa fibers). CARNS1 protein content in these fiber type-specific samples were determined with standard Western blotting technique as described above, with loading of 5 µL per pool. Band densities were expressed as fold changes relative to the average density from type I fibers per blot. Statistical analysis was performed using a Wilcoxon signed-rank test.

CARNS1 mRNA expression

The fiber type-specific RNAseq dataset of Rubenstein *et al.* was downloaded from the GEO repository under accession number GSE130977 (*18*). This dataset consists of RNAseq data of pools of type I or type II fibers from m. vastus lateralis biopsies of 9 healthy, older men. For full details on generation of this dataset, we refer to the original publication. Raw counts were normalized with DESeq2 to allow for between-sample comparisons (*71*). First, normalized counts for *MYH2* (ENSG00000125414, type II fibers) and *MYH7* (ENSG00000092054, type I fibers) were extracted and assessed for each fiber pool as purity quality control. Based on this analysis, the fiber pools of one participant were excluded for further analysis. Next, normalized counts of *CARNS1* (ENSG00000172508) were extracted and compared between type I and type II fiber pools within each participant using a Wilcoxon signed-rank test.

Immunohistochemical detection of CARNS1

Sagittal and frontal cryosections (10 µm) were cut from whole mouse brains and human white matter samples. Following acetone fixation (10 min) and blocking (30 min, 10% donkey serum in PBS with 1% BSA), sections were exposed overnight at 4°C to antibodies detecting CARNS1 (rabbit polyclonal, 1:100 in PBS with 1% BSA, HPA038569, Sigma). The next day, sections were washed and exposed to complementary secondary antibodies for 60 min (1:500 in PBS with 1% BSA, Thermo Fisher). Fluorescence imaging was performed with a Leica DM4000 B LED (Leica Microsystems). For double-labeling, we used the following antibodies: OLIG2 (1:50, goat polyclonal, AF2418, R&D Systems), neurofilament heavy polypeptide (NF-H, rabbit polyclonal, 1:200, ab8135, Abcam), CD68 (mouse monoclonal, 1:100, M0814 KP1 clone, Dako), GFAP (mouse monoclonal, 1:100, G3893, Sigma). Absence of CARNS1 from *Carns1*-KO brain sections was used as negative control.

 Statistical analysis

678 Statistical analyses were performed in GraphPad Prism v9.4 and were described in the appropriate

679 method paragraphs. Significance level was set at $\alpha=0.05$.

681 References

1.   W. Gulewitsch, S. Amiradžibi, Ueber das Carnosin, eine neue organische Base des
Fleischextractes. *Berichte der deutschen chemischen Gesellschaft* **33**, 1902-1903 (1900).
2.   A. A. Boldyrev, G. Aldini, W. Derave, Physiology and pathophysiology of carnosine.
Physiological reviews **93**, 1803-1845 (2013).
3.   G. Aldini, B. de Courten, L. Regazzoni, E. Gilardoni, G. Ferrario, G. Baron, A. Altomare, A.
D'Amato, G. Vistoli, M. Carini, Understanding the antioxidant and carbonyl sequestering
activity of carnosine: Direct and indirect mechanisms. *Free Radical Research* **55**, 321-330
(2021).
4.   J. J. Matthews, M. D. Turner, L. Santos, K. J. Elliott-Sale, C. Sale, Carnosine increases insulin-
stimulated glucose uptake and reduces methylglyoxal-modified proteins in type-2 diabetic
human skeletal muscle cells. *Amino Acids*, 1-8 (2023).
5.   B. Saunders, K. Elliott-Sale, G. G. Artioli, P. A. Swinton, E. Dolan, H. Roschel, C. Sale, B. Gualano,
β-alanine supplementation to improve exercise capacity and performance: a systematic
review and meta-analysis. *British Journal of Sports Medicine* **51**, 658-669 (2017).
6.   G. Aldini, M. Orioli, G. Rossoni, F. Savi, P. Braidotti, G. Vistoli, K. J. Yeum, G. Negrisoli, M. Carini,
The carbonyl scavenger carnosine ameliorates dyslipidaemia and renal function in Zucker
obese rats. *Journal of cellular and molecular medicine* **15**, 1339-1354 (2011).
7.   E. J. Anderson, G. Vistoli, L. A. Katunga, K. Funai, L. Regazzoni, T. B. Monroe, E. Gilardoni, L.
Cannizzaro, M. Colzani, D. De Maddis, A carnosine analog mitigates metabolic disorders of
obesity by reducing carbonyl stress. *The Journal of clinical investigation* **128**, 5280-5293 (2018).
8.   J. Spaas, W. Franssen, C. Keytsman, L. Blancquaert, T. Vanmierlo, J. Bogie, B. Broux, N. Hellings,
J. van Horssen, D. K. Posa, Carnosine quenches the reactive carbonyl acrolein in the central
nervous   system   and   attenuates   autoimmune   neuroinflammation. *Journal of*
neuroinflammation **18**, 1-19 (2021).
9.   I. Everaert, Y. Taes, E. De Heer, H. Baelde, A. Zutinic, B. Yard, S. Sauerhöfer, L. Vanhee, J.
Delanghe, G. Aldini, Low plasma carnosinase activity promotes carnosinemia after carnosine
ingestion in humans. *American Journal of Physiology-Renal Physiology*, (2012).
10.   M. Teufel, V. Saudek, J.-P. Ledig, A. Bernhardt, S. Boularand, A. Carreau, N. J. Cairns, C. Carter,
D. J. Cowley, D. Duverger, Sequence identification and characterization of human carnosinase
and a closely related non-specific dipeptidase. *Journal of Biological Chemistry* **278**, 6521-6531
(2003).
11.   J. Drozak, M. Veiga-da-Cunha, D. Vertommen, V. Stroobant, E. Van Schaftingen, Molecular
identification of carnosine synthase as ATP-grasp domain-containing protein 1 (ATPGD1).
Journal of biological chemistry **285**, 9346-9356 (2010).
12.   J. V. Creighton, L. de Souza Gonçalves, G. G. Artioli, D. Tan, K. J. Elliott-Sale, M. D. Turner, C. L.
Doig, C. Sale, Physiological Roles of Carnosine in Myocardial Function and Health. *Advances in*
Nutrition **13**, 1914-1929 (2022).
13.   G. Aldini, M. Orioli, M. Carini, R. Maffei Facino, Profiling histidine-containing dipeptides in rat
tissues by liquid chromatography/electrospray ionization tandem mass spectrometry. *Journal*
of mass spectrometry **39**, 1417-1428 (2004).

14.   L. Wang-Eckhardt, A. Bastian, T. Bruegmann, P. Sasse, M. Eckhardt, Carnosine synthase
deficiency is compatible with normal skeletal muscle and olfactory function but causes
reduced olfactory sensitivity in aging mice. *Journal of Biological Chemistry*, jbc. RA120. 014188
(2020).
15.   J. Spaas, P. Van Noten, C. Keytsman, I. Nieste, L. Blancquaert, W. Derave, B. O. Eijnde,
Carnosine and skeletal muscle dysfunction in a rodent multiple sclerosis model. *Amino Acids*
53, 1749-1761 (2021).
16.   I. Everaert, S. Stegen, B. Vanheel, Y. Taes, W. Derave, Effect of beta-alanine and carnosine
supplementation on muscle contractility in mice. *Med Sci Sports Exerc* **45**, 43-51 (2013).
17.   L. Luo, W. Ma, K. Liang, Y. Wang, J. Su, R. Liu, T. Liu, N. Shyh-Chang, Spatial metabolomics
reveals skeletal myofiber subtypes. *Science Advances* **9**, eadd0455 (2023).
18.   A. B. Rubenstein, G. R. Smith, U. Raue, G. Begue, K. Minchev, F. Ruf-Zamojski, V. D. Nair, X.
Wang, L. Zhou, E. Zaslavsky, Single-cell transcriptional profiles in human skeletal muscle.
Scientific reports **10**, 1-15 (2020).
19.   N. Nordsborg, M. Mohr, L. D. Pedersen, J. J. Nielsen, H. Langberg, J. Bangsbo, Muscle interstitial
potassium kinetics during intense exhaustive exercise: effect of previous arm exercise.
American Journal of Physiology-Regulatory, Integrative and Comparative Physiology **285**,
R143-R148 (2003).
20.   A. Reddy, L. H. Bozi, O. K. Yaghi, E. L. Mills, H. Xiao, H. E. Nicholson, M. Paschini, J. A. Paulo, R.
Garrity, D. Laznik-Bogoslavski, pH-gated succinate secretion regulates muscle remodeling in
response to exercise. *Cell* **183**, 62-75. e17 (2020).
21.   M. J. Mittenbühler, M. P. Jedrychowski, J. G. Van Vranken, H.-G. Sprenger, S. Wilensky, P. A.
Dumesic, Y. Sun, A. Tartaglia, D. Bogoslavski, A. Mu, Isolation of extracellular fluids reveals
novel secreted bioactive proteins from muscle and fat tissues. *Cell Metabolism*, (2023).
22.   M. A. Kamal, H. Jiang, Y. Hu, R. F. Keep, D. E. Smith, Influence of genetic knockout of Pept2 on
the in vivo disposition of endogenous and exogenous carnosine in wild-type and Pept2 null
mice. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **296**,
R986-R991 (2009).
23.   V. Peters, C. Q. Klessens, H. J. Baelde, B. Singler, K. A. Veraar, A. Zutinic, J. Drozak, J. Zschocke,
C. P. Schmitt, E. de Heer, Intrinsic carnosine metabolism in the human kidney. *Amino acids* **47**,
2541-2550 (2015).
24.   T. Weigand, F. Colbatzky, T. Pfeffer, S. F. Garbade, K. Klingbeil, F. Colbatzky, M. Becker, J.
Zemva, R. Bulkescher, R. Schürfeld, A global Cndp1-knock-out selectively increases renal
carnosine and anserine concentrations in an age-and gender-specific manner in mice.
International journal of molecular sciences **21**, 4887 (2020).
25.   E. Heidenreich, T. Pfeffer, T. Kracke, N. Mechtel, P. Nawroth, G. F. Hoffmann, C. P. Schmitt, R.
Hell, G. Poschet, V. Peters, A Novel UPLC-MS/MS Method Identifies Organ-Specific Dipeptide
Profiles. *International journal of molecular sciences* **22**, 9979 (2021).
26.   D. Abraham, J. J. Pisano, S. Udenfriend, The distribution of homocarnosine in mammals.
Archives of biochemistry and biophysics **99**, 210-213 (1962).
27.   S. J. Kish, T. L. Perry, S. Hansen, Regional distribution of homocarnosine, homocarnosine-
carnosine synthetase and homocarnosinase in human brain. *Journal of neurochemistry* **32**,
1629-1636 (1979).
28.   W. Derave, I. Everaert, S. Beeckman, A. Baguet, Muscle carnosine metabolism and β-alanine
supplementation in relation to exercise and training. *Sports medicine* **40**, 247-263 (2010).
29.   A. Mannion, P. Jakeman, M. Dunnett, R. Harris, P. Willan, Carnosine and anserine
concentrations in the quadriceps femoris muscle of healthy humans. *European journal of*
applied physiology and occupational physiology **64**, 47-50 (1992).
30.   R. C. Harris, M. Dunnett, P. L. Greenhaff, Carnosine and taurine contents in individual fibres of
human vastus lateralis muscle. *Journal of Sports Sciences* **16**, 639-643 (1998).
31.   A. Baguet, I. Everaert, H. De Naeyer, H. Reyngoudt, S. Stegen, S. Beeckman, E. Achten, L.
Vanhee, A. Volkaert, M. Petrovic, Effects of sprint training combined with vegetarian or mixed

diet on muscle carnosine content and buffering capacity. *European journal of applied*
physiology **111**, 2571-2580 (2011).

32. W. Zhou, S. Yang, P. G. Wang. (Future Science, 2017), vol. 9, pp. 1839-1844.

33. F. L. Margolis, Carnosine in the primary olfactory pathway. *Science* **184**, 909-911 (1974).

34. E. Mahootchi, S. Cannon Homaei, R. Kleppe, I. Winge, T.-A. Hegvik, R. Megias-Perez, C. Totland,
F. Mogavero, A. Baumann, J. C. Glennon, GADL1 is a multifunctional decarboxylase with tissue-
specific roles in β-alanine and carnosine production. *Science Advances* **6**, eabb3713 (2020).

35. Y. Zhang, K. Chen, S. A. Sloan, M. L. Bennett, A. R. Scholze, S. O'Keeffe, H. P. Phatnani, P.
Guarnieri, C. Caneda, N. Ruderisch, An RNA-sequencing transcriptome and splicing database
of glia, neurons, and vascular cells of the cerebral cortex. *Journal of Neuroscience* **34**, 11929-
11947 (2014).

36. T. M. Consortium, S. t. w. group, Single-cell transcriptomics of 20 mouse organs creates a
Tabula Muris. *Nature* **562**, 367-372 (2018).

37. J. J. O'Dowd, D. J. Robins, D. J. Miller, Detection, characterisation, and quantification of
carnosine and other histidyl derivatives in cardiac and skeletal muscle. *Biochimica et*
Biophysica Acta (BBA)-General Subjects **967**, 241-249 (1988).

38. K. Yan, Z. Mei, J. Zhao, M. A. I. Prodhan, D. Obal, K. Katragadda, B. Doelling, D. Hoetker, D. K.
Posa, L. He, Integrated Multilayer Omics Reveals the Genomic, Proteomic, and Metabolic
Influences of Histidyl Dipeptides on the Heart. *Journal of the American Heart Association* **11**,
e023868 (2022).

39. J. Zhao, D. J. Conklin, Y. Guo, X. Zhang, D. Obal, L. Guo, G. Jagatheesan, K. Katragadda, L. He, X.
Yin, Cardiospecific Overexpression of ATPGD1 (Carnosine Synthase) Increases Histidine
Dipeptide Levels and Prevents Myocardial Ischemia Reperfusion Injury. *Journal of the*
American Heart Association, e015222 (2020).

40. L. de Souza Gonçalves, L. P. Sales, T. R. Saito, J. C. Campos, A. L. Fernandes, J. Natali, L. Jensen,
A. Arnold, L. Ramalho, L. R. G. Bechara, Histidine dipeptides are key regulators of excitation-
contraction coupling in cardiac muscle: Evidence from a novel CARNS1 knockout rat model.
Redox biology **44**, 102016 (2021).

41. P. Swietach, J.-B. Youm, N. Saegusa, C.-H. Leem, K. W. Spitzer, R. D. Vaughan-Jones, Coupled
Ca2+/H+ transport by cytoplasmic buffers regulates local Ca2+ and H+ ion signaling.
Proceedings of the National Academy of Sciences **110**, E2064-E2073 (2013).

42. P. Swietach, C. H. Leem, K. W. Spitzer, R. D. Vaughan-Jones, Pumping Ca2+ up H+ gradients: A
Ca2+–H+ exchanger without a membrane. *The Journal of physiology* **592**, 3179-3188 (2014).

43. M. Gardner, K. M. Illingworth, J. Kelleher, D. Wood, Intestinal absorption of the intact peptide
carnosine in man, and comparison with intestinal permeability to lactulose. *The Journal of*
physiology **439**, 411-422 (1991).

44. R. C. Harris, M. Tallon, M. Dunnett, L. Boobis, J. Coakley, H. J. Kim, J. L. Fallowfield, C. Hill, C.
Sale, J. A. Wise, The absorption of orally supplied β-alanine and its effect on muscle carnosine
synthesis in human vastus lateralis. *Amino acids* **30**, 279-289 (2006).

45. A. Baguet, I. Everaert, B. Yard, V. Peters, J. Zschocke, A. Zutinic, E. De Heer, T. Podgórski, K.
Domaszewska, W. Derave, Does low serum carnosinase activity favour high-intensity exercise
capacity? *American Journal of Physiology-Heart and Circulatory Physiology*, (2014).

46. V. K. Pandya, B. Sonwane, R. Rathore, A. Unnikrishnan, S. Kumaran, M. J. Kulkarni,
Development of multiple reaction monitoring assay for quantification of carnosine in human
plasma. *RSC advances* **10**, 763-769 (2020).

47. S. de Jager, L. Blancquaert, T. Van der Stede, E. Lievens, S. De Baere, S. Croubels, E. Gilardoni,
L. G. Regazzoni, G. Aldini, J. G. Bourgois, The ergogenic effect of acute carnosine and anserine
supplementation: dosing, timing, and underlying mechanism. *Journal of the International*
Society of Sports Nutrition **19**, 70-91 (2022).

48. H. Oppermann, S. Elsel, C. Birkemeyer, J. Meixensberger, F. Gaunitz, Erythrocytes prevent
degradation of carnosine by human serum carnosinase. *International journal of molecular*
sciences **22**, 12802 (2021).

49. K. Nagai, A. Niijima, T. Yamano, H. Otani, N. Okumra, N. Tsuruoka, M. Nakai, Y. Kiso, Possible
role of L-carnosine in the regulation of blood glucose through controlling autonomic nerves.
Experimental biology and medicine **228**, 1138-1145 (2003).
50. M. C. K. Severinsen, B. K. Pedersen, Muscle–organ crosstalk: the emerging roles of myokines.
Endocrine reviews **41**, 594-609 (2020).
51. B. K. Pedersen, M. A. Febbraio, Muscles, exercise and obesity: skeletal muscle as a secretory
organ. *Nature Reviews Endocrinology* **8**, 457-465 (2012).
52. R. M. Murphy, M. J. Watt, M. A. Febbraio, Metabolic communication during exercise. *Nature*
metabolism **2**, 805-816 (2020).
53. M. E. Nelson, S. Madsen, K. C. Cooke, A. M. Fritzen, I. H. Thorius, S. W. Masson, L. Carroll, F. C.
Weiss, M. M. Seldin, M. Potter, Systems-level analysis of insulin action in mouse strains
provides insight into tissue-and pathway-specific interactions that drive insulin resistance. *Cell*
Metabolism **34**, 227-239. e226 (2022).
54. H. Ihara, Y. Kakihana, A. Yamakage, K. Kai, T. Shibata, M. Nishida, K.-i. Yamada, K. Uchida, 2-
Oxo-histidine–containing dipeptides are functional oxidation products. *Journal of Biological*
Chemistry **294**, 1279-1289 (2019).
55. S. P. Baba, J. D. Hoetker, M. Merchant, J. B. Klein, J. Cai, O. A. Barski, D. J. Conklin, A. Bhatnagar,
Role of aldose reductase in the metabolism and detoxification of carnosine-acrolein
conjugates. *Journal of Biological Chemistry* **288**, 28163-28179 (2013).
56. V. H. Carvalho, A. H. Oliveira, L. F. de Oliveira, R. P. da Silva, P. Di Mascio, B. Gualano, G. G.
Artioli, M. H. Medeiros, Exercise and β-alanine supplementation on carnosine-acrolein adduct
in skeletal muscle. *Redox biology* **18**, 222-228 (2018).
57. L. Wang-Eckhardt, I. Becker, Y. Wang, J. Yuan, M. Eckhardt, Absence of endogenous carnosine
synthesis does not increase protein carbonylation and advanced lipoxidation end products in
brain, kidney or muscle. *Amino Acids*, 1-11 (2022).
58. T. Van der Stede, L. Blancquaert, F. Stassen, I. Everaert, R. Van Thienen, C. Vervaet, L.
Gliemann, Y. Hellsten, W. Derave, Histamine H1 and H2 receptors are essential transducers of
the integrative exercise training response in humans. *Science Advances* **7**, eabf2856 (2021).
59. K. Verboven, K. Wouters, K. Gaens, D. Hansen, M. Bijnen, S. Wetzels, C. D. Stehouwer, G. H.
Goossens, C. G. Schalkwijk, E. E. Blaak, Abdominal subcutaneous and visceral adipocyte size,
lipolysis and inflammation relate to insulin resistance in male obese humans. *Scientific reports*
8, 1-8 (2018).
60. F. Van de Velde, D. M. Ouwens, A.-H. Batens, Y. Van Nieuwenhove, B. Lapauw, Divergent
dynamics in systemic and tissue-specific metabolic and inflammatory responses during weight
loss in subjects with obesity. *Cytokine* **144**, 155587 (2021).
61. R. Chaleckis, I. Murakami, J. Takada, H. Kondoh, M. Yanagida, Individual variability in human
blood metabolites identifies age-related differences. *Proceedings of the National Academy of*
Sciences **113**, 4252-4259 (2016).
62. J. De Brandt, W. Derave, F. Vandenabeele, P. Pomiès, L. Blancquaert, C. Keytsman, M. S.
Barusso-Grüninger, F. F. de Lima, M. Hayot, M. A. Spruit, Efficacy of 12 weeks oral beta-alanine
supplementation in patients with chronic obstructive pulmonary disease: a double-blind,
randomized, placebo-controlled trial. *Journal of cachexia, sarcopenia and muscle* **13**, 2361-
2372 (2022).
63. L. Gliemann, N. Rytter, P. Piil, J. Nilton, T. Lind, M. Nyberg, M. Cocks, Y. Hellsten, The
endothelial mechanotransduction protein platelet endothelial cell adhesion molecule-1 is
influenced by aging and exercise training in human skeletal muscle. *Frontiers in physiology* **9**,
1807 (2018).
64. L. Gliemann, N. Rytter, A. Tamariz-Ellemann, J. Egelund, N. Brandt, H. H. Carter, Y. Hellsten,
Lifelong Physical Activity Determines Vascular Function in Late Postmenopausal Women.
Medicine and Science in Sports and Exercise **52**, 627-636 (2020).

65.   L. Olsen, B. Hoier, C. Hansen, M. Leinum, H. Carter, T. Jorgensen, J. Bangsbo, Y. Hellsten,
877   Angiogenic potential is reduced in skeletal muscle of aged women. *The Journal of Physiology*
878   **598**, 5149-5164 (2020).
879   66.   C. Hansen, S. Møller, T. Ehlers, K. A. Wickham, J. Bangsbo, L. Gliemann, Y. Hellsten, Redox
880   balance in human skeletal muscle-derived endothelial cells-Effect of exercise training. *Free*
881   *Radical Biology and Medicine* **179**, 144-155 (2022).
882   67.   B. Høier, K. Olsen, M. Nyberg, J. Bangsbo, Y. Hellsten, Contraction-induced secretion of VEGF
883   from skeletal muscle cells is mediated by adenosine. *American Journal of Physiology-Heart and*
884   *Circulatory Physiology* **299**, H857-H862 (2010).
885   68.   C. G. Perry, D. A. Kane, C.-T. Lin, R. Kozy, B. L. Cathey, D. S. Lark, C. L. Kane, P. M. Brophy, T. P.
886   Gavin, E. J. Anderson, Inhibiting myosin-ATPase reveals a dynamic range of mitochondrial
887   respiratory control in skeletal muscle. *Biochemical Journal* **437**, 215-222 (2011).
888   69.   S. de Jager, S. Van Damme, S. De Baere, S. Croubels, R. Jäger, M. Purpura, E. Lievens, J. G.
889   Bourgois, W. Derave, No Effect of Acute Balenine Supplementation on Maximal and
890   Submaximal Exercise Performance in Recreational Cyclists. *International Journal of Sport*
891   *Nutrition and Exercise Metabolism* **1**, 1-9 (2023).
892   70.   M. R. Larsen, D. E. Steenberg, J. B. Birk, K. A. Sjøberg, B. Kiens, E. A. Richter, J. F. Wojtaszewski,
893   The insulin-sensitizing effect of a single exercise bout is similar in type I and type II human
894   muscle fibres. *The Journal of Physiology* **598**, 5687-5699 (2020).
895   71.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
896   seq data with DESeq2. *Genome biology* **15**, 1-21 (2014).
897

Acknowledgements

For human brain tissues, all material has been collected from donors for or from whom a written informed consent for a brain autopsy and the use of the material and clinical information for research purposes had been obtained by the Netherlands Brain Bank (NBB). The experiment protocols and methods used for analysing brain samples were conducted with the approval of the NBB and the Medical Ethical Committee of Hasselt University, and carried out according to institutional guidelines. All HCDs analyses were performed using an UHPLC-MS/MS instrument part of the Ghent University MSsmall Expertise Centre for advanced mass spectrometry analysis of small organic molecules. Also, the University Biobank Limburg (UBiLim) is acknowledged for providing storage and release of some human biological material used in this publication. Figure 7D, 7E, 7F and FG (experimental setups) were created in BioRender. The technical assistance of Jens Jung Nielsen, Anneke Volkaert, Thomas Ehlers, Nicklas Frisch and Josephine Rol is greatly appreciated.

Funding

Research Foundation-Flanders – FWO 11B4220N (TVDS)

Research Foundation-Flanders – FWO 1138520N (JS)

Research Foundation-Flanders – FWO 11C0421N (SDJ)

Research Foundation-Flanders – FWO V433222N (TVDS)

Research Foundation-Flanders – FWO G080321N (WD)

Author contributions

TVDS, JS, SDJ and WD designed the study.

TVDS, JS, SDJ, BVer and CH performed the experiments and/or biochemical analyses.

TVDS, JS, SDJ, JDB, RVT, KV, DH, TB, BL, CVP, KD, BVan and LG contributed to tissue collection.

SC, BO, LG, YH and WD supervised the study.

TVDS, JS, SDJ and WD analyzed the data.

TVDS, JS and WD drafted the manuscript.

All authors approved the final version of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All data needed to evaluate the conclusions in the paper are present in the manuscript and supplementary materials. Additional data related to this paper can be obtained upon reasonable request to the author.