Hemi Manganese Exchangers 1 and 2 enable manganese import at the plasma membrane

in cyanobacteria

Mara Reis[1,2*], Fabian Brandenburg[1,3,*], Michael Knopp[4], Samantha Flachbart[1], Andrea

Bräutigam[2], Sabine Metzger[5], Sven B. Gould[4], Marion Eisenhut[1,2]

[1]   Institute of Plant Biochemistry, Cluster of Excellence on Plant Sciences (CEPLAS),

Heinrich-Heine University, Düsseldorf, Germany.

[2]   Computational Biology, Faculty of Biology, CeBiTec, Bielefeld University, Bielefeld,

Germany.

[3]   Department of Solar Materials, Helmholtz-Centre for Environmental Research - UFZ,

Leipzig, Germany.

[4]   Institute for Molecular Evolution, Heinrich-Heine University, Düsseldorf, Germany.

[5]   MS-Platform, Cluster of Excellence on Plant Sciences (CEPLAS), Botanical Institute,

University of Cologne, Cologne, Germany.

* These authors contributed equally.

Correspondence

Marion Eisenhut,

Email: marion.eisenhut@uni-bielefeld.de

Funding information

Deutsche Forschungsgemeinschaft (DFG), Grant/Award Numbers:

EI 945/3-2, CRC1535 - Project ID 458090666, MAdLand SPP2237 – Project ID 440043394,

CRC1208 - Project ID267205415

ABSTRACT

Manganese (Mn) is key to oxygenic photosynthesis as it catalyzes the splitting of water in photosystem II and functions as cofactor of multiple enzymes. A single ABC-type transporter, MntCAB, is so far established for the uptake of the metal under limiting conditions in cyanobacteria. It is unknown how Mn is imported under replete conditions. We identified two proteins in the cyanobacterium *Synechocystis* sp. PCC 6803, which are homologous to the unknown protein family 0016 (UPF0016) member manganese exporter (Mnx). In contrast to Mnx, which consists of six transmembrane domains, the new candidate proteins contain three transmembrane domains. Hence, we named them hemi manganese exchangers (Hmx) 1 and 2. Knock-out mutants in *hmx1* and/or *hmx2* showed sensitivity toward low Mn supplementation, and reduced intracellular Mn pools. Additional deletion of *mntC* hindered the cells to thrive unless the medium was supplemented with Mn to compensate for the depletion of their intracellular Mn pool. In accordance with the observed localization of Hmx1 and Hmx2 in the plasma membrane, we postulate a Mn uptake function for heteromeric Hmx1/2 across the plasma membrane under a wide range of Mn concentrations and a supporting role for the MntCAB system under Mn-limiting conditions. On the basis of their phylogenies, we propose that Hmx1 and Hmx2 are the ancestral progenitors of eukaryote-type UPF0016 proteins with six transmembrane domains. The Mn transport function of Hmx1/2 underscores this as a fundamental and ancient feature of the UPF0016 family. Potentially, Hmx1 and Hmx2 coevolved with the internalization of the oxygen-evolving complex.

INTRODUCTION

The transition metal manganese (Mn) is of crucial importance across all kingdoms of life. Mn associates with proteins, metabolites, and nucleic acids (reviewed in Bosma et al., 2021), fulfills an activating role for several proteins, and serves as a direct cofactor of a variety of enzymes (Hänsch & Mendel, 2009; Schmidt & Husted, 2019). Mn-dependent superoxide dismutase is central in scavenging of reactive oxygen species, and also Mn^{2+}-ions associated with uncharacterized small molecules are suggested to defend against oxidative stress (Jakubovics & Jenkinson, 2001; Marschner, 2012). Diverse enzymes of carbon metabolism (Marschner, 2012), glucosyltransferases (Breton et al., 2006), oxalate decarboxylase (Tottey et al., 2008), and enzymes involved in isoprenoid and amino acid biosynthesis are strictly Mn-requiring (Schmidt et al., 2020). In oxygenic photosynthetic organisms, Mn is of even higher importance, since it functions as the inorganic catalyst in the oxidation of water. Together with oxygen and calcium atoms, Mn ions form the Mn_4O_5Ca cluster and catalyze the light-driven splitting of water molecules into electrons, protons, and molecular oxygen, as part of photosystem II (Nelson & Junge, 2015). Consequently, photoautotrophic cyanobacteria have a 100-fold higher demand for Mn than non-photosynthetic bacteria (Keren et al., 2002).

Critical for proper provision of the cell with Mn is a sufficient uptake of the metal from the environment into the cytoplasm, and further integration into the target proteins and molecules. In bacteria, two different types of Mn importers are known: the ATP-binding cassette (ABC) type transporter and the Natural Resistance-Associated Macrophage Protein (NRAMP) type transporter (Bosma et al., 2021). Though NRAMP proteins, such as MntH (Kehres et al., 2000), are encoded by cyanobacterial genomes, their specific relevance in metal transport has not been investigated so far. In contrast, the ABC-type Mn transporter MntCAB (Bartsevich & Pakrasi, 1995; Bartsevich a& Pakrasi, 1996) is well studied in the model cyanobacterium *Synechocystis* sp. PCC 6803 (*Synechocystis*). Expression of the *mntCAB*-operon occurs under the control of the two-component system ManS/ManR (Ogawa et al., 2002; Yamaguchi et al.,

2002). Sensing of Mn limitation induces the expression and installation of the high-affinity

transporter in the plasma membrane and enables cell growth under such conditions (Bartsevich

& Pakrasi, 1995). Intriguingly, a deletion mutant in *mntC* was still able to accumulate Mn inside

the cells when (sub)micromolar amounts of Mn were present in the growth medium. This hinted

at the presence of a second, yet unidentified Mn uptake system in *Synechocystis* (Bartsevich a&

Pakrasi, 1996).

Another class of Mn transporter occurring in cyanobacteria and also eukaryotes, was

only recently identified. The cyanobacterial founding member is the Mn exporter (Mnx)

(Brandenburg et al., 2017b), also named SynPAM71 (Gandini et al., 2017). Mnx resides in the

thylakoid membrane and shuttles Mn from the cytoplasm into the thylakoid lumen to (i) assist in

Mn provision to PSII and (ii) prevent detrimental overaccumulation of Mn in the cytoplasm

(Brandenburg et al., 2017b). Mnx belongs to the Uncharacterized Protein Family (UPF) 0016.

Most members of the family consist of two repetitions of a domain that contains a conserved

ExGD motif in the first of three predicted transmembrane domains (TMDs), making them 6-TMD

proteins (Demaegd et al., 2014). Transport of Mn was demonstrated for the UPF0016 members

PAM71 (Schneider et al., 2016), CMT1 (Eisenhut et al., 2018; Zhang et al., 2018), PML1

(Hoecker et al., 2020), PML2 (Hoecker et al., 2020), and PML3 (Yang et al., 2021; He et al.,

2022) in *Arabidopsis thaliana* (Arabidopsis). In other analyses, also Ca was determined to be a

transport substrate (Wang et al., 2016; Frank et al., 2019).

Here, we identified and characterized two additional members of the UPF0016 in

Synechocystis. Hemi manganese exchanger (Hmx) 1 and 2 are each 3-TMD proteins and jointly

function as a constitutive Mn importer in the plasma membrane of *Synechocystis.* They likely

represent the ancestral state from which the 6-TMD proteins, such as Mnx or PAM71, which is

the only version of UPF0016 proteins occurring in eukaryotes, evolved through gene fusion.

MATERIAL AND METHODS

Synechocystis Strains and Growth Conditions

The glucose-tolerant Japanese strain of *Synechocystis* sp. PCC 6803 obtained from Martin

Hagemann (University of Rostock, Germany) served as wild type (WT). Axenic cultures were

routinously grown in BG11 medium adjusted with 20 mM HEPES-KOH to pH 7.5 (Rippka et al.,

1979) in a shaker at 30 °C and 200 rpm, illuminated with 100 µmol photons m^{-2} s^{-1} constant

white light. Growth medium of the mutant lines was supplemented with appropriate antibiotics,

50 µg mL^{-1} kanamycin (Km), 20 µg mL^{-1} spectinomycin (Sp), or 12.5 µg mL^{-1} gentamycin (Gm).

Generation of *Synechocystis* Knockout and Double-Knockout Lines

The Δ*hmx1* knockout mutant was generated by introduction of the plasmid pUC-4S including a

Sp resistance cassette after HincII digestion into the NaeI restriction site of the PCR-amplified

(primers FB69 and FB70; Table S1) open reading frame of *slr*1170. The cloning vector pJET1.2

(ThermoFisher) served as vector backbone. To generate the knockout construct for *hmx2,* the

open reading frame for *ssr*1558 including upstream and downstream sequences was amplified

using the primers ME368 and ME369 (Table S1). The Km resistance cassette derived from the

plasmid pUC-4K after digestion with HincII was inserted into the SmaI restriction site of *hmx2*.

The construct for generating a Δ*mntC* mutant was produced by amplification of the open

reading frame *sll*1598 using the primers ME163 and ME164 (Table S1). The Sp resistance

cassette from pUC-4S was introduced into the MscI site. All restriction enzymes were

purchased from New England Biolabs, USA.

Transformation, selection on Sp- or Km-containing BG11 plates and segregation of independent

clones was verified by PCR analysis as described in (Eisenhut et al., 2006). The Δ*hmx1*/Δ*hmx2,*

and Δ*hmx2*/Δ*mntC* double knockout mutants were generated by transformation of single

mutants with the desired knockout-construct, following the same protocol.

Subcellular localization experiments

For the generation of CFP-fusion protein constructs, the CFP-coding sequence and Gm resistance cassette from a vector as described in (Heinz et al., 2016) was utilized. Using the primers FB105 and FB106 (Table S1) XhoI and NheI restriction sites were added by PCR to the *hmx1* open reading frame including its 800 bp upstream region. Similarly, the 800 bp downstream region of *hmx1* was PCR amplified with EcoRI restriction sites added to both ends (primers FB107 and FB108; Table S1). Restriction digest and T4 DNA Ligase (all NEB enzymes) were used to first clone the 3'-region of *hmx1* downstream of a CFP and Gm resistance cassette, before *hmx1* including the 5'-region of *hmx1* was cloned upstream of the *cfp* gene. A GSGSG peptide linker separates the gene of interest and *hmx1* to allow proper folding of both proteins. As vector backbone, pJET1.2 (ThermoFisher) was used. A *hmx2:cfp* fusion was generated in the same way using primers FB109, FB110 and FB111, FB112 (Table S1). *Synechocystis* Δ*hmx1* and Δ*hmx2* cells were transformed with the constructs as described above, using Gm for selection. The antibiotics Km and Sp used for selection of the knockout of *hmx1* and *hmx2*, respectively, were omitted from the medium to allow replacement of the knockout alleles by the expression alleles. Successful transformation and segregation was verified by PCR using primers FB69/FB70 (*hmx1:cfp*) or ME368/ME369 (*hmx2:cfp*). For primer sequences see Supplemental Table S1.

For imaging, the cells were immobilized on microscopic glass slides by a thin layer of solid BG11 medium (1:1 mixture of 2-fold concentrated BG11 medium, with 24 mM sodium thiosulfate added and 3% [w/v] bacto agar). A Leica TCS SP8 STED 3X microscope with a HC PL APO CS2 100x/1.40 OIL objective was used. An argon laser at 488 nm and 70 W output intensity was used for excitation. Emission was detected using Leica HyD hybrid detectors from 470-530 nm (CFP) and 660-700 nm (chlorophyll). Microscopy was performed at the Center of Advanced Imaging, Heinrich-Heine University Düsseldorf.

Drop Tests

The effect of varying amounts of $MnCl_2$ on the different lines was tested on solid BG11 medium.

Cultures were grown until mid-log phase and 5 d starved for Mn by cultivating in BG11 medium

without $MnCl_2$ (BG11 -Mn). Then, 2 µL of culture with an OD_{750} of 0.25 and subsequent 1:10,

1:100, and 1:1000 dilutions were spotted onto agar plates (BG11, pH 7.5 with 24 mM sodium

thiosulfate added; solidified with 1.5% [w/v] bacto agar). The plates were supplemented with

$MnCl_2$ as indicated (1x $MnCl_2$ = 9 µM $MnCl_2$) and did not contain antibiotics. Plates were

incubated under continuous white-light illumination of 100 µmol photons m^{-2} s^{-1} at 30 °C for 5 d.

ICP-MS Measurements

Cells were washed with EDTA (20 mM HEPES-KOH, pH 7.5, and 5 mM EDTA) (Keren et al.,

2002) before and after pre-cultivation under Mn-limiting conditions (BG11 -Mn) for 5 d, to ensure

similar intracellular Mn concentrations in all lines. Before the experiment, cells were adjusted to

an OD_{750} of 0.8 and treated with $MnCl_2$ concentrations as given in the respective experiments.

Before the experiment and 4 h after $MnCl_2$ treatment, samples were taken and washed as

described in (Brandenburg et al., 2017a). In short, to determine the total content of Mn in a

sample, cells were washed two times with ice-cold HEPES (20 mM HEPES-KOH, pH 7.5) to

preserve the periplasmic Mn storage. Additionally, the samples were washed two times with 4

mL Milli-Q grade (18 MΩ cm) water before further processing. To release the periplasmic Mn

pool, a second set of samples was washed initially two times with HEPES containing 5 mM

EDTA and subsequentially washed two times with Milli-Q grade water. The washed samples

were re-suspended afterwards in 0.4 mL 65% nitric acid and digested for 3 h at 70 °C. The

digested samples were diluted to ~4 % nitric acid with 6.5 mL Milli-Q grade water. Elemental

composition of the samples was determined by ICP-MS (Agilent 7700) at the CEPLAS Plant

Metabolism and Metabolomics Facility, University of Cologne. The cell numbers of the samples

were estimated using a cell counter (Beckman Coulter Z2).

Sequence Analysis and Identification of Candidate Genes

Proteins of the UPF0016 were identified using Pfam (http://pfam.xfam.org) (Finn et al., 2016)

and BlastP (https://blast.ncbi.nlm.nih.gov/Blast.cgi) analysis (Altschul et al., 1990). DNA and

protein sequences were obtained from the genome database CyanoBase

(http://genome.microbedb.jp/cyanobase).   Clustal   Omega

(https://www.ebi.ac.uk/Tools/msa/clustalo) was used for sequence alignment (Sievers et al.,

2011).

Expression Analysis

The coefficient of variation (standard deviation divided by the mean to account and correct for

transcript abundance level) was calculated for all genes in *Synechocystis* sp. PCC 6803 (Wulf

et al., 2024) by mapping 46 publicly available expression datasets (SRX10706739,

SRX10706741, SRX10706742, SRX10706743, SRX10706745, SRX10706747, SRX10706750,

SRX3580390, SRX5980004, SRX5980006, SRX5980011, SRX5980016, SRX5980017,

SRX5980018, SRX5980022, SRX7098395, SRX8102890, SRX8102891, SRX8102892,

SRX8102893, SRX8102894, SRX8102895, SRX8102896, SRX8102897, SRX8102898,

SRX8102899, SRX8102900, SRX8102901, SRX8844415, SRX8844416, SRX8844417,

SRX8844420, SRX8844421, SRX8844428, ERX3642241, ERX3642242, ERX3642243,

ERX3642245, ERX3642247, ERX3642249, ERX3642250, ERX3642253, ERX3642254,

SRX7873832, SRX7873837, SRX9234496) using kallisto version 0.44 in stranded mode onto

the transcriptome of *Synechocystis* sp. PCC 6803 (accession number NC_000911, downloaded

on January 03, 2023, from NCBI https://www.ncbi.nlm.nih.gov/nuccore/NC_000911). Transcript

per million values were exported to Excel and mean and standard deviation calculated using

AVERAGE and ST.DEVP.

Phylogenetic Analysis

A database of 5,655 complete prokaryotic genomes of the RefSeq database (O'Leary et al., 2016) was search via diamond blastp (Buchfink et al., 2015) using the "--very-sensitive" option. Since Mnx, Hmx1, and Hmx2 share significant sequence identity, exact pairwise global alignments were produced to determine whether queries with multiple hit are a Hmx protein or the fusion Mnx protein (Rice et al., 2000). Homologs were visualized using a presence-and-absence matrix, color-coding their pairwise local identity. The same set of seed sequences was used to search for eukaryotic homologs within a database of 150 eukaryotes (Ku et al., 2015). All significant hits with a maximum e-value of 1×10^{-10} and at least 25% sequence identity were retained for further analysis (Table S2).

RESULTS

The genome of *Synechocystis* encodes two half-sized proteins with respect to UPF0016 homologs

Transporters of the Mnx family (UPF0016) are critical for proper Mn distribution within the cyanobacterial (Brandenburg et al., 2017b; Gandini et al., 2017) and plant cell (Schneider et al., 2016; Eisenhut et al., 2018; Zhang et al., 2018; Hoecker et al., 2020; Yang et al., 2021; He et al., 2022). While we identified two plastid-targeted Mnx proteins for all members of the green lineage (Schneider et al., 2016; Hoecker et al., 2017; Eisenhut et al., 2018), in cyanobacteria just a single homolog was found. Closer inspection of cyanobacterial genomes, however, revealed the existence of two additional genes encoding more distantly related UPF0016 proteins. These proteins are half-size versions of the canonical Mnx proteins and contain only one cluster consisting of three TMDs including the conserved ExGD motif, of which two in tandem are characteristic for the UPF0016 family (Figure 1A). Accordingly, we named these

proteins Hemi manganese exchanger (Hmx) 1 and 2. In *Synechocystis,* Hmx1 is encoded by

*slr*1170 and Hmx2 by *ssr*1558. A BlastP run shows that Slr1170 and Ssr1558 have a 35% and

31% sequence identity to *Synechocystis* Mnx, respectively. Furthermore, we found Hmx1/Hmx2

orthologues to be encoded by neighboring genes in the majority (90 %) of the 172

cyanobacterial species we analyzed, while in only a few (10%) species, such as in

Synechocystis, they were dispersed across the genome (Table S3). In those cases, where

hmx1 and *hmx2* genes are organized as a transcriptional unit, we frequently detected a third

gene being part of the operon (Figure 1B). They are members of the UPF0153, yet their function

remains unknown. One exception is *psb28-2* of *Cyanothece* sp. ATCC 51142, which was

identified upstream of the *slr*1170 orthologue. Psb28-2 is involved in PSII and chlorophyll

biosynthesis (Dobáková et al., 2009; Nowaczyk et al., 2012).

Deletion of *hmx1* or *hmx2* impairs intracellular Mn accumulation

Due to the frequent genomic appearance of the genes in transcriptional units, we hypothesized

that Hmx1 and Hmx2 interact and function in Mn transport, as other members of the Mnx family

do. To test this hypothesis, single and double mutants of *hmx1* or/and *hmx2* were generated by

insertional inactivation in *Synechocystis*. In the case of *hmx1,* a Sp resistance cassette and in

the case of *hmx2,* a Km resistance cassette was inserted into the coding region leading to an

interruption of the respective reading frame (Figure 2A). As verified by PCR, we obtained fully

segregated mutant lines for Δ*hmx1,* Δ*hmx2,* and Δ*hmx1*/Δ*hmx2* (Figure 2B). All lines were

tested for their susceptibility toward different Mn concentrations (Figure 2C). In contrast to the

Δ*mnx* mutant, which is sensitive toward elevated Mn supply (Brandenburg et al., 2017b; Gandini

et al., 2017), mutants in *hmx* genes were not susceptible to high but low Mn concentrations in

the medium. Without any and with standard (1x, 9 μM $MnCl_2$) Mn supply, single and double

mutants displayed retarded growth. This impairment was improved by elevated (5x) Mn

concentration in the medium. We also tested the mutants´ sensitivity towards reduced

supplementation with calcium (Ca) and iron (Fe) (Figure S1), since other known Mn

transporters, such as members of the NRAMP, IRT, or the CAX family, are known to also use

these substrates (Socha & Guerinot, 2014). While omitting Fe from the medium fully disabled

growth of all strains, including that of the WT, depleting Ca only did not affect growth of any

strain. Also, all other combinations did not result in different growth behavior of Δ*hmx1* and

Δ*hmx2* mutants in comparison to the WT.

The results support the assumption that Hmx1 and Hmx2 facilitate Mn transport, likely

Mn import with high specificity. We additionally examined the intracellular Mn concentrations

and found that all mutant lines, whether single or double mutant, accumulated significantly less

Mn inside the cell (excluding the periplasmic space) after the addition of 1x $MnCl_2$ (Figure 2D).

The equal growth and accumulation phenotype of single and double mutants indicated that

Hmx1 and Hmx2 need to be both active and probably functionally assemble as heteromers *in*

vivo.

Hmx1 and Hmx2 reside in the plasma membrane

The low Mn sensitive phenotype and the reduced intracellular Mn pools of the mutants pointed

toward an import function for Hmx1/2, prompting the investigation of the subcellular localization

of the proteins. We generated mutant lines expressing either Hmx1 (*hmx1:cfp*) or Hmx2

(*hmx2:cfp*) fused to a C-terminal cyan fluorescent protein (CFP). The fusion constructs were

designed in such a way that they replaced the original gene. That is, *hmx1:cfp* and *hmx2:cfp*

were introduced into the native genomic context under control of the respective native promoter.

To use these constructs as complementation lines and thus demonstrate that the observed

phenotypes are only due to the deletion of the specific gene, we transformed Δ*hmx1* and Δ*hmx2*

single-mutant cells. Full segregation of the expression lines demonstrated that the fused version

fully displaced the knockout alleles (Figure S2A). In comparison to Δ*hmx1/*Δ*hmx2*, the *hmx1:cfp*

and *hmx2:cfp* lines are not sensitive toward low Mn concentrations (Figure S2B). On the one

hand this demonstrates that the CFP fusion proteins functionally assembled in the correct native

cellular site, and on the other hand that the expression lines could be used to complement the

mutants. Confocal fluorescent microscopy (Figure 3) revealed that the CFP signals of both

Hmx1 and Hmx2 did not fully overlap with the signal of the chlorophyll autofluorescence, but

were rather oriented toward the outside of the cell. This suggests that both Hmx1 and Hmx2

reside in the plasma membrane. Furthermore, the CFP signals had a patchy, dotted

appearance and showed local maxima at regions of low chlorophyll fluorescence (e.g., Figure

3B: ROI6 and ROI9; Figure 3D: ROI6 and ROI8), which are typical for PSII biogenesis centers

(Heinz et al., 2016).

Double mutants in *hmx2* and *mntC* are not viable under Mn-limiting conditions

Previous work identified MntCAB as the major Mn importer at the plasma membrane under Mn-

limiting conditions (Bartsevich & Pakrasi, 1996). Since in those experiments Mn uptake was not

fully abolished in a Δ*mntC* mutant, the presence of a second high-affinity transport system was

postulated. Hmx1/2 also reside in the plasma membrane and likely facilitate Mn transport. To

provide further evidence for their transporter function, we generated a single mutant in *mntC,*

Δ*mntC* (Figure 4A) and a double mutant with an additional defect in *hmx2*, Δ*mntC/*Δ*hmx2.* In the

Δ*mntC* mutant, zero MntCAB transport activity was observed (Bartsevich & Pakrasi, 1996). After

genotype verification (Figure 4B), the mutant lines were studied for their Mn sensitivity (Figure

4C). As expected, Δ*mntC* showed strong growth retardation only on medium lacking Mn. A

concentration of 0.5x (4.5 μM) $MnCl_2$ in the BG11 medium was sufficient to fully compensate the

phenotype. The Δ*hmx2* mutant underperformed under reduced Mn availability conditions (w/o to

2.5x $MnCl_2$), but could be rescued by elevated (5x) Mn concentrations. The double mutant

mounted the strongest Mn sensitivity phenotype. A $MnCl_2$ concentration below 1x was not

sufficient to allow Δ*mntC/*Δ*hmx2* to grow. However, 5x $MnCl_2$ in the medium fully rescued the

phenotype. Likewise, the determination of intracellular Mn demonstrated that Δ*mntC* and

$\Delta mntC/\Delta hmx2$ contained significantly depleted Mn pools after a 5-day period of Mn limitation

(Figure 4D). Furthermore, the addition of 0.5x $MnCl_2$ resulted in disturbed intracellular

accumulation in all mutant lines, with the most pronounced impairment observed in

$\Delta mntC/\Delta hmx2$. The accumulation was significantly reduced even in comparison to both single

mutants, $\Delta hmx2$ and $\Delta mntC$ (Figure 4D).

hmx1 and hmx2 are constitutively expressed

To test for house-keeping or inducible gene expression, the coefficient of variation was

calculated from 46 publicly available data sets in the short read data archive (Wulf et al., 2024)

and showed that *hmx1* and *hmx2* belong to the transcripts of low variation, whereas *mntCAB*

were highly variable transcripts (Figure 5A). Mn-independent expression of *hmx1* and *hmx2* was

additionally supported by analyzing transcriptional profiles of *Synechocystis* WT cells from Mn

limitation (Sharon et al., 2014) and Mn excess (Reis et al., 2024) experiments. While transcript

abundances of the *mntCAB* operon were elevated under Mn-limiting conditions and reduced

under Mn excess conditions, *hmx1* and *hmx2* expression was not significantly changed under

either condition (Figure 5B).

Hmx1 and Hmx2 are conserved across and are almost exclusive to cyanobacteria

To determine the evolutionary origin of the investigated Mn transporters, a database of 5,455

bacterial and 212 archaeal complete genomes was screened via diamond BLASTp. Since Mnx,

Hmx1, and Hmx2 share substantial sequence identity, additional pairwise global alignments

were used to determine the annotation of all hits. The gene distribution suggests a

cyanobacterial origin of Hmx1 and Hmx2, since Hmx-type homologs with three TMDs are rarely

found outside of cyanobacteria, independent of the chosen e-value cutoff (Figure 6, Figure S3,

Table S2). Exceptions are single Hmx1/Hmx2 homologs in *e.g.*, *Desulfovibrio desulfuricans*

ND132 or *Desulfovibrio magneticus* RS-1. In contrast, Mnx homologs with six TMDs are present

in a wider range of Proteobacteria, Actinobacteria, Chlorobi, Clostridia, and one member of the

Negativicutes. Eukaryotic genomes appear to encode only Mnx homologs having six TMDs

conserved (Figure S4, Table S2). Focusing on the cyanobacterial genomes (Figure 6), it

becomes apparent that some strains contain all three homologs (Hmx1, Hmx2, and Mnx; *e.g.*,

Synechocystis sp. PCC 6803), some only the pair of Hmx1 and Hmx2 (*e.g.*, *Synechococcus*

WH 7803), and a few only one Hmx homolog (*e.g.*, *Gloeobacter violaceus* PCC7421).

Cyanobacterial genomes hence offer a window into the stages of Hmx/Mnx evolution, ranging

from the single half-sized, 3-TMD homolog of the UPF0016 family in an early branching

cyanobacterium, such as *Gloeobacter*, via two 3-TMD Hmx homologs in the majority of

branching cyanobacteria, to the 6-TMD version of UPF0016 homologs in about half of them.

DISCUSSION

Members of the UPF0016 family have recently emerged as important representatives of

secondary transporters with Ca^{2+} and, most importantly, Mn^{2+} as substrates (Stribny et al.,

2020). Typically, the 6-TMD transporters occur in a wide range of eukaryotic groups (Figure S4)

and also cyanobacteria (Figure 6). In eukaryotes the transporters serve in Ca^{2+} and Mn^{2+} uptake

into the Golgi apparatus, where Mn-dependent glucosyltransferases are located. A functional

defect of the human homolog TMEM165 is linked to cases of congenital disorders of

glycosylation, an inheritable disease leading to various pathological symptoms (Foulquier et al.,

2012; Stribny et al., 2020).

In photosynthetic eukaryotes, genes encoding 6-TMD members of the UPF0016 family

have undergone multiple duplication events (Hoecker et al., 2017). The genome of *Arabidopsis*

for instance encodes five UPF0016 proteins (Hoecker et al. 2017). CHLOROPLAST

MANGANESE TRANSPORTER 1 (CMT1) and PHOTOSYNTHESIS-AFFECTED MUTANT 71

(PAM71) reside in plastid membranes. CMT1 imports Mn across the plastid inner envelope into

the plastid stroma, while PAM71 facilitates uptake into the thylakoid lumen to provide Mn for its

incorporation into the oxygen evolving complex of PSII (Schneider et al., 2016; Eisenhut et al.,

2018; Zhang et al., 2018). PHOTOSYNTHESIS-AFFECTED MUTANT 71 LIKE 3 (PML3), also

known as BIVALENT CATION TRANSPORTER 3 (BICAT3), is *trans*-Golgi localized and plays a

critical role in glycosylation reactions under Mn limitation conditions, in cell wall biosynthesis

(Yang et al., 2021), and the allocation of Mn between Golgi apparatus and chloroplast (He et al.,

2022). PML4 and PML5 localize to the endoplasmic reticulum and likely fine-tune the uptake of

Mn (Hoecker et al., 2020). We identified half-sized versions of the UPF0016 proteins, Hmx1 and

Hmx2, in the cyanobacterium *Synechocystis* that each contain three instead of six TMDs

(Figure 1A).

Hmx1 and Hmx2 enable constitutive Mn uptake as heteromers at the plasma membrane

The vast majority of cyanobacterial genomes encode two genes encoding 3-TMD proteins,

where *hmx1* and *hmx2* are usually arranged in an operon (Figure 1B, Table S3). Since such

regulatory systems typically contain genes coding for functionally related partners, we expected

both proteins Hmx1 and Hmx2 to assemble as heteromers and function as Mn transporter in the

same way as their 6-TMD homologs of the UPF0016 family. Though a direct physiological

interaction was not investigated by us, our results support this hypothesis.

We were able to delete the genes in single and double knock-out mutants (Figure 2B).

The full genome segregation indicates that Hmx1 and Hmx2 are not essential for survival under

standard conditions with 1x (9 µM) $MnCl_2$ supplementation. The reduced growth performance at

this concentration and even more so in the absence of Mn (Figure 2C), however, clearly

demonstrates its important role in Mn uptake. The decreased ability to accumulate Mn inside of

cells after a 9 µM $MnCl_2$ pulse (Figure 2D) furthermore corroborates the Mn uptake function for

Hmx1/2.

The only Mn ABC-type transporter MntCAB that has been described for cyanobacteria

until now, serves Mn import at the plasma membrane under Mn limitation (Bartsevich & Pakrasi,

1995; Bartsevich & Pakrasi, 1996). Intriguingly, *mntCAB* knockout lines continued to show Mn

uptake when supplemented with micromolar amounts of Mn, indicating the presence of a

second, yet unidentified Mn uptake system (Bartsevich & Pakrasi, 1996). Our analysis of double

mutants with deletions in both *mntC* and *hmx2* revealed that both transporters assist each other

at Mn scarcity, since Δ*mntC/*Δ*hmx2* mutants are not able to survive without any Mn

supplementation (Figure 4C), while also showing the lowest levels of intracellular Mn

accumulation (Figure 4D). Bartsevich and Pakrasi (1996) furthermore postulated that the

additional Mn transport system should be active at different, likely micromolar extracellular Mn

conditions. This holds true for Hmx1/2. While the Δ*mntC* mutant shows the strongest growth

retardation at 0x $MnCl_2$, the phenotype is already fully recued at 0.5x (4.5 µM) $MnCl_2$. This result

was expected, since expression of *mntCAB* occurs under the control of the ManS/ManR two-

component system (Ogawa et al., 2002; Yamaguchi et al., 2002) and is known to be induced

only at extracellular Mn concentrations below 1 µM (Yamaguchi et al., 2002; Eisenhut, 2020).

Thus, at 4.5 µM $MnCl_2$ MntCAB is not expressed and another Mn uptake system must

compensate for its absence, which is Hmx1/2. Δ*hmx2* shows impaired growth performance at

concentrations up to 5x $MnCl_2$ (Figure 2C, Figure 4C) and the Δ*hmx1* and Δ*hmx2* mutants both

display Mn sensitivity over a rather broad range of $MnCl_2$ concentrations (0 - 9 µM) (Figure 2C).

Together with the observation that their gene expression pattern mirrors that of house-keeping

genes rather than being inducible (Figure 5A), and hardly reacts to changes in Mn

supplementation (Figure 5B), we postulate that Hmx1/2 functions as a constitutive Mn

transporter.

It was furthermore proposed (Bartsevich & Pakrasi, 1996) that the unidentified Mn

transporter should be highly specific for Mn and our results support this suggestion, too.

Besides Mn, only Ca has been demonstrated to be a substrate for some UPF0016 proteins

(Demaegd et al., 2013; Colinet et al., 2016; Wang et al. 2016; Frank et al., 2019). For its 6-TMD

homolog Mnx, specifically Mn-dependent effects were shown (Brandenburg et al., 2017b;

Gandini et al., 2017). Our experiments with medium depleted of Ca or Fe (Figure S1) revealed

unaffected growth of Δ*hmx1* and Δ*hmx2* mutants. Though we cannot rule out transport of other

untested metals, such as copper or zinc, our results point towards a high substrate specificity of

Hmx1/2 for Mn.

Finally, to serve Mn import, the proteins are expected to localize to the plasma

membrane. In both lines expressing *hmx1:cfp* or *hmx2:cfp* we observed that the CFP

fluorescence did not entirely overlap with the chlorophyll autofluorescence (Figure 3), which

would be indicative for a thylakoid membrane localization. The halo-like appearance is different

from thylakoid membrane proteins, such as Mnx (Brandenburg et al., 2017b) and rather

indicates Hmx1 and Hmx2 residing in the plasma membrane. To minimize experimental

artifacts, the CFP fusion constructs were introduced into the genes´ native chromosomal sites,

so that expression was guided by the native promoters and to rule out potential overexpression

artifacts. Furthermore, since the expression lines did not show Mn sensitivity (Figure S2B), we

argue that Hmx1:CFP and Hmx2:CFP reside in their designated locations and are functional.

Uptake across the plasma membrane serves Mn delivery to cytoplasmic Mn-requiring

proteins and also further passage via the thylakoid membrane transporter Mnx to assist Mn

incorporation into PSII. The additional and distinct patchiness of the CFP fluorescence signal is

very similar to that observed for CurT (Heinz et al., 2016; Ostermeier et al., 2022). CurT locates

to thylakoid convergence zones, which enable contact sites between the plasma and thylakoid

membrane. These zones are postulated to house early steps in PSII assembly (Nickelsen &

Rengstl, 2013; Heinz et al., 2016). Thus, it is conceivable that Hmx1/2 resides at these contact

sites and enables the Mn uptake for efficient provision of the metal cofactor at the site of pD1

translation, PratA-assisted Mn incorporation, and PSII biogenesis. Further studies, however, will

be needed to prove this hypothetical function.

Strikingly, the compensation of the Mn sensitive phenotype *of* Δ*hmx2* (Figure 2C) and

the viability of the Δ*mntC/*Δ*hmx2* double mutant (Figure 4C) at elevated Mn concentrations

suggest the presence of an additional Mn uptake system at the plasma membrane. However,

this transporter likely has only a low-affinity for Mn as indicated by the compensation with 5x

$MnCl_2$ (at minimum). A good candidate for the low-affinity Mn transport system is the Fe(III)

ABC-type transporter FutABC (Katoh et al., 2001). Expression of the transporter subunits is

slightly enhanced at Mn limitation (Sharon et al., 2014). FutABC might transport Mn as co-

substrate by a piggybacking mechanism (Brandenburg et al., 2017b; Eisenhut, 2020).

Hmx1 and Hmx2 are rather small proteins (117 and 92 amino acids, respectively). They

both comprise three TMDs and contain the signature ExGD motive, which is suggested to

participate in forming the pore of the transporter (Stribny et al., 2020). According to our

experimental results, they are only functional if both are present, presumably assembling as

heteromers: single and double mutants in *hmx1* and *hmx2* have congruent phenotypes. That is,

they show reduced growth on medium with low Mn concentrations (Figure 2C) and have smaller

intracellular Mn pools in comparison to the WT (Figure 2D). If homomers were active in Mn

transport, single deletions of either *hmx1* or *hmx2* would not have resulted in Mn sensitivity and

additive characteristics in the Δ*hmx1/*Δ*hmx2* double mutant would have been expected. Also,

hmx1 and *hmx2* are with 90 % (Table S3) predominantly encoded as operons, which supports a

functional interaction of both 3-TMD proteins and single 3-TMD units are too small for building a

transporter pore. Thus, we suggest that Hmx1/2 is required to assemble as multimeric

heteromers to enable Mn transport. Schneider et al. observed interaction of PAM71, a 6-TMD

protein of the UPF0016, with itself (Schneider et al., 2016). Likely, as demonstrated for

SWEETs (Xuan et al., 2013) or MFS eukaryote sugar transporters (Abramson et al., 2003),

UPF0016 proteins form a functional transporter, if assembling into 12-TMD complexes. Future

interaction and structural studies could provide evidence that the 3-TMDs and their conserved

ExGD motif, as represented by Hmx1 and Hmx2, in all cases act as the principle building block.

In summary, through the characterization of Hmx1/2 we propose to having identified the

so far only biochemically evidenced Mn transporter of cyanobacteria. It allows constitutive Mn

uptake and in collaboration with the MntCAB system ensures proper Mn provision under both

Mn-limiting and sufficient conditions. A model with Hmx1/2 integrated into the Mn homeostasis

network is provided in Figure 7. Like chloroplasts of the green lineage (Chloroplastida), also

cyanobacteria employ two transporters of the UPF0016 family for sequential uptake of Mn. A

comparable example is the occurrence of Cu-specific P-type ATPases in both cyanobacteria

(CtaA and PacS) (Kanamaru et al., 1994; Phung et al., 1994; Tottey et al., 2001) and plants

(PAA1 and PAA2) (Abdel-Ghany et al., 2005). In both cases the homologs act in tandem to

transport Cu^+-ions via the plasma membrane/chloroplast envelope and the thylakoid membrane.

Cyanobacterial Hmx1 and Hmx2 represent the ancestral form of UPF0016 proteins

Pairs of Hmx proteins are almost exclusively encoded in cyanobacterial genomes (Figure 6).

Isolated *hmx* genes, so called singletons, are found in some cyanobacteria, including

Gloeobacter, but also in other microorganisms (Demaegd et al., 2013). *hmx1* and *hmx2* likely

evolved by duplication of a single *hmx* gene within the cyanobacterial clade and were passed on

via horizontal gene transfer, also into non-cyanobacterial prokaryotes. Even screening with a

fairly low e-value (Figure S3), resulted in the detection of Hmx homologs with three TMDs (also

as singletons) only in a very few cases outside of the cyanobacterial group (Figure 6, Table S2).

Cyanobacteria, however, frequently encode the fusion form (UPF0016 proteins) with six TMDs.

In contrast to the 3-TMD proteins, this 6-TMD type is also observed in several other bacteria,

such as Gammaproteobacteria and the MneA of *e.g., Vibrio cholerae* (Fisher et al., 2016),

Deltaproteobacteria, or Actinobacteria (Figure 6). An ancestral origin of UPF0016 proteins and

subsequential loss in the vast majority of lineages cannot be ruled out, but parsimony suggests

a cyanobacterial origin of Hmx proteins and its duplication accompanied by a fusion event to

produce 6-TMD proteins of the Mnx-type, which was then horizontally transferred in some cases

(Table S2). Fusion forms of proteins, such as Mnx, have a higher probability to be retained after

horizontal gene transfer. Since our experiments with the knockout mutants in *hmx1* and/or *hmx2*

demonstrated that both proteins are essential for the function as Mn transporter, Hmx proteins

only add an advantage, if both proteins are transferred. Thus, their individual probability of being

retained after lateral gene transfer is rather low. It is conceivable that Hmx1 and Hmx2

represent the ancestral progenitor of the fused 6-TMD form of UPF0016 proteins, which was –

through endosymbiosis – carried into the eukaryotic tree of life (Hoecker et al., 2021). This is a

good time to remember that eukaryote genomes only contain 6-TMD versions of UPF0016

members (Figure S4).

The genomic occurrence of a *hmx* singleton in *Gloeobacter violaceus* allows to

speculate about a coevolutionary scenario of the UPF0016 Mn transporters with the

internalization of the OEC. *Gloeobacter violaceus* is an early-branching cyanobacterium that

lacks thylakoid membranes. Its components of the oxygenic photosynthesis apparatus are

located in the plasma membrane and the OEC resides in the periplasm (Raven & Sánchez-

Baracaldo, 2021). Accordingly, Mn uptake via the plasma membrane is only necessary to

provide Mn for intracellular Mn-dependent enzymes, not for the incorporation into the OEC. For

this purpose, the single Hmx protein in *Gloeobacter violaceus,* which possibly functions as

homomer, might be sufficient. However, with the emergence of thylakoid-forming cyanobacteria,

and the consequent internalization of the OEC into the thylakoid system, more efficient Mn

uptake became indispensable. It is conceivable that the duplication of a singleton *hmx* gene into

hmx1 and *hmx2* benefits the number of Mn importer that can be present (gene dosage effect),

which assures a sufficient and constitutive Mn uptake at the plasma membrane and at the PSII

biogenesis centers. The additional fusion form Mnx resides in the thylakoid membrane and

serves Mn import into the thylakoid lumen to assist Mn provision to the OEC. This sequential

uptake system specifically for Mn was endosymbiontically passed on to plastid containing

eukaryotes.

In conclusion, cyanobacteria employ all organizational structures of UPF0016 proteins,

from basic 3-TMD units to fused 6-TMD proteins. Hence, cyanobacteria are the most suitable

model to study the evolution and function of this important class of transport proteins. The Mn

transport function of Hmx1/2 highlights this as fundamental ancient feature of the UPF0016

family. We suggest that Hmx1/2 coevolved as an essential consequence of the cellular

internalization of the OEC (reviewed in Martin et al., 2018) at the basis of cyanobacteria

performing oxygenic photosynthesis.

AUTHOR CONTRIBUTIONS

MR, FB, ME designed the research. MR, FB, MK, SF, AB, SM, and ME performed the research.

All authors contributed to data analysis and discussion. MR, FB, MK, SBG, and ME wrote the

paper. All authors have read and agreed to this version of the manuscript.

ACKNOWLEDGEMENTS

We thank Stefanie Weidtkamp-Peters and Thomas Zobel (Center of Advanced Imaging,

Heinrich-Heine University Düsseldorf) for excellent technical assistance. We thank Jörg

Nickelsen and his team (LMU Munich) for providing us with the codon optimized CFP vector.

We acknowledge Lisa M. Heihoff for computational support. This work was funded by the

Deutsche Forschungsgemeinschaft (DFG) through the grant EI 945/3-2 and SFB1535 - Project

ID 458090666 to ME. SBG is grateful for DFG support through MAdLand SPP2237

(440043394) and the CRC1208 (267205415). We acknowledge support for the publication costs

by the Open Access Publication Fund of Bielefeld University and the DFG.

DATA AVAILABILITY STATEMENT

541 All relevant data can be found in the manuscript or in the Supporting Information.

SUPPORTING INFORMATION

Table S1: List of oligonucleotides used in this study

Table S2: Occurrence of Mnx, Hmx1, and Hmx2 homologs across bacteria, archaea, and

eukaryotes

Table S3: Occurrence and organization of *UPF0016* genes in cyanobacterial genomes

Figure S1: Growth of mutant strains on medium depleted of Ca or Fe

Figure S2: Examination of *hmx1:cfp* and *hmx2:cfp* expression lines

Figure S3: Occurrence of UPF0016 genes in prokaryotic genomes across varying e-value cut-

offs

Figure S4: Occurrence of UPF0016 genes in eukaryotic genomes with cut-off e-value ≦ 1E-10

REFERENCES

Abdel-Ghany SE, Müller-Moulé P, Niyogi KK, Pilon M, Shikanai T (2005) Two P-Type ATPases

Are Required for Copper Delivery in *Arabidopsis thaliana* Chloroplasts. *Plant Cell,* 17:

1233–1251

Abramson J, Smirnova I, Kasho V, Verner G, Kaback HR, Iwata S (2003) Structure and

Mechanism of the Lactose Permease of *Escherichia coli*. *Science,* 301: 610–615

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool.

Journal of Molecular Biology, 215: 403–410

Bartsevich VV, Pakrasi HB (1995) Molecular identification of an ABC transporter complex for

manganese: analysis of a cyanobacterial mutant strain impaired in the photosynthetic

oxygen evolution process. *EMBO Journal,* 14: 1845–1853

Bartsevich VV, Pakrasi HB (1996) Manganese transport in the cyanobacterium *Synechocystis*

sp. PCC 6803. *Journal of Biological Chemistry*, 271: 26057–26061

Bosma EF, Rau MH, van Gijtenbeek LA, Siedler S (2021) Regulation and distinct physiological

roles of manganese in bacteria. *FEMS Microbiology Reviews*, 45: fuab028

Brandenburg F, Schoffman H, Keren N, Eisenhut M (2017a) Determination of Mn

Concentrations in *Synechocystis* sp. PCC6803 Using ICP-MS. *Bio-Protocol,* 7: 1–7

Brandenburg F, Schoffman H, Kurz S, Krämer U, Keren N, Weber APM, Eisenhut M (2017b)

The *Synechocystis* manganese exporter mnx is essential for manganese homeostasis in

cyanobacteria. *Plant Physiology*, 173: 1798–1810

Breton C, Šnajdrová L, Jeanneau C, Koča J, Imberty A (2006) Structures and mechanisms of

glycosyltransferases. *Glycobiology*, 16: 29R-37R

Buchfink B, Xie C, Huson DH (2015) Fast and sensitive protein alignment using DIAMOND.

Nature Methods, 12: 59–60

Colinet A-S, Sengottaiyan P, Deschamps A, Colsoul M-L, Thines L, Demaegd D, Duchêne M-C,

Foulquier F, Hols P, Morsomme P (2016) Yeast Gdt1 is a Golgi-localized calcium

transporter required for stress-induced calcium signaling and protein glycosylation.

Scientific Reports, 6: 24282

Demaegd D, Colinet AS, Deschamps A, Morsomme P (2014) Molecular evolution of a novel

family of putative calcium transporters. *PLoS One,* 9: e100851

Demaegd D, Foulquier F, Colinet A-S, Gremillon L, Legrand D, Mariot P, Peiter E, Van

Schaftingen E, Matthijs G, Morsomme P (2013) Newly characterized Golgi-localized family

of proteins is involved in calcium and pH homeostasis in yeast and human cells.

Proceedings of the National Academy of Sciences of the United States of America, 110:

6859–6864

Dobáková M, Sobotka R, Tichý M, Komenda J (2009) Psb28 Protein Is Involved in the

Biogenesis of the Photosystem II Inner Antenna CP47 (PsbB) in the Cyanobacterium

Synechocystis sp. PCC 6803. *Plant Physiology*, 149: 1076–1086

Eisenhut M (2020) Manganese Homeostasis in Cyanobacteria. *Plants,* 9: 18

Eisenhut M, Hoecker N, Schmidt SB, Basgaran RM, Flachbart S, Jahns P, Eser T, Geimer S,

Husted S, Weber APM, Leister D, Schneider A (2018) The Plastid Envelope

CHLOROPLAST MANGANESE TRANSPORTER1 Is Essential for Manganese

Homeostasis in Arabidopsis. *Molecular Plant,* 11: 955–969

Eisenhut M, Kahlon S, Hasse D, Ewald R, Lieman-Hurwitz J, Ogawa T, Ruth W, Bauwe H,

Kaplan A, Hagemann M (2006) The Plant-Like C2 Glycolate Cycle and the Bacterial-Like

Glycerate Pathway Cooperate in Phosphoglycolate Metabolism in Cyanobacteria. *Plant*

Physiology, 142: 333–342

Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi

M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A (2016) The Pfam protein families

database: towards a more sustainable future. *Nucleic Acids Research*, 44: D279–D285

Fisher CR, Wyckoff EE, Peng ED, Payne SM (2016) Identification and Characterization of a

Putative Manganese Export Protein in *Vibrio cholerae*. *Journal of Bacteriology*, 198: 2810–

2817

Foulquier F, Amyere M, Jaeken J, Zeevaert R, Schollen E, Race V, Bammens R, Morelle W,

Rosnoblet C, Legrand D, Demaegd D, Buist N, Cheillan D, Guffon N, Morsomme P,

Annaert W, Freeze HH, Van Schaftingen E, Vikkula M, Matthijs G (2012) TMEM165

Deficiency Causes a Congenital Disorder of Glycosylation. *American Journal of Human*

Genetics, 91: 15–26

Frank J, Happeck R, Meier B, Hoang MTT, Stribny J, Hause G, Ding H, Morsomme P, Baginsky

S, Peiter E (2019) Chloroplast-localized BICAT proteins shape stromal calcium signals and

are required for efficient photosynthesis. *New Phytologist,* 221: 866–880

Gandini C, Schmidt SB, Husted S, Schneider A, Leister D (2017) The transporter SynPAM71 is

located in the plasma membrane and thylakoids, and mediates manganese tolerance in

Synechocystis PCC6803. *New Phytologist*, 215: 256–268

Hänsch R, Mendel RR (2009) Physiological functions of mineral micronutrients (Cu, Zn, Mn, Fe,

Ni, Mo, B, Cl). *Current Opinion in Plant Biology*, 12: 259–266

He J, Yang B, Hause G, Rössner N, Peiter-Volk T, Schattat MH, Voiniciuc C, Peiter E (2022)

The trans-Golgi-localized protein BICAT3 regulates manganese allocation and matrix

polysaccharide biosynthesis. *Plant Physiology*, 190: 2579–2600

Heinz S, Rast A, Shao L, Gutu A, Gügel IL, Heyno E, Labs M, Rengstl B, Viola S, Nowaczyk

MM, Leister D, Nickelsen J (2016) Thylakoid Membrane Architecture in *Synechocystis*

Depends on CurT, a Homolog of the Granal CURVATURE THYLAKOID1 Proteins. *Plant*

Cell, 28: 2238–2260

Hoecker N, Hennecke Y, Schrott S, Marino G, Schmidt SB, Leister D, Schneider A (2021) Gene

Replacement in Arabidopsis Reveals Manganese Transport as an Ancient Feature of

Human, Plant and Cyanobacterial UPF0016 Proteins. *Frontiers in Plant Science,* 12:

697848

Hoecker N, Honke A, Frey K, Leister D, Schneider A (2020) Homologous Proteins of the

Manganese Transporter PAM71 Are Localized in the Golgi Apparatus and Endoplasmic

Reticulum. *Plants*, 9: 239

Hoecker N, Leister D, Schneider A (2017) Plants contain small families of UPF0016 proteins

including the PHOTOSYNTHESIS AFFECTED MUTANT71 transporter. *Plant Signaling &*

Behavior, 12: 1–4

Jakubovics NS, Jenkinson HF (2001) Out of the iron age: new insights into the critical role of

manganese homeostasis in bacteria. *Microbiology* 147: 1709–1718

Kanamaru K, Kashiwagi S, Mizuno T (1994) A copper-transporting P-type ATPase found in the

thylakoid membrane of the cyanobacterium *Synechococcus* species PCC7942. *Molecular*

Microbiology, 13: 369–377

Katoh H, Hagino N, Grossman AR, Ogawa T (2001) Genes Essential to Iron Transport in the

Cyanobacterium *Synechocystis* sp. Strain PCC 6803. *Journal of Bacteriology*, 183: 2779–

2784

Kehres DG, Zaharik ML, Finlay BB, Maguire ME (2000) The NRAMP proteins of *Salmonella*

typhimurium and *Escherichia coli* are selective manganese transporters involved in the

response to reactive oxygen. *Molecular Microbiology*, 36: 1085–1100

Keren N, Kidd MJ, Penner-Hahn JE, Pakrasi HB (2002) A light-dependent mechanism for

massive accumulation of manganese in the photosynthetic bacterium *Synechocystis* sp.

PCC 6803. *Biochemistry*, 41: 15085–15092

Ku C, Nelson-Sathi S, Roettger M, Sousa FL, Lockhart PJ, Bryant D, Hazkani-Covo E,

McInerney JO, Landan G, Martin WF (2015) Endosymbiotic origin and differential loss of

eukaryotic genes. *Nature*, 524: 427–432

Marschner H (2012) Marschner's Mineral Nutrition of Higher Plants. 3rd ed, New York,

Academic Press

Martin WF, Bryant DA, Beatty JT (2018) A physiological perspective on the origin and evolution

of photosynthesis. *FEMS Microbiology Reviews*, 42: 205–231

Nelson N, Junge W (2015) Structure and Energy Transfer in Photosystems of Oxygenic

Photosynthesis. *Annual Review of Biochemistry*, 84: 659–683

Nickelsen J, Rengstl B (2013) Photosystem II assembly: from cyanobacteria to plants. *Annual*

Review of Plant Biology, 64: 609-635

Nowaczyk MM, Krause K, Mieseler M, Sczibilanski A, Ikeuchi M, Rögner M (2012) Deletion of

psbJ leads to accumulation of Psb27–Psb28 photosystem II complexes in

Thermosynechococcus elongatus. Biochimica et Biophysica Acta, Bioenergetics, 1817:

1339–1345

O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B, Robbertse B,

Smith-White B, Ako-Adjei D, et al (2016) Reference sequence (RefSeq) database at NCBI:

current status, taxonomic expansion, and functional annotation. *Nucleic Acids Research*,

44: D733–D745

Ogawa T, Bao DH, Katoh H, Shibata M, Pakrasi HB, Bhattacharyya-Pakrasi M (2002) A two-

component signal transduction pathway regulates manganese homeostasis in

Synechocystis 6803, a photosynthetic organism. *Journal of Biological Chemistry*, 277:

28981–28986

Ostermeier M, Heinz S, Hamm J, Zabret J, Rast A, Klingl A, Nowaczyk MM, Nickelsen J (2022)

Thylakoid attachment to the plasma membrane in *Synechocystis* sp. PCC 6803 requires

the AncM protein. *Plant Cell,* 34: 655-678

Phung LT, Ajlani G, Haselkorn R (1994) P-type ATPase from the cyanobacterium

Synechococcus 7942 related to the human Menkes and Wilson disease gene products.

Proceedings of the National Academy of Sciences of the United States of America, 91:

9651–9654

Raven JA, Sánchez-Baracaldo P (2021) *Gloeobacter* and the implications of a freshwater origin

of Cyanobacteria. *Phycologia*, 60: 402–418

Reis M, Zenker S, Viehöver P, Niehaus K, Bräutigam A, Eisenhut M (2024) Study of excess

manganese stress response highlights the central role of manganese exporter Mnx for

holding manganese homeostasis in the cyanobacterium *Synechocystis* sp. PCC 6803.

bioRxiv 2024.08.16.608223. [Preprint] Available from:

https://doi.org/10.1101/2024.08.16.608223. [Accessed: 27[th] August 2024]

Rice P, Longden I, Bleasby A (2000) EMBOSS: The European Molecular Biology Open

Software Suite. *Trends in Genetics*, 16: 276–277

Rippka R, Deruelles J, Waterbury JB (1979) Generic assignments, strain histories and

properties of pure cultures of cyanobacteria. *Journal of General Microbiology*, 111: 1–61

Schmidt SB, Husted S (2019) The Biochemical Properties of Manganese in Plants. *Plants*, 8:

381

Schmidt SB, Eisenhut M, Schneider A (2020) Chloroplast Transition Metal Regulation for

Efficient Photosynthesis. *Trends in Plant Science*, 25: 817–828

Schneider A, Steinberger I, Herdean A, Gandini C, Eisenhut M, Kurz S, Morper A, Hoecker N,

Rühle T, Labs M, et al (2016) The evolutionarily conserved protein PHOTOSYNTHESIS

AFFECTED MUTANT71 is required for efficient manganese uptake at the thylakoid

membrane in Arabidopsis. *Plant Cell,* 28: 892-910

Sharon S, Salomon E, Kranzler C, Lis H, Lehmann R, Georg J, Zer H, Hess WR, Keren N

(2014) The hierarchy of transition metal homeostasis: Iron controls manganese

accumulation in a unicellular cyanobacterium. *Biochimica et Biophysica Acta,*

Bioenergetics, 1837: 1990–1997

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M,

Söding J, Thompson JD, Higgins DG (2011) Fast, scalable generation of high-quality

protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology*, 7:

539

Socha AL, Guerinot ML (2014) Mn-euvering manganese: the role of transporter gene family

members in manganese uptake and mobilization in plants. *Frontiers in Plant Science*,

5:106.

Stengel A, Gügel IL, Hilger D, Rengstl B, Jung H, Nickelsen J (2012) Initial steps of

photosystem II de novo assembly and preloading with manganese take place in biogenesis

centers in *Synechocystis. Plant Cell,* 24: 660-675

Stribny J, Thines L, Deschamps A, Goffin P, Morsomme P (2020) The human Golgi protein

TMEM165 transports calcium and manganese in yeast and bacterial cells. *Journal of*

Biological Chemistry, 295: 3865–3874

Tottey S, Rich PR, Rondet SAM, Robinson NJ (2001) Two Menkes-type ATPases Supply

Copper for Photosynthesis in *Synechocystis* PCC 6803. *Journal of Biological Chemistry*,

276: 19999–20004

Tottey S, Waldron KJ, Firbank SJ, Reale B, Bessant C, Sato K, Cheek TR, Gray J, Banfield MJ,

Dennison C, Robinson NJ (2008) Protein-folding location can regulate manganese-binding

versus copper- or zinc-binding. *Nature,* 455: 1138–1142

Wang C, Xu W, Jin H, Zhang T, Lai J, Zhou X, Zhang S, Liu S, Duan X, Wang H, Peng C, Yang

C (2016) A Putative Chloroplast-Localized Ca^{2+} /H^+ Antiporter CCHA1 Is Involved in

Calcium and pH Homeostasis and Required for PSII Function in Arabidopsis. *Molecular*

Plant, 9: 1183–1196

Wulf D, Bräutigam A, Eisenhut M (2024) Using Publicly Available RNA-seq Data for Expression

Analysis of Genes of Interest. *Methods in Molecular Biology* (Clifton, N.J.), 2792: 241–250

Xuan YH, Hu YB, Chen L-Q, Sosso D, Ducat DC, Hou B-H, Frommer WB (2013) Functional role

of oligomerization for bacterial and plant SWEET sugar transporter family. *Proceedings of*

the National Academy of Sciences of the United States of America, 110: E3685-E3694

Yamaguchi K, Suzuki I, Yamamoto H, Lyukevich A, Bodrova I, Los DA, Piven I, Zinchenko V,

Kanehisa M, Murata N (2002) A two-component Mn^{2+}-sensing system negatively regulates

expression of the *mntCAB* operon in *Synechocystis*. *Plant Cell*, 14: 2901–2913

Yang C, Wang C, Singh S, Fan N, Liu S, Zhao L, Cao H, Xie W, Yang C, Huang C (2021)

Golgi-localised manganese transporter PML3 regulates *Arabidopsis* growth through

modulating Golgi glycosylation and cell wall biosynthesis. *New Phytologist*, 231: 2200–

2214

Zhang B, Zhang C, Liu C, Jing Y, Wang Y, Jin L, Yang L, Fu A, Shi J, Zhao F, Lan W, Luan S

(2018) Inner Envelope CHLOROPLAST MANGANESE TRANSPORTER 1 Supports

Manganese Homeostasis and Phototrophic Growth in Arabidopsis. *Molecular Plant*, 11:

943–954

Figure legends

Figure 1: A) Amino acid alignment of UPF0016 members of *Synechocystis*. Grey boxes

indicate predicted transmembrane domains. The conserved ExGD motif is highlighted in red.

Asterisks (*) indicate strictly conserved residues and a colon (:) depicts residues with similar

properties. B) Genomic organization of *hmx1* (orange) and *hmx2* (red) orthologs across some

selected cyanobacteria. Additional genes as part of the transcriptional unit are depicted in grey.

The direction of arrows indicates the orientation in the genome.

Figure 2: Analysis of $\Delta hmx1$ and $\Delta hmx2$ mutants. A) Scheme for the generation of mutants in

hmx1 ($\Delta hmx1$) and *hmx2* ($\Delta hmx2$) by insertional inactivation. B) Verification of mutations in

$\Delta hmx1$, $\Delta hmx2$, $\Delta hmx1/\Delta hmx2$, and $\Delta hmx2/\Delta hmx1$ by genotyping. PCR analysis was performed

with gDNA and gene specific primers *(hmx1*: primers FB69/FB70, WT = 954 bp; Mu = 5,002 bp.

hmx2: primers ME368/ME369, WT = 1,279 bp; Mu = 2,531 bp). C) Droptest to monitor Mn

sensitivity. Cells were washed with BG11 -Mn and adjusted to an OD_{750} of 0.25. Cells were

diluted 1:10, 1:100, 1:1000 with BG11 -Mn and 2 µL of these cell suspensions were dropped

onto BG11 medium supplemented without (w/o) $MnCl_2$ (0 µM), 1x $MnCl_2$ (9 µM), or 5x $MnCl_2$ (45

µM). Pictures were taken after 5 d growth at 30 °C, 70 µmol photons m^{-2} s^{-1}. D) Determination of

780 intracellular Mn concentrations before (0 h) and 4 h after addition of 9 µM MnCl$_2$. Cells were

781 precultivated for 5 d in BG11 -Mn and then treated with 9 µM MnCl$_2$. Samples were taken before

782 (0 h) and 4 h after the addition of MnCl$_2$. To only determine the intracellular Mn pool, periplasmic

783 Mn was eliminated by two washing steps with 5 mM EDTA in 20 mM HEPES buffer (pH 7.5).

784 Data obtained for the WT measurement at 0 h was set to 100 % and subsequent values were

785 normalized to this point. Typical values obtained for the WT at 0 h were 1.4*10^6 Mn atoms/cell

786 (= 100%). Shown are averages and standard deviations of three biological (with five technical)

787 replicates each. Asterisks indicate significant differences between the reference value and

788 specified knockout mutant according to a Student´s t-test (*: $P \leq 0.05$; **: $P \leq 0.01$).

790 **Figure 3:** Subcellular localization of Hmx1 and Hmx2. The subcellular localization of Hmx1 and

791 Hmx2 was determined by confocal fluorescence microscopy. CFP was C-terminally fused to

792 Hmx1 and Hmx2. The fusion constructs were introduced into the original *hmx1* and *hmx2* locus,

793 resulting in the strains *hmx1:cfp* and *hmx2:cfp* and expression under the endogenous promoter.

794 Typical results for *hmx1:cfp* are shown in A and B and for *hmx2:cfp* in C and D. CFP

795 fluorescence is shown in orange, chlorophyll autofluorescence in blue, and a merged image

796 shows both signals at the same time (A and C). Intensities of the signals from the outside to the

797 inside of the cell at various positions of the cell circumference were analyzed as indicated by the

798 regions of interest (ROIs) (B and D). Intensities of the signals along the cell circumference were

799 analyzed as indicated by the ROIs (B and D).

801 **Figure 4:** Analysis of Δ*hmx2* and Δ*mntC* mutants. A) Scheme for the generation of a mutant in

802 *mntC* (Δ*mntC*) by deletion (Δ = 35 bp) and insertion of a spectinomycin resistance cassette

803 (Sp^R). B) Verification of mutations in Δ*hmx2*, Δ*mntC,* and Δ*hmx2/*Δ*mntC* by genotyping. PCR

804 analysis was performed with gDNA and gene specific primers (*hmx2*: primers ME368/ME369,

805 WT = 1,279 bp; Mu = 2,531 bp. *mntC*: primers ME163/ME164, WT = 664 bp; Mu = 2,712 bp). C)

Droptest to monitor Mn sensitivity. Cells were washed with BG11-Mn and adjusted to an OD_{750}

of 0.25. Cells were diluted 1:10, 1:100, 1:1000 with BG11-Mn and 2 µL of these cell

suspensions were dropped onto BG11 medium supplemented without (w/o) $MnCl_2$ (0 µM), 0.5x

$MnCl_2$ (4.5 µM), 0.75x $MnCl_2$ (6.75 µM), 1x $MnCl_2$ (9 µM), 2.5x $MnCl_2$ (22.5 µM), or 5x $MnCl_2$ (45

µM). Pictures were taken after 5 d growth at 30 °C, 70 µmol photons m^{-2} s^{-1}. D) Determination of

intracellular Mn concentrations before (0 h) and 4 h after addition of 4.5 µM $MnCl_2$. Cells were

precultivated for 5 d in BG11 -Mn and then treated with 4.5 µM $MnCl_2$. Samples were taken

before (0 h) and 4 h after the addition of $MnCl_2$. To exclusively determine the intracellular Mn

pool, periplasmic Mn was eliminated by two washing steps with 5 mM EDTA in 20 mM HEPES

buffer (pH 7.5). Data obtained for the WT measurement at 0 h was set to 100 %, and

subsequent values were normalized to this point. Typical values obtained for the WT at 0 h were

$1*10^6$ Mn atoms/cell (= 100%). Shown are averages and standard deviations of 4 or 5 biological

with 5 technical replicates each. Asterisks indicate significant differences between referring WT

value and specified knockout mutant according to a Student´s t-test (*: $P ≤ 0.05$; **: $P ≤ 0.01$).

Figure 5: Expression analysis of *hmx1* and *hmx2.* A) Coefficient of variation for *hmx1* and

hmx2. As representatives for inducible gene expression, *flv2*, *flv4*, *isiA*, *isiB*, *mntA*, *mntB*, and

mntC were visualized. *petB* is used as a control for a house-keeping gene. B) Mn-dependent

transcript accumulation in *Synechocystis* WT cells. Given are log2 fold changes of transcript

abundances 48 h after application of Mn limitation (w/o $MnCl_2$) conditions (-Mn; Sharon et al.,

2014) or 24 h of Mn excess (10x $MnCl_2$) conditions (+Mn, Reis et al., 2024) versus Mn control

(1x $MnCl_2$) conditions. Asterisks indicate significant changes ($q < 0.01$, q-value calculated

according to Benjamini-Hochberg).

Figure 6: Occurrence of UPF0016 genes in prokaryotic genomes. A database of 5,655

complete prokaryotic genomes was searched for homologs of Mnx, Hmx1, and Hmx2 via

DIAMOND. The presence-and-absence pattern was color coded based on sequence identity

and only hits with at least 25 % sequence identity and a maximum e-value of 1E-10 were

plotted. Occurrence of Mnx, Hmx1, and Hmx2 homologs in cyanobacterial genomes only are

shown in the inlet.

Figure 7: Biological role of Hmx1 and Hmx2 in cellular Mn homeostasis of cyanobacteria. For

Synechocystis, two distinct Mn pools were observed: up to 80% of the cellular content was

found to accumulate in the periplasm, while only 20% reside inside the intracellular space

(Keren et al., 2002). In the periplasmic space, Mn is most likely loosely bound to the outer

membrane by the negative membrane potential or bound to soluble Mn-binding proteins, such

as MncA (Keren et al., 2002; Tottey et al., 2008). Inside the cell, roughly 80% of the Mn is

associated with the Mn-cluster of the OEC. This major sink of Mn is assumed to being supplied

via two routes. i) The tetratricopeptide repeat protein PratA delivers Mn from the periplasm

directly to the precursor of the D1 (pD1) reaction center protein in the biogenesis centers of the

cell (Stengel et al., 2012). ii) The manganese exporter Mnx (Brandenburg et al., 2017b), also

known as SynPAM71 (Gandini et al., 2017) transports Mn from the cytoplasm into the thylakoid

lumen and thus serves as an alternative delivery option. Besides the ABC-type transporter

MntCAB, which is only expressed under Mn-limiting conditions (Bartsevich & Pakrasi, 1996), the

here studied UPF0016 proteins Hmx1 and Hmx2 serve Mn import at the plasma membrane and

biogenesis centers. They form heteromers and are constitutively active. The ferric Fe importer

FutABC likely has a low-affinity Mn uptake activity.

A

```
Mnx   MLTAFTAGLLLITVSELGDKTFFIAMILAMRYPRRWVLVGVVGGLAAMTILSVLMGQIFT
Hmx1  ------------------------------------------------------------
Hmx2  ------------------------------------------------------------

Mnx   FLPTRYINYAEVALFLIFGTKLLWDARRIKATANLEEMEDAEKAIASGEKKLKIVPRGWG
Hmx1  ----------------------------------MSPPLPLLLPSS-QTAVSQDLPASPNYFRP
Hmx2  ----------------------------------------------------------MDWQ

Mnx   IVVESFALTFVAEWGDRTQIATIALAAS-NNAWGVSAGAILGHTICAVIAVMGGKFVAGR
Hmx1  VFFSTFLTIFLAEMGDKTQLSTLLISAESQSPWVVFAGSALALISTSLLGVSLGYWIARR
Hmx2  LFGLSFITVFLAEIGDKSQLAAIALGGSAKSPRAVFFGSVTALILASFLGVLAGGSLAQF
      :.   :*    *:** **::*:::: :... :.  *  *:  .  :.:.*  *  :*

Mnx   ISEKTVTLIGGLLFYLFAVVSWWTKIA--
Hmx1  LDPQILDFSVALLLLLIA-GLLMGDVVSA
Hmx2  LPTKLLKALAALGFTIMALRLLWPNQED-
      :  : :    .* : ::*       .
```

B

Anabaena sp. 90

Prochlorococcus marinus str. MIT 9211

Synechococcus sp. PCC 7002

Chroococcidiopsis thermalis PCC 7203

Cyanobium gracile PCC 6307

Cyanothece sp. ATCC 51142

Geitlerinema sp. PCC 7407

200 bp

ANA_C20442   ANA_C20443

P9211_09121   P9211_09131   P9211_09141

SYNPCC7002_A2755   SYNPCC7002_A2757   SYNPCC7002_A2758

Chro_1096   Chro_1097   Chro_1098

Cyagr_0711   Cyagr_0712   Cyagr_0713

psb28-2   cce_2793   cce_2794

GEI7407_3823   GEI7407_3824   GEI7407_3825

gene   hmx1   hmx2

A

Sp^R + vector

NaeI

hmx1 (slr1170)

5′ — 3′

WT: 954 bp
Mu: 5,002 bp

Km^R

SmaI

hmx2 ssr1558

5′ — 3′

WT: 1,279 bp
Mu: 2,531 bp

500 bp

B

Ladder WT Δ*hmx1* Δ*hmx2* Δ*hmx1*/Δ*hmx2* H₂O

kb
5.0
3.0

1.0
0.75

hmx1

Ladder WT Δ*hmx1* Δ*hmx2* Δ*hmx1*/Δ*hmx2* H₂O

3.0
2.0
1.5
1.0

hmx2

C

WT Δ*hmx1* Δ*hmx2* Δ*hmx1*/Δ*hmx2*   WT Δ*hmx1* Δ*hmx2* Δ*hmx1*/Δ*hmx2*   WT Δ*hmx1* Δ*hmx2* Δ*hmx1*/Δ*hmx2*

undil.
1:10
1:100
1:1000

w/o $MnCl_2$ 1x $MnCl_2$ 5x $MnCl_2$

D

Intracellular Mn concentration [% relative to WT, 0 h]

0 h   4 h + $MnCl_2$

WT Δ*hmx1* Δ*hmx2* Δ*hmx1*/Δ*hmx2*   WT Δ*hmx1* Δ*hmx2* Δ*hmx1*/Δ*hmx2*