Filament formation drives catalysis by glutaminase enzymes important in cancer progression

Shi Feng[1], Cody Aplin[1], Thuy-Tien T. Nguyen[1], Shawn K. Milano[1], and Richard A. Cerione[1,2*]

[1]Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York, 14853

[2]Department of Molecular Medicine, Cornell University, Ithaca, New York, 14853

*Corresponding author: rac1@cornell.edu

The glutaminase enzymes GAC and GLS2 catalyze the hydrolysis of glutamine to glutamate, satisfying the 'glutamine addiction' of cancer cells. They are the targets of anti-cancer drugs; however, their mechanisms of activation and catalytic activity have been unclear. Here we demonstrate that the ability of GAC and GLS2 to form filaments is directly coupled to their catalytic activity and present their cryo-EM structures which provide an unprecedented view of the conformational states essential for catalysis. Filament formation guides an 'activation loop' to assume a specific conformation that works together with a 'lid' to close over the active site and position glutamine for nucleophilic attack by an essential serine. Our findings highlight how ankyrin repeats on GLS2 regulate enzymatic activity, while allosteric activators stabilize, and clinically relevant inhibitors block, filament formation that enables glutaminases to catalyze glutaminolysis and support cancer progression.

Metabolic reprogramming is well-established as a hallmark of cancer progression[1-3]. A shift in the metabolic requirements of cancer cells compared to normal cells, known as the Warburg effect[4-6], results in their dependence on glutamine as an energy source in addition to glucose.

Glutamine catabolism, which starts with the hydrolysis of glutamine to glutamate catalyzed by the

glutaminase family of enzymes, becomes highly upregulated in cancer cells to compensate for the

uncoupling of the glycolytic pathway from the TCA cycle due to the Warburg effect, thus

providing the biosynthetic precursors necessary for their enhanced proliferation and survival[7-10].

Given the resulting glutamine addiction of cancer cells, the development of allosteric small

molecule inhibitors that target glutaminase activity is being examined as a potential therapeutic

strategy to block tumor progression in various malignancies including aggressive breast, lung, and

pancreatic cancers, as well as for their effectiveness when combined with other anti-cancer

drugs[11-16]. The glutaminases are encoded by two independent genes, *gls* and *gls2*. *Gls* encodes

kidney-type glutaminase (KGA) and its C-terminal truncated splice variant glutaminase C (GAC)

which is highly expressed in many cancers. *Gls2* encodes only one active isozyme, liver-type

glutaminase (LGA; from hereon designated as GLS2). While GLS2 has been suggested to serve

as a tumor suppressor by directly binding the Rac GTPase and promoting ferroptosis in

hepatocellular carcinoma[17,18], it has also been shown to be necessary for cancer progression as we

and others have found that it is essential for the growth and survival of luminal subtype receptor-

positive breast cancer cells[9,19]. Moreover, GLS2 is highly expressed in ovarian, lung and colorectal

cancers and has been suggested to compensate for GAC downregulation or inhibition[10,20].

All glutaminase enzymes share a catalytic domain containing identical active site residues and

a conserved mechanism for hydrolyzing glutamine to glutamate[21-23]. The catalytic efficiency of

the glutaminase enzymes is markedly increased in the presence of inorganic phosphate *in vitro*[24,25],

although the concentration of this allosteric activator required to promote their enzymatic activity

may not be attainable in many physiological contexts suggesting that other anionic activators likely

exist. Understanding more about the catalytic mechanism used by these enzymes and the structural

changes that they need to undergo to become activated will be important for further developing drug candidates that effectively block their enzymatic activity. Potent inhibitors targeting GAC are available, the most effective being the BPTES/CB-839 group of compounds with CB-839 and others such as UPGL004 having nanomolar binding affinity[26,27]. However, the development of specific and potent allosteric inhibitors of GLS2 has not been as successful, as currently the best examples are the 968-class of molecules[9,15] which have binding affinities for the enzyme only in the micromolar range.

It has been known for some time that a number of metabolic enzymes assemble into filament-like structures which may play a key regulatory role in their catalytic activity, but in the majority of cases, the importance of these higher order structures for enzyme regulation and function has not been definitively established[28]. Recently it has been shown that GAC can assemble into long filaments in cells to facilitate efficient glutamine consumption[29]. While the formation of these oligomeric supramolecular assemblies has been suggested to be associated with the *in vitro* activation of GAC by inorganic phosphate[30,31], an essential question concerns if the assembly of such higher-order oligomers is in fact directly coupled to catalytic activity. Moreover, it has yet to be established whether filament formation is a common mechanism for all glutaminase isozymes. This is an especially important question for GLS2 given that it might be differentially regulated due to the presence of ankyrin repeats, which are absent on the C-terminal truncated splice variant GAC[32,33], and because it can sometimes exert the opposite functional effects from GAC on cancer progression.

In this study, we show that GAC as well as GLS2 undergoes filament formation in the presence of the substrate glutamine and the activator inorganic phosphate, with the assembly of these higher-order oligomers being directly coupled to glutamine hydrolysis and product (glutamate) release.

We further present the high-resolution structures for a GAC filament bound to its anionic activator

inorganic phosphate (Pi), and for a constitutively active GLS2 filament, determined by cryo-

electron microscopy (cryo-EM). These novel filamentous structures now provide us with several

important new insights into how key regions of the glutaminases and their essential residues are

positioned to achieve maximal catalytic activity, as well as shed light on how anionic activators,

allosteric inhibitors, and ankyrin repeats can impart their regulatory effects on enzymes that play

critical roles in cancer progression.

RESULTS

General structural features of the glutaminase enzymes

Several high-resolution X-ray structures for KGA and GAC have been reported; however,

thus far, there has not been a structural determination for the full-length apo-GLS2 protein.

Therefore, we obtained a cryo-EM structure for wildtype, full-length GLS2. Figs. 1a and 1b show

the 3.1 Å structure of the human apo-GLS2 tetramer and the relative positions of the key regions

for catalytic activity (i.e., the 'activation loop', the catalytic site, and the 'lid loop' that opens and

closes over the catalytic domain)[23]. Fig. 1a also highlights the three major domains of a monomeric

unit of GLS2, namely, the N-terminal region, C-terminal ankyrin repeats, and the highly conserved

catalytic site. The overall structure of the apo-GLS2 tetramer strongly resembles those of the apo-

forms of GAC and KGA solved by X-ray crystallography (Fig. 1a)[24,33], and the active sites within

each GLS2 subunit show the same general features as a previously reported crystal structure of the

enzyme's limit catalytic domain as well as those of GAC[22] (Fig. 1c). The fact that these structural

features are highly conserved among the glutaminase enzymes indicate that they share a common

catalytic mechanism. We analyzed several key structural elements in the catalytic domain of GLS2

that have been suggested to exert an allosteric regulation of glutaminase activity. The stretch of

residues Gly248-Glu258, which has been shown to play an essential role in catalysis and thus

referred to as the 'activation loop'[23,34], is disordered and flexible in the GLS2 structure (Fig. 1b).

In GAC, this loop contains the binding site for the BPTES class of allosteric inhibitors which

includes CB-839, a drug candidate that has been in clinical trials[26]. The loop forms a stable,

inactive conformation when bound to BPTES and to its more potent analogs, CB-839 (not shown)

and UPGL0004[35,36] (Extended Data Fig. 1). However, the activation loop of GLS2 differs from

KGA and GAC by two residues, rendering GLS2 insensitive to these inhibitors[24]. In addition,

residues Tyr182-Lys188 in GLS2, which correspond to a loop in KGA and GAC forming a 'lid'

that needs to close over bound substrates for catalysis to occur, are also flexible like the activation

loops in the GLS2 structure. This is in contrast to the previously reported crystal structure of the

GLS2 limit catalytic domain, which showed the lid region adopting an "open" conformation[22].

Glutaminase filament formation is directly coupled to catalytic activity

The glutaminase enzymes have been shown to undergo dimer-to-tetramer transitions upon

binding activators such as inorganic phosphate (Pi)[25,37], as a necessary step for enzymatic activity.

However, high-resolution X-ray crystal structures of the tetrameric forms of glutaminase enzymes

have failed to shed light on the critical changes in their catalytic sites that promote the hydrolysis

of glutamine to glutamate. In fact, the positions of the catalytic residues in the structures obtained

for glutaminase dimers and tetramers are virtually identical[25,38]. Therefore, these findings further

raised the possibility that GAC needs to form a higher-order oligomeric state in cells, resulting in

a filament-like structure, in order for catalysis to occur[29,31,39]. Given the structural similarities

between GAC and GLS2, it might be expected that GLS2 would undergo a similar structural transition, although GLS2 can in some cancers exhibit the opposite effects from GAC by acting as a tumor suppressor rather than a promoter raising the possibility that it might exhibit distinct structural features. Thus, we set out to see if a filament-like structure is indeed formed by GLS2 as well as GAC and whether its formation is coupled to catalysis.

Using negative stain electron microscopy (EM), we visualized the structural changes that recombinant wild-type human GAC and GLS2 undergo upon the addition of the substrate glutamine and the anionic activator Pi under catalytic conditions (Figs. 2a and 2b). GAC immediately formed filaments with the addition of the substrate and activator, similar in appearance to those described in a previous report[31], and the same was true for GLS2, thus indicating that filament formation is common to all glutaminase enzymes. However, once the substrate glutamine was fully converted to product (glutamate), the filaments disassembled into tetramers as shown for GLS2 in Fig. 2c, supporting the idea that filament formation is coupled to catalytic turnover.

To further establish a direct connection between filament formation and enzyme catalysis, we took advantage of right-angle light scattering (RALS) to monitor the assembly of these higher order oligomers in real time. Fig. 2d shows an example for GLS2. Upon adding Pi and glutamine to the enzyme, there was an immediate, marked increase in light scattering. The increase in RALS peaked within minutes and was followed by a rapid decline, with the decrease in the peak matching the time scale for the dissociation of GLS2 filaments as visualized by negative stain EM. The maximal increase in RALS required both the substrate glutamine and the anionic activator Pi and was not detected in the absence of substrate (Extended Data Fig. 2a), demonstrating that the changes in RALS accurately reflect the association and dissociation of filaments. Subsequent

additions of glutamine to the enzyme re-established filament formation, with substrate

consumption again resulting in filament disassembly (Fig. 2d). The addition of the product,

glutamate, was unable to trigger filament formation. Taken together, these results further indicate

that filament assembly-disassembly is directly coupled to substrate binding and catalytic turnover.

Mutations that result in the constitutive activation of the glutaminase enzymes were then

examined by negative stain EM and RALS for their effects on filament formation. Changing a key

lysine residue (i.e., Lys320 in GAC and Lys253 in GLS2) to alanine within the activation loop

gives rise to constitutively active enzymes, with their activity matching that of the wild-type

enzyme measured in the presence of the anionic activator Pi[38]. The RALS profile and negative

stain images showed that glutamine addition alone was sufficient to drive constitutively active

forms of human GAC (K320A) and GLS2 (K253A) to form filaments, as the activation loop

substitution effectively substitutes for the requirement of an anionic activator (Extended Data Figs.

2b-d). Again, the product glutamate was unable to drive filament assembly, nor did the presence

of high concentrations of glutamate prior to glutamine addition have any effect on the ability of

substrate to induce filament formation (Extended Data Fig. 2d). The lifetime of the higher order

oligomeric filaments, as readout by RALS, was a direct function of the substrate (glutamine)

concentration, as shown for GLS2 (Extended Data Fig. 2e). Likewise, the length of the filaments

that formed were a function of substrate concentration and consisted of as many as 30 tetramers

(Extended Data Fig. 2f), most likely because the greater the substrate concentration, the longer the

time required for its total consumption and thus the ability to increase the growth of the filaments.

This is again consistent with filament assembly-disassembly being directly coupled to enzyme

activation and catalytic turnover. We then reasoned that conditions that allow substrate binding

and filaments to form, but do not permit product (glutamate) formation, should result in the

persistence of these higher order oligomers. Therefore, we tested this prediction by taking

advantage of our earlier finding that substituting an essential tyrosine residue within the enzyme

active site of GAC does not disrupt substrate (glutamine) binding but prevents its hydrolysis to

glutamate[25,34]. As shown in Fig. 2e, the RALS signal for GAC (Y466W) did not diminish over

time.

Structural determinations of GAC and GLS2 filaments

We determined a 3.3 Å cryo-EM structure of the human GAC (Y466W) mutant which,

because it is defective for hydrolyzing glutamine to glutamate, assembled into a stable filament

upon binding substrate and the anionic activator phosphate (Pi)[25,34]. Each successive tetramer is

rotated ~51 degrees relative to the filament axis with the helical rise of ~68 Å (Figs. 3a, b). We

also obtained a 3.3 Å cryo-EM structure of the constitutively active recombinant human GLS2

(K253A) mutant that formed a filament immediately after the addition of glutamine alone (and

prior to substrate depletion), as this mutant does not require an anionic activator to be catalytically

competent (Figs. 3c, d). The tetramers are arranged in a side-by-side architecture and each

successive tetramer is rotated ~48 degrees relative to the filament axis, with the helical rise of ~66

Å (Fig. 3d). These structures for activated glutaminase enzymes differed from the previously

proposed end-to-end double helix model for GAC filament formation[31]. In both the GAC and

GLS2 filament structures, each tetramer interacts with its immediate neighbor through an interface

of six alpha helices on the lateral side of the catalytic domain near the lid loop that closes over the

substrate binding site (Figs. 3e, f). Within this highly conserved interface burying about 710 Å^2

and 752 Å^2 of solvent-accessible area on GAC and GLS2, respectively, Phe355 in GAC (Phe288

in GLS2), together with Phe373 (Phe306 in GLS2), Phe378 (Phe311 in GLS2), Asp412 (Asp345

in GLS2), and Gln416 (Gln349 in GLS2) from one subunit form an intricate interacting network

with Asn375 (Asn308 in GLS2) and Gln379 (Gln312 in GLS2) from the subunit of an adjacent

tetramer (Figs. 3g, h).

When comparing the filament structures with those for the corresponding tetrameric apo-

enzymes described above (Figs. 1, and Extended Data Fig. 1), we found striking differences in the

conformation of the activation loops and the lids. The activation loops in both the GAC and GLS2

tetramers, either in their apo-, substrate-bound, or product-bound states, are highly flexible and

cannot be resolved, except when GAC is bound to the BTPES class of inhibitors (Extended Data

Fig. 1). This led us to originally suspect that a flexible rather than stabilized activation loop was

essential for the catalytic activity of these enzymes. However, in the substrate- and Pi-bound GAC

filament, as well as in the constitutively active substrate-bound GLS2 filament structure, the

activation loops are well resolved and adopt a conformation distinct from the inhibitor-induced

state in GAC (Figs. 3e, f, and Extended Data Fig. 1).

The first significant conformational change involving the activation loops of the GAC and

GLS2 filaments, compared to their positions in the apo-enzymes, resulted in a new motif

designated here as the 'phenylalanine-tyrosine lock' in GAC and the 'tyrosine lock' in GLS2.

These residues lock-in the substrate at the conserved active sites (Figs. 4a, b). In each dimer within

a tetrameric unit comprising the filament structures of GAC and GLS2, Glu325 in GAC (Glu258

in GLS2) within the activation loop of one monomer subunit forms a new salt bridge with Arg317

(Arg250 in GLS2) from the activation loop of the adjacent monomer subunit (Figs. 4c, d). In

solving the cryo-EM structure for the GAC(Y466W) filament in the presence of glutamine and Pi,

we were able to identify the Pi binding site within the dimer-dimer interface of each tetrameric

unit where it interacts with Lys320 and thereby stabilizes the activation loop (Fig. 4c, and Extended

Data Fig. 3, left panel). This helps to explain how the binding of Pi to a GAC(F327W) mutant is

able to induce an increase in Trp327 fluorescence within the activation loops of the enzyme[40].

Conversely, in the absence of Pi and substrate, the activation loops of the enzyme are flexible and

therefore are not visualized. Thus, the activation loops of each dimer within the filament fold in a

compact manner, thereby bringing Phe318 of GAC (Tyr251 of GLS2) from each loop closer to the

active site (Fig. 4c). In the apo-enzyme structures, Arg387 of GAC and Arg320 in GLS2 face

toward the active site and prevent glutamine from being 'locked in'. However, in the filament

structure of GAC, Pi interacts with Arg387, enabling it to form a salt bridge with Glu397 at the

dimer-dimer interface (Figs. 4c, d), which together with the interactions of Pi with Tyr394 and

Lys398 (Extended Data Fig. 4a), may be necessary for the glutaminase enzymes to undergo the

dimer-to-tetramer transition, a prerequisite for filament formation and catalytic activity.

Interestingly, in the filament structure for the constitutively active GLS2 mutant, Arg320 (which

corresponds to Arg 387 in GAC) forms the same type of salt bridge in the absence of Pi, as that

seen when Pi binds to GAC(Y466W). The formation of these Arg-Glu salt bridges, together with

the substitution of a less bulky alanine residue for lysine, allows the movement of the activation

loops into a space that otherwise would only occur when Pi binds, and likely explains why the

Lys-to-Ala substitutions within the loops yield enzymes that no longer require the binding of an

anionic activator.

When comparing the filament structures with those of the apo-enzyme, a second significant

conformational change involves the lid loops, resulting in the insertion of Tyr249 of GAC (Tyr182

of GLS2) into the active site pockets. These two conformational rearrangements involving the

activation and lid loops fully close off the active site in each subunit, acting as a 'Phe-Tyr lock' in

GAC and a 'Tyr lock' in GLS2, effectively locking the substrate glutamine into an optimal

orientation proximal to the residues essential for catalysis (Figs. 4a, b). Moreover, the insertion of

the phenyl ring of Phe318 of GAC (Tyr251 in GLS2) brings it within ~5 Å of Lys289 (Lys222 in

GLS2), as shown for GAC in the right panel of Extended Data Fig. 3 and GLS2 in Fig. 4e, thereby

enabling a cation-π interaction that orients and increases the basicity of a key active site lysine

residue. This enables the lysine residue to effectively deprotonate an active site serine (i.e., Ser286

in GAC and Ser219 in GLS2), which undergoes a nucleophilic attack and forms a tetrahedral

intermediate with glutamine. Hydrolysis can then occur following this initial reaction to form

glutamate and ammonia[41].

An additional conformational change that occurs specifically in the constitutively active GLS2

filament structure, when compared to the tetrameric apo-enzyme, results in a new hydrogen

bonding network introduced by the activation loops. Hydrogen bonds that form between Arg250

and Asp400 of GLS2 appear to 'stretch' the alpha helix containing Asp400, causing Tyr399 on the

N-terminal end of the alpha helix to insert further into the active site where the substrate binds

(Fig. 4f). Following the insertion of Tyr399, Glu258 from the adjacent dimer forms a hydrogen

bond with Lys440, which displaces the loop containing Lys440, and therefore makes space for the

insertion of Tyr399. As a result, Tyr399 is now in close proximity to both Lys222 and the substrate,

which is necessary for catalysis. Interestingly, the corresponding tyrosine residue in apo-GAC

(Y466W) is already inserted into the active site (Extended Data Fig. 4b), i.e., identical to its

position in the GAC and GLS2 filament structures. Because this catalytic residue is properly

positioned in the active site even in the apo- or 'resting' state of GAC, it might contribute, together

with the absence of ankyrin repeats (see below), to GAC having a significantly higher specific

activity compared to GLS2. Collectively, these unprecedented visualizations of the interactions

within the active sites of the GAC and GLS2 filaments upon the binding of substrate now provide

new insights into the mechanisms by which the glutaminase enzymes catalyze glutaminolysis.

A mechanism for glutaminase activation coupled to filament formation

The high-resolution glutaminase filament structures and RALS data presented here show how

glutaminase filament formation is coupled to catalysis. In the apo-enzymes, the lid loops do not

adopt a fixed conformation and are highly flexible. Such flexibility would likely introduce steric

clashes at the tetramer-tetramer interfaces within the filament, thereby preventing filament

formation (Fig. 5a, right panel). Upon binding to the enzyme, the substrate forms hydrogen bonds

with Glu381 in GAC and Glu314 in GLS2, resulting in the movement of these residues toward the

active site (Extended Data Figs. 5a, b, left panels). As shown for GLS2, this enables the alpha helix

containing residues Asn308 to Thr317 to move inward, resulting in additional interactions between

Thr310 and Gln185, and between Ser313 and Pro184 on the lid loop (Extended Data Fig. 5b, left

panel). These two sets of interactions, and those which occur between Tyr249 and Phe318 in GAC,

and between Tyr182 and Tyr251 in GLS2, with glutamine stabilize the lid loop in a conformation

that together with the activation loop effectively locks-in the substrate. This is necessary for

catalysis, as well as exposes the filament interface, therefore allowing additional tetramers to

contribute to the growth of the higher-order oligomer (Fig. 5a, left panel). Reciprocally, it is the

binding of tetramers to form the enzyme filament that restricts the flexibility of the lids and allows

them to adopt a conformation that closes over the substrate binding sites, thus enabling catalysis

to occur. Following a catalytic turnover, the product glutamate no longer interacts with the active

site tyrosine and glutamic acid residues and thus has a relatively low affinity for the enzyme. Upon

its dissociation, the activation loops and the lids becoming highly flexible again to await the next

substrate binding event. When all the available substrate is consumed, the lids are not able to be

restrained which disrupts the filament interface and gives rise to the dissociation of the filament

back to tetramers (Fig. 5a, right panel). The missing density of the lid loop of the terminal tetramer

in the cryo-EM map for the constitutively active GLS2 (K253A) filament represents a snapshot of

this detachment (Extended Data Fig. 5b, right panel) that occurs following catalytic turnover and

product release. However, in the case of the GAC (Y466W) mutant, which binds Pi and substrate

but cannot undergo catalysis, the filament persists (Fig. 2e) and now the density of the lid loop

within the terminal tetramer of the filament structure remains intact (Extended Data Fig. 5a, right

panel).

Given that filament formation is directly coupled to glutaminase activation, it might then be

expected that small molecule inhibitors that prevent the glutaminases from undergoing the

conformational transitions necessary for their activation would negatively impact their ability to

assemble into these higher order oligomeric structures. We therefore examined the effects of two

allosteric glutaminase inhibitors, UPGL0004 and 968, on filament formation. UPGL004 is a potent

member of the BPTES-class of compounds and specifically targets GAC[27,36] by binding to the

activation loops (Extended Data Fig. 1) and trapping them in a stable conformational state that

prevents their movements into the active sites and from working together with the lid loops to

establish the filament interface and lock-in the substrate to undergo catalytic turnover. Thus, we

found that while the vehicle control (DMSO) did not interfere with the ability of GAC (Y466W)

to form filaments within 30 seconds which persisted through 30 minutes as visualized by negative

stain EM (Figs. 5b, c), UPGL004 addition immediately blocked their assembly (Fig. 5d). The

BPTES-class of compounds are ineffective against GLS2 because of subtle differences in their

activation loops from those of GAC[24]. However, 968 is a small molecule inhibitor that effectively

inhibits GLS2 catalytic activity[9]. It appears to bind at the activation loops of GLS2 based on the

density we observed for the cryo-EM structure obtained for the 968-GLS2 complex, but without

overlapping the binding sites for either the BPTES-class of inhibitors nor Pi (Extended Data Fig.

6a), consistent with kinetic analyses from our laboratory[14]. Rather, upon binding to GLS2, 968

would block the salt bridge that forms between Arg258 on one activation loop and Glu250 on an

adjacent activation loop, thereby preventing the substrate from being locked in at the catalytic site.

This could also potentially weaken the interactions between GLS2 dimers to form tetramers, as we

have observed by cryo-EM 2D classifications where 968 treatment results in the appearance of

some dimeric enzyme particles (Extended Data Fig. 6b). Collectively, these effects would be

expected to prevent the formation of filaments that are required for optimal catalysis, which is

what we observed by negative stain EM in Figs. 5e and 5f. Likewise, mutation of a key residue

within the tetramer-tetramer interface necessary for filament formation (e.g. Asn 308 in GLS2),

significantly reduces their size (Fig. 5g) with a corresponding decrease in catalytic activity

(Extended Data Fig. 6c). Together, these findings further reinforce the conclusion that the

formation of glutaminase filaments is an essential step for enzyme activation and to achieve

maximal catalytic turnover.

The role of the ankyrin repeat motif in GLS2

The C-terminal region of GLS2 contains an ankyrin repeat motif consisting of three pairs of

helical bundles located on the distal ends of the tetramer, which is absent in GAC (Figs. 6a, b).

The ankyrin repeats in GLS2 appear to exert a negative regulation of catalytic activity, despite

being approximately 70 Å away from the active sites in each subunit. The KGA isoform of GLS

also contains ankyrin repeats and has a much lower specific activity than its C-terminal truncated

splice variant GAC; moreover, truncated forms of KGA lacking these repeats show increased

specific activity[33]. Each GLS2 dimer within a tetrameric unit shares an ankyrin repeat bundle

connected to the catalytic domain through a flexible linker. Interestingly, the ankyrin repeats are

highly asymmetric. Instead of showing a symmetrical occupancy of the area between the N-

terminal motifs of adjacent monomeric units within a tetramer, they show a preferential occupancy

on one particular side (Extended Data Fig. 7). As described above, the activation loops within each

dimer unit comprising the filament structures need to form an Arg-Glu salt bridge that enables the

activation loops to be correctly positioned for catalytic activity (i.e., Arg317-Glu325 in GAC and

Arg250-Glu258 in GLS2). This requires a scissor-like movement between two monomers within

the same dimer to position their activation loops into close proximity (Figs. 6a, b, insets at the

bottom; these movements are indicated by arrows; Extended Data Supplementary movie 1).

However, within the GLS2 structure, the ankyrin repeats appear to hinder this movement and

thereby may prevent the activation loops from adopting the positions necessary for maximal

catalytic capability (Fig. 6b). These effects may help contribute to GLS2 having a lower specific

activity compared to GAC.

DISCUSSION

The glutaminase enzymes GAC and GLS2 have both been implicated in cancer progression

due to the upregulation of their expression and catalytic activities[9,10,14,19,20,42]. The elevations in

glutamine metabolism triggered by these metabolic enzymes serve to compensate for the

uncoupling of glycolysis from the TCA cycle, known as the Warburg effect, by providing the

carbon sources necessary for the biosynthetic processes that enable cancer cells to undergo rapid

proliferation and survive various types of cellular stress that would otherwise kill normal healthy

cells. However, in some cancer contexts, GLS2 has been proposed to function as a tumor

suppressor[17,18]. While this raised the question of whether GLS2 might exhibit distinct structural

features from GAC, here we show that like GAC, GLS2 assumes a higher order filament-like

structure which is directly coupled to catalytic turnover.

In our efforts to learn more about the mechanisms by which the glutaminase enzymes catalyze

glutamine hydrolysis we had developed fluorescence spectroscopic readouts to monitor directly

the binding of the substrate glutamine in the presence and absence of the anionic activator Pi, as

well as probe the interactions of allosteric inhibitors while simultaneously assaying enzyme

activity[27,37,38,40]. These studies showed how the binding of substrate and anionic activators helped

to drive the transition between enzyme dimers to tetramers[25]. Given that the addition of an anionic

activator like Pi could induce the formation of glutaminase tetramers and was necessary for

enzymatic activity, it was assumed that the dimer-to-tetramer transition was responsible for

catalysis. Still, it was puzzling that when comparing the active sites of the dimeric and tetrameric

GAC species, there were no apparent differences in the positioning of the key catalytic residues,

nor was there any indication of how the activation loops contributed to enzyme activity. However,

we have been able to obtain important new insights into the mechanisms of glutaminase activation

and catalysis when examining the ability of GAC and GLS2 to form filament-like structures under

catalytic conditions, both through the use of cryo-EM and RALS. By performing cryo-EM

structural analyses of the GAC (Y466W) mutant which is still able to bind glutamine and Pi but is

defective in hydrolysis, we were able to capture a snapshot of the substrate- and activator-induced

conformational changes that lead to filament assembly. We found by negative stain EM, cryo-EM,

and RALS that the constitutively active GLS2 (K253A) mutant also formed higher-order filament-

like structures comprising several enzyme tetramers under catalytic turnover conditions, similar to

GAC. Importantly, we demonstrated that the lifetimes of the filament structures for both GAC and

GLS2 were directly coupled to catalysis, as they persisted as long as sufficient substrate was

available but then rapidly dissociated when the substrate was ultimately used up, confirming

suggestions that the formation of glutaminase filaments is directly linked to their catalytic

activity[29,31,39].

For all prior structures of the glutaminase isozymes, the activation loops were only visible

when the enzymes were bound to allosteric inhibitors of the BPTES family of compounds and

there were no indications of how they play a critical role in enzyme catalysis, especially given

their apparent distance from the active sites. There also was little indication of how the activation

loops communicated with the lids that form over the bound substrate, which is necessary for

enzyme catalysis[34,38]. However, the cryo-EM structures of activated GAC and GLS2 filaments

show activation loops that are well resolved and move toward the active site of each enzyme

subunit, positioning an essential tyrosine residue sufficiently close to a catalytically important

lysine residue to form a cation-π interaction. This leads to the de-protonation of an active site serine

residue, which is then poised for nucleophilic attack on the substrate glutamine. The glutaminase

filament structures also show that the lids undergo conformational changes relative to their

positions in the apo-enzyme (tetramer), resulting in the insertion of a key tyrosine residue into the

active site. These changes effectively close off each active site, locking the substrate glutamine

into the proper orientation for catalysis. Several additional hydrogen bonding interactions are

evident within the filament structures that help stabilize the interactions between the tetrameric

units and enable the activation loops to be inserted into the active sites. Upon catalytic turnover,

the release of the product glutamate relieves constraints on the activation loops and introduces

flexibility to the lid loops, which disrupts the interface between tetramers and results in the

dissociation of the filaments, thus accounting for the tight correlation between enzyme activity and

filament formation/dissociation.

Our structure-function studies of glutaminase filaments have shed light on how anionic

activators like Pi bind to activation loops of the enzymes and drive a dimer-to-tetramer transition

and filament assembly which are essential steps for catalysis. They also highlight how allosteric

inhibitors targeting the glutaminase enzymes and the presence of ankyrin repeats prevent the

activation loops from undergoing the conformational changes necessary for catalytic activity and

filament formation. Moreover, these findings now open the way to several interesting new lines of

study. For example, new therapeutic strategies might be developed to block the formation of these

higher-order oligomers. Identifying other potential activators of both GAC and GLS2 in cancer

cells that can induce filament formation will be important, given that the Pi levels necessary to

stimulate enzyme activity and filament formation might not be achievable under many

physiological contexts. It will also be interesting to see how post-translational modifications such

as lysine succinylation, which occur at specific sites on GAC and have been suggested to both

promote enzyme activation by stimulating oligomer (tetramer) formation[29] and to enhance

ubiquitination and degradation[43], affect filament formation, as well as whether the ankyrin repeats

are responsible for conferring GLS2 with functional capabilities distinct from those of GAC in

different cancers. Finally, an intriguing possibility will be to see whether the glutaminase enzymes

co-assemble in filamentous structures with other enzymes suggested to form higher order

oligomers such as glutamate dehydrogenase[28]. Such assemblies might more efficiently couple the

activities of members of a metabolic pathway by substrate channeling, with the products of one

enzyme being channeled to the next enzyme in the pathway to use as substrates.

REFERENCES

1.   Pavlova, N.N. and C.B. Thompson, *The Emerging Hallmarks of Cancer Metabolism.* Cell Metab, 2016. **23**(1): p. 27-47.

2.   Faubert, B., A. Solmonson, and R.J. DeBerardinis, *Metabolic reprogramming and cancer progression.* Science, 2020. **368**(6487).

3.   Hanahan, D., *Hallmarks of Cancer: New Dimensions.* Cancer Discov, 2022. **12**(1): p. 31-46.

4.   Warburg, O., *On the origin of cancer cells.* Science, 1956. **123**(3191): p. 309-14.

5.   Vander Heiden, M.G., L.C. Cantley, and C.B. Thompson, *Understanding the Warburg effect: the metabolic requirements of cell proliferation.* Science, 2009. **324**(5930): p. 1029-33.

6.   Lukey, M.J., W.P. Katt, and R.A. Cerione, *Targeting amino acid metabolism for cancer therapy.* Drug Discov Today, 2017. **22**(5): p. 796-804.

7.   Yang, C., et al., *Glutamine oxidation maintains the TCA cycle and cell survival during impaired mitochondrial pyruvate transport.* Mol Cell, 2014. **56**(3): p. 414-424.

8.   Leone, R.D., et al., *Glutamine blockade induces divergent metabolic programs to overcome tumor immune evasion.* Science, 2019. **366**(6468): p. 1013-1021.

9.   Lukey, M.J., et al., *Liver-Type Glutaminase GLS2 Is a Druggable Metabolic Node in Luminal-Subtype Breast Cancer.* Cell Rep, 2019. **29**(1): p. 76-88 e7.

10.   Mendez-Lucas, A., et al., *Identifying strategies to target the metabolic flexibility of tumours.* Nat Metab, 2020. **2**(4): p. 335-350.

11. Altman, B.J., Z.E. Stine, and C.V. Dang, *From Krebs to clinic: glutamine metabolism to cancer therapy.* Nat Rev Cancer, 2016. **16**(11): p. 749.

12. Harding, J.J., et al., *A Phase I Dose-Escalation and Expansion Study of Telaglenastat in Patients with Advanced or Metastatic Solid Tumors.* Clin Cancer Res, 2021. **27**(18): p. 4994-5003.

13. Shen, Y.A., et al., *Inhibition of glutaminolysis in combination with other therapies to improve cancer treatment.* Curr Opin Chem Biol, 2021. **62**: p. 64-81.

14. Wang, J.B., et al., *Targeting mitochondrial glutaminase activity inhibits oncogenic transformation.* Cancer Cell, 2010. **18**(3): p. 207-19.

15. Katt, W.P., et al., *Dibenzophenanthridines as inhibitors of glutaminase C and cancer cell proliferation.* Mol Cancer Ther, 2012. **11**(6): p. 1269-78.

16. Byun, J.K., et al., *Inhibition of Glutamine Utilization Synergizes with Immune Checkpoint Inhibitor to Promote Antitumor Immunity.* Mol Cell, 2020. **80**(4): p. 592-606 e8.

17. Zhang, C., et al., *Glutaminase 2 is a novel negative regulator of small GTPase Rac1 and mediates p53 function in suppressing metastasis.* Elife, 2016. **5**: p. e10727.

18. Suzuki, S., et al., *GLS2 Is a Tumor Suppressor and a Regulator of Ferroptosis in Hepatocellular Carcinoma.* Cancer Res, 2022. **82**(18): p. 3209-3222.

19. Dias, M.M., et al., *GLS2 is protumorigenic in breast cancers.* Oncogene, 2020. **39**(3): p. 690-702.

20. Saha, S.K., et al., *Multiomics Analysis Reveals that GLS and GLS2 Differentially Modulate the Clinical Outcomes of Cancer.* J Clin Med, 2019. **8**(3).

21. Katt, W.P., M.J. Lukey, and R.A. Cerione, *A tale of two glutaminases: homologous enzymes with distinct roles in tumorigenesis.* Future Med Chem, 2017. **9**(2): p. 223-243.

22.   Ferreira, I.M., et al., *Structure and activation mechanism of the human liver-type*
glutaminase GLS2. Biochimie, 2021. **185**: p. 96-104.

23.   Nguyen, T.T., W.P. Katt, and R.A. Cerione, *Alone and together: current approaches to*
targeting glutaminase enzymes as part of anti-cancer therapies. Future Drug Discov, 2023.
4(4): p. FDD79.

24.   DeLaBarre, B., et al., *Full-length human glutaminase in complex with an allosteric*
inhibitor. Biochemistry, 2011. **50**(50): p. 10764-70.

25.   Nguyen, T.T., et al., *High-resolution structures of mitochondrial glutaminase C tetramers*
indicate conformational changes upon phosphate binding. J Biol Chem, 2022. **298**(2): p.
101564.

26.   Gross, M.I., et al., *Antitumor activity of the glutaminase inhibitor CB-839 in triple-negative*
breast cancer. Mol Cancer Ther, 2014. **13**(4): p. 890-901.

27.   Milano, S.K., et al., *New insights into the molecular mechanisms of glutaminase C*
inhibitors in cancer cells using serial room temperature crystallography. J Biol Chem,
2022. **298**(2): p. 101535.

28.   Park, C.K. and N.C. Horton, *Structures, functions, and mechanisms of filament forming*
enzymes: a renaissance of enzyme filamentation. Biophys Rev, 2019. **11**(6): p. 927-994.

29.   Jiang, B., et al., *Filamentous GLS1 promotes ROS-induced apoptosis upon glutamine*
deprivation via insufficient asparagine synthesis. Mol Cell, 2022. **82**(10): p. 1821-1835 e6.

30.   Cassago, A., et al., *Mitochondrial localization and structure-based phosphate activation*
mechanism of Glutaminase C with implications for cancer metabolism. Proc Natl Acad Sci
U S A, 2012. **109**(4): p. 1092-7.

31.   Ferreira, A.P., et al., *Active glutaminase C self-assembles into a supratetrameric oligomer*

that can be disrupted by an allosteric inhibitor. J Biol Chem, 2013. **288**(39): p. 28009-20.

32.   Li, J., A. Mahajan, and M.D. Tsai, *Ankyrin repeat: a unique motif mediating protein-*

protein interactions. Biochemistry, 2006. **45**(51): p. 15168-78.

33.   Pasquali, C.C., et al., *The origin and evolution of human glutaminases and their atypical*

C-terminal ankyrin repeats. J Biol Chem, 2017. **292**(27): p. 11572-11585.

34.   Li, Y., et al., *The activation loop and substrate-binding cleft of glutaminase C are*

allosterically coupled. J Biol Chem, 2020. **295**(5): p. 1328-1337.

35.   Robinson, M.M., et al., *Novel mechanism of inhibition of rat kidney-type glutaminase by*

bis-2-(5-phenylacetamido-1,2,4-thiadiazol-2-yl)ethyl sulfide (BPTES). Biochem J, 2007.

406(3): p. 407-14.

36.   Huang, Q., et al., *Characterization of the interactions of potent allosteric inhibitors with*

glutaminase C, a key enzyme in cancer cell glutamine metabolism. J Biol Chem, 2018.

293(10): p. 3535-3545.

37.   Stalnecker, C.A., et al., *Mechanism by which a recently discovered allosteric inhibitor*

blocks glutamine metabolism in transformed cells. Proc Natl Acad Sci U S A, 2015. **112**(2):

p. 394-9.

38.   Li, Y., et al., *Mechanistic Basis of Glutaminase Activation: A KEY ENZYME THAT*

PROMOTES GLUTAMINE METABOLISM IN CANCER CELLS. J Biol Chem, 2016.

291(40): p. 20900-20910.

39.   Moller, M., et al., *Small angle X-ray scattering studies of mitochondrial glutaminase C*

reveal extended flexible regions, and link oligomeric state with enzyme activity. PLoS One,

2013. **8**(9): p. e74783.

40.   Stalnecker, C.A., J.W. Erickson, and R.A. Cerione, *Conformational changes in the*

activation loop of mitochondrial glutaminase C: A direct fluorescence readout that

distinguishes the binding of allosteric inhibitors from activators. J Biol Chem, 2017.

292(15): p. 6095-6107.

41.   Brown, G., et al., *Functional and structural characterization of four glutaminases from*

Escherichia coli and Bacillus subtilis. Biochemistry, 2008. **47**(21): p. 5724-35.

42.   Gao, P., et al., *c-Myc suppression of miR-23a/b enhances mitochondrial glutaminase*

expression and glutamine metabolism. Nature, 2009. **458**(7239): p. 762-5.

43.   Greene, K.S., et al., *SIRT5 stabilizes mitochondrial glutaminase and supports breast*

cancer tumorigenesis. Proc Natl Acad Sci U S A, 2019.

METHODS

Recombinant glutaminase expression and purification

An N-terminal His-tagged form of the full-length human liver-type glutaminase isoform GLS2 without the mitochondrial localization sequence (residue 38-602) was cloned into the pET28a plasmid as previously described[36]. Site-directed mutagenesis was performed using Phusion DNA polymerase (New England Biolabs, NEB). Protein expression was carried out first by transforming constructs into *E. Coli* BL21 (DE3) competent cells (NEB), which were then grown in LB media overnight with 50 μg/mL kanamycin. The starter cultures were subsequently inoculated at ~1:100 ratio in 6 L cultures with the same antibiotic concentration and left shaking at 37°C, 180 rpm for 3-4 hours until the OD600 reached between 0.6 and 0.8. The flasks were then chilled at 4°C for 1-2 hours before induction with 30 μM IPTG (isopropyl β-D-1-thiogalactopyranoside) and shaking at room temperature (RT), 180 rpm for 16-18 hours. Cells were collected by centrifugation at 5,000xg for 10 minutes and frozen before resuspension in 150 mL lysis buffer (50 mM Tris-HCl pH 8.5, 500 mM NaCl, 10% glycerol), supplemented with protease inhibitor cocktail (Roche). Cells were lysed with lysozyme and mechanical sonication, and DNase I was added to reduce the mixture's viscosity. The soluble fractions were separated from the debris by ultra-centrifugation (Ti45 rotor, 40,000 rpm for 45 minutes). The lysate was then loaded onto Co^{2+}-charged TALON resin (GoldBio), equilibrated previously with the wash buffer (50 mM Tris-HCl pH 8.5, 10 mM NaCl, 10 mM imidazole). The protein that bound to the column was washed with the wash buffer and eluted with wash buffer supplemented with 320 mM imidazole. Further purification was performed by anion exchange chromatography using HiTrap Q HP column (Cytiva) and size-exclusion chromatography using Superdex 200 pg 16/600 column (GE Healthcare). Human GAC preparations were prepared as previously described[25,27]. Proteins were kept in 20 mM Tris-HCl pH

8.5, 150 mM NaCl, snap-frozen in liquid nitrogen, and stored at -80°C. Protein concentrations

were determined by absorbance at 280 nm using extinction coefficients calculated using the

Expasy ProtParam tool.

Right-angle light scattering (RALS)

Frozen glutaminase aliquots were thawed on ice and diluted to a final concentration of 2 μM

in 1 mL of gel filtration buffer (20 mM Tris-HCl pH 8.5, 150 mM NaCl) added in a 1.2 mL cuvette

and inserted into a Varian Cary Eclipse fluorimeter. The cuvette temperature was set to 25°C and

stirred with a magnetic stir bar. The signal was recorded using excitation and emission wavelengths

of 340 nm (5 nm bandpass). The samples were incubated until the signal was stable (~2-3 mins),

then 100 μL of glutamine from a 200 mM stock solution were added to initiate the reaction.

Subsequently, 100 μL of inorganic phosphate (1 M) or glutamate (200 mM) were added for further

analysis. The raw scattering traces were filtered to improve the signal-to-noise ratio using SciPy

(version 1.9.2) and plotted using Matplotlib (version 3.7.1).

Negative stain Electron Microscopy

Formvar/carbon film 200 mesh copper grids (Electron Microscopy Sciences, EMS) were

plasma cleaned by using PELCO easiGlow system (TED PELLA). Glutaminase (final

concentration: 1 μM) was mixed with the substrate glutamine (final concentration: 20 mM).

Inorganic phosphate (K_2HPO_4 final concentration: 50 mM) was added to the mixture when needed.

The mixture was incubated at RT for 30 seconds. Then, 10 μL of the mixture was applied onto the

grids for a 60-second incubation, and the excess protein solution was blotted with filter paper. Ten

μL of 2% uranyl acetate were then applied to the grid twice for two 30-second stainings followed

by blotting. The grids were air-dried for 5 minutes and visualized by Thermo Fischer F200Ci

electron microscope at 120 keV.

Cryo-EM grids preparation and images acquisition for apo-GLS2

Four μL of 5 μM GLS2 were applied to plasma-cleaned grids (300 mesh UltraAu Foil

R1.2/1.3, Electron Microscopy Sciences, EMS) for a 30-second incubation. The excess solution

was blotted for 4-6 seconds with filter paper by a Vitrobot mark IV (Thermo Fischer Scientific).

Grids were subsequently vitrified in liquid ethane and stored in liquid nitrogen. Cryo-EM single

particle micrographs of apo-GLS2 were collected on a Talos Arctica (Thermo Fischer Scientific)

at the Cornell Center for Materials Research. The microscope was operated at 200 keV at 63,000x

nominal magnification using a Gatan K3 direct electron camera in super-resolution mode, with a

Gatan GIF Quantum LS Imaging energy filter, corresponding to a physical pixel size of 1.23 Å/pix.

Given the orientation bias of the sample, a non-tilted dataset of 1,171 images and a 30-degree tilted

dataset of 488 images were collected and merged for further data processing. The images were

obtained with a defocus range of -0.8 to -2.0 um by EPU (Thermo Fischer Scientific). Each stack

movie was recorded with 50 frames for a total dose of ~50 e-/Å^2.

Cryo-EM grids preparation and images acquisition for GLS2 (K253A) filament

Human GLS2 (K253A), 3 μM, was mixed with 20 mM glutamine to initiate filament

formation. The mixture was applied to plasma-cleaned grids (300 mesh Quantifoil Au R1.2/1.3,

EMS) for 30-second incubations. The excess solution was blotted for 5-7 seconds with filter paper

by a Vitrobot mark IV. Grids were subsequently vitrified in liquid ethane and stored in liquid

nitrogen. Cryo-EM single particle micrographs of apo-GLS2 were collected on a Titan Krios

(Thermo Fischer Scientific) at The New York Structural Biology Center. The microscope was

operated at 300 keV at 81,000x nominal magnification using a Gatan K3 direct electron camera in

counting mode, corresponding to a physical pixel size of 1.058 Å/pix. Thirty-eight hundred and

four images were obtained with a defocus range of -0.8 to -2.0 mm by Leginon[44]. Each stack movie

was recorded with 40 frames for a total dose of ~50 e-/Å^2.

Cryo-EM grids preparation and images acquisition for GAC (Y466W) filament

Human GAC (Y466W), 8 μM, was mixed with 20 mM glutamine and 1M inorganic phosphate

to initiate filament formation. The mixture was applied to plasma-cleaned grids (300 mesh

Quantifoil Cu R1.2/1.3, EMS) for 30-second incubations. The excess solution was blotted for 2-3

seconds with filter paper by a Vitrobot mark IV. Grids were subsequently vitrified in liquid ethane

and stored in liquid nitrogen. Cryo-EM single particle micrographs of GAC (Y466W) filament

were collected on a Talos Arctica (Thermo Fischer Scientific) at the Cornell Center for Materials

Research. The microscope was operated at 200 keV at 63,000x nominal magnification using a

Gatan K3 direct electron camera in counted mode, with a Gatan GIF Quantum LS Imaging energy

filter, corresponding to a physical pixel size of 1.31 Å/pix. Thirty-nine hundred images were

obtained with a defocus range of -0.8 to -2.0 mm by SerialEM[45]. Each stack movie was recorded

with 50 frames for a total dose of ~50 e-/Å^2.

Cryo-EM data processing

For the apo-GLS2 map, dose-fractionated image stacks were subjected to motioncor2 wrapped

in RELION 4.0.0.[46-48] through SBGrid, followed by patch CTF estimation in CryoSPARC

V4.0.0.[49]. The motion-corrected micrographs with CTF better than 4 Å were passed for later

processing. A blob picker in CryoSPARC was used to generate the templates for further Topaz

picking[50], where ~1M particles were picked. The particles were subjected to 3D classification and

heterogenous refinement, and only the best classes were kept. The remaining ~329K particles were

subjected to homogenous and heterogenous refinement. The final reconstruction with C2

symmetry yields a map with a resolution of 3.12 Å determined by FSC. The workflow is provided

in Extended Data Fig. 8. Cryo-EM maps are visualized in ChimeraX[51].

For the GAC (Y466W) filament structure, dose-fractionated image stacks were subjected to

patch motion correction, followed by patch CTF estimation in CryoSPARC V4.2.1. The motion-

corrected micrographs with CTF better than 4 Å were passed for later processing. The template-

free filament trace job in CryoSPARC picks ~397K particles for 2D classification. About 180K

particles were extracted and subjected to 3D classification. The best ~45k particles were

reconstructed with C2 symmetry and refined to yield a 3.35 Å map determined by FSC. The

workflow is provided in Extended Data Fig 9.

For the GLS2 (K253A) filament structure, dose-fractionated image stacks were subjected to

patch motion correction, followed by patch CTF estimation in CryoSPARC V4.0.0. The motion-

corrected micrographs with CTF better than 4 Å were passed for later processing, which yielded

3,640 micrographs. Manual picking in CryoSPARC was used on 40 micrographs to pick ~600

particles manually, which yielded five classes of templates by 2D classification. The selected

template was used for the filament trace job in CryoSPARC to template pick particles. About 333K

particles were extracted and subjected to 3D classification. The best ~48k particles were

reconstructed and refined to yield a 3.3 Å map determined by FSC. The workflow is provided in

Extended Data Fig 10.

Model building and refinement

The initial apo-GLS2, GAC (Y466W), and GLS2 (K253A) model was adapted from the PDB entry 4BQM for the catalytic domain and 5U0K for the ankyrin repeats. First, all models were manually docked into the map by UCSF ChimeraX. Next, the model was fitted into the map by ISOLDE for the overall adjustment[52], then underwent multiple rounds of automatic refinement using Phenix real space refinement[53], and manual adjusting and building ligands using Coot[54]. Finally, validation was performed with MolProbity[55]. The final refinement statistics is provided in Extended Data Table 1.

Data availability

The cryo-EM density maps and the atomic model for apo-GLS2, GAC (Y466W) and the GLS2 (K253A) filament structure reported in this paper have been deposited in EM Data Bank (EMDB: EMD-40920, EMD-40918, and EMD-40950) and the Protein Data Bank (PDB: 8SZL, 8SZJ, and 8T0Z), respectively.

Code availability

No custom computer code was used in this study.

44.   Suloway, C., et al., *Automated molecular microscopy: the new Leginon system.* J Struct Biol, 2005. 151(1): p. 41-60.

45.   Mastronarde, D.N., *Automated electron microscope tomography using robust prediction of specimen movements.* J Struct Biol, 2005. **152**(1): p. 36-51.

46.   Morin, A., et al., *Collaboration gets the most out of software.* Elife, 2013. **2**: p. e01456.

47.   Scheres, S.H., *RELION: implementation of a Bayesian approach to cryo-EM structure*
determination. J Struct Biol, 2012. **180**(3): p. 519-30.

48.   Zheng, S.Q., et al., *MotionCor2: anisotropic correction of beam-induced motion for*
improved cryo-electron microscopy. Nat Methods, 2017. **14**(4): p. 331-332.

49.   Punjani, A., et al., *cryoSPARC: algorithms for rapid unsupervised cryo-EM structure*
determination. Nat Methods, 2017. **14**(3): p. 290-296.

50.   Bepler, T., et al., *Positive-unlabeled convolutional neural networks for particle picking in*
cryo-electron micrographs. Nat Methods, 2019. **16**(11): p. 1153-1160.

51.   Pettersen, E.F., et al., *UCSF ChimeraX: Structure visualization for researchers, educators,*
and developers. Protein Sci, 2021. **30**(1): p. 70-82.

52.   Croll, T.I., *ISOLDE: a physically realistic environment for model building into low-*
resolution electron-density maps. Acta Crystallogr D Struct Biol, 2018. **74**(Pt 6): p. 519-
530.

53.   Liebschner, D., et al., *Macromolecular structure determination using X-rays, neutrons and*
electrons: recent developments in Phenix. Acta Crystallogr D Struct Biol, 2019. **75**(Pt 10):
p. 861-877.

54.   Emsley, P., et al., *Features and development of Coot.* Acta Crystallogr D Biol Crystallogr,
2010. **66**(Pt 4): p. 486-501.

55.   Williams, C.J., et al., *MolProbity: More and better reference data for improved all-atom*
structure validation. Protein Sci, 2018. **27**(1): p. 293-315.

Acknowledgments We thank Drs. Katherine Spoth, Mariena Silvestry-Ramos, Kasahun Neselu,

and Edward Eng for help with EM instrumentation. We would also like to thank Drs. Brian R.

Crane and Sekar Ramachandran for the helpful feedback on the structural analysis, and Samuel L.

Deck and Megan Xu for helping produce glutaminase mutants. This work was supported by grants

from the NIH (R35GM122575 and R01CA201402). The acquisition of the human apo-GLS2 and

mouse GAC (Y466W) structures used the Cornell Center for Materials Research Shared Facilities,

which is supported by NSF (DMR- 1719875). The collection of negative stain images relied on

using an instrument supported by the NIH through award S10OD030470-01. The acquisition of

the GLS2 (K253A) structure was performed at the National Center for CryoEM Access and

Training (NCCAT) and the Simons Electron Microscopy Center located at the New York

Structural Biology Center, supported by the NIH Common Fund Transformative High-Resolution

Cryo-Electron Microscopy program (U24 GM129539) and by grants from the Simons Foundation

(SF349247) and NY State Assembly.

Author Contributions All authors contributed to each aspect of the study, but primarily, T-

T.T.N., S.K.M, C.A. and S.F. developed methods to express and purify the glutaminase and

mutants; C.A. and S.F. developed and performed the RALS assay; S.F. performed the activity

assay; S.F. prepared the samples for negative stain EM and cryo-EM, and collected and processed

the EM data. S.F. built the apo-GLS2, GAC Y466W filament and GLS2 K253A filament

structures; S.F., C.A., T-T.T.N., S.K.M. and R.A.C. analysed the structures; S.F., C.A., T-T.T.N.,

S.K.M. and R.A.C. wrote the manuscript; R.A.C. conceived of and oversaw the project.

Competing interests The authors declare no competing interests.

Additional information

Supplementary information is available for the paper.

Correspondence and requests for materials should be addressed to Richard A. Cerione.

Reprints and permissions information is available at www.nature.com/reprints.

Figure 1. The cryo-EM structure of human full-length apo-GLS2 and its comparison with

the X-ray crystal structures of the GLS isoforms.

(a) The structures of tetrameric GAC, KGA, and GLS2 (from left to right). One monomer in each

tetramer is highlighted in solid color. The glutaminase catalytic domain (dashed box in (A)). **(b)**

Zoomed in view of the catalytic domain showing the activation loop (light yellow), the catalytic

site (dashed box) and the lid loop (dark yellow). **(c)** The conserved catalytic residues of GLS2

(blue) and GAC (PDB ID: 3UNW, light purple). Residues are numbered according to human

GLS2 and GAC numbering is shown in parentheses.

a GAC + Pi + Q (at 1 min)
Start of reaction, filaments form

b GLS2 + Pi + Q (at 1 min)
Start of reaction, filaments form

c GLS2 + Pi +Q (at 10 mins)
End of reaction, filaments dissociate

Figure 2. The formation of the glutaminase filaments is directly coupled to catalytic

activity.

(a) Negative stain EM of GAC (1 μM) with inorganic phosphate (Pi, 100 mM) one minute after

adding glutamine (Q, 20 mM). **(b)** Negative stain EM of GLS2 (1 μM) with Pi (100 mM) one

minute after adding glutamine (20 mM). **(c)** Negative stain EM of GLS2 (1 μM) with Pi (100

mM) 10 minutes after adding glutamine (20 mM). **(d)** Right-angle light scattering (RALS) of

GLS2 (2 μM) with Pi (100 mM) after subsequent additions of glutamine (20 mM), or glutamate

(20 mM). **(e)** RALS of GAC (Y466W) with Pi (100 mM) and glutamine (20 mM).

Figure 3. Structures of the human GAC (Y466W) filament and the human GLS2 (K253A) filament captured under catalytic turnover conditions.

(a) Structure of the GAC (Y466W) filament showing a five tetramer stretch of the higher-order oligomer with a helical rise of 68 Å. Tetramers are colored alternately either orange and red or green. The orange and red tetramers are colored to distinguish between monomers in each tetramer. The filament interface and the N-terminal contacts are highlighted. **(b)** The GAC (Y466W) filament viewed down the filament axis. The first two tetramers are outlined with red and black silhouettes, sequentially. The dashed line indicates the longest dimension of each tetramer and the 51° rotation around the filament axis between tetramers. **(c)** Structure of the GLS2 (K253A) filament showing a five tetramer stretch of the higher-order oligomer with a helical rise of 66 Å. Tetramers are colored alternately either purple and yellow or gray. The purple and yellow tetramers are colored to distinguish between monomers in each tetramer. The filament interface is highlighted. **(d)** The GLS2 (K253A) filament viewed down the filament axis. The first two tetramers are outlined with purple and black silhouettes, sequentially. The dashed line indicates the longest dimension of each tetramer and the 48° rotation around the filament axis between tetramers. **(e-f)** The position of the helical filament interface, the activation loop, the catalytic site, and the lid loop in GAC (**e**, red) and GLS2 (**f**, yellow). The next tetramer in the filament is colored green in GAC and gray in GLS2. The helical filament interface is shown in solid colors. **(g-h)** The interaction network at the helical filament interface between tetramers for GAC (**g**, red and green) and GLS2 (**h**, yellow and gray). The close proximity of the catalytic site to the helical filament interface is highlighted with a dashed box.

Figure 4. Structures of the GAC and GLS2 filaments reveal allosteric conformational changes within the glutaminase tetrameric units during catalysis.

(a-b) The phenylalanine-tyrosine lock in GAC (Phe318 and Tyr249) and the tyrosine lock in GLS2 (Tyr251 and Tyr182). Conserved catalytic residues are labeled, and the electron densities of the activation loop and the lid loop are shown as mesh. **(c)** The movement of Arg387 between apo-GAC (PDB ID: 3UNW, Arg387 is shown in light gray) and the GAC (Y466W) filament (pink) is shown with an arrow. The interactions between inorganic phosphate (Pi) and residues Lys320 and Arg387 near the activation loop in the GAC (Y466W) filament are highlighted. Salt bridges between GAC dimers within a tetramer (pink and salmon) are labeled as dashed yellow lines. **(d)** The movement of Arg320 between apo-GLS2 (Arg320 is shown in cyan) and the GLS2 (K253A) filament (yellow) is shown with an arrow. Salt bridges between GLS2 dimers within a tetramer (yellow and purple) are labeled as dashed yellow lines. **(e)** The tyrosine lock in GLS2 consists of Tyr251 from the activation loop (yellow) and Tyr182 from the lid loop (orange). Glutamine is shown in green and the cation-π interaction between Lys222 and Tyr251 is labeled. **(f)** The movement of conserved catalytic residue Tyr399 between apo-GLS2 (light blue) and the GLS2 filament (light yellow) is shown with an arrow. The activation loops of two adjacent monomers within the GLS2 filament are colored yellow (subunit A) and purple (subunit B). Interactions between key residues consisting of hydrogen bonds and salt bridges are shown as yellow dashed lines.

a

Substrate bound

Filament forms

adjacent tetramer

Stablized lid loop

Lid lock

Substrate lock

Exposed interface

Product released (apo)

adjacent tetramer

Destablized lid loop

Destablized activation loop

putative exit of product

Clash at interface

Filament dissociates

b 200 nm

GAC Y466W + Pi + DMSO
(30 seconds after adding substrate)

c 100 nm

GAC Y466W + Pi + DMSO
(30 minutes after adding substrate)

d 100 nm

GAC Y466W + Pi + 1 uM UPGL0004
(30 seconds after adding substrate)

e 200 nm

GLS2 + Pi
(30 seconds after adding substrate)

f 100 nm

GLS2 + Pi + 10 uM 968
(30 seconds after adding substrate)

g 200 nm

GLS2 N308A + Pi
(30 seconds after adding substrate)

Figure 5. The proposed mechanism of glutaminase filament formation and dissociation.

(a) The proposed mechanism for glutaminase filament formation and dissociation. Left: When substrate binds, the lid (yellow) forms a 'lid lock', and the helical interface is exposed (shown in gray), allowing the adjacent tetramer to extend the filament. Right: When product is released, the lid loop becomes flexible, displacing the adjacent tetramer, and blocking the helical interface. **(b)** Negative stain EM of GAC (Y466W) (1 µM) with inorganic phosphate (Pi, 200 mM) and DMSO 30 seconds after adding glutamine (substrate, 20 mM). **(c)** Negative stain EM of GAC (Y466W) (1 µM) with Pi (200 mM) and DMSO 30 minutes after adding substrate (20 mM). **(d)** Negative-stain EM of GAC (Y466W) (1 µM) with Pi (200 mM) and UPGL0004 (1 µM) 30 seconds after adding substrate (20 mM). **(e)** Negative stain EM of GLS2 (1 µM) with Pi (50 mM) 30 seconds after adding substrate (20 mM). **(f)** Negative stain EM of GLS2 (1 µM) with Pi (50 mM) and compound 968 (10 µM) 30 seconds after adding substrate (20 mM). **(g)** Negative stain EM of GLS2 (N308A) (1 µM) with Pi (50 mM) 30 seconds after adding substrate (20 mM).

Figure 6. The proposed mechanism for regulation of GLS2 catalytic activity by the C-

terminal ankyrin repeats.

(a) The unrestricted scissor-like movement of the N-terminal domain between monomers within a

dimer of the GAC filament (orange and magenta) and apo-GAC (light gray) after filament

formation. The arrows indicate the direction of conformational changes. Inset on the bottom: the

scissor-like movement brings Glu325 and Arg 317 from adjacent GAC monomers within a dimer

into close proximity and helps them form a salt bridge to stabilize the activation loops. **(b)** The

scissor-like movement of the N-terminal domain between monomers within a dimer of the GLS2

filament (purple and yellow) and apo-GLS2 (light blue) is restricted by the ankyrin repeats (solid

box). The arrows indicate the direction of conformational changes. The formation of the salt bridge

between Glu258 and Arg250 from adjacent GLS2 monomers within a dimer is burdened by the

blocking of ankyrin repeats, which prevents the activation loops contact at close proximity. Also,

see Extended Data Supplementary movie 1.

Substrate

GLS2 Y182

GAC Y249

GLS2 Y251

GAC F318

Substrate lock

Activation loop

GAC Y466W filament (this study)

GLS2 K253A filament (this study)

GLS inhibitors

F318

KGA+BPTES 3VOZ
GAC+BPTES+Glu 3UO9
GAC+UPGL00004 5WJ6

Extended Data Figure 1. The activation loops of the GAC (Y466W) filament (salmon), GLS2

(K253A) filament (yellow) and GAC bound to the allosteric inhibitors BPTES, alone or together

with glutamate, and UPGL0004 (purple, lavender and pink, respectively).

Extended Data Figure 2. (a) Right-angle light scattering (RALS) of GLS2 (2 µM) after the

addition of glutamine (Q, 20 mM), inorganic phosphate Pi (100 mM), or Q (20 mM) and Pi (100

mM). **(b)** Negative stain EM of GAC (K320A) (1 µM) after adding Q (20 mM). **(c)** Negative stain

EM of GLS2 (K253A) (1 µM) after adding Q (20 mM). **(d)** RALS of GLS2 (K253A) (2 µM) after

the addition of glutamate (E, 20 mM) and Q (20 mM) (black), Q (20 mM) (red), or Q (20 mM)

and Pi (100 mM) (blue). **(e)** RALS of GLS2 (2 µM) with Pi (100 mM) after the addition of

increasing concentrations of glutamine. **(f)** The average number of tetramers per filament plotted

against substrate (glutamine) concentration. The filament length was determined through an

analysis of negative stain electron microscopy images (n=3) with ImageJ. The width of each GLS2

tetramer is 64 Å, as measured from the high resolution cryo-EM structure of the GLS2 filament.

X-axis: the concentration of substrate (glutamine) in mM; left Y-axis: the number of tetramers per

filament; right Y-axis: the average length of filaments in Å. Circles indicate the mean value, and

error bars are shown for the standard error.

821 **Extended Data Figure 3.** Left panel: the electron density of inorganic phosphate (Pi) bound near

822 the activation loop in the GAC (Y466W) filament. Right panel: the phenylalanine-tyrosine lock

823 in GAC consisting of Phe318 from the activation loop (magenta) and Tyr249 from the lid loop

824 (magenta). The substrate, glutamine, is shown in green. Lys289 interacts with Phe318 from the

825 activation loop through a cation-π interaction.

827   **Extended Figure 4. (a)** The anionic activator, inorganic phosphate (Pi), interacts with Lys320 of

828   the activation loop, and Arg387, Tyr394, and Lys398 from the dimer-dimer interface of GAC

829   (Y466W), shown as yellow dashed lines. **(b)** The conserved residue in the active site, Tyr466, is

830   already inserted into the active site even without the new interaction between the Glu325 with

831   Asp467 and Arg317 that occurs within the filament structure.

Extended Data Figure 5. (a) Left panel: the lid loop of GAC is stabilized in the active conformation by a lid lock involving the lid loop and residues adjacent to the filament helical interface between tetramers within the filament. Hydrogen bonds are indicated by yellow dashed lines. Apo-GAC is light gray, and the GAC filament is shown in magenta. The adjacent tetramer in the filament is shown in green. The arrow indicates the direction of the conformational change. Right panel: the cryo-EM density map of the lid loop of the terminal subunit in the GAC filament. The lid loop in the catalytic-deficient GAC (Y466W) filament has strong density in its terminal subunit. **(b)** Left panel: the lid loop of GLS2 is stabilized in the active conformation through a lid lock. Hydrogen bonds are indicated by yellow dashed lines. Apo-GLS2 is light blue and the GLS2 filament is shown in yellow and orange. The adjacent tetramer that is forming the helical interface is shown in dark gray. The arrow indicates the direction of the conformational change. Right panel: the cryo-EM density map of the lid loop of the terminal subunit in the GLS2 filament. The weak density of the lid loop in the terminal tetramer of GLS2 suggests increased flexibility compared to the lid loops in the other tetramers that make up the filament.

a

GLS2 (wild type) dimer with 968

90°

968 density

b

Untreated tetramer → +968 → Treated dimer

Activation loop (subunit A)
Activation loop (subunit B)
GLS2 R250
GLS2 E258
968
Pi
BPTES (3UO9)

c

GAC (K320A)
GLS2 (wild type)
GLS2 (N308A)
GAC (Y466W)

Absorbance intensity (A.U.)

Time (mins)

Extended Data Figure 6. (a) Upper left: the 3D reconstruction of wild type GLS2 bound to 968. Bottom left inset: The electron density for 968 suggests that it interacts with the activation loops. Bottom right inset: The relative positions of 968, BPTES, and inorganic phosphate (Pi) are shown. Notably, 968 interferes with the interaction between Arg250 on one activation loop and Glu258 on an adjoining activation loop. **(b)** cryo-EM 2D classification of GLS2 treated with 968 shows that approximately 10% of the particles can be found as dimers in the presence of 968, whereas no dimers are detected in the absence of 968. Each monomer depicting the GLS2 tetramer and dimer is outlined in red. **(c)** The fluorescence trace of the activity assay of GLS2 (N308A) compared with wild type GLS2, GAC (Y466W), and GAC (K320A). Assays of glutaminase activity were performed as previously described[25]. X-axis: time in minutes; Y-axis: the intensity of the absorbance at 340 nm, which is proportional to the yield of catalytic product. The concentrations of GLS2 (N308A), wild type GLS2 and GAC (Y466W) used in these experiments were 0.6 μM, and the concentration of GAC (K320A) was 0.12 μM. The concentration of glutamine in these assays was 20 mM. The concentration of inorganic phosphate was 50 mM when assaying the catalytic activity of GLS2 and GLS2 (N308A), and 200 mM when assaying GAC (Y466W) and GAC (K320A).

868   **Extended Data Figure 7.** The asymmetric positioning of the ankyrin repeats in apo-GLS2 from

869   different subunits shown in orange and yellow.

1,171 0˚-tilt movies

488 30˚-tilt movies

Patch motion-correction → Patch CTF estimation → Keep <4Å → Blob picking / Topaz picking

~1M particles

2D classification

0˚-tilt dataset

30˚-tilt dataset

3D classification

Class 01 ~261K

Class 02 ~218K

Class 03 ~315K

Class 04 ~219K

Heterogenous refinement

Class 01 ~108K

Class 02 ~329K

Multiple rounds of homo/non-uniform refinement
Symmetry: C2

Representative micrograph

GSFSC Resolution: 3.12Å

No Mask (3.7Å)
Spherical (3.6Å)
Loose (3.4Å)
Tight (3.1Å)
Corrected (3.1Å)

3.12 Å map (EMDB-xxxxx)

Local resolution (Å)

3.2 3.4 3.6 3.8 4.0 4.1

873   **Extended Data Figure 8.** The cryo-EM data processing workflow of apo-GLS2.

3,900 movies →(Patch motion-correction)→ Patch CTF estimation →(Keep <4Å)→ Manual picking / Filament tracer

~397K particles → 2D classification

~180K particles → 3D classification

Class 01 ~12K   Class 02 ~112K   Class 03 ~13K   Class 04 ~42K

Homogenous refinement
Non-uniform refinement
Helical refinement

3.6 Å map (not shown) →(3D classification)→

Class 01 ~4K   Class 02 ~3K   Class 03 ~60K   Class 04 ~45K

Homogenous refinement
Non-uniform refinement
Helical refinement (C2 Symmetry)
CTF refinement

GSFSC Resolution: 3.35Å
No Mask (4.1Å)
Loose (3.7Å)
Tight (3.3Å)
Corrected (3.4Å)

3.35 Å map (EMDB-xxxxx)

Representative micrograph

877 **Extended Data Figure 9.** The cryo-EM data processing workflow of GAC Y466W filament.

880   **Extended Data Figure 10.** The cryo-EM data processing workflow of GLS2 K253A filament.

881   **Extended Data Table 1. Cryo-EM data collection, refinement and validation statistics.**
882

	Apo GLS2 (EMD-40920) (PDB 8SZL)	GAC(Y466W) filament (EMD-40918) (PDB 8SZJ)	GLS2(K253A) filament (EMD-40950) (PDB 8T0Z)
Data collection and processing			
Magnification	63kx	63kx	81kx
Voltage (kV)	200	200	300
Electron exposure (e–/Å^2)	50	50	50
Defocus range (μm)	-0.8~-2.0	-0.8~-2.0	-0.8~-2.0
Physical pixel size (Å)	1.23	1.31	1.058
Symmetry imposed	C2	C2	C1
Initial particle images (no.)	~1M	~397K	~333K
Final particle images (no.)	~329K	~45K	~49K
Map resolution (Å)	3.12	3.35	3.26
FSC threshold	0.143	0.143	0.143
Map resolution range (Å)	3.0-5.0	3.1-5.0	3.0-5.0
Refinement			
Initial model used (PDB code)	4BQM	4BQM	4BQM
Model resolution (Å)	3.12	3.35	3.26
FSC threshold	0.143	0.143	0.143
Model resolution range (Å)	3.0-5.0	3.0-5.0	3.0-5.0
Map sharpening B factor (Å^2)	164.90	112.1	211.56
Model composition			
Non-hydrogen atoms	15,095	36,204	47,436
Protein residues	1,961	4,656	6,108
Ligands	0	24	12
B factors (Å^2)			
Protein	164.90	189.44	211.56
Ligand	N/A	173.39	N/A
R.m.s. deviations			
Bond lengths (Å)	0.003	0.002	0.005
Bond angles (°)	0.553	0.647	0.671
Validation			
MolProbity score	2.01	1.89	2.02
Clashscore	11.86	10.20	11.00
Poor rotamers (%)	0.06	0.67	0.69
Ramachandran plot			
Favored (%)	93.70	94.74	92.59
Allowed (%)	6.30	5.13	7.31
Disallowed (%)	0.00	0.13	0.10

883