1   <u>Article type (Letter)</u>
2
3   <u>Article title</u>
4   Facultative symbiont virulence determines horizontal transmission rate without host strain
5   specificity
6

7   Suegene Noh[1]*, Emily R. Larson[1], Rachel M. Covitz[2], Anna Chen[1], Prachee R. Mazumder[1], Ron
8   F. Peck[1], Marisa C. Hamilton[3], Robert A. Dettmann[4]
9
[1] Biology Department, Colby College, Waterville, Maine, USA
[2] Department of Developmental, Molecular, and Chemical Biology, Tufts University School of
Medicine, Boston, Massachusetts, USA
[3] University Program in Genetics and Genomics, Duke University, Durham, North Carolina, USA
[4] Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA
* Corresponding author: suegene.noh@colby.edu

16
17   <u>Running title</u>
18   Virulence-transmission tradeoff in a facultative symbiosis
19
20   <u>Keywords</u>
21   Coevolution, facultative symbiosis, fitness, flow cytometry, horizontal transmission, protist,
22   virulence factor
23
24   <u>Abstract (211)</u>
25
<u>Total word count (4978)</u>

Abstract (214)
In facultative symbioses, only a fraction of hosts are associated with a symbiont. Understanding
why specific host and symbiont strains are associated can inform us of the evolutionary forces
affecting facultative symbioses. Possibilities include ongoing host-symbiont coevolution driven
by reciprocal selection, or priority effects that are neutral in respect to the host-symbiont
interaction. We hypothesized that ongoing host-symbiont coevolution would lead to higher
fitness estimates for naturally co-occurring (native) host and symbiont combinations compared
to nonnative combinations. We used the *Dictyostelium discoideum* – *Paraburkholderia bonniea*
system to test this hypothesis. *P. bonniea* features a reduced genome size relative to another
Paraburkholderia symbiont of *D. discoideum*, indicating a significant history of coevolution with
its host. Facultative symbionts may experience continued genome reduction if coevolution is
ongoing, or their genome size may have reached a stable state if the symbiosis has also
stabilized. Our work demonstrates that ongoing coevolution is unlikely for *D. discoideum* and *P.*
bonniea. The system instead represents a stable facultative symbiosis. Specifically associated
host and symbiont strains in this system are the result of priority effects, and presently
unassociated hosts are simply uncolonized. We find evidence for a virulence-transmission
trade-off without host strain specificity, and identify candidate virulence factors in the genomes
of *P. bonniea* strains that may contribute to variation in benevolence.
45

Lay summary
Symbiotic relationships between hosts and their microbial partners are prolonged and intimate
associations. Some of these relationships are obligatory for both a host and symbiont to survive,
while others are facultative and each partner can survive without the other. In the latter case,
some host individuals may be associated with a symbiont while others are not. Specific host and
symbiont combinations can be the result of reciprocal adaptation between host and symbiont
partners so that naturally co-occurring combinations are best suited for each other in terms of
their biological fitness. On the other hand, the symbiont that a host is associated with may
simply be the symbiont that arrived first, in what is called a priority effect. We sought to
determine which possibility best explained naturally co-occurring combinations of host and
symbiont strains of the social amoeba *Dictyostelium discoideum* and its symbiont
Paraburkholderia bonniea. Our work demonstrates that *D. discoideum* and *P. bonniea* are in a
stable facultative relationship. Specifically associated host and symbiont combinations are the
result of priority effects, and *D. discoideum* hosts without symbionts are simply uncolonized.
This work fills a gap in our understanding of the evolutionary forces affecting facultative
symbiotic relationships. We also show for the first time that *P. bonniea* symbionts can spread
among amoeba hosts when they aggregate together during the social stage of their life cycle.
63
Introduction
Host-symbiont associations range from obligate or facultative depending on the degree of host-
symbiont dependency (Sachs et al. 2011; Fisher et al. 2017). For obligate symbioses, when
host and symbiont need each other to survive, we expect significant coadaptation between host
and symbiont over a longer period of coevolution (Law and Dieckmann 1998). In contrast, for
facultative symbioses in which host and symbiont can each survive in a free-living state, we
expect coadaptation to a lesser degree or relationships to be more recent (Lo et al. 2016).
Significant coadaptation in obligate symbioses, especially those that feature strict vertical
transmission of symbionts, often result in symbionts with highly reduced genome sizes (Moran
et al. 2008; McCutcheon and Moran 2012). In facultative symbioses, many of which include
horizontal modes of symbiont transmission, symbiont genomes tend to be intermediate in size
(Toft and Andersson 2010; Sachs et al. 2011; Lo et al. 2016; Fisher et al. 2017). While ongoing
coevolution can lead to continued genome reduction for facultative symbionts, genome size may

have reached a stable state if the facultative symbiosis has also stabilized. For example,
Glomeribacter and Glomeromycota fungi are an example of an ancient facultative symbiosis
supported by phylogenetic codivergence patterns (Mondo et al. 2012).

Many aspects of facultative symbioses are relatively poorly understood, including which
factors determine distribution patterns of facultative symbionts among individuals of a host
species (Niepoth et al. 2018). In many facultative symbioses, only a fraction of hosts are
associated with a symbiont. Understanding why specific individual host and symbiont strains are
associated with each other can inform us of the forces contributing to the evolution of facultative
symbioses. Possibilities include ongoing host-symbiont coevolution driven by reciprocal
selection, or priority effects that are neutral in respect to the host-symbiont interaction itself
(Ganesan et al. 2022). Mutualistic coevolution could lead to specific combinations of host-
symbiont strains that enhance each other's fitness relative to other combinations (Rafaluk-Mohr
et al. 2018). In contrast, priority effects in a symbiosis context have to do with the order of
symbiont colonization and are typically considered when multiple symbiont species or strains
coexist within the same environment as the host (Ganesan et al. 2022). If we extend priority
effects to unassociated individual hosts in a facultative symbiosis, presently unassociated
individuals may simply be uncolonized. In this sense, the question of what drives host-symbiont
association patterns within a facultative symbiosis is related to questions of how host-microbe
associations originate (Sieber et al. 2021) and how microbial communities are assembled
(Tucker and Fukami 2014).

The potential for, and outcome of, long-term association between a host and any
potential symbionts is determined by intrinsic factors such as host and symbiont genetics, as
well as various extrinsic environmental factors (Schmid-Hempel 2011). For this work, we
focused on the intrinsic factors of host defense and symbiont virulence and their potential
contributions to fitness consequences of host-symbiont interactions. Host resistance (the ability
to limit the extent of an infection) and tolerance (the ability to tolerate the physiological
consequences of an infection) are the two large categories of host defense components that
affect evolutionary outcomes for symbioses (Simms and Triplett 1994; Råberg et al. 2007; Ayres
and Schneider 2012). The equivalent symbiont virulence components are proliferation (the
ability to proliferate to a certain extent in an infection) and benevolence (the ability to cause
beneficial or harmful physiological consequences of an infection) (Wollein Waldetoft et al. 2020).
Given a specific combination of host and symbiont, variable host resistance and symbiont
proliferation will lead to variation in symbiont density, while variable host tolerance and symbiont
benevolence will lead to variation in host fitness consequences relative to symbiont density
(Figure 1). Differentiating components of host defense or symbiont virulence is important
because of their downstream evolutionary effects. For example, host resistance often limits the
spread of infections while host tolerance can do the opposite and cause infections to increase in
frequency in a population (Roy and Kirchner 2000; Råberg 2014). Targeting pathogen
benevolence rather than proliferation may be more likely to prevent counter adaptation such as
the evolution of antimicrobial resistance (Wollein Waldetoft et al. 2020).

We used the *Dictyostelium discoideum – Paraburkholderia bonniea* system to determine
whether ongoing host-symbiont coevolution affects which host-symbiont strain pairs naturally
co-occur (are "native") within a facultative symbiosis, and whether host defense and symbiont
virulence contribute. The amoeba *D. discoideum* is an established model for understanding
intracellular pathogen infections (Steinert and Heuner 2005). It forms a facultative symbiosis
with three species of *Paraburkholderia*: *P. agricolaris*, *P. bonniea*, and *P. hayleyella* (DiSalvo et
al. 2015; Brock et al. 2020). Roughly one quarter of soil-isolated *D. discoideum* strains carry one
or occasionally multiple species of these symbiotic *Paraburkholderia* (Haselkorn et al. 2019).
The three symbionts also provide an opportunity to contrast the effect of different evolutionary
histories of association. Comparative genomics suggest that *P. bonniea* and *P. hayleyella* have

evolved genomes that are half the size of other *Paraburkholderia* in the amoeba host
environment (Noh et al. 2022). Patterns of host-symbiont coadaptation were previously tested
for *P. agricolaris* and *P. hayleyella*, indicating that reciprocal adaptation is ongoing for native *P.*
hayleyella hosts and symbionts. Native *P. hayleyella* hosts had a fitness advantage over
nonnative hosts when cured and reinfected with *P. hayleyella*, while native *P. agricolaris* hosts
did not have a similar advantage for *P. agricolaris* infections (Shu et al. 2018). Inside *D.*
discoideum hosts, *P. hayleyella* grew faster than on its own while *P. agricolaris* did not (Garcia
et al. 2019).

We hypothesized that ongoing host-symbiont coevolution would lead to higher fitness
estimates for native host and symbiont combinations compared to nonnative combinations. We
predicted coadapted intrinsic host and symbiont factors would contribute to fitness differences in
host-symbiont associations. Using the fact that *D. discoideum* form social groups and symbiont
horizontal transmission should be facilitated by these social groups, we examined host and
symbiont fitness at the group level to better consider the evolutionary consequences of variation
in fitness (Alizon and Michalakis 2015). We expected to find evidence of ongoing coevolution for
D. discoideum – P. bonniea based on previous results from *P. hayleyella*, the sister species to
P. bonniea, that contrasted from *P. agricolaris*.

Methods

D. discoideum has single-cell and multi-cell stages in its life cycle. Vegetative single-cell
amoebas feed on bacteria until food density becomes low and starvation begins. Next, in their
social stage, starving amoebas aggregate into a multicellular form that ultimately forms fruiting
bodies. The amoebas that survive the social stage become spores contained within the fruiting
body sorus. We used the vegetative stage to expose *D. discoideum* amoebas to *P. bonniea*
symbionts, and used spore counts after an unmanipulated social stage as a measure of host
fitness. We then used a subsequent social stage to expose uninfected amoebas to pre-infected
amoebas, and estimated symbiont horizontal transmission after this manipulated social stage as
an important aspect of symbiont fitness.

Focal host and symbiont strains

We used three native host strains of *D. discoideum* (QS395, QS433, QS859), three
nonnative *D. discoideum* strains (QS4, QS17, QS18), and three *P. bonniea* symbiont strains
(bb395, bb433, bb859). All *D. discoideum* strains were previously isolated from Mountain Lake
Biological Station in Virginia, USA. The native host strains had been cured of their symbionts
using tetracycline and verified as symbiont-free using PCR. For each replicate, host (with food
bacteria *Klebsiella pneumoniae*) and symbiont strains were grown from freezer stock on SM/5
plates (2□g glucose, 2□g BactoPeptone (Oxoid), 2□g yeast extract (Oxoid), 0.2□g MgCl$_2$,
1.9□g KH$_2$PO$_4$, 1□g K$_2$HPO$_4$ and 15□g agar per liter). KK2 buffer (2.2 g KH2PO4 monobasic
and 0.7 g K2HPO4 dibasic per liter) was used throughout for handling bacteria, *D. discoideum*
spores and amoebas.

Host fitness

We estimated host fitness at a range of infection prevalence for each host-symbiont
combination (6 host strains x 3 symbiont strains). We estimate host fitness at the social group
level rather than at the individual amoeba host level. Instead of symbiont density (e.g. Figure 1)
we estimate symbiont infection prevalence within a group of amoebas, and instead of host
fitness as a direct measure of individual host mortality we estimate group productivity given a
degree of symbiont prevalence. *Experiment*: For each combination, *D. discoideum* spores from

freshly grown fruiting bodies were collected and deposited on SM/5 plates at a density of 2×10^5
spores per plate and RFP-labeled symbionts at multiplicities of infection (MOI) of 0 (control),
0.6, 3, and 15. We made triplicate sample plates per combination and left these plates to fruit at
21°C for 5-7 days. We then collected all fruiting bodies and estimated the number of spores per
sample by counting 50x diluted spores on a hemocytometer and multiplying these counts by the
total volume of spores. *Analysis*: Our models examine the relationship between host fitness and
infection prevalence (not MOI). We estimated infection prevalence (percent of RPF+ infected
spores in a sample of 100,000) by running a sample of spores from each plate on a BD
FACSCalibur flow cytometer (Becton, Dickinson and Company, Franklin Lakes, NJ). FCS files
were imported into FlowJo v10.8.1 for analysis. For host fitness, we took the percent of spores
from each host-symbiont combination relative to the uninfected control that was prepared at the
same time. For example, host strain QS4 was infected with bb395 at MOI of 0 (control), 0.6, 3,
and 15. Spore counts of each individual sample plate were divided by the average spore count
of the three uninfected control plates.
189
Symbiont transmission
We estimated symbiont horizontal transmission for each host-symbiont combination (6
host strains x 3 symbiont strains). We estimate symbiont transmission at the social group level
rather than at the individual amoeba host level using the social stage of the *D. discoideum* life
cycle. For symbiont transmission we estimate the rate of transmission that occurs within a social
group of interacting hosts given a previous degree of symbiont infection prevalence, rather than
transmission from one host to another. *Pre-infection*: *D. discoideum* spores from freshly grown
fruiting bodies were collected and deposited on SM/5 plates at a density of 2 x 10^5 spores per
plate with RFP-labeled symbionts at MOI of 0 (control), plus an additional 3-5 MOI ranging from
0.3 to 30. Once these plates had fruited and amoebas were pre-infected, we collected fresh
spores and deposited them on SM/5 plates at densities of 1 x 10^5 and 2 x 10^5 spores per
plate with only food bacteria. *Experiment*: When amoebas were at log-phase growth roughly 36
hours later, uninfected amoebas were dyed with CellTracker Green CMFDA (Invitrogen)
dissolved in DMSO. Pre-infected amoebas (RFP+) were carried along with the dyed amoebas,
only exposed to DMSO, and treated in the same way through room temperature incubation and
washes. Afterward, dyed amoebas were combined with infected amoebas at 1:0 (dyed only
negative control), 0:1 (infected only), and 1:1 (mixed) ratios and placed on nitrocellulose filters
to aggregate and develop into fruiting bodies in triplicate. 5-7 days later, we collected all fruiting
bodies and their spores from these filters. *Analysis*: Our models examine the relationship
between symbiont transmission and infection prevalence (not MOI). We estimated infection
prevalence from the infected only sample and horizontal transmission from the mixed sample as
indicated by the co-occurrence of dye and infection in a spore. As above, we ran spore samples
on a BD FACSCalibur flow cytometer, and imported FCS files into FlowJo v10.8.1 for analysis.
We applied gates for spores, then dyed cells from the negative control, and infected cells from
one of the highest MOI infected only samples. A logical (AND) gate was applied to the mixed
sample to find the percent of spores that were previously uninfected dyed amoebas that were
now positive for symbiont infection (Figure 2).
217
Statistical analysis
Statistical analyses were performed in R v3.6.0 and with packages car v.3.0-12 (Fox and
Weisberg 2019) and lme4 v.1.1-28 (Bates et al. 2015). For both host fitness and symbiont
transmission, we fit linear models with mixed effects. For host fitness, we tested how percent
spores (relative percent of spores produced) were affected by symbiont strain and host type
(native or nonnative). We tested the effect of host type by coding the native combinations either

strictly (e.g. only QS859 infected with bb859 would be considered native) or leniently (e.g. any
native host of *P. bonniea* infected with any *P. bonniea* strain is considered native). We used
infection prevalence as a continuous predictor and experiment date and host strain as random
effects. For symbiont transmission, we fit a similar model but with percent transmission (percent
newly infected spores that were previously uninfected amoebas) as the dependent variable. For
all models we fit the most complex model first, then removed nonsignificant terms and
compared the simpler model with the complex one using a chi-squared test. We examined
effect sizes of factors in the final model using the package effectsize v.0.7.0 (Ben-Shachar et al.
2020) and its epsilon_squared() function. We also compared infection prevalence across the
two experiments. We fit linear models with host strain as a random effect and the different
number of social stages (nonnumeric factor; one for host fitness, two for symbiont transmission)
and symbiont identity as fixed effects. We examined residuals of the final model for any
indications that we had violated model assumptions. We used the package emmeans v.1.7.2
(Searle et al. 1980) and its emtrends() function for post-hoc tests of significant differences
between pairwise slopes of infection prevalence by symbiont strain. We also used the
emmeans() function for post-hoc tests of significant differences between means of infection
prevalence by symbiont strain. *P*-values were adjusted using Tukey's method.
241
Genome analysis
We sequenced and assembled the genomes of bb395 and bb433 as follows. High
molecular weight DNA was extracted using Lucigen MasterPure^{TM} Complete DNA and RNA
purification kits (LGC, Teddington, UK). Extracted DNA was sent to University of Washington
PacBio Sequencing Services for PacBio HiFi sequencing and MiGS for Oxford Nanopore (ONT)
and illumina sequencing. Raw reads were cleaned using filtlong v0.2.1 (Wick 2023) and fastp
v0.23.1 (Chen et al. 2018), then assembled and polished using Trycycler v0.5.3 (Wick et al.
2021) and Polypolish v0.5.0 (Wick and Holt 2022). Trycycler input files were created using Flye
v2.9.1-b1780 (Kolmogorov et al. 2019) for both types of reads, Hifiasm v0.18.5-r499 (Cheng et
al. 2022) for PacBio reads, and Raven v1.8.1 (Vaser and Šikić 2021) for ONT reads. Assembled
contigs were re-oriented with Circulator v1.5.5 (Hunt et al. 2015). Genes were predicted using
Prokka v1.14.6 (Seemann 2014). Pseudofinder v1.0 (Syberg-Olsen et al. 2022) was used to
remove pseudogenes. Pan-genome analysis of the three *P. bonniea* genomes was performed
using Roary v3.13.0 (Page et al. 2015). For examination of the differences between genomes,
whole genomes were aligned using progressiveMauve (Darling et al. 2010) within Geneious
Prime v2020.2.5 (https://www.geneious.com).
258
259
260 <u>Results</u>
261
Host fitness
We estimated the relationship between host fitness and symbiont infection prevalence
by exposing native and nonnative *D. discoideum* hosts to three strains of *P. bonniea* at a range
of multiplicities of infection. Native hosts were *D. discoideum* strains harboring their own *P.*
bonniea symbiont strain when they were isolated from the wild. Host fitness decreased with
increasing infection prevalence across all combinations of hosts and symbionts, but there was
no difference between the native and nonnative host-symbiont combinations (Figure 3; Table 1).
This was regardless of whether native status was defined using a strict or lenient definition
(Supplementary material Table S1). Instead, symbiont strain identity significantly affected how
host fitness responded to infection prevalence. For example, bb859 infections resulted in a
significantly shallower slope in host fitness decline (β= -0.118, SE= 0.0172) compared to bb395

(β= -0.180, SE= 0.0158) and bb433 infections (β= -0.231, SE= 0.0157) (Figure 3). The effects of
bb395 and bb433 infections on host fitness were not significantly different from each other.
These results support a significant effect of symbiont benevolence variation on host fitness
during initial infection. Based on our results, bb859 is more benevolent while bb395 and bb433
are more malevolent to all hosts.
278
Symbiont transmission

We estimated the relationship between symbiont transmission and symbiont infection
prevalence by exposing uninfected amoebas to pre-infected amoebas during a manipulated
social stage of the host life cycle. Horizontal transmission was detected when previously
uninfected spores harbored *P. bonniea* at the end of this social stage. The degree of symbiont
horizontal transmission increased with the infection prevalence of pre-infected amoebas that
entered the social stage (Figure 4). This positive correlation confirms that the increase in
infection is due to horizontal transmission rather than vertical transmission (Ebert 2013). This is
consistent with previous reports that very few amoebas go through cell division during the social
stage (Muramoto and Chubb 2008). As with host fitness above, native and nonnative host-
symbiont combinations were no different in overall patterns of symbiont transmission (Table 2),
regardless of how native status was defined (Supplementary material Table S2).

In this experiment, infected bb859 reached higher infection prevalence and showed a
higher rate of horizontal transmission compared to bb395 and bb433 (Figure 4). Pre-infection
was established in one social stage, and the experiment was performed in a subsequent social
stage. The three *P. bonniea* strains used in this experiment do not significantly differ in their per
amoeba-spore bacterial density (Miller et al. 2020). Therefore, higher infection prevalence in
bb859 compared to the other strains does not necessarily indicate that bb859 is better at
proliferation. However, bb859 symbionts transmitted at a significantly higher rate (β= 0.525, SE=
0.0244) beyond that simply due to bb859 infections occurring at higher infection prevalence
compared to the other two strains bb395 (β= 0.361, SE= 0.0343) and bb433 (β= 0.249, SE=
0.0289) (Figure 4). When native hosts were leniently defined but not when strictly defined,
infection prevalence and host type interacted to reveal that nonnative hosts became infected by
bb859 at a slightly higher density compared to native hosts and therefore also transmitted
symbionts at a higher level.
304
Synthesis of host and symbiont fitness

We observed a significant difference in infection prevalence between the two
experiments that is likely due to differences in symbiont strain-specific virulence (Table 3; Figure
5). The main difference between the design of the two experiments is that for *Host fitness* we
used newly-infected amoebas, but for *Symbiont transmission* we used amoebas that had been
infected in a previous vegetative stage, to minimize the potential for symbionts to be newly
acquired from the environment and to maximize our ability to detect horizontal transmission
among amoebas. In addition, we infected amoebas for the symbiont transmission experiment at
a wider range of MOI (0.3-30) compared to the host fitness experiment (0.6-15) in order to
achieve as wide a range of infection prevalence as possible. We found that we were unable to
push the maximum of these ranges higher by using a higher MOI for the initial infection.

Combined observations from both experiments suggest that variation in symbiont
benevolence leads to variation in symbiont prevalence among *P. bonniea* strains infecting
amoeba hosts in the longer term. We observed no differences in infection prevalence after the
first vegetative stage – social stage cycle (Figure 5(a)). But more malevolent strains bb395 (μ=
36.5, SE= 2.38) and bb433 (μ = 33.8, SE= 2.29) become significantly less prevalent among
amoeba hosts compared to more benevolent strain bb859 (μ= 53.5, SE= 2.17) over the course

of two vegetative stage - social stage cycles (Figure 5(b)). A higher rate of horizontal
transmission in benevolent strains compared to malevolent strains during social stages
contributes to this result (Figure 4).

Candidate virulence factors

The three *P. bonniea* strains used in this experiment do not significantly differ in their per
spore bacterial density (Miller et al. 2020). Yet the fact that we observe virulence variation in
benevolence suggests that bb395 and bb433 possess malevolent virulence factors that bb859
lacks, or bb859 possesses one or more benevolent virulence factors that the other strains do
not. Whole genome alignment revealed two structural variants each approximately 13 kilobases
in size present in strains bb395 (1:108428 – 122213 and 1:2654181 – 2667432) and bb433
(1:108428 – 122213 and 1:2658228 – 2671479) and absent in bb859. These variants likely
resulted from integration of plasmids as both regions have genes encoding putative plasmid
recombinases near the 3' end. We also identified an approximately 6 kilobase structural variant
in strain bb859 (2:708878 – 715517) that lacks closely-related homologs in the other sequenced
genomes of *Paraburkholderia* symbionts of *D. discoideum* (Brock et al. 2020)*.

<u>Discussion</u>

The reduced-genome symbiont *P. bonniea* is able to persistently infect *D. discoideum*
and form a facultative symbiosis in which only a fraction of hosts are associated with this
symbiont (DiSalvo et al. 2015; Haselkorn et al. 2019; Miller et al. 2020). Based on our results, *D.
discoideum* and *P. bonniea* are in a stable facultative symbiosis, where host-symbiont
coevolution is primarily historical rather than ongoing. Host and symbiont strains responded
similarly to association in native and nonnative combinations. We observed variation in
virulence among *P. bonniea* strains but its effect did not vary for native and nonnative *D.
discoideum* hosts.

Strain level variation in symbiont virulence – virulence-transmission trade-off

Horizontal transmission of *Paraburkholderia* symbionts was previously assumed, but we
demonstrate it for the first time in this study. In addition, we found evidence that strain level
variation in symbiont virulence affects transmission by enabling higher rates of horizontal
transmission in benevolent strains compared to malevolent strains. The virulence-transmission
trade-off hypothesis posits that if there is a biological link between symbiont virulence and
transmission (e.g. because symbionts use host resources to replicate), symbionts should evolve
toward a level of virulence where transmission is greatest (Anderson and May 1982). This
hypothesized trade-off has proven difficult to observe for several reasons, including the
influences of host range, multiple symbiont infections, and host population structure that can
affect host-symbiont interactions in complex ways (Alizon et al. 2009; Leggett et al. 2013).
However, recent meta-analyses indicate that there is partial support for the trade-off hypothesis
in several biological systems (Acevedo et al. 2019). The three core predictions of the trade-off
hypothesis are as follows: (i) within-host symbiont replication rates have a positive relationship
with symbiont virulence; (ii) within-host symbiont replication rates have a positive relationship
with symbiont transmission rates; and (iii) symbiont virulence and transmission have a trade-off
through the negative relationship between symbiont virulence and host recovery rates.

Because we estimated host fitness and symbiont transmission at the social group level
rather than at the individual amoeba host level, our evidence that supports each of these points
may be an indirect test of the virulence-transmission trade-off, as the original theory is

formulated based on investigating symbiont density in individual hosts. However, considering
fitness at the population level can facilitate our understanding of how host defense and symbiont
virulence evolve (Alizon and Michalakis 2015). We find support for each of the core predictions
of the virulence-transmission trade-off hypothesis. Symbiont infection prevalence had a negative
relationship with host fitness (i; Figure 3), and symbiont infection prevalence had a positive
relationship with symbiont horizontal transmission (ii; Figure 4). Lastly, variation in symbiont
benevolence led to increased horizontal transmission of benevolent strains compared to
malevolent strains (iii; Figure 4).

Transmission dynamics are an important missing piece of information in further
understanding the evolution of the facultative symbiosis between *D. discoideum* and
Paraburkholderia. Given the persistence of *Paraburkholderia* symbiont infections in the lab and
the facultative aspect of the symbiosis itself, we expect these symbionts to transmit among
hosts using both vertical and horizontal routes. We did not test for vertical transmission and
specifically designed our experiment to use the social stage so that only horizontal transmission
was possible. Our results clearly show the presence of significant horizontal symbiont
transmission among *D. discoideum* amoebas in the social stage of their life cycle. The
coexistence and relative influences of both types of transmission routes are better understood in
other amoeba-bacteria symbioses (Herrera et al. 2020).

Lack of variation in host defense – ecology and geography

Our results indicate that native hosts of *P. bonniea* do not possess enhanced host
defenses to *P. bonniea* compared to nonnative hosts. These results contrast with evidence of
host-symbiont coadaptation in *P. hayleyella* and its native hosts (Shu et al. 2018; Garcia et al.
2019). Both *P. hayleyella* and *P. bonniea* have reduced genomes and are sister species to each
other (Brock et al. 2020). The lack of host strain-specific counteradaptation to *P. bonniea*
virulence variation may be related to the ecology of host and symbiont. *P. bonniea* is the rarest
of *Paraburkholderia* symbionts of *D. discoideum* (Haselkorn et al. 2019; DuBose et al. 2022).
Rare encounters between new hosts and *P. bonniea* may not have a significant impact on *D.*
discoideum populations, particularly if *D. discoideum* itself is sparsely distributed across soil
landscapes. The fitness cost of infection for rare *P. bonniea* may be insufficient for hosts to
evolve defenses against it (Anderson and May 1982).

Another potential reason why we did not find evidence of host defense variation may be
because our native and nonnative hosts were collected from the same geographical locality. In
other words, coevolution in this system may occur at the population level rather than at the
strain level. Examples of strain-level coadaptation in facultative symbionts are limited. In pea
aphids, coadaptation with their symbionts goes in both directions. Pea aphids from a population
(*Lotus pedunculatus* biotype) that typically carry *Hamiltonella defensa* were more likely to
establish symbiosis when newly infected with *Hamiltonella* compared to hosts from another
population (*Lotus corniculatus* biotype) (Niepoth et al. 2018). However, in a tripartite host-
symbiont-pathogen system, host-by-symbiont and symbiont-by-pathogen genotype by genotype
interactions affected how protective *Regiella* symbionts were to pea aphids against the fungal
pathogen *Pandora*, but not in an way that supported coadaptation between native host and
symbiont genotype combinations (Parker et al. 2017). *Xenorhabdus* spp. are obligate symbionts
to various *Steinernema* spp. nematodes. Evidence of host-symbiont coadaptation has been
found in a couple different pairings, including between strains of *Xenorhabdus nematophila* and
Steinernema carpocapsae (Chapuis et al. 2009), and between *Xenorhabdus bovienii* strains
and their various *Steinernema* species hosts (Murfin et al. 2015).

Alternatively, the difference in results between *P. bonniea* and *P. hayleyella* may be due
to previous experimental approaches that do not account for host or symbiont fitness as a

function of infection prevalence. We plan to apply the experimental approach of the current
study to the relationship between *D. discoideum* and *P. hayleyella*, which is known to cause
more detrimental fitness consequences to nonnative hosts (Shu et al. 2018; Miller et al. 2020). If
previous results hold for *P. hayleyella,* it would provide support for the evolution of reduced
antagonism in *P. bonniea*. Amoeba hosts experience only a mild reduction in fitness to *P.*
bonniea infection compared to *P. agricolaris* or *P. hayleyella* (Miller et al. 2020). Reduced
antagonism is a potential outcome of symbiosis that is favored when virulence-transmission
tradeoffs are present and new hosts are rare (Yamamura 1993; Johnson et al. 2021). Both of
these conditions appear to hold for *P. bonniea* and support our interpretation that the *D.*
discoideum – P. bonniea relationship is a stable facultative symbiosis.

Candidate virulence factors for benevolence variation

We identified two structural variants that are present in the genomes of bb395 and
bb433 but absent in bb859, as well as an additional structural variant that is present in the
genome of bb859 but absent in the other two strains. We examined these regions for potential
virulence factors that might contribute to the variation in benevolence we observed among *P.*
bonniea strains.

Several candidate genes may confer increased malevolence to bb395 and bb433.
Among these, PB395_00119/ PB433_00119 encode a putative member of the peptidase S8
family (also called subtilisin-related peptidases) that are known contributors to the pathogenesis
of *Streptococcus pneumoniae* (Ali et al. 2021). Interestingly, there are no closely-related
homologs in other sequenced *Paraburkholderia* genomes, but proteins with the highest
similarity in Genbank are found in *Burkholderia pseudomallei* (MBF3536330.1, 93% identity
over the length of the protein) and *Ralstonia solanacearum* (NKA33280.1, 93% identity over the
length of the protein), which are respectively pathogens of mammals and plants. Potentially, this
peptidase may be introduced into *D. discoideum* cells and modify host responses to infection.

In the other putative plasmid integration site, PB395_02359/ PB433_02360 is a member
of the Xenobiotic Response Element (XRE) family of transcriptional regulators. It is most similar
to a homolog in the opportunistic human pathogen *Burkholderia cepacia* complex
(WP_060080935.1, 80% identity over the length of the protein). In these proteins, the putative
DNA-binding domain is at the N terminus of the protein and the C-terminal portion of the protein
contains a predicted peptidase domain. This architecture is similar to the AlpR gene in
Pseudomonas aeruginosa which is involved in the pathway of programmed cell death and has
been demonstrated as a virulence factor (McFarland et al. 2015).

One gene in particular appears to be an interesting candidate for a benevolence factor
for strain bb859. PBONN_03430 encodes a putative member of a Type I secretion system
known to secrete a wide variety of proteins from Gram negative bacteria (Spitz et al. 2019). An
interesting possibility is that the PBONN_03403 protein may be part of a complex that secretes
proteins that modulate activities in *D. discoideum* cells to prevent host cell death and allow
bb859 to transmit to other host cells. We are actively working to assess the role of these genes
in *P. bonniea* virulence.

Conclusions

Our work demonstrates that ongoing coevolution is unlikely for *D. discoideum* and *P.*
bonniea. The system instead represents a stable facultative symbiosis. Specifically associated
host and symbiont strains in this system are the result of priority effects rather than ongoing
host-symbiont coevolution, and presently unassociated hosts are simply uncolonized. In
addition, we found evidence for a virulence-transmission trade-off without host strain specificity.
A benevolent strain had a higher horizontal transmission rate compared to malevolent strains

during the social stage of the host life cycle, regardless of host strain. Lastly, we identified
candidate virulence factors in *P. bonniea* genomes, paving the way for future research in
determinants of strain level variation in benevolence.
471
472 <u>Data availability</u>
473   Full genome sequences for *P. bonniea* strains are available through NCBI Genomes
474 (txid2152891). Strain bb859 (SAMN09651436) was submitted previously and bb395 and bb433
475 were newly submitted (SAMN35686367, SAMN35686368). Additional data and code supporting
476 this work can be accessed at: https://github.com/noh-lab/social-infection
477

<u>Acknowledgements</u>
We thank Susanne DiSalvo, who generously shared the RFP-labeled *P. bonniea* strains,
and the Strassmann-Queller lab who generously shared the *D. discoideum* strains used in this
work. We thank past and current members of the Noh lab (Caroline Lunt, Gus Shuster) who
contributed moral support for this project during its development. Lily Khadempour, Laura
Runyen-Janecky, Chris Moore, and Judy Stone provided helpful comments that improved
previous drafts.
This work was supported by the National Institutes of Health and its National Institute of
General Medical Sciences by an Institutional Development Award (IDeA) under Grant Number
P20GM103423 (subaward to SN) and Colby College startup funds.

References
Acevedo, M. A., F. P. Dillemuth, A. J. Flick, M. J. Faldyn, and B. D. Elderd. 2019. Virulence-
driven trade-offs in disease transmission: A meta-analysis*. Evolution 73:636–647.
Ali, M. Q., T. P. Kohler, L. Schulig, G. Burchhardt, and S. Hammerschmidt. 2021.
Pneumococcal extracellular serine proteases: molecular analysis and impact on
colonization and disease. Frontiers in Cellular and Infection Microbiology 11.
Alizon, S., A. Hurford, N. Mideo, and M. Van Baalen. 2009. Virulence evolution and the trade-off
hypothesis: history, current state of affairs and the future. Journal of Evolutionary Biology
22:245–259.
Alizon, S., and Y. Michalakis. 2015. Adaptive virulence evolution: the good old fitness-based
approach. Trends in Ecology & Evolution 30:248–254.
Anderson, R. M., and R. M. May. 1982. Coevolution of hosts and parasites. Parasitology
85:411–426. Cambridge University Press.
Ayres, J. S., and D. S. Schneider. 2012. Tolerance of infections. Annual Review of Immunology
30:271–294.
Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using
lme4. Journal of Statistical Software 67:1–48.
Ben-Shachar, M. S., D. Lüdecke, and D. Makowski. 2020. effectsize: Estimation of effect size
indices and standardized parameters. Journal of Open Source Software 5:2815.
Brock, D. A., S. Noh, A. N. M. Hubert, T. S. Haselkorn, S. DiSalvo, M. K. Suess, A. S. Bradley,
M. Tavakoli-Nezhad, K. S. Geist, D. C. Queller, and J. E. Strassmann. 2020.
Endosymbiotic adaptations in three new bacterial species associated with *Dictyostelium*
discoideum: *Paraburkholderia agricolaris* sp. nov., *Paraburkholderia hayleyella* sp. nov.,
and *Paraburkholderia bonniea* sp. nov. PeerJ 8:e9151.
Chapuis, É., V. Emelianoff, V. Paulmier, N. Le Brun, S. Pagès, M. Sicard, and J.-B. Ferdy.
2009. Manifold aspects of specificity in a nematode–bacterium mutualism. Journal of
Evolutionary Biology 22:2104–2117.
Chen, S., Y. Zhou, Y. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics 34:i884–i890.

Cheng, H., E. D. Jarvis, O. Fedrigo, K.-P. Koepfli, L. Urban, N. J. Gemmell, and H. Li. 2022. Haplotype-resolved assembly of diploid genomes without parental data. Nat Biotechnol 40:1332–1335. Nature Publishing Group.

Darling, A. E., B. Mau, and N. T. Perna. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. PLOS ONE 5:e11147. Public Library of Science.

DiSalvo, S., T. S. Haselkorn, U. Bashir, D. Jimenez, D. A. Brock, D. C. Queller, and J. E. Strassmann. 2015. *Burkholderia* bacteria infectiously induce the proto-farming symbiosis of *Dictyostelium* amoebae and food bacteria. PNAS 112:E5029–E5037.

DuBose, J. G., M. S. Robeson, M. Hoogshagen, H. Olsen, and T. S. Haselkorn. 2022. Complexities of inferring symbiont function: *Paraburkholderia* symbiont dynamics in social amoeba populations and their impacts on the amoeba microbiota. Applied and Environmental Microbiology 0:e01285-22. American Society for Microbiology.

Ebert, D. 2013. The epidemiology and evolution of symbionts with mixed-mode transmission. Annu. Rev. Ecol. Evol. Syst. 44:623–643.

Fisher, R. M., L. M. Henry, C. K. Cornwallis, E. T. Kiers, and S. A. West. 2017. The evolution of host-symbiont dependence. Nat Commun 8:15973. Nature Publishing Group.

Fox, J., and S. Weisberg. 2019. An R companion to applied regression. 3rd ed. Sage, Thousand Oaks, CA.

Ganesan, R., J. C. Wierz, M. Kaltenpoth, and L. V. Flórez. 2022. How it all begins: bacterial factors mediating the colonization of invertebrate hosts by beneficial symbionts. Microbiology and Molecular Biology Reviews 86:e00126-21. American Society for Microbiology.

Garcia, J. R., T. J. Larsen, D. C. Queller, and J. E. Strassmann. 2019. Fitness costs and benefits vary for two facultative *Burkholderia* symbionts of the social amoeba, *Dictyostelium discoideum*. Ecology and Evolution 9:9878–9890.

Haselkorn, T. S., S. DiSalvo, J. W. Miller, U. Bashir, D. A. Brock, D. C. Queller, and J. E. Strassmann. 2019. The specificity of *Burkholderia* symbionts in the social amoeba farming symbiosis: prevalence, species, genetic and phenotypic diversity. Molecular Ecology 28:847–862.

Herrera, P., L. Schuster, C. Wentrup, L. König, T. Kempinger, H. Na, J. Schwarz, S. Köstlbacher, F. Wascher, M. Zojer, T. Rattei, and M. Horn. 2020. Molecular causes of an evolutionary shift along the parasitism–mutualism continuum in a bacterial symbiont. Proceedings of the National Academy of Sciences 117:21658–21666. Proceedings of the National Academy of Sciences.

Hunt, M., N. D. Silva, T. D. Otto, J. Parkhill, J. A. Keane, and S. R. Harris. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. Genome Biology 16:294.

Johnson, C. A., G. P. Smith, K. Yule, G. Davidowitz, J. L. Bronstein, and R. Ferrière. 2021. Coevolutionary transitions from antagonism to mutualism explained by the Co-Opted Antagonist Hypothesis. Nat Commun 12:2867. Nature Publishing Group.

Kolmogorov, M., J. Yuan, Y. Lin, and P. A. Pevzner. 2019. Assembly of long, error-prone reads using repeat graphs. Nat Biotechnol 37:540–546. Nature Publishing Group.

Law, R., and U. Dieckmann. 1998. Symbiosis through exploitation and the merger of lineages in evolution. Proceedings of the Royal Society of London. Series B: Biological Sciences 265:1245–1253. Royal Society.

Leggett, H. C., A. Buckling, G. H. Long, and M. Boots. 2013. Generalism and the evolution of parasite virulence. Trends in Ecology & Evolution 28:592–596.

Lo, W.-S., Y.-Y. Huang, and C.-H. Kuo. 2016. Winding paths to simplicity: genome evolution in facultative insect symbionts. FEMS Microbiology Reviews 40:855–874.

McCutcheon, J. P., and N. A. Moran. 2012. Extreme genome reduction in symbiotic bacteria.
Nat Rev Microbiol 10:13–26. Nature Publishing Group.
McFarland, K. A., E. L. Dolben, M. LeRoux, T. K. Kambara, K. M. Ramsey, R. L. Kirkpatrick, J.
D. Mougous, D. A. Hogan, and S. L. Dove. 2015. A self-lysis pathway that enhances the
virulence of a pathogenic bacterium. Proceedings of the National Academy of Sciences
112:8433–8438. Proceedings of the National Academy of Sciences.
Miller, J. W., C. R. Bocke, A. R. Tresslar, E. M. Schniepp, and S. DiSalvo. 2020.
Paraburkholderia symbionts display variable infection patterns that are not predictive of
amoeba host outcomes. Genes 11:674. Multidisciplinary Digital Publishing Institute.
Mondo, S. J., K. H. Toomer, J. B. Morton, Y. Lekberg, and T. E. Pawlowska. 2012. Evolutionary
stability in a 400-million-year-old heritable facultative mutualism. Evolution 66:2564–
2576.
Moran, N. A., J. P. McCutcheon, and A. Nakabachi. 2008. Genomics and evolution of heritable
bacterial symbionts. Annual Review of Genetics 42:165–190.
Muramoto, T., and J. R. Chubb. 2008. Live imaging of the *Dictyostelium* cell cycle reveals
widespread S phase during development, a G2 bias in spore differentiation and a
premitotic checkpoint. Development 135:1647–1657.
Murfin, K. E., M.-M. Lee, J. L. Klassen, B. R. McDonald, B. Larget, S. Forst, S. P. Stock, C. R.
Currie, and H. Goodrich-Blair. 2015. *Xenorhabdus bovienii* strain diversity impacts
coevolution and symbiotic maintenance with *Steinernema* spp. nematode hosts. mBio
6:e00076-15. American Society for Microbiology.
Niepoth, N., J. Ellers, and L. M. Henry. 2018. Symbiont interactions with non-native hosts limit
the formation of new symbioses. BMC Evolutionary Biology 18:27.
Noh, S., B. J. Capodanno, S. Xu, M. C. Hamilton, J. E. Strassmann, and D. C. Queller. 2022.
Reduced and nonreduced genomes in *Paraburkholderia* symbionts of social amoebas.
mSystems 0:e00562-22. American Society for Microbiology.
Page, A. J., C. A. Cummins, M. Hunt, V. K. Wong, S. Reuter, M. T. G. Holden, M. Fookes, D.
Falush, J. A. Keane, and J. Parkhill. 2015. Roary: rapid large-scale prokaryote pan
genome analysis. Bioinformatics 31:3691–3693.
Parker, B. J., J. Hrček, A. H. C. McLean, and H. C. J. Godfray. 2017. Genotype specificity
among hosts, pathogens, and beneficial microbes influences the strength of symbiont-
mediated protection. Evolution 71:1222–1231.
Råberg, L. 2014. How to live with the enemy: understanding tolerance to parasites. PLoS Biol
12.
Råberg, L., D. Sim, and A. F. Read. 2007. Disentangling genetic variation for resistance and
tolerance to infectious diseases in animals. Science 318:812–814.
Rafaluk-Mohr, C., B. Ashby, D. A. Dahan, and K. C. King. 2018. Mutual fitness benefits arise
during coevolution in a nematode-defensive microbe model. Evolution Letters 2:246–
256.
Roy, B. A., and J. W. Kirchner. 2000. Evolutionary dynamics of pathogen resistance and
tolerance. Evolution 54:51–63.
Sachs, J. L., C. J. Essenberg, and M. M. Turcotte. 2011. New paradigms for the evolution of
beneficial infections. Trends in Ecology & Evolution 26:202–209.
Schmid-Hempel, P. 2011. Evolutionary Parasitology. Oxford University Press, New York, New
York.
Searle, S. R., F. M. Speed, and G. A. Milliken. 1980. Population marginal means in the linear
model: an alternative to least squares means. The American Statistician 34:216–221.
Taylor & Francis.
Seemann, T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069.

Shu, L., D. A. Brock, K. S. Geist, J. W. Miller, D. C. Queller, J. E. Strassmann, and S. DiSalvo.
2018. Symbiont location, host fitness, and possible coadaptation in a symbiosis between
social amoebae and bacteria. eLife 7:e42660.
Sieber, M., A. Traulsen, H. Schulenburg, and A. E. Douglas. 2021. On the evolutionary origins
of host–microbe associations. Proceedings of the National Academy of Sciences
118:e2016487118. Proceedings of the National Academy of Sciences.
Simms, E. L., and J. Triplett. 1994. Costs and benefits of plant responses to disease: resistance
and tolerance. Evolution 48:1973–1985.
Spitz, O., I. N. Erenburg, T. Beer, K. Kanonenberg, I. B. Holland, and L. Schmitt. 2019. Type I
secretion systems — one mechanism for all? Microbiology Spectrum
7:10.1128/microbiolspec.psib-0003–2018. American Society for Microbiology.
Steinert, M., and K. Heuner. 2005. *Dictyostelium* as host model for pathogenesis. Cell.
Microbiol. 7:307–314.
Syberg-Olsen, M. J., A. I. Garber, P. J. Keeling, J. P. McCutcheon, and F. Husnik. 2022.
Pseudofinder: detection of pseudogenes in prokaryotic genomes. Molecular Biology and
Evolution msac153.
Toft, C., and S. G. E. Andersson. 2010. Evolutionary microbial genomics: insights into bacterial
host adaptation. Nature Reviews Genetics 11:465–475.
Tucker, C. M., and T. Fukami. 2014. Environmental variability counteracts priority effects to
facilitate species coexistence: evidence from nectar microbes. Proceedings of the Royal
Society B: Biological Sciences 281:20132637. Royal Society.
Vaser, R., and M. Šikić. 2021. Time- and memory-efficient genome assembly with Raven. Nat
Comput Sci 1:332–336. Nature Publishing Group.
Wick, R. R. 2023. https://github.com/rrwick/Filtlong.
Wick, R. R., and K. E. Holt. 2022. Polypolish: Short-read polishing of long-read bacterial
genome assemblies. PLOS Computational Biology 18:e1009802. Public Library of
Science.
Wick, R. R., L. M. Judd, L. T. Cerdeira, J. Hawkey, G. Méric, B. Vezina, K. L. Wyres, and K. E.
Holt. 2021. Trycycler: consensus long-read assemblies for bacterial genomes. Genome
Biology 22:266.
Wollein Waldetoft, K., L. Råberg, and R. Lood. 2020. Proliferation and benevolence—A
framework for dissecting the mechanisms of microbial virulence and health promotion.
Evolutionary Applications 13:879–888.
Yamamura, N. 1993. Vertical transmission and evolution of mutualism from parasitism.
Theoretical Population Biology 44:95–109.
652
653

654   Tables

656   Table 1. Analysis of Deviance table for Host fitness with lenient native definition

Variable	X^2	df	P-value	η^2_p	95% CI
Infection prevalence	376.002	1	< 0.001	0.63	(0.56, 1)
Symbiont strain	17.998	2	< 0.001	- 0.06	(0, 1)
Host type	< 0.001	1	0.981	- 0.37	(0, 1)
Infection prevalence : Symbiont strain	23.886	2	< 0.001	0.09	(0.04, 1)

660 Table 2. Analysis of Deviance table for Symbiont transmission with lenient native definition

Variable	X^2	df	P-value	η^2_p	95% CI
Infection prevalence	484.965	1	< 0.001	0.77	(0.71, 1)
Symbiont strain	149.430	2	< 0.001	0.43	(0.20, 1)
Host type	2.092	1	0.148	0.53	(0.25, 1)
Infection prevalence : Symbiont strain	109.289	2	< 0.001	0.65	(0.52, 1)
Infection prevalence : Host type	35.991	1	< 0.001	0.30	(0.17, 1)
Symbiont strain : Host type	9.509	2	0.009	0.14	(0.01, 1)

661
662
663

664   Table 3. Analysis of Deviance table for infection prevalence across experiments

Variable	X^2	df	P-value	η^2_p	95% CI
Number of social stages	71.196	1	< 0.001	0.16	(0.11, 1)
Symbiont strain	17.789	2	< 0.001	0.02	(0.00, 1)
Number of social stages : Symbiont strain	39.197	2	< 0.001	0.09	(0.05, 1)

665
666

<u>Figure legends</u>

Figure 1. Theoretical differences between resistance-based and tolerance-based host defense,
and proliferation-based and benevolence-based symbiont virulence. Both types of variation can
be detected by measuring host fitness and detecting when (a) different host genotypes vary in
responses to infection by the same symbiont, or (b) the same host genotype varies in responses
to infection by different symbiont genotypes. (a) Hosts may vary in **resistance** to the same
symbiont. More resistant hosts (a – black line; lower average symbiont density) will have overall
higher fitness compared to less resistant hosts (b – gray line; higher average symbiont density).
But the fitness cost per increase in symbiont density is similar for both hosts, as indicated by the
similar slopes. On the other hand, hosts may vary in **tolerance** to the same symbiont. More
tolerant hosts (a – black line; shallower slope) may have a similar average symbiont density to
less tolerant hosts (b – gray line; steeper slope), but will suffer a lower fitness cost per increase
in symbiont density. (b) The same host may suffer different fitness effects due to symbionts that
vary in **proliferation**. The better proliferating symbiont (b – gray line; higher average symbiont
density) will impart higher fitness costs to the host than the worse proliferating symbiont (a –
black line; lower average symbiont density). But the fitness cost to the host per increase in
symbiont density is similar for both symbionts. The same host may be infected with symbionts
that vary in **benevolence**. More benevolent symbionts (a – black line; shallower slope) will
impart a lower fitness cost per increase in symbiont density compared to more malevolent
symbionts (b – gray line; steeper slope), though both symbionts may reach similar average
densities. (Framework adapted from Raberg et al 2007 and Wollein Waldetoft *et al.*, 2020.)

Figure 2. Representative samples from symbiont transmission experiment to demonstrate how
symbiont transmission during a social stage is detected. (a) Dyed cells (green population) are
gated on uninfected controls treated with Celltracker. (b) Infected cells with RFP-labeled
symbionts (pink population) are gated on pre-infected controls. (c) Newly infected cells (blue
population) are detected by applying logical gates to the experimental samples that mixed
uninfected cells and pre-infected cells.

Figure 3. Host fitness was negatively correlated with infection prevalence and differed in slope
but not average infection prevalence among symbiont strains after a single social stage.
Infection prevalence was estimated per host-symbiont-MOI sample as the percent of infected
spores that had RFP labeled symbionts in them. Host fitness was estimated by the percent of
spores produced per host-symbiont-MOI sample, relative to uninfected hosts prepared at the
same time as the infected host-symbiont combinations. There was no significant difference
between native (top row; lenient definition) vs. nonnative (bottom row) host strains.

Figure 4. The rate of horizontal transmission was positively correlated with pre-infection
prevalence and also differed in slope and average infection prevalence among symbiont strains
after a second social stage. Infection prevalence was estimated per host-symbiont-MOI sample
as the percent of RPF+ infected spores of the pre-infected control. Horizontal transmission was
estimated by the percent of RFP+ infected spores in the test sample that were also positive for
a green membrane dye. These dyed cells were uninfected prior to the experiment, during which
they were combined with pre-infected (and undyed) cells. There was no significant difference in
symbiont transmission among native (top row; lenient definition) and nonnative (bottom row)
host strains.

Figure 5. Infection prevalence was significantly different when amoebas were observed after
one (a) vs. two social stages (b) after infection. More malevolent strains (bb395 and bb433)

reach lower infection prevalence among amoeba hosts compared to more benevolent strains
(bb859) over time.

(a) Host defense components

Resistance | Tolerance

Host fitness

Symbiont density

Host genotype
a
b

(b) Symbiont virulence components

Proliferation | Benevolence

Host fitness

Symbiont density

Symbiont genotype
a
b