1 *Paraburkholderia bonniea* symbiont strain identity impacts amoeba host fitness and rate of
2 horizontal transmission
3

4 Suegene Noh[1]*, Emily R. Larson[1], Rachel M. Covitz[2], Anna Chen[1], Prachee R. Mazumder[1],
5 Marisa C. Hamilton[3], Robert A. Dettmann[4]
6
[1] Biology Department, Colby College, Waterville, Maine, USA
[2] Department of Developmental, Molecular, and Chemical Biology, Tufts University School of
Medicine, Boston, Massachusetts, USA
[3] University Program in Genetics and Genomics, Duke University, Durham, North Carolina, USA
[4] Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA
* Corresponding author: suegene.noh@colby.edu

<u>Abstract (250)</u>

Whether coevolution determines distribution patterns of facultative symbionts among individuals
of a host species is an open question. We used the *Dictyostelium discoideum*-*Paraburkholderia*
system to investigate the role of host fitness and symbiont transmission in co-occurrence
patterns of naturally occurring (native) host-symbiont strain pairs. *P. bonniea* is able to
persistently infect *D. discoideum* in the lab, yet few wild *D. discoideum* isolates natively host this
symbiont. If native host-symbiont combinations lead to higher fitness for host and symbiont
compared to nonnative combinations, we assumed a stronger chance that coevolution had led
to their co-occurrence. We approached the factors that influence the fitness consequences of
host-symbiont interactions in terms of host defense mechanisms and symbiont virulence
mechanisms. Given a specific combination of host and symbiont, variable host resistance and
symbiont proliferation lead to variation in symbiont density, while variable host tolerance and
symbiont benevolence lead to variation in host fitness relative to symbiont density. We used
both vegetative and social stages of the host life cycle to approximate interactions between *D.*
discoideum and *P. bonniea*, when hosts form social groups and symbiont horizontal
transmission is facilitated by these social groups. We found no difference in host defense
mechanisms between native and nonnative hosts of *P. bonniea*, but *P. bonniea* possesses
variable phenotypes in proliferation and benevolence. More importantly, variation in symbiont
benevolence led to increased horizontal transmission of benevolent strains compared to
malevolent strains during the social stage of the host life cycle.

<u>Importance (150)</u>

Hosts and symbionts in a facultative symbiosis do not depend on each other for survival.
Perhaps due to this labile relationship, we lack a solid understanding of why only some
individual hosts carry facultative symbionts rather than all. We used the social amoeba
Dictyostelium discoideum and its bacterial symbiont *Paraburkholderia bonniea* to test whether
naturally occurring (native) host-symbiont strain pairs are the result of coevolution. We find that
native combinations of hosts and symbionts do not enjoy improved fitness compared to
nonnative combinations. Instead, both host and symbiont fitness depend on symbiont identity,
as *P. bonniea* strains varied in their ability to proliferate within groups of hosts and in the degree

of fitness loss they caused given a similar infection prevalence. More importantly, variation in
the latter property, known as benevolence, led to increased horizontal transmission of
benevolent strains compared to malevolent strains during the social stage of the host life cycle.
47
48 <u>Running title</u>
49 Symbiont benevolence impacts horizontal transmission
50
<u>Introduction</u>
Symbiosis can be broadly defined as an intimate and prolonged association among different
organisms, inclusive of positive, neutral, or negative fitness outcomes to the partners involved.
Host-symbiont relationships can be obligate or facultative. Obligate symbioses, when host and
symbiont need each other to survive, are expected to be the result of reciprocal selection
between host and symbiont over a long period of evolutionary time (Frank 1995; Law and
Dieckmann 1998). In contrast, facultative symbioses, in which host and symbiont can each
survive in a free-living state, are often assumed to be the result of shorter term relationships
(Moran et al. 2008). However, stable long-term facultative symbiotic relationships may be more
common than expected. For example, phylogenetic codivergence patterns support the stability
of the ancient facultative symbiosis between *Glomeribacter* and Glomeromycota fungi (Mondo
et al. 2012). Recurring co-infection patterns of specific symbionts support the stability of
heritable facultative microbial communities that infect the pea aphid *Acyrthosiphon pisum* (Rock
et al. 2018). Stable facultative symbioses are well suited to reveal how symbiotic relationships
are initiated and maintained over evolutionary time, because both host and symbiont can
independently evolve in response to any changes in the ecological context of the symbiotic
relationship. However, many aspects of facultative symbioses are relatively poorly understood
by virtue of the labile relationship. This includes which factors determine distribution patterns of
facultative symbionts among individuals within a host species (Niepoth et al. 2018). For
example, we lack a solid understanding of why some individual hosts carry facultative symbionts
while others do not.
The potential for, and outcome of, long-term association between a host and any
potential symbionts is determined by host genetics, symbiont genetics, and various
environmental factors (Schmid-Hempel 2011). We approached the specific factors that influence
the fitness consequences of host-symbiont interactions in terms of host defense mechanisms
and symbiont virulence mechanisms. Host resistance (the ability to limit the extent of an
infection) and tolerance (the ability to tolerate the physiological consequences of an infection)
are the two large categories of host defense mechanisms that affect evolutionary outcomes for
symbioses (Simms and Triplett 1994; Råberg et al. 2007; Ayres and Schneider 2012). The
equivalent symbiont virulence mechanisms are proliferation (the ability to proliferate to a certain
extent in an infection) and benevolence (the ability to cause beneficial or harmful physiological
consequences of an infection) (Wollein Waldetoft et al. 2020). Given a specific combination of
host and symbiont, variable host resistance and symbiont proliferation lead to variation in
symbiont density, while variable host tolerance and symbiont benevolence lead to variation in
host fitness consequences relative to symbiont density (Figure 1). The evolution of host
resistance and tolerance have epidemiological consequences, as host resistance often limits
the spread of infections while host tolerance can do the opposite and cause infections to

increase in frequency in a population (Roy and Kirchner 2000; Råberg 2014). The evolution of
symbiont benevolence and proliferation have significance in therapeutic drug development, as
targeting pathogen benevolence rather than proliferation has been suggested as a potential way
to prevent counter adaptation such as the evolution of antimicrobial resistance (Wollein
Waldetoft et al. 2020).

The amoeba *Dictyostelium discoideum* is an established model for understanding
intracellular pathogen infections (Steinert and Heuner 2005). It forms a facultative symbiosis
with three species of *Paraburkholderia*: *P. agricolaris*, *P. bonniea*, and *P. hayleyella* (Brock et al.
2020). Roughly one quarter of soil-isolated *D. discoideum* strains carry one or occasionally
multiple species of these symbiotic *Paraburkholderia* (Haselkorn et al. 2019). All three species
of *D. discoideum*-associated *Paraburkholderia* are unique among close relatives in their ability
to persistently infect *D. discoideum* (DiSalvo et al. 2015; Haselkorn et al. 2019). But two of the
Paraburkholderia symbionts have genomes that are half the size of the third (Brock et al. 2020;
Noh et al. 2022). Longer term symbionts tend to have smaller genomes compared to free-living
relatives (Mira et al. 2001; Weinert and Welch 2017). While *P. agricolaris* is difficult to
distinguish from other *Paraburkholderia* species based on its genome size and content,
comparative genomics analyses indicate that *P. bonniea* and *P. hayleyella* have evolved
reduced genomes in the amoeba host environment (Noh et al. 2022).

The fitness outcomes of *D. discoideum*-*Paraburkholderia* association appear to be
context-dependent as with most facultative host-microbe symbioses (Drew et al. 2021; Husnik
et al. 2021). *D. discoideum* hosts generally suffer negative fitness consequences of association.
Amoebas infected with *Paraburkholderia* symbionts in the lab tend to eat less food bacteria
during vegetative growth, migrate shorter distances as slugs during their multicellular social
stage, form shorter and smaller volume fruiting bodies, produce fewer spores, and carry other
bacteria alive (secondary carriage) into their next vegetative growth stage (Brock et al. 2011;
DiSalvo et al. 2015; Shu et al. 2018; Miller et al. 2020). Potential context-dependent fitness
benefits to the host appear to be an improved competitive ability against other *D. discoideum*
strains, hypothetically by passing on symbiont infections or releasing *Paraburkholderia*
secretions in a defensive manner (Brock et al. 2013). We know less about fitness outcomes of
association for *Paraburkholderia* symbionts but *P. hayleyella* (though not *P. agricolaris*) reaches
higher densities in the presence of *D. discoideum* compared to on its own in soil medium
(Garcia et al. 2019).

Our goal was to understand whether naturally occurring (native) host-symbiont strain
pairs within one of the reduced genome species, *P. bonniea*, is the result of host-symbiont
coevolution. We use the term native host to indicate a *D. discoideum* strain from which one of
the three *Paraburkholderia* symbiont species were isolated or identified. If native host-symbiont
combinations lead to higher fitness estimates for host and symbiont compared to nonnative
combinations, we assumed a stronger chance that coevolution had led to their co-occurrence.
Patterns of coevolution were previously tested for the two other *D. discoideum* symbionts, *P.*
agricolaris and *P. hayleyella*, but not for *P. bonniea*. In response to infection with *P. hayleyella*,
nonnative and native *P. agricolaris* hosts had a higher fitness cost compared to native *P.*
hayleyella hosts. In contrast, nonnative hosts, native *P. agricolaris* hosts, and native *P.*
hayleyella hosts respond with similarly decreased fitness to *P. agricolaris* infection (Shu et al.
2018). *P. hayleyella* had higher abundance in soil with *D. discoideum* than soil without, while *P.*

agricolaris abundance was not affected by *D. discoideum* presence (Garcia et al. 2019). Both
studies suggest that coevolution has occurred for *P. hayleyella* hosts and symbionts but not for
P. agricolaris hosts and symbionts. Because *P. hayleyella* has a reduced genome, as does *P.*
bonniea, we expected to see support for coevolution in our study.
D. discoideum has single cell and multi-cell stages in its life cycle. Vegetative single cell
amoebas feed on bacteria until food density becomes low and starvation begins. Now in their
social stage, starving amoebas secrete a chemical signal that causes amoebas within range to
aggregate toward higher concentrations of this signal. This leads to aggregation, which is
followed by transformation of the aggregated group into a multicellular body that ultimately
forms fruiting bodies. The amoebas that survive the social stage become spores contained
within the fruiting body sorus. In the lab, both stages of the life cycle occur as *D. discoideum*
spores are allowed to hatch on petri dishes with food bacteria. Vegetatively multiplying amoebas
will eventually exhaust their available food, aggregate, and ultimately form fruiting bodies after
going through a social stage *in situ*.
We used both vegetative and social stages of this life cycle to approximate interactions
between *D. discoideum* hosts and *P. bonniea* symbionts in an environment in which hosts form
social groups and symbiont horizontal transmission is facilitated by these social groups. We
used the vegetative stage to expose *D. discoideum* amoebas to *P. bonniea* symbionts, and
used spore counts after an unmanipulated social stage as a measure of host fitness. We then
used a subsequent social stage to expose uninfected amoebas to pre-infected amoebas, and
estimated symbiont horizontal transmission after this manipulated social stage as an important
aspect of symbiont fitness. We estimate host fitness and symbiont transmission at the social
group level rather than at the individual amoeba host level using the social stage of the *D.*
discoideum life cycle. Our estimates of symbiont density (e.g. Figure 1) is therefore better
described as symbiont infection prevalence within a group of amoebas, and host fitness is also
not a direct measure of individual host mortality but instead a relative measure of group
productivity given a degree of symbiont prevalence. The same is true for symbiont transmission
and is an estimate of transmission that occurs within a social group of interacting hosts rather
than transmission from one host to another, as our estimates are a measure of newly added
symbiont infection prevalence in a group of amoebas given a previous degree of symbiont
infection prevalence.
163
<u>Methods</u>
Focal host and symbiont strains
We used three native host strains of *P. bonniea* (QS395, QS433, QS859) and three
nonnative *D. discoideum* strains (QS4, QS17, QS18). All *D. discoideum* strains were previously
isolated from Mountain Lake Biological Station in Virginia, USA. The native host strains had
been cured of their symbionts using tetracycline and verified as symbiont-free using PCR. We
used three *P. bonniea* symbiont strains (BbQS395, BbQS433, BbQS859). For each replicate,
host (with food bacteria *Klebsiella pneumoniae*) and symbiont strains were grown from freezer
stock on SM/5 plates (2 g glucose, 2 g BactoPeptone (Oxoid), 2 g yeast extract (Oxoid), 0.2 g
$MgCl_2$, 1.9 g KH_2PO_4, 1 g K_2HPO_4 and 15 g agar per liter). KK2 buffer (2.2 g KH2PO4
monobasic and 0.7 g K2HPO4 dibasic per liter) was used throughout for handling bacteria, *D.*
discoideum spores and amoebas.

Host fitness

To estimate host fitness at a range of infection prevalence, *D. discoideum* spores from
freshly grown fruiting bodies were collected and placed on SM/5 plates at a density of 2 x 10^5
spores per plate and RFP-labeled symbionts at multiplicities of infection (MOI) of 0 (control),
0.6, 3, and 15. We made three replicate plates per combination and left these plates to fruit at
21 C for 5-7 days. We then collected all fruiting bodies from each plate and estimated the
number of spores by counting diluted spores on a hemocytometer and multiplying these counts
by the total volume of spores. We then took the percent of spores from each host-symbiont
combination relative to the uninfected control that was prepared at the same time. For example,
host strain QS4 was infected with BbQS395 at MOI of 0 (control), 0.6, 3, and 15. Spore counts
of each individual sample plate were divided by the average spore count of the three uninfected
control plates. Because our models examine the relationship between host fitness and infection
prevalence (not MOI), we estimated infection prevalence (percent of RPF+ infected spores in a
sample of 100,000) by running a sample of spores on a BD FACSCalibur flow cytometer
(Becton, Dickinson and Company, Franklin Lakes, NJ) to.

Symbiont transmission

To estimate symbiont horizontal transmission during a social stage, *D. discoideum*
spores from freshly grown fruiting bodies were collected and placed on SM/5 plates at a density
of 2 x 10^5 spores per plate with RFP-labeled symbionts at MOI of 0 (control), plus an additional
3-5 MOI ranging from 0.3 to 30. Once these plates had fruited and amoebas were pre-infected,
we collected fresh spores and placed them on SM/5 plates at densities of 1 x 10^5 and 2 x 10^5
spores per plate with only food bacteria. When amoebas were at log-phase growth roughly 36
hours later, uninfected amoebas were dyed with CellTracker Green CMFDA (Invitrogen)
dissolved in DMSO. Pre-infected amoebas (RFP+) were carried along with the dyed amoebas,
only exposed to DMSO, and treated in the same way through room temperature incubation and
washes. Afterward, dyed uninfected amoebas were combined with infected amoebas at 1:0
(dyed only negative control), 0:1 (infected only), and 1:1 (mixed) ratios and placed on moistened
nitrocellulose filters to aggregate and develop into fruiting bodies in replicates of three. 5-7 days
later, we collected all fruiting bodies from these filters and ran these samples on a BD
FACSCalibur flow cytometer. We estimated infection prevalence from the infected only sample
and horizontal transmission from the mixed sample as indicated by the co-occurrence of dye
and infection in a spore. FCS files were imported into FlowJo v10.8.1 for analysis. We applied
gates for spores, then dyed cells from the negative control, and infected cells from the highest
MOI sample. A logical (AND) gate was applied to the 1:1 mixed sample to find the percent of
spores that were previously uninfected amoebas but were now positive for symbiont infection
(Figure 2).

Statistical analysis

Statistical analyses were performed in R v3.6.0 and with packages car v.3.0-12 (Fox and
Weisberg 2019) and lme4 v.1.1-28 (Bates et al. 2015). For both host fitness and symbiont
transmission, we fit linear models with mixed effects. For host fitness, we tested how percent
spores (relative percent of spores produced) were affected by symbiont strain and host type

(native or nonnative). We tested the effect of host type by coding the native combinations either
strictly (e.g. only QS859 infected with BbQS859 would be considered native) or leniently (e.g.
any native host of *P. bonniea* infected with any *P. bonniea* strain is considered native). We used
infection prevalence as a continuous predictor and experiment date and host strain as random
effects. For symbiont transmission, we fit a similar model but with percent transmission (percent
newly infected spores that were previously uninfected amoebas) as the dependent variable. We
fit the most complex model first, then removed nonsignificant terms and compared the simpler
model with the complex one using a chi-squared test. We accepted the simpler model if no
significant difference was found in model fit. We examined residuals of the final model for any
indications that we had violated model assumptions. We examined effect sizes of factors in the
final model using the package effectsize v.0.7.0 (Ben-Shachar et al. 2020) and its
epsilon_squared() function. We also compared infection prevalence across the two
experiments. We fit linear models with host strain as a random effect and the different number
of social stages (nonnumeric factor; one for host fitness, two for symbiont transmission) and
symbiont identity as fixed effects. We examined residuals of the final model for any indications
that we had violated model assumptions and examined effect sizes of factors.
236
237   *Data availability*
238   Data and R code supporting this work can be accessed at:
239   https://github.com/noh-lab/social-infection
240
241   <u>Results</u>
242
Host fitness
We estimated the relationship between host fitness and symbiont infection prevalence
by exposing native and nonnative *D. discoideum* hosts to three strains of *P. bonniea* at a range
of multiplicities of infection. Host fitness decreased with increasing infection prevalence across
all combinations of hosts and symbionts that we tested, but there was no difference between
native and nonnative hosts (Figure 3; Table 1). This was regardless of whether native host
status was defined using a strict or lenient definition (Table S1). Instead, symbiont strain identity
significantly affected host fitness. For example, BbQS859 infections resulted in a shallower
slope in host fitness decline compared to BbQS433 infections which resulted in a steeper slope
(Figure 3). These results support a significant effect of symbiont benevolence variation on host
fitness during initial infection. Based on our results, BbQS859 is more benevolent while
BbQS433 is more malevolent to any type of host. BbQS395 appears to show an intermediate
benevolence phenotype.
256
Symbiont transmission
We estimated the relationship between symbiont transmission and initial symbiont
infection prevalence by exposing uninfected amoebas to pre-infected amoebas during a
manipulated social stage of the host life cycle and detecting the percentage of newly infected
spores. The degree of symbiont horizontal transmission increased with the infection prevalence
of pre-infected amoebas that entered each social stage (Figure 4). This positive correlation
confirms that the increase in infection is due to horizontal transmission rather than vertical

transmission (Ebert 2013), which is what we expected because very few amoebas go through
cell division during the social stage (Muramoto and Chubb 2008). As with host fitness above,
native and nonnative hosts were no different in overall patterns of symbiont transmission,
regardless of how native hosts are defined (Table 2). However, the degree and slope of
symbiont transmission was affected by symbiont strain identity, indicating that both symbiont
proliferation and benevolence affect symbiont transmission (Figure 4). For example, over more
than a single social stage, BbQS859 has higher proliferation compared to BbQS395 and
BbQS433. In addition, BbQS859 symbionts were transmitted at significantly higher levels
compared to the other symbiont strains, and at a steeper slope beyond that simply due to
BbQS859 infections occurring at higher infection prevalence. We also detected some
differences in how infection prevalence, and host type interacted when we leniently defined
native hosts. Howeve, much of this difference may be attributed to BbQS859, where nonnative
hosts became infected at a slightly higher density compared to native hosts and therefore also
transmitted symbionts at a higher level.
278
Synthesis
We observed a significant difference between infection prevalence in these two
experiments. This is likely due to differences in symbiont strain-specific virulence, both in terms
of proliferation and benevolence (Table 3; Figure 5). The main difference between the design of
the two experiments is that for *Host fitness* we used newly-infected amoebas, but for *Symbiont*
transmission we used pre-infected amoebas that had been infected in a previous vegetative
stage so as to minimize the potential for symbionts to be newly acquired from the environment
and to maximize our ability to detect horizontal transmission among amoebas. We infected
amoebas for the symbiont transmission experiment at a wider range of MOI (0.3-30) compared
to the host fitness experiment (0.6-15) in order to achieve as wide a range of infection
prevalence as possible. We found that we were unable to push the maximum of these ranges
higher by using a higher MOI for the initial infection. In fact, typically one of the intermediate
MOI resulted in the highest observed infection prevalence after the amoebas had gone through
a social stage.
Our combined observations from both experiments suggest that variation in symbiont
benevolence leads to variation in symbiont proliferation among *P. bonniea* strains infecting
amoeba hosts. More malevolent strains (BbQS395 and BbQS433) proliferate less among
amoeba hosts compared to more benevolent strains (BbQS859) over the course of two
vegetative stage - social stage cycles (Figure 5). This leads to less transmission of malevolent
strains compared to benevolent strains during the next social stage (Figure 4).
299
Discussion
Host and microbe partners engaged in a facultative symbiosis are not mutually
dependent on each other for survival as are partners in an obligate symbiosis. Yet stable
associations between hosts and facultative symbionts clearly exist (Mondo et al. 2012; Werner
et al. 2015). *P. bonniea*, along with *P. hayleyella* and *P. agricolaris*, is able to persistently infect
D. discoideum and form a facultative symbiosis (DiSalvo et al. 2015; Haselkorn et al. 2019;
Miller et al. 2020). Both *P. bonniea* and *P. hayleyella* have reduced genomes that are roughly
half the size of *P. agricolaris*, leading to the hypothesis that both reduced-genome species may

have associated with *D. discoideum* for a longer period of evolutionary time compared to *P.*
agricolaris (Brock et al. 2020; Noh et al. 2022), and thus forming a stable facultative symbiosis.
Whereas we had expected that the presence of reduced-genome symbionts may likely be due
to host-symbiont coevolution, our results indicate this may not be the case for hosts and *P.*
bonniea symbionts. Our results from *P. bonniea* show the presence of variation in symbiont
virulence mechanisms and a lack of variation in *D. discoideum* host defense mechanisms in
response to *P. bonniea* infection. The fitness outcomes of infected hosts were no different
between native and nonnative hosts. Overall, when more individual hosts were infected, the
collective number of spores produced decreased for both types of hosts, indicating that infection
with *P. bonniea* is a detriment to amoeba host fitness. We did find evidence for strain level
variation in symbiont virulence mechanisms that affect symbiont transmission, by enabling
increased horizontal transmission in benevolent strains compared to malevolent strains. Overall,
when more individual hosts were infected symbionts were better able to infect additional hosts
via horizontal transmission, suggesting that *P. bonniea* benefits from infecting more amoeba
hosts.
323
Lack of variation in host defense mechanisms – ecology and host range of Paraburkholderia
symbionts
Our results indicate that native hosts of *P. bonniea* have not evolved improved defense
mechanisms in response to association with *P. bonniea* compared to nonnative hosts. We find
this to be a curious result. Both *P. hayleyella* and *P. bonniea* have reduced genomes and are
each other's closest relatives (Brock et al. 2020). Some evidence of host-symbiont coevolution
has been observed for *P. hayleyella* and its native hosts, though with different experimental
approaches (Shu et al. 2018; Garcia et al. 2019). *P. bonniea* had remained untested until now.
Our post-hoc hypothesis for the lack of defense mechanisms in hosts of *P. bonniea* is related to
the ecology of both host and symbiont. *P. bonniea* is the rarest of *Paraburkholderia* symbionts
of *D. discoideum* (Haselkorn et al. 2019; DuBose et al. 2022) and it also appears to be the
symbiont species that is least detrimental to host fitness on average (Miller et al. 2020). Rare
encounters between new hosts and *P. bonniea* may not have a significant impact on *D.*
discoideum populations, particularly if *D. discoideum* itself is sparsely distributed across soil
landscapes. If both host and symbiont are rare, we expect the fitness cost of infection for the
relatively benign *P. bonniea* to be insufficient for hosts to evolve defense mechanisms against it
(Anderson and May 1982).
The apparent lack of evolved host defense mechanisms in native hosts of *P. bonniea*
brings us to an interesting question regarding *Paraburkholderia* symbionts of *D. discoideum*
social amoebas: are they generalist or specialist symbionts to *D. discoideum*? *P. agricolaris* and
other *Burkholderia* sensu lato appear to frequently infect Dictyostelid and Acytostelid social
amoeba species (Sallinger et al. 2021). Comparison between soil microbiomes and amoeba
microbiomes found that *Burkholderia* are present in soil samples but are orders of magnitude
more abundant in amoebas. This increased abundance suggests that bacteria that can
associate with social amoebas are benefiting from being separated from various environmental
stresses and competition from other soil bacteria, and potentially also adapted to living inside
social amoebas. These pieces of evidence indicate that *P. agricolaris* may be a generalist
symbiont of social amoebas, and *P. agricolaris* infection from the environment and horizontal

infection between amoebas may be common. The rarer *P. bonniea* and *P. hayleyella* are still
elusive in terms of their natural host ranges (DuBose et al. 2022). But given their relative
scarcity and difference in genome size, we predict that both reduced-genome species are more
likely to be specialists of *D. discoideum*. The host range of a symbiont is affected by ecological
factors including host density and relatedness, symbiont transmission dynamics, and the
presence of additional symbionts competing for access to host resources (Barrett et al. 2009;
Leggett et al. 2013). The projected ecological differences between *P. agricolaris* and the
reduced genome species *P. bonniea* and *P. hayleyella* appear to be consistent with previous
observations that generalist symbionts (potentially *P. agricolaris*) are more likely to have
environmental origins while specialists (potentially *P. bonniea* and *P. hayleyella*) are more likely
to have been commensals evolving within a host species (Brown et al. 2012; Bäumler and Fang
2013).
364
Strain level variation in symbiont virulence mechanisms – virulence-transmission trade-off
We find evidence that strains of *P. bonniea* vary in both symbiont virulence mechanisms
of proliferation and benevolence. We consider the virulence-transmission trade-off to
contextualize this observed variation in the interaction between *D. discoideum* and *P. bonniea*.
Despite the commonly-held expectation that relatively benign symbionts will result from long-
term host-symbiont coevolution, the virulence-transmission trade-off hypothesis does not have a
guaranteed expectation of what will happen to symbiont virulence per se. Instead, the trade-off
hypothesis posits that if there is a biological link between symbiont virulence and transmission
(because symbionts use host resources to replicate), symbionts should evolve toward a level of
virulence where transmission is greatest (Anderson and May 1982). This hypothesized trade-off
has proven difficult to observe for several reasons, including the influences of host range,
multiple symbiont infections, and host population structure that can affect host-symbiont
interactions in complex ways (Alizon et al. 2009; Leggett et al. 2013). However, recent meta-
analyses indicate that there is partial support for the trade-off hypothesis in several biological
systems (Acevedo et al. 2019). The three core predictions of the trade-off hypothesis are as
follows: (i) within-host symbiont replication rates have a positive relationship with symbiont
virulence; (ii) within-host symbiont replication rates have a positive relationship with symbiont
transmission rates; and (iii) symbiont virulence and transmission have a trade-off through the
negative relationship between symbiont virulence and host recovery rates. Diverse study
systems show support for (i) and (ii), but only a nonsignificant trend for (iii) (Acevedo et al.
2019).
Because we estimated host fitness and symbiont transmission at the social group level
rather than at the individual amoeba host level, our evidence that supports each of these points
may be considered an indirect test of the virulence-transmission trade-off as the original theory
is built on investigating symbiont density in individual hosts. That said, we find support for each
of the core predictions of the virulence-transmission trade-off hypothesis. Symbiont infection
prevalence had a negative relationship with host fitness (i; Figure 3), and symbiont infection
prevalence had a positive relationship with symbiont horizontal transmission (ii; Figure 4).
Lastly, variation in symbiont benevolence led to increased horizontal transmission of benevolent
strains compared to malevolent strains (iii; Figure 4). The patterns of symbiont virulence
mechanisms we observed also predict that more benevolent *P. bonniea* strains should be found

at higher frequencies given similar conditions compared to more malevolent strains. Because of
the relative scarcity of *P. bonniea* in nature, we may never observe the ecological consequence
of this trade-off we have detected in the lab. However the presence and extent of horizontal
transmission in *P. bonniea* indicates that this symbiont benefits from infecting *D. discoideum*
hosts by both proliferating within a given host and transmitting to new hosts.
We expect transmission dynamics to be an important factor in further understanding *P.*
bonniea and *P. hayleyella* evolution because of the complex interaction between host
specificity, symbiont virulence, and symbiont transmission. Given the persistence of
Paraburkholderia symbiont infection in the lab and the facultative aspect of the symbiosis itself,
we expect these symbionts to transmit among hosts using both vertical and horizontal routes.
While we did not test for vertical transmission, our results clearly show the presence of
significant horizontal symbiont transmission among *D. discoideum* amoebas in the social stage
of their life cycle. The coexistence and relative influences of both types of transmission routes
are better understood in other amoeba-bacteria symbioses. For example, *Acanthamoeba* hosts
interact with *Parachlamydia acanthamoebae* symbionts across the cyclical life cycle of the
symbiont (Herrera et al. 2020). The symbiont can replicate within a host-derived vacuole in the
form of reticulate bodies that are also vertically transmitted from parent to daughter cells.
Parachlamydia symbionts are released into the environment in the form of elementary bodies
that do not replicate but can horizontally infect new hosts.
415
Coevolution or random co-occurrence of host and symbiont in a facultative symbiosis
Our results from *P. bonniea* show the presence of variation in symbiont virulence
mechanisms and a lack of variation in *D. discoideum* host defense mechanisms between native
and nonnative hosts. Therefore, naturally occurring host-*P. bonniea* strain pairs do not appear
to be the result of evolved host defense mechanisms in native hosts compared to nonnative
hosts. The presence of phenotypic variation among symbiont strains and in some host-symbiont
combinations indicates that adaptation is potentially ongoing for *P. bonniea* symbionts. In
contrast the lack of variation among *D. discoideum* host types can indicate a lack of adaptation
potential but can also be the result of host defense mechanism fixation that has occurred post-
adaptation. A lack of adaptation potential in host defense mechanisms against *P. bonniea* may
be because *P. bonniea* presents only a weak selection pressure, due to the mild reduction in
fitness experienced by amoeba hosts even at high infection prevalence and what we expect is
the relatively infrequent encounter rate of *P. bonniea* by environmental *D. discoideum*. We plan
to apply the experimental approach of the current study to the relationship between *D.*
discoideum and its other reduced-genome symbiont species *P. hayleyella*, which is known to
cause much more detrimental fitness consequences to newly infected hosts (Shu et al. 2018;
Miller et al. 2020). Additional results from *P. hayleyella* should provide more insight into the
patterns we observed using *P. bonniea*.
Attempts to find individual strain-level coevolution in facultative symbionts are rare.
Niepoth and colleagues (2018) found that one particular *Hamiltonella defensa* symbiont strain
caused detrimental fitness effects to nonnative pea aphid hosts compared to the native host.
However, the detrimental fitness effects to nonnative hosts appeared to be transient, and did not
continue after infection establishment. In addition, hosts from a population (*Lotus pedunculatus*
biotype) that typically carry *H. defensa* were more likely to establish symbiosis when newly

infected compared to hosts from another population (*Lotus corniculatus* biotype). These results
suggest that a potential reason why we did not find evidence of host adaptation is because our
native and nonnative hosts were collected from the same geographical locality. Therefore, it is
possible that *D. discoideum* hosts from a locality where *P. bonniea* symbionts are found have
already adapted to their infection and host defense mechanisms have become fixed. At this
time, it is unclear how to test whether the patterns we observed with *P. bonniea* are the result of
post-adaptation fixation because we do not have a way to determine whether an amoeba
collected from the field has encountered a *Paraburkholderia* previously.
448
Conclusions
The *D. discoideum-Paraburkholderia* system continues to provide opportunities for
investigating the evolution of a naturally occurring facultative symbiosis between a protist host
and bacterial symbionts. In addition, we find this system to be particularly valuable in that it
presents an opportunity to study symbiosis in the context of social groups of hosts, so that we
can readily assess the extent and impact of symbiont horizontal transmission on both host and
symbiont. In summary, our work shows that *P. bonniea* possesses variable phenotypes in
symbiont proliferation and benevolence, whereas native and nonnative hosts of *P. bonniea* do
not differ in host defense mechanisms in response to association with *P. bonniea*. It is difficult to
extrapolate what kind of evolutionary consequences may arise based on differences in
experimentally estimated fitness effects in microbes, because variation in microbial population
size can easily range across several orders of magnitude (Sheppard et al. 2018). However, the
presence and extent of horizontal transmission in *P. bonniea* suggests that this symbiont
benefits from infecting *D. discoideum* hosts by both proliferating within a given host and
transmitting to new hosts. More importantly, we find support for the virulence-transmission
trade-off, as variation in symbiont benevolence led to increased horizontal transmission of
benevolent strains compared to malevolent strains during the social stage of the host life cycle.
In contrast, native *D. discoideum* hosts do not appear to have a fitness advantage over
nonnative *D. discoideum* hosts in their interactions with *P. bonniea*. Therefore, naturally
occurring host-*P. bonniea* strain pairs do not appear to be the result of coevolution between
host and symbiont.
470

Acknowledgements
We thank past and current members of the Noh lab (Caroline Lunt, Gus Shuster) who
contributed moral support for this project during its development. Ron Peck, Laura Runyen-
Janecky, and Chris Moore provided helpful comments that improved previous drafts.
This work was supported by the National Institutes of Health and its National Institute of
General Medical Sciences by an Institutional Development Award (IDeA) under Grant Number
P20GM103423 (subaward to SN) and Colby College startup funds.
478

References
Acevedo, M. A., F. P. Dillemuth, A. J. Flick, M. J. Faldyn, and B. D. Elderd. 2019. Virulence-

driven trade-offs in disease transmission: A meta-analysis*. Evolution 73:636–647.

Alizon, S., A. Hurford, N. Mideo, and M. Van Baalen. 2009. Virulence evolution and the trade-off

hypothesis: history, current state of affairs and the future. Journal of Evolutionary Biology

22:245–259.

Anderson, R. M., and R. M. May. 1982. Coevolution of hosts and parasites. Parasitology

85:411–426. Cambridge University Press.

Ayres, J. S., and D. S. Schneider. 2012. Tolerance of infections. Annual Review of Immunology

30:271–294.

Barrett, L. G., J. M. Kniskern, N. Bodenhausen, W. Zhang, and J. Bergelson. 2009. Continua of

specificity and virulence in plant host–pathogen interactions: causes and consequences.

New Phytologist 183:513–529.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using

lme4. Journal of Statistical Software 67:1–48.

Bäumler, A., and F. C. Fang. 2013. Host specificity of bacterial pathogens. Cold Spring Harb

Perspect Med 3:a010041. Cold Spring Harbor Laboratory Press.

Ben-Shachar, M. S., D. Lüdecke, and D. Makowski. 2020. effectsize: Estimation of effect size

indices and standardized parameters. Journal of Open Source Software 5:2815.

Brock, D. A., T. E. Douglas, D. C. Queller, and J. E. Strassmann. 2011. Primitive agriculture in a

social amoeba. Nature 469:393–396.

Brock, D. A., S. Noh, A. N. M. Hubert, T. S. Haselkorn, S. DiSalvo, M. K. Suess, A. S. Bradley,

M. Tavakoli-Nezhad, K. S. Geist, D. C. Queller, and J. E. Strassmann. 2020.

Endosymbiotic adaptations in three new bacterial species associated with *Dictyostelium*

discoideum: *Paraburkholderia agricolaris* sp. nov., *Paraburkholderia hayleyella* sp. nov.,

and *Paraburkholderia bonniea* sp. nov. PeerJ 8:e9151.

Brock, D. A., S. Read, A. Bozhchenko, D. C. Queller, and J. E. Strassmann. 2013. Social

amoeba farmers carry defensive symbionts to protect and privatize their crops. Nature

Communications 4:2385.

Brown, S. P., D. M. Cornforth, and N. Mideo. 2012. Evolution of virulence in opportunistic

pathogens: generalism, plasticity, and control. Trends in Microbiology 20:336–342.

DiSalvo, S., T. S. Haselkorn, U. Bashir, D. Jimenez, D. A. Brock, D. C. Queller, and J. E.

Strassmann. 2015. *Burkholderia* bacteria infectiously induce the proto-farming symbiosis

of *Dictyostelium* amoebae and food bacteria. PNAS 112:E5029–E5037.

Drew, G. C., E. J. Stevens, and K. C. King. 2021. Microbial evolution and transitions along the

parasite–mutualist continuum. Nat Rev Microbiol 19:623–638. Nature Publishing Group.

DuBose, J. G., M. S. Robeson, M. Hoogshagen, H. Olsen, and T. S. Haselkorn. 2022.

Complexities of inferring symbiont function: *Paraburkholderia* symbiont dynamics in

social amoeba populations and their impacts on the amoeba microbiota. Applied and

Environmental Microbiology 0:e01285-22. American Society for Microbiology.

Ebert, D. 2013. The epidemiology and evolution of symbionts with mixed-mode transmission.

Annu. Rev. Ecol. Evol. Syst. 44:623–643.

Fox, J., and S. Weisberg. 2019. An R companion to applied regression. 3rd ed. Sage, Thousand

Oaks, CA.

Frank, S. A. 1995. The origin of synergistic symbiosis. Journal of Theoretical Biology 176:403–

410.

Garcia, J. R., T. J. Larsen, D. C. Queller, and J. E. Strassmann. 2019. Fitness costs and

benefits vary for two facultative *Burkholderia* symbionts of the social amoeba,

Dictyostelium discoideum. Ecology and Evolution 9:9878–9890.

Haselkorn, T. S., S. DiSalvo, J. W. Miller, U. Bashir, D. A. Brock, D. C. Queller, and J. E.

Strassmann. 2019. The specificity of *Burkholderia* symbionts in the social amoeba

farming symbiosis: prevalence, species, genetic and phenotypic diversity. Molecular

Ecology 28:847–862.

Herrera, P., L. Schuster, C. Wentrup, L. König, T. Kempinger, H. Na, J. Schwarz, S.

Köstlbacher, F. Wascher, M. Zojer, T. Rattei, and M. Horn. 2020. Molecular causes of an

evolutionary shift along the parasitism–mutualism continuum in a bacterial symbiont.

Proceedings of the National Academy of Sciences 117:21658–21666. Proceedings of

the National Academy of Sciences.

Husnik, F., D. Tashyreva, V. Boscaro, E. E. George, J. Lukeš, and P. J. Keeling. 2021. Bacterial

and archaeal symbioses with protists. Current Biology 31:R862–R877.

Law, R., and U. Dieckmann. 1998. Symbiosis through exploitation and the merger of lineages in

evolution. Proceedings of the Royal Society of London. Series B: Biological Sciences

265:1245–1253. Royal Society.

Leggett, H. C., A. Buckling, G. H. Long, and M. Boots. 2013. Generalism and the evolution of

parasite virulence. Trends in Ecology & Evolution 28:592–596.

Miller, J. W., C. R. Bocke, A. R. Tresslar, E. M. Schniepp, and S. DiSalvo. 2020.

Paraburkholderia symbionts display variable infection patterns that are not predictive of

amoeba host outcomes. Genes 11:674. Multidisciplinary Digital Publishing Institute.

Mira, A., H. Ochman, and N. A. Moran. 2001. Deletional bias and the evolution of bacterial

genomes. Trends in Genetics 17:589–596.

Mondo, S. J., K. H. Toomer, J. B. Morton, Y. Lekberg, and T. E. Pawlowska. 2012. Evolutionary

stability in a 400-million-year-old heritable facultative mutualism. Evolution 66:2564–

2576.

Moran, N. A., J. P. McCutcheon, and A. Nakabachi. 2008. Genomics and evolution of heritable

bacterial symbionts. Annual Review of Genetics 42:165–190.

Muramoto, T., and J. R. Chubb. 2008. Live imaging of the *Dictyostelium* cell cycle reveals

widespread S phase during development, a G2 bias in spore differentiation and a

premitotic checkpoint. Development 135:1647–1657.

Niepoth, N., J. Ellers, and L. M. Henry. 2018. Symbiont interactions with non-native hosts limit

the formation of new symbioses. BMC Evolutionary Biology 18:27.

Noh, S., B. J. Capodanno, S. Xu, M. C. Hamilton, J. E. Strassmann, and D. C. Queller. 2022.

Reduced and nonreduced genomes in *Paraburkholderia* symbionts of social amoebas.

mSystems 0:e00562-22.

Råberg, L. 2014. How to live with the enemy: understanding tolerance to parasites. PLoS Biol

12.

Råberg, L., D. Sim, and A. F. Read. 2007. Disentangling genetic variation for resistance and

tolerance to infectious diseases in animals. Science 318:812–814.

Rock, D. I., A. H. Smith, J. Joffe, A. Albertus, N. Wong, M. O'Connor, K. M. Oliver, and J. A.

Russell. 2018. Context-dependent vertical transmission shapes strong endosymbiont

community structure in the pea aphid, *Acyrthosiphon pisum.* Molecular Ecology

27:2039–2056.

Roy, B. A., and J. W. Kirchner. 2000. Evolutionary dynamics of pathogen resistance and

tolerance. Evolution 54:51–63.

Sallinger, E., M. S. Robeson, and T. S. Haselkorn. 2021. Characterization of the bacterial

microbiomes of social amoebae and exploration of the roles of host and environment on

microbiome composition. Environmental Microbiology 23:126–142.

Schmid-Hempel, P. 2011. Evolutionary Parasitology. Oxford University Press, New York, New

York.

Sheppard, S. K., D. S. Guttman, and J. R. Fitzgerald. 2018. Population genomics of bacterial

host adaptation. Nat Rev Genet 19:549–565. Nature Publishing Group.

Shu, L., D. A. Brock, K. S. Geist, J. W. Miller, D. C. Queller, J. E. Strassmann, and S. DiSalvo.

2018. Symbiont location, host fitness, and possible coadaptation in a symbiosis between

social amoebae and bacteria. eLife 7:e42660.

Simms, E. L., and J. Triplett. 1994. Costs and benefits of plant responses to disease: resistance

and tolerance. Evolution 48:1973–1985.

Steinert, M., and K. Heuner. 2005. *Dictyostelium* as host model for pathogenesis. Cell.

Microbiol. 7:307–314.

Weinert, L. A., and J. J. Welch. 2017. Why might bacterial pathogens have small genomes?

Trends in Ecology & Evolution 32:936–947.

Werner, G. D. A., W. K. Cornwell, J. H. C. Cornelissen, and E. T. Kiers. 2015. Evolutionary

signals of symbiotic persistence in the legume–rhizobia mutualism. Proceedings of the

National Academy of Sciences 112:10262–10269. Proceedings of the National Academy

of Sciences.

Wollein Waldetoft, K., L. Råberg, and R. Lood. 2020. Proliferation and benevolence—A

framework for dissecting the mechanisms of microbial virulence and health promotion.

Evolutionary Applications 13:879–888.

595
596
597
598

599   <u>Tables and Figures</u>

601   Table 1. Analysis of Deviance table for Host fitness with lenient native definition

Variable	X^2	df	P-value	η^2_p	95% CI
Infection prevalence	376.002	1	< 0.001	0.63	(0.56, 1)
Symbiont strain	17.998	2	< 0.001	- 0.06	(0, 1)
Host type	< 0.001	1	0.981	- 0.37	(0, 1)
Infection prevalence : Symbiont strain	23.886	2	< 0.001	0.09	(0.04, 1)

604   Table 2. Analysis of Deviance table for Symbiont transmission with lenient native definition

Variable	X^2	df	P-value	η^2_p	95% CI
Infection prevalence	484.965	1	< 0.001	0.77	(0.71, 1)
Symbiont strain	149.430	2	< 0.001	0.43	(0.20, 1)
Host type	2.092	1	0.148	0.53	(0.25, 1)
Infection prevalence : Symbiont strain	109.289	2	< 0.001	0.65	(0.52, 1)
Infection prevalence : Host type	35.991	1	< 0.001	0.30	(0.17, 1)
Symbiont strain : Host type	9.509	2	0.009	0.14	(0.01, 1)

607   Table 3. Analysis of Deviance table for infection prevalence across experiments

Variable	X^2	df	P-value	η^2_p	95% CI
Number of social stages	71.196	1	< 0.001	0.16	(0.11, 1)
Symbiont strain	17.789	2	< 0.001	0.02	(0.00, 1)
Number of social stages : Symbiont strain	39.197	2	< 0.001	0.09	(0.05, 1)

(a) Host defense mechanisms

(b) Symbiont virulence mechanisms

610
611

Figure 1. Theoretical differences between resistance-based and tolerance-based host defense
mechanisms, and proliferation-based and benevolence-based symbiont virulence mechanisms.
Both types of variation can be detected by measuring host fitness and detecting when (a)
different host genotypes vary in responses to infection by the same symbiont, or (b) the same
host genotype varies in responses to infection by different symbiont genotypes. (a) Hosts may
vary in **resistance** to the same symbiont. More resistant hosts (a – black line; lower average
symbiont density) will have overall higher fitness compared to less resistant hosts (b – gray line;
higher average symbiont density). But the fitness cost per increase in symbiont density is similar
for both hosts, as indicated by the similar slopes. On the other hand, hosts may vary in
tolerance to the same symbiont. More tolerant hosts (a – black line; shallower slope) may have
a similar average symbiont density to less tolerant hosts (b – gray line; steeper slope), but will
suffer a lower fitness cost per increase in symbiont density. (b) The same host may suffer
different fitness effects due to symbionts that vary in **proliferation**. The better proliferating
symbiont (b – gray line; higher average symbiont density) will impart higher fitness costs to the
host than the worse proliferating symbiont (a – black line; lower average symbiont density). But
the fitness cost to the host per increase in symbiont density is similar for both symbionts. The
same host may be infected with symbionts that vary in **benevolence**. More benevolent
symbionts (a – black line; shallower slope) will impart a lower fitness cost per increase in
symbiont density compared to more malevolent symbionts (b – gray line; steeper slope), though
both symbionts may reach similar average densities. (Framework adapted from Raberg et al
2007 and Wollein Waldetoft *et al.*, 2020.)

Figure 2. Representative samples from symbiont transmission experiment to demonstrate how
symbiont transmission during a social stage is detected. (a) Dyed cells (green population) are
gated on uninfected controls treated with Celltracker. (b) Infected cells with RFP-labeled
symbionts (pink population) are gated on pre-infected controls. (c) Newly infected cells (blue
population) are detected by applying logical gates to the experimental samples that mixed
uninfected cells and pre-infected cells.

640
641
Figure 3. Host fitness was negatively correlated with infection prevalence and differed in slope
but not average infection prevalence among symbiont strains after a single social stage.
Infection prevalence was estimated per host-symbiont-MOI sample as the percent of infected
spores that had RFP labeled symbionts in them. Host fitness was estimated by the percent of
spores produced per host-symbiont-MOI sample, relative to uninfected hosts prepared at the
same time as the infected host-symbiont combinations. There was no significant difference
between native (top row; lenient definition) vs. nonnative (bottom row) hosts.
649

Figure 4. The rate of horizontal transmission was positively correlated with pre-infection prevalence and also differed in slope and average infection prevalence among symbiont strains after a second social stage. Infection prevalence was estimated per host-symbiont-MOI sample as the percent of RPF+ infected spores of the pre-infected control. Horizontal transmission was estimated by the percent of RFP+ infected spores in the test sample that were also positive for a green membrane dye. These dyed cells were uninfected prior to the experiment, during which they were combined with pre-infected (and undyed) cells. There was no significant difference in symbiont transmission among native (top row; lenient definition) and nonnative (bottom row) hosts.

664
665
666   Figure 5. Infection prevalence was significantly different when amoebas were observed after
667   one (a) vs. two social stages (b) after infection. More malevolent strains (BbQS395 and
668   BbQS433) appear to proliferate less among amoeba hosts compared to more benevolent
669   strains (BbQS859) over time.
670
671

(a) Host defense mechanisms

Resistance | Tolerance | Both

Host fitness

Symbiont density

Host genotype
- a
- b

(b) Symbiont virulence mechanisms

Proliferation | Benevolence | Both

Host fitness

Symbiont density

Symbiont genotype
- a
- b

(a) Host fitness exp.

(b) Symbiont transmission exp.

Infection prevalence

Symbiont strain

Host type
○ native
□ nonnative