Can constraints on recombination reestablishment explain the evolution of heteromorphic sex chromosomes?

Authors: Thomas Lenormand[1]*, Denis Roze[2,3]

Affiliations:

[1] CEFE, Univ Montpellier, CNRS, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France.

[2] CNRS, IRL 3614, Roscoff, France.

[3] Sorbonne Université, Station Biologique de Roscoff, France.

*Correspondence to: thomas.lenormand@cefe.cnrs.fr

Short title: Sex chromosomes and recombination reestablishment

Abstract

Y and W chromosomes often stop recombining and degenerate. Most work on recombination suppression has focused on the mechanisms favoring recombination arrest in the short term. Yet, the long-term maintenance of recombination suppression is critical to evolve heteromorphic sex chromosomes. This long-term maintenance has been little investigated. In the long term, recombination suppression may be maintained for selective reasons (e.g. involving the emergence of nascent dosage compensation), or due to mechanistic constraints preventing the reestablishment of recombination, for instance when complex chromosomal rearrangements evolve on the Y. In this paper, we investigate these 'constraint' theories. We show that they face a series of theoretical difficulties: they are not robust to extremely low rates of recombination restoration; they would rather cause population extinction than Y degeneration; they are less efficient at producing a non-recombining and degenerate Y than scenarios adding a selective pressure against recombination, whatever the rate of recombination restoration. Finally, whether such very high constraint really exist is questionable. Very low rates of recombination reestablishment are sufficient to prevent Y degeneration, given the large fitness advantage to recover a non-degenerate Y or W for the heterogametic sex. The assumption of a lack of genetic variation to restore recombination seems also implausible given known mechanisms to restore a recombining pair of sex chromosomes.

Introduction

How could the degeneration of whole Y or W chromosomes evolve without being opposed by natural selection? Several theories have been put forward to explain this problem, often separating it into several subproblems: Why does recombination initially stop between sex chromosomes? Why does degeneration occur on a nonrecombining part of a genome? Why is recombination suppression maintained in the long term, once degeneration has already caused substantial degradation of the Y or W? Why isn't recombination eventually reestablished to limit maladaptation? When considering the suppression of recombination, it is useful to distinguish between the initial arrest and its long-term maintenance. We first briefly mention the different ideas that have been proposed for the initial arrest, which has attracted the most attention. We then present the different mechanisms that could lead to a long-term maintenance of recombination suppression, which has been much less investigated. We discuss the XY case throughout the paper, but all arguments apply to ZW systems as well.

The initial recombination arrest on sex chromosomes

Six main ideas have been proposed to explain shot term recombination arrest on sex chromosomes. Even if our focus on the long term maintenance of recombination suppression, it is useful to list these ideas to clarify the difference between the short and long term suppression of recombination.

The first proposes that the sex determination (SD) locus happened to arise in a region of the genome where recombination was already suppressed [1,2] or in a non-recombining genome. In achiasmate species, in particular, the heterogametic sex does not recombine (the so-called Haldane-Huxley rule [3,4]), so that, wherever the SD locus is located, the Y or W will not recombine. The direction of causality between the evolution of suppressed recombination on sex chromosomes and the evolution of achiasmy is however difficult to establish [4].

The second idea proposes that recombination is selected against to prevent the production of neuter individuals in species where sex is determined by a combination of a male-sterility and a female-sterility locus [5,6]. This explanation is applicable to species with such specific combinations of SD loci, and is likely to apply to the evolution of separate sexes from dioecy or genetic sex determination from environmental sex determination. However, it cannot account for later events of recombination suppression beyond the region of the genome containing the SD loci.

The third idea is the "sexually antagonistic selection" scenario [6–12], where suppression of recombination is selectively favored due to the occurrence of loci with sexually-antagonistic (SA) effects on sex chromosomes. In this case, a non-recombining Y benefits from a selective advantage, by permanently combining male determining and male-beneficial alleles. A variant of this scenario involves sex-differences in the intensity of selection (but not in the direction of selection as considered

in SA theory) against deleterious mutations [13]. There is plenty of evidence for SA variation [14,15],

making this hypothesis very plausible. However little evidence for this scenario has been obtained

despite intensive research [1,10,16], partly due to the difficulty of definitively establishing the role of

SA variation in driving recombination suppression.

The fourth scenario involves "lucky" inversions on the Y capturing the SD locus [17–19]. Inversions

capture at birth a sample of segregating deleterious variation. Some "lucky" inversions can have a

selective advantage because they initially capture a portion of the proto-Y that carries fewer or milder

deleterious mutations, compared to the population average. Their initial fitness advantage decays

through time but can be sufficient to allow them to fix [17,18]. Much of this decay is caused by a

deterministic return to the population average load [18,20], but selective interference also

contributes. This decay rate is larger for larger inversions [17,18], but inversions with a large initial

advantage are likely to be sampled among larger inversions [17]. These opposing effects determine

the distribution of fixed inversion sizes, with a mode biased toward smaller sizes compared to their

input size distribution [17,18]. Contrary to the fixation of 'lucky' inversions on autosomes [20], the

fixation of inversions capturing the SD locus on the proto-Y is expected to cause recombination

suppression on the inverted segment of the Y [16,21]. This process can involve any Y variant

suppressing recombination around the SD locus and closely linked to this region, not only inversions

(we use the term inversion only for simplifying the presentation). For instance, it could involve changes

in chromatin structure or the loss of recombination hotspots. It solely depends on the presence of

deleterious mutations and variation in recombination rates, and is thus expected to be widespread.

However, due to its recent description, it has not been investigated empirically.

The fifth mechanism involves recombination suppression near the pseudo-autosomal region boundary

due to the neutral accumulation of sequence divergence that decreases homology between the X and

Y, and suppresses recombination as a side effect [22]. The idea is that strict homology is required for

recombination to occur. However, while this effect has been documented in mitotic lineages [23–25,

and other references cited in 22], it is not clear whether small amounts of divergence prevent

recombination during meiosis. In particular, data from tetraploid rye and from crosses among strains

of *Arabidopsis thaliana* with varying levels of divergence suggest that heterozygosity may enhance

rather than inhibit crossovers [26–28].

The last mechanism proposes that XY recombination suppression is favored because it increases

heterozygosity around the SD locus in the heterogametic sex. Recent versions of this idea proposed

that tight linkage between the SD locus and overdominant mutations, or combinations of recessive

deleterious mutations (generating pseudo-overdominance) would be favored because these

mutations would then be more often present in the heterozygous state. In partially inbred populations,

linkage to the SD locus indeed increases heterozygosity, and this can favor recombination suppression

in the presence of overdominant mutations [29], while more work is needed for the case of recessive

deleterious mutations and random mating [18].

The long-term maintenance of recombination arrest on sex chromosomes

It has been long established that in the absence of recombination, deleterious mutations will tend to

accumulate in the Y due to selective interference [11,12,30–35], which should generate some

maladaptation, especially on males. Why then is recombination not reestablished when degeneration

becomes too strong? Three main ideas can be distinguished.

The first is that SA effects are sufficiently strong to maintain recombination arrest despite the

accumulation of deleterious mutations. This has nevertheless been questioned [36–38]. Indeed, the

deleterious effect of Y degeneration may eventually offset the selective advantage of linking male

determining and male-beneficial alleles. The restoration of recombination, if it is possible, may then

become favorable. However, new SA mutations may continue to appear and accumulate on sex

chromosomes, giving time for the population to evolve Y silencing / dosage compensation (DC) limiting

maladaptation caused by Y degeneration.

The second idea is based on regulatory evolution. Once recombination is suppressed, cis regulators of

gene expression may diverge between the X and Y, for genes located in the non-recombining portion

of the Y. This regulatory instability may lead to the evolution of early DC (through the joint evolution

of cis and trans acting factors), concomitant with Y early silencing and degeneration. In the model

proposed by [17], the emergence of this DC builds up pervasive SA regulatory effects, selectively

preventing the long-term reestablishment of recombination. Note that regulatory evolution may lead

to the accumulation of deleterious mutations and degeneration even in conditions where selective

interference is inoperative [39].

The third possibility, that we term the "constraint" scenario, involves cases where recombination

suppression is maintained in the long term despite the fact that it has become disadvantageous, due

to mechanistic constraints preventing the re-establishment of recombination. Most models for the

evolution of sex chromosomes ignore the possibility that recombination can be reestablished,

implicitly assuming that a constraint maintains recombination arrest on the long term. Few models

present a more detailed reasoning about this constraint. For instance in Jeffries et al.'s simulation

model of neutral arrest of recombination [22], crossovers are assumed to be fully suppressed once

sequence divergence becomes too high. The constraint emerges from the loss of homology. However,

the authors note that, in reality, rare recombination events could occasionally occur at high sequence

divergence. While Jeffries et al.'s model does not include deleterious mutations, degeneration would

generate strong selection to restore recombination and these rare events would be highly beneficial.

Another idea is that reestablishing recombination might be difficult once complex chromosomal

rearrangements have occurred on the Y. For instance, Jay et al.'s model [19] assumes that

recombination is suppressed as soon an inversion occurs, and that the occurrence of secondary

inversions (overlapping or occurring within a first one) prevents reversion of the first one. Hence,

recombination could only be reestablished in a region if all inversions are exactly reinversed,

irrespectively of the actual colinearity (or lack thereof) between the X and Y. This can allow lucky

inversions to persist in the long term, irrespective of the process of degeneration.

In this paper, we revisit this constraint scenario. We focus on the case where the short-term

recombination arrest is caused by lucky inversions. The initial arrest is not the factor of interest here,

so that we use the simplest model (the lucky inversion process only requires the occurrence of

deleterious mutations, and variation in recombination rates). We contend that explanations based on

the mechanistic constraint that recombination cannot be restored on the Y chromosome face several

theoretical challenges, rendering them unlikely, in our view, to account for the evolution of sex

chromosomes. Furthermore, we argue that mechanistic constraints on recombination restoration may

often not be sufficiently strong to lead to stable heteromorphic sex chromosomes.

Results

Modelling sex chromosome evolution

We analyze a model of sex chromosome evolution where recombination arrest is caused by lucky

inversions, and explore the constraint scenario by varying the rate of recombination restoration.

Specifically, we use the general model of sex chromosome evolution that we previously introduced to

explore the regulatory theory [19,20, which should be consulted for more details], but removing the

regulatory effects. This model considers a sex chromosome pair with a large number of genes (here

500) subject to deleterious mutations occurring in their coding sequences. Fitness is determined

multiplicatively across loci by the effect of deleterious mutations with a dominance coefficient equal

to 0.25, as observed on average for mildly deleterious mutations [40]. For simplicity, the SD locus is

located at one extremity of the chromosome. Recombination variation is modeled by introducing

mutations suppressing recombination in a region around the SD locus. These mutations can be thought

as being inversions (and we will refer to them as such) although other types of mechanisms are

possible, as already mentioned. For instance, the removal of recombination hotspots, or the addition

of a recombination suppressor sequence would work too. Specifically, as described in [17], we assume

that inversions occur on the Y at a rate U_{inv} per chromosome per generation (we use $U_{inv} = 10^{-5}$).

We only consider inversions that include the SD locus (or extend the non-recombining region of the Y

carrying the SD locus). Other inversions are not confined to males and can fix in the population, which

does not lead to recombination suppression on the Y (homozygous inversions recombine normally).

We denote the non-recombining fraction of the Y by z (between 0 and 1). This variable is also used to

measure the endpoint of each inversion on the chromosome. When $z = 0$, X and Y chromosomes

recombine freely, but otherwise X-Y recombination only occurs within the chromosomal segment [z,

1] (the SD locus being located at position 0). When $z = 1$, the X and Y do not recombine at all. When

a new inversion occurs, its size is drawn as a uniform fraction of the non-recombining part of the Y.

Specifically, on a Y where recombination is already stopped between 0 and z_i, after a new inversion i+1

the non-recombining region will extend to $z_{i+1} = z_i + (1 - z_i)u$, where u is a uniform deviate

between 0 and 1. To allow for the possibility that recombination may be reestablished, we assume that

reversions can also occur (at a rate U_{rev} per chromosome per generation), reverting the last inversion

on the non-recombining part of the Y. We investigate the dynamics of Y evolution in this model by

supposing that reversion rates are much smaller than inversion rates, with $U_{rev} =$

$10^{-6}, 10^{-7}, 10^{-8}, 10^{-9}$, i.e., from 1 to 4 orders of magnitude lower than the rate of occurrence of

inversions ($U_{inv} = 10^{-5}$). At the start of a simulation, each individual carries a pair of fully recombining

chromosomes, with the SD locus located at one extremity. Note that we do not perform a full

exploration of the parameter space here, but rather use the simulations to illustrate the different

points that are developed below concerning the effect of the reversion rate on sex chromosome

evolutionary dynamics. Note that parameters scaled by the population size are likely to be the

determinants of the evolutionary process, so that the scenario can be extended to different population

sizes by the appropriate rescaling of mutation rates, selection coefficients and times. The simulations

assume a constant population size (10^4 offspring individuals are drawn each generation, irrespective

of the average absolute fitness of male and female individuals in the previous generation). However,

the population was considered to be extinct when the average fitness of males became a thousand

times lower than the average fitness of females.

Very low rates of reversion can prevent long-term recombination suppression

To evaluate how strong the constraint on reversions should be for recombination suppression to be

maintained in the long term, we investigated cases where the rate of reversion was much lower than

the rate of inversion. We ran replicated simulations lasting four million generations. With our standard

parameters (Table 1), approximately 0.76 inversions fix per million generations. Typical outcomes are

illustrated on Fig 1 (taken from runs with $U_{rev} = 10^{-8}$). The majority of inversions that reach fixation

remain relatively short-lived (Fig 2A) and most of them occur one at a time (i.e. they are reverted

before a second one fixes). Before fixation, the marginal fitness of Y inversion decreases through time,

as they tend to accumulate deleterious mutations (returning to the equilibrium load and being exposed

to selective interference). After they fix, Y inversions continue degenerating because of selective

interference, which reduces male fitness and eventually offsets their initial fitness advantage. At this point, they become deleterious, and it is just a matter of time before a reversion occurs that would be selectively favored (Fig 2B shows the marginal fitness at birth and at the time of reversion for fixed inversions, for different reversion rates). This is the example illustrated on Fig 1A. In a few cases, another inversion fixes on top of the first one, before the first has reversed (this corresponds to the example illustrated on Fig 1B). When the rate of reversion is very low, several inversions may stack on the first one. In these cases, the lifespan of the first inversion is prolonged because it can be reversed only after the second one (or third one etc.) is reversed. This is due to the rather stringent hypothesis of our model that reversions can only occur on the last stratum present on the Y. This assumption protects the first inversion from reversion (until all other strata have reverted), and considerably reduces effective reversion rates (as it becomes zero for all but the last stratum on a Y). Considering reversions that could fully reestablish recombination on the Y at once would greatly reduce this (potentially unrealistic) effect. Whether or not stacking occurs, however, all inversions become reversed at some point (or the population goes extinct as we discuss below). There is no long-term maintenance of recombination suppression. Typically, Y chromosomes transiently carry one or two strata (Fig S1) for a relatively short time (Fig 2A). We can therefore conclude that the constraint scenario cannot explain the long-term maintenance of recombination suppression for rates of reversion up to 4 orders of magnitude lower than rates of inversion. This conclusion is very conservative, as our model of reversion does not allow the reversion of a first inversions if a second inversion occurs extending the non-recombining region (the first one can only be reversed after the second one is reversed). Without this constraint, it would be even more difficult to maintain recombination suppression

Very low rates of reversion can prevent degeneration

Many old Y chromosomes are largely non-recombining and degenerate, mutations having accumulated up to the point where genes have become nonfunctional or have been lost. In our model, degeneration corresponds to the situation where a gene has accumulated deleterious mutations up to a maximum fitness effect of s_{max}, (corresponding to the fitness drop caused by the loss of function of a gene, here set to 0.3). When reversion rates are low, some inversions can fix and persist in the population for some time, especially when secondary inversions also occur prolonging the lifespan of the first one. Usually, this does not correspond to a large fraction of the Y, but on rare occasions, this can be significant (in particular, when several fixed inversions are stacked, which reduces the rate of reversion of all but the last one of them, as explained above). For example, in the example illustrated on Fig 1B, about a third of the Y stopped recombining for nearly a million generations. Even in these extreme cases, however, degeneration remains moderate: even with extremely low rates of reversion

(10^{-9}), almost no gene accumulates deleterious mutations up to s_{max}. Yet, we use a relatively high rate

of deleterious mutation per gene (U_g = 2 x 10^{-4}), a distribution of fitness effects of mutations with a

relatively high mean (s_{mean} = 0.05) and a large proportion of small effect mutations (the distribution of

effects is exponential). In the vast majority of cases, almost no loss-of-function is detectable (Fig S2A).

The reason for this lack of loss of function is that many weakly deleterious mutations accumulate in all

genes present on a fixed inversion. Collectively, their impact on the marginal fitness of the inversion

starts to be strong long before any gene in particular becomes fully nonfunctional. Hence, inversions

become selectively disfavored (and therefore selectively eliminated as soon as a reversion arises), long

before they exhibit any gene loss. We can therefore conclude that the constraint scenario cannot

explain strong degeneration for rates of reversion up to 4 orders of magnitude lower than rates of

inversion. Degeneration may occur under very low rates of reversion if carrying nonfunctional genes

on the Y would only cause a very small fitness cost for males (a situation that would be represented by

setting s_{max} to a small value in our model). This situation seems unlikely in the absence of a mechanism

silencing impaired genes on the Y, however, while letting gene expression evolve would lead to the

regulatory scenario, under which recombination arrest can be maintained even in the absence of any

constraint on recombination restoration [17].

The constraint scenario is more likely to lead to extinction than Y degeneration

It may be argued that reversion rates are even smaller than the ones we considered, making the

constraint scenario a possibility, at least theoretically. There is a strong argument against this

possibility. When an inversion fixes and starts accumulating deleterious mutations, it depresses male

fitness. Initially, a lucky inversion is selectively favored because it captures a fraction of the Y carrying

fewer deleterious mutations (or deleterious mutations with smaller effects) compared to the average

Y population. Deleterious mutations start accumulating within the inversion due to the fact that the

inversion tends to return towards the average mutation load [18], and to selective interference. As

explained above, a first threshold is reached when the accumulation of deleterious mutations

depresses the marginal fitness of this portion of the Y below the average marginal fitness of the

homologous portion of the X in the population. Reversions then become selectively favored, and it is

just a matter of time before one occurs and eliminates the inversion. If the reversion rate is extremely

low, this can indeed take a long time. However, a second threshold will be reached relatively quickly,

corresponding to the non-viability or sterility of males, and hence to the extinction of the

population/species. This cannot happen in our model as we assume a constant population size, i.e.,

the absolute number of individuals in the population does not depend on the fitness of individuals

(soft selection), but the simulations show a crash in male fitness relative to female fitness when

inversions are maintained for a sufficiently long time (Fig S2B). It is not easy to determine the male

fitness threshold that would lead to population extinction in nature. We used a relatively conservative

threshold equal to 10^{-3} (meaning that the fitness of males is three orders of magnitude lower than the

fitness of females). Such a distortion of male vs. female fitness would be particularly conspicuous in

natural conditions. In the few cases where some degeneration occurs (e.g. when a large inversion

unfortunately fixes), the population reaches this limit quickly and becomes extinct. Reversions can

rescue the population and prevent extinction, but if they are too rare, they do not occur quickly enough

to prevent it. For instance, with very low rates of reversions (U_{rev} = 10^{-9}), this threshold was often

reached in our simulations (extinction occurred in 70% of cases, 14 replicates out of 20 within the first

4 million generations of evolution). This estimate is conservative, since considering that the number of

males in the population may be much lower than assumed under our soft selection regime would lead

to an even faster accumulation of deleterious mutations on the Y (due to stronger drift). Note that our

model does not include back mutations, which would eventually stop the decline in fitness caused by

deleterious mutation accumulation. However, previous work has shown that in the absence of

recombination, mean fitness reaches very low values even when back mutations do occur, unless the

mean fitness effect of deleterious mutations is extremely weak [41,42]. Hence, a theory based on

constraints alone cannot explain both degeneration and the persistence of populations/species. For

degeneration and persistence to occur, Y silencing and DC must also evolve, which can be a powerful

selective mechanism that stabilizes recombination arrest. We show in supplementary material that,

under the same parameters, Y degeneration and DC indeed occur relatively fast when regulators are

free to evolve. These results also point out that heteromorphic sex chromosomes may readily evolve

under some scenarios. Even when reversion rates are extremely low, models involving a selective

pressure maintaining recombination arrest predict complete and permanent recombination

suppression and degeneration (without a major drop in male fitness), while models solely based on

constraints only produce partial and transient degeneration and recombination suppression.

Discussion

The arguments and results presented in this article imply that, in the absence of regulatory evolution,

the decreased fitness of the heterogametic sex due to mutation accumulation on the Y should lead to

two possible outcomes: (i) the restoration of recombination if reversions can occur, even at very low

rates, or (ii) the extinction of populations if constraints on reversions are sufficiently strong. In the

following, we discuss the constraint theory in the light of those results and possible mechanisms of

recombination reestablishment, before indicating future avenues for theoretical and empirical

research concerning the initial steps of recombination suppression, the mechanisms responsible for

the long-term maintenance of recombination suppression and the extension of these theories.

What is the degree of constraint on recombination reestablishment?

The possibility of recombination restoration should depend on the mechanism of recombination arrest. Recent theoretical work has emphasized the possible role of inversions in suppressing recombination on sex chromosomes [17–19,43], although these models also apply to other mechanisms of stepwise recombination arrest. Inversions do indeed occur frequently within populations and may be caused by ectopic recombination between repeated sequences [44–47]; they are often observed on sex chromosomes [10,48–50], although some of these inversions may have occurred after recombination arrest. Inversions are well known to reduce recombination rates in heterokaryotes. This reduction is not necessarily because inversions inhibit homologous pairing, but because an odd number of crossovers within the inversion loop leads to the production of unbalanced chromosomes. It is important to note that recombination may still occur when the number of crossovers falling within the inversion is even (especially in the case of relatively large inversions where crossover interference is weaker [51–54]), while gene conversion events may also allow for genetic exchanges between inverted and non-inverted segments [54,55].

In a previous model of inversion dynamics on sex chromosomes, Jay et al. [19] consider that recombination restoration is possible only when an inversion with exactly the same breakpoints ("reinversion") restores the exact original gene order before another inversion overlaps or occurs within the first one. If a second nested or overlapping inversion occurs, it can also be reversed, but only before a third nested/overlapping inversion occurs and so on. In this model, the chance of reinversion becomes vanishing low as the number of breakpoints increases. Indeed, with N breakpoints on the chromosome and with a first inversion spanning k breakpoints, the chance of reinversion is $\sim 1/N^2$, while the chance of a second overlapping / nested inversion is $\sim k/N$. In the results we presented, this level of constraint is achieved for the very low rates of reversions. For instance, with N = 100 breakpoints and a first inversion spanning k = 10 breakpoints, reinversions are 3 orders of magnitude less likely than inversions. An interesting feature of this model is that it mechanistically represents inversion and reinversions. It also captures the idea that reestablishing the exact gene order with random inversions becomes increasingly difficult as they accumulate. This phenomenon may indeed occur and constrain the reestablishment of recombination.

However, several processes could largely limit the constraint imposed by the accumulation of overlapping and nested inversions. First, recombination may occasionally occur even if gene collinearity is not exact, as shown with ectopic recombination [56–58], especially after a second inversion restoring the original direction on a portion of the chromosome. Fig S4 and S5 illustrates such possibilities for nested or overlapping inversions. With imperfect collinearity, recombined chromosomes have low fitness in general, but here, with a partially degenerated Y, the question is

more subtle, as the loss of some (already) degenerated genes may be compensated by the acquisition

of non-degenerated portion of the X. What matters is the relative fitness of recombined Y compared

to the current (partially degenerated) low fitness Y. Hence, many favorable cases of imperfect

recombination could occur and be favored, which could largely increase rates of recombination

reestablishment.

Another possibility is that recombining sex chromosomes may be reestablished by moving the SD locus

out of the non-recombining region. This mechanism can always occur, even when complex

rearrangements have taken place on the Y. This may occur for instance by recruiting a new master

switch gene for sex-determination, or following a duplication of the existing SD locus into a new

location. Gene duplications are frequent events in eukaryotic genomes [59]. Rates in the range 10^{-5} –

10^{-7} per gene per generation have been reported in animals [60–62], i.e. a much higher rate than the

rate of reversions that we investigated. If the SD locus moves to another (recombining) location on the

sex chromosome, a new Y recently re-derived from the X (and fully recombining) can evolve easily.

Examples of such events are often reported [63–65], indicating that they may be common. The SD

locus may also move to another autosomal pair, leading to the evolution of a new pair of fully

recombining sex chromosomes. Examples of such turnovers abound [66], indicating that they are also

common. Models have also shown that the accumulation of deleterious mutations would select for

these turnover events [37,38]. In some species, another possibility for restoring recombination on the

Y involves environmental sex reversal when recombination is sex-dependent [36].

Often, the idea that recombination restoration is strongly constrained stems from the idea that

'reinversions' are not observed. While the rate of reinversion is probably low, this view is not entirely

accurate. Reinversions have been occasionally shown to occur in laboratory populations of Drosophila

[67] at rates 10^{-3} – 10^{-4}: while most of them were caused by X-ray irradiation [68,69], one occurred

spontaneously in a stock population [70,71] and it was proposed that reinversion may be favored by

the physical proximity of the breakpoints during loop formation [69]. More empirical work on this issue

is needed in order to assess to what extent such a process may occur (and at which rate) despite the

repression of crossing-over in the vicinity of inversion breakpoints. However, the observation of an

accumulation of nested and overlapping inversions alone is not an indication that the absence of

recombination restoration was due to a constraint or to a selection pressure against recombination.

Chromosomal rearrangements can secondarily accumulate in a non-recombining region [1].

Hence, while the accumulation of complex rearrangements is certainly a way to suppress

recombination, the maintenance of recombination arrest on the Y by a constraint alone requires a very

high level of constraint, given the maladaptation caused by mutation accumulation on the Y. This level

of constraint is in fact a rather strong assumption. It requires that reinversions are very rare, that rare

recombination event involving imperfectly colinear chromosomes do not occur, and that the SD locus

cannot move outside the non-recombination region.

Conclusion and perspectives

After recombination suppression, the accumulation of deleterious mutations on the Y then generates

a selective pressure to restore recombination and purge those alleles. This selective pressure becomes

stronger as male fitness declines, soon making recombination restoration events highly favorable. This

explains that even with extremely low rates of recombination reestablishment, recombination

suppression cannot persist in the long-term when only deleterious mutations are considered. In this

case, recombination restoration events are rescuing the population from extinction, and even if they

are rare, play a disproportionate role on the outcome. This issue is even more acute in models where

recombination arrest is caused by neutral divergence [22] as recombination suppression occurs

gradually, rather than quickly as in the lucky inversion scenario. This is the central theoretical argument

against theories of sex chromosome evolution solely based on mechanistic constraints [19,22]: the

accumulation of deleterious alleles caused by recombination arrest should eventually lead to

population extinction or to the re-establishment of recombination (via reinversion(s) or sex

chromosome turnover), rather than the long-term maintenance of degenerate Y or W chromosomes.

Several issues remain to be investigated in more detail. The different processes possibly involved in

the evolution of Y chromosomes need to be better integrated. In particular, the conditions allowing

the long-term maintenance of recombination suppression under the SA theory should be investigated

further [following 37,38] and combined with models of regulatory evolution. A better integration of

the different mechanisms of regulatory variation may also be useful, such as mechanisms based on

imprinting [72] or based on reallocation of transcription factors to X-linked genes [11]. While the

widespread occurrence of DC [72,73] indicates that it is needed at least for some genes, it would be

interesting to introduce heterogeneity among genes in selection on dosage, and varying the genetic

architecture of DC (local vs global). Several cases of interest should be investigated further, notably

cases involving non-random mating [18,29,74] or UV and mating-type chromosomes. Analytical

models are needed to generalize the recent findings that have mostly been explored by simulation.

Finally, from an empirical perspective, more data are needed on patterns of recombination

suppression, degeneration and mechanisms of early DC evolution in young sex chromosome systems.

Overall, the level of constraint on recombination restoration may not be the key parameter to

understand why heteromorphic or homomorphic sex chromosomes occur in a given species. The ease

to evolve dosage compensation is likely to be the main driver: if regulatory evolution is difficult, sex

chromosomes will remain homomorphic (recombination will be restored in some way or the species

will go extinct). If regulatory evolution and DC can evolve relatively easily, stable heteromorphic sex
chromosomes may persist on the long term [17].

Acknowledgments

404 We thank A. Muyle for useful comments.

Funding

406 This work is supported by the ANR Genasex and CisTransEvol.

Code availability

408 The simulation code is available at:

409 T. Lenormand, D. Roze, Zenodo (2021), doi:10.5281/zenodo.5504423.

References

1. Charlesworth D. When and how do sex-linked regions become sex chromosomes? Evolution.
2021;75: 569–581. doi:10.1111/evo.14196

2. Rifkin JL, Beaudry FEG, Humphries Z, Choudhury BI, Barrett SCH, Wright SI. Widespread
recombination suppression facilitates plant sex chromosome evolution. Molecular Biology and
Evolution. 2021;38: 1018–1030. doi:10.1093/molbev/msaa271

3. Bell G. The masterpiece of nature: the evolution and genetics of sexuality. Berkeley: University
of California Press; 1982.

4. Lenormand T, Dutheil J. Recombination difference between sexes: a role for haploid selection.
Hurst LD, editor. PLoS Biology. 2005;3: 396–403. doi:10.1371/journal.pbio.0030063

5. Charlesworth B, Charlesworth D. A model for the evolution of dioecy and gynodioecy. The
American Naturalist. 1978;112: 975–997. doi:10.1086/283342

6. Bull JJ. Evolution of sex determining mechanisms. Menlo Park, CA: Benjamin Cummings; 1983.

7. Rice WR. Sex chromosomes and the evolution of sexual dimorphism. Evolution. 1984;38: 735–
742.

8. Rice W. The accumulation of sexually antagonistic genes as a selective agent promoting the
evolution of reduced recombination between primitive sex chromosomes. Evolution. 1987;41:
911–914.

9. Charlesworth B. The evolution of sex-chromosomes. Science. 1991;251: 1030–1033.

10. Charlesworth D. Evolution of recombination rates between sex chromosomes. Phil Trans R Soc
B. 2017;372: 20160456. doi:10.1098/rstb.2016.0456

11. Charlesworth B. Model for evolution of Y chromosomes and dosage compensation. Proceedings
of the National Academy of Sciences of the United States of America. 1978;75: 5618–5622.

12. Bachtrog D. Y-chromosome evolution: Emerging insights into processes of Y-chromosome
degeneration. Nature Reviews Genetics. 2013;14: 113–124. doi:10.1038/nrg3366

13.   Lenormand T. The evolution of sex dimorphism in recombination. Genetics. 2003;163: 811–822.

14.   Rowe L, Chenoweth SF, Agrawal AF. The genomics of sexual conflict. American Naturalist.
2018;192: 274–286. doi:10.1086/698198

15.   Bonduriansky R, Chenoweth SF. Intralocus sexual conflict. Trends in Ecology and Evolution.
2009;24: 280–288. doi:10.1016/j.tree.2008.12.005

16.   Ironside JEE. No amicable divorce? Challenging the notion that sexual antagonism drives sex
chromosome evolution. BioEssays. 2010;32: 718–726. doi:10.1002/bies.200900124

17.   Lenormand T, Roze D. Y recombination arrest and degeneration in the absence of sexual
dimorphism. Science. 2022;375: 663–666. doi:10.1101/2021.05.18.444606

18.   Olito C, Ponnikas S, Hansson B, Abbott JK. Consequences of partially recessive deleterious genetic
variation for the evolution of inversions suppressing recombination between sex chromosomes.
Evolution. 2022;76: 1320–1330. doi:10.1111/evo.14496

19.   Jay P, Tezenas E, Véber A, Giraud T. Sheltering of deleterious mutations explains the stepwise
extension of recombination suppression on sex chromosomes and other supergenes. PLOS
Biology. 2022;20: e3001698. doi:10.1371/journal.pbio.3001698

20.   Nei M, Kojima KI, Schaffer HE. Frequency changes of new inversions in populations under
mutation-selection equilibria. Genetics. 1967;57: 741–750.

21.   Bowen ST. The genetics of *Artemia salina*. V. Crosssing over between the X and Y chromosomes.
Genetics. 1965;52: 695–710.

22.   Jeffries DL, Gerchen JF, Scharmann M, Pannell JR. A neutral model for the loss of recombination
on sex chromosomes. Philosophical Transactions of the Royal Society B: Biological Sciences.
2021;376. doi:10.1098/rstb.2020.0096

23.   Datta A, Hendrix M, Lipsitch M, Jinks-Robertson S. Dual roles for DNA sequence identity and the
mismatch repair system in the regulation of mitotic crossing-over in yeast. Proceedings of the
National Academy of Sciences. 1997;94: 9757–9762. doi:10.1073/pnas.94.18.9757

24.   Li L, Jean M, Belzile F. The impact of sequence divergence and DNA mismatch repair on
homeologous recombination in *Arabidopsis*. The Plant Journal. 2006;45: 908–916.
doi:10.1111/j.1365-313X.2006.02657.x

25.   Do AT, Brooks JT, Le Neveu MK, LaRocque JR. Double-strand break repair assays determine
pathway choice and structure of gene conversion events in *Drosophila melanogaster*. G3
Genes|Genomes|Genetics. 2014;4: 425–432. doi:10.1534/g3.113.010074

26.   Benavente E, Sybenga J. The relation between pairing preference and chiasma frequency in
tetrasomics of rye. Genome. 2004;47: 122–133. doi:10.1139/g03-134

27.   Blackwell AR, Dluzewska J, Szymanska-Lejman M, Desjardins S, Tock AJ, Kbiri N, et al. MSH 2
shapes the meiotic crossover landscape in relation to interhomolog polymorphism in
Arabidopsis. EMBO J. 2020;39. doi:10.15252/embj.2020104858

28. Ziolkowski PA, Berchowitz LE, Lambing C, Yelina NE, Zhao X, Kelly KA, et al. Juxtaposition of heterozygous and homozygous regions causes reciprocal crossover remodelling via interference during *Arabidopsis* meiosis. eLife. 2015;4: e03708. doi:10.7554/eLife.03708

29. Charlesworth B, Wall JD. Inbreeding, heterozygote advantage and the evolution of neo–X and neo–Y sex chromosomes. Proc R Soc Lond B. 1999;266: 51–56. doi:10.1098/rspb.1999.0603

30. Rice WR. Genetic hitchhiking and the evolution of reduced genetic activity of the Y sex chromosome. Genetics. 1987;116: 161–167. doi:10.1093/genetics/116.1.161

31. Peck JR. A ruby in the rubbish: beneficial mutations, deleterious mutations and the evolution of sex. Genetics. 1994;137: 597–606.

32. Charlesworth B. The evolution of chromosomal sex determination and dosage compensation. Current Biology. 1996;6: 149–162. doi:http://dx.doi.org/10.1016/S0960-9822(02)00448-7

33. Gordo I, Charlesworth B. The speed of Muller's ratchet with background selection, and the degeneration of Y chromosomes. Genetical research. 2001;78: 149–161.

34. Charlesworth B, Charlesworth D. Rapid fixation of deleterious alleles can be caused by Muller's ratchet. Genetical Research. 1997;70: 63–73. doi:10.1017/S0016672397002899

35. Engelstädter J. Muller's ratchet and the degeneration of Y chromosomes: A simulation study. Genetics. 2008;180: 957–967. doi:10.1534/genetics.108.092379

36. Grossen C, Neuenschwander S, Perrin N. The evolution of XY recombination: Sexually antagonistic selection versus deleterious mutation load. Evolution. 2012;66: 3155–3166. doi:10.1111/j.1558-5646.2012.01661.x

37. Blaser O, Grossen C, Neuenschwander S, Perrin N. Sex-chromosome turnovers induced by deleterious mutation load. Evolution. 2013;67: 635–645. doi:10.1111/j.1558-5646.2012.01810.x

38. Blaser O, Neuenschwander S, Perrin N. Sex-chromosome turnovers: The hot-potato model. The American Naturalist. 2014;183: 140–146. doi:10.1086/674026

39. Lenormand T, Fyon F, Sun E, Roze D. Sex chromosome degeneration by regulatory evolution. Current Biology. 2020;30: 3001-3006.e5. doi:10.1016/j.cub.2020.05.052

40. Manna F, Martin G, Lenormand T. Fitness landscape: an alternative theory for the dominance of mutations. Genetics. 2011;189: 923–937.

41. Charlesworth B, Betancourt AJ, Kaiser VB, Gordo I. Genetic recombination and molecular evolution. Cold Spring Harbor Symposia on Quantitative Biology. 2009;74: 177–186. doi:10.1101/sqb.2009.74.015

42. Kaiser VB, Charlesworth B. Muller's ratchet and the degeneration of the *Drosophila miranda* neo-*Y* chromosome. Genetics. 2010;185: 339–348. doi:10.1534/genetics.109.112789

43. Olito C, Abbott JK. The evolution of suppressed recombination between sex chromosomes and the lengths of evolutionary strata. Evolution. 2023; qpad023. doi:10.1093/evolut/qpad023

44. Porubsky D, Höps W, Ashraf H, Hsieh P, Rodriguez-Martin B, Yilmaz F, et al. Recurrent inversion
polymorphisms in humans associate with genetic instability and genomic disorders. Cell.
2022;185: 1986-2005.e26. doi:10.1016/j.cell.2022.04.017

45. Cáceres M, Puig M, Ruiz A. Molecular characterization of two natural hotspots in the *Drosophila*
buzzatii genome induced by transposon insertions. Genome Res. 2001;11: 1353–1364.
doi:10.1101/gr.174001

46. Feuk L, MacDonald JR, Tang T, Carson AR, Li M, Rao G, et al. Discovery of human inversion
polymorphisms by comparative analysis of human and chimpanzee DNA sequence assemblies.
PLoS Genet. 2005;1: e56. doi:10.1371/journal.pgen.0010056

47. Dobzhansky TH. Genetics of the evolutionary process. New York: Columbia University Press;
1970.

48. Lemaitre C, Braga MDV, Gautier C, Sagot M-F, Tannier E, Marais GAB. Footprints of inversions at
present and past pseudoautosomal boundaries in human sex chromosomes. Genome Biology
and Evolution. 2009;1: 56–66. doi:10.1093/gbe/evp006

49. Zhang W, Wai CM, Ming R, Yu Q, Jiang J. Integration of genetic and cytological maps and
development of a pachytene chromosome-based karyotype in papaya. Tropical Plant Biol.
2010;3: 166–170. doi:10.1007/s12042-010-9053-2

50. Peichel CL, McCann SR, Ross JA, Naftaly AFS, Urton JR, Cech JN, et al. Assembly of the threespine
stickleback Y chromosome reveals convergent signatures of sex chromosome evolution. Genome
Biology. 2020;21: 1–31. doi:10.1186/s13059-020-02097-x

51. Stevison LS, Hoehn KB, Noor MAF. Effects of inversions on within- and between-species
recombination and divergence. Genome Biology and Evolution. 2011;3: 830–841.
doi:10.1093/gbe/evr081

52. Levine RP. Crossing over and inversions in coadapted systems. The American Naturalist. 1956
[cited 6 Feb 2023]. doi:10.1086/281905

53. Ishii T, Nagai A, Hirono T, Nakamura K. Hibernation of mosquitoes in rock caves on Miyato island.
Sci Rep Tôhoku Univ Ser IV. 1964;30: 159–165.

54. Korunes KL, Noor MAF. Pervasive gene conversion in chromosomal inversion heterozygotes. Mol
Ecol. 2019;28: 1302–1315. doi:10.1111/mec.14921

55. Navarro A, Betrán E, Barbadilla A, Ruiz A. Recombination and gene flux caused by gene
conversion and crossing over in inversion heterokaryotypes. Genetics. 1997;146: 695–709.
doi:10.1093/genetics/146.2.695

56. Kupiec M, Petes TD. Allelic and ectopic recombination between Ty elements in yeast. Genetics.
1988;119: 549–559. doi:10.1093/genetics/119.3.549

57. Montgomery EA, Huang SM, Langley CH, Judd BH. Chromosome rearrangement by ectopic
recombination in <i>Drosophila melanogaster<i>: genome structure and evolution. Genetics.
1991;129: 1085–1098. doi:10.1093/genetics/129.4.1085

58.   Lambert S, Mizuno K, Blaisonneau J, Martineau S, Chanet R, Fréon K, et al. Homologous
544   recombination restarts blocked replication forks at the expense of genome rearrangements by
545   template exchange. Molecular Cell. 2010;39: 346–359. doi:10.1016/j.molcel.2010.07.015

546   59.   Li W-H, Gu Z, Cavalcanti ARO, Nekrutenko A. Detection of gene duplications and block
547   duplications in eukaryotic genomes. J Struct Funct Genomics. 2003;3: 27–34.

548   60.   Watanabe Y, Takahashi A, Itoh M, Takano-Shimizu T. Molecular spectrum of spontaneous de
549   novo mutations in male and female germline cells of *Drosophila melanogaster*. Genetics.
550   2009;181: 1035–1043. doi:10.1534/genetics.108.093385

551   61.   Turner DJ, Miretti M, Rajan D, Fiegler H, Carter NP, Blayney ML, et al. The rates of de novo meiotic
552   deletions and duplications causing several genomic disorders in the male germline. Nat Genet.
553   2008;40: 90–95. doi:10.1038/ng.2007.40

554   62.   Lipinski KJ, Farslow JC, Fitzpatrick KA, Lynch M, Katju V, Bergthorsson U. High spontaneous rate
555   of gene duplication in *Caenorhabditis elegans*. Curr Biol. 2011;21: 306–310.
556   doi:10.1016/j.cub.2011.01.026

557   63.   Charlesworth D, Bergero R, Graham C, Gardner J, Keegan K. How did the guppy Y chromosome
558   evolve? PLOS Genetics. 2021;17: e1009704. doi:10.1371/journal.pgen.1009704

559   64.   Sharma A, Heinze SD, Wu Y, Kohlbrenner T, Morilla I, Brunner C, et al. Male sex in houseflies is
560   determined by Mdmd, a paralog of the generic splice factor gene CWC22. Science. 2017;356:
561   642–645. doi:10.1126/science.aam5498

562   65.   Meisel RP, Olafson PU, Adhikari K, Guerrero FD, Konganti K, Benoit JB. Sex chromosome evolution
563   in Muscid flies. G3 Genes|Genomes|Genetics. 2020;10: 1341–1352. doi:10.1534/g3.119.400923

564   66.   Furman BLS, Metzger DCH, Darolti I, Wright AE, Sandkam BA, Almeida P, et al. Sex chromosome
565   evolution: So many exceptions to the rules. Genome Biology and Evolution. 2020;12: 750–763.
566   doi:10.1093/gbe/evaa081

567   67.   Krimbas CB, Powell JR. Drosophila inversion polymorphism. CRC Press; 1992.

568   68.   Hinton T. A correlation of phenotypic changes and chromosomal rearrangements at the two ends
569   of an inversion. Genetics. 1950;35: 188–205. doi:10.1093/genetics/35.2.188

570   69.   Novitski E. The regular reinversion of the roughest3 inversion. Genetics. 1961;46: 711–717.
571   doi:10.1093/genetics/46.7.711

572   70.   Emmens CW. Salivary gland cytology of Roughest3 inversion and reinversion, and Roughest2.
573   Journ of Genetics. 1937;34: 191–202. doi:10.1007/BF02982262

574   71.   Grüneberg H. The position effect proved by a spontaneous reinversion of the X-chromosome in
575   *Drosophila melanogaster*. Journ of Genetics. 1937;34: 169–189. doi:10.1007/BF02982261

576   72.   Muyle A, Marais G, Bačovský V, Hobza R, Lenormand T. Dosage compensation evolution in plants:
577   theories, controversies and mechanisms. Philosophical Transactions of the Royal Society B.
578   2022;377: 20210222.

579   73.   Gu L, Walters JR. Evolution of sex chromosome dosage compensation in animals: A beautiful
580   theory, undermined by facts and bedeviled by details. Genome Biology and Evolution. 2017;9:
581   2461–2476. doi:10.1093/gbe/evx154

582   74.   Tezenas E, Giraud T, Véber A, Billiard S. The fate of recessive deleterious or overdominant
583   mutations near mating-type loci under partial selfing. Peer Community Journal. 2023;3: e14.
584   doi:10.24072/pcjournal.238

Fig 1. Examples of inversion-reversion dynamics. Each example is illustrated with three panels. The top panel shows the average non recombining fraction of the Y in the population ($\overline{z}$) through time (x axis in million generations). Colored lines correspond to fixed inversions (i.e., all inversions reaching a frequency of 1 during the simulation); different fixed inversions have a different color. The colored line extends between the time of occurrence of the inversion and the time when it becomes extinct. The middle panel shows the (per gene) average cumulative fitness effect of deleterious mutations on these inversions through time (same color code as in top panel). The bottom panel shows the ratio of the average fitness of males / females in the population through time.

Fig 2. Characteristics of inversions under the constraint scenario. Panel A indicates the lifespan of fixed inversions (taken on 20 replicates) for different reversion rates (on y-axis). Time was cut off at the end of the simulation (after 4 million generations) or if the population became extinct (by reaching a male/female fitness ratio < 0.001). Panel B shows the marginal fitness at birth of inversions (x-axis) versus marginal fitness at last recorded time (y-axis, log scale). The latter most often corresponds to a reversion, but in a few cases, it corresponds to the end of the simulation, or population extinction. Color codes correspond to different reversion rates as indicated in the legend.

Fig 3. Overall evolution of the Y chromosome under the constraint scenario. The x-axis (in log scale)

gives the fraction of the Y that is non recombining (averaged over all males in the population) after

two million generations. The y-axis (in log scale) gives the fraction of knock-out genes on the Y

(averaged over all males in the population; a knock out is defined as a gene having accumulated

deleterious mutations up to s_{max} = 0.3), after two million generations. Each dot represents a replicated

population. Color codes indicates different rates of reversion (U_{rev}). The rate of inversion is 10^{-5} in all

cases. In a few cases (with U_{rev} = 10^{-9}, filled yellow disks) the population became extinct before 2 million

generations. In these cases (marked with a skull), the x and y axes values are taken at the time of

extinction.

Parameters	Values
Population size	10000
Number of genes	500
Average effect of deleterious mutations (the distribution is exponential)	0.05
Dominance coefficient of deleterious mutations	0.25
Maximum fitness drop caused by mutations in a gene, corresponding to a knock-out	0.3
Mutation rate of genes	0.0002
Recombination rate between consecutive genes (R_g)	0.0005
Rate of inversion mutations	0.00001
Rate of reversion mutations (10 to 10^4 lower than rate of inversions)	variable

621   **Table 1. Parameter values used in simulations.**

Supplementary material

1. Comparing the constraint scenario to a scenario including a selective pressure against recombination

It is not because reversion rates are low (representing strong constraints on recombination restoration) that explanations of long-term recombination suppression solely based on constraints are likely to hold. Quantitatively, the question is rather, for given reversion rates, to determine the most likely scenario for observing Y chromosomes with non-recombining and degenerate strata within a realistic timeframe. To illustrate this point, we simulated the evolution of Y chromosomes under the same low reversion rates used above, but allowing regulators to evolve using the model described in [17], which should be consulted for more details. We use the same simulations than in the main text, but we consider that the expression of each gene is controlled by a cis-regulator and two trans-regulators (one only expressed in males and the other only expressed in females). These regulators determine quantitative traits that control the total and allele-specific level of expression of each gene. Total gene expression is supposed to be under stabilizing selection for all genes. Fitness is determined multiplicatively across loci by the effect of deleterious mutations (whose dominance depends on the relative strength of cis-regulators in heterozygotes, with a baseline dominance in the absence of cis-regulatory variation equal to 0.25) and by the departure from optimal expression at each locus. Cis and trans regulators mutate at a fixed rate per generation (with Gaussian variation in trait values). Table S1 indicates the additional parameters and their value for the simulations with evolution of regulators.

In this case, we observe much faster Y recombination suppression and degeneration than in the absence of regulatory evolution, for all reversion rates investigated. This is probably also true for scenarios involving SA loci, which generate a selective pressure against recombination accelerating the process. Low reversion rates are favorable to any theory on the maintenance of recombination arrest, not only those solely based on constraints. Indeed, low reversion rates give more time for other SA loci to accumulate, or nascent DC to emerge, and more time for degeneration to happen. In our model of regulatory evolution, and with very low reversion rates, almost any fixed inversion has time to develop nascent DC, generating sexually-antagonistic regulatory effects that effectively disfavor recombination. Fig S1 shows that many more strata accumulate on the Y in this case than in the absence of regulatory evolution, while Fig S2C shows that most of these strata are long lived and fully degenerated (while almost none is degenerated for the same parameter values in the absence of regulatory evolution, Fig S2A). Fig S2D shows that this degeneration is not associated to a large drop in male to female fitness ratio, while this drop is considerable in the absence of regulatory evolution. When regulators evolve, reversions occur but are not selectively favored. Indeed, the reestablishment of X-Y recombination on a given stratum causes X cis-regulators to move to the Y, creating recombinant

(low fitness) Y that cause a departure from optimal gene expression in males. This is a case of an
evolved selective constraint. The ultimate cause of long-term recombination suppression is not the
absence of genetic variation for reestablishing recombination once an inversion has fixed (mechanistic
constraint), but that it is selectively unfavorable to re-establish it. Overall, for the parameter values
considered here, we see that after 2 million generations a large fraction of the Y has stopped
recombining and degenerated in the presence of regulatory evolution with the lowest reversion rates
(for high reversion rates the process is still ongoing with only few small stabilized and degenerated
strata, Fig S3). Nothing nearly comparable occurs without regulatory evolution, where at most, in a
few cases with extremely low rates of reversion, some recombination suppression evolves and drives
the population to extinction without leading to significant Y degeneration (Fig S3). We conclude that
even if reversion rates were extremely low, so that a scenario solely based on constraints could
produce partial and transient degeneration and recombination suppression, a scenario involving a
selective pressure maintaining recombination arrest is orders of magnitude more likely to produce
complete and permanent recombination suppression and degeneration.

2. Recombination reestablishment after secondary nested or overlapping inversions

A difficulty to model recombination evolution by inversions and reinversions is that it is difficult to
model the fact that the recombination process is not 'perfect' in the sense that it can occur between
regions that are only locally homologous, as in the case of ectopic recombination. In the case of Y
evolution, recombination may occasionally occur even in the absence exact collinearity between the X
and Y. This is illustrated on Fig S4 for an overlapping inversion that includes the SD locus, and on Fig S5
for nested inversions. In both cases, the secondary inversion has a homologous region on the X and
can pair at meiosis, allowing recombination to occur around the SD locus. If a crossover occurs within
this region, a new Y can be produced which may not carry the exact complement compared to the
original chromosome (with either deleted or duplicated positions). In each case, such a crossover will
eliminate parts of the former Y on the first stratum, i.e., regions that may have already accumulated a
load of deleterious mutations. This recombined Y could be particularly favorable (compared to the
degenerated Y), even if some sequences are duplicated or missing compared to the X (see Figs S4, S5).
Indeed, whether the recombined Y can invade depends on its marginal fitness relative to the marginal
fitness of the potentially highly degenerated Y chromosomes present in the population (and not to the
marginal fitness of a hypothetical mutation-free Y chromosome with full gene content). Furthermore,
the recombined Y may be "improved" in further steps since it can now recombine more easily with the
X around the SD locus after this first recombination event. In particular, a second crossover near the
SD locus can further improve collinearity with the X and eliminate further degenerated parts of the Y
from the first stratum that are still present (see Figs S4, S5). Alternatively, recombination may also be

restored if an inversion arises on the X, facing the inversion on the Y [17]. Again, more empirical work

is needed to assess whether recombination may indeed occur in such scenarios despite non-perfect

collinearity. However, the occurrence of ectopic recombination between repeated sequences indicates

that it is possible in principle [45,57], while the results of the present article show that even very low

rates would be sufficient to maintain recombination in the long term.

Fig S1. Fraction of the time during which the Y carries a given number of fixed strata. During a replicate simulation, there are times during which an inversion is fixed, perhaps including several strata, and times during which no fixed inversion is present (see examples on Fig 1). The bar chart gives this %time, across all replicates when a given number of strata are fixed in the population, for different reversion rates (without regulator evolution as in the constraint scenario, four bars on the left, or with regulator evolution as in the regulatory scenario, four bars on the right). This % is computed excluding the first 500 000 generation (to cut the initial phase influenced by the initial condition where the chromosome starts fully recombining and without any fixed inversion). The gray level corresponds to the number of fixed strata present as indicated in the legend on the right. For instance, in the constraint scenario with reversion rate equal to 10^{-6}, there is no fixed inversion 93.5% of the time, and one fixed inversion 6.3% of the time. In contrast, in the regulatory scenario with the same reversion rate, there is no fixed inversion 7.3% of the time, one fixed inversion 16% of the time, 2 fixed inversions 23% of the time, 3 fixed inversions 23.7% of the time and more than 3 fixed inversions 30% of the time (over the 3.5 million last generations of a simulation).

Fig S2. Detailed characteristics of fixed inversions in the constraint (panels A, B) and regulatory (panels C, D)

scenarios. In all panels, the x-axis gives the lifespan of inversions (as defined in Fig 2A) and the y-axis the size

of the inversion (both in log scale). Each dot represents a different fixed inversion that occurred across all

replicates. The color of the dot indicates the reversion rate of the simulation during which the inversion was

observed (as given in the legend). On panels A and C, a gray disk is added around each inversion. The gray level

indicates the % of gene knocked out (%KO) on that inversion at the last time when the inversion is observed

(i.e., just before it is reversed, at the end of the simulation at 4 million generations, or at population extinction).

On panel A (constraint scenario), the gray level is light as this %KO never exceeds 5%. On panel C (regulatory

scenario), this gray level is darker as this %KO reaches very high values (being either close to zero or above

98%). On panel B and D, a gray disk is also added around each inversion, this time representing the drop in the

male / female fitness ratio caused by this inversion between the first and last time it is observed. Noting r(t)

this male / female ratio at time t, this drop is computed as $r(t_1)/r(t_2)$ between times t_1 and t_2. When several

inversions are simultaneously present in a given time interval, the log(drop) is portioned proportionally to the

relative size of each inversion $s_1/(s_1+s_2)$, i.e. with two inversions of size s_1 and s_2, the drop accrued to the first

inversion is Exp(s1/(s1+s2) Log $r(t_1)/r(t_2)$). For instance, with two inversions of equal size, each is assigned the

square root of the fitness drop on the interval (so that the product of the fitness drop of each inversion gives

the overall fitness drop). With this correction, the fitness drop associated to an inversion is more

representative of what is happening on this inversion (rather than being caused by the presence of another

inversion). On panel B (constraint scenario), these fitness drops can reach large values (more than 99%

reduction in the male/female fitness ratio), but they remain very low in the regulatory scenario (never

exceeding 10% on panel D).

Fig S3. Overall evolution of the Y chromosomes under the constraint and regulatory scenarios. Same

figure than Fig. 3, but with results of regulatory simulations for comparison (open symbols).The x-axis

(in log scale) gives the fraction of the Y that is non recombining (averaged over all males in the

population) after two million generations. The y-axis (in log scale) gives the fraction of knock-out genes

on the Y (averaged over all males in the population; a knock out is defined as a gene having

accumulated deleterious mutations up to $s_{max} = 0.3$), after two million generations. Each dot represents

a replicated population. Open symbols: regulators evolve (regulatory scenario); filled symbol:

regulators do not evolve (constraint scenario). Color codes indicates different rates of reversion (U_{rev}).

The rate of inversion is 10^{-5} in all cases. In a few cases (with $U_{rev} = 10^{-9}$ in the constraint scenario, filled

yellow disks) the population became extinct before 2 million generations. In these cases (marked with

a skull), the x and y axes values are taken at the time of extinction.

Fig. S4. Restoration of recombination on the Y with overlapping inversions. (a) depicts the fully
recombining XY pair with the SD locus in red. The orange dashing indicates where homologous pairing
is possible. A first inversion occurs (purple) on the Y between positions 3 and 8 leading to the situation
in (b). Pairing and recombination do not occur around the SD locus. This first stratum on the Y
chromosome can start degenerating in the absence of recombination. This is shown by the brown color
of the position numbers. A second overlapping inversion occurs (green) on the Y between positions 2
and 5. The resulting Y in (c) can now pair with the X between positions 5 and 8. Crossing-over on the
left of the SD locus (black cross) can generate recombinant Y with duplicated positions 2-4 (note that
this Y recovers functional copies in positions 3-5, and has two functional copies in position 2). Crossing-
over on the right of the SD locus (gray cross) can generate recombinant Y with missing positions 2-4
(note that the lack of these positions may not reduce fitness a lot if they were degenerated). (e) In
both cases, a second crossover with the X occurring near the SD locus (shown with a small cross) can
restore full collinearity with the X and get rid of other degenerated copies (on the right and left position
of the Y, respectively).

Fig. S5. Restoration of recombination on the Y with nested inversions. (a) depicts the fully recombining XY pair with the SD locus in red. The orange dashing indicates where homologous pairing is possible. A first inversion occurs (purple) on the Y between positions 3 and 8 leading to the situation in (b). Pairing and recombination do not occur around the SD locus. This first stratum on the Y chromosome can start degenerating in the absence of recombination. This is shown by the brown color of the position numbers. A second nested inversion occurs (green) on the Y between positions 4 and 7. The resulting Y in (c) can now pair with the X between positions 4 and 7. Crossing-over on the left of the SD locus (black cross) can generate a recombinant Y with duplicated positions 3 and missing 8. Note that this Y can have a higher fitness by recovering functional copies at positions 3-5. Position 8 is missing, but this may not be consequential since it was a degenerated position. Crossing-over on the right of the SD locus (gray cross) can generate a recombinant Y with duplicated position 8 and missing 3. Again, the fitness of this Y may increase, since it recovers functional copies at positions 7-8, while losing a degenerated copy at position 3. (e) In both cases, a second crossover with the X occurring near the sex-determining locus (shown with a small cross) can reconstitute a Y chromosome fully colinear with the X.

Parameters	Values
Standard deviation of mutational effects on cis and trans regulatory traits	0.2
Number of gene, cis-regulator, male-limited and female-limited trans-regulators	500
Intensity of stabilizing selection on expression levels	0.1
Mutation rates on cis-regulators	0.0002
Mutation rates on trans-regulators	0.0001
Recombination rate between cis-regulator and its gene (R_c)	0.00005

816 **Table S1. Additional parameter values used in regulatory simulations.**