The differentiation and integration of the hippocampal dorsoventral axis are controlled by two nuclear receptor genes

Xiong Yang[1#], Rong Wan[1#], Zhiwen Liu[2,3], Su Feng[2,3], Jiaxin Yang[1], Naihe Jing[2,3*], Ke Tang[1*]

1, Precise Genome Engineering Center, School of Life Sciences, Guangzhou University, Guangzhou, Guangdong 510006, China

2, Guangzhou Laboratory/Bioland Laboratory, Guangzhou, 510005, China

3, CAS Key Laboratory of Regenerative Biology, Guangdong Provincial Key Laboratory of Stem Cell and Regenerative Medicine, Guangdong Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China.

#, Co-first authors

*, Co-corresponding authors

Correspondence: njing@sibcb.ac.cn or ktang.sc@gmail.com.

Abstract

The hippocampus executes imperative functions from declarative memory to adaptive

behaviors associated with cognition and emotion. However, the mechanisms of how

morphogenesis and functions along the hippocampal dorsoventral axis are differentiated and

integrated are still largely unclear. Here, we show that *COUP-TFI* and *-TFII* genes are

distinctively expressed in the dorsal and ventral hippocampus, respectively. The loss of

COUP-TFII results in ectopic CA1/CA3 domains in the ventral hippocampus. The deficiency

of *COUP-TFI* leads to the failed specification of dorsal CA1, among which there are place

cells. The deletion of both *COUP-TF* genes causes almost agenesis of the hippocampus with

abnormalities of trisynaptic circuit and adult neurogenesis. Moreover, *COUP-TFI/-TFII* may

cooperate to guarantee appropriate morphogenesis and function of the hippocampus by

regulating the *Lhx5-Lhx2* axis. Our findings revealed a novel mechanism that *COUP-TFI* and

COUP-TFII converge to govern the differentiation and integration of distinct characteristics

of the hippocampus in mice.

Keywords

Memory; Hippocampus; *COUP-TF* Gene; Place Cell; Trisynaptic Circuit

Introduction

Memory, including declarative and nondeclarative memory, unifies our mental world to ensure the quality of life for people of all ages, from newborns to elderly individuals (Eichenbaum & Cohen, 2014; Kandel, Dudai, & Mayford, 2014). The pioneering studies of Milner and her colleagues revealed that the hippocampus is required for declarative memory but not nondeclarative memory (Penfield & Milner, 1958; Scoville & Milner, 1957). The discovery of activity-dependent long-term potentiation and place cells provides the neurophysiological basis of hippocampal function (Bliss & Lomo, 1973; O'Keefe & Dostrovsky, 1971). The rodent hippocampus can be divided into the dorsal and ventral domains, corresponding to the posterior and anterior hippocampus in humans, respectively. In recent decades, numerous studies have supported the Moser theory that the hippocampus is a heterogeneous structure with distinct characteristics of gene expression, connectivity, and functions along its dorsoventral axis (Bast, 2007; Fanselow & Dong, 2010; Moser & Moser, 1998; Strange, Witter, Lein, & Moser, 2014). The dorsal hippocampus, which connects and shares similar gene expression with the neocortex (Fanselow & Dong, 2010), serves the "cold" cognitive function associated with declarative memory and spatial navigation. The ventral hippocampus, which connects and generates similar gene expression with the amygdala and hypothalamus (Cenquizca & Swanson, 2007; Kishi et al., 2000; Pitkanen, Pikkarainen, Nurminen, & Ylinen, 2000), corresponds to the "hot" affective states related to emotion and anxiety (Fanselow & Dong, 2010; Tyng, Amin, Saad, & Malik, 2017). Nonetheless, to date, the molecular and cellular mechanisms by which the morphogenesis, connectivity, and functions along the dorsoventral axis of the hippocampus are differentiated and integrated are largely unknown.

The hippocampus, a medial temporal lobe structure in the adult rodent forebrain, originates from the medial pallium (MP) in the medial line of the early dorsal telencephalon. The

cortical hem (CH), which is located ventrally to the MP, functions as an organizer for

hippocampal development (Hebert & Fishell, 2008; Schuurmans & Guillemot, 2002). It has

been demonstrated that both extrinsic signals, such as *Wnts* and *Bmps*, and intrinsic factors,

including *Emx1*, *Emx2*, *Lef1*, *Lhx2*, and *Lhx5*, are involved in the regulation of early

morphogenesis of the hippocampus. As the earliest *Wnt* gene to be exclusively expressed in

the cortical hem, *Wnt3a* is required for the genesis of the hippocampus (S. M. Lee, Tole,

Grove, & McMahon, 2000); in addition, *Lef1* is downstream of Wnt signaling, and the

hippocampus is completely absent in *Lef1-neo/neo* null mutant mice (Galceran, Miyashita-

Lin, Devaney, Rubenstein, & Grosschedl, 2000). Wnt signaling is essential for early

development of the hippocampus. *Emx1* and *Emx2* are mouse homologs of *Drosophila empty*

spiracles (Simeone et al., 1992). Interestingly, the dorsal hippocampus is smaller in an *Emx1*

null mutant (Yoshida et al., 1997), while *Emx2* is required for the growth of the hippocampus

but not for the specification of hippocampal lineages (Tole, Goudreau, Assimacopoulos, &

Grove, 2000). Moreover, *Lhx5*, which encodes a LIM homeobox transcription factor and is

specifically expressed in the hippocampal primordium, is necessary for the formation of the

hippocampus (Zhao et al., 1999). *Lhx2*, encoding another LIM homeobox transcription factor,

is required for the development of both the hippocampus and neocortex (Mangale et al., 2008;

Monuki, Porter, & Walsh, 2001). Intriguingly, deficiency of either *Lhx5* or *Lhx2* results in

agenesis of the hippocampus, and more particularly, these genes inhibit each other (Hebert &

Fishell, 2008; Mangale et al., 2008; Roy, Gonzalez-Gomez, Pierani, Meyer, & Tole, 2014;

Zhao et al., 1999), indicating that the *Lhx5* and *Lhx2* genes may generate an essential

regulatory axis to ensure the appropriate hippocampal development. Nevertheless, whether

there are other intrinsic genes that participate in the regulation of morphogenesis and function

of the hippocampus has not been fully elucidated.

COUP-TF genes, including *COUP-TFI* (also known as *NR2F1*) and *COUP-TFII* (also

known as *NR2F2*), encode two transcription factor proteins belonging to the nuclear receptor

superfamily (Yang, Feng, & Tang, 2017). Mutations of *COUP-TFI* are highly related to

neurodevelopmental disorders (NDD), such as intellectual disability (ID) and autism

spectrum disorders (ASD) (Bertacchi et al., 2020; Bosch et al., 2014; Contesse, Ayrault,

Mantegazza, Studer, & Deschaux, 2019), and mutations of the *COUP-TFII* gene are

associated with congenital heart defects (CHD) (High et al., 2016). By using animal models,

our studies and others have demonstrated that *COUP-TF* genes participate in the regulation

of the development of the central nervous system (Zhang et al., 2020). The *COUP-TFI* plays

an essential role in the differentiation of cortical excitatory projection neurons and inhibitory

interneurons, the development of the dorsal hippocampus, and cortical arealization

(Armentano et al., 2007; Flore et al., 2017; Lodato et al., 2011; C. Zhou et al., 1999; C. Zhou,

Tsai, & Tsai, 2001). *COUP-TFII* plays a vital role in the development of the amygdala,

hypothalamus, and cerebellum (S. Feng et al., 2017; Kim, Takamoto, Yan, Tsai, & Tsai, 2009;

Tang, Rubenstein, Tsai, & Tsai, 2012). Nevertheless, whether and how *COUP-TFI* and/or

COUP-TFII genes regulate the differentiation and integration of hippocampal morphogenesis,

connectivity, and function is still largely unclear.

Here, our data show that *COUP-TFI* and *-TFII* genes are differentially expressed along the

dorsoventral axis of the adult hippocampus. The loss of *COUP-TFII* results in ectopic CA1

and CA3 domains in the ventral hippocampus. In addition, the deficiency of *COUP-TFI* leads

to not only dysplasia of the dorsal hippocampus but also failed specification and

differentiation of the dorsal CA1 pyramidal neuron lineage. Furthermore, the deletion of both

genes in the *RXCre*/+; *COUP-TFI*$^{F/F}$; *COUP-TFII*$^{F/F}$ double mutant mouse causes almost

agenesis of the hippocampus, accompanied by compromised specification of the CA1, CA3,

and dentate gyrus (DG) domains. The components of the trisynaptic circuit are abnormal in

the corresponding single-gene or double-gene mutant model. Moreover, *COUP-TF* genes

may cooperate to guarantee the appropriate morphogenesis and function of the hippocampus

by regulating the *Lhx5-Lhx2* axis.

Results

Differential expression profiles of *COUP-TFI* and *-TFII* genes along the dorsoventral

axis in the developing and adult hippocampus

To investigate the functions of the *COUP-TFI* and *COUP-TFII* genes in the hippocampus,

immunofluorescence staining was first performed to examine their expression in adult wild-

type mice at postnatal month 1 (1M). In both the coronal and sagittal sections, COUP-TFI

exhibited a septal/dorsal high-temporal/ventral low expression pattern along the hippocampus

(Figure 1Aa, d, c, f, g, j, m, i, l, and o), and its expression is highest in the dorsal CA1 region

(Figure 1Aa, d, c, f, j, and l), where place cells are mainly located (O'Keefe & Conway, 1978;

O'Keefe & Dostrovsky, 1971), and the dorsal DG, where there are adult neural stem cells

(NSCs) (Gould & Cameron, 1996). However, the expression of COUP-TFII was high in the

temporal/ventral hippocampus but was barely detected in the septal/dorsal part of the

hippocampus (Figure 1Ab, c, e, f, h, i, k, l, n, and o). The dorsal-high COUP-TFI and ventral-

high COUP-TFII expression profiles were further verified in the adult hippocampi by western

blotting assays (Figure 1Ap, q). Interestingly, COUP-TFI and -TFII generated

complementary expression patterns in the hippocampal primordium with COUP-TFI in the

dorsal MP and COUP-TFII in the ventral CH at embryonic days 12.5 (E12.5) and 14.5 (E14.5)

(Figure 1—figure supplement 1Aa-f). Additionally, septal/dorsal-high COUP-TFI and

temporal/ventral-high COUP-TFII expression patterns were observed at postnatal day 0 (P0)

(Figure 1—figure supplement 1Ag-l). The data above revealed that the differential expression

patterns of *COUP-TFI* and *-TFII* genes were generated and maintained along the

dorsoventral axis in the early hippocampal primordium, the developing and adult

hippocampus, indicating that they could play distinct roles in the mediation of the

morphogenesis and functions of the hippocampus.

Next, to investigate the roles of *COUP-TF* genes in the hippocampus, an *RXCre* mouse was

used to excise the expression of the *COUP-TFI* and/or *COUP-TFII* genes in the hippocampal

regions. The deletion efficiency of *RXCre* recombinase was verified by immunofluorescence

assays. Compared with control mice, either *COUP-TFII* or *COUP-TFI* could be excised in

the adult hippocampus of corresponding single-gene mutants at 1M (Figure 1—figure

supplement 1Ba-i). In addition, compared with control mice, both the *COUP-TFI* and *-TFII*

genes were almost completely deleted in the hippocampal primordium at E14.5 in mutant

mice (Figure 1—figure supplement 1Bj-o). All the data above show that *RXCre* recombinase

could efficiently excise *COUP-TFI* and/or *COUP-TFII* in the early developing and adult

hippocampus.

The *COUP-TFII* gene is required for the appropriate morphogenesis of the ventral

hippocampus but not of the dorsal hippocampus

Given that the *COUP-TFII* gene is highly and specifically expressed in the adult ventral

hippocampus and the CH of the hippocampal primordium (Figure 1 and Figure 1—figure

supplement 1), we asked whether the *COUP-TFII* gene is required for the appropriate

morphogenesis of the hippocampus, particularly the ventral hippocampus. To answer this

question, we conducted Nissl staining with samples from the *COUP-TFII* single-gene

(*RXCre/+*; *COUP-TFII^{F/F}* or *RXCre/+*; *CII^{F/F}*) knockout mouse model. In coronal sections,

compared with the control at 1M, the septal hippocampus was normal; unexpectedly, an

ectopic CA-like region was observed medially in the temporal hippocampus in the *COUP-*

TFII mutant, where there was the prospected posterior part of the medial amygdaloid nucleus

(MeP), indicated by the star (Figure 1Ba-f). The presence of the ectopic CA-like region in the

ventral but not dorsal hippocampus of the mutant was further confirmed at the prospected

MeP and prospected amygdalohippocampal area (AHi) in sagittal sections, as indicated by

the star (Figure 1Bg-l). Furthermore, immunofluorescence assays were performed to verify

whether specific lineages of the hippocampus were altered with sagittal sections, in which the

subregions of both the dorsal and ventral hippocampus were well displayed and distinguished.

Ctip2 is a marker for CA1 pyramidal neurons and DG granule neurons, and HuB is a marker

for CA3 pyramidal cells (Sugiyama, Osumi, & Katsuyama, 2014). The dorsal hippocampus

appeared normal in both the control and mutant mice at 1M (Figure 1Ca-l). Nonetheless,

compared with the observations in control mice, an ectopic HuB-positive CA3 pyramidal

neuron lineage, indicated by the star, and a duplicated Ctip2-positive CA1 pyramidal neuron

lineage, indicated by the arrowhead, were observed in the ventral hippocampal area in the

mutant (Figure 1Ca-f, m-r), revealing that there were ectopic CA1 and CA3 lineages in the

COUP-TFII mutants.

Consistent with the previous report (Leid et al., 2004), the expression of Ctip2 was detected

in the amygdala including the AHi and posteromedial cortical amygdaloid nucleus (PMCo);

in addition, Ctip2 was also highly expressed in the dorsal part of the MeP (MePD) in the

control (Figure 1Cb, n).

Intriguingly, compared with the controls at 1M, there were ectopic CA domains in the mutant

ventral hippocampus with the expense of the Ctip2 positive AHi and MePD amygdaloid

nuclei (Figure 1Ce, q), indicated by the arrows. Clearly, all the data above suggested that the

COUP-TFII gene is necessary to ensure the appropriate morphogenesis of the ventral

hippocampus.

At early embryonic stages, *COUP-TFII* was preferentially expressed in the CH (Figure 1—

figure supplement 1Ab,e), the organizer of the hippocampus, and at the adult stage, *COUP-*

TFII was also highly expressed in some amygdala nuclei such as the AHi and medial

amygdaloid nucleus, which are adjacent to the ventral/temporal hippocampus (Figure 1Ae,h,n)

(Tang et al., 2012). We would like to investigate the correlation of the CH and/or amygdala

anlage with the duplicated ventral hippocampal domains in the *COUP-TFII* mutant in detail

in our future study. The observations above suggest that *COUP-TFII* is not only specifically

expressed in the ventral hippocampus but is also required for morphogenesis and probably

the function of the ventral hippocampus. Since the ventral hippocampus participates in the

regulation of emotion and stress, mutations in the *COUP-TFII* gene lead to CHDs, and the

formation of the ventral hippocampus is disrupted in *COUP-TFII* adult mutant mice, we

wondered whether CHD patients with *COUP-TFII* mutations also exhibit symptoms

associated with psychiatric disorders such as depression, anxiety, or schizophrenia.

The *COUP-TFI* gene is required for the specification and differentiation of the dorsal

CA1 identity

Next, we asked whether the deletion of the *COUP-TFI* gene by *RXCre* also affected the

development of the hippocampus. Consistent with the previous finding in *Emx1Cre/+*;

COUP-TFI$^{F/F}$ (*Emx1Cre/+*; *CI$^{F/F}$*) mutant mice (Flore et al., 2017), it was the septal/dorsal

hippocampus, not the temporal/ventral hippocampus, that was specifically shrunken in both

coronal (Figure 2Aa-f) and sagittal sections (Figure 2Ag-i) of *RXCre/+*; *COUP-TFI$^{F/F}$*

(*RXCre/+*; *CI$^{F/F}$*) mutant mice. Then, we asked whether the loss of the *COUP-TFI* gene

caused abnormal specification and differentiation of hippocampal lineages. Compared with

the control mice, *COUP-TFI* mutant mice had fewer HuB-positive CA3 pyramidal neurons,

as indicated by the star; intriguingly, Ctip2-positive CA1 pyramidal neurons failed to be

detected, as indicated by the arrowhead, with Ctip2-positive DG granule neurons unaltered in

the dorsal hippocampus (Figure 2Ba-l). The loss of the dorsal CA1 pyramidal neuron identity

in mutant mice was further confirmed by Wfs1, another dorsal CA1 pyramidal neuron-

specific marker (Takeda et al., 2001) (Figure 2Ca-r). Nonetheless, the HuB-positive and

Ctip2-positive lineages were comparable in the ventral hippocampus between the control and

mutant mice (Figure 2Ba-f, m-r), even though the low expression of COUP-TFI was detected

there. Indeed, *COUP-TFI* is not only expressed at the highest level in the dorsal CA1 but is

also required for the specification and differentiation of dorsal CA1 pyramidal neurons,

among which place cells are essential for learning and memory (O'Keefe & Conway, 1978;

O'Keefe & Dostrovsky, 1971).

Given that dysplasia of the dorsal hippocampus was generated in both the *Emx1Cre* and

RXCre models (Flore et al., 2017) (Figure 2Aa-l), we asked whether the development of

dorsal CA1 pyramidal neurons was also abolished in *Emx1Cre/+*; *COUP-TFI$^{F/F}$* mutant mice.

To answer this question, immunofluorescence staining was conducted first. Compared with

that in the control mice, the proportions of either the Wfs1- or Ctip2-positive CA1 domain

were reduced in the mutant mice at 3M (Figure 2—figure supplement 1Aa-h), indicating that

the differentiation of the dorsal CA1 pyramidal neurons was also compromised in the

Emx1Cre model, although it was less severe than that in the *RXCre* model. To make our

findings more consistent with previous studies, we further conducted experiments with the

Emx1Cre model. Afterward, to investigate the fine structure of the dorsal CA1 pyramidal

neurons, Golgi staining was performed. Compared with those of control mice, the numbers of

secondary dendrites and branch points of both the apical and basal dendrites were

significantly reduced in the dorsal CA1 pyramidal neurons of mutant mice at 3M (Figure 2—

figure supplement 1Ba-e). Then, the dorsal hippocampus-related spatial learning and memory

behavior test, the Morris water maze, was performed (Vorhees & Williams, 2006). Consistent

with a previous report (Flore et al., 2017), spatial learning and memory function was

significantly impaired in adult *Emx1Cre*/+; *COUP-TFI$^{F/F}$* mice, compared with the control

mice (Figure 2—figure supplement 1C). The data above suggest that *COUP-TFI* is

imperative for the morphogenesis, lineage specification, and spatial learning and memory of

the dorsal hippocampus, and particularly, the compromised dorsal CA1 lineage could

contribute to the phenotypes associated with neurodevelopmental disorders, including ID or

ASD.

The *COUP-TFI* and *-TFII* genes coordinate to ensure the genesis of the hippocampus

Given that the loss of either *COUP-TFI* or *-TFII* leads to dysplasia of the dorsal or ventral

hippocampus, respectively, we asked whether these genes compensate for each other to

regulate the morphogenesis of the hippocampus. To answer this question, the *RXCre/+*;

COUP-TFI^{F/F}; *COUP-TFII^{F/F}* (*RXCre/+*; *CI^{F/F}*; *CII^{F/F}*) double-mutant mouse was generated,

and a few homozygous double-gene mutant mice survived for approximately 3 weeks (3W).

Nonetheless, the reason for the lethality of the double-mutant mice is still unknown. Nissl

staining data showed that compared with that of control mice, the septal hippocampus was

severely shrunken, as indicated by the star, and the temporal hippocampus was barely

observed in the double-mutant mouse brains (Figure 3Aa-h). Unexpectedly, an ectopic

nucleus was observed in the region of the prospected temporal hippocampus, indicated by the

arrowhead, in the double-mutant mice (Figure 3Ag, h). In addition, compared with those of

controls, the regions with HuB-positive CA3 pyramidal neurons and Ctip2-positive or Prox1-

positive DG granule neurons were diminished in the double mutants; in particular, Ctip2-

positive dorsal CA1 pyramidal neurons could not be detected in the double mutants (Figure

3Ba-l). Furthermore, compared with the domains of controls, no HuB-positive, Ctip2-positive,

or Prox1-positive domains could be detected in the prospected temporal hippocampus in the

double mutants (Figure 3Ca-l). The results above suggest that the *COUP-TFI* and *COUP-*

TFII genes coordinate with each other to mediate the appropriate morphogenesis of the entire

hippocampus.

***COUP-TF* genes and adult neurogenesis in the hippocampus**

Given that the *COUP-TFI* or *-TFII* gene was highly expressed in the dorsal or ventral DG,

respectively (Figure 1A), and that RXCre recombinase could efficiently delete either gene in

the DG (Figure 1—figure supplement 1Ba-i), we asked whether the loss of the *COUP-TFI*

or/and *-TFII* gene in the DG would affect hippocampal adult neurogenesis. To answer this

question, the ventral DG, dorsal DG, and septal DG were chosen to perform

immunofluorescence assays in the *COUP-TFII* mutant, *COUP-TFI* mutant, and double

mutant models, respectively. Adult NSCs in the subgranular zone (SGZ) of the DG express

both GFAP and Nestin, and newborn granule neurons express Dcx (Gao, Arlotta, Macklis, &

Chen, 2007). The numbers of NSCs and newborn neurons in the SGZ of the DG were

comparable between control mice and either the *COUP-TFII* or *COUP-TFI* mutant mice

(Figure 3—figure supplement 1Aa-p, Ba-h); nonetheless, compared with those of control

mice, the numbers of the NSCs and newborn granule neurons in the SGZ of the DG were

reduced in the double mutants (Figure 3—figure supplement 1Aq-x, Bi-l), and the reduction

in both lineages was significant (Figure 3—figure supplement 1Ca, b). The data above

suggest that *COUP-TF* genes may coordinate with each other to execute essential functions

for appropriate hippocampal adult neurogenesis in the DG.

Hippocampal trisynaptic connectivity was impaired in adult *COUP-TFII* single-, *COUP-

***TFI* single-, and double-mutant mice**

Given that dysplasia of the hippocampus was observed in all three mouse models, we asked

whether the connectivity of the hippocampal trisynaptic circuit associated with the DG, CA3,

and CA1 regions (Amaral, 1993) was normal in these models. To answer this question, the

components of the trisynaptic circuit were characterized in the ventral hippocampus of

COUP-TFII mutants, the dorsal hippocampus of *COUP-TFI* mutants, and the septal

hippocampus of double mutants. Calretinin is a marker of mossy cells, Calbindin is a marker

of mossy fibers, and SMI312 is a marker of Schafer collaterals (Flore et al., 2017). Compared

with those of controls (Figure 4Aa, b, e, f, i, and j), the numbers of Calretinin-positive mossy

cells were reduced, Calbindin-positive mossy fibers were longer but thinner, and SMI312-

positive Schafer collaterals were thinner and discontinued in the ventral hippocampus of

COUP-TFII mutants at 1M (Figure 4Ac, d, g, h, k, and l). In addition, similar to the previous

report (Flore et al., 2017), the numbers of Calretinin-positive mossy cells were decreased,

Calbindin-positive mossy fibers were shorter and thinner, and SMI312-positive Schafer

collaterals were barely detected in the dorsal hippocampus of *COUP-TFI* mutants (Figure

4Ac, d, g, h, k, and l), compared with those of controls at 1M (Figure 4Ba, b, e, f, i, and j).

Moreover, compared with those of control mice (Figure 4Ca, b, e, f, i, and j), the numbers of

Calretinin-positive mossy cells were reduced; both Calbindin-positive mossy fibers and

SMI312-positive Schafer collaterals were barely detected in the prospected septal

hippocampus of the double mutants at 3W (Figure 4Cc, d, g, h, k, and l). Clearly, without

both *COUP-TF* genes, the connectivity of the hippocampal trisynaptic circuit was abolished

more severely. The observations above revealed that the formation of the trisynaptic circuit,

which is one of the fundamental characteristics of hippocampal neurophysiology (Basu &

Siegelbaum, 2015), was abnormal in all three mouse models, indicating that both the

morphology and functions of the hippocampus are most likely compromised in the loss of the

COUP-TFI and/or *COUP-TFII* gene.

The expression of several essential regulatory genes associated with early hippocampal development was abnormal in double mutants

Given that the hippocampus was almost completely diminished in double mutants, we asked how *COUP-TF* genes participated in the regulation of the early morphogenesis of the hippocampus. To answer this question, total RNA isolated from the whole telencephalons of control (n=5) and double-mutant-(n=3) embryos at E11.5 was used to generate cDNA, and then real-time quantitative PCR (RT–qPCR) assays were performed. As expected, compared with that of control mice, the expression of *COUP-TFI* and *COUP-TFII* was reduced significantly in the double mutant mice (Figure 5A). Then, we mainly focused on the intrinsic regulatory networks by analyzing the expression profiles of two groups of transcription factor genes. The *Foxg1*, *Gli3*, *Lhx2*, *Otx1*, *Otx2*, and *Pax6* genes, which are highly related to the early patterning of the dorsal telencephalon (Hebert & Fishell, 2008), were in the first group; *Axin2*, *Emx1*, *Emx2*, *Lef1*, *Lhx5*, and *Tcf4* genes, which are associated with early hippocampal development (Galceran et al., 2000; Moore & Iulianella, 2021; Tole et al., 2000; Yoshida et al., 1997; Zhao et al., 1999), were in the other group. The expression of the *Foxg1*, *Gli3*, *Lhx2*, *Otx1*, *Otx2*, and *Pax6* genes was comparable between the controls and double mutants (Figure 5A), indicating that the early patterning of the dorsal telencephalon is largely unaltered. Compared with that of control mice, the expression of the *Axin2*, *Emx2*, *Lef1*, and *Tcf4* genes was normal in the double mutants; interestingly, the expression of the *Emx1* and *Lhx5* transcripts was decreased significantly in the double mutants at E11.5 compared to that in control mice (Figure 5A). In addition, the reduced expression of Lhx5 protein was confirmed in the hippocampal primordium in the double mutants at E14.5 by immunofluorescence assay (Figure 5Ba-d, a'-d'). The expression of *Lhx2* was expanded ventrally into the choroid plexus in the *Lhx5* null mutant mice (Zhao et al., 1999), indicating that *Lhx5* could inhibit *Lhx2* expression locally. Interestingly, the expression of the Lhx2

protein was increased in the hippocampal primordium, indicated by the arrowhead, in the

COUP-TF double-mutant mice at E14.5 (Figure 5Be-h, e'-h').

Next, we asked whether neural precursor cells (NPCs), intermediate progenitor cells (IPCs),

or newborn neurons were affected in the early development of the hippocampus in double-

mutant mice. Sox2 is a marker for NPCs, Tbr2 is a marker for IPCs, and NeuroD1 is a marker

for newborn neurons (Yu, Marchetto, & Gage, 2014). The expression of Sox2 in the

hippocampal regions was comparable between the control and double-mutant mice at E14.5

(Figure 5Ca-d, a'-d'), indicating that the generation of NPCs was normal. Nevertheless,

compared with the control embryos, the numbers of Tbr2-positive IPCs and NeuroD1-

positive newborn neurons were reduced in the double-mutant embryos (Figure 5Ce-l, 5Ce'-

l'), and the reduction was significant (Figure 5D). Our observations were consistent with

previous findings in *Lhx5* null mutant mice that the specification of the hippocampal NPCs

was normal, but the later differentiation event was abolished (Zhao et al., 1999). All the data

above suggest that *COUP-TF* genes may cooperate to ensure the early morphogenesis of the

hippocampus by regulating the appropriate expression levels of *Lhx5* and *Lhx2* genes.

Nevertheless, we could not exclude other possibilities that *COUP-TF* genes could also

participate in the modulation of hippocampal development through *Emx1* or other genes.

Discussion

In our present study, we observed dorsal-high COUP-TFI and ventral-high COUP-TFII

expression profiles in the adult hippocampus. The deletion of the *COUP-TFII* gene led to

duplicated CA1 and CA3 domains of the ventral hippocampus. The loss of *COUP-TFI*

resulted in the failed specification and differentiation of the dorsal CA1 pyramidal neuron

lineage with a diminished dorsal hippocampus. Furthermore, the deficiency of both *COUP-*

TF genes caused atrophy of almost the entire hippocampus, accompanied by compromised

generation of the CA1, CA3, and DG identities. In addition, the dorsal trisynaptic

components, ventral trisynaptic components, or entire trisynaptic components were abolished

in the corresponding *COUP-TFI* gene mutant, *COUP-TFII* gene mutant, or *COUP-TFI/-TFII*

double-gene mutant mice. Moreover, *COUP-TF* genes may cooperate to ensure the

appropriate morphogenesis and function of the hippocampus by regulating the *Lhx5-Lhx2*

axis.

1. *COUP-TFII* governs the distinct characteristics of the ventral hippocampus

Sixty years ago, the pioneering work of Milner and her colleagues discovered the essential

role of the hippocampus in declarative memory (Penfield & Milner, 1958; Scoville & Milner,

1957). Recently, accumulating evidence has supported the Moser theory that the

hippocampus is a heterogeneous structure with distinct characteristics of gene expression,

connectivity, and function along its dorsoventral axis (Bast, 2007; Fanselow & Dong, 2010;

Moser & Moser, 1998; Strange et al., 2014). The dorsal hippocampus marked in blue, in

which gene expression is similar to the neocortex, serves the "cold" cognitive function

associated with declarative memory and spatial navigation, and the ventral hippocampus

marked in red, in which gene expression is close to the hypothalamus and amygdala,

corresponds to the "hot" affective states related to emotion and anxiety (Figure 5—figure

supplement 1). The ventral hippocampus generates direct connectivity with the amygdala,

hypothalamus, medial prefrontal cortex (mPFC), and olfactory bulb (Cenquizca & Swanson,

2007; Hoover & Vertes, 2007; Kishi et al., 2000; Pitkanen et al., 2000; Roberts et al., 2007).

Nonetheless, thus far, the molecular and cellular mechanism of how the morphogenesis,

connectivity, and function of the ventral hippocampus is achieved has been largely unclear.

COUP-TFII, a nuclear receptor gene associated with heart disease (High et al., 2016), was

highly and exclusively expressed in the ventral hippocampus in adult mice and was expressed

ventrally in the CH of the hippocampal primordium in mouse embryos (Figure 1, Figure 1—

figure supplement 1, Figure 5Ea), indicating that the *COUP-TFII* gene may participate in the

regulation of the development and function of the ventral hippocampus. First, deficiency of

the *COUP-TFII* gene led to the duplication of the CA1 and CA3 domains of the ventral

hippocampus but not the dorsal hippocampus, which was confirmed both morphologically

and molecularly (Figure 1). Second, the formation of the trisynaptic circuit was specifically

abolished in the ventral hippocampus of *COUP-TFII* mutants (Figure 4), indicating that the

intrahippocampal circuit, information transfer, and function of the ventral hippocampus could

be compromised. Third, the ventral hippocampus generates neural circuits with the mPFC,

amygdala, nucleus accumbens, and hypothalamus, which are associated with

anxiety/behavioral inhibition, fear processing, pleasure/reward seeking, and the

neuroendocrine system, respectively (Anacker & Hen, 2017; Baik, 2020; Bryant & Barker,

2020; Cenquizca & Swanson, 2007; Herman et al., 2016; Kishi et al., 2000; O'Leary & Cryan,

2014; Pitkanen et al., 2000). These ventral hippocampal projections may be important for

processing information related to emotion and anxiety. Intriguingly, our previous studies

revealed that *COUP-TFII* is required for the development of the hypothalamus, amygdala,

and olfactory bulb (S. Feng et al., 2017; Tang et al., 2012; X. Zhou et al., 2015), all of which

generate functional neural circuits with the ventral hippocampus (Fanselow & Dong, 2010).

Particularly, both the hypothalamus and amygdala are also diminished in *RXCre/+*; *COUP-*

TFII^{F/F} mutant mice (S. Feng et al., 2017; Tang et al., 2012), indicating that their

interconnectivities with the ventral hippocampus are abnormal. Thus, all the findings above

suggest that *COUP-TFII* is a novel and essential intrinsic regulator that controls the

morphogenesis, connectivity, and function of the ventral hippocampus.

Given that mutations of *COUP-TFII* are highly associated with CHDs and are required for

the distinct characteristics of the ventral hippocampus, we wondered whether CHD patients

carrying mutations of *COUP-TFII* also display symptoms of psychiatric disorders, such as

depression, anxiety, or schizophrenia, related to the ventral hippocampus. In our future study,

we would like to generate hippocampus-specific or hippocampal subdomain-specific

conditional knockout models to dissect distinct roles of the *COUP-TFII* gene in the

hippocampus, particularly in the ventral hippocampus, in detail.

2. The *COUP-TFI* gene is required for the specification and differentiation of dorsal

CA1 pyramidal neurons

The expression of *COUP-TFI*, another orphan nuclear receptor gene associated with

neurodevelopmental disorders (Bertacchi et al., 2020; Bosch et al., 2014; Contesse et al.,

2019), is high in the dorsal MP of the hippocampal primordium and is higher in the dorsal

hippocampus than in the ventral hippocampus (Figure 1, Figure 1—figure supplement 1,

Figure 5Ea) (Flore et al., 2017). Consistent with previous observations in *Emx1Cre/+*;

COUP-TFI^{F/F} mutant mice (Flore et al., 2017), the dorsal hippocampus but not the ventral

hippocampus was specifically shrunken in *RXCre/+; COUP-TFI^{F/F}* mice (Figure 2, Figure

5—figure supplement 1). *COUP-TFI* is expressed at the highest level in dorsal CA1

pyramidal neurons (Figure 1), indicating that the *COUP-TFI* gene may play a role in the

specification and differentiation of dorsal CA1 pyramidal neurons. As expected, the

expression of Ctip2 and Wfs1, two markers for dorsal CA1 pyramidal neurons, could not be

detected in the prospected dorsal CA1 domain in *RXCre/+; COUP-TFI^{F/F}* mutant mice

(Figure 2); furthermore, *Emx1Cre/+; COUP-TFI^{F/F}* mutant mice partially phenocopied the

compromised development of the dorsal CA1 lineage (Figure 2—figure supplement 1). It

seems that the spatiotemporal activity of RXCre recombinase is better or broader than that of

the Emx1Cre recombinase during the critical period of the specification of the dorsal CA1

pyramidal neuron identity. In addition, the Golgi staining assay revealed that the

development of the dendrites of the dorsal CA1 pyramidal neurons was abnormal (Figure 2—

figure supplement 1). All the observations above indicate that the *COUP-TFI* gene is not only

necessary for the morphogenesis of the dorsal hippocampus but is also required for the

specification and differentiation of the dorsal CA1 pyramidal neurons, among which there are

place cells. The identification of place cells fifty years ago was one of the most important

breakthroughs to understand the memory and hippocampus (O'Keefe & Dostrovsky, 1971).

Except for spatial information, place cells in the dorsal CA1 may also encode nonspatial

representations, such as time (Eichenbaum, 2017; Lisman et al., 2017). Notably, 95% of

patients carrying *COUP-TFI* mutations are associated with ID. Here, our observations

support the notion that the *COUP-TFI* gene is a novel intrinsic regulator that specifies the

dorsal CA1 pyramidal cell identity, which will benefit the understanding of both

neurophysiological functions of the hippocampus and the etiology of NDD including ID and

ASD.

3. *COUP-TFI* and *-TFII* cooperate to ensure the appropriate morphogenesis of the

hippocampus by regulating the *Lhx5-Lhx2* axis in mice

In wild-type mice, *COUP-TFI* and *-TFII* genes generated complementary expression profiles

in the embryonic hippocampal primordium with *COUP-TFI* in the dorsal MP, marked in

green; *COUP-TFII* in the ventral CH, marked in red (Figure 5Ea, Figure 1—figure

supplement 1); and in the adult hippocampus with high-COUP-TFI expression in the dorsal,

marked in green; and high-COUP-TFII expression in the ventral, marked in red (Figure 5Ea,

Figure 1). These findings indicated that *COUP-TF* genes may coordinate to regulate

hippocampal development. Indeed, as discussed above, the loss of either *COUP-TFI* or *-TFII*

only leads to dysplasia of the dorsal hippocampus (Flore et al., 2017) (Figure 2, Figure 2—

figure supplement 1) or ventral hippocampus (Figure 1), respectively; intriguingly, while

both genes are efficiently excised by *RXCre* in the hippocampal primordium (Figure 1—

figure supplement 1), more severely shrunken hippocampi developed in the adult double

knockout mice (Figure 3). The dosage-dependent severity of hippocampal abnormalities

suggested that two nuclear receptor genes, *COUP-TFI* and *-TFII* could cooperate with each

other to execute an imperative and intrinsic function in the development of the hippocampus.

It is known that both extrinsic signals and intrinsic factors participate in the regulation of the

early development of the hippocampus. Notably, mutations of *Wnt3a* and *Lef1* eliminate the

entire hippocampus (Galceran et al., 2000; S. M. Lee et al., 2000). Given that the expression

of *Axin2*, *Lef1* and *Tcf4*, three Wnt-responsive transcription factor genes, was not altered in

the *COUP-TF* double mutant (Figure 5A), it is unlikely that abnormal Wnt signaling is the

cause of the compromised hippocampus. *Lhx5* is specifically expressed in the hippocampal

primordium and is required for the morphogenesis of the hippocampus (Zhao et al., 1999).

Lhx2 is necessary for hippocampal development by repressing cortical hem fate (Mangale et

al., 2008; Monuki et al., 2001). Agenesis of the hippocampus is observed in either *Lhx5* or

Lhx2 null mutant mice, and these genes particularly repress each other (Hebert & Fishell,

2008; Mangale et al., 2008; Roy et al., 2014; Zhao et al., 1999), indicating that the proper

expression levels of *Lhx5* and *Lhx2* genes are imperative to maintain the appropriate

development of the hippocampus (Figure 5Eb). The transcriptional expression of *Lhx5* but

not *Lhx2* was reduced in *COUP-TF* double-mutant mice at E11.5, and three days later,

decreased expression of the Lhx5 protein and enhanced expression of the Lhx2 protein were

detected in the hippocampal primordium of double-mutant mice (Figure 5A-B). In addition,

consistent with the observations in the *Lhx5* null mutant (Li et al., 2021; Miquelajáuregui et

al., 2010), the generation of Sox2-positive hippocampal NPCs was not affected, but the

development of Tbr2-positive IPCs and NeuroD1-positive newborn neurons was abnormal in

COUP-TF double-mutant mice (Figure 5). Thus, our findings reveal a novel intrinsic

regulatory mechanism that *COUP-TFI* and *COUP-TFII*, two disease-associated nuclear

receptor genes, may cooperate with each other to ensure proper hippocampal morphogenesis

by regulating the *Lhx5-Lhx2* axis. Intriguingly, compared with the adult *Emx1Cre/+*; *COUP-*

TFI^{F/F} mutant mice, the hippocampus was much smaller in the adult *Emx1Cre/+*; *COUP-*

TFI^{F/F}; *COUP-TFII^{F/F}* double-gene mutant mice; nevertheless, both the dorsal and ventral

hippocampus were readily detected in double-gene mutants with *Emx1Cre* (our unpublished

observations). In addition, the discrepancy between the shrunken dorsal hippocampus

associated with the loss of *COUP-TFI* and the duplicated CA domains of the ventral

hippocampus associated with the deficiency of *COUP-TFII* suggested that the regulatory

network related to *COUP-TF* genes during the early morphogenesis of the hippocampus

could be much more complicated than suspected and should be investigated in our future

study.

4. *COUP-TF* genes are imperative for the formation of the trisynaptic circuit

The hippocampus and entorhinal cortex (EC) are interconnected through various neural

circuits to mediate the flow of the information associated with declarative memory (Basu &

Siegelbaum, 2015). Both direct and indirect glutamatergic circuits are involved in the relay of

information from the EC to the hippocampal CA1, and the trisynaptic pathway is the most

well-characterized indirect circuit. The EC sends sensory signals from association cortices

via the perforant path to the DG, then the DG granule cells send excitatory mossy fiber

projections to CA3 pyramidal neurons, and CA3 pyramidal neurons project to CA1 via the

Schaffer collaterals (H. Lee, GoodSmith, & Knierim, 2020). Consistent with the high

expression of COUP-TFI in the dorsal hippocampus and COUP-TFII in the ventral

hippocampus (Figure 1), the dorsal trisynaptic circuit is specifically damaged in *COUP-TFI*

mutants, as is the ventral trisynaptic circuit in *COUP-TFII* mutant mice. Moreover, the

hippocampal trisynaptic circuit was almost completely absent in the double mutants (Figure 4,

and our unpublished observations). The information transfer associated with the trisynaptic

circuits should be abolished particularly in the dorsal and/or the ventral hippocampus in the

above corresponding genetic mouse models. Interestingly, *COUP-TFI* is also required to

specify the medial EC cell fate (J. Feng et al., 2021). Therefore, the impaired formation and

function of trisynaptic circuits could be caused by the abnormal development of CA1, CA3,

DG or EC lineages. Nonetheless, given that newborn granule neurons are continuously

generated in the adult DG to integrate into the existing neural circuits essential for declarative

memory (Toda, Parylak, Linker, & Gage, 2019; Tuncdemir, Lacefield, & Hen, 2019) and that

hippocampal adult neurogenesis was severely compromised in the *COUP-TF* double mutant

(Figure 3—figure supplement 1), we could not exclude the possibility that impaired adult

neurogenesis may also contribute to the malformation and impaired function of the

trisynaptic pathway in double mutants.

The hippocampus is heterogeneous along its dorsoventral axis, and either the dorsal or

ventral hippocampus generates unique and distinguishable characteristics of gene expression

and connectivity, which enable the hippocampus to execute an integrative function from the

encoding and retrieval of certain declarative memory to adaptive behaviors. Lesions of the

dorsal hippocampus, which are essential for the cognitive process of learning and memory,

lead to amnesia and ID; while damage to the ventral hippocampus, which is imperative for

emotion and affection, is highly associated with psychiatric disorders including depression,

anxiety, and schizophrenia. Our findings in this study reveal novel intrinsic mechanisms by

which two nuclear receptor genes, *COUP-TFI* and *-TFII*, which are associated with NDD or

CHD, converge to govern the differentiation and integration of the hippocampus along the

dorsoventral axis morphologically and functionally. Furthermore, our present study provides

novel genetic model systems to investigate the crosstalk among the hippocampal complex in

gene expression, morphogenesis, cell fate specification and differentiation, connectivity,

functions of learning/memory and emotion/anxiety, adaptive behaviors, and the etiology of

neurological diseases. Nevertheless, many enigmas, such as whether and how the

abnormalities of either the dorsal or ventral hippocampus affect the characteristics of the

other, remain unsolved. In addition to the excitatory lineages and circuits, interneurons and

inhibitory circuits play vital roles in maintaining the plasticity and functions of the

hippocampus. We also wonder whether and how defects in interneurons and inhibitory

circuits could contribute to the compromised morphogenesis, connectivity, and functions of

the hippocampus and the etiology of psychiatric and neurological conditions including ID,

ASD, depression, anxiety, and schizophrenia.

544
Materials and Methods

Animals

COUP-TFI-floxed mice, *COUP-TFII-floxed* mice, *RXCre* mice and *Emx1Cre* mice used in

the study were of the C57B6/129 mixed background. The noon of vaginal plug day was set as

the embryonic day 0.5 (E0.5). All animal protocols were approved by the Animal Ethics

Committee of the Shanghai Institute of Biochemistry and Cell Biology. All methods were

performed in accordance with the relevant guidelines and regulations. Only the littermates

were used for the comparison.

Nissl staining

We used xylene to dewax paraffin sections, followed by rinsing with 100%, 95%, and 70%

ethanol. The slides were stained in 0.1% Cresyl Violet solution for 25 mins. Then the

sections were washed quickly in the water and differentiated in 95% ethanol. We used 100%

ethanol to dehydrate the slides, followed by rinsing with the xylene solution. Finally, the

neutral resin medium was used to mount the slides.

Immunohistochemical (IHC) staining

The paraffin sections were dewaxed and rehydrated as described above for Nissl staining.

The slides were boiled in 1×antigen retrieval solution (DAKO) under microwave conditions

for 15 min. After cooled to room temperature (RT), the slides were incubated with 3% H2O2

for 30 min. Then, the slides were treated with blocking buffer for 60 min at RT and then

incubated with the primary antibody in the hybridization buffer (10×diluted blocking buffer)

overnight (O/N) at 4 °C. The next day, the tyramide signal amplification kit (TSA)

(Invitrogen) was used according to the manufacturer's protocol. After being incubated with

1%TSA blocking buffer, the sections were treated with a biotinylated secondary antibody for

60 min at RT. After being washed with 1×PBS three times, the slides were incubated with

1×HRP-conjugated streptavidin for 1 h. Next, the tyramide working solution was prepared,

including the 0.15‰ H_2O_2 in distilled water, the 100×diluted tyramide substrate solution

(tyramide-488 or tyramide-594), and the amplification buffer. The sections were incubated

with the working solution for 10 min. Then the slides were counterstained with DAPI and

mounted with the antifade mounting medium (Southern Biotech) (S. Feng et al., 2017; Zhang

et al., 2020). Finally, the sections were observed and images were captured with a digital

fluorescence microscope (Zeiss).

The following primary antibodies were used in the study: mouse anti-COUP-TFI (1:1000,

R&D), mouse anti-COUP-TFII (1:2000, R&D), rabbit anti-HuB (1:500, Abcam), rat anti-

Ctip2 (1:500, Abcam), rabbit anti-Wfs1 (1:500, ProteinTech), goat anti-Prox1 (1:500, R&D),

rabbit anti-Calretinin (1:500, Sigma), rabbit anti-Calbindin (1:500, Swant), mouse anti-

SMI312(1:200, Covance), rabbit anti-Sox2 (1:500, Affinity BioReagents), rat anti-Tbr2

(1:500,Thermo Fisher), goat anti-NeuroD1 (1:200, Santa Cruz), goat anti-Lhx2 (1:200, Santa

Cruz), rabbit anti-Lhx5 (1:200, Millipore), mouse anti-GFAP(1:500, Sigma), rabbit anti-

Nestin (1:200, Santa Cruz), goat anti-Dcx (1:500, Santa Cruz). The following secondary

antibodies were used in the study: donkey anti-mouse IgG biotin-conjugated (1:400,

JacksonImmuno), donkey anti-rabbit IgG biotin-conjugated (1:400, JacksonImmuno), donkey

anti-goat IgG biotin-conjugated (1:400, JacksonImmuno), donkey anti-rat IgG biotin-

conjugated (1:400, JacksonImmuno).

Western blotting

We homogenized the isolated dorsal and ventral hippocampus tissues respectively in the

RIPA buffer (Applygen) with protease inhibitor cocktail (Sigma) and phosphatase inhibitors

(Invitrogen) and then centrifuged at the speed of 12,000 rpm for 30 mins. We collected the

supernatants and analyzed the total concentrations by the BCA kit (Applygen). Gradient

SDS-PAGE gels were used to separate the same amounts of protein sample (40 μg/lane), and

then the proteins were transferred to the PVDF membranes (Millipore). After being blocked

by the 3% BSA (Sigma) for 2h, the membranes that contained proteins were incubated by

primary antibodies at 4 °C O/N. The membranes were rinsed with 1×PBST three times for 10

mins and then treated with biotinylated secondary antibodies for 2h at RT. After washing

with 1×PBST, membranes were treated with the HRP-conjugated streptavidin for 1h at RT.

Finally, we used the chemiluminescence detection system (Tanon) to detect the bands. The

density of the protein band was analyzed by the software Image J.

The primary antibodies were used in the experiment as below: mouse anti-COUP-TFI

(1:2000, R&D), rabbit anti-COUP-TFII (1:3000, R&D). The following secondary antibodies

were applied in the study, including goat anti-mouse IgG biotin-conjugated (1:1000, KPL),

goat anti-rabbit IgG biotin-conjugated (1:1000, KPL).

Golgi staining

Deep anesthesia was performed before sacrificing the control and mutant mice, and the brains were immediately isolated. The FD rapid GolgiStain kit (FD NeuroTech) was used to process the brain tissue samples, which were immersed in an equal volume of immersion solution mixed with solutions A and B and stored in the dark for two weeks at RT. At least 5ml of immersion solution is used for each cubic meter of the tissue. For the best results, the container of tissue was gently swirled from side to side twice a week during the incubating period. And then the brain tissue was transferred to solution C in the dark at RT for at least 72 hours (up to 1 week). Finally, the tissues were cut into 100 μm thick slices with a cryostat at -20°C to -22°C and transferred to gelatin-coated microscope slides containing solution C using a sample retriever. The slices were dried naturally at RT. The concrete staining procedure of the kit was followed using the manufactory's protocol. Then, the sections were rinsed twice with double-distilled water for 4 minutes each time. The slices were placed in a mixture of one volume of solution D, one volume of solution E, and two volumes of double distilled water for 10 minutes and were rinsed twice with distilled water for 4 minutes each time. Then the sections were dehydrated in 50%, 75%, and 95% ethanol for 4 minutes respectively. Next, the slices were dehydrated in 100% ethanol 4 times for 4 minutes each time. Finally, the sections were cleared in xylene and mounted with a neutral resin medium.

Morris water maze

By recording the time spent by the mice swimming in the water tank and finding the escape platform hidden underwater, and the swimming trajectory, the Morris water maze test can objectively reflect the spatial learning and memory ability of the mice. We poured tap water into the water maze tank and added an appropriate amount of milky white food dye that was

mixed well. The height of the liquid level was about 1 cm higher than the escape platform,

and the water temperature was kept at about 25±1℃. At the same time, four markers of

different shapes were pasted on the four directions of the inner wall above the water tank to

distinguish different directions. The Morris water maze test was divided into the training

phase and the probe trial phase. The training phase lasted for 6 days, 4 times a day, and the

interval between each training was about 30 minutes. During training, the mice were placed

into the tank from the entry points of four different quadrants facing the inner wall, and their

latency was recorded from the time from entering the water to finding a hidden underwater

platform and standing on it. After the mouse found the platform, let it stay on the platform for

10 seconds before removing it. If the mouse failed to discover the platform 60 seconds after

entering the water, it was guided to find the platform and stayed for 10 seconds. Each mouse

was placed into the water tank from four water entry points and recorded as one training

session. The probe trial was carried out on the seventh day and the underwater platform was

removed. Each experimental mouse was put into the water tank at the same water entry point,

and allowed to move for 60 seconds. The time that each mouse spent in the quadrant, where

the platform was originally placed, was recorded.

RNA isolation and quantitative real-time PCR

Total RNAs were prepared from the whole telencephalon of the control (n=5) and double

mutant (n=3) mice at E11.5 respectively, with the TRIzol Reagent (Invitrogen) by following

the manufactory's protocol. Reverse-transcription PCR and real-time quantitative PCR assays

were performed as described previously (Tang et al., 2012). A student's t-test was used to

compare the means of the relative mRNA levels between the control group and mutant group.

Primer sequences are as follows:

Axin2-f, 5'-ctgctggtcaggcaggag-3', Axin2-r, 5'-tgccagtttctttggctctt-3'; Coup-tfI-f, 5'-

caaagccatcgtgctattca-3', Coup-tfI-r, 5'-cctgcaggctttcgatgt-3'; Coup-tfII-f, 5'-cctcaaagtgggcatgagac-

3', Coup-tfII-r, 5'-tgggtaggctgggtaggag-3'; Emx1-f, 5'-ctctccgagacgcaggtg-3', Emx1-r, 5'-

ctcagactccggcccttc-3'; Emx2-f, 5'-cacgcttttgagaagaacca-3', Emx2-r, 5'-gttctccggttctgaaacca-3';

Foxg1-f, 5'-gaaggcctccacagaacg-3', Foxg1-r, 5'-ggcaaggcatgtagcaaaag-3'; Gli3-f, 5'-

tgatccatctcctattcctcca-3', Gli3-r, 5'-tctggatacgtcgggctact-3'; Lef1-f, 5'-tcctgaaatccccaccttct-3', Lef1-

r, 5'-tgggataaacaggctgacct-3'; Lhx2-f, 5'-cagcttgcgcaaaagacc-3', Lhx2-r, 5'-taaaaggttgcgcctgaact-

3'; Lhx5-f, 5'-tgtgcaataagcagctatcca-3'; Lhx5-r, 5'-caaactgcggtccgtaca-3'; Otx1-f, 5'-

ccagagtccagagtccaggt-3', Otx1-r, 5'-ccgggttttcgttccatt-3'; Otx2-f, 5'-ggtatggacttgctgcatcc-3', Otx2-r,

5'-cgagctgtgccctagtaaatg-3'; Pax6-f, 5'-gttccctgtcctgtggactc-3', Pax6-r, 5'-accgcccttggttaaagtct-3';

Tcf4-f, 5'-aaatggccactgcttgatgt-3', Tcf4-r, 5'-gcaccaccggtactttgttc-3'.

Quantification and statistical analysis

The number of specified immunofluorescent marker-positive cells was assessed by Image J

Cell Counter in full image fields. Three brain sections per mouse were counted for each index.

GraphPad Prism 7.0 (GraphPad) was used to perform statistical analysis. The data analysis

used one-way analysis of variance (ANOVA), Dunnett's or Tukey's post hoc tests, and

student's unpaired t-test. The data were expressed as the mean ± SEM. The data obtained

from at least three independent replicates were used for statistical analysis. $P < 0.05$ was

considered the significant statistical difference.

Acknowledgements

We thank Ms. Emerald Tang for her assistance with the manuscript. This work was supported

by National Natural Science Foundation of China (31671508) and Guangdong Provincial

Basic and Applied Basic Research Fund (2021A1515011299) to K.T; and in part by the

National Key Basic Research and Development Program of China (2018YFA0108500,

2019YFA0801402, 2018YFA0800100, 2018YFA0108000, 2018YFA0107200,

2017YFA0102700), "Strategic Priority Research Program" of the Chinese Academy of

Sciences, Grant No. (XDA16020501, XDA16020404) to N. J.

Author contributions

X.Y., W.R. designed and conducted the experiments. X.Y. and K.T. organized and wrote the

manuscript. Z.W.L., S.F. and J.X.Y. assisted in conducting the experiments and analyzed the

data. N.J. and K.T. conceived the project and approved the manuscript.

Competing interests

The authors declare no competing interests.

References
696

Amaral, D. G. (1993). Emerging principles of intrinsic hippocampal organization. *Current*
Opinion in Neurobiology, 3(2), 225-229. doi:https://doi.org/10.1016/0959-
4388(93)90214-J
Anacker, C., & Hen, R. (2017). Adult hippocampal neurogenesis and cognitive flexibility -
linking memory and mood. *Nat Rev Neurosci, 18*(6), 335-346.
doi:10.1038/nrn.2017.45
Armentano, M., Chou, S. J., Tomassy, G. S., Leingärtner, A., O'Leary, D. D., & Studer, M.
(2007). COUP-TFI regulates the balance of cortical patterning between frontal/motor
and sensory areas. *Nat Neurosci, 10*(10), 1277-1286. doi:10.1038/nn1958
Baik, J. H. (2020). Stress and the dopaminergic reward system. *Exp Mol Med, 52*(12), 1879-
1890. doi:10.1038/s12276-020-00532-4
Bast, T. (2007). Toward an integrative perspective on hippocampal function: from the rapid
encoding of experience to adaptive behavior. *Rev Neurosci, 18*(3-4), 253-281.
doi:10.1515/revneuro.2007.18.3-4.253
Basu, J., & Siegelbaum, S. A. (2015). The Corticohippocampal Circuit, Synaptic Plasticity,
and Memory. *Cold Spring Harb Perspect Biol, 7*(11).
doi:10.1101/cshperspect.a021733
Bertacchi, M., Romano, A. L., Loubat, A., Tran Mau-Them, F., Willems, M., Faivre, L., . . .
Studer, M. (2020). NR2F1 regulates regional progenitor dynamics in the mouse
neocortex and cortical gyrification in BBSOAS patients. *Embo j, 39*(13), e104163.
doi:10.15252/embj.2019104163
Bliss, T. V., & Lomo, T. (1973). Long-lasting potentiation of synaptic transmission in the
dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J*
Physiol, 232(2), 331-356. doi:10.1113/jphysiol.1973.sp010273

Bosch, D. G., Boonstra, F. N., Gonzaga-Jauregui, C., Xu, M., de Ligt, J., Jhangiani, S., . . . Schaaf, C. P. (2014). NR2F1 mutations cause optic atrophy with intellectual disability. *Am J Hum Genet, 94*(2), 303-309. doi:10.1016/j.ajhg.2014.01.002

Bryant, K. G., & Barker, J. M. (2020). Arbitration of Approach-Avoidance Conflict by Ventral Hippocampus. *Front Neurosci, 14*, 615337. doi:10.3389/fnins.2020.615337

Cenquizca, L. A., & Swanson, L. W. (2007). Spatial organization of direct hippocampal field CA1 axonal projections to the rest of the cerebral cortex. *Brain Res Rev, 56*(1), 1-26. doi:10.1016/j.brainresrev.2007.05.002

Contesse, T., Ayrault, M., Mantegazza, M., Studer, M., & Deschaux, O. (2019). Hyperactive and anxiolytic-like behaviors result from loss of COUP-TFI/Nr2f1 in the mouse cortex. *Genes Brain Behav, 18*(7), e12556. doi:10.1111/gbb.12556

Eichenbaum, H. (2017). On the Integration of Space, Time, and Memory. *Neuron, 95*(5), 1007-1018. doi:10.1016/j.neuron.2017.06.036

Eichenbaum, H., & Cohen, N. J. (2014). Can we reconcile the declarative memory and spatial navigation views on hippocampal function? *Neuron, 83*(4), 764-770. doi:10.1016/j.neuron.2014.07.032

Fanselow, M. S., & Dong, H. W. (2010). Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron, 65*(1), 7-19. doi:10.1016/j.neuron.2009.11.031

Feng, J., Hsu, W. H., Patterson, D., Tseng, C. S., Hsing, H. W., Zhuang, Z. H., . . . Chou, S. J. (2021). COUP-TFI specifies the medial entorhinal cortex identity and induces differential cell adhesion to determine the integrity of its boundary with neocortex. *Sci Adv, 7*(27). doi:10.1126/sciadv.abf6808

Feng, S., Xing, C., Shen, T., Qiao, Y., Wang, R., Chen, J., . . . Tang, K. (2017). Abnormal Paraventricular Nucleus of Hypothalamus and Growth Retardation Associated with Loss of Nuclear Receptor Gene COUP-TFII. *Sci Rep, 7*(1), 5282. doi:10.1038/s41598-017-05682-6

Flore, G., Di Ruberto, G., Parisot, J., Sannino, S., Russo, F., Illingworth, E. A., . . . De Leonibus, E. (2017). Gradient COUP-TFI Expression Is Required for Functional Organization of the Hippocampal Septo-Temporal Longitudinal Axis. *Cereb Cortex, 27*(2), 1629-1643. doi:10.1093/cercor/bhv336

Galceran, J., Miyashita-Lin, E. M., Devaney, E., Rubenstein, J. L., & Grosschedl, R. (2000). Hippocampus development and generation of dentate gyrus granule cells is regulated by LEF1. *Development, 127*(3), 469-482. doi:10.1242/dev.127.3.469

Gao, X., Arlotta, P., Macklis, J. D., & Chen, J. (2007). Conditional knock-out of beta-catenin in postnatal-born dentate gyrus granule neurons results in dendritic malformation. *J Neurosci, 27*(52), 14317-14325. doi:10.1523/jneurosci.3206-07.2007

Gould, E., & Cameron, H. A. (1996). Regulation of neuronal birth, migration and death in the rat dentate gyrus. *Dev Neurosci, 18*(1-2), 22-35. doi:10.1159/000111392

Hebert, J. M., & Fishell, G. (2008). The genetics of early telencephalon patterning: some assembly required. *Nat Rev Neurosci, 9*(9), 678-685. doi:10.1038/nrn2463

Herman, J. P., McKlveen, J. M., Ghosal, S., Kopp, B., Wulsin, A., Makinson, R., . . . Myers, B. (2016). Regulation of the Hypothalamic-Pituitary-Adrenocortical Stress Response. *Compr Physiol, 6*(2), 603-621. doi:10.1002/cphy.c150015

High, F. A., Bhayani, P., Wilson, J. M., Bult, C. J., Donahoe, P. K., & Longoni, M. (2016). De novo frameshift mutation in COUP-TFII (NR2F2) in human congenital diaphragmatic hernia. *Am J Med Genet A, 170*(9), 2457-2461. doi:10.1002/ajmg.a.37830

Hoover, W. B., & Vertes, R. P. (2007). Anatomical analysis of afferent projections to the medial prefrontal cortex in the rat. *Brain Struct Funct, 212*(2), 149-179. doi:10.1007/s00429-007-0150-4

Kandel, E. R., Dudai, Y., & Mayford, M. R. (2014). The molecular and systems biology of memory. *Cell, 157*(1), 163-186. doi:10.1016/j.cell.2014.03.001

Kim, B. J., Takamoto, N., Yan, J., Tsai, S. Y., & Tsai, M. J. (2009). Chicken Ovalbumin Upstream Promoter-Transcription Factor II (COUP-TFII) regulates growth and patterning of the postnatal mouse cerebellum. *Dev Biol, 326*(2), 378-391. doi:10.1016/j.ydbio.2008.11.001

Kishi, T., Tsumori, T., Ono, K., Yokota, S., Ishino, H., & Yasui, Y. (2000). Topographical organization of projections from the subiculum to the hypothalamus in the rat. *J Comp Neurol, 419*(2), 205-222. doi:10.1002/(sici)1096-9861(20000403)419:2<205::aid-cne5>3.0.co;2-0

Lee, H., GoodSmith, D., & Knierim, J. J. (2020). Parallel processing streams in the hippocampus. *Curr Opin Neurobiol, 64*, 127-134. doi:10.1016/j.conb.2020.03.004

Lee, S. M., Tole, S., Grove, E., & McMahon, A. P. (2000). A local Wnt-3a signal is required for development of the mammalian hippocampus. *Development, 127*(3), 457-467. doi:10.1242/dev.127.3.457

Leid, M., Ishmael, J. E., Avram, D., Shepherd, D., Fraulob, V., & Dolle, P. (2004). CTIP1 and CTIP2 are differentially expressed during mouse embryogenesis. *Gene Expr Patterns, 4*(6), 733-739. doi:10.1016/j.modgep.2004.03.009

Li, J., Sun, L., Peng, X. L., Yu, X. M., Qi, S. J., Lu, Z. J., . . . Shen, Q. (2021). Integrative genomic analysis of early neurogenesis reveals a temporal genetic program for differentiation and specification of preplate and Cajal-Retzius neurons. *PLoS Genet, 17*(3), e1009355. doi:10.1371/journal.pgen.1009355

Lisman, J., Buzsaki, G., Eichenbaum, H., Nadel, L., Ranganath, C., & Redish, A. D. (2017). Viewpoints: how the hippocampus contributes to memory, navigation and cognition. *Nat Neurosci, 20*(11), 1434-1447. doi:10.1038/nn.4661

Lodato, S., Tomassy, G. S., De Leonibus, E., Uzcategui, Y. G., Andolfi, G., Armentano, M., . . . Studer, M. (2011). Loss of COUP-TFI alters the balance between caudal ganglionic eminence- and medial ganglionic eminence-derived cortical interneurons and results in resistance to epilepsy. *J Neurosci, 31*(12), 4650-4662. doi:10.1523/jneurosci.6580-10.2011

Mangale, V. S., Hirokawa, K. E., Satyaki, P. R., Gokulchandran, N., Chikbire, S., Subramanian, L., . . . Monuki, E. S. (2008). Lhx2 selector activity specifies cortical identity and suppresses hippocampal organizer fate. *Science, 319*(5861), 304-309. doi:10.1126/science.1151695

Miquelajáuregui, A., Varela-Echavarría, A., Ceci, M. L., García-Moreno, F., Ricaño, I., Hoang, K., . . . Zhao, Y. (2010). LIM-homeobox gene Lhx5 is required for normal development of Cajal-Retzius cells. *J Neurosci, 30*(31), 10551-10562. doi:10.1523/jneurosci.5563-09.2010

Monuki, E. S., Porter, F. D., & Walsh, C. A. (2001). Patterning of the dorsal telencephalon and cerebral cortex by a roof plate-Lhx2 pathway. *Neuron, 32*(4), 591-604. doi:10.1016/s0896-6273(01)00504-9

Moore, S. A., & Iulianella, A. (2021). Development of the mammalian cortical hem and its derivatives: the choroid plexus, Cajal-Retzius cells and hippocampus. *Open Biol, 11*(5), 210042. doi:10.1098/rsob.210042

Moser, M. B., & Moser, E. I. (1998). Functional differentiation in the hippocampus. *Hippocampus, 8*(6), 608-619. doi:10.1002/(SICI)1098-1063(1998)8:6<608::AID-HIPO3>3.0.CO;2-7

O'Keefe, J., & Conway, D. H. (1978). Hippocampal place units in the freely moving rat: why
they fire where they fire. *Exp Brain Res, 31*(4), 573-590. doi:10.1007/BF00239813
O'Keefe, J., & Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary
evidence from unit activity in the freely-moving rat. *Brain Res, 34*(1), 171-175.
doi:10.1016/0006-8993(71)90358-1
O'Leary, O. F., & Cryan, J. F. (2014). A ventral view on antidepressant action: roles for adult
hippocampal neurogenesis along the dorsoventral axis. *Trends Pharmacol Sci, 35*(12),
675-687. doi:10.1016/j.tips.2014.09.011
Penfield, W., & Milner, B. (1958). Memory deficit produced by bilateral lesions in the
hippocampal zone. *AMA Arch Neurol Psychiatry, 79*(5), 475-497.
doi:10.1001/archneurpsyc.1958.02340050003001
Pitkanen, A., Pikkarainen, M., Nurminen, N., & Ylinen, A. (2000). Reciprocal connections
between the amygdala and the hippocampal formation, perirhinal cortex, and
postrhinal cortex in rat. A review. *Ann N Y Acad Sci, 911*, 369-391.
doi:10.1111/j.1749-6632.2000.tb06738.x
Roberts, A. C., Tomic, D. L., Parkinson, C. H., Roeling, T. A., Cutter, D. J., Robbins, T. W.,
& Everitt, B. J. (2007). Forebrain connectivity of the prefrontal cortex in the
marmoset monkey (Callithrix jacchus): an anterograde and retrograde tract-tracing
study. *J Comp Neurol, 502*(1), 86-112. doi:10.1002/cne.21300
Roy, A., Gonzalez-Gomez, M., Pierani, A., Meyer, G., & Tole, S. (2014). Lhx2 regulates the
development of the forebrain hem system. *Cereb Cortex, 24*(5), 1361-1372.
doi:10.1093/cercor/bhs421
Schuurmans, C., & Guillemot, F. (2002). Molecular mechanisms underlying cell fate
specification in the developing telencephalon. *Curr Opin Neurobiol, 12*(1), 26-34.
doi:10.1016/s0959-4388(02)00286-6
Scoville, W. B., & Milner, B. (1957). Loss of recent memory after bilateral hippocampal
lesions. *J Neurol Neurosurg Psychiatry, 20*(1), 11-21. doi:10.1136/jnnp.20.1.11
Simeone, A., Gulisano, M., Acampora, D., Stornaiuolo, A., Rambaldi, M., & Boncinelli, E.
(1992). Two vertebrate homeobox genes related to the Drosophila empty spiracles
gene are expressed in the embryonic cerebral cortex. *EMBO J, 11*(7), 2541-2550.
doi:10.1002/j.1460-2075.1992.tb05319.x
Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of
the hippocampal longitudinal axis. *Nat Rev Neurosci, 15*(10), 655-669.
doi:10.1038/nrn3785
Sugiyama, T., Osumi, N., & Katsuyama, Y. (2014). A novel cell migratory zone in the
developing hippocampal formation. *J Comp Neurol, 522*(15), 3520-3538.
doi:10.1002/cne.23621
Takeda, K., Inoue, H., Tanizawa, Y., Matsuzaki, Y., Oba, J., Watanabe, Y., . . . Oka, Y.
(2001). WFS1 (Wolfram syndrome 1) gene product: predominant subcellular
localization to endoplasmic reticulum in cultured cells and neuronal expression in rat
brain. *Hum Mol Genet, 10*(5), 477-484. doi:10.1093/hmg/10.5.477
Tang, K., Rubenstein, J. L., Tsai, S. Y., & Tsai, M. J. (2012). COUP-TFII controls amygdala
patterning by regulating neuropilin expression. *Development, 139*(9), 1630-1639.
doi:10.1242/dev.075564
Toda, T., Parylak, S. L., Linker, S. B., & Gage, F. H. (2019). The role of adult hippocampal
neurogenesis in brain health and disease. *Mol Psychiatry, 24*(1), 67-87.
doi:10.1038/s41380-018-0036-2
Tole, S., Goudreau, G., Assimacopoulos, S., & Grove, E. A. (2000). Emx2 is required for
growth of the hippocampus but not for hippocampal field specification. *J Neurosci,*
20(7), 2618-2625. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/10729342

Tuncdemir, S. N., Lacefield, C. O., & Hen, R. (2019). Contributions of adult neurogenesis to
dentate gyrus network activity and computations. *Behav Brain Res, 374*, 112112.
doi:10.1016/j.bbr.2019.112112
Tyng, C. M., Amin, H. U., Saad, M. N. M., & Malik, A. S. (2017). The Influences of
Emotion on Learning and Memory. *Front Psychol, 8*, 1454.
doi:10.3389/fpsyg.2017.01454
Vorhees, C. V., & Williams, M. T. (2006). Morris water maze: procedures for assessing
spatial and related forms of learning and memory. *Nat Protoc, 1*(2), 848-858.
doi:10.1038/nprot.2006.116
Yang, X., Feng, S., & Tang, K. (2017). COUP-TF Genes, Human Diseases, and the
Development of the Central Nervous System in Murine Models. *Curr Top Dev Biol,*
125, 275-301. doi:10.1016/bs.ctdb.2016.12.002
Yoshida, M., Suda, Y., Matsuo, I., Miyamoto, N., Takeda, N., Kuratani, S., & Aizawa, S.
(1997). Emx1 and Emx2 functions in development of dorsal telencephalon.
Development, 124(1), 101-111. doi:10.1242/dev.124.1.101
Yu, D. X., Marchetto, M. C., & Gage, F. H. (2014). How to make a hippocampal dentate
gyrus granule neuron. *Development, 141*(12), 2366-2375. doi:10.1242/dev.096776
Zhang, K., Yu, F., Zhu, J., Han, S., Chen, J., Wu, X., . . . Tang, K. (2020). Imbalance of
Excitatory/Inhibitory Neuron Differentiation in Neurodevelopmental Disorders with
an NR2F1 Point Mutation. *Cell Rep, 31*(3), 107521. doi:10.1016/j.celrep.2020.03.085
Zhao, Y., Sheng, H. Z., Amini, R., Grinberg, A., Lee, E., Huang, S., . . . Westphal, H. (1999).
Control of hippocampal morphogenesis and neuronal differentiation by the LIM
homeobox gene Lhx5. *Science, 284*(5417), 1155-1158.
doi:10.1126/science.284.5417.1155
Zhou, C., Qiu, Y., Pereira, F. A., Crair, M. C., Tsai, S. Y., & Tsai, M. J. (1999). The nuclear
orphan receptor COUP-TFI is required for differentiation of subplate neurons and
guidance of thalamocortical axons. *Neuron, 24*(4), 847-859. doi:10.1016/s0896-
6273(00)81032-6
Zhou, C., Tsai, S. Y., & Tsai, M. J. (2001). COUP-TFI: an intrinsic factor for early
regionalization of the neocortex. *Genes Dev, 15*(16), 2054-2059.
doi:10.1101/gad.913601
Zhou, X., Liu, F., Tian, M., Xu, Z., Liang, Q., Wang, C., . . . Yang, Z. (2015). Transcription
factors COUP-TFI and COUP-TFII are required for the production of granule cells in
the mouse olfactory bulb. *Development, 142*(9), 1593-1605. doi:10.1242/dev.115279
903
904

Figure legends

Figure 1. Duplicated CA1 and CA3 domains are generated in the ventral hippocampus of $RXCre/+$; $CII^{F/F}$ mutant mice.

A, The expression of COUP-TFI (**a, d, g, j, m**) and COUP-TFII (**b, e, h, k, n**) in coronal sections (**a-f**) and sagittal sections (**g-o**) of the hippocampus at postnatal month 1 (1M); representative Western blots and quantitative densitometry data for the expression of COUP-TFI and COUP-TFII in the dorsal and ventral hippocampus at 1M (**p-q**). **B,** In coronal sections along the rostrocaudal axis (**a-f**) and sagittal sections along the lateral-medial axis (**g-l**) of the hippocampus in mutant mice, compared with that in control mice (**a-c, g-i**), the ectopic CA-like structure, indicated by the star, was observed in the ventral region in $COUP$-$TFII$ gene mutant ($RXCre/+$; $CII^{F/F}$) mice at 1M (**d-f, j-l**). **C,** The expression of HuB and Ctip2 in the corresponding inserted area in **a-f** under a high magnification objective lens at 1M (**g-r**); compared with those of control mice (**a-c, g-i, m-o**), the duplicated HuB-positive CA3 domain, indicated by the star, and Ctip2-positive domains, indicated by the arrowhead, were specifically observed in the ventral hippocampus (**d-f, p-r**) but not in the dorsal hippocampus (**d-f, j-l**) of $COUP$-$TFII$ mutant mice at 1M; Ctip2 positive AHi and MePD amygdaloid nuclei were barely observed in the $COUP$-$TFII$ mutant mice, indicated by the arrows, instead of the ectopic CA domains at the prospected amygdaloid regions (**e-f, q-r**). Data are expressed as the mean ± SEM. *Student's t test*, [*]$P<0.05$, [**]$P<0.01$. dCA1, dorsal CA1; dCA3, dorsal CA3; dDG, dorsal dentate gyrus; vCA1, ventral CA1; vCA3, ventral CA3; vDG, ventral dentate gyrus; PMCo, posteromedial cortical amygdaloid nucleus; AHi, amygdalohippocampal area; dHPC, dorsal hippocampus; vHPC, ventral hippocampus; AMY, amygdala nuclei; CTX, cortex; MePD, posterodorsal part of the medial amygdaloid nucleus. Scale bars, **Aa-c**, **Ad-f**, **Aj-o**, **Cg-r**, 100 µm; **Ag-i**, **Ba-l**, **Ca-f**, 200µm.

Figure 1—figure supplement 1. The expression of *COUP-TF* genes in the early developing hippocampus and different conditional knock mouse models.

A, The expression of *COUP-TFI* and *COUP-TFII* genes in the developing hippocampus at E12.5 (**a-c**), E14.5 (**d-f**), and P0 (**g-l**). **B,** Compared with that of control mice (**a-c**), *COUP-TFII* is efficiently deleted by RXCre recombinase in the hippocampus of *COUP-TFII* mutant mice at 1M (**d-f**); *COUP-TFI* is clearly deleted by RXCre recombinase in the hippocampus of *COUP-TFI* mutant mice at 1M (**g-i**). Compared with that of control mice (**j-l**), COUP-TFI and COUP-TFII were efficiently deleted by RXCre recombinase at the hippocampal primordium, including MP and CH, in *COUP-TFI/-TFII* double-mutant mice at E14.5 (**m-o**). MP, medial pallium; CH, cortical hem; HPC, hippocampus; dHPC, dorsal hippocampus; vHPC, ventral hippocampus. Scale bars, **Aa-f**, **Bj-o**, 200 μm; **Ag-l**, **Ba-i**, 100 μm.

Figure 2. The specification and differentiation of the dorsal CA1 lineage failed with the dysplastic dorsal hippocampus in *RXCre/+*; *CI$^{F/F}$* mutant mice.

A, In coronal sections along the rostrocaudal axis (**a-f**) and sagittal sections along the lateral-medial axis (**g-l**) of the hippocampus, compared with that of control mice (**a-c**, **g-i**), the dorsal hippocampus was shrunken, indicated by the star, in *COUP-TFI* gene mutant (*RXCre/+;CI$^{F/F}$*) mice at 1M (**d-f**, **j-l**). **B,** The expression of HuB and Ctip2 in the corresponding inserted area in **a-f** under a high magnification objective lens at 1M (**g-r**); compared with that of control mice (**a-c**, **g-i**, **m-o**), the HuB-positive CA3 domain was reduced in the dorsal hippocampus, especially the Ctip2-positive dorsal CA1, which was barely detected in *COUP-TFI* mutant mice at 1M (**d-f**, **j-l**), while their expression in the ventral hippocampus was comparable between the controls and mutants (**d-f**, **p-r**). **C,** The expression of HuB and Wfs1 in the corresponding inserted area in **a-f** under a high magnification objective lens at 1M (**g-r**); the expression of HuB and the dCA1 marker Wfs1 in the control (**a-c**, **g-i**, **m-o**) and *COUP-TFII*

mutant mice (**d-f**, **j-l**, **p-r**) at 1M. Wfs1-positive dorsal CA1 could not be detected in *COUP-*

TFI mutant mice at 1M, as indicated by the arrowhead. HPC, hippocampus; dHPC, dorsal

hippocampus; vHPC, ventral hippocampus. Scale bars, **Aa-l**, **Ba-f**, **Ca-f,** 200 μm; **Bg-r**, **Cg-r,**

100 μm.

Figure 2—figure supplement 1. Defects in *Emx1Cre/+*; *COUP-TFI^{F/F}* mutant mice.

A, Immunofluorescence staining data showed that compared with those of control mice (**a, c,**

e, g), the proportions of either the Wfs1- or Ctip2-positive dorsal CA1 domain were reduced

in *Emx1Cre/+*; *COUP-TFI^{F/F}* mutant mice at 3M (**b, d, f, h**). **B,** Golgi staining showed that

compared with those of controls, the numbers of branch points and secondary dendrites of

both apical and basal dendrites were significantly reduced in the dorsal hippocampal CA1

pyramidal neurons of *Emx1Cre/+*; *COUP-TFI^{F/F}* mutant mice (**a-c**). **C,** The Morris water

maze behavior test showed that compared with that of controls, spatial learning and memory

were significantly damaged in *Emx1Cre/+*; *COUP-TFI^{F/F}* mutant mice. Data are expressed as

the mean ± SEM. *Student's t test,* $^{*}P<0.05$, $^{**}P<0.01$, $^{***}P<0.001$. Scale bars, **Aa-h,** 100 μm;

Ba-b, 50 μm.

Figure 3. Defects in the hippocampus in *RXCre/+*; *CI^{F/F}*; *CII^{F/F}* double-gene mutant

mice.

A, In coronal sections along the rostrocaudal axis, compared with control mice (**a-d**), the

hippocampus was atrophic in *RXCre/+*; *CI^{F/F}; CII^{F/F}* double mutant mice, indicated by the

star, and an ectopic unknown nucleus was observed in the caudal plates, indicated by the

arrowhead (**e-h**). **B,** Compared with that of control mice (**a-c**, **g-i**), the expression of HuB,

Ctip2, and Prox1 was decreased in the hippocampus of *COUP-TFI/-TFII* double-gene mutant

mice at 3 weeks postnatal (3W) (**d-f**, **j-l**). **C,** Compared with that of control mice (**a-c**, **g-i**),

the expression of HuB could not be detected in the presumptive CA3 domain, and the

expression of Ctip2 or Prox1 could not be detected in the presumptive DG domain of the

prospected ventral hippocampus of *RXCre/+*; *CI^{F/F}*; *CII^{F/F}* double mutant mice. Scale bars,

Aa-h, 200 μm; **Ba-l**, **Ca-l**, 100 μm.

Figure 3—figure supplement 1. Adult neurogenesis was abnormal in the hippocampi of

COUP-TFI/-TFII **double-gene mutant mice.**

A, The expression of GFAP and Nestin, markers of NSCs, in the SGZ of the vDG in control

and *COUP-TFII* mutant mice at 1M (**a-h**), in the SGZ of the dDG in control and *COUP-TFI*

mutant mice at 1M (**i-p**) and in the SGZ of the DG in control and *COUP-TFI/-TFII* double-

gene mutant mice at 3W (**q-x**). **B**, The expression of Dcx, a marker of newborn neurons, in

the SGZ of the vDG in control and *COUP-TFII* mutant mice at 1M (**a-d**), in the SGZ of the

dDG in control and *COUP-TFI* mutant mice at 1M (**e-h**) and in the SGZ of the DG in control

and *COUP-TFI/-TFII* double-gene mutant mice at 3W (**i-l**). **C,** Quantitative analysis of

GFAP/Nestin-positive cells (**a**) and Dcx-positive cells (**b**) in the SGZ of the DG in control

and *COUP-TFI/-TFII* double-gene mutant mice at 3W. The numbers of GFAP and Nestin

double-positive NSCs and Dcx-positive newborn neurons were significantly reduced in

double-mutant mice. Data are expressed as the mean ± SEM. *Student's t test*, **$P<0.01$,

***$P<0.001$. vDG, ventral DG; dDG, dorsal DG; NSC, neural stem cell; SGZ, subgranular

zone. Scale bars, **Aa-x**, 50 μm; **Ba-l**, 100 μm.

1001

1002 **Figure 4. The impairment of hippocampal trisynaptic connectivity in *COUP-TFII***

1003 **single-gene, *COUP-TFI* single-gene, and *COUP-TFI/-TFII* double-gene mutant mice.**

1004 **A,** The expression of Calretinin, Calbindin, and SMI312 in the ventral hippocampus of the

1005 control (**a-b**, **e-f**, **i-j**) and *COUP-TFII* single-gene mutant mice (**c-d**, **g-h**, **k-l**). **B,** The

expression of Calretinin, Calbindin, and SMI312 in the dorsal hippocampus of the control (**a-b**, **e-f**, **i-j**) and *COUP-TFI* single-gene mutant mice (**c-d**, **g-h**, **k-l**). **C,** The expression of Calretinin, Calbindin, and SMI312 in the hippocampus of the control (**a-b**, **e-f**, **i-j**) and *COUP-TFI/-TFII* double-gene mutant mice (**c-d**, **g-h**, **k-l**). vHPC, ventral hippocampus; dHPC, dorsal hippocampus; HPC, hippocampus. Scale bars, **Aa-l**, **Ba-l**, **Ca-l**, 100 μm.

Figure 5. *COUP-TF* genes regulate the expression of key genes associated with early hippocampal development.

A, The expression profiles of genes involved in hippocampal development in control and the double mutant mice at E11.5. **B,** Compared with that of control mice (**a-b**, **a'-b'**), the expression of Lhx5 was reduced in double-mutant mice at E14.5 (**c-d**, **c'-d'**); compared with that of control mice (**e-f**, **e'-f'**), the expression of Lhx2 was increased in double-mutant mice at E14.5 (**g-h**, **g'-f'**). **C,** Compared with that of control mice (**a-b**, **a'-b'**), the expression of Sox2 was normal in double-mutant mice at E14.5 (**c-d**, **c'-d'**); compared with that of control mice (**e-f**, **e'-f'**), the expression of Tbr2 was decreased in *COUP-TF* mutant mice at E14.5 (**g-h**, **g'-h'**); compared with that of control mice (**i-j**, **i'-j'**), the expression of NeuroD1 was reduced in double-mutant mice at E14.5 (**k-l**, **k'-l'**). **D,** Quantitative analysis of Tbr2-positive cells and NeuroD1-positive cells in **Ce'-h'** and **Ci'-l'**. **E,** In the hippocampal primordium of the early embryo, *COUP-TFI* is expressed dorsally in the MP, and *COUP-TFII* is expressed ventrally in the CH. In the mature adult hippocampus, the expression of *COUP-TFI* is higher in the dorsal hippocampus, which is related to spatial learning and memory, and the expression of *COUP-TFII* is mainly in the ventral hippocampus, which is associated with emotion and anxiety (**a**). Our findings support a novel molecular mechanism by which *COUP-TFI* and *COUP-TFII* may cooperate to ensure the appropriate morphogenesis and functions of the hippocampus by modulating the *Lhx5-Lhx2* axis (**b**). Data are expressed as

1031 the mean ± SEM. *Student's t test*, [**]*P*<0.01. HP, hippocampal primordium; DP, dorsal

1032 pallium; MP, medial pallium; CH, cortical hem; ChP, choroid plexus. Scale bars, **Ba-d**, **Be-h**,

1033 **Ca-d**, **Ce-h**, **Ci-l**, 200 μm; **Ba'-d'**, **Be'-h'**, **Ce'-h'**, **Ci'-l'**, 50 μm.

1034
Figure 5—figure supplement 1. *COUP-TFI* **and** *-TFII* **genes coordinate to control**

distinct characteristics of the hippocampus.

Roles of *COUP-TFI* and *COUP-TFII* genes in the development and function of the

hippocampus and the association with neurological diseases. *COUP-TFI* is required for the

morphogenesis of the dorsal hippocampus and the specification of dorsal CA1 pyramidal

neuron lineage, which are associated with neurodevelopmental disorders, including ID and

ASDs. *COUP-TFII* is required to prevent the duplication of the CA1 and CA3 lineages of the

ventral hippocampus, which may be related to psychiatric diseases such as depression,

anxiety, or schizophrenia. The *COUP-TFI* and *COUP-TFII* genes are novel intrinsic

regulatory genes, which cooperate with each other to ensure the early morphogenesis of the

hippocampus.

1046

Figure 1, Yang et al.

Figure 1—figure supplement 1, Yang et al.

A

B

Figure 3, Yang et al.

Figure 3—figure supplement 1, Yang et al.

Figure 4, Yang et al.

Figure 5, Yang et al.

Figure 5—figure supplement 1, Yang et al.