The differentiation and integration of the hippocampal dorsoventral axis are controlled

by two nuclear receptor genes

Xiong Yang[1#], Rong Wan[1#], Zhiwen Liu[2,3], Su Feng[2,3], Jiaxin Yang[1], Naihe Jing[2,3*], Ke

Tang[1*]

1, Precise Genome Engineering Center, School of Life Sciences, Guangzhou University,

Guangzhou, Guangdong 510006, China

2, Guangzhou Laboratory/Bioland Laboratory, Guangzhou, 510005, China

3, CAS Key Laboratory of Regenerative Biology, Guangdong Provincial Key Laboratory of

Stem Cell and Regenerative Medicine, Guangdong Institutes of Biomedicine and Health,

Chinese Academy of Sciences, Guangzhou 510530, China.

#, Co-first authors

*, Co-corresponding authors

Correspondence: njing@sibcb.ac.cn or ktang.sc@gmail.com.

17
18

Abstract

The hippocampus executes crucial functions from declarative memory to adaptive behaviors

associated with cognition and emotion. However, the mechanisms of how morphogenesis and

functions along the hippocampal dorsoventral axis are differentiated and integrated are still

largely unclear. Here, we show that *Nr2f1* and *Nr2f2* genes are distinctively expressed in the

dorsal and ventral hippocampus, respectively. The loss of *Nr2f2* results in ectopic CA1/CA3

domains in the ventral hippocampus. The deficiency of *Nr2f1* leads to the failed specification

of dorsal CA1, among which there are place cells. The deletion of both *Nr2f* genes causes

almost agenesis of the hippocampus with abnormalities of trisynaptic circuit and adult

neurogenesis. Moreover, *Nr2f1/2* may cooperate to guarantee appropriate morphogenesis and

function of the hippocampus by regulating the *Lhx5-Lhx2* axis. Our findings revealed a novel

mechanism that *Nr2f1* and *Nr2f2* converge to govern the differentiation and integration of

distinct characteristics of the hippocampus in mice.

Keywords

Memory; Hippocampus; *Nr2f* Gene; Place Cell; Trisynaptic Circuit

Introduction

Memory, including declarative and nondeclarative memory, unifies our mental world to ensure the quality of life for people of all ages, from newborns to elderly individuals (Eichenbaum & Cohen, 2014; Kandel, Dudai, & Mayford, 2014). The pioneering studies of Milner and her colleagues revealed that the hippocampus is required for declarative memory but not nondeclarative memory (Penfield & Milner, 1958; Scoville & Milner, 1957). The discovery of activity-dependent long-term potentiation and place cells provides the neurophysiological basis of hippocampal function (Bliss & Lomo, 1973; O'Keefe & Dostrovsky, 1971). The rodent hippocampus can be divided into the dorsal and ventral domains, corresponding to the posterior and anterior hippocampus in humans, respectively. In recent decades, numerous studies have supported the Moser theory that the hippocampus is a heterogeneous structure with distinct characteristics of gene expression, connectivity, and functions along its dorsoventral axis (Bast, 2007; Fanselow & Dong, 2010; Moser & Moser, 1998; Strange, Witter, Lein, & Moser, 2014). The dorsal hippocampus, which connects and shares similar gene expression with the neocortex (Fanselow & Dong, 2010), serves the "cold" cognitive function associated with declarative memory and spatial navigation. The ventral hippocampus, which connects and generates similar gene expression with the amygdala and hypothalamus (Cenquizca & Swanson, 2007; Kishi et al., 2000; Pitkanen, Pikkarainen, Nurminen, & Ylinen, 2000), corresponds to the "hot" affective states related to emotion and anxiety (Fanselow & Dong, 2010; Tyng, Amin, Saad, & Malik, 2017). Nonetheless, to date, the molecular and cellular mechanisms by which the morphogenesis, connectivity, and functions along the dorsoventral axis of the hippocampus are differentiated and integrated are largely unknown.

The hippocampus, a medial temporal lobe structure in the adult rodent forebrain, originates from the medial pallium (MP) in the medial line of the early dorsal telencephalon. The cortical hem (CH), which is located ventrally to the MP, functions as an organizer for hippocampal

development (Hebert & Fishell, 2008; Schuurmans & Guillemot, 2002). It has been

demonstrated that both extrinsic signals, such as *Wnts* and *Bmps*, and intrinsic factors,

including *Emx1*, *Emx2*, *Lef1*, *Lhx2*, and *Lhx5*, are involved in the regulation of early

morphogenesis of the hippocampus. As the earliest *Wnt* gene to be exclusively expressed in

the cortical hem, *Wnt3a* is required for the genesis of the hippocampus (S. M. Lee, Tole, Grove,

& McMahon, 2000); in addition, *Lef1* is downstream of Wnt signaling, and the hippocampus

is completely absent in *Lef1^{neo/neo}* null mutant mice (Galceran, Miyashita-Lin, Devaney,

Rubenstein, & Grosschedl, 2000). Wnt signaling is essential for early development of the

hippocampus. *Emx1* and *Emx2* are mouse homologs of *Drosophila empty spiracles* (Simeone

et al., 1992). Interestingly, the dorsal hippocampus is smaller in an *Emx1* null mutant (Yoshida

et al., 1997), while *Emx2* is required for the growth of the hippocampus but not for the

specification of hippocampal lineages (Tole, Goudreau, Assimacopoulos, & Grove, 2000).

Moreover, *Lhx5*, which encodes a LIM homeobox transcription factor and is specifically

expressed in the hippocampal primordium, is necessary for the formation of the hippocampus

(Zhao et al., 1999). *Lhx2*, encoding another LIM homeobox transcription factor, is required for

the development of both the hippocampus and neocortex (Mangale et al., 2008; Monuki, Porter,

& Walsh, 2001; Porter et al., 1997). Intriguingly, deficiency of either *Lhx5* or *Lhx2* results in

agenesis of the hippocampus, and more particularly, these genes inhibit each other (Hebert &

Fishell, 2008; Mangale et al., 2008; Roy, Gonzalez-Gomez, Pierani, Meyer, & Tole, 2014;

Zhao et al., 1999), indicating that the *Lhx5* and *Lhx2* genes may generate an essential regulatory

axis to ensure the appropriate hippocampal development. Nevertheless, whether there are other

intrinsic genes that participate in the regulation of morphogenesis and function of the

hippocampus has not been fully elucidated.

Nr2f genes, including *Nr2f1* and *Nr2f2,* encode two transcription factor proteins belonging to

the nuclear receptor superfamily (Yang, Feng, & Tang, 2017). Mutations of *Nr2f1* are highly

related to neurodevelopmental disorders (NDD), such as intellectual disability (ID) and autism

spectrum disorders (ASD) (Bertacchi et al., 2020; Bosch et al., 2014; Contesse, Ayrault,

Mantegazza, Studer, & Deschaux, 2019), and mutations of the *Nr2f2* gene are associated with

congenital heart defects (CHD) (Al Turki et al., 2014). By using animal models, our studies

and others have demonstrated that *Nr2f* genes participate in the regulation of the development

of the central nervous system (Zhang et al., 2020). The *Nr2f1* plays an essential role in the

differentiation of cortical excitatory projection neurons and inhibitory interneurons, the

development of the dorsal hippocampus, and cortical arealization (Armentano et al., 2007;

Bertacchi et al., 2020; Del Pino et al., 2020; J. Feng et al., 2021; Flore et al., 2017; Lodato et

al., 2011; C. Zhou et al., 1999; C. Zhou, Tsai, & Tsai, 2001). *Nr2f2* plays a vital role in the

development of the amygdala, hypothalamus, and cerebellum (S. Feng et al., 2017; Kim,

Takamoto, Yan, Tsai, & Tsai, 2009; Tang, Rubenstein, Tsai, & Tsai, 2012). Nevertheless,

whether and how *Nr2f1* and/or *Nr2f2* genes regulate the differentiation and integration of

hippocampal morphogenesis, connectivity, and function is still largely unclear.

Here, our data show that *Nr2f1* and *Nr2f2* genes are differentially expressed along the

dorsoventral axis of the postnatal hippocampus. The loss of *Nr2f2* results in ectopic CA1 and

CA3 domains in the ventral hippocampus. In addition, the deficiency of *Nr2f1* leads to not only

dysplasia of the dorsal hippocampus but also failed specification and differentiation of the

dorsal CA1 pyramidal neuron lineage. Furthermore, the deletion of both genes in the RX$^{Cre/+}$;

Nr2f1$^{F/F}$; *Nr2f2*$^{F/F}$ double mutant mouse causes almost agenesis of the hippocampus,

accompanied by compromised specification of the CA1, CA3, and dentate gyrus (DG) domains.

The components of the trisynaptic circuit are abnormal in the corresponding single-gene or

double-gene mutant model. Moreover, *Nr2f* genes may cooperate to guarantee the appropriate

morphogenesis and function of the hippocampus by regulating the *Lhx5-Lhx2* axis.

Results

Differential expression profiles of *Nr2f1* and *Nr2f2* genes along the dorsoventral axis in

the developing and postnatal hippocampus

To investigate the functions of the *Nr2f1* and *Nr2f2* genes in the hippocampus,

immunofluorescence staining was first performed to examine their expression in wild-type

mice at postnatal month 1 (1M). In both the coronal and sagittal sections, NR2F1 exhibited a

septal/dorsal high-temporal/ventral low expression pattern along the hippocampus (Figure 1Aa,

d, c, f, g, j, m, i, l, and o), and its expression is highest in the dorsal CA1 region (Figure 1Aa,

d, c, f, j, and l), where place cells are mainly located (O'Keefe & Conway, 1978; O'Keefe &

Dostrovsky, 1971), and the dorsal DG, where there are adult neural stem cells (NSCs) (Gould

& Cameron, 1996). However, the expression of NR2F2 was high in the temporal/ventral

hippocampus but was barely detected in the septal/dorsal part of the hippocampus (Figure 1Ab,

c, e, f, h, i, k, l, n, and o). The dorsal-high NR2F1 and ventral-high NR2F2 expression profiles

were further verified in the postnatal hippocampi at 1M by western blotting assays (Figure 1Ap,

q). At embryonic day 10.5 (E10.5), NR2F1 was detected in the dorsal pallium (DP) laterally

and NR2F2 was expressed in the MP and CH medially (Figure 1—figure supplement 1Aa-b).

At E11.5 and E12.5, the expression of NR2F2 remained in the CH (Figure 1—figure

supplement 1Ac-d, Bb-c). Interestingly, NR2F1 and NR2F2 generated complementary

expression patterns in the hippocampal primordium with NR2F1 in the dorsal MP and NR2F2

in the ventral CH at E14.5 (Figure 1—figure supplement 1Ba-f). Additionally, septal/dorsal-

high NR2F1 and temporal/ventral-high NR2F2 expression patterns were observed at postnatal

day 0 (P0) (Figure 1—figure supplement 1Bg-l). The data above revealed that the differential

expression patterns of *Nr2f1* and *Nr2f2* genes were generated and maintained along the

dorsoventral axis in the early hippocampal primordium, the developing and postnatal

hippocampus, indicating that they could play distinct roles in the mediation of the

morphogenesis and functions of the hippocampus.

Next, to investigate the roles of *Nr2f* genes in the hippocampus, an RXCre mouse was used

to excise the expression of the *Nr2f1* and/or *Nr2f2* genes (Swindell et al., 2006; Tang et al.,

2012). The deletion efficiency of RXCre recombinase was verified by immunofluorescence

assays. Compared with control mice, either *Nr2f2* or *Nr2f1* could be excised in the postnatal

hippocampus of corresponding single-gene mutants at 1M (Figure 1—figure supplement 1Ca-

i). In addition, compared with control mice, both the *Nr2f1* and *Nr2f2* genes were almost

completely deleted in the hippocampal primordium in mutant mice at E14.5 (Figure 1—figure

supplement 1Cj-o). Since the LacZ expression serves as an indicator for the deletion of *Nr2f2*

(Swindell et al., 2006; Tang et al., 2012), we performed immunofluorescence staining with

antibodies against NR2F2 and LacZ on the sagittal sections of $RX^{Cre/+}$; $Nr2f2^{F/+}$ and $RX^{Cre/+}$;

$Nr2f2^{F/F}$ mice at E11.5. NR2F2 was readily detected at the hippocampal primordium of the

heterozygous mutant embryo at E11.5 (Figure 1—figure supplement 1Da, c, g); in contrast, the

expression of *Nr2f2* was significantly reduced in the homozygous mutant (Figure 1—figure

supplement 1Dd, f, j). In addition, compared with the heterozygous mutant embryo (Figure 1—

figure supplement 1Db-c, h), the LacZ signals clearly increased in the hippocampal

primordium of the homozygous mutant embryo at E11.5 (Figure 1—figure supplement 1De-f,

k), suggesting that RX-Cre recombinase can efficiently excise the *Nr2f2* gene in the

hippocampal primordium as early as E11.5. Intriguingly, we observed that the expression of

Nr2f1 increased in the caudal hippocampal primordium of the *Nr2f2* homozygous mutant

embryo at E11.5 (Figure 1—figure supplement 1Di, l), indicating that similar to the

observations in the early optic cup (Tang et al., 2010), *Nr2f1* and *Nr2f2* genes could be partially

compensate with each other in the developing hippocampal primordium. All the data above

show that RXCre recombinase could efficiently excise *Nr2f1* and/or *Nr2f2* in the early

developing and postnatal hippocampus.

The *Nr2f2* gene is required for the appropriate morphogenesis of the ventral

hippocampus but not of the dorsal hippocampus

Given that the *Nr2f2* gene is highly and specifically expressed in the postnatal ventral

hippocampus and the CH of the hippocampal primordium (Figure 1 and Figure 1—figure

supplement 1), we asked whether the *Nr2f2* gene is required for the appropriate morphogenesis

of the hippocampus, particularly the ventral hippocampus. To answer this question, we

conducted Nissl staining with samples from the *Nr2f2* single-gene (RX$^{Cre/+}$; *Nr2f2$^{F/F}$*) knockout

mouse model. In coronal sections, compared with the control at 1M, the septal hippocampus

was normal; unexpectedly, an ectopic CA-like region was observed medially in the temporal

hippocampus in the *Nr2f2* mutant, where the prospective posterior part of the medial

amygdaloid (MeP) nucleus was situated, indicated by the star (Figure 1Ba-f). The presence of

the ectopic CA-like region in the ventral but not dorsal hippocampus of the mutant was further

confirmed by the presence of the prospective MeP and amygdalohippocampal area (AHi) in

sagittal sections, as indicated by the star (Figure 1Bg-l). Furthermore, immunofluorescence

assays were performed to verify whether specific lineages of the hippocampus were altered

with sagittal sections, in which the subregions of both the dorsal and ventral hippocampus were

well displayed and distinguished. Ctip2 is a marker for CA1 pyramidal neurons and DG granule

neurons, and HuB is a marker for CA3 pyramidal cells (Sugiyama, Osumi, & Katsuyama,

2014). The dorsal hippocampus appeared normal in both the control and mutant mice at 1M

(Figure 1Ca-l). Nonetheless, compared with the observations in control mice, an ectopic HuB-

positive CA3 pyramidal neuron lineage, indicated by the star, and a duplicated Ctip2-positive

CA1 pyramidal neuron lineage, indicated by the arrowhead, were observed in the ventral

hippocampal area in the mutant (Figure 1Ca-f, m-r), revealing that there were ectopic CA1 and

CA3 lineages in the *Nr2f2* mutants.

Consistent with the previous report (Leid et al., 2004), the expression of Ctip2 was detected in

the amygdala including the AHi and posteromedial cortical amygdaloid nucleus (PMCo); in

addition, Ctip2 was also highly expressed in the dorsal part of the MeP (MePD) in the control

(Figure 1Cb, n).

Intriguingly, compared with the controls at 1M, there were ectopic CA domains in the mutant

ventral hippocampus with the expense of the Ctip2 positive AHi and MePD amygdaloid nuclei

(Figure 1Ce, q), indicated by the arrows. Clearly, all the data above suggested that the *Nr2f2*

gene is necessary to ensure the appropriate morphogenesis of the ventral hippocampus.

At early embryonic stages, *Nr2f2* was preferentially expressed in the CH (Figure 1—figure

supplement 1Ab, d, 1Bb, e), the organizer of the hippocampus, and at postnatal 1-month-old

(1M) stage, *Nr2f2* was also highly expressed in some amygdala nuclei such as the AHi and

medial amygdaloid nucleus, which are adjacent to the ventral/temporal hippocampus (Figure

1Ae, h, n) (Tang et al., 2012). We would like to investigate the correlation of the CH and/or

amygdala anlage with the duplicated ventral hippocampal domains in the *Nr2f2* mutant in detail

in our future study. The observations above suggest that *Nr2f2* is not only specifically

expressed in the ventral hippocampus but is also required for morphogenesis and probably the

function of the ventral hippocampus. Since the ventral hippocampus participates in the

regulation of emotion and stress, mutations in the *Nr2f2* gene lead to CHDs, and the formation

of the ventral hippocampus is disrupted in *Nr2f2* mutant mice at 1M, we wondered whether

CHD patients with *Nr2f2* mutations also exhibit symptoms associated with psychiatric

disorders such as depression, anxiety, or schizophrenia.

The *Nr2f1* gene is required for the specification and differentiation of the dorsal CA1 identity

Next, we asked whether the deletion of the *Nr2f1* gene by RX-Cre also affected the development of the hippocampus. Consistent with the previous finding in *Emx1$^{Cre/+}$*; *Nr2f1$^{F/F}$* mutant mice (Flore et al., 2017), it was the septal/dorsal hippocampus, not the temporal/ventral hippocampus, that was specifically shrunken in both coronal (Figure 2Aa-f) and sagittal sections (Figure 2Ag-i) of RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant mice. Then, we asked whether the loss of the *Nr2f1* gene caused abnormal specification and differentiation of hippocampal lineages. Compared with the control mice, *Nr2f1* mutant mice had fewer HuB-positive CA3 pyramidal neurons, as indicated by the star; intriguingly, Ctip2-positive CA1 pyramidal neurons failed to be detected, as indicated by the arrowhead, with Ctip2-positive DG granule neurons unaltered in the dorsal hippocampus (Figure 2Ba-l). The loss of the dorsal CA1 pyramidal neuron identity in mutant mice was further confirmed by Wfs1, another dorsal CA1 pyramidal neuron-specific marker (Takeda et al., 2001) (Figure 2Ca-r). Nonetheless, the HuB-positive and Ctip2-positive lineages were comparable in the ventral hippocampus between the control and mutant mice (Figure 2Ba-f, m-r), even though the low expression of NR2F1 was detected there. Indeed, *Nr2f1* is not only expressed at the highest level in the dorsal CA1 but is also required for the specification and differentiation of dorsal CA1 pyramidal neurons, among which place cells are essential for learning and memory (O'Keefe & Conway, 1978; O'Keefe & Dostrovsky, 1971).

Given that dysplasia of the dorsal hippocampus was generated in both the *Emx1-Cre* and RX-Cre models (Flore et al., 2017) (Figure 2Aa-l), we asked whether the development of dorsal CA1 pyramidal neurons was also abolished in *Emx1$^{Cre/+}$*; *Nr2f1$^{F/F}$* mutant mice. To answer this question, immunofluorescence staining was conducted first. Compared with that in the control mice, the proportions of either the Wfs1- or Ctip2-positive CA1 domain were reduced in the

mutant mice at 3M (Figure 2—figure supplement 1Aa-h), indicating that the differentiation of

the dorsal CA1 pyramidal neurons was also compromised in the *Emx1-Cre* model, although it

was less severe than that in the RX-Cre model. To make our findings more consistent with

previous studies, we further conducted experiments with the *Emx1-Cre* model. Afterward, to

investigate the fine structure of the dorsal CA1 pyramidal neurons, Golgi staining was

performed. Compared with those of control mice, the numbers of secondary dendrites and

branch points of both the apical and basal dendrites were significantly reduced in the dorsal

CA1 pyramidal neurons of mutant mice at 3M (Figure 2—figure supplement 1Ba-e). Then, the

dorsal hippocampus-related spatial learning and memory behavior test, the Morris water maze,

was performed (Vorhees & Williams, 2006). Consistent with a previous report (Flore et al.,

2017), spatial learning and memory function was significantly impaired in adult *Emx1*$^{Cre/+}$;

Nr2f1$^{F/F}$ mice, compared with the control mice (Figure 2—figure supplement 1C). The data

above suggest that *Nr2f1* is vital for the morphogenesis, lineage specification, and spatial

learning and memory of the dorsal hippocampus, and particularly, the compromised dorsal

CA1 lineage could contribute to the phenotypes associated with neurodevelopmental disorders,

including ID or ASD.

The *Nr2f1* and *Nr2f2* genes coordinate to ensure the genesis of the hippocampus

Given that the loss of either *Nr2f1* or *Nr2f2* leads to dysplasia of the dorsal or ventral

hippocampus, respectively, we asked whether these genes compensate for each other to

regulate the morphogenesis of the hippocampus. To answer this question, the RX$^{Cre/+}$; *Nr2f1*$^{F/F}$;

Nr2f2$^{F/F}$ double-mutant mouse was generated, and a few homozygous double-gene mutant

mice survived for approximately 3 weeks (3W). Nonetheless, the reason for the lethality of the

double-mutant mice is still unknown. Nissl staining data showed that compared with that of

control mice, the septal hippocampus was severely shrunken, as indicated by the star, and the

temporal hippocampus was barely observed in the double-mutant mouse brains (Figure 3Aa-

h). Unexpectedly, an ectopic nucleus was observed in the region of the prospective temporal

hippocampus, indicated by the arrowhead, in the double-mutant mice (Figure 3Ag-h). In

addition, compared with those of controls, the regions with HuB-positive CA3 pyramidal

neurons and Ctip2-positive or Prox1-positive DG granule neurons were diminished in the

double mutants; in particular, Ctip2-positive dorsal CA1 pyramidal neurons could not be

detected in the double mutants (Figure 3Ba-l). Furthermore, compared with the domains of

controls, no HuB-positive, Ctip2-positive, or Prox1-positive domains could be detected in the

prospective temporal hippocampus in the double mutants (Figure 3Ca-l). The results above

suggest that the *Nr2f1* and *Nr2f2* genes coordinate with each other to mediate the appropriate

morphogenesis of the entire hippocampus.

Nr2f genes and adult neurogenesis in the hippocampus

Given that the *Nr2f1* or *Nr2f2* gene was highly expressed in the dorsal or ventral DG,

respectively (Figure 1A), and that RX-Cre recombinase could efficiently delete either gene in

the DG (Figure 1—figure supplement 1Ca-i), we asked whether the loss of the *Nr2f1* or/and -

Nr2f2 gene in the DG would affect hippocampal adult neurogenesis. To answer this question,

the ventral DG, dorsal DG, and septal DG were chosen to perform immunofluorescence assays

in the *Nr2f2* mutant, *Nr2f1* mutant, and double mutant models, respectively. Adult NSCs in the

subgranular zone (SGZ) of the DG express both GFAP and Nestin, and newborn granule

neurons express Dcx (Gao, Arlotta, Macklis, & Chen, 2007). The numbers of NSCs and

newborn neurons in the SGZ of the DG were comparable between control mice and either the

Nr2f2 or *Nr2f1* mutant mice (Figure 3—figure supplement 1Aa-p, Ba-h); nonetheless,

compared with those of control mice, the numbers of the NSCs and newborn granule neurons

in the SGZ of the DG were reduced in the double mutants (Figure 3—figure supplement 1Aq-

x, Bi-l), and the reduction in both lineages was significant (Figure 3—figure supplement 1Ca,

b). The data above suggest that *Nr2f* genes may coordinate with each other to execute essential

functions for appropriate hippocampal adult neurogenesis in the DG.

Hippocampal trisynaptic connectivity was impaired in postnatal *Nr2f2* single-, *Nr2f1*

single-, and double-mutant mice at about 1M

Given that dysplasia of the hippocampus was observed in all three mouse models, we asked

whether the connectivity of the hippocampal trisynaptic circuit associated with the DG, CA3,

and CA1 regions (Amaral, 1993) was normal in these models. To answer this question, the

components of the trisynaptic circuit were characterized in the ventral hippocampus of *Nr2f2*

mutants, the dorsal hippocampus of *Nr2f1* mutants, and the septal hippocampus of double

mutants. Calretinin is a marker of mossy cells, Calbindin is a marker of mossy fibers, and

SMI312 is a marker of Schafer collaterals (Flore et al., 2017). Compared with those of controls

(Figure 4Aa, b, e, f, i, and j), the numbers of Calretinin-positive mossy cells were reduced,

Calbindin-positive mossy fibers were longer but thinner, and SMI312-positive Schafer

collaterals were thinner and discontinued in the ventral hippocampus of *Nr2f2* mutants at 1M

(Figure 4Ac, d, g, h, k, and l). In addition, similar to the previous report (Flore et al., 2017), the

numbers of Calretinin-positive mossy cells were decreased, Calbindin-positive mossy fibers

were shorter and thinner, and SMI312-positive Schafer collaterals were barely detected in the

dorsal hippocampus of *Nr2f1* mutants (Figure 4Ac, d, g, h, k, and l), compared with those of

controls at 1M (Figure 4Ba, b, e, f, i, and j). Moreover, compared with those of control mice

(Figure 4Ca, b, e, f, i, and j), the numbers of Calretinin-positive mossy cells were reduced; both

Calbindin-positive mossy fibers and SMI312-positive Schafer collaterals were barely detected

in the prospective septal hippocampus of the double mutants at 3W (Figure 4Cc, d, g, h, k, and

l). Clearly, without both *Nr2f* genes, the connectivity of the hippocampal trisynaptic circuit was

abolished more severely. The observations above revealed that the formation of the trisynaptic

circuit, which is one of the fundamental characteristics of hippocampal neurophysiology (Basu

& Siegelbaum, 2015), was abnormal in all three mouse models, indicating that both the

morphology and functions of the hippocampus are most likely compromised in the loss of the

Nr2f1 and/or *Nr2f2* gene.

The expression of several essential regulatory genes associated with early hippocampal

development was abnormal in double mutants

Given that the hippocampus was almost completely diminished in double mutants, we asked

how *Nr2f* genes participated in the regulation of the early morphogenesis of the hippocampus.

To answer this question, total RNA isolated from the whole telencephalons of control (n=5)

and double-mutant-(n=3) embryos at E11.5 was used to generate cDNA, and then real-time

quantitative PCR (RT-qPCR) assays were performed. As expected, compared with that of

control mice, the expression of *Nr2f1* and *Nr2f2* was reduced significantly in the double mutant

mice (Figure 5A). Then, we mainly focused on the intrinsic regulatory networks by analyzing

the expression profiles of two groups of transcription factor genes. The *Foxg1*, *Gli3*, *Lhx2*,

Otx1, Otx2, and *Pax6* genes, which are highly related to the early patterning of the dorsal

telencephalon (Hebert & Fishell, 2008), were in the first group; *Axin2*, *Emx1*, *Emx2*, *Lef1*, *Lhx5*,

and *Tcf4* genes, which are associated with early hippocampal development (Galceran et al.,

2000; Moore & Iulianella, 2021; Tole et al., 2000; Yoshida et al., 1997; Zhao et al., 1999),

were in the other group. The expression of the *Foxg1*, *Gli3*, *Lhx2*, *Otx1, Otx2*, and *Pax6* genes

was comparable between the controls and double mutants (Figure 5A), indicating that the early

patterning of the dorsal telencephalon is largely unaltered. Compared with that of control mice,

the expression of the *Axin2*, *Emx2*, *Lef1*, and *Tcf4* genes was normal in the double mutants;

interestingly, the expression of the *Emx1* and *Lhx5* transcripts was decreased significantly in

the double mutants at E11.5 compared to that in control mice (Figure 5A). Consistent with the

downregulated expression of *Lhx5* transcripts in the double mutant, the expression of the Lhx5

protein was reduced in the CH in the double mutants at E11.5; moreover, the number of Lhx5-

positive Cajal-Retzius cells decreased in the double mutant embryos at E11.5, E13.5 and E14.5

(Figure 5Ba-d, a'-d', a''-d'', i-l, i'-l', q-t, q'-t'). The expression of *Lhx2* was expanded

ventrally into the choroid plexus in the *Lhx5* null mutant mice (Zhao et al., 1999), indicating

that *Lhx5* could inhibit *Lhx2* expression locally. Consistent with RT-qPCR data, the expression

of Lhx2 was comparable between the control and double-mutant mice at E11.5 (Figure 5Be-h,

e'-h'). Interestingly, the expression of the Lhx2 protein was increased in the hippocampal

primordium in the *Nr2f* double-mutant mice at E13.5 and E14.5 (Figure 5Bm-p, m'-p', u-x, u'-

x'). The upregulation of Lhx2 expression is most likely associated with the reduced expression

of the *Lhx5* gene.

Next, we asked whether neural precursor cells (NPCs), intermediate progenitor cells (IPCs),

or newborn neurons were affected in the early development of the hippocampus in double-

mutant mice. Sox2 is a marker for NPCs, Tbr2 is a marker for IPCs, and NeuroD1 is a marker

for newborn neurons (Yu, Marchetto, & Gage, 2014). The expression of Sox2 in the

hippocampal regions was comparable between the control and double-mutant mice at E14.5

(Figure 5Ca-d, a'-d'), indicating that the generation of NPCs was normal. Nevertheless,

compared with the control embryos, the numbers of Tbr2-positive IPCs and NeuroD1-positive

newborn neurons were reduced in the double-mutant embryos (Figure 5Ce-l, e'-l'), and the

reduction was significant (Figure 5D). Our observations were consistent with previous findings

in *Lhx5* null mutant mice that the specification of the hippocampal NPCs was normal, but the

later differentiation event was abolished (Zhao et al., 1999). All the data above suggest that

Nr2f genes may cooperate to ensure the early morphogenesis of the hippocampus by regulating

the appropriate expression levels of *Lhx5* and *Lhx2* genes. Nevertheless, we could not exclude

other possibilities that *Nr2f* genes could also participate in the modulation of hippocampal

development through *Emx1* or other genes.

Discussion

In our present study, we observed dorsal-high NR2F1 and ventral-high NR2F2 expression

profiles in the postnatal hippocampus. The deletion of the *Nr2f2* gene led to duplicated CA1

and CA3 domains of the ventral hippocampus. The loss of *Nr2f1* resulted in the failed

specification and differentiation of the dorsal CA1 pyramidal neuron lineage with a diminished

dorsal hippocampus. Furthermore, the deficiency of both *Nr2f* genes caused atrophy of almost

the entire hippocampus, accompanied by compromised generation of the CA1, CA3, and DG

identities. In addition, the dorsal trisynaptic components, ventral trisynaptic components, or

entire trisynaptic components were abolished in the corresponding *Nr2f1* gene mutant, *Nr2f2*

gene mutant, or *Nr2f1/2* double-gene mutant mice. Moreover, *Nr2f* genes may cooperate to

ensure the appropriate morphogenesis and function of the hippocampus by regulating the *Lhx5-*

Lhx2 axis.

1. *Nr2f2* governs the distinct characteristics of the ventral hippocampus

Sixty years ago, the pioneering work of Milner and her colleagues discovered the essential role

of the hippocampus in declarative memory (Penfield & Milner, 1958; Scoville & Milner, 1957).

Recently, accumulating evidence has supported the Moser theory that the hippocampus is a

heterogeneous structure with distinct characteristics of gene expression, connectivity, and

function along its dorsoventral axis (Bast, 2007; Fanselow & Dong, 2010; Moser & Moser,

1998; Strange et al., 2014). The dorsal hippocampus marked in blue, in which gene expression

is similar to the neocortex, serves the "cold" cognitive function associated with declarative

memory and spatial navigation, and the ventral hippocampus marked in red, in which gene

expression is close to the hypothalamus and amygdala, corresponds to the "hot" affective states

related to emotion and anxiety (Figure 5—figure supplement 1). The ventral hippocampus

generates direct connectivity with the amygdala, hypothalamus, medial prefrontal cortex

(mPFC), and olfactory bulb (Cenquizca & Swanson, 2007; Hoover & Vertes, 2007; Kishi et

al., 2000; Pitkanen et al., 2000; Roberts et al., 2007). Nonetheless, thus far, the molecular and

cellular mechanism of how the morphogenesis, connectivity, and function of the ventral

hippocampus is achieved has been largely unclear.

Nr2f2, a nuclear receptor gene associated with heart disease(Al Turki et al., 2014) (High et al.,

2016), was highly and exclusively expressed in the ventral hippocampus in 1-month-old mice

and was expressed ventrally in the CH of the hippocampal primordium in mouse embryos

(Figure 1, Figure 1—figure supplement 1, Figure 5Ea), indicating that the *Nr2f2* gene may

participate in the regulation of the development and function of the ventral hippocampus. First,

deficiency of the *Nr2f2* gene led to the duplication of the CA1 and CA3 domains of the ventral

hippocampus but not the dorsal hippocampus, which was confirmed both morphologically and

molecularly (Figure 1). Second, the formation of the trisynaptic circuit was specifically

abolished in the ventral hippocampus of *Nr2f2* mutants (Figure 4), indicating that the

intrahippocampal circuit, information transfer, and function of the ventral hippocampus could

be compromised. Third, the ventral hippocampus generates neural circuits with the mPFC,

amygdala, nucleus accumbens, and hypothalamus, which are associated with

anxiety/behavioral inhibition, fear processing, pleasure/reward seeking, and the

neuroendocrine system, respectively (Anacker & Hen, 2017; Baik, 2020; Bryant & Barker,

2020; Cenquizca & Swanson, 2007; Herman et al., 2016; Kishi et al., 2000; O'Leary & Cryan,

2014; Pitkanen et al., 2000). These ventral hippocampal projections may be important for

processing information related to emotion and anxiety. Intriguingly, our previous studies

revealed that *Nr2f2* is required for the development of the hypothalamus, amygdala, and

olfactory bulb (S. Feng et al., 2017; Tang et al., 2012; X. Zhou et al., 2015), all of which

generate functional neural circuits with the ventral hippocampus (Fanselow & Dong, 2010).

Particularly, both the hypothalamus and amygdala are also diminished in RX$^{Cre/+}$; Nr2f2$^{F/F}$

mutant mice (S. Feng et al., 2017; Tang et al., 2012), indicating that their interconnectivities

with the ventral hippocampus are abnormal. Thus, all the findings above suggest that *Nr2f2* is

a novel and essential intrinsic regulator that controls the morphogenesis, connectivity, and

function of the ventral hippocampus.

Given that mutations of *Nr2f2* are highly associated with CHDs (Al Turki et al., 2014), the

expression of *Nr2f2* is also confined to the ventral hippocampus in human embryos (Alzu'bi et

al., 2017), and *Nr2f2* gene is required for the distinct characteristics of the ventral hippocampus

in mouse (Figure 1, Figure 4), we wondered whether CHD patients carrying mutations of *Nr2f2*

also display symptoms of psychiatric disorders, such as depression, anxiety, or schizophrenia,

related to the ventral hippocampus. In our future study, we would like to generate

hippocampus-specific or hippocampal subdomain-specific conditional knockout models to

dissect distinct roles of the *Nr2f2* gene in the hippocampus, particularly in the ventral

hippocampus, in detail.

2. The *Nr2f1* gene is required for the specification and differentiation of dorsal CA1

pyramidal neurons

The expression of *Nr2f1*, another orphan nuclear receptor gene associated with

neurodevelopmental disorders (Bertacchi et al., 2020; Bosch et al., 2014; Contesse et al., 2019),

is high in the dorsal MP of the hippocampal primordium and is higher in the dorsal

hippocampus than in the ventral hippocampus (Figure 1, Figure 1—figure supplement 1, Figure

5Ea) (Flore et al., 2017). Consistent with previous observations in *Emx1$^{Cre/+}$*; Nr2f1$^{F/F}$ mutant

mice (Flore et al., 2017), the dorsal hippocampus but not the ventral hippocampus was

specifically shrunken in RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mice (Figure 2, Figure 5—figure supplement 1).

Nr2f1 is expressed at the highest level in dorsal CA1 pyramidal neurons (Figure 1), indicating

that the *Nr2f1* gene may play a role in the specification and differentiation of dorsal CA1

pyramidal neurons. As expected, the expression of Ctip2 and Wfs1, two markers for dorsal

CA1 pyramidal neurons, could not be detected in the prospective dorsal CA1 domain in RX$^{Cre/+}$;

Nr2f1$^{F/F}$ mutant mice (Figure 2); furthermore, *Emx1$^{Cre/+}$*; *Nr2f1$^{F/F}$* mutant mice partially

phenocopied the compromised development of the dorsal CA1 lineage (Figure 2—figure

supplement 1). It seems that the spatiotemporal activity of RX-Cre recombinase is better or

broader than that of the Emx1-Cre recombinase during the critical period of the specification

of the dorsal CA1 pyramidal neuron identity. In addition, the Golgi staining assay revealed that

the development of the dendrites of the dorsal CA1 pyramidal neurons was abnormal (Figure

2—figure supplement 1). All the observations above indicate that the *Nr2f1* gene is not only

necessary for the morphogenesis of the dorsal hippocampus but is also required for the

specification and differentiation of the dorsal CA1 pyramidal neurons, among which there are

place cells. The identification of place cells fifty years ago was one of the most important

breakthroughs in understanding the role of the hippocampus in memory (O'Keefe &

Dostrovsky, 1971). Except for spatial information, place cells in the dorsal CA1 may also

encode nonspatial representations, such as time (Eichenbaum, 2017; Lisman et al., 2017).

Notably, 95% of patients carrying *Nr2f1* mutations are associated with ID. Here, our

observations support the notion that the *Nr2f1* gene is a novel intrinsic regulator that specifies

the dorsal CA1 pyramidal cell identity, which will benefit the understanding of both

neurophysiological functions of the hippocampus and the etiology of NDD including ID and

ASD.

3. *Nr2f1* and *Nr2f2* cooperate to ensure the appropriate morphogenesis of the hippocampus by regulating the *Lhx5-Lhx2* axis in mice

In wild-type mice, *Nr2f1* and *Nr2f2* genes generated complementary expression profiles in the embryonic hippocampal primordium with *Nr2f1* in the dorsal MP, marked in green; *Nr2f2* in the ventral CH, marked in red (Figure 5Ea, Figure 1—figure supplement 1); and in the postnatal hippocampus with high-NR2F1 expression in the dorsal, marked in green; and high-NR2F2 expression in the ventral, marked in red (Figure 5Ea, Figure 1). These findings indicated that *Nr2f* genes may coordinate to regulate hippocampal development. Indeed, as discussed above, the loss of either *Nr2f1* or *Nr2f2* only leads to dysplasia of the dorsal hippocampus (Flore et al., 2017) (Figure 2, Figure 2—figure supplement 1) or ventral hippocampus (Figure 1), respectively; intriguingly, while both genes are efficiently excised by RX-Cre in the hippocampal primordium (Figure 1—figure supplement 1), more severely shrunken hippocampi developed in the 3-week-old double knockout mice (Figure 3). The dosage-dependent severity of hippocampal abnormalities suggested that two nuclear receptor genes, *Nr2f1* and *Nr2f2* could cooperate with each other to execute an essential and intrinsic function in the development of the hippocampus.

It is known that both extrinsic signals and intrinsic factors participate in the regulation of the early development of the hippocampus. Notably, mutations of *Wnt3a* and *Lef1* eliminate the entire hippocampus (Galceran et al., 2000; S. M. Lee et al., 2000). Given that the expression of *Axin2*, *Lef1* and *Tcf4*, three Wnt-responsive transcription factor genes, was not altered in the *Nr2f* double mutant (Figure 5A), it is unlikely that abnormal Wnt signaling is the cause of the compromised hippocampus. *Lhx5* is specifically expressed in the hippocampal primordium and is required for the morphogenesis of the hippocampus (Zhao et al., 1999). *Lhx2* is necessary for hippocampal development by repressing cortical hem fate (Mangale et al., 2008; Monuki et al., 2001). Agenesis of the hippocampus is observed in either *Lhx5* or *Lhx2* null mutant mice,

and these genes particularly repress each other (Hebert & Fishell, 2008; Mangale et al., 2008;

Roy et al., 2014; Zhao et al., 1999), indicating that the proper expression levels of *Lhx5* and

Lhx2 genes are critical to maintain the appropriate development of the hippocampus (Figure

5Eb). The transcriptional and protein expression levels of *Lhx5* but not *Lhx2* were first reduced

in the hippocampal primordium of *Nr2f* double-mutant mice at E11.5; later, enhanced

expression of the Lhx2 protein was detected in the hippocampal primordium of double-mutant

mice at E13.5 and E14.5 (Figure 5A-B). Moreover, the number of Lhx5-positive Cajal-Retzius

cells was clearly reduced in the double mutant embryos at E11.5, E13.5 and E14.5; consistent

with the observations in the *Lhx5* null mutant (Li et al., 2021; Miquelajáuregui et al., 2010),

the generation of Sox2-positive hippocampal NPCs was not affected, but the development of

Tbr2-positive IPCs and NeuroD1-positive newborn neurons was abnormal in *Nr2f* double-

mutant mice (Figure 5). Thus, our findings reveal a novel intrinsic regulatory mechanism that

Nr2f1 and *Nr2f2*, two disease-associated nuclear receptor genes, may cooperate with each other

to ensure proper hippocampal morphogenesis by regulating the *Lhx5-Lhx2* axis. Intriguingly,

compared with the adult $Emx1^{Cre/+}$; $Nr2f1^{F/F}$ mutant mice, the hippocampus was much smaller

in the adult $Emx1^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-gene mutant mice; nevertheless, both the

dorsal and ventral hippocampus were readily detected in double-gene mutants with *Emx1-Cre*

(our unpublished observations). In addition, the discrepancy between the shrunken dorsal

hippocampus associated with the loss of *Nr2f1* and the duplicated CA domains of the ventral

hippocampus associated with the deficiency of *Nr2f2* suggested that the regulatory network

related to *Nr2f* genes during the early morphogenesis of the hippocampus could be much more

complicated than suspected and should be investigated in our future study.

4. *Nr2f* genes are imperative for the formation of the trisynaptic circuit

The hippocampus and entorhinal cortex (EC) are interconnected through various neural circuits

to mediate the flow of the information associated with declarative memory (Basu &

Siegelbaum, 2015). Both direct and indirect glutamatergic circuits are involved in the relay of

information from the EC to the hippocampal CA1, and the trisynaptic pathway is the most well-

characterized indirect circuit. The EC sends sensory signals from association cortices via the

perforant path to the DG, then the DG granule cells send excitatory mossy fiber projections to

CA3 pyramidal neurons, and CA3 pyramidal neurons project to CA1 via the Schaffer

collaterals (H. Lee, GoodSmith, & Knierim, 2020). Consistent with the high expression of

NR2F1 in the dorsal hippocampus and NR2F2 in the ventral hippocampus (Figure 1), the dorsal

trisynaptic circuit is specifically damaged in *Nr2f1* mutants, as is the ventral trisynaptic circuit

in *Nr2f2* mutant mice. Moreover, the hippocampal trisynaptic circuit was almost completely

absent in the double mutants (Figure 4, and our unpublished observations). The information

transfer associated with the trisynaptic circuits should be abolished particularly in the dorsal

and/or the ventral hippocampus in the above corresponding genetic mouse models.

Interestingly, *Nr2f1* is also required to specify the medial EC cell fate (J. Feng et al., 2021).

Therefore, the impaired formation and function of trisynaptic circuits could be caused by the

abnormal development of CA1, CA3, DG or EC lineages. Nonetheless, given that newborn

granule neurons are continuously generated in the adult DG to integrate into the existing neural

circuits essential for declarative memory (Toda, Parylak, Linker, & Gage, 2019; Tuncdemir,

Lacefield, & Hen, 2019) and that hippocampal adult neurogenesis was severely compromised

in the *Nr2f* double mutant (Figure 3—figure supplement 1), we could not exclude the

possibility that impaired adult neurogenesis may also contribute to the malformation and

impaired function of the trisynaptic pathway in double mutants. We would like to investigate

what sort of synaptic circuitry is compromised either physiologically or morphologically in the

trisynaptic circuit of individual animal model in detail in future studies.

The hippocampus is heterogeneous along its dorsoventral axis, and either the dorsal or ventral

hippocampus generates unique and distinguishable characteristics of gene expression and

connectivity, which enable the hippocampus to execute an integrative function from the

encoding and retrieval of certain declarative memory to adaptive behaviors. Lesions of the

dorsal hippocampus, which are essential for the cognitive process of learning and memory,

lead to amnesia and ID; while damage to the ventral hippocampus, which is central for emotion

and affection, is highly associated with psychiatric disorders including depression, anxiety, and

schizophrenia. Our findings in this study reveal novel intrinsic mechanisms by which two

nuclear receptor genes, *Nr2f1* and *Nr2f2*, which are associated with NDD or CHD, converge

to govern the differentiation and integration of the hippocampus along the dorsoventral axis

morphologically and functionally. Furthermore, our present study provides novel genetic

model systems to investigate the crosstalk among the hippocampal complex in gene expression,

morphogenesis, cell fate specification and differentiation, connectivity, functions of

learning/memory and emotion/anxiety, adaptive behaviors, and the etiology of neurological

diseases. Nevertheless, many enigmas, such as whether and how the abnormalities of either the

dorsal or ventral hippocampus affect the characteristics of the other, remain unsolved. In

addition to the excitatory lineages and circuits, interneurons and inhibitory circuits play vital

roles in maintaining the plasticity and functions of the hippocampus. We also wonder whether

and how defects in interneurons and inhibitory circuits could contribute to the compromised

morphogenesis, connectivity, and functions of the hippocampus and the etiology of psychiatric

and neurological conditions including ID, ASD, depression, anxiety, and schizophrenia.

552
Materials and Methods

Animals

Nr2f1-floxed mice, *Nr2f2-floxed* mice, *Emx1^Cre* mice and RX^Cre mice (Swindell et al., 2006)

(PMID: 16850473) used in the study were of the C57B6/129 mixed background. The noon of

vaginal plug day was set as the embryonic day 0.5 (E0.5). Only male mice at age of 10 weeks

or older were used in the Morris water maze. For other experiments, both male and female

mice were used. All animal protocols were approved by the Institutional Animal Care and Use

Committee (IACUC) at the Shanghai Institute of Biochemistry and Cell Biology, Chinese

Academy of Sciences (Protocols: SIBCB-NAF-14-001-S308-001). All methods were

performed in accordance with the relevant guidelines and regulations. Only the littermates were

used for the comparison.

Nissl staining

We used xylene to dewax paraffin sections, followed by rinsing with 100%, 95%, and 70%

ethanol. The slides were stained in 0.1% Cresyl Violet solution for 25 mins. Then the sections

were washed quickly in the water and differentiated in 95% ethanol. We used 100% ethanol to

dehydrate the slides, followed by rinsing with the xylene solution. Finally, the neutral resin

medium was used to mount the slides.

Immunohistochemical (IHC) staining

The paraffin sections were dewaxed and rehydrated as described above for Nissl staining. The

slides were boiled in 1×antigen retrieval solution (DAKO) under microwave conditions for 15

min. After cooled to room temperature (RT), the slides were incubated with 3% H_2O_2 for 30

min. Then, the slides were treated with blocking buffer for 60 min at RT and then incubated

with the primary antibody in the hybridization buffer (10×diluted blocking buffer) overnight

(O/N) at 4 °C. The next day, the tyramide signal amplification kit (TSA) (Invitrogen) was used

according to the manufacturer's protocol. After being incubated with 1%TSA blocking buffer,

the sections were treated with a biotinylated secondary antibody for 60 min at RT. After being

washed with 1×PBS three times, the slides were incubated with 1×HRP-conjugated

streptavidin for 1 h. Next, the tyramide working solution was prepared, including the 0.15‰

H_2O_2 in distilled water, the 100×diluted tyramide substrate solution (tyramide-488 or tyramide-

594), and the amplification buffer. The sections were incubated with the working solution for

10 min. Then the slides were counterstained with DAPI and mounted with the antifade

mounting medium (Southern Biotech) (S. Feng et al., 2017; Zhang et al., 2020). Finally, the

sections were observed and images were captured with a digital fluorescence microscope

(Zeiss).

The following primary antibodies were used in the study: mouse anti-NR2F1 (1:1000, R&D,

Cat # PP-H8132-00), mouse anti-NR2F2 (1:2000, R&D, Cat # PP-H7147-00), rabbit anti-

NR2F2 (1:2000, a gift from Dr. Zhenzhong Xu, Zhejiang University, China), rabbit anti-HuB

(1:500, Abcam, Cat # ab204991), rat anti-Ctip2 (1:500, Abcam, Cat # ab18465), rabbit anti-

Wfs1 (1:500, ProteinTech, Cat # 11558-1-AP), goat anti-Prox1 (1:500, R&D, Cat # AF2727),

rabbit anti-Calretinin (1:500, Sigma, Cat # C7479), rabbit anti-Calbindin (1:500, Swant, Cat #

CB38), mouse anti-SMI312 (1:200, Covance, Cat # SMI-312R), rabbit anti-Sox2 (1:500,

Affinity BioReagents, Cat # PA1-16968), rat anti-Tbr2 (1:500, Thermo Fisher, Cat # 12-4875-

82), goat anti-NeuroD1 (1:200, Santa Cruz, Cat # sc-1084), goat anti-Lhx2 (1:200, Santa Cruz,

Cat # sc-19344), goat anti-Lhx5 (1:200, R&D, Cat # AF6290), goat anti-β-galactosidase (LacZ)

(1:400, Biogenesis, Cat # 4600-1409), mouse anti-GFAP (1:500, Sigma, Cat # G3893), rabbit

anti-Nestin (1:200, Santa Cruz, Cat # sc-20978), goat anti-Dcx (1:500, Santa Cruz, Cat # sc-

8066). The following secondary antibodies were used in the study: donkey anti-mouse IgG

biotin-conjugated (1:400, JacksonImmuno, Cat # 715-065-150), donkey anti-rabbit IgG biotin-

conjugated (1:400, JacksonImmuno, Cat # 711-065-152), donkey anti-goat IgG biotin-

conjugated (1:400, JacksonImmuno, Cat # 705-066-147), donkey anti-rat IgG biotin-

conjugated (1:400, JacksonImmuno, Cat # 712-065-150).

Western blotting

We homogenized the isolated dorsal and ventral hippocampus tissues from 1-month-old mice

respectively in the RIPA buffer (Applygen) with protease inhibitor cocktail (Sigma) and

phosphatase inhibitors (Invitrogen) and then centrifuged at the speed of 12,000 rpm for 30 mins.

We collected the supernatants and analyzed the total concentrations by the BCA kit (Applygen).

Gradient SDS-PAGE gels were used to separate the same amounts of protein sample (40

μg/lane), and then the proteins were transferred to the PVDF membranes (Millipore). After

being blocked by the 3% BSA (Sigma) for 2h, the membranes that contained proteins were

incubated by primary antibodies at 4 °C O/N. The membranes were rinsed with 1×PBST three

times for 10 mins and then treated with biotinylated secondary antibodies for 2h at RT. After

washing with 1×PBST, membranes were treated with the HRP-conjugated streptavidin for 1h

at RT. Finally, we used the chemiluminescence detection system (Tanon) to detect the bands.

The density of the protein band was analyzed by the software Image J.

The primary antibodies were used in the experiment as below: mouse anti-NR2F1 (1:2000,

R&D, Cat # PP-H8132-00), rabbit anti-NR2F2 (1:3000, a gift from Dr. Zhenzhong Xu,

Zhejiang University, China), mouse anti-GAPDH (1:1000, Santa Cruz, Cat # sc-32233). The

following secondary antibodies were applied in the study, including goat anti-mouse IgG

biotin-conjugated (1:1000, KPL, Cat # 16-18-06), goat anti-rabbit IgG biotin-conjugated

(1:1000, KPL, Cat # 16-15-06).

Golgi staining

Deep anesthesia was performed before sacrificing the control and mutant mice, and the brains

were immediately isolated. The FD rapid GolgiStain kit (FD NeuroTech) was used to process

the brain tissue samples, which were immersed in an equal volume of immersion solution

mixed with solutions A and B and stored in the dark for two weeks at RT. At least 5mL of

immersion solution was used for each cubic meter of the tissue. To achieve the best results, the

container of tissue was gently swirled from side to side twice a week during the incubating

period. Afterwards, the brain tissue was transferred to solution C in the dark at RT for at least

72 hours (up to 1 week). Finally, the tissues were cut into 100 μm thick slices with a cryostat

at -20°C to -22°C and transferred to gelatin-coated microscope slides containing solution C

using a sample retriever. The slices were dried naturally at RT. The concrete staining procedure

of the kit was followed using the manufactory's protocol. Then, the sections were rinsed twice

with double-distilled water for 4 minutes each time. The slices were placed in a mixture of one

volume of solution D, one volume of solution E, and two volumes of double distilled water for

10 minutes and were rinsed twice with distilled water for 4 minutes each time. The sections

were later dehydrated in 50%, 75%, and 95% ethanol for 4 minutes respectively. Next, the

slices were dehydrated in 100% ethanol 4 times for 4 minutes each time. Finally, the sections

were cleared in xylene and mounted with a neutral resin medium.

Morris water maze

By recording the time spent by the mice swimming in the water tank and finding the escape

platform hidden underwater, and the swimming trajectory, the Morris water maze test can

objectively reflect the spatial learning and memory ability of the mice. The test was divided

into the control and mutant group with at least eight mice in each group. We poured tap water

into the water maze tank and added an appropriate amount of well-mixed, milky white food

dye. The height of the liquid level was about 1 cm higher than the escape platform, and the

water temperature was kept at about 25±1C°. At the same time, four markers of different shapes

were pasted on the four directions of the inner wall above the water tank to distinguish different

directions. The Morris water maze test was divided into the training phase and the probe trial

phase. The training phase lasted for 6 days, 4 times a day, and the interval between each training

was about 30 minutes. During training, the mice were placed into the tank from the entry points

of four different quadrants facing the inner wall, and their latency was recorded from the time

they entered the water to the time they found a hidden underwater platform and stood on it.

After the mouse found the platform, we let it stay on the platform for 10 seconds before

removing it. If the mouse failed to discover the platform 60 seconds after entering the water, it

was guided to find the platform and left to stay for 10 seconds. Each mouse was placed into

the water tank from four water entry points and recorded as one training session. The probe

trial was carried out on the seventh day, and the underwater platform was removed. Each

experimental mouse was put into the water tank at the same water entry point and allowed to

move for 60 seconds. The time that each mouse spent in the quadrant, where the platform was

originally placed, was recorded.

RNA isolation and quantitative real-time PCR

Total RNAs were prepared from the whole telencephalon of the control (n=5) and double

mutant (n=3) mice at E11.5 respectively, with the TRIzol Reagent (Invitrogen) by following

the manufactory's protocol. Reverse-transcription PCR and real-time quantitative PCR assays

were performed as described previously (Tang et al., 2012). A student's t-test was used to

compare the means of the relative mRNA levels between the control group and mutant group.

Primer sequences are as follows:

Axin2-f, 5'-*CTGCTGGTCAGGCAGGAG*-3', *Axin2-r*, 5'-*TGCCAGTTTCTTTGGCTCTT*-3';

Nr2f1-f, 5'-*CAAAGCCATCGTGCTATTCA*-3', *Nr2f1-r*, 5'-*CCTGCAGGCTTTCGATGT*-3';

Nr2f2-f, 5'-*CCTCAAAGTGGGCATGAGAC*-3', *Nr2f2-r*, 5'-*TGGGTAGGCTGGGTAGGAG*-

3'; *Emx1-f*, 5'-*CTCTCCGAGACGCAGGTG*-3', *Emx1-r*, 5'-*CTCAGACTCCGGCCCTTC*-3';

Emx2-f, 5'-*CACGCTTTTGAGAAGAACCA*-3', *Emx2-r*, 5'-*GTTCTCCGGTTCTGAAACCA*-3';

Foxg1-f, 5'-*GAAGGCCTCCACAGAACG*-3', *Foxg1-r*, 5'-*GGCAAGGCATGTAGCAAAAG*-3';

Gli3-f, 5'-*TGATCCATCTCCTATTCCTCCA*-3', *Gli3-r*, 5'-*TCTGGATACGTCGGGCTACT*-3';

Lef1-f, 5'-*TCCTGAAATCCCCACCTTCT*-3', *Lef1-r*, 5'-*TGGGATAAACAGGCTGACCT*-3';

Lhx2-f, 5'-*CAGCTTGCGCAAAAGACC*-3', *Lhx2-r*, 5'-*TAAAAGGTTGCGCCTGAACT*-3';

Lhx5-f, 5'-*TGTGCAATAAGCAGCTATCCA*-3', *Lhx5-r*, 5'-*CAAACTGCGGTCCGTACA*-3';

Otx1-f, 5'-*CCAGAGTCCAGAGTCCAGGT*-3', *Otx1-r*, 5'-*CCGGGTTTTCGTTCCATT*-3';

Otx2-f, 5'-*GGTATGGACTTGCTGCATCC*-3', *Otx2-r*, 5'-*CGAGCTGTGCCCTAGTAAATG*-3';

Pax6-f, 5'-*GTTCCCTGTCCTGTGGACTC*-3', *Pax6-r*, 5'-*ACCGCCCTTGGTTAAAGTCT*-3';

Tcf4-f, 5'-*AAATGGCCACTGCTTGATGT*-3', *Tcf4-r*, 5'-*GCACCACCGGTACTTGTTC*-3'.

Quantification and statistical analysis

The number of specified immunofluorescent marker-positive cells was assessed by Image J

Cell Counter in full image fields. Three brain sections per mouse were counted for each index.

GraphPad Prism 7.0 (GraphPad) was used to perform statistical analysis. The data analysis

used one-way analysis of variance (ANOVA), Dunnett's or Tukey's post hoc tests, and

student's unpaired t-test. The data were expressed as the mean ± SEM. The data obtained from

at least three independent replicates were used for statistical analysis. $P < 0.05$ was considered

the significant statistical difference.

Acknowledgements

We thank Ms. Emerald Tang for her assistance with the manuscript. This work was supported

by National Natural Science Foundation of China (31671508) and Guangdong Provincial Basic

and Applied Basic Research Fund (2021A1515011299) to K.T; and in part by the National Key

Basic Research and Development Program of China (2018YFA0108500, 2019YFA0801402,

2018YFA0800100, 2018YFA0108000, 2018YFA0107200, 2017YFA0102700), "Strategic

Priority Research Program" of the Chinese Academy of Sciences, Grant No. (XDA16020501,

XDA16020404) to N. J.

Data availability

710   Numerical data are available in the manuscript and supporting files.

Competing interests

713   The authors declare no competing interests.

References

716
Al Turki, S., Manickaraj, A. K., Mercer, C. L., Gerety, S. S., Hitz, M. P., Lindsay, S., . . .
Hurles, M. E. (2014). Rare variants in NR2F2 cause congenital heart defects in
humans. *Am J Hum Genet, 94*(4), 574-585. doi:10.1016/j.ajhg.2014.03.007
Alzu'bi, A., Lindsay, S. J., Harkin, L. F., McIntyre, J., Lisgo, S. N., & Clowry, G. J. (2017).
The Transcription Factors COUP-TFI and COUP-TFII have Distinct Roles in
Arealisation and GABAergic Interneuron Specification in the Early Human Fetal
Telencephalon. *Cereb Cortex, 27*(10), 4971-4987. doi:10.1093/cercor/bhx185
Amaral, D. G. (1993). Emerging principles of intrinsic hippocampal organization. *Current*
Opinion in Neurobiology, 3(2), 225-229. doi:https://doi.org/10.1016/0959-
4388(93)90214-J
Anacker, C., & Hen, R. (2017). Adult hippocampal neurogenesis and cognitive flexibility -
linking memory and mood. *Nat Rev Neurosci, 18*(6), 335-346.
doi:10.1038/nrn.2017.45
Armentano, M., Chou, S. J., Tomassy, G. S., Leingärtner, A., O'Leary, D. D., & Studer, M.
(2007). COUP-TFI regulates the balance of cortical patterning between frontal/motor
and sensory areas. *Nat Neurosci, 10*(10), 1277-1286. doi:10.1038/nn1958
Baik, J. H. (2020). Stress and the dopaminergic reward system. *Exp Mol Med, 52*(12), 1879-
1890. doi:10.1038/s12276-020-00532-4
Bast, T. (2007). Toward an integrative perspective on hippocampal function: from the rapid
encoding of experience to adaptive behavior. *Rev Neurosci, 18*(3-4), 253-281.
doi:10.1515/revneuro.2007.18.3-4.253
Basu, J., & Siegelbaum, S. A. (2015). The Corticohippocampal Circuit, Synaptic Plasticity,
and Memory. *Cold Spring Harb Perspect Biol, 7*(11).
doi:10.1101/cshperspect.a021733

Bertacchi, M., Romano, A. L., Loubat, A., Tran Mau-Them, F., Willems, M., Faivre, L., . . . Studer, M. (2020). NR2F1 regulates regional progenitor dynamics in the mouse neocortex and cortical gyrification in BBSOAS patients. *EMBO J, 39*(13), e104163. doi:10.15252/embj.2019104163

Bliss, T. V., & Lomo, T. (1973). Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J Physiol, 232*(2), 331-356. doi:10.1113/jphysiol.1973.sp010273

Bosch, D. G., Boonstra, F. N., Gonzaga-Jauregui, C., Xu, M., de Ligt, J., Jhangiani, S., . . . Schaaf, C. P. (2014). NR2F1 mutations cause optic atrophy with intellectual disability. *Am J Hum Genet, 94*(2), 303-309. doi:10.1016/j.ajhg.2014.01.002

Bryant, K. G., & Barker, J. M. (2020). Arbitration of Approach-Avoidance Conflict by Ventral Hippocampus. *Front Neurosci, 14*, 615337. doi:10.3389/fnins.2020.615337

Cenquizca, L. A., & Swanson, L. W. (2007). Spatial organization of direct hippocampal field CA1 axonal projections to the rest of the cerebral cortex. *Brain Res Rev, 56*(1), 1-26. doi:10.1016/j.brainresrev.2007.05.002

Contesse, T., Ayrault, M., Mantegazza, M., Studer, M., & Deschaux, O. (2019). Hyperactive and anxiolytic-like behaviors result from loss of COUP-TFI/Nr2f1 in the mouse cortex. *Genes Brain Behav, 18*(7), e12556. doi:10.1111/gbb.12556

Del Pino, I., Tocco, C., Magrinelli, E., Marcantoni, A., Ferraguto, C., Tomagra, G., . . . Studer, M. (2020). COUP-TFI/Nr2f1 Orchestrates Intrinsic Neuronal Activity during Development of the Somatosensory Cortex. *Cereb Cortex, 30*(11), 5667-5685. doi:10.1093/cercor/bhaa137

Eichenbaum, H. (2017). On the Integration of Space, Time, and Memory. *Neuron, 95*(5), 1007-1018. doi:10.1016/j.neuron.2017.06.036

Eichenbaum, H., & Cohen, N. J. (2014). Can we reconcile the declarative memory and spatial navigation views on hippocampal function? *Neuron, 83*(4), 764-770. doi:10.1016/j.neuron.2014.07.032

Fanselow, M. S., & Dong, H. W. (2010). Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron, 65*(1), 7-19. doi:10.1016/j.neuron.2009.11.031

Feng, J., Hsu, W. H., Patterson, D., Tseng, C. S., Hsing, H. W., Zhuang, Z. H., . . . Chou, S. J. (2021). COUP-TFI specifies the medial entorhinal cortex identity and induces differential cell adhesion to determine the integrity of its boundary with neocortex. *Sci Adv, 7*(27). doi:10.1126/sciadv.abf6808

Feng, S., Xing, C., Shen, T., Qiao, Y., Wang, R., Chen, J., . . . Tang, K. (2017). Abnormal Paraventricular Nucleus of Hypothalamus and Growth Retardation Associated with Loss of Nuclear Receptor Gene COUP-TFII. *Sci Rep, 7*(1), 5282. doi:10.1038/s41598-017-05682-6

Flore, G., Di Ruberto, G., Parisot, J., Sannino, S., Russo, F., Illingworth, E. A., . . . De Leonibus, E. (2017). Gradient COUP-TFI Expression Is Required for Functional Organization of the Hippocampal Septo-Temporal Longitudinal Axis. *Cereb Cortex, 27*(2), 1629-1643. doi:10.1093/cercor/bhv336

Galceran, J., Miyashita-Lin, E. M., Devaney, E., Rubenstein, J. L., & Grosschedl, R. (2000). Hippocampus development and generation of dentate gyrus granule cells is regulated by LEF1. *Development, 127*(3), 469-482. doi:10.1242/dev.127.3.469

Gao, X., Arlotta, P., Macklis, J. D., & Chen, J. (2007). Conditional knock-out of beta-catenin in postnatal-born dentate gyrus granule neurons results in dendritic malformation. *J Neurosci, 27*(52), 14317-14325. doi:10.1523/jneurosci.3206-07.2007

Gould, E., & Cameron, H. A. (1996). Regulation of neuronal birth, migration and death in the rat dentate gyrus. *Dev Neurosci, 18*(1-2), 22-35. doi:10.1159/000111392

Hebert, J. M., & Fishell, G. (2008). The genetics of early telencephalon patterning: some
assembly required. *Nat Rev Neurosci, 9*(9), 678-685. doi:10.1038/nrn2463
Herman, J. P., McKlveen, J. M., Ghosal, S., Kopp, B., Wulsin, A., Makinson, R., . . . Myers,
B. (2016). Regulation of the Hypothalamic-Pituitary-Adrenocortical Stress Response.
Compr Physiol, 6(2), 603-621. doi:10.1002/cphy.c150015
High, F. A., Bhayani, P., Wilson, J. M., Bult, C. J., Donahoe, P. K., & Longoni, M. (2016).
De novo frameshift mutation in COUP-TFII (NR2F2) in human congenital
diaphragmatic hernia. *Am J Med Genet A, 170*(9), 2457-2461.
doi:10.1002/ajmg.a.37830
Hoover, W. B., & Vertes, R. P. (2007). Anatomical analysis of afferent projections to the
medial prefrontal cortex in the rat. *Brain Struct Funct, 212*(2), 149-179.
doi:10.1007/s00429-007-0150-4
Kandel, E. R., Dudai, Y., & Mayford, M. R. (2014). The molecular and systems biology of
memory. *Cell, 157*(1), 163-186. doi:10.1016/j.cell.2014.03.001
Kim, B. J., Takamoto, N., Yan, J., Tsai, S. Y., & Tsai, M. J. (2009). Chicken Ovalbumin
Upstream Promoter-Transcription Factor II (COUP-TFII) regulates growth and
patterning of the postnatal mouse cerebellum. *Dev Biol, 326*(2), 378-391.
doi:10.1016/j.ydbio.2008.11.001
Kishi, T., Tsumori, T., Ono, K., Yokota, S., Ishino, H., & Yasui, Y. (2000). Topographical
organization of projections from the subiculum to the hypothalamus in the rat. *J
Comp Neurol, 419*(2), 205-222. doi:10.1002/(sici)1096-
9861(20000403)419:2<205::aid-cne5>3.0.co;2-0
Lee, H., GoodSmith, D., & Knierim, J. J. (2020). Parallel processing streams in the
hippocampus. *Curr Opin Neurobiol, 64*, 127-134. doi:10.1016/j.conb.2020.03.004
Lee, S. M., Tole, S., Grove, E., & McMahon, A. P. (2000). A local Wnt-3a signal is required
for development of the mammalian hippocampus. *Development, 127*(3), 457-467.
doi:10.1242/dev.127.3.457
Leid, M., Ishmael, J. E., Avram, D., Shepherd, D., Fraulob, V., & Dolle, P. (2004). CTIP1
and CTIP2 are differentially expressed during mouse embryogenesis. *Gene Expr
Patterns, 4*(6), 733-739. doi:10.1016/j.modgep.2004.03.009
Li, J., Sun, L., Peng, X. L., Yu, X. M., Qi, S. J., Lu, Z. J., . . . Shen, Q. (2021). Integrative
genomic analysis of early neurogenesis reveals a temporal genetic program for
differentiation and specification of preplate and Cajal-Retzius neurons. *PLoS Genet,
17*(3), e1009355. doi:10.1371/journal.pgen.1009355
Lisman, J., Buzsaki, G., Eichenbaum, H., Nadel, L., Ranganath, C., & Redish, A. D. (2017).
Viewpoints: how the hippocampus contributes to memory, navigation and cognition.
Nat Neurosci, 20(11), 1434-1447. doi:10.1038/nn.4661
Lodato, S., Tomassy, G. S., De Leonibus, E., Uzcategui, Y. G., Andolfi, G., Armentano,
M., . . . Studer, M. (2011). Loss of COUP-TFI alters the balance between caudal
ganglionic eminence- and medial ganglionic eminence-derived cortical interneurons
and results in resistance to epilepsy. *J Neurosci, 31*(12), 4650-4662.
doi:10.1523/jneurosci.6580-10.2011
Mangale, V. S., Hirokawa, K. E., Satyaki, P. R., Gokulchandran, N., Chikbire, S.,
Subramanian, L., . . . Monuki, E. S. (2008). Lhx2 selector activity specifies cortical
identity and suppresses hippocampal organizer fate. *Science, 319*(5861), 304-309.
doi:10.1126/science.1151695
Miquelajáuregui, A., Varela-Echavarría, A., Ceci, M. L., García-Moreno, F., Ricaño, I.,
Hoang, K., . . . Zhao, Y. (2010). LIM-homeobox gene Lhx5 is required for normal
development of Cajal-Retzius cells. *J Neurosci, 30*(31), 10551-10562.
doi:10.1523/jneurosci.5563-09.2010

Monuki, E. S., Porter, F. D., & Walsh, C. A. (2001). Patterning of the dorsal telencephalon
and cerebral cortex by a roof plate-Lhx2 pathway. *Neuron, 32*(4), 591-604.
doi:10.1016/s0896-6273(01)00504-9
Moore, S. A., & Iulianella, A. (2021). Development of the mammalian cortical hem and its
derivatives: the choroid plexus, Cajal-Retzius cells and hippocampus. *Open Biol,*
11(5), 210042. doi:10.1098/rsob.210042
Moser, M. B., & Moser, E. I. (1998). Functional differentiation in the hippocampus.
Hippocampus, 8(6), 608-619. doi:10.1002/(SICI)1098-1063(1998)8:6<608::AID-
HIPO3>3.0.CO;2-7
O'Keefe, J., & Conway, D. H. (1978). Hippocampal place units in the freely moving rat: why
they fire where they fire. *Exp Brain Res, 31*(4), 573-590. doi:10.1007/BF00239813
O'Keefe, J., & Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary
evidence from unit activity in the freely-moving rat. *Brain Res, 34*(1), 171-175.
doi:10.1016/0006-8993(71)90358-1
O'Leary, O. F., & Cryan, J. F. (2014). A ventral view on antidepressant action: roles for adult
hippocampal neurogenesis along the dorsoventral axis. *Trends Pharmacol Sci, 35*(12),
675-687. doi:10.1016/j.tips.2014.09.011
Penfield, W., & Milner, B. (1958). Memory deficit produced by bilateral lesions in the
hippocampal zone. *AMA Arch Neurol Psychiatry, 79*(5), 475-497.
doi:10.1001/archneurpsyc.1958.02340050003001
Pitkanen, A., Pikkarainen, M., Nurminen, N., & Ylinen, A. (2000). Reciprocal connections
between the amygdala and the hippocampal formation, perirhinal cortex, and
postrhinal cortex in rat. A review. *Ann N Y Acad Sci, 911*, 369-391.
doi:10.1111/j.1749-6632.2000.tb06738.x
Porter, F. D., Drago, J., Xu, Y., Cheema, S. S., Wassif, C., Huang, S. P., . . . Westphal, H.
(1997). Lhx2, a LIM homeobox gene, is required for eye, forebrain, and definitive
erythrocyte development. *Development, 124*(15), 2935-2944.
doi:10.1242/dev.124.15.2935
Roberts, A. C., Tomic, D. L., Parkinson, C. H., Roeling, T. A., Cutter, D. J., Robbins, T. W.,
& Everitt, B. J. (2007). Forebrain connectivity of the prefrontal cortex in the
marmoset monkey (Callithrix jacchus): an anterograde and retrograde tract-tracing
study. *J Comp Neurol, 502*(1), 86-112. doi:10.1002/cne.21300
Roy, A., Gonzalez-Gomez, M., Pierani, A., Meyer, G., & Tole, S. (2014). Lhx2 regulates the
development of the forebrain hem system. *Cereb Cortex, 24*(5), 1361-1372.
doi:10.1093/cercor/bhs421
Schuurmans, C., & Guillemot, F. (2002). Molecular mechanisms underlying cell fate
specification in the developing telencephalon. *Curr Opin Neurobiol, 12*(1), 26-34.
doi:10.1016/s0959-4388(02)00286-6
Scoville, W. B., & Milner, B. (1957). Loss of recent memory after bilateral hippocampal
lesions. *J Neurol Neurosurg Psychiatry, 20*(1), 11-21. doi:10.1136/jnnp.20.1.11
Simeone, A., Gulisano, M., Acampora, D., Stornaiuolo, A., Rambaldi, M., & Boncinelli, E.
(1992). Two vertebrate homeobox genes related to the Drosophila empty spiracles
gene are expressed in the embryonic cerebral cortex. *EMBO J, 11*(7), 2541-2550.
doi:10.1002/j.1460-2075.1992.tb05319.x
Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of
the hippocampal longitudinal axis. *Nat Rev Neurosci, 15*(10), 655-669.
doi:10.1038/nrn3785
Sugiyama, T., Osumi, N., & Katsuyama, Y. (2014). A novel cell migratory zone in the
developing hippocampal formation. *J Comp Neurol, 522*(15), 3520-3538.
doi:10.1002/cne.23621

Swindell, E. C., Bailey, T. J., Loosli, F., Liu, C., Amaya-Manzanares, F., Mahon, K. A., . . .
Jamrich, M. (2006). Rx-Cre, a tool for inactivation of gene expression in the
developing retina. *Genesis, 44*(8), 361-363. doi:10.1002/dvg.20225
Takeda, K., Inoue, H., Tanizawa, Y., Matsuzaki, Y., Oba, J., Watanabe, Y., . . . Oka, Y.
(2001). WFS1 (Wolfram syndrome 1) gene product: predominant subcellular
localization to endoplasmic reticulum in cultured cells and neuronal expression in rat
brain. *Hum Mol Genet, 10*(5), 477-484. doi:10.1093/hmg/10.5.477
Tang, K., Rubenstein, J. L., Tsai, S. Y., & Tsai, M. J. (2012). COUP-TFII controls amygdala
patterning by regulating neuropilin expression. *Development, 139*(9), 1630-1639.
doi:10.1242/dev.075564
Toda, T., Parylak, S. L., Linker, S. B., & Gage, F. H. (2019). The role of adult hippocampal
neurogenesis in brain health and disease. *Mol Psychiatry, 24*(1), 67-87.
doi:10.1038/s41380-018-0036-2
Tole, S., Goudreau, G., Assimacopoulos, S., & Grove, E. A. (2000). Emx2 is required for
growth of the hippocampus but not for hippocampal field specification. *J Neurosci,*
20(7), 2618-2625. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/10729342
Tuncdemir, S. N., Lacefield, C. O., & Hen, R. (2019). Contributions of adult neurogenesis to
dentate gyrus network activity and computations. *Behav Brain Res, 374*, 112112.
doi:10.1016/j.bbr.2019.112112
Tyng, C. M., Amin, H. U., Saad, M. N. M., & Malik, A. S. (2017). The Influences of
Emotion on Learning and Memory. *Front Psychol, 8*, 1454.
doi:10.3389/fpsyg.2017.01454
Vorhees, C. V., & Williams, M. T. (2006). Morris water maze: procedures for assessing
spatial and related forms of learning and memory. *Nat Protoc, 1*(2), 848-858.
doi:10.1038/nprot.2006.116
Yang, X., Feng, S., & Tang, K. (2017). COUP-TF Genes, Human Diseases, and the
Development of the Central Nervous System in Murine Models. *Curr Top Dev Biol,*
125, 275-301. doi:10.1016/bs.ctdb.2016.12.002
Yoshida, M., Suda, Y., Matsuo, I., Miyamoto, N., Takeda, N., Kuratani, S., & Aizawa, S.
(1997). Emx1 and Emx2 functions in development of dorsal telencephalon.
Development, 124(1), 101-111. doi:10.1242/dev.124.1.101
Yu, D. X., Marchetto, M. C., & Gage, F. H. (2014). How to make a hippocampal dentate
gyrus granule neuron. *Development, 141*(12), 2366-2375. doi:10.1242/dev.096776
Zhang, K., Yu, F., Zhu, J., Han, S., Chen, J., Wu, X., . . . Tang, K. (2020). Imbalance of
Excitatory/Inhibitory Neuron Differentiation in Neurodevelopmental Disorders with
an NR2F1 Point Mutation. *Cell Rep, 31*(3), 107521. doi:10.1016/j.celrep.2020.03.085
Zhao, Y., Sheng, H. Z., Amini, R., Grinberg, A., Lee, E., Huang, S., . . . Westphal, H. (1999).
Control of hippocampal morphogenesis and neuronal differentiation by the LIM
homeobox gene Lhx5. *Science, 284*(5417), 1155-1158.
doi:10.1126/science.284.5417.1155
Zhou, C., Qiu, Y., Pereira, F. A., Crair, M. C., Tsai, S. Y., & Tsai, M. J. (1999). The nuclear
orphan receptor COUP-TFI is required for differentiation of subplate neurons and
guidance of thalamocortical axons. *Neuron, 24*(4), 847-859. doi:10.1016/s0896-
6273(00)81032-6
Zhou, C., Tsai, S. Y., & Tsai, M. J. (2001). COUP-TFI: an intrinsic factor for early
regionalization of the neocortex. *Genes Dev, 15*(16), 2054-2059.
doi:10.1101/gad.913601
Zhou, X., Liu, F., Tian, M., Xu, Z., Liang, Q., Wang, C., . . . Yang, Z. (2015). Transcription
factors COUP-TFI and COUP-TFII are required for the production of granule cells in
the mouse olfactory bulb. *Development, 142*(9), 1593-1605. doi:10.1242/dev.115279
941

Figure legends

Figure 1. Duplicated CA1 and CA3 domains are generated in the ventral hippocampus of RX$^{Cre/+}$; *Nr2f2$^{F/F}$* mutant mice.

A, The expression of NR2F1 (**a, d, g, j, m**) and NR2F2 (**b, e, h, k, n**) in coronal sections (**a-f**) and sagittal sections (**g-o**) of the hippocampus at postnatal month 1 (1M); representative Western blots and quantitative densitometry data for the expression of NR2F1 and NR2F2 in the dorsal and ventral hippocampus at 1M (**p-q**). **B,** In coronal sections along the rostrocaudal axis (**a-f**) and sagittal sections along the lateral-medial axis (**g-l**) of the hippocampus in mutant mice, compared with that in control mice (**a-c, g-i**), the ectopic CA-like structure, indicated by the star, was observed in the ventral region in *Nr2f2* gene mutant (RX$^{Cre/+}$; *Nr2f2$^{F/F}$*) mice at 1M (**d-f**, **j-l**). **C,** The expression of HuB and Ctip2 in the corresponding inserted area in **a-f** under a high magnification objective lens at 1M (**g-r**); compared with those of control mice (**a-c, g-i, m-o**), the duplicated HuB-positive CA3 domain, indicated by the star, and Ctip2-positive domains, indicated by the arrowhead, were specifically observed in the ventral hippocampus (**d-f, p-r**) but not in the dorsal hippocampus (**d-f**, **j-l**) of *Nr2f2* mutant mice at 1M; Ctip2 positive AHi and MePD amygdaloid nuclei were barely observed in the *Nr2f2* mutant mice, indicated by the arrows, instead of the ectopic CA domains at the prospective amygdaloid regions (**e-f**, **q-r**). Data are expressed as the mean ± SEM. *Student's t test,* $^*P$<0.05, $^{**}P$<0.01. AHi, amygdalohippocampal area; AMY, amygdala nuclei; CTX, cortex; dCA1, dorsal CA1; dCA3, dorsal CA3; dDG, dorsal dentate gyrus; dHPC, dorsal hippocampus; MePD, posterodorsal part of the medial amygdaloid nucleus; PMCo, posteromedial cortical amygdaloid nucleus; vCA1, ventral CA1; vCA3, ventral CA3; vDG, ventral dentate gyrus; vHPC, ventral hippocampus. Scale bars, **Aa-c**, **Ad-f**, **Aj-o**, **Cg-r**, 100 μm; **Ag-i**, **Ba-l**, **Ca-f**, 200μm.

Figure 1—figure supplement 1. The expression of *Nr2f* genes in the early developing

hippocampus and different conditional knock mouse models.

A, The expression of *Nr2f1* and *Nr2f2* genes in the forebrain at E10.5 (**a-b**) and E11.5 (**c-d**).

B, The expression of *Nr2f1* and *Nr2f2* genes in the developing hippocampus at E12.5 (**a-c**),

E14.5 (**d-f**), and P0 (**g-l**). **C,** Compared with that of control mice (**a-c**), *Nr2f2* is efficiently

deleted by RXCre recombinase in the hippocampus of *Nr2f2* mutant mice at 1M (**d-f**); *Nr2f1*

is clearly deleted by RXCre recombinase in the hippocampus of *Nr2f1* mutant mice at 1M (**g-**

i). Compared with that of control mice (**j-l**), NR2F1 and NR2F2 were efficiently deleted by

RXCre recombinase at the hippocampal primordium, including the MP and CH, in *Nr2f1/2*

double-mutant mice at E14.5 (**m-o**). **D,** Compared with that of the control mice (**a-c, g-h**), the

expression of *Nr2f2* was significantly decreased in the hippocampal primordium of the

homozygous mutant mice at E11.5; meanwhile, the LacZ signals obviously increased in the

Nr2f2 homozygous mutant mice at E11.5 (**d-f, j-k**). Compared with that of the control mice (**i**),

the expression of NR2F1 is activated in the caudal hippocampal primordium of the

homozygous mutant mice at E11.5 (**l**). CH, cortical hem; dHPC, dorsal hippocampus; HP,

hippocampal primordium; HPC, hippocampus; MP, medial pallium; vHPC, ventral

hippocampus. Scale bars, **Aa-d, Ba-f, Cj-o**, 200 μm; **Bg-l, Ca-i, Dg-l**, 100 μm; **Da-f**, 250 μm.

Figure 2. The specification and differentiation of the dorsal CA1 lineage failed with the

dysplastic dorsal hippocampus in RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant mice.

A, In coronal sections along the rostrocaudal axis (**a-f**) and sagittal sections along the lateral-

medial axis (**g-l**) of the hippocampus, compared with that of control mice (**a-c, g-i**), the dorsal

hippocampus was shrunken, indicated by the star, in *Nr2f1* gene mutant (RX$^{Cre/+}$; *Nr2f1$^{F/F}$*)

mice at 1M (**d-f, j-l**). **B,** The expression of HuB and Ctip2 in the corresponding inserted area

in **a-f** under a high magnification objective lens at 1M (**g-r**); compared with that of control

mice (**a-c**, **g-i**, **m-o**), the HuB-positive CA3 domain was reduced in the dorsal hippocampus, especially the Ctip2-positive dorsal CA1, which was barely detected in *Nr2f1* mutant mice at 1M (**d-f**, **j-l**), while their expression in the ventral hippocampus was comparable between the controls and mutants (**d-f**, **p-r**). **C,** The expression of HuB and Wfs1 in the corresponding inserted area in **a-f** under a high magnification objective lens at 1M (**g-r**); the expression of HuB and the dCA1 marker Wfs1 in the control (**a-c**, **g-i**, **m-o**) and *Nr2f2* mutant mice (**d-f**, **j-l**, **p-r**) at 1M. Wfs1-positive dorsal CA1 could not be detected in *Nr2f1* mutant mice at 1M, as indicated by the arrowhead. dHPC, dorsal hippocampus; HPC, hippocampus; vHPC, ventral hippocampus. Scale bars, **Aa-l**, **Ba-f**, **Ca-f,** 200 μm; **Bg-r**, **Cg-r,** 100 μm.

Figure 2—figure supplement 1. Defects in *Emx1$^{Cre/+}$*; *Nr2f1$^{F/F}$* mutant mice.

A, Immunofluorescence staining data showed that compared with those of control mice (**a, c, e, g**), the proportions of either the Wfs1- or Ctip2-positive dorsal CA1 domain were reduced in *Emx1$^{Cre/+}$*; *Nr2f1$^{F/F}$* mutant mice at 3M (**b, d, f, h**). **B,** Golgi staining showed that compared with those of controls, the numbers of branch points and secondary dendrites of both apical and basal dendrites were significantly reduced in the dorsal hippocampal CA1 pyramidal neurons of *Emx1$^{Cre/+}$*; *Nr2f1$^{F/F}$* mutant mice (**a-c**). **C,** The Morris water maze behavior test showed that compared with that of controls, spatial learning and memory were significantly damaged in *Emx1$^{Cre/+}$*; *Nr2f1$^{F/F}$* mutant mice. Data are expressed as the mean ± SEM. *Student's t test*, *$P<0.05$, **$P<0.01$, ***$P<0.001$. Scale bars, **Aa-h,** 100 μm; **Ba-b,** 50 μm.

Figure 3. Defects in the hippocampus in RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$* double-gene mutant mice.

A, In coronal sections along the rostrocaudal axis, compared with control mice (**a-d**), the hippocampus was atrophic in RX$^{Cre/+}$; *Nr2f1$^{F/F}$; Nr2f2$^{F/F}$* double mutant mice, indicated by the

star, and an ectopic unknown nucleus was observed in the caudal plates, indicated by the arrowhead (**e-h**). **B,** Compared with that of control mice (**a-c**, **g-i**), the expression of HuB, Ctip2, and Prox1 was decreased in the hippocampus of *Nr2f1/2* double-gene mutant mice at 3 weeks postnatal (3W) (**d-f, j-l**). **C,** Compared with that of control mice (**a-c**, **g-i**), the expression of HuB could not be detected in the presumptive CA3 domain, and the expression of Ctip2 or Prox1 could not be detected in the presumptive DG domain of the prospective ventral hippocampus of RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$* double mutant mice. Scale bars, **Aa-h**, 200 μm; **Ba-l**, **Ca-l,** 100 μm.

Figure 3—figure supplement 1. Adult neurogenesis was abnormal in the hippocampi of *Nr2f1/2* double-gene mutant mice.

A, The expression of GFAP and Nestin, markers of NSCs, in the SGZ of the vDG in control and *Nr2f2* mutant mice at 1M (**a-h**), in the SGZ of the dDG in control and *Nr2f1* mutant mice at 1M (**i-p**) and in the SGZ of the DG in control and *Nr2f1/2* double-gene mutant mice at 3W (**q-x**). **B,** The expression of Dcx, a marker of newborn neurons, in the SGZ of the vDG in control and *Nr2f2* mutant mice at 1M (**a-d**), in the SGZ of the dDG in control and *Nr2f1* mutant mice at 1M (**e-h**) and in the SGZ of the DG in control and *Nr2f1/2* double-gene mutant mice at 3W (**i-l**). **C,** Quantitative analysis of GFAP/Nestin-positive cells (**a**) and Dcx-positive cells (**b**) in the SGZ of the DG in control and *Nr2f1/2* double-gene mutant mice at 3W. The numbers of GFAP and Nestin double-positive NSCs and Dcx-positive newborn neurons were significantly reduced in double-mutant mice. Data are expressed as the mean ± SEM. *Student's t test*, **P<0.01, ***P<0.001. dDG, dorsal DG; NSC, neural stem cell; SGZ, subgranular zone; vDG, ventral DG. Scale bars, **Aa-x**, 50 μm; **Ba-l**, 100 μm.

Figure 4. The impairment of hippocampal trisynaptic connectivity in *Nr2f2* single-gene, *Nr2f1* single-gene, and *Nr2f1/2* double-gene mutant mice.

A, The expression of Calretinin, Calbindin, and SMI312 in the ventral hippocampus of the control (**a-b**, **e-f**, **i-j**) and *Nr2f2* single-gene mutant mice (**c-d**, **g-h**, **k-l**). **B,** The expression of Calretinin, Calbindin, and SMI312 in the dorsal hippocampus of the control (**a-b**, **e-f**, **i-j**) and *Nr2f1* single-gene mutant mice (**c-d**, **g-h**, **k-l**). **C,** The expression of Calretinin, Calbindin, and SMI312 in the hippocampus of the control (**a-b**, **e-f**, **i-j**) and *Nr2f1/2* double-gene mutant mice (**c-d**, **g-h**, **k-l**). dHPC, dorsal hippocampus; HPC, hippocampus; vHPC, ventral hippocampus. Scale bars, **Aa-l**, **Ba-l**, **Ca-l**, 100 μm.

Figure 5. *Nr2f* genes regulate the expression of key genes associated with early hippocampal development.

A, The expression profiles of genes involved in hippocampal development in control and the double mutant mice at E11.5. **B,** Compared with that of control mice (**a-b**, **a'-b'**, **a''-b''**, **i-j**, **i'-j'**, **q-r**, **q'-r'**), the expression of Lhx5 was reduced in double-mutant mice at E11.5 (**c-d**, **c'-d'**, **c''-d''**), E13.5 (**k-l**, **k'-l'**) and E14.5 (**s-t**, **s'-t'**); the expression of Lhx2 was comparable between the control and double-mutant mice at E11.5 (**e-h**, **e'-h'**); and compared with that of control mice (**m-n**, **m'-n'**, **u-v**, **u'-v'**), the expression of Lhx2 was increased in double-mutant mice at E13.5 (**o-p**, **o'-p'**) and E14.5 (**w-x**, **w'-x'**). **C,** Compared with that of control mice (**a-b**, **a'-b'**), the expression of Sox2 was normal in double-mutant mice at E14.5 (**c-d**, **c'-d'**); compared with that of control mice (**e-f**, **e'-f'**), the expression of Tbr2 was decreased in *Nr2f* mutant mice at E14.5 (**g-h**, **g'-h'**); compared with that of control mice (**i-j**, **i'-j'**), the expression of NeuroD1 was reduced in double-mutant mice at E14.5 (**k-l**, **k'-l'**). **D,** Quantitative analysis of Tbr2-positive cells and NeuroD1-positive cells in **Ce'-h'** and **Ci'-l'**. **E,** In the hippocampal primordium of the early embryo, *Nr2f1* is expressed dorsally in the MP, and *Nr2f2* is expressed

ventrally in the CH. In the mature hippocampus, the expression of *Nr2f1* is higher in the dorsal

hippocampus, which is related to spatial learning and memory, and the expression of *Nr2f2* is

mainly in the ventral hippocampus, which is associated with emotion and anxiety (**a**). Our

findings support a novel molecular mechanism by which *Nr2f1* and *Nr2f2* may cooperate to

ensure the appropriate morphogenesis and functions of the hippocampus by modulating the

Lhx5-Lhx2 axis (**b**). Data are expressed as the mean ± SEM. *Student's t test,* **P*<0.01. CH,

cortical hem; ChP, choroid plexus; DP, dorsal pallium; HP, hippocampal primordium; MP,

medial pallium. Scale bars, **Ba-x, Ca-l**, 200 μm.

1074

Figure 5—figure supplement 1. *Nr2f1* and *Nr2f2* genes coordinate to control distinct

characteristics of the hippocampus.

Roles of *Nr2f1* and *Nr2f2* genes in the development and function of the hippocampus and the

association with neurological diseases. *Nr2f1* is required for the morphogenesis of the dorsal

hippocampus and the specification of dorsal CA1 pyramidal neuron lineage, which are

associated with neurodevelopmental disorders, including ID and ASD. *Nr2f2* is required to

prevent the duplication of the CA1 and CA3 lineages of the ventral hippocampus, which may

be related to psychiatric diseases such as depression, anxiety, or schizophrenia. The *Nr2f1* and

Nr2f2 genes are novel intrinsic regulatory genes, which cooperate with each other to ensure the

early morphogenesis of the hippocampus.

1085

1086 **The information of the source data files**

1087 **Figure 1-source data 1.** The Nissl staining results of the control and RX$^{Cre/+}$; *Nr2f2$^{F/F}$* mutant

1088 mice at 1M (part 1); the expression of HuB and Ctip2 in the hippocampus of the control and

1089 RX$^{Cre/+}$; *Nr2f2$^{F/F}$* mutant mice at 1M (part 1).

1090

1091 **Figure 1-source data 2.** The expression of NR2F1 and NR2F2 in coronal sections and sagittal

1092 sections of the mouse brain at 1M (part 1); the expression of HuB and Ctip2 in the hippocampus

1093 of the control and RX$^{Cre/+}$; *Nr2f2$^{F/F}$* mutant mice at 1M (part 2).

1094

1095 **Figure 1-source data 3.** The expression of NR2F1 and NR2F2 in coronal sections and sagittal

1096 sections of the mouse brain at 1M (part 2).

1097

1098 **Figure 1-source data 4.** The expression of NR2F1 and NR2F2 in sagittal sections of the mouse

1099 brain at 1M (part 1).

1100

1101 **Figure 1-source data 5.** The expression of NR2F1 and NR2F2 in sagittal sections of the mouse

1102 brain at 1M (part 2); the Nissl staining results of the control and RX$^{Cre/+}$; *Nr2f2$^{F/F}$* mutant mice

1103 at 1M (part 2).

1104

1105 **Figure 1-source data 6.** The expression of NR2F1 and NR2F2 in sagittal sections of the mouse

1106 brain at 1M (part 3); Western blots data for the expression of NR2F1 and NR2F2 in the dorsal

1107 and ventral hippocampus at 1M; the Nissl staining results of the control and RX$^{Cre/+}$; *Nr2f2$^{F/F}$*

1108 mutant mice at 1M (part 3).

1109

1110 **Figure 1-figure supplement 1-source data 1.** The expression of *Nr2f1* and *Nr2f2* genes in the

1111 developing hippocampus at E12.5; the deletion efficiency of RXCre recombinase in the

1112 hippocampus of RX$^{Cre/+}$; *Nr2f2$^{F/F}$*, RX$^{Cre/+}$; *Nr2f1$^{F/F}$*, and RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$* mice (part

1113 1).

1114

1115 **Figure 1-figure supplement 1-source data 2.** The expression of *Nr2f1* and *Nr2f2* genes in the

1116 telencephalon at E10.5, E11.5, E14.5, and P0; the deletion efficiency of RXCre recombinase

1117 in the hippocampus of RX$^{Cre/+}$; *Nr2f2$^{F/F}$*, RX$^{Cre/+}$; *Nr2f1$^{F/F}$*, and RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$* mice

1118 (part 2).

1119

1120 **Figure 2-source data 1.** The Nissl staining results of the control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant

1121 mice at 1M (part 1).

1122

1123 **Figure 2-source data 2.** The Nissl staining results of the control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant

1124 mice at 1M (part 2); the expression of HuB and Wfs1 in the hippocampus of the control and

1125 RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant mice at 1M.

1126

1127 **Figure 2-source data 3.** The expression of HuB and Ctip2 in the hippocampus of the control

1128 and RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant mice at 1M.

1129

1130 **Figure 2-source data 4.** The expression of HuB, Wfs1, and Ctip2 in the hippocampus of the

1131 control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant mice at 1M.

1132

1133 **Figure 2-figure supplement 1-source data 1.** The expression of Wfs1 and Ctip2 in the dorsal

1134 hippocampus of the control and *Emx1$^{Cre/+}$*; *Nr2f1$^{F/F}$* mutant mice at 3M; Golgi staining results

1135   of the dorsal hippocampal CA1 pyramidal neurons and Morris water maze behavior test data

1136   of the control and $Emx1^{Cre/+}$; $Nr2f1^{F/F}$ mutant mice.

1137

1138   **Figure 3-source data 1.** The Nissl staining results of the control and $RX^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$

1139   double-gene mutant mice at 3W.

1140

1141   **Figure 3-source data 2.** The expression of HuB and Ctip2 in the hippocampus of the control

1142   and $RX^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-gene mutant mice at 3W.

1143

1144   **Figure 3-source data 3.** The expression of HuB and Prox1 in the hippocampus of the control

1145   and $RX^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-gene mutant mice at 3W.

1146

1147   **Figure 3-source data 4.** The expression of HuB, Ctip2, and Prox1 in the hippocampus of the

1148   control and $RX^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-gene mutant mice at 3W.

1149

1150   **Figure 3-figure supplement 1-source data 1.** The expression of GFAP, Nestin, and Dcx in

1151   the SGZ of vDG in the control and $RX^{Cre/+}$; $Nr2f2^{F/F}$ mutant mice at 1M, in the SGZ of dDG

1152   in the control and $RX^{Cre/+}$; $Nr2f1^{F/F}$ mutant mice at 1M, and in the SGZ of DG in the control

1153   and $RX^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-gene mutant mice at 3W (part 1).

1154

1155   **Figure 3-figure supplement 1-source data 2.** The expression of GFAP, Nestin, and Dcx in

1156   the SGZ of vDG in the control and $RX^{Cre/+}$; $Nr2f2^{F/F}$ mutant mice at 1M, in the SGZ of dDG

1157   in the control and $RX^{Cre/+}$; $Nr2f1^{F/F}$ mutant mice at 1M, and in the SGZ of DG in the control

1158   and $RX^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-gene mutant mice at 3W (part 2).

1159

Figure 3-figure supplement 1-source data 3. The expression of GFAP and Nestin in the SGZ

of vDG in the control and RX$^{Cre/+}$; *Nr2f2$^{F/F}$* mutant mice at 1M, in the SGZ of dDG in the

control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant mice at 1M, and in the SGZ of DG in the control and

RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$* double-gene mutant mice at 3W; quantitative analysis of

GFAP/Nestin-positive cells in the SGZ of DG in the control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$*

double-gene mutant mice at 3W.

1166

Figure 3-figure supplement 1-source data 4. The expression of GFAP, Nestin, and Dcx in

the SGZ of vDG in the control and RX$^{Cre/+}$; *Nr2f2$^{F/F}$* mutant mice at 1M, in the SGZ of dDG

in the control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$* mutant mice at 1M, and in the SGZ of DG in the control

and RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$* double-gene mutant mice at 3W (part 3); quantitative analysis

of Dcx-positive cells in the SGZ of DG in the control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$* double-

gene mutant mice at 3W.

1173

1174   **Figure 4-source data 1.** The expression of Calretinin, Calbindin, and SMI312 in the ventral

1175   hippocampus of the control and RX$^{Cre/+}$; *Nr2f2$^{F/F}$* single-gene mutant mice at 1M.

1176

1177   **Figure 4-source data 2.** The expression SMI312 in the dorsal hippocampus of the control and

1178   RX$^{Cre/+}$; *Nr2f1$^{F/F}$* single-gene mutant mice at 1M, and the expression of Calretinin and

1179   Calbindin in the hippocampus of the control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$*; *Nr2f2$^{F/F}$* double-gene

1180   mutant mice at 3W.

1181

1182   **Figure 4-source data 3.** The expression of Calretinin and Calbindin in the dorsal hippocampus

1183   of the control and RX$^{Cre/+}$; *Nr2f1$^{F/F}$* single-gene mutant mice at 1M, and the expression of

1184   SMI312 in the hippocampus of the control and RX$^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-gene mutant

1185   mice at 3W.

1186

1187   **Figure 5-source data 1.** The expression of Lhx5 and Lhx2 in the telencephalon of the control

1188   and RX$^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-mutant mice at E14.5; the expression of Tbr2 and

1189   NeuroD1 in the telencephalon of the control and RX$^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-mutant

1190   mice at E14.5 (part 1).

1191

1192   **Figure 5-source data 2.** The expression of Sox2, Tbr2, and NeuroD1 in the telencephalon of

1193   the control and RX$^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-mutant mice at E14.5.

1194

Figure 5-source data 3. The expression profiles of genes involved in the hippocampal

development of the control and double mutant mice at E11.5; the expression of Lhx5 and Lhx2

in the telencephalon of the control and RX$^{Cre/+}$; $Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-mutant mice at E11.5

and E13.5; the expression of Tbr2 and NeuroD1 in the telencephalon of the control and RX$^{Cre/+}$;

$Nr2f1^{F/F}$; $Nr2f2^{F/F}$ double-mutant mice at E14.5 (part 2); quantitative analysis of Tbr2-positive

and NeuroD1-positive cells in the hippocampal primordium of the control and RX$^{Cre/+}$; $Nr2f1^{F/F}$;

$Nr2f2^{F/F}$ double-mutant mice at E14.5.

1202

1203   **Figure 5-figure supplement 1-source data 1.** Roles of $Nr2f1$ and $Nr2f2$ genes in the

1204   development and function of the hippocampus and the association with neurological disorders.

Figure 1, Yang et al.

Figure 2, Yang et al.

Figure 2—figure supplement 1, Yang et al.

Figure 3, Yang et al.

Figure 3—figure supplement 1, Yang et al.

Figure 4, Yang et al.

Figure 5—figure supplement 1, Yang et al.