Wolbachia has subtle effects on thermal preference in highly inbred Drosophila

melanogaster which vary with life stage and environmental conditions

Anton Strunov[1,*], Charlotte Schoenherr[1], Martin Kapun[1,2,*]

1 Center for Anatomy and Cell Biology, Medical University of Vienna, Austria

2 Central Research Laboratories, Natural History Museum of Vienna, Austria

* co-corresponding authors

Contact:

anton.strunov@meduniwien.ac.at

martin.kapun@meduniwien.ac.at

Running title: *Wolbachia and thermal preference of Drosophila*

Keywords: *Drosophila*, *Wolbachia*, thermal preference, development

Abstract (195 words)

Temperature fluctuations are challenging for ectotherms which are not able to regulate body temperature by physiological means and thus have to adjust their thermal environment via behavior. However, little is yet known about whether microbial symbionts influence thermal preference (T_p) in ectotherms by modulating their physiology. Several recent studies have demonstrated substantial effects of *Wolbachia* infections on host T_p in different *Drosophila* species. These data indicate that the direction and strength of thermal preference variation is strongly dependent on host and symbiont genotypes and highly variable among studies. By employing highly controlled experiments, we investigated the impact of several environmental factors including humidity, food quality, light exposure, and experimental setup that may influence T_p measurements in adult *Drosophila melanogaster* flies. Additionally, we assessed the effects of *Wolbachia* infection on T_p of *Drosophila* at different developmental stages, which has not been done before. We find only subtle effects of *Wolbachia* on host T_p which are strongly affected by experimental variation in adult, but not during juvenile life stages. Our in-depth analyses show that environmental variation has a substantial influence on T_p which demonstrates the necessity of careful experimental design and cautious interpretations of T_p measurements together with a thorough description of the methods and equipment used to conduct behavioral studies.

Introduction

Temperature modulates many physiological processes and has direct effects on development, survival and reproduction of any organism[1,2]. Ectotherms lack the ability to regulate their body temperature by physiological means and are particularly affected by temperature variation[3]. Their thermoregulation is thus often mediated through behavior[4] (Stevenson, 1985) so ectotherms tend to occupy environmental niches close to their optimal thermal conditions to survive and propagate. Every organism exhibits a thermal preference (T_p), which is the preferred body temperature or temperature range that is chosen in the absence of other ecological constraints[5]. However, estimates of T_p are by no means absolute, but may be strongly influenced by interactions with ecological factors and even with symbionts. Endotherms, for example, combat bacterial and viral infections with fever by physiologically increasing their body temperature outside the thermal optimum of their infectious agents. Several studies show that also ectotherms employ similar behavioral strategies by changing their T_p[6–11]. For example, crickets actively seek higher temperatures when infected with the pathogenic bacterium *Serratia marcescens*, which may have physiological effects similar to fever in response to pathogenic infections[12]. However, also the opposite behavior, i.e., behavioral chill, has been reported for poikilotherm organism as a mechanism to fight pathogens, for example, in *Drosophila*[13].

In particular, three recent studies investigated changes of thermal behavior in *Drosophila melanogaster* infected with the bacterial endosymbiont *Wolbachia*[14–16]. Two of these studies found that, depending on the *Wolbachia* variant, infected flies chose colder temperatures than uninfected individuals[14,15]. While such a behavioral response is probably strongly influenced by environmental and experimental conditions and not a general pattern[16], these findings may suggest that the hosts can alleviate detrimental effects of high titer infections by choosing ambient temperatures outside the optimal physiological range of the symbiont. It, however, remains uncertain if such behavioral patterns are restricted to adults or are also found at juvenile life stages. Finding the optimal temperature for development is pivotal since any major disturbance during juvenile stages might severely decrease survival and reproduction chances of the imago[17]. Manipulating thermal preference at this life stage of the host might thus be risky yet beneficial for the microbial symbiont, by "setting up" its thermal environment early in the host development. However, to the best of our knowledge, no study has yet investigated the effects of *Wolbachia* on *Drosophila* T_p at juvenile life stages.

Measuring thermal behaviour in *Drosophila* is complex, and many environmental factors and experimental noise that potentially influence the T_p have to be accounted for. These can include, for example, environmental factors such as humidity, light, olfactory and

acoustic cues, but also properties of the experimental design, such as age of the flies,

developmental temperature, number of specimens and the design of the apparatus that

measures thermal preference, as comprehensively reviewed in Dillon et al.[18]. This issue

becomes evident when comparing the divergent results of the three aforementioned studies

that specifically assessed the T_p of *D. melanogaster* in the presence of *Wolbachia*

infections[14–16]. The absolute T_p values for the same *Wolbachia* strain vary strongly among

the studies (up to 5-6°C), which is potentially influenced by different host genetic

backgrounds, experimental conditions or temperature ranges used in these studies.

Assessing the effects of different environmental factors on thermal preference in the

Wolbachia-Drosophila system is therefore crucial for an informed analysis and interpretation

of behavioral measurements.

Here, we demonstrate that several environmental factors, such as humidity, food,

and the structure of the thermal gradient device have a major impact on T_p in adult *D.*

melanogaster flies in the context of strain-specific *Wolbachia* infections. Additionally, we

investigate - for the first time - the effect of *Wolbachia* on juvenile T_p and find that it is not

affected by infection in early and late fly larvae and at pupariation.

Materials and Methods

Fly lines

In our experiments, we used four highly inbred, long-term laboratory strains of *D.*

melanogaster which were uninfected (*w-*) or infected with either two of the most common

natural *Wolbachia* variants (*w*Mel and *w*MelCS) or with the *w*MelPop lab-variant that were

previously investigated by Truitt et al.[14] for thermal preference. All fly lines were initially

established by Luis Teixeira[19] in a DrosDel w^{1118} isogenic background. Flies were maintained

on a custom fly medium based on agar-agar, molasses, and yeast at 24°C with 12h:12h

light:dark cycle and an average humidity of 50%. Prior to experiments, the infection status

was confirmed by PCR using *Wolbachia*-specific primers amplifying parts of the *wsp* gene:

forward - tggtccaataagtgatgaagaaactagcta and reverse - aaaaattaaacgctactccagcttctgcac[20].

Wolbachia variants infecting the fly strains were distinguished with diagnostic VNTR-141

PCR primers: forward - ggagtattattgatatgcg and reverse - gactaaaggttattgcat. PCR products

vary in length among *Wolbachia* strains[21].

Thermal gradient machine

To carry out T_p measurements in *D. melanogaster* larvae, during pupariation and in

adult flies, we designed and built a new thermal gradient apparatus with two narrow arenas

for precise temperature measurements and with limited space for movement along the

thermal gradient. The "plate" device consisted of an elongated aluminum plate as the bottom
piece resting on two peltier elements, which are either heating or cooling to generate a
temperature gradient. On top, we used an equisized plexiglass plate as the cover, which
contained holes to apply test subjects. Two narrow arenas with 6 mm heights and widths
were established by placing three thin plexiglass dividers between the two plates, which
contained holes for inserting temperature sensors. We restricted the length of the arena to
30 cm by adding foam stoppers at the ends of each lane to prevent flies from escaping and
to maintain airflow. In addition, Whatman paper was used as a bottom layer to cover the
aluminum plate in the arenas. We built two identical devices with the aforementioned design,
which allowed us to investigate T_p in four arenas in parallel during one experimental assay
(see https://github.com/capoony/DrosophilaThermalGradient for a detailed description).
For T_p measurements in adult flies, we additionally designed and built a second
"tube" thermal gradient device with a tubular shape following Rajpurohit and Schmidt[22] and
complementing the design as described in Truitt et al.[14]. This apparatus consisted of a
central cylindrical aluminum rod connected to quadratic aluminum bases at both ends that
conduct heat from two peltier elements, which are either heating or cooling. The aluminum
rod was placed inside a transparent plexiglass tube, which left an open space of
approximately 4 cm in diameter. The plexiglass tube was attached to the aluminum rod with
Teflon covers which enclosed the fly arena on both sides. The plexiglass tube contained
several holes to insert flies and to attach sensors for temperature measurement. A Whatman
paper was inserted right above the aluminum rod to divide the arena into two halves. Flies
were kept in the upper half, which facilitated imaging the location of the flies within the
cylindrical arena around the central rod from atop during T_p measurements (see
https://github.com/capoony/DrosophilaThermalGradient for a detailed description).
Peltier elements (P&N Technology, Xiamen Fujian China) generating heat were
attached to small heat sinks (Fischer Electronic, Germany) connected to a ventilator
(Oezpolat, Germany) to stabilize the temperature. Cooling peltier elements produce excess
waste heat and were therefore placed on top of a bigger heat sink (Fischer Electronic,
Germany), which was partially submerged in a bowl with cold water that was constantly
cooled down by pipes supplying cold water from a distant water bath (fbc630, Fisher
scientific, USA).
20 min prior to each experiment, we attached the peltier elements to power supplies
set to 2 V and 3 V for cooling and heating, respectively, to stabilize the temperature gradient
within the thermal gradient apparatus. We found that the temperature gradient remained
stable for at least 1 h, which generally exceeded the duration of an experimental run (app.
20-30 min). After each experiment, the plexiglass parts of the apparatus were cleaned with

soap and hot water to remove any potential pheromones and other odorants, which might

interfere with thermal preference behavior of flies.

Each experiment was carried out in a dark isolated room with no, or very low noise

distraction. The room was equipped with air-conditioning to keep the ambient temperature

stable at 21-23°C, which was constantly monitored with a data logger. The temperature

within the devices was measured at three points (in the center and at the hot and cold

edges) with digital thermal sensors (Analog devices, USA). The data from thermal sensors

were collected with a custom python script on a Raspberry pi 3B+ computer (Raspberry Pi

Foundation, UK) in 10 sec intervals and saved as text files. The position of the flies within

the devices were assessed by an infrared camera attached atop of the thermal gradient

apparatus. Images were taken every 30 sec and saved to the Raspberry pi computer for

each run. We developed a custom python script to estimate T_p for every fly using information

of an individual's position relative to the linear temperature gradient between the

temperature measurement points. The coordinates of each individual fly and of the thermal

sensors were manually assessed in ImageJ[23] based on infrared images captured 20 min

after the onset of each experiment. The python scripts for quantifying thermal preference

from the coordinates including a test dataset, a detailed description and all raw images from

the thermal preference assays are available at GitHub

(https://github.com/capoony/DrosophilaThermalGradient).

Temperature preference measurements in larvae

To assess larval T_p, we used early and late 3rd instar larvae, 72 h, and 120 h after

egg laying (AEL), respectively, and focused on two fly strains that were either uninfected (w-

) or infected with *w*MelCS for thermal preference experiments (see *Fly lines* section in

Materials and methods). To obtain tightly synchronized cohorts of larvae, we maintained 40-

50 adult female flies at 24°C in glass vials with the medium for 3-6 h for egg laying and then

transferred these to a fresh medium for collecting another replicate. After 72 or 120 h, larvae

were moved to an 18% sucrose solution (in dH_2O) using a spatula. Floating larvae were

gently collected with a cut 1 mL pipette tip and transferred to dH_2O. After washing three

times with dH_2O, larvae were transferred into empty plastic Petri dishes (d=2.5 cm) and kept

for 10-20 min to recover from the stress.

For T_p measurements, we covered the bottom of each lane in the thermal gradient

machine with a 1mm layer of 2% agarose gel to create a habitable environment for larvae

that allowed them to crawl freely. We started the experiment when the thermal gradient was

constant and linear from 18°C to 28°C (1°C per 3cm). Experiments were carried out in a

dark room at 24-25°C, with 30% humidity at different times of the day (from 11 am to 4 pm).

Each run included two replicates of the *w-* and the *w*MelCS lines. 50-100 larvae were

introduced in the middle of each lane (23°C) with a wet brush and a Plexiglass cover was

applied on top of the machine to prevent larvae from escaping. The position of larvae was

analyzed on site after 30 min by quantifying the numbers of larvae located in a certain

temperature zone marked with a pen beforehand.

Measurement of pupariation Tp

For assessing T_p at pupariation, we used late 3rd-instar larvae, 140h after egg laying

(AEL) close to pupariation. Again, we focused on the *w-* and *w*MelCS strains for the

experiments and obtained synchronized cohorts of 140 h-old larvae as explained above.

For T_p measurements, we covered the bottom of the four lanes of the gradient

machine with a 1 mm thick sheet of odorless black plastic to create a contrasting

background and to supply a solid surface for pupariation. We inserted larvae when the

gradient was constant and linear from 18°C to 28°C (1°C per 3cm). Experiments were

carried out in a dark room at 24-25°C with 30% humidity for approximately 12h until every

larva had pupariated. Each run included two replicate lines at the same time (2 times *w-* and

2 times *w*MelCS). 50-100 larvae were introduced in the middle of the lane (23°C) by wet

brush and a plexiglass cover was used to seal the top of the machine to prevent larvae from

crawling out of the arenas. Thermal preference at pupariation was analyzed after 6-12 h by

quantifying the numbers of pupae in certain temperature zones as described above.

Temperature preference measurements in adults

Prior to the experiments, we selected cohorts of 20 females and 20 males, which

were one week old (± two days), for each of the four strains (*w-*, *w*Mel, *w*MelCS and

*w*MelPop). We transferred these to fresh glass vials with medium by gently applying CO_2

and then allowed the flies to recover for two days. To reduce the possibility of an

experimenter bias, we anonymized the ID's of the cohorts prior to experiments by an

independent researcher. For each assay on the "plate" device, we included one replicate of

all four fly lines and placed each of the groups in one of the four lanes. For experiments

using the "tube" device, we carried out two replicate assays in parallel, using randomly

drawn cohorts. Flies were gently collected from the vial with a handmade fly aspirator and

transferred to the thermal gradient apparatus through a hole in the center. Then, the hole

was tightly sealed with cotton, the light was turned off and measurements were taken during

20 min. An image of the whole apparatus was taken every 30 seconds with an infrared

camera attached atop the setup.

Because time of the day can have a strong influence on T_p in *Drosophila*[24], we

performed a maximum of 4 runs per day to avoid potential bias from circadian rhythms. The

time of the runs during the day was usually as follows: first two runs from 10 am to 12 am,

last two runs from 1 pm to 3.30 pm. To quantify T_p of individual flies in the arena, we used

the images of fly locations taken 20 min after the onset of the experiment similar to Truitt et

al.[14].

In addition to the experiments outlined above, we sought to test how environmental

variation influences T_p. We accounted for four different factors: humidity, light, food, and type

of the gradient machine. All four factors were tested in at least three replicates with the

"tube" device, except for the last one where both types were used for comparison. When

testing for the effects of humidity, we used a standard humidifier to increase the humidity

from 30%, which we usually measured without any adjustment, to stable 60% humidity in the

test room. To test the influence of light on thermal preference, we positioned the gradient

machine at the center above a ceiling lamp and performed thermal preference runs with the

light on and off using different biological replicates. When testing for the effect of food quality

on T_p, we kept the flies for at least three generations on different diets before conducting the

experiment. We compared the effects of our in-house food recipe, described above, to

Carolina 4-24 instant food, obtained directly from the manufacturer (Carolina, USA). To test

how the type of device affects T_p, we combined the data of the runs in the two different

devices carried out in the same room with similar ambient temperatures to keep

environmental conditions consistent. The analysis of images taken under infrared light did

not allow to distinguish between sexes. Since differences in T_p has not been reported in

Drosophila melanogaster[25,26], we do not assume that ignoring the factor "sex" would lead to

biased results.

Statistical analyses

The effects of *Wolbachia* infection and other environmental factors on T_p were

analyzed using generalized linear mixed models (GLMMs) with a Poisson error structure in

R[27] based on the "glmer" function in the lme4 package[28]. In our models, we considered T_p of

every individual as the dependent variable, and infection type as the fixed factor. Moreover,

we always included the factor replicate experiments nested within infection type as random

factors in each of our models. For the larval T_p experiment, we additionally included the

larval age (72 h and 120 h) as a fixed factor and the interaction of larval age with infection

status in our model. Moreover, for most of the models, we included the time of the run of

each replicate during the day as a random factor. We used the *anova* function in R to test for

a significant effect of infection status on thermal preference and calculated a Type-III

analysis of deviance comparing nested models to test for significant effects of fixed factors

and interaction in the T_p experiments. Whenever the factor *infection status* had a significant

effect, we performed post-hoc tests using Tukey's HSD method as implemented in the

emmeans package[30]. We also inferred the skewness of T_p distributions using the *moments*

package[31] and compared the average skewness among experimental setups with Wilcoxon

Rank tests in *R*. All *R* code can be found at

https://github.com/capoony/DrosophilaThermalPreference.

Results and Discussion

Wolbachia infection has no effect on T_p in 3^{rd}-instar Drosophila larvae and on pupariation T_p

in D. melanogaster

Using the newly designed thermal gradient apparatus (see Materials and Methods),

we found that T_p in early (72h AEL; n=281) and late (120h AEL; n=658) 3^{rd}-instar larvae

significantly shifted from a median T_p of 22-23°C to a median T_p of 19°C (see **Table S1**) with

progression of development (GLMM: p=2.8e-8, x^2=34.8, df=2; **Table 1A**, **Figure 1A**). Our

results were consistent with previous findings[32,33]. However, we neither observed an effect of

Wolbachia on T_p in 3^{rd}-instar larvae (GLMM; p=0.34, x^2=2.1, df=2; **Table 1A**; **Figure 1A**) nor

an interaction of *Wolbachia* with larval age (GLMM; p=0.17, x^2=1.9, df=1). Since many

larvae clustered at the coldest spot at the wall in the gradient machine (18-19°C), we

repeated the experiment for the 120h age class, which exhibited lower T_p, and extended the

lower temperature range to 15°C. This extended analysis of 529 infected and 332 uninfected

3^{rd}-instar larvae at 120h AEL similarly showed no effect of *Wolbachia* infection on T_p of the

host (GLMM: p=0.38, x^2=0.76, df=1; **Figure S1**, **Table 1B**), although the median T_p

decreased by approximately 1°C with the majority of larvae of both infection types (*w*- and

*w*MelCS) clustering at temperatures between 16°C and 18°C (median T_p of 18°C; **Table S1**).

Table 1. **The influence of *Wolbachia* infections on *Drosophila* T_p at different stages of development.**

Significant effects of generalized mixed models (GLMM) inferred by analysis of deviance are highlighted in bold.

(A) T_p of 72h vs 120h larvae			
Effect	x^2	*df*	*p*-Value
age	**34.8**	**2**	**2.8E-8**
infection	2.1	2	0.34
age*infection	1.9	1	0.17
(B) T_p of 120h larvae			
Effect	x^2	*df*	*p*-Value
infection	0.8	1	0.38
(C) T_p of 140h larvae/pupae			
Effect	x^2	*df*	*p*-Value
infection	0.2	1	0.64
(D) T_p of adult (flat)			
Effect	x^2	*df*	*p*-Value
infection	3.3	3	0.35

Figure 1. Thermal preference (T_p) of early and late *Wolbachia*-infected *D. melanogaster* 3[rd]-instar larvae and larvae at the onset of pupariation. (A) Temperature shift in early and late 3[rd]-instar larvae that were either uninfected (w-; grey) or infected with wMelCS (red), 72 hours and 120 after egg laying (AEL) respectively (see also Figure S1 and Table 1). (B) T_p of *D. melanogaster* larvae at the onset of pupariation (140h AEL) for both infection types (see also Table 1). Note the absence of *Wolbachia* effects on thermal preference at all larval stages.

Pupariation in *Drosophila* is a crucial developmental stage during which the fly is immobilized for several days and undergoes complete metamorphosis. Thus, for larvae to properly develop choosing the optimal pupariation temperature is of great importance[18,34]. We analyzed the T_p of 192 infected and 142 uninfected larvae about to pupariate (140h AEL; **Table S1**). In comparison to T_p in late 3[rd]-instar larvae (median T_p of 18°C for both infection types; **Table S1**), the pupariation T_p shifted to warmer temperatures (**Figure 1B**, median T_p of 20°C for uninfected, and 21°C for infected individuals; **Table S1**). However, similarly to 3[rd]-instar larvae, we found no significant effect of *Wolbachia* on thermal preference at pupariation stage (GLMM: $p=0.64$, $x^2=0.21$, $df=1$; **Table 1C**).

The absence of *Wolbachia* effects on T_p in early and late 3[rd]-instar larvae and on pupariation T_p might indicate that infection does not impose significant costs at least in the developmental stages investigated here. It might be correlated with bacterial titer, which is known to influence the fitness of symbiotic association when it reaches high levels[35–39]. Although little is known about *Wolbachia* titer dynamics during fly development and in spite of yet undiscovered effects of *Wolbachia* infections during larval development, our findings could imply dormancy of infection at juvenile stages of the host. Strunov et al.[37], for example, showed that the titer of the pathogenic *w*MelPop strain in *D. melanogaster* is significantly

lower in larvae than in the imago and stable across development without any effects of
rearing temperature. However, an influence of temperature on *Wolbachia* titer has been
previously observed in adults after metamorphosis[40,41], indicating that bacterial multiplication
and pronounced interaction with the host may take place during late pupariation stage. In
line with this hypothesis, it was shown that *Wolbachia* interferes with pheromone production
in females during pupariation and affects female-to-male communication[42]. Additionally, two
recent papers demonstrated the absence of *Wolbachia* effects on locomotion and fitness at
juvenile stage of wild-caught *D. nigrosparsa*[43] and *D. melanogaster* flies[44]. The absence of
Wolbachia effects on host juvenile traits is challenged by a study, which reported stable
abundance of *Wolbachia* transcripts at all stages of host development[45], although there was
no data on translation of the transcripts. Thus, *Wolbachia* have presumably no or very low
and undetected effects on developing flies. Additionally, other cues like foraging for food[46]
might represent stronger behavioral stimuli, which could mask subtle effects of *Wolbachia* on
thermal behavior.
322
Wolbachia infection has subtle effects on thermal preference in adult flies that are influenced
by the environment and experimental design.
Using our new experimental setup, we sought to reproduce previous experiments
which reported that *Wolbachia* infection can lower thermal preference in adult *D.*
melanogaster flies depending on the *Wolbachia* variant studied[14,15,but see 16]. We used the
same fly lines, initially established by Teixeira et al.[19], as examined by Truitt et al.[14]. We
investigated five replicate cohorts of each fly line (*w-*, *w*Mel, *w*MelCS, and *w*MelPop) with
our new thermal gradient device under highly controlled environmental conditions.
Unexpectedly, we were unable to reproduce the previous results in Truitt et al.[14], since none
of the lines differed in thermal preference (GLMM: $p=0.35$, $x^2=3.29$, $df=3$; **Table1D**; **Figure
2**). Median T_p was 18.3°C, 18.8°C, 18.1°C, and 18.2°C for *w-*, *w*Mel, *w*MelCS, and *w*MelPop,
respectively (**Table S1**). The variation among the replicates was quite high with 2.4 to 3°C
standard deviation among the mean T_p, depending on the infection type studied.
336

Figure 2. Thermal preference of adult *D. melanogaster* with different infection types. Boxplots showing the

median T_p of flies that were either uninfected (w-) or with three different *Wolbachia* types (wMel: blue; wMelCS;

red and wMelPop: orange)

To rule out that the absence of T_p variation among strains infected with different

Wolbachia variants is an artifact of our experimental setup, we newly built a device similar to

the design in Truitt et al.[14] and repeated the T_p assays. Overall, median T_p was

approximately 3°C higher than the measurements from the "plate" device (21.7°C, 22°C,

21.5°C and 21.5°C average T_p for *w-*, *w*Mel, *w*MelCS, and *w*MelPop, respectively; **Table**

S1), which indicates a strong influence of the design on absolute T_p (GLMM: $p<2.2e-16$,

$x^2=748.6$, $df=4$; **Table 2A**). Additionally, we found significant differences among infection

types (GLMM: $p<0.0044$, $x^2=18.88$, $df=3$; **Table 2A**), albeit in the opposite direction

according to our expectations: the T_p of the *w*Mel variant increased by approximately 1°C

compared to *w-*(Tukey's post hoc test: $p<0.003$; **Table S2**). In contrast, *w*MelCS and

*w*MelPop variants did not show any effect on T_p compared to the uninfected control lines

(Tukey's post hoc test: $p>0.05$; **Table S2**), which is in stark contrast to substantial T_p

differences (2-4°C) among variants previously described by Truitt *et al.*[14]. Thus, despite

using a similar design for the thermal gradient device as in Truitt *et al.*[14], we were not able to

reproduce the previously obtained results. Keeping the lines in the lab for 4-5 years with high

inbreeding prior to the repeated thermal gradient assays might have potentially influenced

the results through mutation accumulation, titer reduction or other unknown genetic factors[47].

Moreover, differences in the environmental conditions during our experiment and the

experiments in Truitt *et al.*[14] may strongly confound thermal behavior. For example, the

experiments in Truitt *et al.*[14] were neither controlled for light cues nor for the effects of

humidity nor ambient temperature variation. While the experiments in Arnold *et al.*[15] were

carried out in complete darkness, these assays were also neither controlled for humidity
variation nor ambient temperature variation. Such environmental factors may have a strong
influence on behavior during experimental assays, which could potentially overwhelm subtle
variation in thermal preference.
367
Humidity, diet and light substantially influence thermal preference of flies independent of
Wolbachia infection
The analyses presented above show that the design of the thermal gradient device
plays an important role in T_p measurements. The tubular device consisted of more spare
metal parts than the plate device, which negatively affected the temperature conductivity.
Accordingly, we found that the effective temperature range along the gradient was much
narrower in the tubular compared to the plate device. The minimum temperature in the tube
never decreased below 17°C (maximum temperature: 27°C), whereas the plate reached
temperatures as low as 13°C (maximum temperature: 27°C). We therefore tested if the flies
were limited by the temperature range and were unable to choose preferred temperatures
below the minimum temperature in the tube. In this case, flies may cluster at the lower end
of the gradient in the tube device, which would result in T_p distributions skewed towards the
lower end, compared to the plate device. However, when comparing the skewness of T_p
distributions in replicate experiments among the two devices we did not find significant
differences (Wilcoxon Rank test, W=258, p=0.128) which suggests that the temperature
range within the tube did not limit the behavior of the test flies.
Conversely, cold temperatures as low as 10°C-13°C may cause reduced velocity[48]
which can lead to immobilization[49,50] as a physiological response. This may thus result in
biased T_p measurements due to a clustering of immobilized flies that get trapped at the cold
end of the gradient[18]. Similar to Hague et al.[16], we thus repeated testing for the influence of
Wolbachia on T_p in flies on the plate device by excluding flies with a T_p <15°C, to avoid a
potential bias from reduced mobility in cold temperatures. The analysis based on this
reduced dataset did not indicate a *Wolbachia*-specific influence on host T_p either (GLMM:
p=0.43, x^2=2.76, df=3) and thus did not qualitatively differ from results based on the full
dataset. This suggests that trapping due to immobilization at the cold end of the gradient is
unlikely a confounding factor in our experiments.
We speculate that also other environmental factors associated with the design of the
devices may strongly affect thermal behavior. According to Dillon *et al.*[18], sex, age, humidity,
light, circadian rhythms, feeding status and number of individuals tested at once all influence
thermal behavior. While similar numbers of flies that were tested simultaneously during each
experiment on both devices, the available space for each fly differed dramatically between
the two designs. In particular, the narrower space in the plate device may have led to

stronger interactions among the flies which could result in pronounced gregarism and

clustering due to social interactions at certain regions along the arenas[51]. To further

investigate how experimental design influences T_p measurement, we compared 15 studies

on thermal behavior in *D. melanogaster* flies. This qualitative meta-analysis revealed high

variation in measured T_p, which apparently depends on the design of the gradient device,

the range or the temperature gradient, rearing and experimental conditions, food, age of flies

and number of individuals tested (all summarized in **Table S3**).

Table 2. Statistical analyses of different environmental factors and experimental design influencing T_p in
***Drosophila melanogaster* infected with different *Wolbachia* types.** Significant effects of generalized mixed
models (GLMM) inferred by analysis of deviance are highlighted in bold.

(A) T_p in adults (plate vs tube device)			
Effect	$\chi2$	*df*	*p*-Value
machine	748.6	4	**< 2.2e-16**
infection	18.9	6	**4.4E-03**
machine*infection	4.6	3	0.20
(B) T_p in adults (humidity)			
Effect	$\chi2$	*df*	*p*-Value
humidity	266.8	4	**< 2.2e-16**
infection	15.1	6	**2.0E-02**
humidity*infection	2.2	3	0.52
(C) T_p in adults (food)			
Effect	$\chi2$	*df*	*p*-Value
food	24.1	2	**5.8E-06**
infection	0.2	2	0.92
food*infection	0.2	1	0.68
(D) T_p in adults (light)			
Effect	$\chi2$	*df*	*p*-Value
light	5.7	2	6.0E-02
infection	1.5	2	4.7E-01
light*infection	0.9	1	3.4E-01

To better understand the influence of specific environmental stimuli on thermal

preference assays, we experimentally investigated how humidity, food and light affect T_p

measurement in the context of varying *Wolbachia* infection status. We observed a highly

significant effect of humidity (GLMM: *p*<2.2e-16, x^2=266.93, *df*=4; **Table 2B**; **Figure 3**) on T_p

of adult flies. Higher humidity (60%) resulted in increased thermal preferences, elevated by

1-2°C (**Table S2**). The median T_p at 30% humidity ranged from 21.5°C (*w*MelCs and

*w*MelPop) to 22°C (*w*-), whereas the median T_p at 60% humidity ranged from 22.7°C (*w*-) to

23.5°C (wMel, wMelCS). A positive correlation between humidity and T_p was similarly
observed already in other insects[52,53].
Diet also had a strong influence on thermal behavior. We found that flies on in-house
food preferred 1°C lower temperatures than counterparts maintained on Carolina instant
food (GLMM: p=5.79e-6, x^2=24.2, df=2; **Table 2C**; **Figure 3**). Differences in diet quality or
feeding status (fed versus fasting) are known to influence thermal preference in ectotherms,
where rich food diet increases T_p compared to poor food or fasting conditions[54–57]. Our
results match these previous findings. While we did not assess the exact nutrient
composition in our in-house fly food, the recipe is considered less rich in carbohydrates and
proteins than Carolina 4-24 instant food. Additionally, diet is also known to affect the titer of
Wolbachia in different tissues: Protein-rich food has been found to boost titer levels in
somatic tissues, but not in ovaries. Conversely, carbohydrate-rich food can increase titer
levels in ovaries only[58,59]. If variation in titer levels may have a strong influence on thermal
preference, as speculated previously[14,16], the food quality may indirectly influence T_p by
modulating titer levels. However, we failed to observe a significant influence of light on T_p,
(GLMM: p=0.059, x^2=5.69, df=2; **Table 2D**; **Figure 3**)
Interestingly, in contrast to the experiments performed in our newly designed "plate"
gradient device, the follow-up experiments in the "tube", in particular the experiment testing
for the effects of humidity variation, revealed significant effects of *Wolbachia* infections on T_p
(see **Table 2A**). However, the observed patterns were opposite to expectations. Flies
infected with wMelCS preferred warmer temperatures than uninfected flies in both
experiments (Tukey's post hoc test: p<0.01; **Table S2**; **Figure 3**). These findings contradict
previously observed results where flies infected with wMelCS variant preferred lower
temperatures than uninfected counterparts[14,15]. Our results are, however, in line with the
data observed by Hague et al.[16], showing no shift of preference to higher temperatures in
flies artificially infected with wMelCS. In summary, these results further suggest that T_p
measurement is strongly influenced by the experimental setup and that T_p variation with
respect to *Wolbachia* infections is probably very subtle.
448

450 **Figure 3**. **The influence of environmental and experimental factors on T_p in *Wolbachia*-infected**

451 ***Drosophila melanogaster*. Boxplots showing the** influence of the design of the thermal gradient device,

452 humidity, food, and light on measurement of T_p in *D. melanogaster* and the influence of *Wolbachia* on host

453 thermal preference, where the colors highlight the different *Wolbachia* genotypes

Conclusions

Conclusions

In our thermal behavior assays we found that *Wolbachia* does not influence thermal

preference in early and late 3rd-instar *Drosophila* larvae, consistent with previous studies

suggesting that *Wolbachia* has no, or only very weak effects on juvenile traits of *D.*

melanogaster. In contrast, we found subtle effects of *Wolbachia* on thermal preference in

adult *D. melanogaster* under certain experimental conditions only. These effects were

inconsistent with previous data and indicate that T_p measurement assays are potentially

influenced and confounded by uncontrolled environmental cues. Such unwanted behavioral

stimuli can overwhelm subtle thermal preference differences during behavioral experiments.

As we show in our study, factors such as humidity, diet and design of the gradient device

substantially influence measurements of thermal preference in *D. melanogaster* infected with

different *Wolbachia* variants. Our findings demonstrate the necessity of a careful

experimental design and cautious interpretations of T_p measurements together with a

thorough description of the methods and equipment used to conduct behavioral studies. Our

data and findings thus provide important considerations when planning future behavioral

assays to assess thermal preference not only in *Drosophila* but also in other insects infected

with *Wolbachia*, such as vectors of human diseases like mosquitoes.

Acknowledgments

We are especially grateful to Wolfgang Miller who helped with the design of the experiment, who provided lab space and who conceptually supported this project. We are also indebted to Marcel Freund at the University of Zürich who not only hand-crafted the thermal gradient devices used in this study but also helped with their design. Moreover, we thank Thomas Flatt, Elisabeth Haring and Roman Arguello for helpful comments on previous versions of this manuscript. We are very grateful to Fabian Gstöttenmayr, Janis Jeschgo, Julia Mras, Elina Koivisto and Jasmin Jester who helped with fly maintenance and fly food cooking. This project was funded by a standalone grant from the Austrian Science Fund (FWF P32275) to Martin Kapun.

Authors Contributions

Anton Strunov involved in conceptualization, investigation, data curation, visualization, formal analysis, validation, writing—original draft; Charlotte Schoenherr involved in investigation, data curation and writing—review & editing; Martin Kapun involved in conceptualization, formal analysis, visualization, supervision, funding acquisition, writing— review & editing and project administration.

Data availability

All raw data and the *R* code to carry out the statistical analyses can be found in the Supplementary Information files Strunov_etal_WolbTP_2023_RawData.xlsx and Strunov_etal_WolbTP_2023_Rcode.zip, respectively. A more detailed description of the devices, methods and statistical approaches used in this manuscript can be found online at https://github.com/capoony/DrosophilaThermalGradient

References

1. Huey, R. & Berrigan, D. Temperature, demography, and ectotherm fitness. **158**,

204–10 (2001).

2. Hoffmann, A. A. Physiological climatic limits in Drosophila: patterns and

implications. *J. Exp. Biol.* **213**, 870–880 (2010).

3. Angilletta, M. J. Temperature, Growth Rate, and Body Size in Ectotherms: Fitting

Pieces of a Life-History Puzzle. *Integr. Comp. Biol.* **44**, 498–509 (2004).

4. Stevenson, R. D. The Relative Importance of Behavioral and Physiological

Adjustments Controlling Body Temperature in Terrestrial Ectotherms. *Am. Nat.*

126, 362–386 (1985).

5. Gilbert, A. L. & Miles, D. B. Natural selection on thermal preference, critical

thermal maxima and locomotor performance. *Proc. R. Soc. B Biol. Sci.* **284**,

20170536 (2017).

6. Hart, B. L. Biological basis of the behavior of sick animals. *Neurosci. Biobehav.*

Rev. **12**, 123–137 (1988).

7. Watson, D. W., Mullens, B. A. & Petersen, J. J. Behavioral Fever Response of

Musca domestica (Diptera: Muscidae) to Infection by Entomophthora muscae

(Zygomycetes: Entomophthorales). *J. Invertebr. Pathol.* **61**, 10–16 (1993).

8. Inglis, G. D., Johnson, D. L. & Goettel, M. S. Effects of Temperature and

Thermoregulation on Mycosis byBeauveria bassianain Grasshoppers. *Biol.*

Control **7**, 131 (1996).

9. de Roode, J. C. & Lefèvre, T. Behavioral Immunity in Insects. *Insects* **3**, 789–820

(2012).

10. Poulin, R. Parasite Manipulation of Host Behavior. in *Advances in the Study of*

Behavior vol. 41 151–186 (Elsevier, 2010).

11. Curtis, V. A. Infection-avoidance behaviour in humans and other animals. *Trends*

Immunol. **35**, 457–464 (2014).

12. Adamo, S. A. The specificity of behavioral fever in the cricket Acheta domesticus.

J. Parasitol. **84**, 529–533 (1998).

13. Fedorka, K. M., Kutch, I. C., Collins, L. & Musto, E. Cold temperature preference

in bacterially infected Drosophila melanogaster improves survival but is

remarkably suboptimal. *J. Insect Physiol.* **93–94**, 36–41 (2016).

14. Truitt, A. M., Kapun, M., Kaur, R. & Miller, W. J. *Wolbachia* modifies thermal

preference in *Drosophila melanogaster*. *Environ. Microbiol.* **21**, 3259–3268

(2019).

15. Arnold, P. A., Levin, S. C., Stevanovic, A. L. & Johnson, K. N. Wolbachia-infected

Drosophila prefer cooler temperatures. *Ecol. Entomol.* **44**, 287–290 (2019).

16. Hague, M. T. J., Caldwell, C. N. & Cooper, B. S. Pervasive Effects of Wolbachia

on Host Temperature Preference. *mBio* **11**, e01768-20 (2020).

17. Mirth, C. K., Saunders, T. E. & Amourda, C. Growing Up in a Changing World:

Environmental Regulation of Development in Insects. *Annu. Rev. Entomol.* **66**,

81–99 (2021).

18. Dillon, M. E., Wang, G., Garrity, P. A. & Huey, R. B. Thermal preference in

Drosophila. *J. Therm. Biol.* **34**, 109–119 (2009).

19. Teixeira, L., Ferreira, Á. & Ashburner, M. The Bacterial Symbiont Wolbachia

Induces Resistance to RNA Viral Infections in Drosophila melanogaster. *PLoS*

Biol. **6**, e1000002 (2008).

20. Zhou, W., Rousset, F. & O'Neil, S. Phylogeny and PCR-based classification of

Wolbachia strains using wsp gene sequences. *Proc. R. Soc. B Biol. Sci.* **265**,

509–515 (1998).

21. Riegler, M., Iturbe-Ormaetxe, I., Woolfit, M., Miller, W. J. & O'Neill, S. L. Tandem

repeat markers as novel diagnostic tools for high resolution fingerprinting of

Wolbachia. *BMC Microbiol.* **12**, S12 (2012).

22. Rajpurohit, S. & Schmidt, P. S. Measuring thermal behavior in smaller insects: A

case study in Drosophila melanogaster demonstrates effects of sex, geographic

origin, and rearing temperature on adult behavior. **10**, 1–13 (2016).

23. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25

years of image analysis. *Nat. Methods* **9**, 671–675 (2012).

24. Chen, S.-C. *et al.* Dorsal clock networks drive temperature preference rhythms in

Drosophila. *Cell Rep.* **39**, 110668 (2022).

25. Sayeed, O. & Benzer, S. Behavioral genetics of thermosensation and

hygrosensation in Drosophila. **93**, 6079–6084 (1996).

26. Yamamoto, A. & Ohba, S. Temperature preferences of eleven *Drosophila*

species from Japan: the relationship between preferred temperature and some

ecological characteristics in their natural habitats. *Zoolog. Sci.* **1**, 631–640

(1984).

27. R Core Team. R Foundation for Statistical   Computing. (2019).

28. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects

Models Using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

29. Fox, J. & Weisberg, S. *An R Companion to Applied Regression*. (Sage, 2019).

30. Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. emmeans:

Estimated Marginal Means, aka Least-Squares Means. (2019).

31. Komsta, L. & Novomestky, F. moments: Moments, Cumulants, Skewness,

Kurtosis and Related Tests. (2022).

32. Sokabe, T., Chen, H.-C., Luo, J. & Montell, C. A Switch in Thermal Preference in

Drosophila Larvae Depends on Multiple Rhodopsins. *Cell Rep.* **17**, 336–344

(2016).

33. Tyrrell, J. J., Wilbourne, J. T., Omelchenko, A. A., Yoon, J. & Ni, L. Ionotropic

Receptor-dependent cool cells control the transition of temperature preference in

Drosophila larvae. *PLOS Genet.* **17**, e1009499 (2021).

34. Al-Saffar, Z. Y., Grainger, J. N. R. & Aldrich, J. Temperature and humidity

affecting development, survival and weight loss of the pupal stage of Drosophila

melanogaster, and the influence of alternating temperature on the larvae. *J.*

Therm. Biol. **21**, 389–396 (1996).

35. Min, K.-T. & Benzer, S. Wolbachia, normally a symbiont of Drosophila, can be

virulent, causing degeneration and early death. *Proc. Natl. Acad. Sci.* **94**, 10792–

10796 (1997).

36. Osborne, S. E., Leong, Y. S., O'Neill, S. L. & Johnson, K. N. Variation in Antiviral

Protection Mediated by Different Wolbachia Strains in Drosophila simulans.

PLOS Pathog. **5**, e1000656 (2009).

37. Strunov, A., Kiseleva, E. & Gottlieb, Y. Spatial and temporal distribution of

pathogenic Wolbachia strain wMelPop in Drosophila melanogaster central

nervous system under different temperature conditions. *J. Invertebr. Pathol.* **114**,

22–30 (2013).

38. Chrostek, E. *et al.* Wolbachia Variants Induce Differential Protection to Viruses in

Drosophila melanogaster: A Phenotypic and Phylogenomic Analysis. *PLOS*

Genet. **9**, e1003896 (2013).

39. Martinez, J. *et al.* Symbiont strain is the main determinant of variation in

Wolbachia-mediated protection against viruses across Drosophila species. *Mol.*

Ecol. **26**, 4072–4084 (2017).

40. Strunov, A. A., Ilinskii, Yu. Yu., Zakharov, I. K. & Kiseleva, E. V. Effect of high

temperature on survival of Drosophila melanogaster infected with pathogenic

strain of Wolbachia bacteria. *Russ. J. Genet. Appl. Res.* **3**, 435–443 (2013).

41. Chrostek, E., Martins, N., Marialva, M. S. & Teixeira, L. Wolbachia-Conferred

Antiviral Protection Is Determined by Developmental Temperature. *mBio* **12**,

e02923-20 (2021).

42. Pontier, S. M. & Schweisguth, F. A Wolbachia-Sensitive Communication between

Male and Female Pupae Controls Gamete Compatibility in Drosophila. *Curr. Biol.*

CB **25**, 2339–2348 (2015).

43. Detcharoen, M., Arthofer, W., Jiggins, F. M., Steiner, F. M. & Schlick-Steiner, B.

C. Wolbachia affect behavior and possibly reproductive compatibility but not

thermoresistance, fecundity, and morphology in a novel transinfected host,

Drosophila nigrosparsa. *Ecol. Evol.* **10**, 4457–4470 (2020).

44. Strunov, A., Lerch, S., Blanckenhorn, W. U., Miller, W. J. & Kapun, M. Complex

effects of environment and Wolbachia infections on the life history of Drosophila

melanogaster hosts. *J. Evol. Biol.* **35**, 788–802 (2022).

45. Gutzwiller, F. *et al.* Dynamics of *Wolbachia pipientis* Gene Expression Across the

Drosophila melanogaster Life Cycle. *G3 GenesGenomesGenetics* **5**, 2843–2856

(2015).

46. Sokolowski, M. B., Kent, C. & Wong, J. Drosophila larval foraging behaviour:

Developmental stages. *Anim. Behav.* **32**, 645–651 (1984).

47. Charlesworth, D. & Willis, J. H. The genetics of inbreeding depression. *Nat. Rev.*

Genet. **10**, 783–796 (2009).

48. Crill, W. D., Huey, R. B. & Gilchrist, G. W. Within- and Between-Generation

Effects of Temperature on the Morphology and Physiology of Drosophila

Melanogaster. *Evolution* **50**, 1205–1218 (1996).

49. Davis, H. E., Cheslock, A. & MacMillan, H. A. Chill coma onset and recovery fail

to reveal true variation in thermal performance among populations of Drosophila

melanogaster. *Sci. Rep.* **11**, 10876 (2021).

50. Giraldo, D., Adden, A., Kuhlemann, I., Gras, H. & Geurten, B. R. H. Correcting

locomotion dependent observation biases in thermal preference of Drosophila.

Sci. Rep. **9**, 3974 (2019).

51. Fogleman, J. C. Oviposition site preference for substrate temperature

inDrosophila melanogaster. *Behav. Genet.* **9**, 407–412 (1979).

52. Gunn, D. L. & Cosway, C. A. The Temperature and Humidity Relations of the

Cockroach: V. Humidity Preference. *J. Exp. Biol.* **15**, 555–563 (1938).

53. Deal, J. The Temperature Preferendum of Certain Insects. *J. Anim. Ecol.* **10**, 323

(1941).

54. Thomson, R. C. M. The Reactions of Mosquitoes to Temperature and Humidity.

Bull. Entomol. Res. **29**, 125–140 (1938).

55. Chapman, R. F. The Behaviour of Nymphs of Schistocerca Gregaria (Forskål)

(Orthoptera, Acrididae) in a Temperature Gradient, With Special Reference To

Temperature Preference. *Behaviour* **24**, 283–317 (1965).

56. Angilletta, M. J., Niewiarowski, P. H. & Navas, C. A. The evolution of thermal

physiology in ectotherms. *J. Therm. Biol.* **27**, 249–268 (2002).

57. Pulgar, J. M., Aldana, M., Bozinovic, F. & Ojeda, F. P. Does food quality

influence thermoregulatory behavior in the intertidal fish Girella laevifrons? *J.*

Therm. Biol. **28**, 539–544 (2003).

58. Serbus, L. R. *et al.* The Impact of Host Diet on Wolbachia Titer in Drosophila.

PLOS Pathog. **11**, e1004777 (2015).

59. Ponton, F. *et al.* Macronutrients mediate the functional relationship between

Drosophila and Wolbachia. *Proc. R. Soc. B Biol. Sci.* **282**, (2015).

60. Head, L. M. *et al.* The influence of light on temperature preference in Drosophila.

Curr. Biol. CB **25**, 1063–1068 (2015).