Discovery of multi-state gene cluster switches determining the adaptive mitochondrial and metabolic landscape of breast cancer

Michela Menegollo[1*&] , Robert B. Bentham[2*$], Tiago Henriques[1], Seow Qi Ng[2], Ziyu Ren[2], Clarinde Esculier[2], Sia Agarwal[2], Emily Tong[2], Clement Lo[2], Sanjana Ilangovan[2], Zorka Szabadkai[2], Matteo Suman[1], Neill Patani[2,6], Avinash Ghanate[6], Kevin Bryson[3$], Robert C. Stein[4,5], Mariia Yuneva[6#], Gyorgy Szabadkai[1,2,6#&]

[1]Department of Biomedical Sciences, University of Padova, Padova, 35121, Italy
[2]Department of Cell and Developmental Biology, Consortium for Mitochondrial Research, University College London, London, WC1E 6BT, UK
[3]Department of Computer Sciences, University College London, London, WC1E 6BT, UK
[4]Department of Oncology, University College London Hospitals, London, NW1 2PG, UK
[5]UCL Cancer Institute, University College London, London, WC1E 6DD, UK
[6]The Francis Crick Institute, London, NW1 1AT, UK

* shared first authorship
# shared senior authorship

$ current address: RB: Cancer Genome Evolution Research Group, Cancer Research UK Lung Cancer Centre of Excellence, University College London Cancer Institute, London, WC1E 6BT, UK; KB: University of Glasgow, School of Computing Science, Glasgow G12 8RZ, UK

& correspondence to:
Gyorgy Szabadkai, g.szabadkai@ucl.ac.uk
Michela Menegollo, michela.menegollo@unipd.it
Robert B. Bentham, robert.bentham.11@ucl.ac.uk

Summary

Adaptive metabolic switches are proposed to underlie conversions between cellular states during normal development as well as in cancer evolution, where they represent important therapeutic targets. However, the full spectrum, characteristics and regulation of existing metabolic switches are unknown. We propose that metabolic switches can be recognised by locating large alternating gene expression patterns and associate them with specific metabolic states. We developed a method to identify interspersed genesets by massive correlated biclustering (MCbiclust) and predict their metabolic wiring. Testing the method on major breast cancer transcriptome datasets we discovered a series of gene sets with switch-like behaviour, predicting mitochondrial content, activity and central carbon fluxes in tumours associated with different switch positions. The predictions were experimentally validated by bioenergetic profiling and metabolic flux analysis of ^{13}C-labelled substrates, and were ultimately extended by geneset analysis to link metabolic alterations to cellular states, thus predicting tumour pathology, prognosis and chemosensitivity. The method is applicable to any large and heterogeneous transcriptome dataset to discover metabolic and associated pathophysiological states.

Introduction

Cancer development and evolution entail diverging routes of adaptation in cellular energetics and metabolism to oncogenic signals and the tumour environment[1,2]. These adaptive metabolic rearrangements often result in tumour specific vulnerabilities to pharmacological or dietary targeting, or to the contrary, promote therapy resistance[3]. Thus, evaluating the patterns metabolic adaptation is essential in order to understand tumour biology and to devise novel therapeutic strategies.

However, no comprehensive evaluation of metabolic adaptation patterns has yet been completed. The patterns are presumed to accompany both transient alterations of cellular phenotype or inheritable phenotype switches underlying transitions in cellular states. The states can be determined by cell type, its environment and the cell's position in the differentiation tree, thus transitions in cell state would follow changes in differentiation status, transdifferentiation, e.g. epithelial-mesenchymal transition (EMT), cellular senescence, or reversion to an immature stem or progenitor phenotype[4,5], and to other cell type specific states[6]. Interest has surged recently in metabolic adaptations during cell state transitions[7,8], leading to considerable knowledge of cell state specific metabolic alterations at the metabolite and flux levels[2,9–13]. However, using metabolites and fluxes as exclusive readouts may not address whether the altered metabolic states represent *bona fide* functional adaptations or e.g. simply mass action driven rearrangements of metabolic networks[14]. Moreover, to explore the relationship and eventual co-regulation of the functional adaptation with cellular state, metabolic measurements themselves are insufficient.

Alterations in gene expression patterns (GEPs) are almost invariably associated with cellular state transitions, transformation and bear important prognostic value[15–17]. Accordingly, GEPs can provide information to link cellular state to metabolic adaptation at the gene expression level by exploring the metabolic transcriptome in large scale GEPs, including expression of metabolic pathway components and nuclear encoded mitochondrial genes, together representing >15% of the genome. Large patterns of alteration in the transcriptome, often involving >1000 genes, can arise either from genetic (oncogenic mutations) or non-genetic (environmental) influences, which ultimately determine cellular phenotype and the associated cellular metabolic network[18]. Earlier studies on cancer related metabolic rearrangements mostly focused on the effect of cell autonomous oncogenic signals on specific metabolic pathways[19,20], delineating the metabolic switches required for increased proliferation and growth, or adaptation to hypoxia[1,21]. However, it is now well recognised that other cellular states or features can also be associated with altered gene expression patterns affecting metabolic networks, including not only the fundamental EMT transdifferentiation in cancer metastasis, but other pathological and physiological states[15,16,22,23]. Thus, in order to find patterns of metabolic adaptation, we hypothesised that alterations in metabolic gene expression patterns reflect *bona fide* adaptive transcriptional regulation, driven by a specific cell state. Accordingly, metabolic switches signal transitions between cell states, and apart from being clear pharmacological targets[24], can may well be used for important pathological predictions associated with cell states.

A corollary to such an analysis, and thus our primary aim, is to identify the differential regulation patterns in the full metabolic transcriptome and link it quantitatively with the large scale GEPs associated with different cancer cell states. Further, we aimed to correlate the states with functional metabolic and energy flux measurements as well as to understand their association with tumour biology and their prognostic value. Identifying these patterns

will help (i) the unbiased mapping of potential metabolic adaptation strategies in tumours, (ii) to understand and exploit metabolic vulnerabilities in different cell states, represented as tumour subtypes; and in particular (iii) to recognise and target adaptive patterns resulting in drug-resistant states.

Only a few previous studies have analysed large scale metabolic gene expression data across multiple cancer types, mostly revealing widespread deregulation of metabolic pathways compared to adjacent normal tissue. These approaches consisted of diverse pathway scoring methods based on differential gene expression in numerous cancer types and subtypes[15,17,25,26], or genomic changes associated with transcriptional adaptation to the environment[21]. To discover shared deregulation of individual genes of metabolic pathways, Reznik et al. [16] analysed differential co-expression in a number of cancer types, revealing recurrent loss of co-expression of a set of genes, including mitochondrial electron transport chain components. However, these studies did not account for large correlated expression patterns of broader sets of genes, which inevitably underlies the complete metabolic rewiring and phenotypic (cell state) switching. The patterns are likely to be aligned with specific cell states, thus revealing subtype specific dependencies as novel pharmacological targets, or predict pathological features and clinical behaviour. In addition, defining these patterns will help answer questions such as what drives the deregulation of metabolic pathways, and whether it is adaptive thus targetable, as well as how gene expression and metabolic fluxes correlate. No current approaches to our knowledge addressed this problem.

For the unbiased discovery of large co-regulated gene expression patterns, we simplified our model to include the concerted on and off switching of large gene sets[27], such as bi- and multistate, or bi- and multistable gene switches. The switches thus create anticorrelated gene sets in a selected set of samples representing two different cell states (or cancer subtypes). Here, we created a pipeline for the discovery and further analysis of these large gene expression switches, which can be used as a resource to any large, heterogeneous datasets. To demonstrate the workflow and the value of the resource, we use it to discover metabolic and mitochondrial states (subtypes) of breast cancer, aligning with their cell state determined by the cell of origin. First, using a number of large breast cancer transcriptome datasets, we discover large anticorrelated gene sets in their specific sample set (biclusters), using the central tool of the resource, MCbiclust, finding massive correlated biclusters in large datasets[28]. The biclusters containing the large inverted gene sets define breast cancer metabolic subtypes, dominated by mitochondrial genes, and likely the result of metabolic switches following the cell of origin pattern. Next, we correlated the gene expression patterns with metabolic fluxes, measured in cellular models by mass spectrometry, using heavy carbon labelled substrates. Finally, we show the utility of the resource to predict the metabolic phenotype of cancer cell lines and cancer tissue, to predict metabolic wiring associated with certain cell states and phenotypes, and importantly, to predict chemotherapeutic response of a subset of estrogen receptor (ER) positive tumours, together with overall prognosis.

.

Results

1. Discovery of large gene cluster switches, defining breast tumour mitochondrial and metabolic subtypes

In order to identify co-regulated gene clusters in subgroups of breast cancer samples we used our previously established massive correlating biclustering method, MCbiclust[28]. The method is designed to find subsets of samples in which large genesets, such as sets of metabolic and nuclear encoded mitochondrial genes[29,30], are co-regulated, as measured by the correlation of individual gene expression levels across the selected samples. Technically, an iterative stochastic greedy search method selects the group of samples across which the chosen gene set reaches the maximum absolute correlation score. The samples and gene` set obtained with the highest score define a bicluster.

As an additional feature, MCbiclust using the absolute correlation values for sample selection leads to finding two groups of genes with maximal positive correlations within the groups, but negative correlation between each other. Thus, two genesets with mutually exclusive expression are obtained in each bicluster, defining two groups of samples (see[29] and Figure 1A). Importantly, after defining the bicluster, quantifications of correlations are extended to the (i) whole sample set using the similarity of each sample to the bicluster sample set, ranking them according to the correlation of their bicluster gene set expression with that of the bicluster samples, as well as to the (ii) whole transcriptome, using the correlation of each gene transcript with the average expression of a set of representative bicluster genes (correlation vector values, CV). Thus samples and gene sets can be analysed independently (for the schematics of the method see Figure 1A).

Figure 1

Figure 1. Identification of large scale transcriptome switches

A. Geneset switch identification in bicluster samples. MCbiclust starts with a random geneset in a random sample set, and performs a greedy search aiming for the sample set across which the geneset has the highest correlation. Since the *absolute* value of correlation is maximised during the search, the resulting top correlating samples will display two genesets with sharp anticorrelation (switch) between them. The genesets and samplesets can be extended to the whole transcriptome and sampleset by quantifying correlations (see main text), which provides the basis for gene set enrichment analysis

B. Visualisation of individual switches (MB_1, MB_2, MB_3) in 2D distribution by PC1 and ranking index in the METABRIC dataset. Pam50 classifiers are overlaid to show the differential distribution of samples in the different switches (biclusters). Dashed lines represent the thresholds for belonging to a fork, calculated as 0.04 * forkscale (see Methods).

C. UpSet matrix and plots of aggregates of sample intersections between upper and lower positions of each switch (MB_1, MB_2, MB_3). Unique and overlapping (intersect) samples are shown separately.

D, E. Interswitch relationships are shown in 2D distribution plots of samples by two biclusters (switches). The axes represent the scale between the upper (UF) and lower (LF) of each bicluster. The distribution of samples according to their MB_1 vs MB_2 forkscale value (See Methods) is shown in **D**, whilst the combinations of MB_3 with MB_1 or MB_2 are shown in **E**.

F. Main heatmap shows distribution of the correlation values (CVs, Pearson r) of nearly the whole transcriptome in the MB_1 bicluster; a small set of midrange values were hidden (see gaps in heatmap). Annotations show the locations of mitochondrial and metabolic genes (top), and that of the bicluster genes (bottom). For comparison between the switches, bottom annotation curves show the CV values of each sample in the three biclusters (switches).

G. Fold enrichment of mitochondrial and metabolic (with intersections) genes in the anticorrelated gene sets. Heatmap shows enrichment values of each geneset in each position of all switches. Geneset intersections are illustrated at the bottom.

H. Scheme of switches. MB1 and MB2 share their LF, thus representing a multistate switch, while MB3 is an independent bistable switch, both MB1_2 and MB_3 representing different switch mechanisms (see Main text).

We applied the method to three independent breast cancer transcriptome datasets, METABRIC[31], TCGA[32] and Oslo2[33] and obtained equivalent results. MCbiclust identified three independent biclusters in the METABRIC dataset (termed mitochondrial biclusters MB1, MB2 and MB3, Figure S1A-C), indicating three different large genesets that are strongly co-regulated (including both + or - correlation) in specific subsets of the METABRIC samples. Accordingly, each sample in the bicluster will belong to either of the two positions of a toggle switch, splitting up samples in two groups (termed upper fork, UF and lower fork, LF, see below) driven by their unique gene sets, anti-correlated between the sample groups. To visualise the switches, for each bicluster we plotted the ranking of samples (according to their correlation strength, i.e. the average CV of their bicluster genes) against the differences in the correlation in their gene sets, captured by the first principal component (PC1) of each sample (Figure 1B). The distribution pattern of the samples showed the split nature of the biclusters, with bifurcating PC1 values in highly ranked samples. The same toggle switches were confirmed in the TCGA and Oslo cohorts (Figure S1D,E).

Next, we compared the sample composition across the biclusters and explored the gene sets dominating the clusters. While most forks of the three biclusters were composed of unique samples (Figure 1C), MB1_LF and MB2_LF showed a near complete overlap, and thus were aggregated in a common cluster (MB12_LF). To define the position of the samples in all of the discovered biclusters, we calculated their 'forkscale', indicating how close they are to the tip of the upper (+1) or lower (-1) forks and compared these values between the biclusters (Figure 1D, E). Remarkably, comparison the MB1 and MB2 bicluster switches gives a tripolar distribution, where samples are merged in a common lower fork (MB12_LF) and split between two distinct upper forks, MB1_UF and MB2_UF, indicating the presence of a multi-state switch. Comparison of MB3 with either MB1 or MB2 did not give any significant overlap, indicating independently regulated gene cohorts in the MB3 bicluster.

To define the gene sets dominating the biclusters we have ranked the whole transcriptome based on the CV of each gene, indicating its correlation with the switch pattern. Figure 1F shows the correlation map of the MB1 bicluster. Genes with the highest and lowest CV values were considered the dominant, most strongly regulated genes in the UF and LF switch positions, respectively. Expression levels of these genes are anticorrelated between the UF and LF samples (see Figures 1A and S1F). Importantly, genes with the highest CV values (in both UF and LF switch directions) include not only bicluster genes, indicating that they are part of larger scale gene regulation patterns (Table S1). The patterns and their relationship with the bicluster genes are explored in the next section.

Notably, nuclear encoded mitochondrial and metabolic genes were markedly over-represented in the UF of the MB1 and MB2 bicluster patterns and excluded from the MB12_LF as shown by the metabolic and mitochondrial gene annotation in Figure 1F. The relative enrichment of nuclear encoded mitochondrial and metabolic genes are shown in Figure 1G.

Altogether, MCbiclust identified large scale gene expression switches defining breast cancer subtypes, which include the highly controlled co-regulation of a large group of metabolic and mitochondrial genes. The gene sets identified (see Table S1), showed switch-like behaviour, schematically represented in Figure 1H.

2. Defining metabolic patterns of mitochondrial breast cancer subtypes and their association with cell state

Using the gene and sample sets generated by MCbiclust, we created a pipeline to generate a series of predictions, covering tumour metabolism, oncogenic signalling and clinical outcome. First, in order to understand the transcriptional metabolic signatures (TMS) of the identified switches, we mapped all metabolic genes in all three switches on the global KEGG metabolic pathway map (Figure S2A). These maps indicated that the TMSs comprise all major metabolic modules, including carbohydrate, lipid, nucleotide, amino acid, glycan, cofactor and energy metabolism. Importantly, in each switch, the TMSs of the UF and LF sample sets were associated with either mutually exclusive pathways (e.g. lipid synthesis versus degradation), often with enzyme isotype switches or alternative usage of substrates (e.g. glycolytic versus glutaminolytic energy production). For a more detailed *in silico* analysis of the metabolic phenotypes predicted by the method, we focused on the MB12_LF vs MB1_UF/MB2_UF bi-state switch, using two approaches. First, we compared the CVs of each gene of individual metabolic pathways, highlighting differences in their association with different switch positions. As shown in Figure 2A, in central carbon metabolism, differential correlation (i.e. difference in CVs in the upper and lower switch positions) of glycolytic and glutaminolytic enzyme transcripts demonstrated the transcriptional basis of metabolic phenotype switches, as well as the difference between MB1 and MB2 biclusters (further pathways are shown in Figure S2B). Second, to identify individual genes defining the three states, we used the CVs of the whole genome in the MB1 and MB2 bicluster switches, and identified groups of genes distinguishing the forks (CVs differ by at least 0.4 between the switches and the absolute value of the CV is higher than 0.6), as well as those common to both biclusters (Table S2). The resulting gene sets were then filtered for metabolic and mitochondrial genes (see Figure 1F, G) used for pathway analysis, of which the key findings are shown in Figure 2B. Importantly, using both KEGG pathways and the MitoCarta3.0 gene set and ontology[34], both the MB1_UF and MB2_UF switch positions showed significant activation of mitochondrial biogenesis pathways (KEGG: oxidative phosphorylation, TCA cycle; Mitocarta: mitochondrial central dogma: including mtDNA replication, transcription and translation on the mitochondrial ribosomes), as well as increased OXPHOS components (Figure S2B, Tables S3 and S4). These results indicated that the metabolic switch between the MB12_LF and both the MB1_UF and MB2_UF positions encompass similarly increased mitochondrial biogenesis, and predict higher electron flux through the TCA cycle and OXPHOS as compared to MB12_LF. While most of the intermediary metabolic pathways, with associated nucleotide, lipid and amino acid metabolism were more represented in MB1_UF and MB2_UF, both cellular (KEGG) and mitochondrial (Mitocarta) glutamate metabolism had stronger association with MB12_LF, representing an almost uniquely associated pathway in this switch. These results indicated altered substrate preference. We noted that whilst the MB1_UF and MB2_UF positions both showed higher mitochondrial biogenesis, the pattern of correlation of individual metabolic genes differed in some key points between these switch positions. Nucleotide, fatty acid and glutamate metabolism was strongly regulated in MB1_UF, switching to pyruvate metabolism, TCA cycle with enhanced pentose phosphate shunt in MB2_UF, reflecting the activation of distinct metabolic switch mechanisms, presumably due to differences in cell states and their supporting metabolic configuration, driven by cell state specific signalling.

Figure 2

Figure 2. Decoding the transcriptome switch from individual genes to large scale cell state patterns.

A, B. Metabolic phenotype predictions in the MB_1 and MB_2 bicluster switches. CV values (association with bicluster, see Figure 1 and Methods) of individual genes in the METABRIC breast cancer dataset analysis shown for pathways: glucose, glutamine, pyruvate metabolism and the TCA cycle in **A**. Values are ranked by the MB_1 CV value. The figure allows the identification of high CV genes and where differences are large between the switches (biclusters). The representation of pathways according to the relative number of genes affected is shown in **B** as a heatmap (top: based on top metabolic KEGG pathways; bottom: top MitoCarta3.0 ontology pathways, see Table S5).

C. Heatmap of distribution of the top 250 most variable PARADIGM concepts across the bicluster switches in the TCGA BRCA dataset. The term values were obtained from [35] pan cancer analysis and hierarchically clustered. Annotations show proliferation, tumour purity, switch identity (UF and LF of MB_1, MB_2 or MB_3), Pam50 and histology classifications. Transcriptomic signatures based on PARADIGM concepts were assigned as shown at the right.

D, E. Correlation of known mammary gland cell states and the transcriptomic switches in the METABRIC dataset. 2D distribution plots of samples by two biclusters is shown in **D**, overlaid by either cell of origin marker enrichment values (obtained by gene set variation analysis, GSVA - see Methods; mL: mature luminal, LP: luminal progenitor, MaSC: mammary stem cell, genesets from [36])

Thus, next we asked whether the observed metabolic switch positions with divergent mitochondrial biogenesis follow the activation of specific signalling pathways. To infer signalling pathway activity in individual samples, we analysed the distribution of the results of a previously implemented PARADIGM analysis[37] on a pan-cancer dataset[35] among different switches. Figure 2C shows the clustering of the most divergent PARADIGM pathway entities and their association with different biclusters and forks. Overall, the switch between lower and upper fork positions is dominated by turning on a cell proliferation cluster (MYB, FOXM1, E2F family, *P2*, Figure 2C) in the upper forks (MB1_UF, MB2_UF), or a stress and hypoxia (MAPK, p53, HIF1, *P5-7*) and stemness (p63) signalling cluster in the lower forks (MB12_LF). Interestingly, the MB1_UF and MB2_UF samples clustered separately, likely due to activation of Myc in MB2_UF together with a miRNA pattern (*P1*). While both MB1_UF and MB12_LF switch positions showed strong estrogen receptor signalling (*P4*), MB2_UF contained only ER negative samples (for full gene list see Table S6, for statistical analysis see Figure S3). All bicluster switches contained only a fraction of samples with high immune signature, indicating that differences between the forks is not determined by the immune system.

Taken together with the metabolic gene expression data, our model predicts higher mitochondrial content in MB1_UF and MB2_UF as compared to MB12_LF, which can be thus driven by the cell cycle and proliferation gene network, activated in the upper switch positions (see Figure 2C). The differences between MB1_UF and MB2_UF can be assigned to Myc activation. Conversely, MB12_LF, showed no transcriptional (co-)activation of nuclear encoded mitochondrial genes, but rather demonstrated a clear switch in substrate preference to glutamate from glucose derived pyruvate, a metabolic phenotype associated with stress signalling.

Finally, considering that metabolic switches have been associated with cell states and cell fate decisions, such as stemness and differentiation[38,39], we assessed whether the identified metabolic and signalling switches among groups of breast cancer samples relate to their cell of origin, i.e. their state of stemness or differentiation in a specific direction. Although intrinsic subtypes of breast cancer have been recently associated with specific states of mammary epithelial cell differentiation[40], we have found only a loose association between switch states and intrinsic subtypes (Figure S4A). Thus, we used an unbiased approach by calculating GSVA scores (gene set variance analysis[41]) for gene expression patterns derived from mammary stem cells (MaSC), luminal progenitors (LP) and mature luminal cells (mL)[36], and overlaying the values of each cell-of-origin pattern for each sample on the fork distribution plots (Figures 2D, S4B). Strikingly, there was an apparent association between the three cell states and the MB12_LF / MB1_UF / MB2_UF multi-state switch (Figure 2D). Average scores for all three cell states were significantly higher than for any of the intrinsic (PAM50) subtypes (Figure S4C). Stemness (MaSC) was largely accompanied by MB12_LF features, such as low mitochondrial abundance and glutamate substrate preference, whilst switching to either mL or LP cell state was followed by increased mitochondrial biogenesis driven by proliferation signalling in absence (MB1_UF: mL state) or presence (MB2_UF: LP state) of Myc/miRNA signals (Figure 2E).

3. Mitochondrial and metabolic gene expression profiles predict functional metabolic switches

For functional verification of the predicted metabolic switches, first we aimed to detect the identified gene expression switches in cellular and *in vivo* breast tumour models. Using either our custom or GSVA scoring methods on breast cancer cell line transcriptome datasets[42–44], we identified cell lines belonging to each position of the three bicluster switches, demonstrating that cellular models represent appropriately the full spectrum of transcriptional signatures identified in human samples (Figure 3A, Table S7). On the other hand, in a collection of patient-derived breast cancer xenograft (PDX) models[45], the MB1-2_LF position was not represented among the successfully implanted PDXs, even if all the biclusters and forks were discoverable in the original tumours (Figure 3B). Thus, we used cell line models to explore two hypotheses based on the predictions from computational analysis: cells with MB1_UF phenotype (i) boast higher mitochondrial content and activity, and (ii) have altered substrate preferences as compared with the MB1_LF phenotype. We used MCF7 and T47D cell lines to represent the MB1_UF switch and MDA-MB-436, Hs578t and HCC1134 cell lines for modelling the MB1_LF phenotype.

Figure 3

Figure 3. Mitochondrial function: MB1_UF cells show active functional biogenesis as compared to the MB1_LF phenotype.

A. Cell line scoring to reveal cell lines belonging to the observed switches. Volcano plot of the scores (see Methods, Table S7) and $-\log_{10}$ of the p-values in the MB1_UF / MB1_LF switch, with the cell lines chosen for experimental work highlighted in red.

B. 2D distribution (PC1 vs ranking index, shown for the MB1 bicluster switch) of biopsies obtained for generating PDX samples from the study [45], overlaid in left panel: with Pam50 categories to highlight the similarity to the MB1 switch (see Figure 1B) or in right panel: with growth success whether the sample has grown from the biopsy to PDX status.

C. Oroboros evaluation of ~1M cells of the cell lines grouped by their switch state (MB1_UF / MB1_LF). Reserve, ATP synthesis coupled and uncoupled respirations were quantified using FCCP and oligomycin treatments (See Methods). 2-way ANOVA with Sidak's multiple comparisons test.

D. Average cellular TMRM intensity grouped by switch state (MB1_UF / MB1_LF), data obtained from high content microscopy from > 1K cells belonging to either switch state. Unpaired Student's t-test.

E. Relative expression of respiratory chain subunits quantified from western blot analysis. Results are grouped by switch state (MB1_UF / MB1_LF). 2-way ANOVA with Sidak's multiple comparisons test.

F. High content imaging analysis of mitochondrial biogenesis, structure and function in MB1_UF and MB1_LF cells. Cells were loaded with picoGreen, Hoechst and TMRM and imaged on an ImageXpress Micro XLS widefield high content screening system, as described in Methods. Hoechst normalised picoGreen signal, for amplification of the mitochondrial picoGreen signal, was overlaid and segmented for quantitative analysis on the TMRM signal, denoting thus nucleoid (green) and mitochondrial (red) segments. Below the images, quantifications in cell lines grouped by their switch state (MB1_UF / MB1_LF) of the picoGreen integrated intensity, readout for mtDNA content (left panel); of mitochondrial number, total volume and individual mitochondrial volume, to assess the state of the mitochondrial network (three panels to the right, respectively). Unpaired Student's t-tests.

(ns: $P > 0.05$, *: $P \leq 0.05$, **: $P \leq 0.01$,***: $P \leq 0.001$, ****: $P \leq 0.0001$)

In order to determine the differences in mitochondrial content and activity, we measured oxidative mitochondrial function and used high resolution, high content imaging on large cell populations to quantify overall mtDNA content and morphology in the MB1_UF and MB1_LF fork cell lines (Figure 3C-F). MB1_UF cells had higher oxygen consumption rate and mitochondrial membrane potential, indicating higher redox potential to feed electrons in the respiratory chain (Figure 3C, D). Higher mitochondrial abundance in MB1_UF cells was confirmed by western blot analysis of respiratory chain components (Figure 3E). High resolution image analysis of mitochondria across cell populations (Figure 3F) showed that while the number of mitochondria per cell were equal in the two forks, MB1_UF cells had higher mtDNA content, total and individual mitochondrial area, i.e. larger mitochondrial size and overall content. These experiments confirmed a switch in mitochondrial abundance and function as predicted by gene expression profiling using MCbiclust.

Next, in order to detect the metabolic switch with altered substrate preferences associated with the different mitochondrial switch positions, we used steady state metabolic flux analysis of heavy carbon labelled key cellular fuels, all uniformly labelled ^{13}C-glucose, ^{13}C-glutamine and ^{13}C-pyruvate. MB1_UF and MB1_LF cells were incubated with each substrate in separate experimental sets. Labelled carbons were traced by mass spectrometry analysis to quantify isotopologues of metabolites in main catabolic pathways of the three substrates. In addition, to verify substrate dependencies, we introduced conditions with combinations of reduced glutamine and pyruvate (glutamine[glutamine] H/L = 1mM/0.1mM; [pyruvate] +/- = 1mM/0 mM, Figure 4A).

Using these methods, first the fate of glucose via aerobic glycolysis was assessed by quantifying glucose uptake, lactate production, and relative incorporation of glucose derived heavy carbons into lactate (m+3 isotopologue). We found no significant differences between the MB1_UF and MB1_LF switches in all the conditions tested (Figure 4B), showing that aerobic glycolysis occurs in both MB1 switches at high rate, verified by the nearly complete glucose-derived ^{13}C labelling of glycolytic intermediates (Figure S5A). However, MB1_UF cells produced pyruvate (m+3) from glucose at significantly higher levels than MB1_LF (Figure 4C), suggesting that glucose flux through glycolysis was higher in the MB1_UF switch, presumably delivering pyruvate to be oxidised in the TCA cycle.

Figure 4

Figure 4. Metabolic wiring of the switch positions probed by 13C-labelled substrates

A. Nutrient supply and [13]C-labelling conditions used throughout the experiments presented in this figure. High and low glutamine (H/L = 1mM/0.1mM) levels were combined with the presence and absence of pyruvate (+/- = 1mM/0 mM), glucose was kept at 10mM.

B. Aerobic glycolysis was tested by measuring glucose uptake (first panel) and lactate secretion (second panel) from the mass spectrometry data obtained from cell culture media, as well as the fractional enrichment of m+3 labelled lactate intracellularly (third panel), see Methods. Cells were grouped by switch state (MB1_UF / MB1_LF). 2-way ANOVA with Sidak's multiple comparisons test.

C. Fractional enrichment of the m+3 pyruvate isotopomer, following [U13]C-glucose labelling for 24h, in cells grouped by switch state (MB1_UF / MB1_LF). 2-way ANOVA with Sidak's multiple comparison tests.

D. Overall model of the MB1_UF/MB1_LF metabolic switch in central carbon metabolism. Glucose derived pyruvate is preferentially used in the TCA cycle either by the PDH or PC route in the MB1_UF cells/position, driving high respiration. Glutamine is taken up more in MB1_LF cells, and they use it preferentially in the TCA cycle, and their respiration relies on glutamine derived glutamate, but glutamate is also re-secreted into the extracellular space in high amount in these cells. Extracellular pyruvate does not contribute significantly to the TCA cycle, preferentially used for lactate production and thus setting the cytosolic redox ratio.

E. Relative carbon contribution to citrate following 24h labelling with [U13]C-glucose and [U13]C-glutamine, in cells grouped by switch state (MB1_UF / MB1_LF) The values were calculated as described in Methods. 2-way ANOVA with Sidak's multiple comparison tests.

F. The fate of glucose derived carbons in the TCA cycle. Fractional enrichment of the m+2 (PDH route, left panel) and m+5 (PDH+PC route, right panel) isotopologues in citrate following 24h labelling with [U13]C-glucose, in cells grouped by switch state (MB1_UF / MB1_LF). 2-way ANOVA with uncorrected Fisher's LSD multiple comparison tests.

G. The fate of glutamine in cells grouped by switch state (MB1_UF / MB1_LF). Glutamine uptake and glutamate secretion rates were measured by specific biochemical assays in the culture media prior and following 24h incubation with cells (left two panels). Fractional enrichment of malate m+4 (oxidative route) and citrate m+5 (reductive carboxylation) isotopologues following 24h labelling with [U13]C- glutamine (right two panels). 2-way ANOVA with uncorrected Fisher's LSD multiple comparison tests.

H. Substrate dependence of mitochondrial respiration and cellular ATP production in cells grouped by switch state (MB1_UF / MB1_LF). Left panel shows basal oxygen consumption rate as measured by Seahorse XFe96 analyser (see Methods). Right panel shows total cellular ATP content measured by a luciferase assay following lysis of cells kept in the indicated media for 24h. 2-way ANOVA with uncorrected Fisher's LSD multiple comparison tests.

(ns: $P > 0.05$, *: $P \leq 0.05$, **: $P \leq 0.01$,***: $P \leq 0.001$, ****: $P \leq 0.0001$)

The TCA cycle can be fed either by glucose derived pyruvate or glutamine derived glutamate at any ratio (Figure 4D). Thus, to assess substrate preferences, we quantified the relative carbon contribution of glucose, glutamine and externally added pyruvate carbons to the TCA cycle. The results showed significantly higher glucose-derived pyruvate incorporation into citrate, fumarate and malate in the MB1_UF cells, while more glutamine derived carbons were incorporated into TCA cycle metabolites in MB1_LF cells (Figures 4E, S5B). These results indicated a clear switch between glucose-derived pyruvate versus glutamine feeding into the TCA cycle. Pyruvate entered the TCA cycle both via the PDH and PC routes in MB1_UF cells (Figure 4F). On the other hand, in MB1_LF cells, glutamine was used in both the oxidative and reductive direction at higher levels than in MB1_UF cells, together with increased glutamine uptake and subsequent glutamate secretion (Figure 4G). In summary, whilst in cultured tumour cells both low (MB1_LF) and high (MB1_UF) mitochondrial activity switches are accompanied by aerobic glycolysis, the switches differ in their substrate preference to fuel their TCA cycle activity. The MB1_UF switch promotes the use of glucose as main carbon source, while in MB1_LF glutaminolysis dominates. To functionally verify the switch to glutamine preference in the MB1_LF phenotype, we applied glutamine restriction (see conditions in Figure 4A). The lack of glutamine resulted in reduced bioenergetic competence only in MB1_LF cells, such as reduced mitochondrial oxygen consumption rate and total cellular ATP levels (Figure 4H). Strikingly, glutamine restriction partly reverted the MB1_LF metabolic switch in the MB_UF direction by promoting glucose derived pyruvate entry in the TCA cycle via the PC route (see Figure 4F). Interestingly, externally added pyruvate was incorporated only at low levels into the TCA cycle (Figure S5D), but promoted glucose derived pyruvate utilisation by the PC route in MB1_LF cells, shifting the metabolic phenotype towards the MB1_UF switch (Figure 4F).

Thus we discovered a metabolic switch defining MB1_UF cells, (and presumably in tumours), where a mature luminal (mL) cell-of-origin phenotype, with oestrogen receptor and cell cycle signalling, is associated with an adaptive gene expression pattern, promoting high glycolysis together with high TCA activity and energy production, relying mostly on glucose. In the other position of the switch, in MB1_LF mammary stem cell like cells (MASCs), glutamine is preferred to use to fuel lower mitochondrial activity. In addition, our functional analysis also revealed that the functional metabolic switch is adaptable by environmental changes, the positions can be shifted by altering substrate availability. A switch from glutamine to pyruvate supply in MB1_LF cells not only reverted the low pyruvate entry into the mitochondria and corresponding TCA cycle flux to levels found in MB1_UF cells (Figure 4E, F), but also increased respiration and ATP levels (Figure 4H).

Our prediction assumed that transcriptomic profile switches can determine metabolic routes via modulating the protein expression levels of metabolic pathway components. In order to test this assumption, we performed western blot analysis of enzymes from key pathways by which glucose, glutamine and pyruvate are fed into the TCA cycle (Figures 5A, S6). Expression levels of 29 metabolic enzymes with diverse localisation clustered according to switch positions (UF and LF) across the cell lines (Figure 5B). Higher mitochondrial abundance in the UF was confirmed by increased grp75 expression compared to the cytosolic β-actin marker (Figure 5B, lower panel), and interestingly, changes compared to mitochondrial and cytosolic abundance showed divergent patterns of differential expression (Figure 5B left and right heatmaps, respectively). Moreover, scaled protein levels of key enzymes matched the CV value of their transcripts (Figure 5C). Overall, expression of most metabolic enzymes followed the higher mitochondrial abundance in the MB1_UF state, the ones with the highest difference involved in the malate-aspartate shuttle (GOT1, GOT2) and

pyruvate utilisation (MPC2, pPDH ratio, IDH2)(Figure 5D, E). These results supported the metabolic phenotype described for MB1_UF cells (see Figure 4). On the other hand, a small set of enzymes was overexpressed in the LF, underpinning key characteristics of the LF, e.g. increased glutamine utilisation (GLS1, GFPT2). These data confirmed that transcriptomic patterns can directly predict metabolic pathway activities. Close correlation of transcript and protein levels indicated that transcriptional regulation dominates metabolic pathway activities via the regulation of their key enzymes.

Figure 5

Figure 5. Protein expression profiles match transcriptomic switches

A. Western blot analysis of 29 metabolic enzymes across five cell lines representing MB1_UF and MB1_LF switch positions and cell states. Representative blots from n ≧ 3 independent experiments, used for quantification of signals, normalising density values to either grp75 or β-actin.

B, D, E. Clustering and quantitative analysis of enzyme expression levels obtained from **A.** Values were separately normalised to either mitochondrial (left panels) or cytosolic (right panels) markers, grp75 or β-actin. The comparison of mitochondrial abundance based on the grp75/β-actin ratio is shown in the lower panel in **B.** Heatmaps are based on a pairwise, robust to outliers distance measure of the mean protein expression values. K-means clustering. **D, E.** 2-way ANOVA with uncorrected Fisher's LSD multiple comparison tests.

C. Heatmap and k-means clustering of (i) transcript CV values of the 29 enzymes from the METABRIC dataset (see Figs. 1, 2, positive values predicting higher expression of genes in MB1_UF, negative values corresponding to higher expression in MB1_LF), (ii-iii) the scaled (between -1 and 1) protein expression levels averaged across all cell lines, normalised to grp75 (ii) or β-actin (iii).

(ns: P > 0.05, *:P ≤ 0.05, **: P ≤ 0.01,***: P ≤ 0.001, ****:   P ≤ 0.0001)

4. Pathological and clinical predictions from metabolic gene switches

As metabolic gene switches are associated with cell states, they potentially indicate other, clinically relevant features, genotypes, phenotypes or gene transcription patterns. Thus, we tested the correlation of switch positions with TCGA and METABRIC associated genetic, biological, pathological and clinical data. First, in order to understand whether a specific tumour architecture or histological origin is associated with the switch positions, we used the most recent dataset with comprehensive coverage of histological features and classifications in the full TCGA BRCA sample set[46], and tested the distribution of features among individual forks (Figures 6A, S7, Table S8). A group of features was highly variable between switch states, including enrichment of invasive lobular carcinoma (ILC) in the MB12_LF switch state, or increased signs of nuclear pleomorphism in MB2_UF (Figure 6B). Overall, both MB1_UF and MB2_UF switch positions were associated with increased proliferation (mitosis), higher cellularity (epi_area) as compared to MB12_LF, of which about half of tumours were classified as ILC. In addition, MB2_UF was highly enriched in samples with necrosis, nuclear pleomorphism and immune infiltration (inflam), indicating higher malignity. Next, in order to identify and compare further unique features for each switch position, we have assessed the fraction of samples for each individual feature including both histological and genetic data. As shown in Figure 6C, when analysing the MB1 and MB2 bicluster switches, the MB12_LF, MB1_UF and MB2_UF samples were clustered separately. MB12_LF samples characteristically showed low proliferation, were enriched in LumA intrinsic and ILC histological subtypes, likely determined by specific CN variation, methylation and miRNA clusters (Figure 6D). On the other hand, while both MB1_UF and MB2_UF samples showed increased proliferation, they diverged in their molecular subtype (LumB and Basal, respectively) and the underlying genetic and transcriptional constitution (CN variation, methylation and miRNA clusters).

Finally, we asked whether metabolic switch positions in the MB1 bicluster can predict the clinical outcome of ER positive breast cancer. For this aim, we performed survival analysis of an independent breast cancer dataset[47], using the weighted average of a core geneset of 44 genes determining the MB1_UF and MB1_LF switch positions. The MB1 switch includes almost exclusively ER positive samples, dividing them into two groups, which partly overlap with intrinsic subtypes LumA, LumB and normal-like (see Figure S4A). ER positive tumours overall are sensitive to antihormonal therapy, but adjuvant chemotherapy is usually recommended against tumours progressing to lymph nodes. However, only a fraction of these patients respond to this distressing treatment, motivating large recent efforts to develop gene expression pattern based methods to stratify individual ER positive tumours into chemotherapy sensitive and resistant groups prior to adjuvant treatment[48,49]. The overall survival of patients with MB1_LF pattern expressing tumours was significantly higher as compared to the MB1_UF group (Figure 6E). Moreover, the MB1_LF pattern predicted improved survival in a subgroup of patients with ER positive, Her2 negative and lymph node positive tumours following endocrine and chemotherapy treatment (Figure 6F), indicating higher sensitivity to chemotherapy in this group. The impact of the MB1_LF pattern was independent of previous GEP based methods (e.g. Pam50, see discussion), as its presence in LumB samples was associated with a highly significant increase in survival (Figure 6G).

Figure 6

Figure 6. Pathological and clinical predictions from metabolic gene switches

A. Chi-squared distribution of discrete histological features across the upper, lower and non-fork samples of the three biclusters (MB1-MB3). Heatmap shows k-means clustering of chi-squared test p-values (-log10P), dividing the histological parameters into three groups: high, low and none association with the bicluster.

B. Visualisation of histological feature distribution on the MB1 and MB2 switches. Classifiers of histological tumour type (upper panel) and degree of nuclear pleomorphism are overlaid onto 2D distribution plots of TCGA samples. The axes represent the scale between the upper (UF) and lower (LF) forks of each bicluster.

C. Clustering of histological and genetic features of samples belonging to the MB1 and MB2 switches. Fractions of samples for each feature were calculated and clustered with a custom robust to outliers distance measure and K-means clustering. Genetic features representative of each switch position are highlighted in bold.

D. Key genetic and transcriptional architecture of the MB1 and MB2 switches.

E-G. Survival analysis of samples in the MB1 switch. Kaplan-Meier survival plots of samples with MB1_UF and MB1_LF patterns are shown. **E.** All (1660) samples compiled from the GEO database [47]. **F.** Subset of samples from endocrine and chemotherapy treated ER positive patients (100). **G.** Subset of samples with LumB molecular subtype (51). Menkel Cox logrank test results are shown on the individual plots.

Altogether, these results indicated that large scale mutually exclusive gene expression clusters govern the mitochondrial and metabolic status of breast tumours. These multi-state switches extend beyond the metabolic gene sets, and reveal the biochemistry, architecture and chemosensitivity of tumours.

Discussion

The current availability of large scale genomic and transcriptomic datasets on physiological and pathological conditions escalated the need to devise computational strategies providing statistically sound predictions of how individual genes and gene groups are regulated and contribute to cellular phenotype. The gene expression patterns underlying phenotype or cell state switches[50–52] are complex and have not been comprehensively analysed, although methods to address the question are being developed[53]. These methods are part of pattern recognition, subgroup discovery and dimensionality reduction techniques, and often apply biclustering approaches for enhancing the probability to discover existing patterns of gene expression existing only in a subset of samples. We have recently developed a biclustering method (MCbiclust[28,29]), optimised for discovering large scale anti-correlating patterns to stratify large heterogeneous sample collections. The strategy thoroughly distinguished cancer cell types according to their cell (or organ) of origin, and discovered subgroups of SDH deficient tumour samples partly by their altered metabolic networks[28]. Here we developed a resource where results obtained from massive correlated biclustering are further enriched by advanced pathway analysis as a pipeline to understand metabolic adaptation in >3K heterogeneous breast cancer samples. The method identified large multi-state gene expression pattern switches in signalling and metabolic adaptation, coinciding with differentiation status, i.e the cell of origin of tumour cell types.

Currently available biclustering methods have been productive in finding functionally relevant gene networks, but their applicability also had several limitations[54]. Our previous evaluation of MCbiclust[28] have shown that the algorithm scored better than five mainstream biclustering methods available at the time of the launch of our project, particularly due to its ability to recognise large biclusters. Cell fate is thought to be more often regulated by larger gene expression patterns than activation of a small set of single genes[55–57], which probably explains the readiness of our method to discover patterns in subset of tumours with different cell of origin. In addition, while previous methods mostly used differential gene expression compared to normal samples or pairwise differential correlation[16], in MCbiclust the selection of the bicluster samples are based on a greedy search maximising the average *absolute correlation value* of a large (~1K) geneset, thus covering a large fraction of the whole transcriptome, and allowing the discovery of composite checker pattern biclusters, containing both positively and negatively correlated gene groups, defining two gene sets switching expression in two subsets of samples. Finally, by discovering multiple biclusters, some subsets (MB12_LF) showed near complete overlap, detecting alternative, multi-state switches.

Breast cancer has been intensely surveyed for classification, leading to definition of intrinsic subtypes based on gene expression profiles (Pam50)[58], and integrated clusters using also genomic data[31,59], but further integration with histological data[46] and the exact origins of other emerging subtypes are still debated[33,60]. The stratification of breast cancer provided by MCbiclust only partially overlaps with all previous classifications (see Fig. S4), thus it provides a novel perspective on breast cancer subgroups, in particular from functional aspects, such as metabolic rewiring. Importantly, other recent biclustering approaches to breast cancer datasets do not provide better resolution compared to Pam50 subtyping[61,62]. On the other hand, MCbiclust, apart from demonstrating capability of predicting overall patient survival superior to Pam50 based methods (see Fig. 6), also provided two novel and interesting insights into tumour origin and architecture. First, by comparing previously

described patterns associated with the putative cell of origin of different tumour types, a close correlation with the biclusters was found (see Figs. 2D,E; S4B), illustrating the power of the method to determine cell fate associated gene expression patterns. The lowest correlation was found between MaSC and MB12_LF, which might not be surprising given that MaSC represents an already heterogeneous population of cells[40], of which certain fractions might belong to other switches (in particular MB2_UF). Second, histological analysis of samples belonging to MB12_LF switch showed striking enrichment in ILC, a lobular tumour type[63–66], known for low proliferation, but increased probability of metastasis formation. The signalling pathways associated with the MB12_LF switch might help to explain this elusive property of ILC tumours.

Intriguingly, all biclusters (including the ones arising from random genesets as starting points - see Figs. 1, S1) were significantly enriched in nuclear encoded mitochondrial genes. This may reflect co-regulation of a large set of genes belonging to the same organelle, and indicate the necessity of shaping mitochondrial gene expression patterns according to the actual cell state. Moreover, gene expression and metabolomic analysis have also disclosed strong co-regulation of mitochondrial and extra-mitochondrial metabolic genes, providing a platform for understanding metabolic and bioenergetic adaptation in cancer development. Interestingly, both the MB1_UF and MB2_UF switch positions showed increased transcriptional biogenesis of mitochondria as compared to MB12_LF, where nuclear encoded mitochondrial genes were under-represented. This observation is in line with the currently emerging view of mitochondria in tumours, recognising that normal or high activity of certin mitochondrial metabolic pathways is often required for enhanced tumour proliferation[67]. However, our analysis, focusing only on tumour samples, revelas only relative differences in mitochondrial biogenesis between the switch positions, thus cannot discriminate between upregulation of biogenesis in the MB1_UF and MB2_UF switch positions and suppression of mitochondrial biogenesis and function in the opposite (MB12_LF) position. Recent data indicating loss of mitochondrial mass and function under chronic stress conditions[68] and reduced mitochondrial activity in stem cell states[69] support the latter view.

Overall, we created a method with the potential of discovering large, alternating gene expression patterns, providing a useful resource for exploring physiological and pathological[70] conditions where phenotypic switches play a major role.

References

1. Faubert, B., Solmonson, A., and DeBerardinis, R.J. (2020). Metabolic reprogramming and cancer progression. Science *368*. 10.1126/science.aaw5473.

2. Kreuzaler, P., Panina, Y., Segal, J., and Yuneva, M. (2020). Adapt and conquer: Metabolic flexibility in cancer growth, invasion and evasion. Mol Metab *33*, 83–101. 10.1016/j.molmet.2019.08.021.

3. Fendt, S.-M., Frezza, C., and Erez, A. (2020). Targeting Metabolic Plasticity and Flexibility Dynamics for Cancer Therapy. Cancer Discov. *10*, 1797–1807. 10.1158/2159-8290.CD-20-0844.

4. Intlekofer, A.M., and Finley, L.W.S. (2019). Metabolic signatures of cancer cells and stem cells. Nat Metab *1*, 177–188. 10.1038/s42255-019-0032-0.

5. Boumahdi, S., and de Sauvage, F.J. (2020). The great escape: tumour cell plasticity in resistance to targeted therapy. Nat. Rev. Drug Discov. *19*, 39–56. 10.1038/s41573-019-0044-1.

6. Baryawno, N., Przybylski, D., Kowalczyk, M.S., Kfoury, Y., Severe, N., Gustafsson, K., Kokkaliaris, K.D., Mercier, F., Tabaka, M., Hofree, M., et al. (2019). A Cellular Taxonomy of the Bone Marrow Stroma in Homeostasis and Leukemia. Cell *177*, 1915–1932.e16. 10.1016/j.cell.2019.04.040.

7. García-Jiménez, C., and Goding, C.R. (2019). Starvation and Pseudo-Starvation as Drivers of Cancer Metastasis through Translation Reprogramming. Cell Metab. *29*, 254–267. 10.1016/j.cmet.2018.11.018.

8. Dobson, S.M., García-Prat, L., Vanner, R.J., Wintersinger, J., Waanders, E., Gu, Z., McLeod, J., Gan, O.I., Grandal, I., Payne-Turner, D., et al. (2020). Relapse-Fated Latent Diagnosis Subclones in Acute B Lineage Leukemia Are Drug Tolerant and Possess Distinct Metabolic Programs. Cancer Discov. *10*, 568–587. 10.1158/2159-8290.CD-19-1059.

9. Samanta, D., and Semenza, G.L. (2018). Metabolic adaptation of cancer and immune cells mediated by hypoxia-inducible factors. Biochim. Biophys. Acta Rev. Cancer *1870*, 15–22. 10.1016/j.bbcan.2018.07.002.

10. Jeon, S.-M., Chandel, N.S., and Hay, N. (2012). AMPK regulates NADPH homeostasis to promote tumour cell survival during energy stress. Nature *485*, 661–665. 10.1038/nature11066.

11. Saxton, R.A., and Sabatini, D.M. (2017). mTOR Signaling in Growth, Metabolism, and Disease. Cell *169*, 361–371. 10.1016/j.cell.2017.03.035.

12. Fiore, A.P.Z.P., Ribeiro, P. de F., and Bruni-Cardoso, A. (2018). Sleeping Beauty and the Microenvironment Enchantment: Microenvironmental Regulation of the Proliferation-Quiescence Decision in Normal Tissues and in Cancer Development. Front Cell Dev Biol *6*, 59. 10.3389/fcell.2018.00059.

13. Ye, X., and Weinberg, R.A. (2015). Epithelial-Mesenchymal Plasticity: A Central Regulator of Cancer Progression. Trends Cell Biol. *25*, 675–686. 10.1016/j.tcb.2015.07.012.

14. Li, X., Hui, S., Mirek, E.T., Jonsson, W.O., and Anthony, T.G. (2022). Circulating metabolite homeostasis achieved through mass action. Nature.

15. Rosario, S.R., Long, M.D., Affronti, H.C., Rowsam, A.M., Eng, K.H., and Smiraglia, D.J. (2018). Pan-cancer analysis of transcriptional metabolic dysregulation using The Cancer Genome Atlas. Nat. Commun. *9*, 5330. 10.1038/s41467-018-07232-8.

16. Reznik, E., and Sander, C. (2015). Extensive decoupling of metabolic genes in cancer. PLoS Comput. Biol. *11*, e1004176. 10.1371/journal.pcbi.1004176.

17. Peng, X., Chen, Z., Farshidfar, F., Xu, X., Lorenzi, P.L., Wang, Y., Cheng, F., Tan, L., Mojumdar, K., Du, D., et al. (2018). Molecular Characterization and Clinical Relevance of Metabolic Expression Subtypes in Human Cancers. Cell Rep. *23*, 255–269.e4. 10.1016/j.celrep.2018.03.077.

18. Bradner, J.E., Hnisz, D., and Young, R.A. (2017). Transcriptional Addiction in Cancer. Cell *168*, 629–643. 10.1016/j.cell.2016.12.013.

19. Levine, A.J., and Puzio-Kuter, A.M. (2010). The control of the metabolic switch in cancers by oncogenes and tumor suppressor genes. Science *330*, 1340–1344. 10.1126/science.1193494.

20. Kroemer, G., and Pouyssegur, J. (2008). Tumor cell metabolism: cancer's Achilles' heel. Cancer Cell *13*, 472–482. 10.1016/j.ccr.2008.05.005.

21. Haider, S., McIntyre, A., van Stiphout, R.G.P.M., Winchester, L.M., Wigfield, S., Harris, A.L., and Buffa, F.M. (2016). Genomic alterations underlie a pan-cancer metabolic shift associated with tumour hypoxia. Genome Biol. *17*, 140. 10.1186/s13059-016-0999-8.

22. Reddy, J., Fonseca, M.A.S., Corona, R.I., Nameki, R., Segato Dezem, F., Klein, I.A., Chang, H., Chaves-Moreira, D., Afeyan, L.K., Malta, T.M., et al. (2021). Predicting master transcription factors from pan-cancer expression data. Sci Adv *7*, eabf6123. 10.1126/sciadv.abf6123.

23. Hoadley, K.A., Yau, C., Hinoue, T., Wolf, D.M., Lazar, A.J., Drill, E., Shen, R., Taylor, A.M., Cherniack, A.D., Thorsson, V., et al. (2018). Cell-of-Origin Patterns Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. Cell *173*, 291–304.e6. 10.1016/j.cell.2018.03.022.

24. Li, J., Eu, J.Q., Kong, L.R., Wang, L., Lim, Y.C., Goh, B.C., and Wong, A.L.A. (2020). Targeting Metabolism in Cancer Cells and the Tumour Microenvironment for Cancer Therapy. Molecules *25*. 10.3390/molecules25204831.

25. Nilsson, R., Jain, M., Madhusudhan, N., Sheppard, N.G., Strittmatter, L., Kampf, C., Huang, J., Asplund, A., and Mootha, V.K. (2014). Metabolic enzyme expression highlights a key role for MTHFD2 and the mitochondrial folate pathway in cancer. Nat. Commun. *5*, 3128. 10.1038/ncomms4128.

26. Gaude, E., and Frezza, C. (2016). Tissue-specific and convergent metabolic transformation of cancer correlates with metastatic potential and patient survival. Nat. Commun. *7*, 13041. 10.1038/ncomms13041.

27. Anesiadis, N., Kobayashi, H., Cluett, W.R., and Mahadevan, R. (2013). Analysis and design of a genetic circuit for dynamic metabolic engineering. ACS Synth. Biol. *2*, 442–452. 10.1021/sb300129j.

28. Bentham, R.B., Bryson, K., and Szabadkai, G. (2017). MCbiclust: a novel algorithm to discover large-scale functionally related gene sets from massive transcriptomics data collections. Nucleic Acids Res. *45*, 8712–8730. 10.1093/nar/gkx590.

29. Bentham, R.B., Bryson, K., and Szabadkai, G. (2019). Biclustering Analysis of Co-regulation Patterns in Nuclear-Encoded Mitochondrial Genes and Metabolic Pathways. Methods Mol. Biol. *1928*, 469–478. 10.1007/978-1-4939-9027-6_24.

30. Rath, S., Sharma, R., Gupta, R., Ast, T., Chan, C., Durham, T.J., Goodman, R.P., Grabarek, Z., Haas, M.E., Hung, W.H.W., et al. (2021). MitoCarta3. 0: an updated mitochondrial proteome now with sub-organelle localization and pathway annotations. Nucleic Acids Res. *49*, D1541–D1547.

31. Curtis, C., Shah, S.P., Chin, S.-F., Turashvili, G., Rueda, O.M., Dunning, M.J., Speed, D., Lynch, A.G., Samarajiwa, S., Yuan, Y., et al. (2012). The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. Nature *486*, 346–352. 10.1038/nature10983.

32. Cancer Genome Atlas Network (2012). Comprehensive molecular portraits of human breast tumours. Nature *490*, 61–70. 10.1038/nature11412.

33. Aure, M.R., Vitelli, V., Jernström, S., Kumar, S., Krohn, M., Due, E.U., Haukaas, T.H., Leivonen, S.-K., Vollan, H.K.M., Lüders, T., et al. (2017). Integrative clustering reveals a novel split in the luminal A subtype of breast cancer with impact on outcome. Breast Cancer Res. *19*, 44. 10.1186/s13058-017-0812-y.

34. Yuan, C.C., Calvo, S.E., and Mootha, V.K. (2021). MitoCarta3. 0: an updated mitochondrial proteome now with sub-organelle localization and pathway annotations. Nucleic acids.

35. Berger, A.C., Korkut, A., Kanchi, R.S., Hegde, A.M., Lenoir, W., Liu, W., Liu, Y., Fan, H., Shen, H., Ravikumar, V., et al. (2018). A Comprehensive Pan-Cancer Molecular Study of Gynecologic and Breast Cancers. Cancer Cell *33*, 690–705.e9. 10.1016/j.ccell.2018.03.014.

36. Morel, A.-P., Ginestier, C., Pommier, R.M., Cabaud, O., Ruiz, E., Wicinski, J., Devouassoux-Shisheboran, M., Combaret, V., Finetti, P., Chassot, C., et al. (2017). A stemness-related ZEB1–MSRB3 axis governs cellular pliancy and breast cancer genome stability. Nat. Med. *23*, 568–578. 10.1038/nm.4323.

37. Vaske, C.J., Benz, S.C., Sanborn, J.Z., Earl, D., Szeto, C., Zhu, J., Haussler, D., and Stuart, J.M. (2010). Inference of patient-specific pathway activities from multi-dimensional cancer genomics data using PARADIGM. Bioinformatics *26*, i237–i245. 10.1093/bioinformatics/btq182.

38. Cliff, T.S., and Dalton, S. (2017). Metabolic switching and cell fate decisions: implications for pluripotency, reprogramming and development. Curr. Opin. Genet. Dev. *46*, 44–49. 10.1016/j.gde.2017.06.008.

39. Ly, C.H., Lynch, G.S., and Ryall, J.G. (2020). A Metabolic Roadmap for Somatic Stem Cell Fate. Cell Metab. *31*, 1052–1067. 10.1016/j.cmet.2020.04.022.

40. Fu, N.Y., Nolan, E., Lindeman, G.J., and Visvader, J.E. (2020). Stem Cells and the Differentiation Hierarchy in Mammary Gland Development. Physiol. Rev. *100*, 489–523. 10.1152/physrev.00040.2018.

41. Hänzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis for microarray and RNA-seq data. BMC Bioinformatics *14*, 7. 10.1186/1471-2105-14-7.

42. Neve, R.M., Chin, K., Fridlyand, J., Yeh, J., Baehner, F.L., Fevr, T., Clark, L., Bayani, N., Coppe, J.-P., Tong, F., et al. (2006). A collection of breast cancer cell lines for the study of functionally distinct cancer subtypes. Cancer Cell *10*, 515–527. 10.1016/j.ccr.2006.10.008.

43. Charafe-Jauffret, E., Ginestier, C., Monville, F., Finetti, P., Adélaïde, J., Cervera, N., Fekairi, S., Xerri, L., Jacquemier, J., Birnbaum, D., et al. (2006). Gene expression profiling of breast cell lines identifies potential new basal markers. Oncogene *25*, 2273–2284. 10.1038/sj.onc.1209254.

44. Ghandi, M., Huang, F.W., Jané-Valbuena, J., Kryukov, G.V., Lo, C.C., McDonald, E.R., 3rd, Barretina, J., Gelfand, E.T., Bielski, C.M., Li, H., et al. (2019). Next-generation characterization of the Cancer Cell Line Encyclopedia. Nature *569*, 503–508. 10.1038/s41586-019-1186-3.

45. Bruna, A., Rueda, O.M., Greenwood, W., Batra, A.S., Callari, M., Batra, R.N., Pogrebniak, K., Sandoval, J., Cassidy, J.W., Tufegdzic-Vidakovic, A., et al. (2016). A Biobank of Breast Cancer Explants with Preserved Intra-tumor Heterogeneity to Screen Anticancer Compounds. Cell *167*, 260–274.e22. 10.1016/j.cell.2016.08.041.

46. Heng, Y.J., Lester, S.C., Tse, G.M., Factor, R.E., Allison, K.H., Collins, L.C., Chen, Y.-Y., Jensen, K.C., Johnson, N.B., Jeong, J.C., et al. (2017). The molecular basis of breast cancer pathological phenotypes. J. Pathol. *241*, 375–391. 10.1002/path.4847.

47. Györffy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q., and Szallasi, Z. (2010). An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. Breast Cancer Res. Treat. *123*, 725–731. 10.1007/s10549-009-0674-9.

48. Stein, R.C., Dunn, J.A., Bartlett, J.M.S., Campbell, A.F., Marshall, A., Hall, P., Rooshenas, L., Morgan, A., Poole, C., Pinder, S.E., et al. (2016). OPTIMA prelim: a randomised feasibility study of personalised care in the treatment of women with early breast cancer. Health Technol. Assess. *20*, xxiii – xxix, 1–201. 10.3310/hta20100.

49. Taylor, C., Meisel, J., and Kalinsky, K. (2022). Are we closer to being able to select patients with node-positive hormone receptor-positive breast cancer who can safely omit chemotherapy? Ther. Adv. Med. Oncol. *14*, 17588359221084769. 10.1177/17588359221084769.

50. Xia, K., Xue, H., Dong, D., Zhu, S., Wang, J., Zhang, Q., Hou, L., Chen, H., Tao, R., Huang, Z., et al. (2006). Identification of the proliferation/differentiation switch in the cellular network of multicellular organisms. PLoS Comput. Biol. *2*, e145. 10.1371/journal.pcbi.0020145.

51. Miyamoto, T., Furusawa, C., and Kaneko, K. (2015). Pluripotency, Differentiation, and Reprogramming: A Gene Expression Dynamics Model with Epigenetic Feedback Regulation. PLoS Comput. Biol. *11*, e1004476. 10.1371/journal.pcbi.1004476.

52. Duddu, A.S., Majumdar, S.S., Sahoo, S., Jhunjhunwala, S., and Jolly, M.K. (2022). Emergent dynamics of a three-node regulatory network explain phenotypic switching and heterogeneity: a case study of Th1/Th2/Th17 cell differentiation. Mol. Biol. Cell *33*, ar46. 10.1091/mbc.E21-10-0521.

53. Noronha, M.D.M., Henriques, R., Madeira, S.C., and Zárate, L.E. (2022). Impact of metrics on biclustering solution and quality: A review. Pattern Recognit. *127*, 108612. 10.1016/j.patcog.2022.108612.

54. Joachimiak, M.P., Tuglus, C., Salamzade, R., van der Laan, M., and Arkin, A.P. (2022). Deep surveys of transcriptional modules with Massive Associative K-biclustering (MAK). bioRxiv, 2022.08.26.505372. 10.1101/2022.08.26.505372.

55. Davidson, E.H. (2010). Emerging properties of animal gene regulatory networks. Nature *468*, 911–920. 10.1038/nature09645.

56. Naciri, I., Laisné, M., Ferry, L., Bourmaud, M., Gupta, N., Di Carlo, S., Huna, A., Martin, N., Peduto, L., Bernard, D., et al. (2019). Genetic screens reveal mechanisms for the transcriptional regulation of tissue-specific genes in normal cells and tumors. Nucleic Acids Res. *47*, 3407–3421. 10.1093/nar/gkz080.

57. Stadhouders, R., Filion, G.J., and Graf, T. (2019). Transcription factors and 3D genome conformation in cell-fate decisions. Nature *569*, 345–354. 10.1038/s41586-019-1182-7.

58. Perou, C.M., Sørlie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., Rees, C.A., Pollack, J.R., Ross, D.T., Johnsen, H., Akslen, L.A., et al. (2000). Molecular portraits of human breast tumours. Nature *406*, 747–752. 10.1038/35021093.

59. Russnes, H.G., Lingjærde, O.C., Børresen-Dale, A.-L., and Caldas, C. (2017). Breast Cancer Molecular Stratification: From Intrinsic Subtypes to Integrative Clusters. Am. J. Pathol. *187*, 2152–2162. 10.1016/j.ajpath.2017.04.022.

60. Pommier, R.M., Sanlaville, A., Tonon, L., Kielbassa, J., Thomas, E., Ferrari, A., Sertier, A.-S., Hollande, F., Martinez, P., Tissier, A., et al. (2020). Comprehensive characterization of claudin-low breast tumors reflects the impact of the cell-of-origin on cancer evolution. Nat. Commun. *11*, 3431. 10.1038/s41467-020-17249-7.

61. Rose, T.D., Bechtler, T., Ciora, O.-A., Anh Lilian Le, K., Molnar, F., Köhler, N., Baumbach, J., Röttger, R., and Pauling, J.K. (2022). MoSBi: Automated signature mining for molecular stratification and subtyping. Proc. Natl. Acad. Sci. U. S. A. *119*, e2118210119. 10.1073/pnas.2118210119.

62. Wang, Y.K., Print, C.G., and Crampin, E.J. (2013). Biclustering reveals breast cancer tumour subgroups with common clinical features and improves prediction of disease recurrence. BMC Genomics *14*, 102. 10.1186/1471-2164-14-102.

63. Weigelt, B., Geyer, F.C., Natrajan, R., Lopez-Garcia, M.A., Ahmad, A.S., Savage, K., Kreike, B., and Reis-Filho, J.S. (2010). The molecular underpinning of lobular histological growth pattern: a genome-wide transcriptomic analysis of invasive lobular carcinomas and grade- and molecular subtype-matched invasive ductal carcinomas of no special type. J. Pathol. *220*, 45–57. 10.1002/path.2629.

64. McCart Reed, A.E., Kutasovic, J.R., Lakhani, S.R., and Simpson, P.T. (2015). Invasive lobular carcinoma of the breast: morphology, biomarkers and 'omics. Breast Cancer Res. *17*, 12. 10.1186/s13058-015-0519-x.

65. Michaut, M., Chin, S.-F., Majewski, I., Severson, T.M., Bismeijer, T., de Koning, L., Peeters, J.K., Schouten, P.C., Rueda, O.M., Bosma, A.J., et al. (2016). Integration of genomic, transcriptomic and proteomic data identifies two biologically distinct subtypes of invasive lobular breast cancer. Sci. Rep. *6*, 18517. 10.1038/srep18517.

66. McCart Reed, A.E., Foong, S., Kutasovic, J.R., Nones, K., Waddell, N., Lakhani, S.R., and Simpson, P.T. (2021). The Genomic Landscape of Lobular Breast Cancer. Cancers *13*. 10.3390/cancers13081950.

67. Zong, W.-X., Rabinowitz, J.D., and White, E. (2016). Mitochondria and Cancer. Mol. Cell *61*, 667–676. 10.1016/j.molcel.2016.02.011.

68. Barrett, T.J., Corr, E.M., van Solingen, C., Schlamp, F., Brown, E.J., Koelwyn, G.J., Lee, A.H., Shanley, L.C., Spruill, T.M., Bozal, F., et al. (2021). Chronic stress primes innate immune responses in mice and humans. Cell Rep. *36*, 109595. 10.1016/j.celrep.2021.109595.

69. Sollazzo, M., De Luise, M., Lemma, S., Bressi, L., Iorio, M., Miglietta, S., Milioni, S., Kurelac, I., Iommarini, L., Gasparre, G., et al. (2022). Respiratory Complex I dysfunction in cancer: from a maze of cellular adaptive responses to potential therapeutic strategies. FEBS J. *289*, 8003–8019. 10.1111/febs.16218.

70. Pisco, A.O., and Huang, S. (2015). Non-genetic cancer cell plasticity and therapy-induced stemness in tumour relapse: "What does not kill me strengthens me." Br. J. Cancer *112*, 1725–1732. 10.1038/bjc.2015.146.

71. Gu, Z. (2022). Complex heatmap visualization. Imeta *1*. 10.1002/imt2.43.

72. Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J. (2019). g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). Nucleic Acids Res. *47*, W191–W198. 10.1093/nar/gkz369.

73. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat. Methods *9*, 676–682. 10.1038/nmeth.2019.

74. Gentleman, R.C., Carey, V.J., Bates, D.M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B., Gautier, L., Ge, Y., Gentry, J., et al. (2004). Bioconductor: open software development for computational biology and bioinformatics. Genome Biol. *5*, R80. 10.1186/gb-2004-5-10-r80.

75. Pagliarini, D.J., Calvo, S.E., Chang, B., Sheth, S.A., Vafai, S.B., Ong, S.-E., Walford, G.A., Sugiana, C., Boneh, A., Chen, W.K., et al. (2008). A mitochondrial protein compendium elucidates complex I disease biology. Cell *134*, 112–123. 10.1016/j.cell.2008.06.016.

Supplementary Information

Figure S1.
A-C. Identification and initial characterisation of biclusters from the METABRIC dataset. Silhouette analysis of biclusters identified using the MB1 (**A**) and MB2/MB3 (**B**) genesets (see methods). **C.** Comparison of whole transcriptome CV values between all biclusters found. Density plots of the distribution of CVs across the transcriptome are shown for each bicluster in the central plots. Nuclear encoded mitochondrial genes are shown in red, non-mitochondrial genes are in cyan. The red dashed square indicates the three biclusters identified as independent, and carried over for further analysis.
D-E. Identification of the MB1-MB3 biclusters in independent datasets. **D** shows comparison of whole transcriptome CV values from METABRIC, TCGA-microarray, TCGA-RNAseq and Oslo2 datasets. Density plots of the distribution of CVs across the transcriptome are shown for each bicluster/dataset in the central plots. Nuclear encoded mitochondrial genes are shown in red, non-mitochondrial genes are in cyan. **E** shows the MB1 switch in 2D distribution by PC1 and ranking index in the four different dataset. Pam50 classifiers are overlaid to show the similarity of sample distribution in the different switches (biclusters). The result is representative of all three (MB1-MB3 biclusters).
F. Average gene group expression values in the upper and lower switch position in the MB1 switch in all datasets (METABRIC, Oslo2, TCGA-microarray, TCGA-RNAseq and TCGA-RRPA). Gene group1 (left panel) and gene group2 (right panel) are the two anticorrelated gene groups discovered by MCbiclust, as shown in Figure 1A. Accordingly, gene group1 shows positive correlation, and higher expression in upper fork samples as compared to the lower fork where conversely, group2 shows positive correlation and higher expression. The direction is arbitrarily set and harmonised between the biclusters. Turkey's box and whisker plots of the mean of the average gene group expression values across the indicated sample group (Lower, Upper Forks). The result is representative of all three (MB1-MB3) biclusters.

Figure S2.
A. Mapping of highly scoring metabolic pathways from the METABRIC dataset on the global metabolic network. The MB1_UF (top left panel) vs MB1_LF switch (top right panel, blue) is represented. Pathways where any of the enzyme transcripts had a CV over 0.5 (MB1_UF, orange monocolor scale between CV = 0.5-1.0) or under -0.5 (MB1_LF, blue monocolor scale between CV = -0.5 - -1.0) are highlighted. The lower panels show the main domains of the global metabolic network (KEGG, lower left), and the difference between CVs in the two switch positions (lower right, red (max) - white (0) - blue (min) color scale).
B. Intra-pathway details of individual gene transcript CVs. Supplementary to Figure 2A, showing pathways with high diversity. CV values of individual genes in the METABRIC breast cancer dataset analysis shown for the KEGG pathways: oxidative phosphorylation, fatty acid synthesis and degradation, purine and pyrimidine synthesis. Values are ranked by the MB_1 CV value. The figure allows the identification of high CV genes and where differences are large between the switches (biclusters).

Figure S3.
Statistical analysis of PARADIGM parameters across switch positions. Average PARADIGM values were calculated for each gene in each P cluster (Table 3), and compared between samples of each switch position. Kruskall-Wallis non-parametric test with Holm-corrected pairwise Dunn test was applied.

Figure S4.
A. Overlap between classification of METABRIC breast cancer samples by switch positions (MB1-3 UF and LF) and PAM50 intrinsic subtypes[58].
B. Distribution of different cell-of-origin gene expression scores across the MB1 and MB2 switches.

Figure S5.

A. Heatmap of fractional enrichment (FE) of ^{U13}C-glucose derived carbons in isotopologues of key metabolites analysed in MB1_LF (left panel) and MB1_UF (right panel) cells.

B. Overall carbon contribution into key metabolites following ^{U13}C-glucose (top left panel), ^{U13}C-glutamine (top right panel) and ^{U13}C-pyruvate (bottom panel) labelling in in MB1_LF and MB1_UF cells under different conditions (see Figure 4A).

Figure S6.

A. Quantitative analysis of enzyme expression levels obtained from western blot analysis (Fig. 5A**).** Values were normalised to either mitochondrial (left panel) or cytosolic (right panel) markers, grp75 or β-actin. 2-way ANOVA with uncorrected Fisher's LSD multiple comparison tests.

Figure S7.

A. Visualisation of further histological feature distributions on the MB1 and MB2 switches (see Figure 6B). Classifiers of mitotic activity (left panel), fraction of epithelial area (middle panel) and presence of inflammation (right panel) are overlaid onto 2D distribution plots of TCGA samples along the axes representing the scale between the upper (UF) and lower (LF) forks of each bicluster.

Tables list S1-8

Table S1. METABRIC data: sample characteristics and gene sets

Table S2. Unique and overlapping genesets in the MB1 and MB2 switches, together with CV values of the whole transcriptome

Table S3. Gene set enrichment analysis (KEGG)

Table S4. Gene set enrichment analysis (Mitocarta)

Table S5. Top KEGG and Mitocarta pathways

Table S6. PARADIGM gene sets

Table S7. Breast cancer cell line scoring results

Table S8. TCGA BRCA sample classification data with histological and overall features

Supplementary Figure S1

A

B

Figure S2

Figure S3

Figure S5

A

A

STAR METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit monoclonal anti-Hexokinase 1	Cell Signaling Technology	Cat# 2024, RRID:AB_2116996
Rabbit monoclonal anti-Hexokinase 2	Cell Signaling Technology	Cat# 2867, RRID:AB_2232946
Rabbit polyclonal anti-Triosephosphate isomerase –N-terminal	Abcam	Cat# ab135532
Monoclonal antibody anti-Glyceraldehyde-3-phosphate dehydrogenase	Millipore	Cat# MAB374, RRID:AB_2107445
Rabbit monoclonal anti-Pyruvate kinase M1 isofrom	Cell Signaling Technology	Cat# 7067, RRID:AB_2715534
Rabbit monoclonal anti-Pyruvate kinase M2 isoform	Cell Signaling Technology	Cat# 4053, RRID:AB_1904096
Rabbit monoclonal anti-Lactate dehydrogenase A	Cell Signaling Technology	Cat# 3582, RRID:AB_2066887
Rabbit monoclonal anti-Lactate dehydrogenase B	Abcam	Cat# ab53292, RRID:AB_2234531
Rabbit polyclonal anti-BRP44L (MPC1)	Abcam	Cat# ab74871, RRID:AB_1523226
Rabbit monoclonal anti-MPC2	Cell Signaling Technology	Cat# 46141, RRID:AB_2799295
Mouse monoclonal anti-Pyruvate dehydrogenase E1-alpha subunit	Abcam	Cat# ab110330, RRID:AB_10858459
Rabbit polyclonal anti-pyruvate dehydrogenase E1-alpha subunit (phospho S293)	Abcam	Cat# ab92696, RRID:AB_10711672

Rabbit polyclonal anti-IDH1	Proteintech Group	Cat# 12332-1-AP, RRID:AB_2123159
Mouse monoclonal anti-IDH2	Abcam	Cat# ab55271, RRID:AB_943793
Rabbit polyclonal anti-IDH3A	Abcam	Cat# ab58641, RRID:AB_2295799
Rabbit polyclonal anti-glutaminase	Abcam	Cat# ab93434, RRID:AB_10561964
Rabbit monoclonal anti-GLUD1	Abcam	Cat# ab166618, RRID:AB_2815030
Rabbit polyclonal anti-GOT1	Proteintech Group	Cat# 14886-1-AP, RRID:AB_2113630
Rabbit polyclonal anti-GOT2	Proteintech Group	Cat# 14800-1-AP, RRID:AB_2247898
Rabbit polyclonal anti-ME1	Abcam	Cat# ab97445, RRID:AB_10679994
Rabbit monoclonal anti-ME2	Abcam	Cat# Ab139686
Rabbit monoclonal anti-PCB	Abcam	Cat# ab128952, RRID:AB_11144490
Rabbit polyclonal anti- 3PGDH (PHGDH)	Santa Cruz Biotechnology	Cat# sc292792
Mouse monoclonal anti-PPAT	Abcam	Cat# ab128116, RRID:AB_11143951
Rabbit polyclonal anti-PSAT1	Proteintech Group	Cat# 20180-1-AP, RRID:AB_10665948
Rabbit monoclonal anti-MDH1	Abcam	Cat# ab180152
Mouse anti - GLUL	BD Biosciences	Cat# 610518 RRID:AB_397880
Rabbit monoclonal anti-GFPT1 [EPR4854]	Abcam	Cat# ab125069, RRID:AB_10975709
Recombinant anti-GFPT2	Abcam	Cat# ab190966, RRID:AB_2868470

Total OXPHOS rodent antibody cocktail	Abcam	Cat# ab110413, RRID:AB_2629281
Mouse monoclonal anti-Actin	Santa Cruz Biotechnology	Cat# sc-56459, RRID:AB_830981
Rabbit polyclonal anti-beta tubulin	Santa Cruz Biotechnology	Cat# sc-9104, RRID:AB_2241191
Rabbit polyclonal anti-grp75	Santa Cruz Biotechnology	Cat# sc-13967, RRID:AB_647720
Chemicals, Peptides, and Recombinant Proteins		
[U-^{13}C] glucose	Cambridge Isotope Laboratories, Inc.	CLM-1396
[U-^{13}C] glutamine	Cambridge Isotope Laboratories, Inc.	CLM-1822
[U-^{13}C] pyruvate	Cambridge Isotope Laboratories, Inc.	CLM-1575
Pyruvate	Thermo Fisher Scientific	Cat# 11360070
Glutamine	Thermo Fisher Scientific	Cat# 25030081
Glucose	Thermo Fisher Scientific	Cat# A2494001
Normocin	InvivoGen	ant-nr-1
Hoechst 33342	Thermo Fisher Scientific	Cat# 62249
Tetramethylrhodamine, Methyl Ester, Perchlorate (TMRM)	Thermo Fisher Scientific	Cat# T668
Quant-iT™ PicoGreen™ dsDNA Assay Kits and dsDNA Reagents	Thermo Fisher Scientific	Cat# P11495
Rotenone	Sigma-Aldrich	Cat# R8875
Carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP)	Sigma-Aldrich	Cat# C2920
Antimycin A from Streptomyces sp.	Sigma-Aldrich	Cat# A8674
Oligomycin A	Sigma-Aldrich	Cat# 75351
DMEM, high glucose, pyruvate	Thermo Fisher Scientific (Gibco)	Cat# 41966029

DMEM, high glucose, glutaMAX supplement, pyruvate	Thermo Fisher Scientific (Gibco)	Cat# 31966021
DMEM, no glucose, no glutamine, no phenol red	Thermo Fisher Scientific (Gibco)	Cat# A1443001
Critical Commercial Assays		
Glutamine/Glutamate-Glo Assay	Promega	Cat# J8022
CellTiter-Glo Luminescent Cell Viability Assay	Promega	Cat# G7570
Seahorse XF Cell Mito Stress Test Kit	Agilent	Cat# 103015-100
Experimental Models: Cell Lines		
MCF7	ATCC	HTB-22
T47D	ATCC	HTB-133
Hs578T	ATCC	HTB-126
MDA-MB436	ATCC	HTB-130
HCC1143	ATCC	CRL-2321
Source Data		
METABRIC	Synapse: syn1757063	https://doi.org/10.7303/syn1688369
TCGA	NCBI dbGaP	phs000178.v11.p8
OsloVal	Synapse: syn1710395	https://doi.org/10.7303/syn1688370
Software and Algorithms		
R v4.2.0	R Foundation for Statistical Computing	https://www.r-project.org/
MCbiclust 1.22.0	Bentham et al.[28]	https://doi.org/doi:10.18129/B9.bioc.MCbiclust
ComplexHeatmap	Gu[71]	https://doi.org/doi:10.18129/B9.bioc.ComplexHeatmap
GSVA 1.46.0	Hänzelmann et al[41]	https://doi.org/doi:10.18129/B9.bioc.GSVA
g:Profiler	Raudvere et al[72]	http://biit.cs.ut.ee/gprofiler/gost

Fiji	Schindelin et al.[73]	https://imagej.net/software/fiji/
Prism 9.0	GraphPad Software	https://www.graphpad.com/
Other		
SnakeSkin Dialysis Tubing, 3.5K MWCO, 35 mm dry I.D., 35 feet	Thermo Fisher Scientific	Cat# 88244
Seahorse XF96 cell culture microplates	Agilent	Cat# 102416
Falcon® 96-well Black/Clear Flat Bottom TC-treated Imaging Microplate	Corning Life Sciences	Cat# 353219

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Gyorgy Szabadkai (g.szabadkai@ucl.ac.uk).

Materials availability
This study did not generate new unique reagents.

Data and code availability
All data are available via links provided in the Methods section and Key Resources table. Code for the full analysis pipeline is available from the authors upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell lines
Human breast cancer cell lines MCF7, T47D, Hs578t, MDA436 and HCC1143 were purchased from ATCC. Their identity was confirmed through cell line authentication carried out by BMR Genomics (Italy) by amplification of 23 STR loci (PowerPlex Fusion System kit, Promega). Cells were cultured in Dulbecco's modified Eagle's medium (DMEM, #41966029), supplemented with 10% Fetal Bovine Serum (FBS) (Thermo Fisher Scientific) and Normocin (0.1 mg/mL, InvivoGen) and maintained in a 37°C incubator set with 5% CO2 and 95% humidity. Cell numbers for each experimental setting were counted using a hemocytometer.
For the restricted nutrient condition and stable isotope labelling experiments, cells were cultured using DMEM (no glucose, no glutamine, no phenol red, Thermo Fisher Scientific #A1443001) complemented with glucose, glutamine and pyruvate, as indicated. For the stable isotope labelling, the required amount of labelled compounds, [U-^{13}C] glucose, [U-^{13}C] glutamine and [U-^{13}C] pyruvate, were added. FBS was dialyzed using a SnakeSkin dialysis tubing (Thermo Fisher Scientific) and filtered using 0.22 μm PES filter before being added in 10% proportion, together with Normocin. The conditions were named according to their nutrient concentrations either stably labelled or not (Figure 4A): high and low glutamine (H/L

= 1mM/0.1mM) levels were combined with the presence and absence of pyruvate (+/- = 1mM/0mM), glucose was kept at 10 mM.

METHODS DETAILS

Discovery of large gene cluster switches with MCbiclust

MCbiclust (doi:10.18129/B9.bioc.MCbiclust, current version 1.2.1, R package available in Bioconductor[74]) was used to identify gene cluster switches as described previously[28,29] on the METABRIC[31], TCGA[32] (BRCA) and Oslo2[33] datasets. We have used two defined and two random starting genesets of approximately 1000 genes. The defined genesets were either only mitochondrial (Mitocarta 1.0[75]) or mitochondria related, by choosing the 1000 genes most correlated with the mitochondrial ribosomal gene MRPL58 across the METABRIC dataset. MCbiclust was run for 1000 iterations using each geneset on the University College London high performance cluster. The initial bicluster seeds found are shown in Figure S1, following the selection of the highest correlating gene groups using the HclustGenesHiCor function. The silhouette method was used to determine the number of independent biclusters. Biclusters were then extended (i) by calculating the correlation vector (CV) values for the whole transcriptome using the CVEval function, and (ii) to all samples using SampleSort. The correlation pattern was summarised using principal component analysis (PCA) of the CV values using PC1VecFun. The bifurcating patterns, defining the forks (switch positions), were plotted using the sort order (index) and PC1 values for each sample (see Figures 1B, S1).

Gene set enrichment and pathway analysis

Functional analysis of gene clusters were performed with g:Profiler[72]. The default KEGG data source was used in g:GOSt for functional profiling of all metabolic pathways. The KEGG results were manually filtered to include only metabolic pathways. For mitochondrial functional profiling the custom MitoCarta 3.0 orthology was used as data source[34] in g:Profiler. Signalling pathway activity was assessed using the PARADIGM integrated pathway levels (IPLs) from the PARADIGM[37] analysis on a pan-cancer dataset[35] (https://www.synapse.org/#!Synapse:syn5633407), filtered for BRCA samples. Gene set variation analysis for molecular signature identification was performed using the GSVA 1.46.0 package[41].

Assessment of mitochondrial activity

Oxygen consumption rate was measured with the Seahorse XFe96 bioanalyzer using the Seahorse XF Cell Mito Stress Test Kit (Agilent), or using High-Resolution Respirometry (Oxygraph-2K, Oroboros Instruments, Austria). For the Seahorse measurements cells were seeded on XF96 cell culture microplates (Agilent) 2 days before the experiment (1×10^4 cells/well). For the experiment, the culture medium was replaced with Seahorse XF Base medium (Agilent) supplemented with pyruvate, glutamine and glucose (Thermo Fisher Scientific) as indicated, and incubated for 30 min at 37 °C in a CO_2-free incubator before loading into the Seahorse Analyser. After measuring basal respiration, the drugs oligomycin A (5 µM), FCCP (1 µM), and rotenone/antimycin A (0.5 µM/0.5 µM) were added to each well in sequential order. Data were analysed using the XF Cell Mito Stress Test Report Generator. After the assay, cells were stained with Hoechst 33342 (1 µg/mL; Thermo Fisher Scientific) for 30 min. ImageXpress Micro XL was then used for cell nuclei (cell numbers) counts in each well.

For High-Resolution Respirometry the Oxygraph-2K instrument was used. Prior to the assay, OROBOROS chambers were calibrated with the recording media (DMEM #A1443001, glucose (10 mM), glutamine (1mM), pyruvate (1mM), HEPES (10 mM)). Confluent 10 cm plates of cells were trypsinized and re-suspended in the same media. $2.5*10^6$ cells were added to each chamber, and the O_2 flow signal was allowed to stabilise to basal respiration rate. Drugs were added to chambers using in the following concentrations and order: Oligomycin (2.5µM), FCCP (2µM) and Antimycin A (2.5 µM). Respiratory rate given as oxygen flow in pmol/min/cell was recorded. Data collection and analysis was performed using Datlab 5.0 software.

Imaging

For imaging mitochondrial structure and function, a wide-field high content imaging system (ImageXpress MicroXL, Molecular Devices) was used. Cells were seeded in 96-well black clear thin bottom tissue culture treated imaging plates (Corning). For mitochondrial membrane potential measurements cells were incubated with 30nM TMRM (Thermo Fisher Scientific) in recording media (DMEM #A1443001, glucose (10 mM), glutamine (1mM), pyruvate (1mM), HEPES (10 mM)) for 30 minutes prior to starting imaging and left for the whole time of the experiment at 37ºC. For labelling the nuclei cells were co-loaded with Hoechst 33342 (1 µg/mL; Thermo Fisher Scientific) for 30 min. For TMRM intensity quantification a 20X Nikon (S PLAN FLUOR ELWD 0.45 NA) air objective was used. Images were analysed with the integrated metaXpress imaging and analysis software using the granularity analysis module. Following local background subtraction, the average TMRM intensity/field was used as readout.

For morphology analysis a Nikon 40X (Plan Apo 0.95 NA) air objective was used. Cells were co-loaded with TMRM (30 nM), picoGreen (2.5µL/mL) and Hoechst 33342 (1 µg/mL; Thermo Fisher Scientific) in recording media (DMEM #A1443001, glucose (10 mM), glutamine (1mM), pyruvate (1mM), HEPES (10 mM)) for 30 minutes prior to starting imaging. TMRM was left for the whole time of the experiment at 37ºC. Cells were imaged using a custom protocol to image 3 wavelengths (Lumencor solid state illumination with Semrock [Brightline] filters (nm) ex377/50 em447/60 for Hoechst 3342; ex472/30 em520/35 nm for picoGreen, ex531/40 em593/40 for TMRM), 16 fields/well with a Nikon 40X Plan Apo NA 0.95 air objective, binning 1 with a CMOS detector. TMRM and picoGreen Images were analysed with MetaXpress 5.0 software, using the granularity module, total mitochondrial area and average object size as readout from the TMRM images. For picoGreen analysis, the integrated pit intensity (granularity module) was used as a readout following ratioing with the Hoechst images, background subtraction and thresholding.

Determination of total cellular ATP

Total cellular ATP concentration was measured using CellTiter-Glo® Luminescent Cell Viability Assay (Promega). The method determines the number of viable cells based on the quantification of the ATP present, as readout of metabolically active cells, thus allowing the estimation of total cellular ATP produced by a known number of seeded cells. In detail, 5K cells were seeded in an opaque black 96-well plate and treated with the experimental media (Figure 4A) for 24h. Wells containing the experimental medium only (the blank - for removing the medium background) and 10nM ATP standard (on which the sample's luminescence was normalised to) were included. The protocol was carried out according to the manufacturer's instructions. Briefly, i): plates were equilibrated at room temperature (RT) for approximately 30 minutes, avoiding temperature gradients which could cause uneven signals; ii) a volume

of CellTiter-Glo® Reagent equal to the volume of cell culture medium present in each well (e.g., 100µl of reagent to 100µl of medium containing cells for a 96-well plate) was added; iii) 2 minutes of orbital shaking favouring cell lysis; then, iv) plates were left at RT for 10 minutes to stabilise luminescent signal and v) the luminescence was recorded using PerkinElmer EnVision plate reader. For each plate the triplicate values for the samples total ATP produced, the ATP standard and five replicates of medium luminescence for the background were acquired. The average of the replicates was calculated, and the medium background was subtracted. The resulting values were normalised to the ATP standard previously adjusted for the background and further normalised to the protein content, evaluated using the BCA protein assay (Pierce).

Western Blotting
For the baseline expression of metabolic enzymes and mitochondrial respiratory chain subunits at protein level, proteins were extracted from cells plated in 10 cm dish and cultured in DMEM. Briefly, cells were detached by scraping in cold PBS 1X and pulled down by centrifugation. Whole cell lysate was obtained incubating cell pellet with cold RIPA buffer (150 mM NaCl, 50 mM Tris, 0.5% sodium deossicolate, 0.1% (w/v) SDS, 1% (v/v) Triton) in presence of protease inhibitor cOmplete (Roche), phosphatase inhibitors PhosStop (Roche) and PMSF (Sigma) for 30 minutes on ice and spun down at 4 °C. Protein quantification was done using a BCA protein assay kit (Pierce). Samples were denatured in presence of reducing agents (DTT) at 95°C for 5 minutes, while for probing mitochondrial respiratory chain subunits the optimal temperature was 60°C for 10 minutes. 20 µg of protein lysate was loaded onto a NuPAGE 4-12% bis-tris gel (Thermo Fisher) and run using MOPS 1X buffer. Blotting was done for 2 h at 30 V in a wet system (Invitrogen - Thermo Fisher), by transferring proteins onto nitrocellulose membrane for metabolic enzymes and PVDF for mitochondrial subunits. Membranes were incubated in 5% non-fat dry milk in TBS 1X – 0.1% Tween 20 or 5% BSA in TBS 1X – 0.1% Tween 20 as blocking buffers (according to each antibody datasheet) for 1 h at room temperature. Primary antibody was incubated in blocking buffer overnight at 4°C. The working dilution was 1:1000 for the most of antibodies except for MitoProfile (1:2000), grp75 (1:3000), actin (1:3000). Horseradish-conjugated secondary antibodies (BioRad) were diluted 1:5000 in blocking buffer and membranes were incubated 1 h at room temperature. The blots were visualised using SuperSignal West Pico Chemiluminescent Substrate (Thermo Fisher) and UVITEC Cambridge Mini HD9 Imaging System (Eppendorf). If necessary, membranes were stripped, blocked and probed again with a different primary antibody.

Measurement of glutamine and glutamate concentration in the media
The concentration of glutamine and glutamate in the media were measured using the luminescence-based approach Glutamine/Glutamate-Glo TM Assay kit (Promega). The method is based on two steps: i) glutamine is first converted to glutamate by glutaminase; ii) glutamate is oxidised and NADH produced is used for luminescence reading. Briefly, 10k cells were seeded in an opaque black 96-well plate. On the following day, after one wash with warm PBS1X, cells were treated with the experimental media (Figure 4A) for 24 h. Wells containing the experimental medium only (no cells), 5 points glutamate standard curve (to extrapolate glutamine and glutamate concentration) and wells containing PBS1X (as negative control) were included. Wells with only medium represent a positive control and were used as reference of initial glutamine and glutamate concentration in the medium, on which values subtract the glutamine and glutamate concentration obtained from cells

containing wells, thus calculating the glutamine uptake and glutamate secreted by the cells. The addition and removal of both PBS 1X and media were gently done using a multichannel pipette. Each experimental medium was freshly prepared the day of the treatment. The experiments were carried out according to the manufacturers' instructions. After 24 hours treatment, media were collected and a dilution of 1:25 (in PBS1X) was done in order to fit the linear range of the assay indicated by the commercial kit. The diluted samples were analysed the same day or frozen at -80°C and processed afterwards. The assay was done at room temperature with media and reagent RT equilibrated to avoid uneven signals. Luminescence was recorded using Infinite M200 plate reader (Tecan). Protein content was evaluated using the BCA protein assay (Pierce). The results were presented as nmoles/minutes/grams of protein.

Stable isotope labelling, metabolite extraction and quantification
Cells were grown at standard culturing conditions as described in cell culture methods; the stable-isotope labelling was obtained by culturing cells with media containing either fully labelled [U-^{13}C] glucose, [U-^{13}C] glutamine or [U-^{13}C] pyruvate for 24 h. Before metabolite extraction, plates were taken to a cold room and 500μL of medium from each plate was frozen for later analysis. The remaining media was removed, and the plates placed in Fan ice/water bath before washing twice with ice-cold PBS.
For metabolites extraction, 800μL ice-cold methanol containing an internal standard of 1mM scyllo-inositol was added to each plate and cells were detached by scraping. This mixture was transferred to a 15ml tube, and the plate washed with 800μL of methanol:H_2O (1:1 vol/vol) that was moved to the tube, to which 400μL of ice-cold chloroform was added. The tubes were placed in a water bath sonicator in a cold room for 1 h, with 3x8 minute pulses of sonication and centrifuged for 10 minutes at 16,000rpm at a temperature of 0°C. The supernatant was extracted and dried in a vacuum concentrator. The cell pellet was then re-extracted with 300μL of methanol:H_2O (2:1 vol/vol), this was sonicated, spun and the supernatant added to the previous supernatant tube and dried again in a vacuum concentrator. The remaining cell pellet was used for estimating dry weight and measuring total protein. For phase partitioning, the dried supernatant was resuspended in 350μL of chloroform:methanol:H_2O (1:3:3 vol/vol) and spun for 5 minutes at 0°C and 16,000rpm. The extract is then in a biphasic partition, with the upper phase containing the polar metabolites and the lower phase containing lipidic metabolites. The polar phase fractions were then transferred to GC-MS vial inserts and vacuum-dried. Separate vial inserts had 10μL of the saved cell culture medium added, with 1mM scyllo-inositol, which were also dried in a vacuum concentrator. Each vial insert had 30μL of methanol added, containing 1μL of 5mM nor-leucine as another internal standard, followed by 30μL of methanol without nor-leucine, with the vials being dried in a vacuum concentrator after each addition. Before running samples on the mass spectrometer, polar fractions were derivatised, 20μL methoxyamine (30mg/mL in pyridine) was added to each insert and this was vortexed briefly and incubated at room temperature overnight, Silylation was then done by adding 20μL of BSTFA + TCMS reagent to each insert and incubating for 1 h at room temperature.
For metabolites analysis, an Agilent 7890A GC with a 5975C triple axis detector MSD (Agilent Technologies, Santa Clara, CA) was used. Metabolites were separated on an Agilent J&W 122-5532G DB-5ms capillary column (30m x 0.25mm, 0.25um film thickness), in splitless mode. The injector and transfer line temperatures were 270 and 280°C, respectively. The flow rate of helium carrier gas was 0.7 mL/min. The oven temperature was programmed to hold at 70°C for 2 min, increased to 295°C at a 12.5°C/min ramp rate,

increased from 295°C to 320°C at a 25°C/min ramp rate, and held at 320°C for 3 minutes. The mass spectrometer was operated in scan mode, after a 6 minute solvent delay with a range of 50−565 mass/charge (m/z) and a scan-rate of 2.8 scans per second. Metabolites were identified by matching retention times and fragmentation patterns to commercially available standards. Metabolite peaks were integrated at each isotopologue m/z using MassHunter Workstation software (Agilent Technologies). Peak areas were quantified based on peak areas of known standards using nor-leucine as an internal standard, and then metabolite levels were normalised to protein content. Mass isotopologues were stripped of the contribution from natural abundance, based on the chemical formula of derivatized fragments quantified. Percent enrichment for an isotopologue was calculated by dividing the corrected intensity by the sum of corrected intensities of all isotopologues for that metabolite. Significance of metabolite enrichment between different samples was calculated with one-way ANOVA.

QUANTIFICATION AND STATISTICAL ANALYSIS
Experimental design and statistical analysis are described in the Figure legends.

Acknowledgments

We thank for valuable discussions with C. Barnes, M. Duchen (University College London), K. Bianchi, C. Chelala (Barts Cancer Institute), for the support from all members of the M. Duchen, R. Rizzuto (University of Padova), MY, GS labs, for the PARADIGM data provided by R. Akbani (MD Anderson Cancer Center), and for the support from the Bioinformatics (G. Kelly) and Metabolomics (J. Macrae) STPs of The Francis Crick Institute. GS was supported by the following grants: AIRC (Associazione Italiana per la Ricerca sul Cancro, IG 13447 and 22221), Wellcome Trust Pathfinder Award (204458/Z/16/Z), Biochemical and Biophysical Research Council (BB/L020874/1 and BB/P018726/1) and CRUK Pioneer Award (29264). MY was supported by the Francis Crick Institute, which receives its core funding from Cancer Research UK (FC001223), the UK Medical Research Council (FC001223) and the Wellcome Trust (FC001223, FC0010060); MY and GS was supported by CRUK Grand Challenge Award 2015 C57633/A25043. MM and RB were supported by the Doctoral School Of Research In Biosciences And Biotechnologies at the University of Padova, and a COMPLeX UCL/British Heart Foundation PhD studentship, respectively. ZR is currently supported by the British Heart Foundation (FS/20/4/34958). RCS was supported by the National Institute for Health Research University College London Hospitals Biomedical Research Centre.

Author Contributions

Conceptualization: RB, MM, KB, MY, RCS, GS; Methodology: RB, MM, MY, GS; Software: RB, KB, GS; Formal Analysis: RB, MM, NP, SQN, TH, AG, ZR, CE, GS; Investigation: RB, MM, TH, ZR, CE, SA, CL, SI, ZS, MS, AG, MY, GS; Writing – Original Draft: GS; Writing – Review & Editing: RB, MM, MY, GS; Funding Acquisition: MY, GS.

Declaration of interests

The authors declare no competing interests. RCS is the Chief Investigator of the OPTIMA Trial (ISRCTN42400492) which uses the Prosigna test to make therapeutic decisions in breast cancer treatment. He has no personal financial links with the vendors, NanoString Inc. and Veracyte Inc.