Title: *Diversity in the internal functional feeding elements of sympatric morphs of Arctic charr (Salvelinus alpinus)*

Authors: Jónsdóttir, Guðbjörg Ósk [1], von Elm, Laura-Marie[1], Ingimarsson, Finnur[2], Tersigni, Samuel[1], Snorrason,

Sigurður Sveinn[1], Pálsson, Arnar [1] and Steele, Sarah Elizabeth[1,3]*

Affiliations

1.   Institute of Life- and Environmental Science, University of Iceland.

2.   Kópavogur Nature Center, Kópavogur, Iceland.

3.   Canadian Museum of Nature, Ottawa, Canada.

Corresponding author:

* Steele, S.E, sar.e.steele@gmail.com

Keywords:

Craniofacial morphology, Sympatric polymorphism, Geometric morphometrics, Jaw bones, Dentary, Salmonids.

Abstract

The diversity of functional feeding anatomy is particularly impressive in fishes and correlates with various interspecific ecological specializations. Intraspecific polymorphism can manifest in divergent feeding morphology and ecology, often along a benthic–pelagic axis. Arctic charr (*Salvelinus alpinus*) is a freshwater salmonid known for morphological variation and sympatric polymorphism and in Lake Þingvallavatn, Iceland, four morphs of charr coexist that differ in preferred prey, behaviour, habitat use, and external feeding morphology. We studied variation in six upper and lower jaw bones in adults of these four morphs using geometric morphometrics and univariate statistics. We tested for allometric differences in bone size and shape among morphs, morph effects on bone size and shape, and divergence along the benthic-pelagic axis. We also examined the degree of integration between bone pairs. We found differences in bone size between pelagic and benthic morphs for two bones (dentary and premaxilla). There was clear bone shape divergence along a benthic–pelagic axis in four bones (dentary, articular-angular, premaxilla and maxilla), as well as allometric shape differences between morphs in the dentary. Notably for the dentary, morph explained more shape variation than bone size. Comparatively, benthic morphs possess a compact and taller dentary, with shorter dentary palate, consistent with visible (but less prominent) differences in external morphology. As these morphs emerged in the last 10,000 years, these results indicate rapid functional evolution of specific feeding structures in arctic charr. This sets the stage for studies of the genetics and development of rapid and parallel craniofacial evolution.

Introduction

The origins and maintenance of biodiversity are strongly influenced by natural selection acting upon intraspecific variation with fitness consequences [1-3], with many traits directly affecting survivorship and reproduction exhibiting extensive diversification. Adaptive evolution in vertebrates entails a substantial degree of variation in feeding behaviour and biomechanics of prey capture and processing that may lead to speciation [4-6]. Morphological adaptations in functional feeding elements are associated with rapid diversification of ecology in many vertebrate groups, sometimes resulting in specialization on diverse prey types within lineages [4, 7]. The development of these feeding elements is influenced by an interplay of genes and environmental factors during ontogeny [e.g., 8]. Additionally, developmental remodelling can accommodate ontogenetic niche shifts, for instance within fishes [4, 8], and may contribute to intraspecific and interspecific variation in adult morphology.

Extensive complexity and diversification of skull anatomy among fishes correlates with highly diverse feeding ecologies [4, 9-11]. Evolution of several functional units [9] has given rise to a diversity of feeding methods (e.g., ram, suction, biting) for optimal foraging on specific prey types, varying widely within and among species [9, 10]. Independent evolution of oral jaws, hyoid "jaws" and pharyngeal jaws for prey capture and food processing likely contributed to trophic and morphological diversification [4]. Radiations of ecomorphology and associated prey have occurred widely in fishes, including Pomacentridae [12], Labridae [13], and Cichlidae [14]. While most studies investigate morphological variation among species or higher taxa [9-11, 15-17], extensive variation in cranial morphology also exists within species [16, 18].

Sympatric polymorphism, where two or more morphs within a species inhabit the same geographic area, has been found in many lacustrine fish species and seems to be promoted by vacant niches, habitat variance, differences in spawning preferences, and relaxation of interspecific competition [19, 20]. Resulting morphological divergence is often due to differences in resource use, mainly in benthic vs. pelagic habitats, leading to specializations in traits related to foraging, prey capture and processing [21]. The occurrence of intraspecific morphs that utilize different resources (e.g., prey, habitats) has been widely documented in northern freshwater fishes inhabiting recently de-glaciated systems [21, 22]. Salmonid cranial morphology is less derived compared to more divergent groups [18, 23], yet within this group many cases of sympatric morphological polymorphism have been described [20, 24-26]. Arctic charr (*Salvelinus alpinus*) shows extensive phenotypic variation throughout its

geographic distribution [27] and sympatric polymorphism of lake populations is for instance found in Norway [e.g.,

28, 29], Siberia [e.g., 30], Greenland [e.g., 31] and Iceland [32, 33]. In Iceland, the best example of sympatric

polymorphism is Lake Þingvallavatn, Iceland's largest natural lake [34]. The lake formed ~10,000 years ago in a rift

zone, and is currently a deep lake with lava rocks in most of its banks and shallows [35]. It was colonized by

anadromous charr [36, 37] that subsequently became isolated. The lake now hosts four morphs of arctic charr (Fig

1A and 1B), two benthic: large benthivorous (LB) and small benthivorous (SB) charr and two pelagic: planktivorous

(PL) and piscivorous (PI) charr [38, 39]. The LB-, SB- and PL-charr are genetically distinct [37, 40], while the PI-

charr is more heterogeneous genetically, with similarity to either LB- or PL-charr [40]. The morphs differ in adult

size, diet, habitat use, life history, spawning times, and external and fin morphology [39, 41-45]. Two are large (LB

and PI) and two small (SB and PL), hereafter referred to as the morph size gradient. Benthic morphs live along the

stony littoral bottom and feed mainly on snails [41], and have subterminal mouths (i.e., short lower jaw) and blunt

snouts [45, 46]. Pelagic morphs more resemble anadromous charr, being more fusiform with a terminal mouth (i.e.,

long lower jaw) and pointed snout [45, 46]. PL-charr feed mostly on crustacean zooplankton, and the larger PI-charr

feed mostly on three-spined stickleback [41, 46]. Most studies on arctic charr diversity have examined external

morphology, except studies of gill raker counts in adults [45], fin ray variation in lab reared juveniles [47] and an

unpublished study of internal feeding elements in adults [48]. Notably Ingimarsson (48), using linear measures,

found differences among morphs in maxilla and dentary bone shape, relative size of several skull elements, number

of teeth on the dentary and palatine bones, as well as the angle of teeth on the maxilla, with divergence mainly along

the benthic–pelagic axis. The specific adaptations in cranial morphology by morph and the degree of integration of

variation in bone shape are understudied.

To investigate the functional morphology and impact of prey specialization during divergence of arctic

charr along the benthic-pelagic axis [33, 49], we studied cranial morphology in adults of the four sympatric morphs

in Þingvallavatn. To test for morphological divergence and integration in internal feeding structures among morphs,

we examined variation in the shape of six cranial bones in the feeding apparatus: dentary, articular-angular,

quadrate, premaxilla, maxilla and supramaxilla (Fig 1C). We focus on the following questions and predictions: 1) Is

there an allometric component for size or shape of specific bones, and does this vary by morph? We hypothesize that

size will influence shape, however, do not expect this allometry to differ by morph. 2) Does the shape of any of

those six bones differ among morphs? Given divergence in morph external shape [45], we expect bone shape to

differ among the morphs, most likely in the lower jaw and maxilla. 3) Does morphological differentiation occur

along a benthic-pelagic axis? Again based on external morphology, where benthic morphs have shorter jaws [45],

we expect shortening and compaction of the lower jaw bones (the dentary and/or articular-angular). We also

hypothesize proportionally narrower maxilla (main element of upper jaw) in the pelagic morphs, based on observed

external morphology [50]. 4) Does integration differ among pairs of bones, between the upper and lower jaw, and

across morphs? We expect significant integration between elements, with high integration among bones within the

same jaw, but less integration across the upper and lower jaw due to ecological divergence in the lower jaw bones.

Fig 1. External and internal morphology of Þingvallavatn arctic charr ecomorphs. A) Craniofacial variation; B) Postcranial shape and size variation; C) Primary upper and lower jaw bones studied; D – I) Anatomy of the upper and lower jaw bones with landmarks used to capture shape.

Methods

Sample and data collection

Sexually mature arctic charr from Lake Þingvallavatn were collected (Fall 2020/21) using composite nets and seine nets. Fish were randomly sampled, except we aimed for similar representation of morphs and sexes (LB: 49 (21%), SB: 82 (34%), PL: 61 (25%) and PI-charr: 48 (20%), S1 Appendix). Each fish was measured (weight, fork length (FL)), and photographed from left lateral view (Canon EOS 77D, EFS 18-135 mm lens), and sex, sexual maturity, and stomach contents were evaluated by dissection. Morph classification was based on sampling location and date (morphs differ in spawning times and location), external morphology and stomach contents [45, 50, 51]. Otoliths were extracted from each fish and age determined under 4X magnification in a stereomicroscope. Six jaw bones (dentary, articular-angular, quadrate, premaxilla, maxilla, supramaxilla) were extracted using maceration protocols [51] and photographed (Canon EOS 77D, 50 mm Microlens). The bones from the left side of the head were arranged stereotypically: the dentary and maxilla were photographed and landmarked in medial view, the articular-angular, quadrate and supramaxilla in lateral view, and the premaxilla in fronto-lateral view. To capture external morphology for morph classification, 37 2D landmarks (see S2 Appendix) were digitized for each individual using *tpsdig2* (http://www.sbmorphometrics.org/soft-dataacq.html). To study variation in internal feeding structures, we registered 2D landmarks on the six jaw bones with *tpsdig2* (Fig 1D-I: S3 Appendix). All landmarking (external and internal) was conducted by one person (GOJ). To test repeatability of the bone landmarks 168 specimens were landmarked twice. Two landmarks were excluded after the repeatability analysis, LM 12 for premaxilla (Fig 1G: S3 Appendix) and LM 15 for the articular-angular (Fig 1E: S3 Appendix) because they were added after the first trial. All images of fish and bones are available on Figshare (XXX to YYY).

Statistical analyses

Landmark data were analysed using the R package *geomorph* version 4.0.0 [52, 53]. Landmark coordinates were

aligned using Generalized Procrustes Analysis (GPA) via the function *gpagen*. Aligned shape variables were then

used for all subsequent analyses. Outliers were identified using *plotOutliers* and removed if the bone was damaged.

We tested the repeatability of the bone landmark sets in two ways, using Procrustes ANOVA, using *procD.lm*, to

characterize measurement error (differences between replicates), as well as two-block partial least squares (PLS)

analysis with the function *two.b.pls* to analyze the degree of association between the two replicates (trials). Centroid

size (CS) was calculated separately for each of the six bones (CS_{bone}; indicates use of CS of a bone, and applied to

each bone separately), head (CS_{head}) and whole-body (CS_{body}). As size is an important predictor for shape in fishes

[54, 55] and the morphs are known to differ in adult size and proportions [38], absolute size and the relationship

between CS_{body} and CS_{head} as well as their relationship with each bone (CS_{bone}) were examined. We tested for

differences in size and allometry of size measures between morphs using Procrustes ANCOVA using the function

procD.lm. As salmonid morphology is influenced by sex and age [56, 57], we investigated the influence of these

variables on morph mean and allometry for size and shape. We then used pairwise comparisons using the *pairwise*

function to test for significant differences between groups. External morphology (whole-body and head shape) was

then examined using Procrustes ANOVA to characterize morph differences.

We tested for bone shape differences among morphs and differences in allometric relationships of bone shape and

bone size across morphs using Procrustes ANCOVA, using *procD.lm*, with CS_{bone}, morph and their interaction as

independent variables. A null (shape = CS_{bone}), reduced model (shape = CS_{bone} + morph) and full model (shape =

CS_{bone} x morph) were compared to determine best fit. When the full model had the best fit, differences in allometric

trajectories were investigated using *pairwise* with CS_{bone} as a covariate. We tested for significant differences in both

vector angles (vector correlation) and vector lengths (rate of shape change per unit covariate), considering

significance among both tests as robust support for biological differences. For all tests $\alpha = 0.008$ was chosen, using

the Bonferroni method as a conservative correction for multiple comparisons (six bone datasets). Additionally, we

considered comparisons with Z scores <3 as poorly supported, and interpreted results below that as not biologically

meaningful and chose the reduced model for further interpretation. Differences in allometry among morphs were

investigated using *plotAllometry* and visualized using predicted values from regression analyses. Principal

Component Analysis (PCA, using *plotTangentSpace*) was used to visually represent shape variation among

individuals and morphs. For bones that did not show significant differences in allometry among morphs, residuals

from a regression of shape on size (size-corrected shape variables) via *procD.lm* were used for visualization. Shape

variation along PC axes was summarized with deformation grids and vector plots from *shape.predictor*.

To analyze integration among bones, pairs of bones were compared using two-block partial least squares (PLS)

analysis with the function *two.b.pls*. We compared elements within and among bones of the upper and lower jaws.

The effect sizes were compared to determine if there was significant variation in levels of integration across

different bone pairs or morphs using *compare.pls*. For all analyses we used randomized residuals in 1000

permutation procedures to assess statistical significance and estimate the effect sizes (*RRPP* implemented in

geomorph, [58]. For integration tests $\alpha = 0.003$ was chosen as a conservative threshold (Bonferroni correction for 15

pairwise comparisons among bones). Figures were generated using R and formatted using Inkscape.

Results

Differences in body size, sex and age of sympatric morphs

To capture the morphological diversity in the four sympatric morphs from Lake Þingvallavatn we studied 240

sexually mature individuals. As was expected, morphs differed in size (fork length FL, F = 350.94, p << 0.001), log

weight (F = 381.28, p << 0.001) and age (F = 56.75, p << 0.001, S4, S5 and S6 Appendix). SB- and PL-charr were

smaller at sexual maturity than the LB- and PI-charr. The weight was influenced by sex (p < 0.001), and there was a

significant sex by morph interaction (p = 0.009). The external whole-body (p = 0.001) and head (p = 0.001) shape

differed by morph, with pelagic morphs having a more fusiform body shape and a more pointed snout (S7

Appendix). There was significant sexual dimorphism in whole-body shape ($R^2 = 0.04$, p = 0.001, S8 Appendix),

with males having a more humped back compared to females (not a craniofacial trait). These results corroborate

morphological descriptions of the morphs [45] and known sexual dimorphism [57]. There was weak sexual

dimorphism ($R^2 = 0.01$, p = 0.007) for shape of head and all six bones ($R^2 \leq 0.02$, S8 Appendix) that will not be

discussed further. Age was strongly correlated with FL ($R^2 = 0.7$), but always explained a lower proportion of

variation than FL (for external shape, bone size and shape variation). Therefore, FL was used as a covariate in

allometry analyses of bone size relative to body size.

Allometry of bone size and shape

Examination showed that all six bone landmarks sets were highly repeatable (S9, S10 and S11 Appendix), the

(Line number shifting)

association between replicates was high (r-PLS ~ 0.84 - 0.99) and variation due to measurement error was low

relative to variation among individuals (R^2 = 0.002-0.005, vs. R^2 = 0.89-0.94, S11 Appendix). Thus, the landmarks

developed here for shape of these six bones in salmonids appear to be quite robust.

Predictably, the size of all bones (CS_{bone}) increased with body length ($\log_e$ FL, $R^2 \geq 0.94$, p = 0.001, Fig 2

and S6 and S12 Appendix). Morph and morph by size interaction, while significant, accounted for a small fraction

of the variation ($R^2 \leq 0.04$, S6 Appendix). Additional testing revealed small differences in bone size allometry

between pairs of morphs (S13 and S14 Appendix). These differences mostly reflected differences between larger

(PI-charr) and smaller (SB-, PL-charr) morphs. PI-charr have significantly larger dentary bones than all other

morphs (p = 0.001), as well as significantly larger articular-angular, maxilla, and supramaxilla bones than both LB-

and PL-charr (p < 0.002). PL-charr also have a significantly smaller quadrate, premaxilla and supramaxilla

compared to SB-charr (p = 0.001). As CS_{bone} was highly correlated with body size (CS_{body} and $\log_e$ FL) and a

common allometric relationship was found among morphs, we used CS_{bone} as size measures in subsequent analyses.

Fig 2. Size and shape allometry of dentary, articular-angular premaxilla and maxilla (from top to bottom, inset pictures of each bone). Left panels: relationships between bone size and fork length by morph; Right panels: relationships between bone shape and bone size. Shown are values for individuals (open circles) and the predicted values (filled) of regressions for both bone size (CS_{bone}) vs body size ($\log_e$ FL) or bone-shape versus bone-size. $\log_e$, natural log transformation.

For all bones, shape was significantly affected by size (CS_{bone}), the allometric relationship explained 20-30% of shape variation for dentary, articular-angular, quadrate, and maxilla (R^2 = 0.2-0.3, Table 1, for premaxilla and supramaxilla, bone size only explained 10-15%). Note, for the dentary and articular-angular morph explained a

similar amount of the shape variation as size (Table 1 and S15 Appendix). The allometry of all bones varied by

morph (p = 0.001, Table 1), except for the quadrate, however the effects were small for maxilla and supramaxilla.

Further comparisons of vector correlation angles and lengths indicated that allometric relationships vary only

markedly by morph in the dentary (S16, S17 and S18 Appendix). For this bone, while the common allometry and

morph effects on shape were much larger than the interaction, there was significant support for allometric

differences among several morph pairs (S16 Appendix). Specifically, the PI-charr differed significantly in allometric

trajectory orientation from all other morphs (p ≥ 0.003), converging on a similar shape as LB-charr at larger size.

While the allometric trajectories of LB- and SB-charr had similar lengths, the rate of shape change was higher in

LB- compared to SB-charr (p = 0.001). Therefore, we present below the mean shape differences among morphs

from the full model (CS_{bone} X morph) for the dentary but for all other bones results from the reduced model (CS_{bone} +

morph), as both common allometry and morph affected shape variation significantly (p = 0.001). All visualizations,

unless otherwise noted, have been corrected for size to remove the effects of allometry on shape. We focus first on

the lower jaw bones and then the upper jaw.

219 **Table 1. The influence of size (bone centroid size) and morph on shape variation in 6 jaw bones**. Analysed with
220 Procrustes ANCOVA, testing for main and interaction terms. For all bones, except quadrate, was the full model with
221 interaction the best fit.

Bone Shape	Log$_e$ CS Effect				Morph Effect				Morph x Log$_e$ CS Interaction Effect			
	R^2	F	Z	P	R^2	F	Z	P	R^2	F	Z	P
Dentary	0.237	107.58	5.29	**0.001**	0.228	34.49	6.35	**0.001**	0.026	3.89	5.09	**0.001***
Articular-angular	0.190	72.95	5.26	**0.001**	0.184	23.43	7.43	**0.001**	0.020	2.61	3.56	**0.002**
Quadrate	0.298	102.95	6.32	**0.001**	0.020	2.35	2.98	**0.002**	-	-	-	-
Premaxilla	0.165	55.15	9.14	**0.001**	0.110	12.27	9.53	**0.001**	0.030	3.33	5.19	**0.001**
Maxilla	0.199	67.56	7.75	**0.001**	0.096	10.87	7.54	**0.001**	0.022	2.53	2.83	**0.001**
Supramaxilla	0.099	27.66	6.27	**0.001**	0.074	6.90	5.29	**0.001**	0.027	2.55	2.75	**0.003**

222 *Pairwise tests (S17 and S18 Appendix) indicated only the dentary had true differences in allometry by morph.

223 R^2 = coefficient of determination*; F* = F-statistic; Z = Z-statistic; P = P-values, ≤ 0.008 in bold (significant after
224 Bonferroni correction).

Benthic and pelagic divergence in shape of specific lower jaw bones

For the dentary and articular-angular, the morph effect on shape variation was similar in magnitude as the common

allometric effect (20-30% of shape variation, Table 1). For the other bones the size (allometry) was considerably

stronger.

Most dramatically, pairwise tests revealed differences in dentary shape between all morph pairs across the benthic-

pelagic axis, except among the two smaller morphs (SB- and PL-charr, p = 0.119, Table 2). Also, the comparison

PI-SB-charr was weakly supported (Z < 3). No shape differences were within morphotype (LB-SB-charr, PL-PI-

charr). The dentary shape differences by morph were visualized using PCA (Fig 3), and due to variation in allometry

among morphs, size effects were not removed. The morphs separated along a benthic-pelagic axis on PC1 (50% of

the total variation) with pelagic morphs, on average, having an elongated dentary, increased length of the dental

palate, more acute angle between ridges, and vertically inclined lingual palate relative to benthic morphs, which

aligns with the hypothesis of shorter lower jaw in benthic charr. PC1 represents elongation of the dentary bone, with

positive values representing a more elongate shape, with extension of the dental palate containing the teeth, and

negative PC1 values representing a compact shape with reduction in the length of the dental palate. Positive values

were also associated with a more acute angle between the dorsal and central ridges, reducing the size of the

Meckelian notch, as well as a more vertical inclination of the lingual palate. PC2 explained 13% of the total

variation, and mainly split the morphs by size the larger morphs (LB-, PI-charr) having positive values and taller

bones. The associated shape changes mainly affected the height of the whole bone, with negative PC values

associated with thinner bones overall, reduced height of the mandibular symphysis, coronoid process, and ventral

shelf. Together, PC1 and PC2 separated all morphs clearly in shape space, but they overlap (possibly due to

overlapping sizes of particular individuals and/or more size variation within LB-charr).

Table 2. Morph differences in bone shape estimated by pair-wise distances in mean shape for morphs, from
the best fit ANCOVA models for each bone (Table 1).

Bones	Morph pairs	d	UCL (95%)	Z	P-value
Dentary	LB-PI	0.176	0.129	3.91	**0.001**
	SB-PL	0.106	0.110	1.24	0.119
	LB-PL	0.154	0.131	3.15	**0.001**
	PI-SB	0.131	0.117	2.65	**0.005**
	LB-SB	0.070	0.072	1.47	0.070
	PI-PL	0.042	0.051	0.60	0.270
Articular-angular	LB-PI	0.088	0.023	6.77	**0.001**
	SB-PL	0.063	0.022	4.43	**0.001**
	LB-PL	0.104	0.035	5.12	**0.001**
	PI-SB	0.059	0.038	3.04	**0.001**
	LB-SB	0.055	0.039	2.72	**0.005**

	PI-PL	0.038	0.035	1.98	0.021
Quadrate	LB-PI	0.024	0.025	1.54	0.065
	SB-PL	0.022	0.021	1.73	0.045
	LB-PL	0.037	0.037	1.58	0.050
	PI-SB	0.048	0.038	2.65	**0.006**
	LB-SB	0.038	0.040	1.43	0.080
	PI-PL	0.046	0.035	2.91	**0.004**
Premaxilla	LB-PI	0.094	0.036	5.62	**0.001**
	SB-PL	0.087	0.030	6.30	**0.001**
	LB-PL	0.086	0.054	3.70	**0.001**
	PI-SB	0.098	0.053	4.56	**0.001**
	LB-SB	0.034	0.054	-0.38	0.653
	PI-PL	0.049	0.053	1.31	0.090
Maxilla	LB-PI	0.024	0.019	2.35	0.010
	SB-PL	0.051	0.017	5.93	**0.001**
	LB-PL	0.053	0.028	4.06	**0.001**
	PI-SB	0.048	0.031	3.28	**0.001**
	LB-SB	0.033	0.032	1.83	0.037
	PI-PL	0.050	0.028	4.04	**0.001**
Supramaxilla	LB-PI	0.022	0.038	0.002	0.490
	SB-PL	0.083	0.033	4.60	**0.001**
	LB-PL	0.061	0.056	1.89	0.028
	PI-SB	0.102	0.061	3.41	**0.001**
	LB-SB	0.096	0.059	3.37	**0.001**
	PI-PL	0.051	0.056	1.33	0.092

249   UCL = upper confidence level; Z = Z-statistic; P-values ≤ 0.008 in bold (Bonferroni correction)

251
Fig 3. Strong morph effects on shape variation in the lower jaw bones, dentary (top) and articular angular (bottom). For the dentary, size effects were not removed, but effects of size were removed for articular angular (and other bones, see below). PC-plots of specimens and deformation grids show how shape differences by morphs. None of the shape warps (X- and Y-axis) are magnified. Each dot represents an individual and the ellipses 95% CI for the distribution by morph (large dot represents the mean for each morph in these dimensions). PC1 and 2 explain 50% and 13% of the variation in dentary, respectively. For the articular angular (bottom) PC1 and 2 explain 37% and 13%.

Pairwise tests for articular-angular shape also revealed differences between all morph pairs across the benthic-

pelagic axis. Biologically meaningful differences were not found within morphotypes (Z<3, see Table 2). The

morphs aligned along a benthic-pelagic axis on PC1 which explained 37% of the variation. The two pelagic morphs

were largely overlapping (positive values), with elongated anterior process of the angular "wing", shorter coronoid

process, and narrower quadrate facet. The benthic morphs had more shape variation (especially the LB-charr, as in

dentary) but had a more compact bone shape and wider quadrate facet overall, with LB-charr individuals being most

compact (Fig 3). PC1 represents changes in relative sizes of the anterior process of the angular 'wing', retroarticular,

and quadrate facet. Individuals with positive PC1 values show anterior extension of the anterior process of the

angular "wing", increase in the size of the anterior process of the retroarticular but dorso-ventral compression of the

retroarticular, and narrow quadrate facet. Negative values associate with a more compact shape overall, particularly

with restrictions to the length of the anterior process of the angular "wing", however the coronoid process is

extended anteriorly, a more square retroarticular, and wider quadrate facet. PC2 explains 13% of the variation and

mainly represents differences in the ventral process of the retroarticular, with individuals with negative PC values

with a shorter process. The morphs did not separate along PC2.

In sum, the morph differences in the articular-angular and in particular the dentary were large. In contrast, for the

last lower jaw bone the quadrate morph had weak effects on shape ($R^2 = 0.02$, $Z < 3$, Table 1). PCA and pairwise

tests for mean quadrate shape differences did not reveal shape divergence among morphs (Table 2 and S19

Appendix).

Weaker divergence in shape of upper jaw bones

For the upper jaw bones the magnitude of the morph and size effects on shape were similar (~10-15%, Table 1) but

notably lower than in the dentary and articular-angular. Morph had strongest effect on the premaxilla and weakest

(nearly none) on the supramaxilla. The analyses were done on size corrected data, assuming common allometry.

Pairwise tests for mean premaxilla shape differences among morphs indicated significant differences in all morph

pairs across the benthic-pelagic axis ($p \leq 0.001$, Table 2). No differences were found within morphotype. The

morphs separated along a benthic-pelagic axis on PC1 (25% of the variation). The PL- and PI-charr had positive

PC1 values with elongated and narrow limbs, while SB- and LB-charr overlapped with more negative PC1 values,

having compact and wide limbs (Fig 4). PC1 captures morphological shape differences of the posterior and anterior

limbs. Positive values associated with elongated and narrow posterior and anterior limbs, while negative values with

more compressed and wider posterior and anterior limbs. PC2 explained 14% of the variation, representing variation

due to differences in the curvature of ascending limb but did not separate morphs.

For maxilla shape, pairwise tests of mean shape between morphs revealed significant separation between all morphs

($p = 0.001$), except the SB-LB-charr and LB-PI-charr (Table 2). Therefore, maxilla diverged neither exclusively

along the benthic-pelagic axis nor the size gradient. The morphs separated on the benthic-pelagic axis on PC3 (10%

of variation). This component represents the height of the maxilla (positive: tall and negative: short) aligning with

our hypothesis, but the distributions overlapped substantially (Fig 4). PC1 explained 33% of the total variation, but it

represents technical variation (either due to rotation or tilting of the bone during photographing, that was randomly

distributed with respect to morph). PC2 (23% of total variation), representing biological variation due to differences

in the curvature of the maxillary head and angle of the maxillary neck, and separated the PL slightly from other

morphs. In combination, PC2 and PC3 indicate that PL-charr has reduced morphological variation and appears to

have a thinner maxillary body and caudal lobe than the other morphs.

Pairwise comparisons of mean supramaxilla shape differences among morphs indicated SB deviated from all other

morphs ($p = 0.001$, Table 2). PC1 and PC2 explained 62% of variation, but no combination of PCs revealed clearly

how SB-charr shape deviated (S19 Appendix). We note the morphological variation in SB appeared reduced relative

to other morphs.

Fig 4. Most pronounced divergence in premaxilla of the upper jaw, and subtle changes in maxilla shape. Size corrected PC-plots of specimens and deformation grids showing how the morphs differ in mean (top) premaxilla and (bottom) maxilla shape. None of the warps (X- and Y-axis) are magnified, representation of PC plots and warps as in Fig 3. For premaxilla (top) PC1 and PC2 explain 25% and 14% of the variation respectively. For maxilla (bottom) PC2 and PC3 explain 23% and 10% respectively (PC1 was biased by sampling error, and not depicted).

Covariation in shape change among bones

Lastly, we tested for covariation in shape among bones. Within the upper jaw and within the lower jaw, shape

covaried significantly among all bones (p = 0.001). The degree of covariation was greatest for the dentary and

articular-angular (r-PLS = 0.92; Z = 7.08, S20 Appendix) and much higher compared to the other pairs (r-PLS =

0.54-0.69). Among the jaws, all covariation among bones was significant (p = 0.001) and the degree of covariation

was moderate (r-PLS ~ 0.54-0.78, S20 Appendix). We did not find significant differences in effect size among pairs

in the lower jaw, however in the upper jaw integration between the premaxilla and supramaxilla was found to be

lower (p = 0.006) than all other comparisons, and integration between the maxilla and supramaxilla was found to be

higher (p = 0.002) (S20 Appendix). In addition, analyses did not reveal significant differences in effect size between

morphs for any of the bone pairs (comparisons limited to bones that showed significant covariation or biological

importance as described above, quadrate and supramaxilla excluded, data not shown).

Discussion

Evolutionary divergence along the benthic-pelagic axis occurs repeatedly among fishes [59-62]. Body elongation is

a consistent trend along the benthic-pelagic axis in marine and freshwater fishes [59] as well as fin adaptations for

sustained swimming in pelagic environments and manoeuvrability in benthic habitat [60, 63]. Craniofacial responses

to benthic and pelagic prey resources are quite complex [4, 64-66]. Prey specialization along this axis leads to

morphological adaptation of multiple traits that can be either integrated or modular, with known examples of

functional convergence via a range of morphological solutions [67]. Salmonids are known for extensive variation in

feeding specializations and morphology but to date few studies have investigated variation in internal structures

among or within a salmonid species [except see, 18, 24, 25, 47, 48, 68, 69]. Notably, in addition to pioneering work

of Ingimarsson (2002), one study reported differences in dentary and articular-angular shape in parr, between several

salmonid species and also within rainbow trout (*Oncorhynchus mykiss,* [18]).

We report morph differences in body, head and bone shape, with bones showing more pronounced differences. The

whole-body and head shape differences by morph are consistent with previous studies [45, 57]. We examined six

cranial bones tied directly to feeding and prey capture in recently evolved sympatric morphs of arctic charr, and the

results corroborate our hypothesis (1), of clear allometric shape variation in all six bones. Surprisingly, shape

allometry did differ between morphs for the dentary. In accordance with hypothesis (2) we found the strongest

benthic-pelagic contrast in the most anterior bones in lower jaw bone (dentary). We did not expect premaxilla

divergence however. These bones interact directly with prey, but shape differences were also seen in the articular-

angular and maxilla. Shape differences in dentary and maxilla are consistent with hypothesis (3) (shorter lower jaw

and more round maxilla in benthics). Also, curiously, for both the dentary and articular-angular the morph effects

(benthic-pelagic axis, on PC1 for both) were stronger than the effects of morph size (PC2 for both). In contrast, the

shape of the two posterior bones, quadrate and supramaxilla, did not vary strongly between morphs. Following

hypothesis (4), shape variation in bones were highly correlated with each other, with bones of the same jaw more

integrated than bones among jaws. We discuss these results in the context of functional feeding morphology and its

mechanics, the putative genetic and environmental sources of these traits and suggest future studies of craniofacial

diversity in salmonids. Here we focus on cranial bone shape and potential relationships to prey and mechanics of

prey capture, but acknowledge that postcranial divergence allows access to preferred prey habitat and also aids in

locating, approaching, and capturing individual prey [4, 60].

Bone shape variation and functional divergence

We found strong divergence in the lower jaw along the benthic-pelagic axis. The characteristic subterminal mouth of

the benthic morphs associates with a shortening of the dentary and to lesser extent articular-angular (relative to the

quadrate facet) aligns with our hypothesis (3). Additionally, the anterior region of the dentary around the mandibular

symphysis is angled ventrally relative to pelagic charr (Fig 3). This may aid in creating a more optimal angle to grip

attached prey or perform scraping behaviours on a surface to dislodge prey, such as snails. The benthic morphs may

use a combination of biting and ramming behaviours, as suction forces are likely weak. Taller dentary and articular-

angular bones could be adaptations for feeding on benthic prey, increasing size of the mouth cavity and efficiency of

consuming benthic organisms [70]. The coronoid processes of the dentary and articular-angular are enlarged in

benthic charr, which could associate with a larger mandibular portion of the adductor mandibulae and a larger

aponeurosis maxillaris [71], thus allowing for greater bite forces in benthic morphs which could aid in dislodging

attached prey items [64, 72, 73]. Relatively more elongated lower jaw was expected and observed in pelagic morphs.

Consistently, previous study on lower jaw bone shape in juveniles of *O. mykiss* found the anadromous population

(with presumed pelagic lifestyle) had a more elongated and streamlined lower jaw than the resident population [18].

The authors postulated that elongated, streamlined lower jaws could be adaptations to the pelagic ocean

environment.

In the upper jaw, we found premaxilla shape to diverge along the benthic-pelagic axis more strongly than

the maxilla. PL-charr had smaller premaxilla overall (importantly relative to SB which is of same size), with shorter

ascending limbs and narrow anterior and posterior limbs. Reductions in the height of the ascending process and

elongation of the anterior-posterior axis of the premaxilla has been found in other pelagic fishes [65, 74, 75] that use

ram feeding in the water column [76] while suction feeders with highly protrusible jaws have long ascending limbs

[77]. Moreover, shortening and steeper angles of the ascending limb of the premaxilla can also be associated with

increased biting forces [78] and possibly linked to biting behaviours. The ascending limb of the premaxilla has been

found to extend more dorsally in pelagic fish [79] and in highly protrusible jaws [80], however, in our study the

ascending limb was proportionally longer in the benthic morphs. As the premaxilla, and the skull as a whole, is

relatively rigid in salmonids [sensu 71], the biomechanics of similar morphological changes in fishes with

protrusible jaws could be incomparable. For the maxilla, the third shape axis associated with the benthic-pelagic

axis. The maxilla of fishes is generally involved in the protrusion of the upper jaw maxillary apparatus, while

moveable in salmonids such function may not be present [71] and the proposed movement of these components to

aid in salmonid feeding [81] has been criticized [71]. The shape change seen in the bone matches hypothesis 2,

based on observations done on external photographs in pelagic charr [50].

Neither the quadrate nor supramaxilla diverged along a benthic-pelagic axis (but showed some weak morph

differences). While shape variation in them could reflect biomechanical differences due to size [82, 83], they did not

vary along a morph size gradient. In salmonids, the function of the supramaxilla (if any) is unknown. The morph

differences in supramaxilla may reflect correlations with linked bones (like the maxilla), as the high level of

integration between these bones indicates. In contrast, the quadrate is a critical linkage in the suspensorium, its main

function to allow rotation of the lower jaw. Since changes to the maximum angle of opening potentially needed to

capture different prey would likely only require subtle morphological changes in this bone, it is unlikely we could

detect these differences given our methods, or divergent selection may not have acted strongly on quadrate shape.

While adult morphology should reflect long-term prey consumption (e.g., higher pelagic prey consumption

associates with stronger pelagic features), variation in the mechanisms of capturing and processing of different prey

items within and between sympatric charr morphs, and potential impacts on morphological variation, is poorly

understood. To our knowledge biomechanical studies of salmonids have not yet been done on fishes that specialize

on such divergent prey as snails and zooplankton. Feeding was compared in PL- and SB-charr, revealing reduced

efficiency of zooplankton feeding (pelagic) in SB-charr juveniles due to lower attack rates [41], increased handling

time [84], increased ejection of prey [84] and poorer retention in the branchial apparatus likely due to shorter gill

rakers [41]. Both morphs likely capture zooplankton using a similar strategy, i.e., suction [41], however may exhibit

differences in capture efficiency as a result of morphological differentiation associated with ecological divergence

[41]. Prior studies are limited to biomechanics of piscivorous feeding in brook trout (*Salvelinus fontinalis*) [85] or

theoretical models of suspensorium and lower jaw movement [86]. Salmonid jaw elements are likely to be largely

immobile due to rigid attachments of the components [71]. Therefore, the use and efficiency of suction is likely

limited compared to pelagic fishes with highly mobile jaws [87], and perhaps not highly differentiated among arctic

charr morphs.

405   Benthic-pelagic divergence can manifests in lower jaw length relative to head length [66], as seen in

subterminal mouths of benthic arctic charr [38, 45, 88]. Shelled freshwater snail (*Lymnaea peregra*) is the main prey

of the derived SB- and LB-charr in Þingvallavatn [38, 41]. In general, hypertrophied pharyngeal jaws [89-93],

molariform pharyngeal teeth [94, 95], and decreased gill raker length and density [41, 45, 96, 97] are all

morphological adaptations widely employed across fishes that consume large quantities of hard-shelled benthic

prey. Also, mechanisms for capturing attached prey vary widely among species [64, 72, 73]. Use of strong suction,

gripping and biting, lateral movements, and highly protrusible jaws are commonly utilized traits [64, 72, 73].

However, details of the intersection of these biomechanical and behavioural components for feeding on divergent

prey resources in Þingvallavatn morphs is currently unexplored. In many groups of fishes, strong benthic-pelagic

ecological divergence has led to modification in morphology and behaviour to allow for optimal feeding

performance [17, 59, 60, 64, 66, 75, 77, 79, 98-101]. Clearly, investigation of the behavioural and biomechanical

strategies individuals and morphs use to both capture and process prey, and bone shape in the same individuals, is

needed to reveal how arctic charr specialize on such divergent prey types.

418   While we expected adaptions to divergent prey resources to be expressed in shape divergence of cranial

elements among morphs, other morphological modifications can occur that do not impact bone shape as studied

here. Internal bone size and architecture has been shown to diverge along the benthic-pelagic axis, with benthic

feeders exhibiting more robust hollow cranial bones with complex internal structure, similar to mammalian long-

bones, and planktivores having thin solid bones [102]. Bones of benthic scrapers also have higher rates of

mineralization [103] and are more resistant to bending [102]. We found benthic morphs (LB- and SB-charr) to have

thicker and more robust bones overall (relative to PL- and PI-charr), but variation in their internal structure and

mineralization is unknown. The differences in adult bone size and shape found here may associate with

differentiation in muscle mass and surface area of attachment sites, and as skeletal and musculature may contribute

independently to biomechanical diversity [104], adaptive changes to soft tissues may both be more critical in

foraging divergence (i.e. proportionally larger impact on biomechanics of feeding related to speed, force, and mode

of prey capture) and be more apparent than associated changes in bone shape in these morphs.

431 Influence of extrinsic and intrinsic factors on bone variation within

432 and among morphs

The observed shape variation in the jaw bones within and between morphs could be influenced by differences in

their ecology and food preferences [see above, 41, 42], genetic composition of the morphs [37, 105] and/or

interactions of specific genes and environmental factors.

Adaptive plasticity may contribute to ecological and morphological divergence among sympatric and

allopatric charr [22, 106, 107]. We studied wild caught individuals of the four morphs that mostly operate in

different habitats and feed on different prey items that vary seasonally in abundance and nutrition, all factors that

may influence their development and adult morphology [41, 42, 46]. Because the morphs differ in the location,

timing and synchronicity of spawning [44], their juveniles may receive different maternal provisions and encounter

different juvenile habitats. Arctic charr shows developmental plasticity in body shape and size in response to benthic

vs. pelagic diet treatment in experimental settings [106, 108]. Behavioural traits related to habitat choice, prey

capture and processing could also influence variation in bone traits [98]. For example, in sheepshead wrasse

(*Archosargus probatocephalus*), individuals consuming a higher proportion of hard-shelled prey had larger adductor

mandibulae and jaw bones but also employed biting mode of feeding to dislodge prey as well as buccal

manipulations to crush shells and separate hard and soft parts before ingestion [109]. Thus, some of the

morphological differences between (and/or within) morphs revealed here could reflect this developmental plasticity.

Note however, not all environmentally induced variation in bone shape (plastic response) may be adaptive [110].

Genetic differences by groups can be assessed with experiments and population genetics. Common garden

rearing of offspring of Þingvallavatn morphs indicate genetic influence on many aspects of juvenile and adult

morphology [47, 50, 57, 68, 111], though maternal and/or transgenerational parental effects could also play a role

[112]. Three of the morphs (PL-, SB-, and LB-charr) are genetically separable [37] and genomic data revealed

substantial allele frequency differences between them, on nearly all linkage groups [36, 40]. We saw the strongest

morph differences in the dentary and premaxilla bones of the lower and upper jaws, respectively, that may reflect

alleles influencing these bones specifically. Models indicate these three morphs have been separate for thousands of

generations, with limited gene flow between them [36, 37]. Curiously, the PI-charr is genetically more

heterogeneous, most individuals are genetically similar to PL-charr, but others have either LB-charr or mixed PL-

/LB-charr ancestry [40]. One hypothesis is that PI-charr result from ontogenetic niche shifts of juvenile PL-charr

that learn to catch sticklebacks, abandoning zooplankton feeding, a radical change which may allow individuals to

attain larger size and develop piscivorous characteristics [45]. Another suggestion [40] is that juvenile LB-charr

and/or LB/PI hybrids also shift to piscivory. For most bones studied here, the PI-charr tended to align with PL-charr

(consistent with ancestry), except in the maxilla where PI clustered closer to LB-charr. This heterogeneity of

phenotypes in PI-charr could either reflect variation in genes, or the impact of piscivory on development of specific

bones. Piscivory and associated size increase may place high functional demands on bone shape [113] and enhance

divergence in shape and even suppress morphological variation among piscivorous charr of different genetic origins.

As mapping can identify loci, genes or specific polymorphisms impacting variation in shape [114-116], QTL or

association studies of bone shape variation in PI-, LB- and PL-charr could identify alleles and pathways of relevance

for bone shape variation in salmonids.

Most likely, interactions between particular alleles and specific environmental factors, acting at specific

times or over broader developmental windows [117], produce the observed morph differences in bone shape.

Considering a hypothetical case, SB-charr may have alleles that influence the development of the dentary and

articular-angular (lower jaw bones), that will induce the subterminal morphology if they eat snails as juveniles. Tests

of such hypotheses require joint study of genetic and environmental influences on specific aspects of how bones

develop and acquire their shape and functions. To distinguish between genetic and transgenerational environmental

effects, mapping, crosses between morphs and multigenerational studies are needed. One could test the sensitivity of

shape-alleles to environmental factors, like specific juvenile food types (e.g., benthic or pelagic). Analyses of gene

expression can also reveal developmental pathways that respond differently to environment by morph. Already

RNA-, methylation- and miRNA-sequencing of embryos of these morphs and aquaculture charr have demonstrated

differences by morphs [118, 119]. For instance, in PL-, LB- and SB-charr reared in a common garden ~2000 genes

were differentially expressed by morphs during development (from formation of the gill-arches to establishment of

the viscerocranium primordia) [105]. This proves the morphs differ in genes influencing multiple developmental

systems that could influence these bone shape differences. Specifically, as common garden experiments revealed

ossification of skull elements in juveniles starts earlier in SB- compared to PL-charr [111], it would be interesting to

explore effects of genes and environmental factors on ossification and bone shape. Most probably interactions of

specific environmental attributes and genetic composition cause the observed morph differences in bone shape, for

instance via differences in patterns of growth or differentiation in the bone primordia or linked tissues.

Future avenues in research of bone shape variation in salmonids

We studied variation in six dentary bones in sexually mature fish of derived morphs from one lake. This study could

be expanded in several ways. First, size, age and sex can influence traits [38, 120]. We studied sexually mature

individuals, and average age and size varied quite substantially by morph. It would be interesting to study the

emergence of bone shape variation during ontogeny in these morphs, by sampling age classes and genotyping

individuals (to assign them to morph). For bone shape, size (fork length) had stronger effect on shape than age. Also,

despite significant difference among morphs in allometries (age-size relationship as well as between size measures

of head, body, bones), these effects were rather small, and we concluded bone size was a robust proxy for overall

size (and was used to account for size variation). But with larger samples it might be possible to disentangle and

better estimate these effects. The same applies to potential sexual dimorphism or morph specific sex-dimorphism in

bone size or shape allometry. Secondly, as four derived morphs were studied, the phenotypes could not be polarized

with respect to an outgroup. Studies of anadromous charr could identify ancestral vs. derived character states. Arctic

charr in Iceland have repeatedly evolved along a benthic-pelagic axis and subsequently experienced modification to

allow for capture and processing of diverse prey items (e.g., benthic macroinvertebrates, fish, snails) [32, 33, 36,

88]. It would be interesting to study the degree of parallelism in bone shape variation and integration in more

sympatric benthic-pelagic morphs or the miniaturized SB-charr. Thirdly, the salmonid head contains several dozens

of bones and the cranial case. Ingimarsson (48) found a signal (but weak) of morph differences in linear measures of

several other bones, but geometric morphometrics on these and other bones (in 2D or 3D from CT-scans) may reveal

this more systemically and highlight potentially adaptive changes [55, 121, 122]. Analyses of associated variation in

soft tissues (muscles and ligaments), cartilage and ossification would also be interesting. We found rather high

integration in shape and size variation among bones within and between the upper and lower jaws. However,

covariation of shape and univariate traits (like teeth numbers) or variation in cranial kinesis, how the bones lie

together in the skull and how they move together [85], was not analysed here. The feeding structures of fishes are

highly functionally integrated [4, 87]. Study on the seven dolly varden trout (*Salvelinus malma*) morphs in Lake

Kronotskoe indicated morph specific combinations of morphological traits and differences in benthic-pelagic

ecologies manifest in mouth position and jaw length divergence early in ontogeny [25]. Three cranial bone sets had

with coordinated shifts in ossification: jaws (premaxilla, maxilla and dentary), intraoral tongue-bite apparatus

(vomer and lingual bone, gill arch elements) and skull dermal bones (supraethmoid, frontal and preopercle), most

likely reflecting differences in early ossification [25]. Studies on additional internal skull elements and their

integration are needed to understand the functional capacity of different jaw elements and how they relate to the

remarkable feeding divergence in arctic charr.

520 Conclusions

The diversity of feeding morphology in fishes provides opportunities to examine variation in functional elements, to

determine rates of evolution, integration among morphological units, and how specialization occurs along a benthic-

pelagic axis. We gathered unique data on intraspecific variation in feeding related bones. The results indicate a clear

benthic-pelagic separation that was most prominent in the dentary bone. Further work is needed to understand the

causes of variation in feeding elements within species, how it evolves and develops. Are the axes of jaw bone shape

variation in salmonids shared with other fishes or even other vertebrates? Work along these lines will aid in our

understanding of the processes that still generate the great biodiversity we observe and must protect.

Acknowledgements

We thank Haraldur R. Ingvason and Stefán Már Stefánsson of the now closed Kópavogur Nature Center, and Fia

Finn, Zophonías O. Jónsson and Grégoire Gaillet at the University of Iceland for help with sampling. We thank

Benóný Jónsson and coworkers at the Marine & Freshwater Research Institute of Iceland for providing fish and

data. Members of the Arctic charr-group at the University of Iceland deserve many thanks for help during the

project, discussions, constructive criticism and great social environment. We thank the Institute of Biology at the

University of Iceland for financial support.

References:

1.   Albert AY, Schluter D. Selection and the origin of species. Curr Biol. 2005;15(8):R283-8. Epub 2005/04/28. doi: 10.1016/j.cub.2005.04.015. PubMed PMID: 15854890.

2.   Pigliucci M. Is evolvability evolvable? Nature Reviews Genetics. 2008;9(1):75-82. doi: 10.1038/nrg2278.

3.   Schluter D, Conte GL. Genetics and ecological speciation. Proc Natl Acad Sci U S A. 2009;106 Suppl 1(Suppl 1):9955-62. Epub 2009/06/17. doi: 10.1073/pnas.0901264106. PubMed PMID: 19528639; PubMed Central PMCID: PMCPMC2702799.

4.   Gidmark NJ, Pos K, Matheson B, Ponce E, Westneat MW. Functional Morphology and Biomechanics of Feeding in Fishes. In: Bels V, Whishaw IQ, editors. Feeding in Vertebrates. Switzerland: Springer; 2019. p. 297-332.

5.   Schwenk K. Feeding: Form, Function and Evolution in Tetrapod Vertebrates: Elsevier Science; 2000.

6.   Bels VL, Chardon M, Vandewalle P. Biomechanics of Feeding in Vertebrates. In: Advances in Comparative and Environmental Physiology, vol 18. New York: Springer; 1994.

7.   Grant RB, Grant PR. What Darwin's Finches Can Teach Us about the Evolutionary Origin and Regulation of Biodiversity. BioScience. 2003;53(10):965-75. doi: 10.1641/0006-3568(2003)053[0965:Wdfctu]2.0.Co;2.

8.   Karagic N, Meyer A, Hulsey CD. Phenotypic Plasticity in Vertebrate Dentitions. Integr Comp Biol. 2020;60(3):608-18. Epub 2020/06/17. doi: 10.1093/icb/icaa077. PubMed PMID: 32544244.

9.   Huby A, Parmentier E. Actinopterygians: Head, Jaws and Muscles. In: Ziermann JM, Diaz Jr RE, Diogo R, editors. Heads, Jaws, and Muscles: Anatomical, Functional, and Developmental Diversity in Chordate Evolution. Cham: Springer International Publishing; 2019. p. 93-117.

10.   Liem KF. Ecomorphology of the Teleostean Skull. In: Hanken J, Hall BK, editors. The Skull volume 3. United States of America: The University of Chicago Press; 1993. p. 422-52.

11.   Martinez CM, McGee MD, Borstein SR, Wainwright PC. Feeding ecology underlies the evolution of cichlid jaw mobility. Evolution. 2018. Epub 2018/06/20. doi: 10.1111/evo.13518. PubMed PMID: 29920668.

12.   Cooper WJ, Westneat MW. Form and function of damselfish skulls: rapid and repeated evolution
into a limited number of trophic niches. BMC Evolutionary Biology. 2009;9(1):24. doi: 10.1186/1471-
2148-9-24.
13.   Westneat MW, Alfaro ME, Wainwright PC, Bellwood DR, Grubich JR, Fessler JL, et al. Local
phylogenetic divergence and global evolutionary convergence of skull function in reef fishes of the
family Labridae. Proc Biol Sci. 2005;272(1567):993-1000. Epub 2005/07/19. doi:
10.1098/rspb.2004.3013. PubMed PMID: 16024356; PubMed Central PMCID: PMCPMC1599880.
14.   Mabuchi K, Miya M, Azuma Y, Nishida M. Independent evolution of the specialized pharyngeal
jaw apparatus in cichlid and labrid fishes. BMC Evolutionary Biology. 2007;7(1):10. doi: 10.1186/1471-
2148-7-10.
15.   Konings AF, Wisor JM, Stauffer JR, Jr. Microcomputed tomography used to link head morphology
and observed feeding behavior in cichlids of Lake Malaŵi. Ecol Evol. 2021;11(9):4605-15. Epub
2021/05/13. doi: 10.1002/ece3.7359. PubMed PMID: 33976834; PubMed Central PMCID:
PMCPMC8093705.
16.   Pauers MJ, McMillan SA. Geometric morphometrics reveals surprising diversity in the Lake
Malawi cichlid genus Labeotropheus. Hydrobiologia. 2014;748:145-60.
17.   Ronco F, Matschiner M, Böhne A, Boila A, Büscher HH, El Taher A, et al. Drivers and dynamics of
a massive adaptive radiation in cichlid fishes. Nature. 2021;589(7840):76-81. doi: 10.1038/s41586-020-
2930-4.
18.   Kimmel CB, Watson S, Couture RB, McKibben NS, Nichols JT, Richardson SE, et al. Patterns of
variation and covariation in the shapes of mandibular bones of juvenile salmonids in the genus
Oncorhynchus. Evol Dev. 2015;17(5):302-14. Epub 2015/09/16. doi: 10.1111/ede.12135. PubMed PMID:
26372063; PubMed Central PMCID: PMCPMC4916927.
19.   Smith TB, Skúlason S. Evolutionary Significance of Resource Polymorphisms in Fishes,
Amphibians, and Birds. Annual Review of Ecology and Systematics. 1996;27(1):111-33. doi:
10.1146/annurev.ecolsys.27.1.111.
20.   Markevich G, Esin E. Evolution of the Charrs, Genus Salvelinus (Salmonidae). 2. Sympatric Inner-
lake Diversification (Ecological Peculiarities and Evolutionary Mechanisms Illustrated by Different Groups
of Fish). Journal of Ichthyology. 2018;58. doi: 10.1134/S0032945218030074.
21.   Klemetsen A, Amundsen PA, Dempson J, Jonsson B, Jonsson N, O'connell M, et al. Atlantic
salmon Salmo salar L., brown trout Salmo trutta L. and Arctic charr Salvelinus alpinus (L.): a review of
aspects of their life histories. Ecology of freshwater fish. 2003;12(1):1-59.
22.   Skulason S, Smith TB. Resource polymorphisms in vertebrates. Trends in Ecology & Evolution.
1995;10(9):366-70. doi: https://doi.org/10.1016/S0169-5347(00)89135-1.
23.   Goodrich ES. Structure and Development of Vertebrates. London: Macmillan and Co.; 1931.
24.   Muir AM, Bronte CR, Zimmerman MS, Quinlan HR, Glase JD, Krueger CC. Ecomorphological
Diversity of Lake Trout at Isle Royale, Lake Superior. Transactions of the American Fisheries Society.
2014;143(4):972-87. doi: https://doi.org/10.1080/00028487.2014.900823.
25.   Esin EV, Markevich GN, Pichugin MY. Juvenile divergence in adaptive traits among seven
sympatric fish ecomorphs arises before moving to different lacustrine habitats. J Evol Biol.
2018;31(7):1018-34. Epub 2018/04/20. doi: 10.1111/jeb.13283. PubMed PMID: 29672982.
26.   Jacobs A, Carruthers M, Eckmann R, Yohannes E, Adams CE, Behrmann-Godel J, et al. Rapid
niche expansion by selection on functional genomic variation after ecosystem recovery. Nature Ecology
& Evolution. 2019;3(1):77-86. doi: 10.1038/s41559-018-0742-9.
27.   Klemetsen A. The Charr Problem Revisited: Exceptional Phenotypic Plasticity Promotes
Ecological Speciation in Postglacial Lakes. Freshwater Reviews. 2010;3(1):49-74, 26. doi:
https://doi.org/10.1608/FRJ-3.1.3.

28.   Østbye K, Hagen Hassve M, Peris Tamayo A-M, Hagenlund M, Vogler T, Præbel K. "And if you
gaze long into an abyss, the abyss gazes also into thee": four morphs of Arctic charr adapting to a depth
gradient in Lake Tinnsjøen. Evolutionary Applications. 2020;13(6):1240-61. doi:
https://doi.org/10.1111/eva.12983.
29.   Skoglund S, Siwertsson A, Amundsen PA, Knudsen R. Morphological divergence between three
Arctic charr morphs - the significance of the deep-water environment. Ecol Evol. 2015;5(15):3114-29.
Epub 2015/09/12. doi: 10.1002/ece3.1573. PubMed PMID: 26357540; PubMed Central PMCID:
PMCPMC4559054.
30.   Alekseyev SS, Bajno R, Gordeeva NV, Reist JD, Power M, Kirillov AF, et al. Phylogeography and
sympatric differentiation of the Arctic charr Salvelinus alpinus (L.) complex in Siberia as revealed by
mtDNA sequence analysis. Journal of Fish Biology. 2009;75(2):368-92. doi:
https://doi.org/10.1111/j.1095-8649.2009.02331.x.
31.   Doenz CJ, Krähenbühl AK, Walker J, Seehausen O, Brodersen J. Ecological opportunity shapes a
large Arctic charr species radiation. Proc Biol Sci. 2019;286(1913):20191992. Epub 2019/10/24. doi:
10.1098/rspb.2019.1992. PubMed PMID: 31640512; PubMed Central PMCID: PMCPMC6834057.
32.   Gíslason D, Ferguson MM, Skúlason S, Snorrason SS. Rapid and coupled phenotypic and genetic
divergence in Icelandic Arctic char (Salvelinus alpinus). Canadian Journal of Fisheries and Aquatic
Sciences. 1999;56(12):2229-34. doi: 10.1139/f99-245.
33.   Woods PJ, Skúlason S, Snorrason SS, Kristjánsson BK, Ingimarsson F, Malmquist HJ. Variability in
the functional role of Arctic charr Salvelinus alpinus as it relates to lake ecosystem characteristics.
Environmental Biology of Fishes. 2013;96(12):1361-76. doi: 10.1007/s10641-013-0114-x.
34.   Adalsteinsson H, Jónasson PM, Rist S. Physical Characteristics of Thingvallavatn, Iceland. Oikos.
1992;64(1/2):121-35. doi: 10.2307/3545048.
35.   Saemundsson K. Geology of the Thingvallavatn Area. Oikos. 1992;64(1/2):40-68. doi:
10.2307/3545042.
36.   Brachmann MK, Parsons K, Skúlason S, Ferguson MM. The interaction of resource use and gene
flow on the phenotypic divergence of benthic and pelagic morphs of Icelandic Arctic charr (Salvelinus
alpinus). Ecology and Evolution. 2021;11(12):7315-34. doi: https://doi.org/10.1002/ece3.7563.
37.   Kapralova KH, Morrissey MB, Kristjánsson BK, Ólafsdóttir GÁ, Snorrason SS, Ferguson MM.
Evolution of adaptive diversity and genetic connectivity in Arctic charr (Salvelinus alpinus) in Iceland.
Heredity. 2011;106(3):472-87. doi: 10.1038/hdy.2010.161.
38.   Sandlund OT, Gunnarsson K, Jónasson PM, Jonsson B, Lindem T, Magnússon KP, et al. The Arctic
Charr Salvelinus alpinus in Thingvallavatn. Oikos. 1992;64(1/2):305-51. doi: 10.2307/3545056.
39.   Malmquist HJ, Snorrason SS, Skúlason S. Bleikjan í Þingvallavatni I. Fæðuhættir.
Náttúrufræðingurinn. 1985;55(4):195-217.
40.   Guðbrandsson J, Kapralova KH, Franzdóttir SR, Bergsveinsdóttir ÞM, Hafstað V, Jónsson ZO, et al.
Extensive genetic differentiation between recently evolved sympatric Arctic charr morphs. Ecology and
Evolution. 2019;9(19):10964-83. doi: https://doi.org/10.1002/ece3.5516.
41.   Malmquist HJ. Phenotype-specific feeding behaviour of two arctic charr Salvelinus alpinus
morphs. Oecologia. 1992;92(3):354-61. doi: 10.1007/BF00317461.
42.   Sandlund OT, Jonsson B, Malmquist HJ, Gydemo R, Lindem T, Skúlason S, et al. Habitat use of
arctic charrSalvelinus alpinus in Thingvallavatn, Iceland. Environmental Biology of Fishes.
1987;20(4):263-74. doi: 10.1007/BF00005297.
43.   Jonsson B, Skúlason S, Snorrason SS, Sandlund OT, Malmquist HJ, Jónasson PM, et al. Life History
Variation of Polymorphic Arctic Charr (Salvelinus alpinus) in Thingvallavatn, Iceland. Canadian Journal of
Fisheries and Aquatic Sciences. 1988;45(9):1537-47. doi: 10.1139/f88-182.

44.   Skúlason S, Snorrason SS, Noakes DLG, Ferguson MM, Malmquist HJ. Segregation in spawning
and early life history among polymorphic Arctic charr, Salvelinus alpinus, in Thingvallavatn, Iceland.
Journal of Fish Biology. 1989;35(sA):225-32. doi: https://doi.org/10.1111/j.1095-8649.1989.tb03065.x.
45.   Snorrason SS, Skúlason S, Sandlund OT, Malmquist HJ, Jonsson B, Jónasson PM. Shape
Polymorphism in Arctic charr, *Salvelinus alpinus*, in Thingvallavatn, Iceland. Physiol Ecol Japan.
1989;1:393-404.
46.   Snorrason SS, Skúlason S, Jonsson B, Malmquist HJ, Jónasson PM, Sandlund OT, et al. Trophic
specialization in Arctic charr Salvelinus alpinus (Pisces; Salmonidae): morphological divergence and
ontogenetic niche shifts. Biological Journal of the Linnean Society. 1994;52(1):1-18. doi: 10.1111/j.1095-
8312.1994.tb00975.x.
47.   Eiríksson GM, Skúlason S, Snorrason SS. Heterochrony in skeletal development and body size in
progeny of two morphs of Arctic charr from Thingvallavatn, Iceland. Journal of Fish Biology.
1999;55(sA):175-85. doi: https://doi.org/10.1111/j.1095-8649.1999.tb01054.x.
48.   Ingimarsson F. Breytileiki í höfuðbeinum bleikjuafbrigða í Þingvallavatni. [4th year thesis].
Reykjavík: University of Iceland; 2002.
49.   Noakes DLG. Charr truth: sympatric differentiation in Salvelinus species. Environmental Biology
of Fishes. 2008;83(1):7-15. doi: 10.1007/s10641-008-9379-x.
50.   Skúlason S, Noakes DLG, Snorrason SS. Ontogeny of trophic morphology in four sympatric
morphs of arctic charr Salvelinus alpinus in Thingvallavatn, Iceland. Biological Journal of the Linnean
Society. 1989;38(3):281-301. doi: 10.1111/j.1095-8312.1989.tb01579.x.
51.   Jónsdóttir GÓ. Diversity in the internal functional feeding elements of sympatric morphs of
Arctic charr (Salvelinus alpinus) [Master's thesis]. Reykjavik: University of Iceland; 2022.
52.   Baken EK, Collyer ML, Kaliontzopoulou A, Adams DC. geomorph v4.0 and gmShiny: Enhanced
analytics and a new graphical interface for a comprehensive morphometric experience. Methods in
Ecology and Evolution. 2021;12(12):2355-63. doi: https://doi.org/10.1111/2041-210X.13723.
53.   Adams D, Collyer M, Kaliontzopoulou A, Baken E. Geomorph: Software for geometric
morphometric analyses. R package version 4.0.4. 2022.
54.   Frédrich B, Sheets HD. Evolution of ontogenetic allometry shaping giant species: a case study
from the damselfish genus Dascyllus (Pomacentridae). Biological Journal of the Linnean Society.
2009;99(1):99-117. doi: 10.1111/j.1095-8312.2009.01336.x.
55.   Zelditch ML, Swiderski DL, Sheets HD, Fink WL. Geometric Morphometrics for Biologists: A
Primer: Elsevier Science; 2004.
56.   Pakkasmaa S, Rant E, Piironen J. A morphometric study on four land-locked salmonid species.
Annales Zoologici Fennici. 1998;35.
57.   de la Cámara M, Ponsioen L, Horta-Lacueva QJB, Kapralova KH. The Dynamic Ontogenetic Shape
Patterns of Adaptive Divergence and Sexual Dimorphism. Evolutionary Biology. 2023. doi:
10.1007/s11692-022-09592-y.
58.   Adams DC, Collyer ML. Multivariate Phylogenetic Comparative Methods: Evaluations,
Comparisons, and Recommendations. Systematic Biology. 2017;67(1):14-31. doi:
10.1093/sysbio/syx055.
59.   Friedman ST, Collyer ML, Price SA, Wainwright PC. Divergent Processes Drive Parallel Evolution
in Marine and Freshwater Fishes. Systematic Biology. 2021;71(6):1319-30. doi: 10.1093/sysbio/syab080.
60.   Rincon-Sandoval M, Duarte-Ribeiro E, Davis AM, Santaquiteria A, Hughes LC, Baldwin CC, et al.
Evolutionary determinism and convergence associated with water-column transitions in marine fishes.
Proceedings of the National Academy of Sciences. 2020;117(52):33396-403. doi:
doi:10.1073/pnas.2006511117.

61.   Burress ED, Holcomb JM, Tan M, Armbruster JW. Ecological diversification associated with the
benthic-to-pelagic transition by North American minnows. J Evol Biol. 2017;30(3):549-60. Epub
2016/12/08. doi: 10.1111/jeb.13024. PubMed PMID: 27925684.
62.   Hulsey CD, Roberts RJ, Loh YH, Rupp MF, Streelman JT. Lake Malawi cichlid evolution along a
benthic/limnetic axis. Ecol Evol. 2013;3(7):2262-72. Epub 2013/08/07. doi: 10.1002/ece3.633. PubMed
PMID: 23919168; PubMed Central PMCID: PMCPMC3728963.
63.   Astudillo-Clavijo V, Arbour JH, López-Fernández H. Selection towards different adaptive optima
drove the early diversification of locomotor phenotypes in the radiation of Neotropical geophagine
cichlids. BMC Evol Biol. 2015;15:77. Epub 2015/05/01. doi: 10.1186/s12862-015-0348-7. PubMed PMID:
25928151; PubMed Central PMCID: PMCPMC4435830.
64.   Ferry-Graham LA, Wainwright PC, Westneat MW, Bellwood DR. Mechanisms of benthic prey
capture in wrasses (Labridae). Marine Biology. 2002;141(5):819-30. doi: 10.1007/s00227-002-0882-x.
65.   Buser TJ, Finnegan DL, Summers AP, Kolmann MA. Have Niche, Will Travel. New Means of
Linking Diet and Ecomorphology Reveals Niche Conservatism in Freshwater Cottoid Fishes. Integrative
Organismal Biology. 2019;1(1). doi: 10.1093/iob/obz023.
66.   Cooper WJ, Parsons K, McIntyre A, Kern B, McGee-Moore A, Albertson RC. Bentho-pelagic
divergence of cichlid feeding architecture was prodigious and consistent during multiple adaptive
radiations within African rift-lakes. PLoS One. 2010;5(3):e9551. Epub 2010/03/12. doi:
10.1371/journal.pone.0009551. PubMed PMID: 20221400; PubMed Central PMCID: PMCPMC2833203.
67.   Alfaro ME, Bolnick DI, Wainwright PC. Evolutionary consequences of many-to-one mapping of
jaw morphology to mechanics in labrid fishes. Am Nat. 2005;165(6):E140-54. Epub 2005/06/07. doi:
10.1086/429564. PubMed PMID: 15937739.
68.   Kapralova KH, Jónsson ZO, Palsson A, Franzdóttir SR, le Deuff S, Kristjánsson BK, et al. Bones in
motion: Ontogeny of craniofacial development in sympatric arctic charr morphs. Dev Dyn.
2015;244(9):1168-78. Epub 2015/07/08. doi: 10.1002/dvdy.24302. PubMed PMID: 26150089.
69.   Küttner E, Parsons KJ, Robinson BW, Skúlason S, Danzmann RG, Ferguson MM. Effects of
population, family, and diet on craniofacial morphology of Icelandic Arctic charr (Salvelinus alpinus).
Biological Journal of the Linnean Society. 2013;108(3):702-14. doi: 10.1111/j.1095-8312.2012.02038.x.
70.   Saltykova E, Markevich G, Kuzishchin K. Divergent skull morphology between trophic separated
lacustrine forms of Dolly Varden charr from Lake Kronotskoe, Kamchatka, Russia. Environmental Biology
of Fishes. 2015;98(2):559-70. doi: https://doi.org/10.1007/s10641-014-0287-y. PubMed PMID:
1841104139.
71.   Gromova ES, Makhotin VV. Functional morphology of the visceral apapratus of Atlantic salmon
Salmo salar (Salmonidae). Journal of Ichthyology. 2016;56(4):505-21. doi: 10.1134/S0032945216040044.
72.   Copus JM, Gibb AC. A forceful upper jaw facilitates picking-based prey capture: biomechanics of
feeding in a butterflyfish, Chaetodon trichrous. Zoology. 2013;116(6):336-47. doi:
https://doi.org/10.1016/j.zool.2013.08.005.
73.   Ferry-Graham LA, Gibb AC, Hernandez LP. Premaxillary movements in cyprinodontiform fishes:
An unusual protrusion mechanism facilitates "picking" prey capture. Zoology. 2008;111(6):455-66. doi:
https://doi.org/10.1016/j.zool.2007.11.003.
74.   Yabe M. COMPARATIVE OSTEOLOGY AND MYOLOGY OF THE SUPERFAMILY COTTOIDEA (PISCES:
SCORPAENIFORMES), AND ITS PHYLOGENETIC CLASSIFICATION). MEMOIRS OF THE FACULTY OF
FISHERIES HOKKAIDO UNIVERSITY. 1985;32(1):1-130.
75.   St. John CA, Buser TJ, Kee VE, Kirilchik S, Bogdanov B, Neely D, et al. Diversification along a
benthic to pelagic gradient contributes to fish diversity in the world's largest lake (Lake Baikal, Russia).
Molecular Ecology. 2022;31(1):238-51. doi: https://doi.org/10.1111/mec.16209.

76.   Norton SF. A functional approach to ecomorphological patterns of feeding in cottid fishes. In:
Luczkovich JJ, Motta PJ, Norton SF, Liem KF, editors. Ecomorphology of fishes. Dordrecht: Springer
Netherlands; 1995. p. 61-78.
77.   Cooper WJ, Carter CB, Conith AJ, Rice AN, Westneat MW. The evolution of jaw protrusion
mechanics is tightly coupled to bentho-pelagic divergence in damselfishes (Pomacentridae). J Exp Biol.
2017;220(Pt 4):652-66. Epub 2016/12/04. doi: 10.1242/jeb.143115. PubMed PMID: 27913600.
78.   Witte F. Consistency and Functional Significance of Morphological Differences Between Wild-
Caught and Domestic Haplochromis Squamipinnis (Pisces, Cichlidae). Netherlands Journal of Zoology.
1983;34(4):596-612. doi: https://doi.org/10.1163/002829684X00308.
79.   Munyandamutsa PS, Jere WL, Kassam D, Mtethiwa A. Trophic divergence of Lake Kivu cichlid
fishes along a pelagic versus littoral habitat axis. Ecology and Evolution. 2021;11(4):1570-85. doi:
https://doi.org/10.1002/ece3.7117.
80.   Hulsey CD, Hollingsworth Jr PR, Holzman R. Co-evolution of the premaxilla and jaw protrusion in
cichlid fishes (Heroine: Cichlidae). Biological Journal of the Linnean Society. 2010;100(3):619-29. doi:
https://doi.org/10.1111/j.1095-8312.2010.01468.x.
81.   Sanford C. Salmonid Fish Osteology and Phylogeny (Teleostei: Salmonoidei). Theses Zoologicae.
2000;33:1-264.
82.   Herrel A, Gibb AC. Ontogeny of performance in vertebrates. Physiol Biochem Zool. 2006;79(1):1-
6. Epub 2005/12/29. doi: 10.1086/498196. PubMed PMID: 16380923.
83.   Kolmann MA, Huber DR. Scaling of feeding biomechanics in the horn shark Heterodontus
francisci: ontogenetic constraints on durophagy. Zoology (Jena). 2009;112(5):351-61. Epub 2009/05/12.
doi: 10.1016/j.zool.2008.11.002. PubMed PMID: 19428230.
84.   Skúlason S. Variation in morphology, life history and behaviour among sympatric morphs of
Arctic charr : an experimental approach [doctoral dissertation]. Ottawa: University of Guelph; 1991.
85.   Camp AL, Konow N, Sanford CP. Functional morphology and biomechanics of the tongue-bite
apparatus in salmonid and osteoglossomorph fishes. J Anat. 2009;214(5):717-28. Epub 2009/05/15. doi:
10.1111/j.1469-7580.2009.01056.x. PubMed PMID: 19438765; PubMed Central PMCID:
PMCPMC2707094.
86.   Olsen AM, Westneat MW. Linkage mechanisms in the vertebrate skull: Structure and function of
three-dimensional, parallel transmission systems. Journal of Morphology. 2016;277(12):1570-83. doi:
https://doi.org/10.1002/jmor.20596.
87.   Westneat MW. Skull Biomechanics and Suction Feeding in Fishes. Fish Physiology. 23: Academic
Press; 2005. p. 29-75.
88.   Kristjánsson BK, Skúlason S, Snorrason SS, Noakes DL. Fine-scale parallel patterns in diversity of
small benthic Arctic charr (Salvelinus alpinus) in relation to the ecology of lava/groundwater habitats.
Ecology and evolution. 2012;2(6):1099-112. doi: 10.1002/ece3.235. PubMed PMID: 22833787.
89.   Lauder GV. Functional and morphological bases of trophic specialization in sunfishes (Teleostei,
centrarchidae). J Morphol. 1983;178(1):1-21. Epub 1983/10/01. doi: 10.1002/jmor.1051780102.
PubMed PMID: 30060643.
90.   Huckins CJF. FUNCTIONAL LINKAGES AMONG MORPHOLOGY, FEEDING PERFORMANCE, DIET,
AND COMPETITIVE ABILITY IN MOLLUSCIVOROUS SUNFISH. Ecology. 1997;78(8):2401-14. doi:
https://doi.org/10.1890/0012-9658(1997)078[2401:FLAMFP]2.0.CO;2.
91.   Osenberg CW, Huckins CJF, Kaltenberg A, Martinez A. Resolving within- and between-population
variation in feeding ecology with a biomechanical model. Oecologia. 2004;141(1):57-65. doi:
10.1007/s00442-004-1650-z.
92.   Wainwright PC. Ecomorphology: Experimental Functional Anatomy for Ecological Problems.
American Zoologist. 1991;31(4):680-93. doi: 10.1093/icb/31.4.680.

93.   Wainwright PC. Ecological Explanation through Functional Morphology: The Feeding Biology of
Sunfishes. Ecology. 1996;77(5):1336-43. doi: https://doi.org/10.2307/2265531.
94.   Keast A. Feeding Interrelations Between Age-Groups of Pumpkinseed (Lepomis gibbosus) and
Comparisons with Bluegill (L. macrochirus). Journal of the Fisheries Research Board of Canada.
1978;35(1):12-27. doi: 10.1139/f78-003.
95.   Burress ED. Cichlid fishes as models of ecological diversification: patterns, mechanisms, and
consequences. Hydrobiologia. 2015;748(1):7-27. doi: 10.1007/s10750-014-1960-z.
96.   Amundsen P-A, Bøhn T, Våga GH. Gill raker morphology and feeding ecology of two sympatric
morphs of European whitefish (Coregonus lavaretus). Annales Zoologici Fennici. 2004;41(1):291-300.
97.   Werner EE, Hall DJ. Niche Shifts in Sunfishes: Experimental Evidence and Significance. Science.
1976;191(4225):404-6. doi: doi:10.1126/science.1246626.
98.   Berchtold AE, Colborne SF, Longstaffe FJ, Neff BD. Ecomorphological patterns linking
morphology and diet across three populations of pumpkinseed sunfish (Lepomis gibbosus). Canadian
Journal of Zoology. 2015;93(4):289-97. doi: 10.1139/cjz-2014-0236.
99.   Kusche H, Recknagel H, Elmer KR, Meyer A. Crater lake cichlids individually specialize along the
benthic–limnetic axis. Ecology and Evolution. 2014;4(7):1127-39. doi:
https://doi.org/10.1002/ece3.1015.
100.   Friedman ST, Price SA, Corn KA, Larouche O, Martinez CM, Wainwright PC. Body shape
diversification along the benthic–pelagic axis in marine fishes. Proceedings of the Royal Society B:
Biological Sciences. 2020;287(1931):20201053. doi: doi:10.1098/rspb.2020.1053.
101.   Østbye K, Amundsen P-A, Bernatchez L, Klemetsen A, Knudsen R, Kristoffersen R, et al. Parallel
evolution of ecomorphological traits in the European whitefish Coregonus lavaretus (L.) species complex
during postglacial times. Molecular Ecology. 2006;15(13):3983-4001. doi:
https://doi.org/10.1111/j.1365-294X.2006.03062.x.
102.   Albertson RC, Cooper WJ, Mann KA. More than meets the eye: functionally salient changes in
internal bone architecture accompany divergence in cichlid feeding mode. Int J Evol Biol.
2012;2012:538146. Epub 2012/06/06. doi: 10.1155/2012/538146. PubMed PMID: 22666625; PubMed
Central PMCID: PMCPMC3362014.
103.   Albertson RC, Kocher TD. Assessing morphological differences in an adaptive trait: a landmark-
based morphometric approach. J Exp Zool. 2001;289(6):385-403. Epub 2001/05/15. doi:
10.1002/jez.1020. PubMed PMID: 11351326.
104.   Roberts AS, Farina SC, Goforth RR, Gidmark NJ. Evolution of skeletal and muscular morphology
within the functionally integrated lower jaw adduction system of sculpins and relatives (Cottoidei).
Zoology. 2018;129:59-65. doi: https://doi.org/10.1016/j.zool.2018.06.006.
105.   Guðbrandsson J, Franzdóttir SR, Kristjánsson BK, Ahi EP, Maier VH, Kapralova KH, et al.
Differential gene expression during early development in recently evolved and sympatric Arctic charr
morphs. PeerJ. 2018;6:e4345. doi: https://doi.org/10.7717/peerj.4345.
106.   Kristjánsson BK, Leblanc CA-L, Skúlason S, Snorrason SS, Noakes DLG. Phenotypic plasticity in the
morphology of small benthic Icelandic Arctic charr (Salvelinus alpinus). Ecology of Freshwater Fish.
2018;27(3):636-45. doi: https://doi.org/10.1111/eff.12380.
107.   Skúlason S, Parsons KJ, Svanbäck R, Räsänen K, Ferguson MM, Adams CE, et al. A way forward
with eco evo devo: an extended theory of resource polymorphism with postglacial fishes as model
systems. Biological Reviews. 2019;94(5):1786-808. doi: https://doi.org/10.1111/brv.12534.
108.   Parsons KJ, Sheets HD, Skúlason S, Ferguson MM. Phenotypic plasticity, heterochrony and
ontogenetic repatterning during juvenile development of divergent Arctic charr (Salvelinus alpinus).
Journal of Evolutionary Biology. 2011;24(8):1640-52. Epub 2011/05/24. doi: 10.1111/j.1420-
9101.2011.02301.x. PubMed PMID: 21599773.

109.   Cutwa MM, Turingan RG. Intralocality Variation in Feeding Biomechanics and Prey Use in
Archosargus probatocephalus (Teleostei, Sparidae), with Implications for the Ecomorphology of Fishes.
Environmental Biology of Fishes. 2000;59(2):191-8. doi: 10.1023/A:1007679428331.
110.   Ghalambor CK, McKay JK, Carroll SP, Reznick DN. Adaptive versus non-adaptive phenotypic
plasticity and the potential for contemporary adaptation in new environments. Functional Ecology.
2007;21(3):394-407. doi: https://doi.org/10.1111/j.1365-2435.2007.01283.x.
111.   Eiríksson GM. Heterochrony in Bone Development and Growth in two Morphs of Arctic charr,
Salvelinus alpinus L., from Þingvallavatn, Iceland [Master's thesis]. Reykjavík: University of Iceland; 1999.
112.   Leblanc CA-L, Kristjánsson BK, Skúlason S. The importance of egg size and egg energy density for
early size patterns and performance of Arctic charr Salvelinus alpinus. Aquaculture Research.
2016;47(4):1100-11. doi: https://doi.org/10.1111/are.12566.
113.   Collar DC, O'Meara BC, Wainwright PC, Near TJ. PISCIVORY LIMITS DIVERSIFICATION OF FEEDING
MORPHOLOGY IN CENTRARCHID FISHES. Evolution. 2009;63(6):1557-73. doi:
https://doi.org/10.1111/j.1558-5646.2009.00626.x.
114.   Zimmerman E, Palsson A, Gibson G. Quantitative trait loci affecting components of wing shape in
Drosophila melanogaster. Genetics. 2000;155(2):671-83. Epub 2000/06/03. doi:
10.1093/genetics/155.2.671. PubMed PMID: 10835390; PubMed Central PMCID: PMCPMC1461095.
115.   Palsson A, Dodgson J, Dworkin I, Gibson G. Tests for the replication of an association between
Egfr and natural variation in Drosophila melanogaster wing morphology. BMC Genet. 2005;6:44. Epub
2005/08/17. doi: 10.1186/1471-2156-6-44. PubMed PMID: 16102176; PubMed Central PMCID:
PMCPMC1208880.
116.   Pitchers W, Nye J, Márquez EJ, Kowalski A, Dworkin I, Houle D. A Multivariate Genome-Wide
Association Study of Wing Shape in Drosophila melanogaster. Genetics. 2019;211(4):1429-47. Epub
2019/02/23. doi: 10.1534/genetics.118.301342. PubMed PMID: 30792267; PubMed Central PMCID:
PMCPMC6456314.
117.   Hallgrímsson B, Katz DC, Aponte JD, Larson JR, Devine J, Gonzalez PN, et al. Integration and the
Developmental Genetics of Allometry. Integrative and Comparative Biology. 2019;59(5):1369-81. doi:
10.1093/icb/icz105.
118.   Matlosz S, Sigurgeirsson B, Franzdóttir SR, Pálsson A, Jónsson ZO. DNA methylation differences
during development distinguish sympatric morphs of Arctic charr (Salvelinus alpinus). Molecular
Ecology. 2022;31(18):4739-61. doi: https://doi.org/10.1111/mec.16620.
119.   Kapralova KH, Franzdóttir SR, Jónsson H, Snorrason SS, Jónsson ZO. Patterns of MiRNA
Expression in Arctic Charr Development. PLOS ONE. 2014;9(8):e106084. doi:
10.1371/journal.pone.0106084.
120.   Tamate T, Maekawa K. LATITUDINAL VARIATION IN SEXUAL SIZE DIMORPHISM OF SEA-RUN
MASU SALMON, ONCORHYNCHUS MASOU. Evolution. 2006;60(1):196-201. doi:
https://doi.org/10.1111/j.0014-3820.2006.tb01094.x.
121.   Vidal-García M, Keogh JS. Invasive cane toads are unique in shape but overlap in ecological niche
compared to Australian native frogs. Ecology and Evolution. 2017;7(19):7609-19. doi:
https://doi.org/10.1002/ece3.3253.
122.   Cardini A. Missing the third dimension in geometric morphometrics: how to assess if 2D images
really are a good proxy for 3D structures? Hystrix, the Italian Journal of Mammalogy. 2014;25(2):73-81.
doi: 10.4404/hystrix-25.2-10993.

Supporting information

S1 Appendix. Sampling scheme and summary statistics for the four sympatric charr morphs. Numbers of individuals, average fork length (FL), weight and age and the standard deviation (SD) for length, weight and age of the 240 individuals.

[1]Fork length: length from tip of snout to the posterior tip of middle caudal fin ray.

Log: natural logarithmic transformation.

S2 Appendix. Explanation of 37 landmarks used for landmarking the external shape, X-S indicates sliding landmark.

S3 Appendix. Explanation of landmarks used to capture the shape of the six bones (dentary, articular-angular, quadrate, premaxilla, maxilla and supramaxilla). X-S indicates sliding landmark. See Fig 1D - E for placement of landmarks on bones.

S4 Appendix. Variation in size and age of sexually mature fish of the four sympatric morphs. (A) Histogram, showing age (years) distribution for all morphs by sex (NA indicates SB that could not be sexed). (B) The variation in fork length (FL, cm) by morph by sex represented by a boxplot. (C) The variation in $\log_e$ weight (g) by morph by sex are represented in boxplots.

S5 Appendix. ANOVA results from tests of the influence of morph and sex effects on length (cm FL), $\log_e$ weight (g) and age (years).

F = F-statistic, P = P - values, < 0.05 in bold

S6 Appendix. Results from the Procrustes ANOVA, for effects of morph and $\log_e$ FL on the size (centroid size) of the six bones.

* Despite the p-values, all R^2 for the interaction term were small and no pairwise test of slope by morph were significant.

R^2 = coefficient of determination; F = F-statistic; Z = Z-statistic; P = P-values, < 0.008 in bold.

S7 Appendix. Size corrected PC-plots of specimens and deformation grids showing shape variation in (top) external whole-body shape and (bottom) head shape. Plots of shape warps on X- and Y-axis are unmagnified. Each dot represents an individual and the ellipses represent 95% CI for the distribution by morph (large dot represents the mean of each morph distribution in these two dimensions of shape). For the whole-body (top) PC2 and 3 explain 18% and 8% of the variation respectively (PC1 was biased by sampling error, and not depicted) and for the head shape (bottom) PC3 and 4 explain 40% and 22% respectively (PC1 and PC2 were biased by sampling error, and not depicted).

S8 Appendix. Shape variation in the two external shapes and 6 head bones. Analysed with Procrustes ANOVA, testing for influence of sex and size (centroid size).

R^2 = coefficient of determination; F = F-statistic; Z = Z-statistic; P = P-values, ≤ 0.008 in bold

S9 Appendix. PC-plots of specimens showing shape differences between replicates (trials), for dentary,
premaxilla, articular-angular, maxilla, quadrate and supramaxilla (from top to bottom, left to right). Each dot
represents an individual and the ellipses represent 95% CI for the distribution by replicates (large dot represents the
mean for each morph replicates in these dimensions).

S10 Appendix. Ordination plot (from partial least squares analysis (PLS)) showing the degree of association
between the two replicates (trials) for dentary, premaxilla, articular-angular, maxilla, quadrate and
supramaxilla (from top to bottom, left to right). Block 1 (x-axis) is replicate 1 and Block 2 (y-axis) is replicate 2.
For all bones association between the replicates was always significant (p < 0.001). With the correlation coefficient
being, dentary: 0.989, premaxilla: 0.939, articular-angular: 0.994, maxilla: 0.986, quadrate: 0.942 and supramaxilla:
0.835.

S11 Appendix. ANOVA results from tests of the influence of individual variation and measurement error
(replicates) effects on shape variation in 6 jaw bones. Analysed with Procrustes ANCOVA.

R^2 = coefficient of determination; F = F-statistic; Z = Z-statistic; P = P-values, < 0.008 in bold.
944

S12 Appendix. Size and shape allometry of (top) quadrate and (bottom) supramaxilla). Left panels:
relationships between bone size and fork length by morph; Right panels: relationships between bone shape and bone
size. Shown are values for individuals (open circles) and the predicted values (filled) of regressions for both bone
size (CS_{bone}) vs body size (log_e FL) or bone-shape versus bone-size. Log_e, natural log transformation.

S13 Appendix. Estimates of regression for the relationships between the size of the individual and the size of
the bone, per morph. See also Fig 2 and S6 and S12 Appendix.

S14 Appendix. Results from pairwise comparisons of variation in bone size between morphs for the six bones
(dentary, articular-angular, quadrate, premaxilla, maxilla and supramaxilla). Based on a model assuming no
difference in allometry by morphs.

D = pairwise distances between means; UCL = upper confidence level; Z = Z-statistic; P-values < 0.008 in bold
(Bonferroni).

S15 Appendix. Information on PCA, for all bone shapes, when size effects have not been removed. Symbols
indicate how morphs align, # for when morphs align along morphotype, * when morphs align along the size- gradient.
Double Symbols (* or #) for prominent separation and symbol in brackets for minor.

S16 Appendix. Pairwise distances from tests of dentary shape allometry by morph, between vector angles and
absolute differences between vector lengths. Following ANCOVA model with the interaction of size (bone
centroid size) and morph.

r = slope vector correlations; *angle:* between the two vectors being compared; *d* = pairwise distances between
means; UCL = upper confidence level; Z = Z-statistic; P-values, ≤ 0.008 in bold (Bonferroni correction).

S17 Appendix. Results from pairwise tests of shape by morphs, for differences between absolute vector lengths (D) used to examine possible shape differences by morph in five bones. These bones had significant shape differences according to ANOVA model with main effects and interaction of morph by bone size. Used to verify results from ANOVA and test for true allometry (i.e., different group slopes).

D = pairwise differences between vector lengths; UCL = upper confidence level; Z = Z-statistic; P-values, < 0.008 in bold

S18 Appendix. Results from "Significant Vector Angle test" examining possibly allometric shape differences by morph for five bones, that showed significant allometric differences according to ANOVA model with main and interaction effects (morph x bone size). Used to verify results from ANOVA and test for true allometry (i.e., different group slopes).

R = slope of vector correlation; UCL = upper confidence level; Z = Z-statistic; P-values < 0.008 in bold

S19 Appendix. Size corrected PC-plots of specimens and deformation grids showing shape variation in (top) quadrate and (bottom) supramaxilla. Plots of shape warps on X- and Y-axis are unmagnified. Each dot represents an individual and the ellipses represent 95% CI for the distribution by morph (large dot represents the mean of each morph distribution in these two dimensions of shape). For the quadrate (top) PC1 and 2 explain 30% and 14% of the variation respectively and the supramaxilla (bottom) PC1 and 2 explain 40% and 22% respectively.

S20 Appendix. Tests of integration among craniofacial bones in arctic charr. The six bones were compared using partial least squares (PLS), within anatomical regions (upper and lower jaws) and between regions.

*Effect size of maxilla-supramaxilla were significantly higher than premaxilla-maxilla.

**Effect size of dentary-maxilla and articular-angular were significantly higher than articular-angular-supramaxilla.