Within-species variation in the gut microbiome of medaka (*Oryzias latipes*) is driven by

the interaction of light intensity and genetic background

C. Evangelista[1§*], S. Kamenova[1,2,3], B. Diaz Pauli[1,4], J. Sandkjenn[1], L.A. Vøllestad[1], E.

Edeline[5,6], P. Trosvik[1,†], EJ. de Muinck[1,†]

[1] Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences,

University of Oslo, Oslo, Norway

[2] Departments of Ecology and Natural Resource Management, Norwegian University of Life

Sciences, Ås, Norway

[3] National Museum of Natural History, Sofia, Bulgaria

[4] Department of Biological Science, University of Bergen, Norway

[5] Sorbonne Université, CNRS, IRD, INRA, Institut d'écologie et des sciences de

l'environnement (IEES), 4 place Jussieu, Paris 75005, France.

[6] DECOD (Ecosystem Dynamics and Sustainability), INRAE, IFREMER, Institut Agro,

Rennes, France

[§] Current address: Aquatic Ecology Group, University of Vic - Central University of

Catalonia, 08500, Vic, Spain

[†]co-senior authorship

Corresponding author: charlotte.evangelista0@gmail.com

Abstract

Unravelling evolution-by-environment interactions on the gut microbiome is particularly relevant considering the unprecedented level of human-driven disruption of the ecological and evolutionary trajectories of species. Here, we aimed to evaluate whether an evolutionary response to size-selective mortality influences the gut microbiome of medaka (*Oryzias latipes*), how environmental conditions interact with the genetic background of medaka on their microbiota, and the association between microbiome diversity and medaka growth-related traits. To do so, we studied two lineages of medaka with known divergence in foraging efficiency and life history raised under antagonistic size-selective regimes for 10 generations (i.e. the largest or the smallest breeders were removed to mimic fishing-like or natural mortality). In pond mesocosms, the two lineages were subjected to contrasting population density and light intensity (used as proxies of resource availability). We observed significant differences in the gut microbiome composition and richness between the two lines, and this effect was mediated by light intensity. The bacterial richness of fishing-like medaka (small-breeder line) was reduced by 34% under low-light conditions compared to high-light conditions, while it remained unchanged in natural mortality-selected medaka (large-breeder line). However, the observed changes in bacterial richness did not correlate with changes in adult growth rate or body condition. Given the growing evidence about the gut microbiomes importance to host health, more in-depth studies are required to fully understand the role of the microbiome in size-selected organisms and the possible ecosystem-level consequences.

Introduction

Over the past decade, the expansion of scientific literature investigating the gut microbiome

has been largely motivated by increasing evidence of the microbiome's role in health

maintenance. Studies have highlighted the variety of pathways through which the gut

microbiome can play this role – from influencing nutrient uptake and metabolism to

regulating immune responses and pathogen susceptibility (Hanning and Diaz-Sanchez 2015,

Moran et al. 2019). Gut-associated microbiomes can also incur the consequences of various

anthropogenic stressors (Sepulveda and Moeller 2020, Teyssier et al. 2018, Trosvik et al.

2018, Sonnenburg and Sonnenburg 2019, Fouladi et al. 2020, Varg et al. 2021). These

anthropogenic stressors may lead to imbalances in the gut microbiome (dysbiosis) with

potential implications for individual host fitness. A notable example is the effect of increased

temperature leading to reduced gut bacterial diversity in the common lizard (*Zootoca*

vivipara), also potentially associated with a reduction in survival (Bestion et al. 2017).

Understanding how changes in the gut microbiome are related to host health and survival is

essential to fully understand the mechanisms behind anthropogenic impact on wild

populations.

In teleosts (bony fish), the genetic background (Bolnick et al. 2014a, Smith et al.

2015) and various environmental factors such as diet composition (Bolnick et al. 2014b,

Eckert et al. 2020) and the quality of the surrounding water (Talwar et al. 2018) have been

linked to changes in the gut microbiome composition. However, as genotypes and

environmental conditions are often interacting, disentangling their respective effects on the

microbiome is challenging, and have been mainly studied independently (but see Navarrete et

al. 2012, Sevellec et al. 2014, Sullam et al. 2015). Currently, we lack assessments of

genotype-by-environment effects on fish microbiome (Spor et al. 2011, Talwar et al. 2018,

Piazzon et al. 2020) which limit our understanding about the factors driving variation in gut

microbiome assemblages in teleosts, a comparatively under-studied taxa in microbiome

science (Sullam et al. 2012).

Size-selective harvesting of wild populations is an acute disturbance factor, driving

very fast rates of evolutionary change towards smaller size and faster life history (Darimont

et al. 2009, Nusslé et al. 2012, Heino et al. 2015, Sanderson et al. 2022). Simultaneously,

harvesting reduces population density and hence also increases resource availability. The

evolutionary change driven by size-selective harvesting (van Wijk et al. 2012, Uusi-Heikkilä

et al. 2017) has, in turn, the potential to reshuffle trophic interactions within food webs, with

individuals from heavily harvested populations tending to display narrower diets (Hočevar

and Kuparinen 2021). Assuming that bacterial transmission through diet influences gut

microbial diversity, individuals with narrower diets may have less diverse bacterial

communities (Sevellec et al. 2014). In fish exposed to fishing, the diet and gut microbiome of

fish could be altered by the evolutionary change in size due to size-selective harvesting, the

increase prey abundance due to reduced density by harvesting, and changes in primary

producers that can have a natural cause (algal blooms) or be indirectly linked to harvesting

too, as it was seen that fishing led to higher abundance of primary producers (Frank et al.

2005). Hence, size-selective harvesting provides an ideal context to explore genotype-by-

environment interactions on the gut microbiome but have been largely understudied so far.

In fisheries science, laboratory size-selection experiments are commonly used to

mimic the evolutionary consequences of size-selective fishing, while controlling for

phenotypic plasticity (Conover and Munch 2002; Uusi-Heikkilä et al. 2015). Recently,

Renneville et al. (2020) performed a size-selection experiment using medaka (*Oryzias*

latipes) as a model species. Native to East Asian countries, medaka is a small cyprinodont

fish (adult length = 32 mm) that has a short generation time and is easily reared in the

laboratory, making it an ideal species for selection experiments (Ruzzante and Doyle 1993,

Renneville et al. 2020, Bouffet-Halle et al. 2021). The species is omnivorous with an animal-

based diet preference but can also feed on diatoms and filamentous algae (Edeline et al.

2016). The size-selection procedure consisted of mimicking either fishing mortality where

only small-bodied fish were allowed to reproduce (small-breeder SB line), or a more natural

mortality regime favoring the reproduction of large-bodied fish (large-breeder LB line)

(Reneville et al. 2020, Le Rouzic et al. 2020). As expected from the literature (Stearns 1992,

Conover and Munch 2002), the LB and SB lines evolved opposite life-history traits and

behaviors: small-breeder medaka grew slower, matured earlier and were less efficient

foragers than the large-breeder medaka (Diaz Pauli et al. 2019, Evangelista et al. 2021). The

logical next step is to examine to what extent changes in the gut microbiota of LB and SB

medaka are driven by the interaction between fisheries-induced changes in both evolution

(life-history shift) and environmental conditions (reduced fish density and concomitant

increased resource availability).

In the present study, we assessed how the genetic background of the two medaka lines

interacted with population density and light intensity (used to modulate primary production)

to shape the gut microbiome composition and diversity. For this, SB and LB medaka were

introduced into replicated pond mesocosms for three months. Based on life-history and

foraging traits divergence between the two lines (Diaz Pauli et al. 2019, Evangelista et al.

2020, Evangelista et al. 2021), and based on the assumption that such traits are key drivers of

gut microbiome variation in teleosts (Talwar et al. 2018), we hypothesized divergent gut

microbial communities between SB and LB medaka. We further hypothesized that gut

microbiome differences between the two lines would be more pronounced under suboptimal

conditions, i.e. when access to food resources is limited (Reese and Dunn 2018, Varg et al.

2021). Finally, because microbiome diversity could be important for host fitness, we

evaluated whether fitness proxies (i.e. somatic growth rate and body condition) were

associated with variation in gut microbiome diversity (Bolnick et al. 2014b).

130   **Methods**

Size-dependent selection and fish rearing

The two medaka lines were size-selected over 10 generations under identical laboratory

conditions. Specifically, medaka were kept in 3-L tank at similar density (14 – 17 fish per

tank, 20 tanks per line per generation; Le Rouzic et al. 2020, Renneville et al. 2020), and at

the same temperature (26°C) and photoperiod (14 h Light / 10 h Dark). They were fed *ad*

libitum with a mixed diet of dry food and living brine shrimp *Artemia salina* and/or *Turbatrix*

aceti. These standardized environmental conditions ensured that phenotypic differences

among the selected lines reflected a genetically-based, evolutionary divergence in response to

size-selective harvesting alone (Le Rouzic et al. 2020, Renneville et al. 2020). In addition,

offspring from each breeding pair were transferred to the same tank, so one was able to keep

track of individual pedigrees during the whole selection experiment (Renneville et al. 2020).

The selection procedure consisted in removing the largest or the smallest breeders,

thus producing two lines with distinct life-history strategies: the small-breeder line where

only small-bodied individuals were allowed to reproduce, and the large-breeder lines. At each

generation, on average 88% of fish were removed per line. Size selection was both family-

and individual-based. At 60 day-post-hatching (dph), among a total of at least 20 families per

line (one tank per family to keep track of individual pedigrees), the 10 families with the

largest (large-breeder line) or smallest (small-breeder line) average standard body length (SL)

were kept. At 75 dph, individuals within each of the selected families were measured and the

largest-bodied (large-breeder line) or the smallest-bodied (small-breeder line) mature males

(n = 2 or 3) and females (n = 2 or 3) were used as breeders for the next generation (further

details available in Renneville et al. 2020). On average, at 75 dph, SL was 20.7 mm in small

breeders and 22.0 mm in large breeders (a 5.7 % difference), and the probability of being

mature was 91.7% in small breeders and 77 % in large breeders (a 18.0 % difference)

(Renneville et al. 2020).

In June 2017, experimental populations were created using fish from the eleventh

generation. Specifically, for each line, 180 mature fish (initial standard body length: mean $\pm$

SD; SL_i in small-breeder = 18.9 mm $\pm$ 1.4; SL_i in large-breeder = 19.4 mm $\pm$ 1.4; ANOVA:

$F_{1, 358} = 13.70$, $P < 0.001$) were selected to generate 24 experimental populations composed

of individuals from the same line (48 populations in total), but from distinct families to limit

inbreeding (mean kinship coefficient = 0.23 $\pm$ 0.1 and 0.17 $\pm$ 0.1 SE in LB and SB lines,

respectively; further details available in Le Rouzic et al. 2020). Selected fish were

anaesthetized with MS-222 and marked using visible implant elastomer (VIE; Northwest

Marine Technology, Shaw Island, WA, USA) to render each fish individually identifiable and

to allow the calculation of growth-related traits. Fish from the same experimental population

were pooled in a 3 L tank and maintained at the laboratory until the beginning of the

experiment when they were released into an outdoor mesocosm (48 mesocosms in total).

Outdoor mesocosm experiment

The outdoor experiment was conducted at the CEREEP-Ecotron Ile de France (Saint-Pierre-

les-Nemours, France; cereep.bio.ens.psl.eu) using 48 mesocosms (500 L, 0.8 m deep, 1.0 m

diameter) arranged in 5 blocks. All mesocosms were filled simultaneously from 4 to 6 April

2017 with a mix of dechlorinated tap water (100 L) and oligotrophic water from a local pond

(300 L). The pond water was pre-filtered through 150 μm mesh to remove large benthic

invertebrates, zooplankton and debris. The mesocosms were supplied with 2 L of mature

sediment mixture including benthic invertebrates (mainly Ephemeroptera and Chironomidae

larvae, Planorbidae, Hydrachnidia, Nematoda and Ostracoda) and 2 L of homogenized

mixture of zooplankton (Copepoda and Cladocera) collected from local ponds. In each

mesocosm, two floating shelters made of wool threads (30 cm length) provided spawning

substrate and two floating brushes made of plastic threads provided protection for larvae.

Each mesocosm was then covered with a shading net (see details below) and given 3 months

to mature before fish were introduced. On 12 June, all mesocosms were enriched with 2 mL

of a liquid mixture of 0.32 μg P L^{-1} as KH$_2$PO$_4$ and 6.32 μg N L^{-1} as NaNO$_3$ to favor primary

production.

On 4 July 2017, large- and small-breeder fish were released into the outdoor

mesocosms under contrasting environmental conditions. The experiment consisted of a 2×2

$\times 2$ full factorial design with size-selected line (LB and SB) crossed with density (high HD

and low LD) and light intensity (high HL and low LL). Each treatment combination was

replicated six times (48 mesocosm in total; Fig. 1). High- and low-density treatments

consisted of twelve (3.2 mg fish L-1 ± 0.3 SD) and three (0.9 mg fish L-1 ± 0.1 SD) fish per

mesocosm, respectively, with a female biased sex ratio of 2:1. Light intensity was

manipulated using shade nets with different mesh size that allowed the passage of 92% (high

light intensity HL) and 70% of ambient light (low light intensity LL). Light supply was used

to modulate primary production while avoiding too high growth of filamentous algae. This

factorial design resulted in a total of 8 treatment combinations (Fig. 1), each with 6 replicates.

Figure 1 (a) Design of the mesocosm experiment used to test the effects of Line × Density and Line × Light intensity on fish gut microbiome. Fish from the Large-breeder (LB) line are in orange and fish from the Small-breeder (SB) line are in blue. **(b)** Pictures of the outdoor mesocosms (upper picture) and shade nets used to manipulate light intensity (lower pictures).

Gut microbiome sampling and growth-related trait measurements

On 22 September 2017, marked fish were recaptured with hand nets (survival rate = 92%). A total of 126 marked fish were randomly and homogeneously sub-sampled among the mesocosms (number of fish per treatment: mean ± SD = 15.8 ± 0.9, min = 14, max = 17; number of fish per mesocosm: mean ± SD = 2.6 ± 0.6, min = 1, max = 4). After 24 hours fasting, each selected fish was measured for final standard length (SL_f ± 1 mm), weighed (W_f ± 1 mg), euthanized using MS-222 and dissected using disposable laboratory-grade razor blades. The whole intestine (including potential remaining content because of small size) was sampled and flash-frozen in liquid nitrogen for up to 5 hours, then stored in a -80☐ freezer until DNA extraction. To limit contaminations during dissection, working environment and dissection tools were sterilised between each individual. Scalpels were thoroughly washed using 96% laboratory grade ethanol.

216   Body condition of each selected individual was calculated using the residuals of the

217   relationship between $\log_{10} W_f$ and $\log_{10} SL_f$. The somatic growth rate (mm month^{-1}) of each

218   selected fish was calculated as follows:

$$Growthrate = \left(SL_f - SL_i\right)/t$$

where SL_f and SL_i are the final and initial standard length, and t is the duration of the

experiment (3 months).

DNA extraction and sequencing

After defrosting at room temperature, bacterial DNA from medaka gut samples was extracted

using the DNeasy PowerSoil kit (Qiagen, Germany) according to the manufacturer's

instructions. The quantity and quality of purified DNA was checked using a NanoDrop

spectrophotometer (Thermo Fisher Scientific, USA). Library preparation for Illumina

sequencing was carried out according to the dual indexing protocol described by Fadrosh et

al. (2014). This protocol uses the 319F and 806R primer set to amplify the V3-V4 region of

the 16S rRNA gene. DNA sequencing was done on an Illumina MiSeq apparatus in 300 bp

PE mode. The DNA sequencing was carried out at the Norwegian Sequencing Centre (NSC),

and sequence demultiplexing was done using the custom NSC "demultiplexer" software

(https://github.com/nsc-norway/triple_index-demultiplexing/tree/master/src),   which   also

removes barcode sequences and heterogeneity spacers. Among the 126 samples, seven

displayed amplification failure, and one was removed from the dataset due to mislabelling.

Bioinformatics analysis

Further sequence data processing was performed using the Divisive Amplicon Denoising

Algorithm as implemented in the dada2 v1.16 R-package (Callahan et al. 2016). Taxonomic

classification of amplicon sequence variants (ASVs) was carried out using the Ribosomal

Database Project v16 training set (Wang et al. 2007). In addition to the fish samples, four

DNA extraction and one PCR negative controls (i.e. ultra-pure water instead of DNA) were

sequenced and processed as described above. The quality profiles of the negative controls

after filtering showed that each control contained very low number of sequences of very poor

quality (Evangelista et al., unpublished data). The sequences that did make it through

processing (~290 sequences for all 5 samples combined) corresponded to only 7 unique

sequences, each unique to a negative control sample. The taxonomic classification of these 7

unique sequences referred to taxa that are not expected in the fish gut samples (*Moraxella*,

Ralstonia, *Brevundimonas*, *Massilia*, *Propionibacterium*, *Micrococcus*, and *Knoellia*). This

suggests that overall little contamination occurred.

Using the R package phyloseq (v.1.40.0, McMurdie and Holmes 2013), we further

filtered the fish gut data in order to remove any contaminant or artefactual sequences. First,

ASVs with a Phylum-level assignment probability < 0.80 and those classified as chloroplast

DNA were discarded from the dataset. Second, we excluded all ASVs with a total abundance

lower than 0.005% of the dataset's total abundance as they are most likely sequencing errors

(Bokulich et al. 2013). Finally, samples with a total sequence reads abundance of < 5000

reads were removed from the dataset (n = 15). The dataset consisted of 103 samples,

comprising 627 ASVs for a total of 3,591,039 sequence reads. Sequencing depth ranged from

a minimum of 5588 to 85470 reads per sample, with a mean of 34864 reads per sample.

Between-sample differences in library sequencing depth were standardized to the median

sequencing depth (Appendix S1).

Statistical analyses

All statistical analyses were run with R v.4.2.1 (R Development Core Team, 2022) using the

Family level as taxonomic resolution because it was the best taxonomic level for

discriminating (median bootstrap support is 0.52 and 0.89 for Family and Genus level

taxonomic assignment, respectively). We used a PERMANOVA to test for differences in

community composition according to Line × Density and Line × Light intensity. This was

carried out using the *adonis* function in the vegan package (v.2.6.4, Oksanen et al. 2020), by

implementing weighted (to take into account information on relative abundance) and

unweighted (to take into account taxa presence or absence) UniFrac dissimilarities based on

999 permutations. Statistical tests indicated that there was no deviation from multivariate

dispersion (*betadisper* function from vegan), except for the Line × Light intensity factor in

the unweighted UniFrac distance PERMANOVA. Nonetheless, sample size was similar

between treatment combinations and results should be robust (Anderson and Walsh 2013).

Non-metric multidimensional scaling (NMDS) plots with both weighted and unweighted

UniFrac distances were used to visualize gut microbiome composition.

Based on these community composition analyses, we agglomerated the data to family

level and visualized the relative abundance of normalized data according to the Line

treatment. Significant effects of Line or Line × Environment on gut microbiome community

composition were further investigated using differential abundance analysis based on the

linear discriminant analysis (LDA) on effect size (LEfSe) method. LEfSe was implemented

in the microeco package (v.0.12.1, Liu et al. 2021) using a non-parametric Kruskal-Wallis

test to detect differences in Family abundance (bootstrap test number = 100, significance

threshold = 0.01).

Gut microbiome diversity was estimated using the first three Hill numbers (^{q}D; Chiu

and Chao 2014, Alberdi and Gilbert 2019a) calculated using the R package hilldiv (v.1.5.1,

Alberdi and Gilbert 2019b): $q = 0$ (species richness), $q = 1$ (the exponential of Shannon's

entropy index) and $q = 2$ (the inverse of the Simpson's diversity index). Linear models were

used to test the effect of Line × Density and Line × Light intensity on each Hill number.

When significant, the interactions were further investigated using post hoc Tukey's pairwise

comparison using the emmeans package (v.1.8.1.1, Lenth 2021). Spearman correlations

(adjusted for multiple testing using false discovery rate (fdr) correction) were used to test for

associations between bacterial richness and diversity (i.e. the first three Hill numbers) with

medaka growth-related traits (i.e. body condition and somatic growth rate), as well as with

medaka final standard length. These correlations were performed using the *corr.test* function

from the psych package (v.2.2.9, Revelle 2021). Finally, we applied a Canonical Correlation

Analysis (CCA) to explore the relationships between gut microbiome composition and

medaka's traits. CCA was conducted using the mixOmics package (v.6.23.4, Rohart et al.

2017). Results from the CCA were visualised using a correlation circle plot.

Results

Composition of the whole gut microbiome

After standardization of the data, we identified a total of 3,189,868 sequence reads (mean =

30,969 reads per sample) and 627 ASVs for 103 samples. As expected, dominant phylum of

the medaka gut microbiome included Proteobacteria (60.5% and 59.2% of reads in LB and

SB medaka, respectively), followed by Bacteroidetes (2.5 and 2.4%, respectively),

Verrucomicrobia (2.0 and 3.2%, respectively) and Actinobacteria (2.2 and 3.9%,

respectively). Bacterial communities were also characterized by large relative abundance of

Cyanobacteria (29.2 and 28.8%, respectively), while Firmicutes represented only 0.7% and

0.5% of reads in LB and SB medaka, respectively (Fig. S2).

Genotype-driven variation in gut microbiome

The two medaka lines had distinct gut microbial communities (weighted UniFrac distance

PERMANOVA: F = 2.30, P = 0.025, R^2 = 0.022; Fig. 2a). Overall, 5 families (i.e.

Aeromonadaceae, Neisseriaceae, Family_II, Rhodobacteraceae and unspecified

Cyanobacteria) dominated the gut microbiome of all samples, comprising together 62% and

59% of the total bacterial relative abundance of LB and SB medaka lines, respectively (Fig.

2b). Of these families, the relative abundance of both *Aeromonadaceae* and *Neisseriaceae*

were significantly higher in LB than in SB medaka (LEfSe; P = 0.001 and P = 0.016,

respectively; Fig. 2c), while *Rhodobacteraceae* was more abundant (relative abundance) in

the gut of SB than LB medaka (LEfSe: P = 0.017; Fig. 2c). The two other families did not

significantly differ between the two lines. Unweighted UniFrac distance PERMANOVA

revealed no significant difference between SB and LB medaka (F = 1.32, P = 0.109, R^2 =

0.013), indicating that low-abundance taxa did not differ between the two lines.

Figure 2 (a) NMDS ordination (NMDS stress = 0.13) of variation in bacterial community
composition of fish from the Large-breeder (LB; orange dots) and Small-breeder (SB; blue
dots) lines. Data represent ordination based on weighted UniFrac distances among the 103 fish
individuals. **(b)** Gut microbial taxonomic composition (Family level) of fish according to its
selective background (Large-breeder line, LB; Small-breeder line SB). **(c)** Linear discriminant
analysis (LDA; LDA score > 2) effect sizes representing the twelve ASVs (Family level) that

significantly differ in relative abundance between the Large-breeder (orange) and Small-
breeder (blue) lines.

Differential relative abundance analyses showed that *Family IV*, *Oxalobacteraceae*,

and *Rhodocyclaceae* were significantly more abundant in the gut of LB than SB medaka (P =

0.030, P = 0.006, P = 0.018, respectively). In contrast, the gut microbiome of SB medaka had

more *Microbacteriaceae* (P = 0.009), *Rhizobiaceae* (P = 0.027), *Rhizobiales* (*unspecified*

Family, P = 0.033), *Methylobacteriaceae* (P = 0.017), *Family I* (P = 0.020) and

Hyphomicrobiaceae (P = 0.030) than that of LB ones (Fig. 2c).

Table 1 Analysis-of-variance table derived from the linear models used to assess the effect of
size-selected line, fish density, and light intensity on gut microbiome diversity (${}^{q=0}$D, ${}^{q=1}$D
and ${}^{q=2}$D). Significant P values are highlighted in bold.
343

Responses	Predictors	Sum of Sq.	F_{df}	*P*
${}^{q=0}$D	Intercept	338340	111.04_1	**< 0.001**
	Line	21006	6.89_1	**0.010**
	Density	1187	0.39_1	0.534
	Light intensity	148	0.05_1	0.826
	Line × Density	11306	3.71_1	0.057
	Line × Light intensity	22238	7.30_1	**0.008**
	Residuals	295573	$—_{97}$	—
${}^{q=1}$D	Line	1424	3.66_1	0.059
	Density	196	0.50_1	0.479
	Light intensity	173	0.45_1	0.506
	Line × Density	531	1.37_1	0.245
	Line × Light intensity	126	0.32_1	0.570
	Residuals	37732	$—_{97}$	—
${}^{q=2}$D	Line	408.2	3.40_1	0.068
	Density	5.7	0.05_1	0.828
	Light intensity	1.4	0.01_1	0.915
	Line × Density	105.4	0.88_1	0.351
	Line × Light intensity	0.0	0.00_1	0.995
	Residuals	116642.1	$—_{97}$	—

Environment-driven variation in gut microbiome

Weighted and unweighted UniFrac distance PERMANOVA showed no effect of light

intensity (F = 1.23, P = 0.265, R^2 = 0.012; F = 1.45, P = 0.061, R^2 = 0.014) and fish density

(F = 1.35, P = 0.195, R^2 = 0.013; F = 1.31, P = 0.111, R^2 = 0.013) on the gut microbial

community (Fig. 3). Similarly, the gut microbiome diversity of medaka (species richness,

exponential Shannon or inverse Simpson) were not influenced by the environmental

conditions alone (Table 1).

352
353 **Figure 3** NMDS ordination (NMDS stress = 0.13) of variation in bacterial community
354 composition **(a)** in low or high light intensity (grey and yellow dots, respectively), **(c)** in low or
355 high fish density (pink and green dots, respectively. Data represent ordination based on
356 weighted UniFrac distances among the 103 fish individuals.
357

Gene-by-environment interaction on gut microbiome

Weighted UniFrac analyses showed that the interactions between medaka line and the

environment (i.e. Line × Density and Line × Light intensity) had no significant effect on gut

microbiome composition (PERMANOVA: F = 1.08, P = 0.330, R^2 = 0.010 and F = 0.82, P =

0.561, R^2 = 0.001, respectively). In contrast, unweighted UniFRac analyses revealed that

Line × Light intensity significantly influenced the gut microbial composition (F = 1.66, P =

0.020, R^2 = 0.016). Indeed, the gut microbiome of LB medaka depicted greater compositional

diversity than that of SB medaka in the high-light treatment, while the opposite pattern was

observed in the low-light treatment (Fig. 4a).

Gut microbial richness was modulated by the interaction between the medaka line and

light intensity (Table 1). Specifically, SB medaka had a higher bacterial richness (when q =

0) in the high-light intensity compared to the low-light treatment (Line × Light: Tukey post

hoc: $^{q=0}D_{SB-HL}$ vs. $^{q=0}D_{SB-LL}$: t_{97} = 3.59, P < 0.001; mean $^{q=0}D$ ± SE: 169 ± 11 and 114 ± 8,

respectively), while bacterial richness of LB medaka did not change with light variation (Fig.

4b; $^{q\,=\,0}\mathrm{D}_{LB\text{-}HL} = 141 \pm 14$, $^{q\,=\,0}\mathrm{D}_{LB\text{-}LL} = 145 \pm 10$). However, LB medaka seemed to have a

higher bacterial species richness than SB medaka, but only in the low-light treatment (Line ×

Light: Tukey post hoc: $^{q\,=\,0}\mathrm{D}_{LB\text{-}LL}$ vs. $^{q\,=\,0}\mathrm{D}_{SB\text{-}LL}$: $t_{97} = 1.97$, P = 0.052). Other metrics of gut

microbiome diversity (when $q = 1$ or 2) were not influenced by the interaction between Line

and environmental conditions (Table 1).

Figure 4 (a) NMDS ordination (NMDS stress = 0.15) of variation in bacterial community
composition of SB (blue) and LB (orange) fish in the different light conditions. Data represent
ordination based on unweighted (d;) UniFrac distances among the 103 fish individuals. **(b)**
Raincloud plot showing the light-intensity effect of Line on bacterial richness ($^{q\,=\,0}\mathrm{D}$). Dots
represent the fish (n = 103), boxplots and half violin plots illustrate the probability density of
the data.

Lack of correlation between microbiome and growth-related traits

No significant correlation was observed between gut microbiome diversity (estimated using

the first three Hill numbers) and medaka's body condition, somatic growth, and final standard

length (Spearman correlations: adjusted $P > 0.876$ for all; Appendix 3). Correlations between

medaka's traits and gut microbial Families were not detected (Fig. 5).

392
393 **Figure 5** Correlation circle plot from the CCA applied between medaka's traits (grey circles)
394 and gut microbial composition (Family level, black squares). The farther from the center a
395 bacterial family or traits is, the greater the association with the component. Variables
396 projected in the same direction of the plot are positively correlated.
397

Discussion

Due to its pivotal role for host fitness and health, there has been a growing research interest in

the factors driving gut microbiome variation in animal species. Yet, studies focusing on the

genotype-by-environment effects on the gut microbiome of fish remain limited (Sevellec et

al. 2014, Piazzon et al. 2020). Using size-selected medaka lines (large-breeder LB and small-

breeder SB) in a pond mesocosm experiment, we found that the composition of the core gut

microbial community differed between the two lines. In addition, differences in rare taxa

(unweighted UniFrac) between the lines were modulated by light intensity, with more

dispersion for LB medaka than SB medaka in the high-light treatment and the opposite

pattern in the low-light treatment. The microbiome richness of SB medaka was influenced by

light intensity, while that of LB remained unchanged regardless of the environmental

conditions. Together, this is consistent with our prediction that evolutionary changes due to

size-selective harvesting have the potential to shape the gut microbiome assemblage within

harvested populations. Our results also suggest that the interaction between the genetic

background of medaka (i.e. the selected line) and the environmental conditions could be

important, as previously reported in the gut microbiome communities of two guppy ecotypes,

sampled across different streams (Sullam et al. 2015). However, contrary to our prediction,

variation in microbiome diversity or composition was not associated with any of the

measured growth-related traits of adult medaka.

Our findings confirm observations from literature showing that genotype could drive

gut microbiome composition among fish groups (Sevellec et al. 2018, Small et al. 2019,

Smith et al. 2015). In our case, the largest difference in relative read abundance was found for

the *Aeromonadaceae* family which was almost 3 times more abundant in LB than in SB

medaka (28.9% and 10.1%, respectively). Our previous findings from the same pond

experiment suggested that adult and juvenile LB medaka foraged more on benthic prey

hidden in the sediments than the SB medaka (Evangelista et al. 2021). Although further

investigation is required to back-up this hypothesis, increased relative abundance of

Aeromonadaceae in LB might reflect distinct foraging strategies between the two lines as

Aeromonadaceae are known to be facultative aerobes and are mainly found in anoxic

sediments (Tomás 2012, Laviad and Halpern 2016). Additionally, some *Aeromonas* display

cellulolytic activity, which can be useful for the digestion of plant-based diet (Li et al. 2016).

One could hypothesize that a higher proportion of *Aeromonadaceae* in LB medaka could be

associated with a more omnivorous feeding habit compared to SB medaka (Liu et al. 2016).

In addition, *Neisseriaceae* found in greater quantities in LB was also found to be more

abundant in rainbow trout fed with fish-based meal as to opposed to those fed with insect

meal (Terova et al. 2021), potentially suggesting that LB fish predate more on fish larvae

than SB medaka. Interestingly, *Rhodobacteraceae*, which were found in higher proportion in

SB medaka here, has been associated with higher levels of environmentally induced stress in

the host (Pootakham et al. 2019 and references herein). This might add another line of

evidence about the lesser capacity of SB medaka to cope with a less favourable surrounding

environment, as previously reported (Evangelista et al. 2020, Evangelista et al. 2021).

Altogether, our results suggest that even if the gut microbiome composition between the two

lines differs, the mere description of microbiome diversity and composition is not sufficient

to precisely identify the underlying causes of gut microbiome variation between SB and SL

medaka. In fact, more targeted diet manipulation experiments between lines (e.g. Sullam et

al. 2015) would be required in order to clearly identify whether adaptation to size selection

could directly affect the gut microbiome, or indirectly through changes in diet.

Under low-light intensity, the gut microbiome of SB medaka showed a 34% decrease

of bacterial richness compared to the high-light intensity treatment, and also presented a

somewhat lower richness than in LB medaka, though this difference was not significant (P =

0.052). This suggests that the gut microbiome diversity of fish selected for earlier maturation

and slower growth rate (as is often the case with size-selective harvesting) can be reduced

under sub-optimal environmental conditions. The underlying mechanisms of such genotype-

by-environment effects are hard to pinpoint and can be the results of many interacting factors

– light-induced changes in the SB medaka behaviour or physiology, combined with changes

in diet or in the bacterial composition of water, ultimately altering the gut microbiome.

However, as we did not sequence the microbiome from the water used in the experimental

ponds, we are not able to assess the extent to which variation in water composition or even

potential co-amplification of bacterial taxa from the environment (Talwar et al. 2018) could

influence our results. In term of bias, previous studies suggest that fish gut microbiome

composition is not influenced by the bacterial composition of the surrounding waters

(Schmidt et al. 2015, Wang et al. 2018), thus implying water should not be a confounding

factor in the context of our study. The present study focused on the deterministic processes

that influenced the gut microbiome composition of medaka, but it is important to note that

stochastic processes may also explained substantial variability in the bacterial composition of

medaka (Jones et al. 2022).

Whether high microbial diversity matters for the host remains a central question in

microbiome studies. For instance, Bolnick et al. (2014b) found a positive effect of the gut

microbial diversity on the body condition of laboratory-reared stickleback (*Gasterosteus*

aculeatus), but no such effect on condition of wild stickleback. In our experiment, gut

microbiome diversity was not associated with growth-related traits of adult medaka, perhaps

because low diversity does not entail the loss of essential microbiome-mediated functions.

Therefore, the lack of associations could simply reflect our yet limited understanding about

the taxonomic identity and functional role of gut bacteria in non-model organisms. Thus,

changes in bacterial diversity might be associated with either positive or negative impact for

the host, according to the degree of decoupling between taxonomic identity, functional role

and the environmental context (Bolnick et al. 2014b). Overall, our results highlight the fact

that our perception about gut microbiome benefits is also probably biased by data based on a

very limited range of species (Hammer et al. 2019). Nonetheless, variation in the microbiome

can impact the digestive capacity and body condition, and this diversity may act as an

underlying mechanism for phenotypic plasticity of the host (Alberdi et al. 2016). How

changes in bacterial diversity translate into functional changes will require further

investigation.

Our study reveals that the gut microbiome of fish can be influenced by interactions

between their genetic background and the environment. Studying genotype-by-environment

interactions on the gut microbiome may bring new perspectives into the role of microbiomes

in eco-evolutionary dynamics, as changes in gut microbial communities could potentially

reverberate into changes in ecosystem functioning and services (Graham et al. 2016, Dutton

et al. 2021), including fisheries productivity (Gallo et al. 2020, Diwan et al. 2021). As the demand for fish for human consumption is increasing, we also claim that more research is needed to enhance our understanding of the possible effects of fisheries-induced evolution on the gut microbiome. Comparison of the gut microbiome of fish in relation to different management strategies (e.g. balanced versus size-selective fishing versus protected areas) may reveal important mechanisms influencing populations' adaptability and resilience, and thus help restoring highly impacted fish stocks (Gallo et al. 2020). It is also important for future studies to reveal the functional consequence of changes in gut microbiome (Ghanbari et al. 2015, Tarnecki et al. 2017), especially functions directly responsible for fish behaviour and fitness so that the target preservation of highly beneficial gut microbiomes within harvested populations could be incorporated as part of more sustainable fisheries practices.

Acknowledgments

We are grateful to Clémentine Renneville and Arnaud Le Rouzic for initiating the medaka lines; David Carmignac and Romain Péronnet for their help maintaining the fish; Julia Dupeu, Anders Herland and Jacques Meriguet for field assistance; the platform PLANAQUA and the CEREEP Ecotron Ile-De-France for the access to the experimental facilities; and Natacha Nikolic for valuable comments on a previous version of this manuscript. We also thank Konstantinos Kormas, Marco Basili, Laetitia Wilkins and one anonymous reviewer for their insightful comments that improved the manuscript.

Data, script, code, and supplementary information availability

Data and R codes that support the findings of this study are hosted in the Figshare repository (https://figshare.com/s/d5235f25f3d7b15a0e47). SRA accession for sequences data is PRJNA929943

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of

interest in relation to the content of the article. The authors declare the non-financial conflict

of interest: SK is a recommender for PCI Ecology.

Funding

This work was supported by The Research Council of Norway (projects 251307/F20 and

255601/E40) and its mobility program (projects 272354 to CE and 268218/MO to BDP). EE

was supported by IDEX SUPER (project Convergences J14U257 MADREPOP) and by

Rennes Métropole (AIS 18C0356). The experiments realized in the CEREEP Ecotron Ile-De-

France benefited from the support received by an "Investissements d'Avenir" program from

the Agence Nationale de la Recherche (ANR-10-EQPX-13-01 Planaqua and ANR-11-INBS-

0001 AnaEE France).

Ethics

The experiment was approved by the Darwin Ethical committee (case file #Ce5/2010/041)

from the French Ministry of Education, Higher Education and Research.

Author's contributions

CE conceived and coordinated the study with input from SK, EM and PT; CE, JD and BDP

collected the samples; SK and EM carried out the molecular analyses; PT carried out the

bioinformatic analyses; CE analyzed the data and wrote the initial draft of the manuscript

535 with input from SK, PT and EM; All authors contributed to revisions and approved the final

536 version of the manuscript.

538 **References**
539

Alberdi A., Aizpurua O., Bohmann K., Zepeda-Mendoza M.L. & Gilbert M.T.P. (2016). Do
vertebrate gut metagenomes confer rapid ecological adaptation? *Trends in Ecology &*
Evolution **31**, 689–699. https://doi.org/10.1016/j.tree.2016.06.008
Alberdi A. & Gilbert M.T.P. (2019a). A guide to the application of Hill numbers to
DNA☐based diversity analyses. *Molecular Ecology Resources* **19**, 804–817.
https://doi.org/10.1111/1755-0998.13014
Alberdi A, & Gilbert M.T.P. (2019b). hilldiv: an R package for the integral analysis of
diversity based on Hill numbers. bioRxiv: 545665.
Anderson MJ, Walsh DCI (2013) PERMANOVA, ANOSIM, and the Mantel test in the face
of heterogeneous dispersions: what null hypothesis are you testing? *Ecological*
Monographs **83**, 557–574. https://doi.org/10.1890/12-2010.1
Bestion E., Jacob S., Zinger L., Di Gesu L., Richard M., White J., *et al.* (2017). Climate
warming reduces gut microbiota diversity in a vertebrate ectotherm. *Nature Ecology &*
Evolution **1**, 0161. https://doi.org/10.1038/s41559-017-0161
Bolnick DI, Snowberg LK, Caporaso JG, et al (2014a) Major Histocompatibility Complex
class II b polymorphism influences gut microbiota composition and diversity. *Molecular*
Ecology **23,** 4831–4845. https://doi.org/10.1111/mec.12846
Bolnick D.I., Snowberg L.K., Hirsch P.E., Lauber C.L., Knight R., Caporaso J.G., *et al.*
(2014b). Individuals' diet diversity influences gut microbial diversity in two freshwater
fish (threespine stickleback and eurasian perch). *Ecology Letters* **17**, 979–987.
https://doi.org/10.1111/ele.12301
Bokulich N.A., Subramanian S., Faith J.J., Gevers D., Gordon J.I., Knight R., *et al.* (2013).
Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing.
Nature Methods **10**, 57–59. https ://doi.org/10.1038/nmeth.2276
Bouffet☐Halle A., Mériguet J., Carmignac D., Agostini S., Millot A., Perret S., *et al.* (2021).
Density☐dependent natural selection mediates harvest☐induced trait changes. *Ecology*
Letters, ele.13677. https://doi.org/10.1111/ele.13677
Callahan B., McMurdie P., Rosen M., Han A.W., Johnson A.Jo.A & Holmes S.P. (2016)
DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*
13, 581–583. https://doi.org/10.1038/nmeth.3869
Chiu C.-H. & Chao A. (2014). Distance-based functional diversity measures and their
decomposition: a framework based on Hill numbers. *PloS ONE* **9**, e100014.
https://doi.org/10.1371/journal.pone.0100014
Conover D.O. & Munch S.B. (2002) Sustaining fisheries yields over evolutionary time
scales. *Science* **297**, 94–96. https://doi.org/10.1126/science.1074085

Darimont CT, Carlson SM, Kinnison MT, et al (2009) Human predators outpace other agents
of trait change in the wild: Fig. 1. *Proceedings of the National Academy of Sciences* **106**,
952–954. https://doi.org/10.1073/pnas.0809235106

Diaz Pauli B., Garric S., Evangelista C., Vøllestad L.A. & Edeline E. (2019). Selection for
small body size favours contrasting sex-specific life histories, boldness and feeding in
medaka, Oryzias latipes. *BMC Evolutionary Biology* **19**, 127.
https://doi.org/10.1186/s12862-019-1460-x

Diwan A.D., Harke S.N., Gopalkrishna & Panche A.N. (2022). Aquaculture industry
prospective from gut microbiome of fish and shellfish: An overview. *Journal of Animal*
Physiology and Animal Nutrition **106**, 441–469. https://doi.org/10.1111/jpn.13619

Dutton C.L., Subalusky A.L., Sanchez A., Estrela S., Lu N., Hamilton S.K., *et al.* (2021). The
meta-gut: community coalescence of animal gut and environmental microbiomes.
Scientific Reports **11**, 23117. https://doi.org/10.1038/s41598-021-02349-1

Edeline E., Terao O. & Naruse K. (2016). Empirical evidence for competition-driven
semelparity in wild medaka. *Population Ecology* **58**, 371–383.
https://doi.org/10.1007/s10144-016-0551-4

Eckert E.M., Anicic N. & Fontaneto D. (2021). Freshwater zooplankton microbiome
composition is highly flexible and strongly influenced by the environment. *Molecular*
Ecology **30**, 1545–1558. https://doi.org/10.1111/mec.15815

Evangelista C., Dupeu J., Sandkjenn J., Pauli B.D., Herland A., Meriguet J., *et al.* (2021).
Ecological ramifications of adaptation to size-selective mortality. *Royal Society Open*
Science **8**, 210842. https://doi.org/10.1098/rsos.210842

Evangelista C., Vøllestad L.A., Diaz Pauli B. & Edeline E. (2020). Density-dependent
consequences of size-selective induced life-history changes to population fitness in
medaka (*Oryzias latipes*). *Canadian Journal of Fisheries and Aquatic Sciences*, 1–8.
https://doi.org/10.1139/cjfas-2019-0406

Fadrosh D.W., Ma B., Gajer P., Sengamalay N., Ott S., Brotman R.M., *et al.* (2014). An
improved dual-indexing approach for multiplexed 16S rRNA gene sequencing on the
Illumina MiSeq platform. *Microbiome* **2**, 6. https://doi.org/10.1186/2049-2618-2-6

Fouladi F., Bailey M.J., Patterson W.B., Sioda M., Blakley I.C., Fodor A.A., *et al.* (2020).
Air pollution exposure is associated with the gut microbiome as revealed by shotgun
metagenomic sequencing. *Environment International* **138**, 105604.
https://doi.org/10.1016/j.envint.2020.105604

Frank KT, Petrie B, Choi JS, Leggett WC (2005) Trophic cascades in a formerly cod-
dominated ecosystem. *Science* **308,** 1621–1623. https ://doi.org/10.1126/science.1113075

Gallo B.D., Farrell J.M. & Leydet B.F. Fish gut microbiome: a primer to an emerging
discipline in the fisheries sciences. *Fisheries*, **45**, 271–282.
https://doi.org/10.1002/fsh.10379

Ghanbari M, Kneifel W, Domig KJ (2015) A new view of the fish gut microbiome: advances
from next-generation sequencing. *Aquaculture* **448,** 464–475.
https://doi.org/10.1016/j.aquaculture.2015.06.033

Graham E.B., Knelman J.E., Schindlbacher A., Siciliano S., Breulmann M., Yannarell A., *et*
al. (2016). Microbes as engines of ecosystem function: when does community structure

enhance predictions of ecosystem processes? *Frontiers in Microbiology* **7**, 1–10.
https://doi.org/10.3389/fmicb.2016.00214

Hammer T.J., Sanders J.G. & Fierer N. (2019). Not all animals need a microbiome. *FEMS*
Microbiology Letters **366**, fnz117. https://doi.org/10.1093/femsle/fnz117

Hanning I. & Diaz-Sanchez S. (2015). The functionality of the gastrointestinal microbiome in
non-human animals. *Microbiome* **3**, 51. https://doi.org/10.1186/s40168-015-0113-6

Heino M, Díaz Pauli B, Dieckmann U (2015) Fisheries-induced evolution. *Annual Review of*
Ecology, Evolution, and Systematics **46**, 461–480. https://doi.org/10.1146/annurev-
ecolsys-112414-054339

Hočevar S. & Kuparinen A. (2021). Marine food web perspective to fisheries⬜induced
evolution. *Evolutionary Applications* **14**, 2378–2391. https://doi.org/10.1111/eva.13259

Jones E.W., Carlson J.M. & Ludington W.B. (2022). Stochastic microbiome assembly
depends on context. *Proceedings of the National Academy of Science* **119**
(7)e2115877119. https://doi.org/10.1073/pnas.211587711

Laviad S. & Halpern M. (2016). Chironomids' relationship with *Aeromonas* species.
Frontiers in Microbiology **7**. https://doi.org/10.3389/fmicb.2016.00736

Lenth R.V. (2021). emmeans: estimated marginal means, aka least-squares means. R package
version 1.6.3. https://CRAN.R-project.org/package=emmeans

Le Rouzic A., Renneville C., Millot A., Agostini S., Carmignac D. & Édeline É. (2020).
Unidirectional response to bidirectional selection on body size II. Quantitative genetics.
Ecology and Evolution, ece3.6783. https://doi.org/10.1002/ece3.6783

Li H., Wu S., Wirth S., Hao Y., Wang W., Zou H., *et al.* (2016). Diversity and activity of
cellulolytic bacteria, isolated from the gut contents of grass carp (*Ctenopharyngodon*
idellus) (Valenciennes) fed on Sudan grass (*Sorghum sudanense*) or artificial feedstuffs.
Aquaculture Research **47**, 153–164. https://doi.org/10.1111/are.12478

Liu C., Cui Y., Li X. & Yao M. (2021). *microeco*: an R package for data mining in microbial
community ecology. *FEMS Microbiology Ecology* **97**, fiaa255.
https://doi.org/10.1093/femsec/fiaa255

Liu H., Guo X., Gooneratne R., Lai R., Zeng C., Zhan F., *et al.* (2016). The gut microbiome
and degradation enzyme activity of wild freshwater fishes influenced by their trophic
levels. *Scientific Reports* **6**, 24340. https://doi.org/10.1038/srep24340

McMurdie P.J. & Holmes S. (2013). phyloseq: An R package for reproducible interactive
analysis and graphics of microbiome census data. *PLoS ONE* **8**, e61217.
https://doi.org/10.1371/journal.pone.0061217

Moran N.A., Ochman H. & Hammer T.J. (2019). Evolutionary and ecological consequences
of gut microbial communities. *Annual Review of Ecology, Evolution, and Systematics* **50**,
451–475. https://doi.org/10.1146/annurev-ecolsys-110617-062453

Navarrete P., Magne F., Araneda C., Fuentes P., Barros L., Opazo R., *et al.* (2012). PCR-
TTGE analysis of 16S rRNA from rainbow trout (*Oncorhynchus mykiss*) gut microbiota
reveals host-specific communities of active bacteria. *PLoS ONE* **7**, e31335.
https://doi.org/10.1371/journal.pone.0031335

Nusslé S, Hendry AP, Carlson SM (2016) When should harvest evolution matter to
population dynamics? *Trends in Ecology & Evolution* **31**, 500–502.
https://doi.org/10.1016/j.tree.2016.04.001

Oksanen J., Blanchet F.G., Friendly M., Kindt R, Legendre P., McGlinn D., *et al.* (2020). vegan: community ecology package. R package version 2.5-7. https://CRAN.R-project.org/package=vegan

Piazzon M.C., Naya-Català F., Perera E., Palenzuela O., Sitjà-Bobadilla A. & Pérez-Sánchez J. (2020). Genetic selection for growth drives differences in intestinal microbiota composition and parasite disease resistance in gilthead sea bream. *Microbiome* **8**, 168. https://doi.org/10.1186/s40168-020-00922-w

Pootakham W, Mhuantong W, Yoocha T, et al (2019) Heat☐induced shift in coral microbiome reveals several members of the Rhodobacteraceae family as indicator species for thermal stress in *Porites lutea*. *Microbiology Open* **8**, e935. https://doi.org/10.1002/mbo3.935

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reese A.T. & Dunn R.R. (2018). Drivers of microbiome biodiversity: a review of general rules, feces, and ignorance. *mBio* **9**, e01294–18. https://doi.org/10.1128/mBio.01294-18

Renneville C., Millot A., Agostini S., Carmignac D., Maugars G., Dufour S., *et al.* (2020). Unidirectional response to bidirectional selection on body size. I. Phenotypic, life☐history, and endocrine responses. *Ecology and Evolution* ece3.6713. https://doi.org/10.1002/ece3.6713

Revelle W. (2021). psych: Procedures for personality and psychological research, Northwestern University, Evanston, Illinois, USA, https://CRAN.R-project.org/package=psych Version = 2.1.9.

Rohart F, Gautier B, Singh A, Lê Cao K-A (2017) mixOmics: An R package for 'omics feature selection and multiple data integration. *PLoS Computational Biology* **13**, e1005752. https://doi.org/10.1371/journal.pcbi.1005752

Ruzzante D.E. & Doyle R.W. (1993). Evolution of social behaviour in a resource-rich, structured environment: selection experiments with medaka (*Oryzias latipes*). *Evolution* **47**, 456–470. https://doi.org/10.1111/j.1558-5646.1993.tb02106.x

Sanderson S., Beausoleil M., O'Dea R.E., Wood Z.T., Correa C., Frankel V., *et al.* (2022). The pace of modern life, revisited. *Molecular Ecology* **31**, 1028–1043. https://doi.org/10.1111/mec.16299

Schmidt V.T., Smith K.F., Melvin D.W. & Amaral-Zettler L.A. (2015). Community assembly of a euryhaline fish microbiome during salinity acclimation. *Molecular Ecology* **24**, 2537–2550. https://doi.org/10.1111/mec.13177

Sepulveda J. & Moeller A.H. (2020). The effects of temperature on animal gut microbiomes. *Frontiers in Microbiology* **11**, 384. https://doi.org/10.3389/fmicb.2020.00384

Sevellec M, Pavey SA, Boutin S, et al (2014) Microbiome investigation in the ecological speciation context of lake whitefish (*Coregonus clupeaformis*) using next-generation sequencing. *Journal of Evolutionary Biology* **27**, 1029–1046. https://doi.org/10.1111/jeb.12374

Sevellec M., Derome N. & Bernatchez L. (2018). Holobionts and ecological speciation: the intestinal microbiota of lake whitefish species pairs. *Microbiome* **6**, 47. https://doi.org/10.1186/s40168-018-0427-2

Small C.M., Currey M., Beck E.A., Bassham S. & Cresko W.A. (2019). Highly reproducible
16S sequencing facilitates measurement of host genetic influences on the stickleback gut
microbiome. *mSystems* **4**, e00331-19. https://doi.org/10.1128/mSystems.00331-19
Smith C., Snowberg L.K., Caporaso G.J., Knight R. & Bolnick D. (2015) Dietary input of
microbes and host genetic variation shape among-population differences in stickleback gut
microbiota. *The ISME Journal* **9**, 2515–2526. https://doi.org/10.1038/ismej.2015.64
Sonnenburg J.L. & Sonnenburg E.D. (2019). Vulnerability of the industrialized microbiota.
Science **366**, eaaw9255. https://doi.org/10.1126/science.aaw9255
Spor A., Koren O. & Ley R. (2011). Unravelling the effects of the environment and host
genotype on the gut microbiome. *Nature Reviews Microbiology* **9**, 279–290.
https://doi.org/10.1038/nrmicro2540
Stearns, S. 1992. The evolution of life histories. Oxford University Press, Oxford.
Sullam KE, Essinger SD, Lozupone CA, et al (2012) Environmental and ecological factors
that shape the gut bacterial communities of fish: a meta-analysis. *Molecular Ecology* **21**,
3363–3378. https://doi.org/10.1111/j.1365-294X.2012.05552.x
Sullam KE, Rubin BE, Dalton CM, et al (2015) Divergence across diet, time and populations
rules out parallel evolution in the gut microbiomes of Trinidadian guppies. *ISME Journal*
9, 1508–1522. https://doi.org/10.1038/ismej.2014.231
Talwar C., Nagar S., Lal R. & Negi R.K. (2018) Fish gut microbiome: current approaches
and   future   perspectives.   *Indian   Journal   of   Microbiology*   **58**,   397–414.
https://doi.org/10.1007/s12088-018-0760-y
Tarnecki A.M., Burgos F.A., Ray C.L. & Arias C.R. (2017). Fish intestinal microbiome:
diversity and symbiosis unravelled by metagenomics. *Journal of Applied Microbiology*,
123(1), 2–17. https://doi.org/10.1111/jam.13415
Terova G, Gini E, Gasco L, et al (2021) Effects of full replacement of dietary fishmeal with
insect meal from *Tenebrio molitor* on rainbow trout gut and skin microbiota. *Journal of
Animal Science and Biotechnology* **12,** 30. https://doi.org/10.1186/s40104-021-00551-9
Teyssier A., Rouffaer L.O., Saleh Hudin N., Strubbe D., Matthysen E., Lens L., *et al.* (2018).
Inside the guts of the city: urban-induced alterations of the gut microbiota in a wild
passerine.   *Science   of   The   Total   Environment*   **612**,   1276–1286.
https://doi.org/10.1016/j.scitotenv.2017.09.035
Tomás J.M. (2012). The Main *Aeromonas* Pathogenic Factors. *ISRN Microbiology* **2012**, 1–
22. https://doi.org/10.5402/2012/256261
Trosvik P., Rueness E.K., de Muinck E.J., Moges A. & Mekonnen A. (2018). Ecological
plasticity in the gastrointestinal microbiomes of Ethiopian *Chlorocebus* monkeys.
Scientific Reports **8**, 20. https://doi.org/10.1038/s41598-017-18435-2
Uusi-Heikkilä S, Whiteley AR, Kuparinen A, et al (2015) The evolutionary legacy of size-
selective harvesting extends from genes to populations. *Evolutionary Applications* **8**, 597–
620. https://doi.org/10.1111/eva.12268
Uusi-Heikkilä S., Sävilammi T., Leder E., Arlinghaus R. & Primmer C.R. (2017). Rapid,
broad-scale gene expression evolution in experimentally harvested fish populations.
Molecular Ecology **26**, 3954–3967. https://doi.org/10.1111/mec.14179
van Wijk S.J., Taylor M.I., Creer S., Dreyer C., Rodrigues F.M., Ramnarine I.W., *et al.*
(2013). Experimental harvesting of fish populations drives genetically based shifts in body

size and maturation. *Frontiers in Ecology and the Environment* **11**, 181–187.
https://doi.org/10.1890/120229

Varg J.E., Kunce W., Outomuro D., Svanbäck R. & Johansson F. (2021). Single and
combined effects of microplastics, pyrethroid and food resources on the life-history traits
and microbiome of *Chironomus riparius*. *Environmental Pollution* **289**, 117848.
https://doi.org/10.1016/j.envpol.2021.117848

Wang Q., Garrity G.M., Tiedje J.M. & Cole J.R. (2007). Naïve Bayesian Classifier for Rapid
Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Applied and*
Environmental Microbiology **73**, 5261–5267. https://doi.org/10.1128/AEM.00062-07

Wang C., Sun G., Li S., Li X. & Liu Y. (2018). Intestinal microbiota of healthy and
unhealthy Atlantic salmon *Salmo salar L.* in a recirculating aquaculture system. *Journal of*
Oceanology and Limnology **36**, 414–426. https://doi.org/10.1007/s00343-017-6203-5

Appendix 1: Rarefaction curves

Figure S1 Rarefaction curves showing the number of ASV with increasing number of reads (Sequence sample size) **(a)** before and **(b)** after

standardization to the median sequencing depth. Fish from the Large-breeder (LB) line are in orange (n = 52) and fish from the Small-breeder

(SB) line are in blue (n = 51).

Appendix 2: Relative abundance of bacterial phyla in gut samples

Figure S2 Gut microbial taxonomic composition (Phylum level) of fish according to its

selective background (Large-breeder line, LB; Small-breeder line SB).

 Appendix 3: Correlations between medaka growth-related traits and gut microbiome

diversity

Figure S3 Correlations between gut microbial diversity estimated using the first three Hill numbers (^{q} D; species richness, exponential Shannon index and Reciprocal Simpson index) and **(a-c)** body condition estimated as the residuals of the relationship between $\log_{10}W_f$ and $\log_{10}SL_f$, **(d-f)** somatic growth rate [mm month^{-1}]) and **(g-i)** final standard length (mm) of fish from the Large-breeder (LB; orange dots, n = 52) and Small-breeder (SB; blue dots, n = 51) lines. None of the correlations were significant.