Land-cover change in Cuba may favor biodiversity: An example using

Omphalea* (Angiosperma: Euphorbiaceae) and *Urania boisduvalii

(Lepidoptera: Uraniidae)

Land-cover change in Cuba may favor biodiversity: An example using

Omphalea* and *Urania boisduvalii

Claudia Nuñez-Penichet[1*], Juan Maita-Chamba[2], and Jorge Soberón[1]

[1]Biodiversity Institute and Department of Ecology & Evolutionary Biology, University of

Kansas.

[2]Carrera de Ingeniería Forestal, Centro de Investigaciones Tropicales del Ambiente y

Biodiversidad, Universidad Nacional de Loja, Loja 110111, Ecuador

[*]Corresponding Author: Claudia Nuñez-Penichet

Biodiversity Institute and Department of Ecology & Evolutionary Biology, University of

Kansas, 1345 Jayhawk Blvd., Lawrence, Kansas 66045, USA. claununez199o@gmail.com

ORCID:

CNP: https://orcid.org/0000-0001-7442-8593

JMC: https://orcid.org/0000-0001-5058-8613

JS: https://orcid.org/0000-0003-2160-4148

ABSTRACT

Changes in land cover directly affect biodiversity. Here we assessed land-cover change in

Cuba in the past 35 years and analyzed how this change may affect the distribution of

Omphalea plants and *Urania boisduvalii* moths. We analyzed the vegetation cover of the

Cuban archipelago between 1985 and 2020. We used Google Earth Engine to classify two

satellite image compositions into seven cover types: forest and shrubs, mangrove, soil

without vegetation cover, wetlands, pine forest, agriculture, and water bodies. We considered

three different areas for quantifications of land-cover change: (1) protected areas, (2) areas of

potential distribution of *Omphalea*, and (3) areas of potential distribution of the plant within

the protected areas. We found that the category of "forest and shrubs" have increased

significantly in Cuba in the past 35 years and that most of the gained forest and shrub areas

were agricultural land in the past. This same pattern was observed in the areas of potential

distribution of *Omphalea*; whereas almost all cover types were mostly stables inside the

protected areas. The transformation of agricultural areas into forest and shrubs could

represent an interesting opportunity for biodiversity conservation in Cuba. Other detailed

studies about biodiversity composition in areas of forest and shrubs gain would greatly

benefit our understanding on the value of such areas for conservation.

Keywords:

afforestation, conservation, global change, Google Earth Engine, land-use change, remote

sensing

Introduction

Landscape degradation is one of the most important anthropogenic causes of

biodiversity loss, driven mostly by changes in land use [1]. This is a problem for most

species, but it may be especially important for those with a restricted distribution or those that

depend on a specific climate or habitat for survival [2]. How habitat modifications are

affecting species has been the focus of attention in several studies [3]. Those studies have

been mainly focused on plants, birds, and mammals [4–6]. However, other groups like

insects, are underrepresented [7], in spite of documented examples of extinctions due to

habitat loss [8].

The neotropical genus *Urania* (Uraniidae) includes four species of diurnal moths, *U.*

sloanus (endemic to Jamaica, presumed extinct; [9]), *U. fulgens* (from Mexico to northern

Colombia), *U. leilus* (from southern Colombia to Bolivia), and *U. boisduvalii* (endemic to

Cuba; [10]). These moths feed, during their larvae stages, exclusively on *Omphalea*

(*Euphorbiaceae*) plants [9,11–13], which produce secondary metabolites as a defense

mechanism against *Urania* herbivory [9]. These secondary compounds make the plants toxic

for the moths, forcing them to move to a different host-plant population [14], therefore,

Urania needs to have several patches of *Omphalea* available to guarantee its survival.

Previous studies have also suggested that patches of *Omphalea* need to be structurally

complex to support the moths' larvae, as different larval stages use plants with different

levels of maturity [15].

In Cuba, there are three species of *Omphalea* plants [9] mainly distributed in coastal

and other karstic zones. *Omphalea trichotoma* (distributed in western and eastern Cuba) and

O. hypoleuca (reported only from Viñales, Pinar del Río) are endemic [16], while *O. diandra*

is widely distributed in the neotropics [9]. In this archipelago, these plants are present in areas

that have been under pressure from the development of infrastructure for tourism and oil

extraction activities [17] enhancing the importance of studying their populations and how

they may be affected.

Given the direct dependance on host plant availability, *Urania* moths appear to be

highly sensitive to changes in land use [9]. For instance, *U. sloanus* is now considered extinct

due to habitat degradation [9]. However, no study has explored how distributional areas of

these moths and plants are being affected over time due to land-cover change. Here, we

aimed (1) to evaluate the change in land use in Cuba in the past 35 years, and (2) to analyze

how these land use changes may affect the distribution of *Omphalea* plants and *Urania*

boisduvalii moths.

Methods

To evaluate land use change in Cuba (S1 Appendix 1), we analyzed vegetation cover

of this archipelago between 1985 and 2020 using Landsat images and Google Earth Engine

[18] for geospatial processing. Our study area was covered by sixteen scenes (paths: 10 to 17

and rows: 44 to 46) of the Landsat 5 ETM sensor (1985) and the Landsat 8 OLI/TIRS sensor

(2020), with images at 30 meter resolution. To minimize temporal, spatial, and spectral

varying scattering and absorbing effects of atmospheric gasses and aerosols, we used the

collection 2 of Land Surface Reflectance [19]. Cloud-free images of Cuba were produced by

using those with cloud cover ≤50% and a temporal filter including the years 1984-1988 for

the period 1985 and the years 2020-2021 for 2020. Cloud and cloud shadows were masked

using pixel quality attributes generated from the C Function of Mask (CFMASK) algorithm

(QA_PIXEL Bitmask) for both sensors [20,21].

The resulting images (1985 and 2020) were classified into seven cover types to
represent forest and shrubs (considered as suitable) and, mangrove, soil without vegetation
cover, wetlands, pine forest, agriculture, and water bodies (considered as unsuitable areas for
Omphalea plants; [22]). The classification of cover types was done by selecting regions of
interest (ROI) for each of the categories on the Google Earth Engine. The number of ROIs
selected for each category varied to represent how diverse they were (300 for forest and
shrubs, 100 for mangrove, 50 for soil without vegetation cover, 100 for wetlands, 100 for
pine forest, 100 for agriculture, and 100 for water bodies). All ROIs were selected manually
based on visual interpretation of the respective Landsat images from Google Earth and were
divided randomly into training (70%) and testing (30%).

We selected Random Forest as the classification algorithm as it has been described to
be good at handling outliers and noisy datasets [23], have good performance with complex
datasets [24] and higher accuracy than other algorithms [25,26], and its high processing speed
[23]. The classified images were post-processed in QGIS [27] to correct misclassified pixels
using a raster calculator and elevational considerations (S1 Appendix 2). We assessed the
accuracy of classifications via: 1) overall accuracy (OA) which is calculated dividing the total
number of correctly classified values by the total number of values [28,29], and 2) kappa
coefficient [30,31] for measuring the agreement between classification and truth values. A
kappa value of 1 represents perfect agreement, while a value of 0 represents no agreement.
These steps were done on the Google Earth Engine platform, using the resulting images from
the post-processing analyses and 100 new validation ROIs created randomly for each of the
considered cover types but water bodies in which we used only 50.

The resulting classified images for Cuba were masked to: (1) terrestrial protected
areas [32]; (2) potential distribution areas of *Omphalea* (from [22], hereafter OD; and (3)
potential distribution areas of the plant within protected areas. We quantified the percentage

of each of the seven cover types and their changes between periods for all Cuba and in each

of the masked areas (areas of interest). All these analyses were done in R [33] using the

packages rgdal [34] and raster [35]. All the code needed to reproduce these analyses is openly

available at https://github.com/claununez/Land-coverChangeCuba. All the data needed is

openly available at https://figshare.com/articles/dataset/Data_for_the_manuscript_Land-

cover_change_in_Cuba_may_favor_biodiversity_An_example_using_Omphalea_plants_and

_Urania_boisduvalii/21779129.

Results

We obtained an overall accuracy (OA) of 92.7% and 87.6% for 1985 and 2020

classifications, with a Kappa coefficient of 0.91 and 0.85, respectively. The values of OA for

each land cover type ranged from 80% (pine forest) to 100% (agriculture) in 1985, and from

71% (wetlands) to 99% (forest and shrubs) in 2020 (S1 Table).

We found that, in the past 35 years, forest and shrub areas have increased across the

entire Cuban territory (from 23.5% in 1985 to 38.89% in 2020; Table 1). Agricultural lands

and soils without vegetation cover, on the other hand, showed a reduction from 50.18% to

39.12% and from 6.03% to 1.85%, respectively (Table 1; Fig 1). The other four types of

cover considered (mangrove, wetland, pine forest, and water bodies) were present in similar

proportions in the two scenarios studied (Table 1). Most areas that changed from one cover

type to another were located inland, especially in areas of low elevation (Fig 1; S1 Appendix

1).

Table 1. Percentage of area classified as one of the seven classification types considered for

the years 1985 and 2020. *Omphalea* includes all the species of *Omphalea* genus distributed in

Cuba combined and the areas of potential distribution of *Omphalea* is referring to these

species' potential distribution in Cuba. The protected areas in Cuba are referring to the

terrestrial protected areas only.

Classification type	Cuba		Protected areas in Cuba		Areas of potential distribution of *Omphalea*		Areas of potential distribution of *Omphalea* inside protected areas	
	1985	2020	1985	2020	1985	2020	1985	2020
Forest and shrubs	23.50	38.89	33.80	36.42	57.67	73.26	71.24	74.96
Mangrove	4.03	4.16	13.05	12.82	1.66	1.91	3.47	3.86
Soil without vegetation cover	6.03	1.85	4.62	0.88	2.61	0.55	2.01	0.33
Wetland	5.84	6.08	17.13	19.30	2.68	2.02	4.34	3.86
Pine forest	2.28	1.24	2.43	2.07	5.60	3.44	6.42	5.46
Agriculture	50.18	39.12	7.92	6.86	26.92	15.64	8.04	6.70
Water bodies	8.14	8.65	21.05	21.65	2.86	3.19	4.48	4.82

145   Figure 1. Vegetation types in Cuba for the years 1985 and 2020.

147   In the areas of potential distribution of *Omphalea* plants, we found a similar pattern to

the one described above for Cuba (Table 1). Most of these changes were concentrated in

central Cuba, whereas stable areas were mostly in Guanahacabibes (westernmost part of

Cuba), the southern part of Isla de la Juventud, and highlands (S1 Appendix 1; S1 Figure).

The percentage of area of each cover type was similar between 1985 and 2020 inside the

protected areas in Cuba and in the OD that were inside the protected areas (Table 1, S2

Figure, S3 Figure). The only exception was soils without vegetation cover, which decreased

from 4.62% to 0.88% inside protected areas in Cuba, and from 2.01% to 0.33% in the OD

inside protected areas (Table 1, S2 Figure, S3 Figure).

We found that only 19.16% of Cuba classified as forest and shrubs in 2020, was

classified under the same category in 1985 (Table 2). The low stability of this cover type was

also found when quantifying in the other areas of interest, with 29.68% of the protected areas

in Cuba, 52.52% of the OD, and 64.49% of the OD inside protected areas found to be forest

and shrubs in both time periods analyzed (Table 2). Most of the forest and shrub areas gained

in Cuba in 2020 were agricultural lands in the past (Table 2). The proportion of change from

agricultural land to forest and shrubs was also high in the OD and the other areas of interest

(Table 2). The agricultural lands that changed to forest and shrubs by 2020 were distributed

in small patches across the Cuban archipelago and were less predominant in the eastern

region (Fig 2). The areas that changed to forest and shrubs in OD were also small patches, but

their spatial distribution helped increase the connectivity among larger stable forest and

shrubs areas (Fig 2, Table 2).

Table 2. Percentage of change between the classification types considered, during the period

1985 - 2020. *Omphalea* includes all the species of the genus *Omphalea* distributed in Cuba

combined and the areas of potential distribution of *Omphalea* is referring to these species'

potential distribution in Cuba.

Type of change	Cuba	Protected areas in Cuba	Areas of potential distribution of *Omphalea*	Areas of potential distribution of *Omphalea* inside protected areas
Stable forest and shrubs	19.16	29.68	52.52	64.49
Mangrove to forest and shrubs	0.19	0.65	0.30	0.80
Soil without vegetation cover to forest and shrubs	0.92	0.12	0.74	0.10
Wetlands to forest and shrubs	0.75	1.31	1.02	1.34
Pine forest to forest and shrubs	1.41	1.45	3.66	4.03
Agriculture to forest and shrubs	16.45	3.21	15.46	4.20
Water bodies to forest and shrubs	0.01	0.01	0.01	0.02
Stable mangrove	2.88	8.94	0.40	1.86
Forest and shrubs to mangrove	0.35	1.08	0.40	1.06
Soil without vegetation cover to mangrove	0.03	0.07	0.02	0.02
Wetlands to mangrove	0.70	2.28	0.38	0.77
Pine forest to mangrove	0.01	0.03	0.02	0.05
Agriculture to mangrove	0.06	0.09	0.05	0.05

Water bodies to mangrove	0.13	0.34	0.04	0.05
Stable soil without vegetation cover	1.05	0.58	0.35	0.23
Forest and shrubs to soil without vegetation cover	0.03	0.04	0.03	0.04
Mangrove to soil without vegetation cover	0.00	0.00	0.00	0.00
Wetlands to soil without vegetation cover	0.06	0.17	0.02	0.04
Pine forest to soil without vegetation cover	0.00	0.00	0.00	0.00
Agriculture to soil without vegetation cover	0.70	0.07	0.15	0.03
Water bodies to soil without vegetation cover	0.00	0.01	0.00	0.00
Stable wetlands	3.77	12.23	1.00	1.76
Forest and shrubs to wetlands	0.23	0.49	0.13	0.20
Mangrove to wetlands	0.80	2.98	0.28	0.65
Soil without vegetation cover to wetlands	0.81	2.67	0.44	0.99
Pine forest to wetlands	0.00	0.00	0.00	0.00
Agriculture to wetlands	0.29	0.27	0.09	0.07
Water bodies to wetlands	0.19	0.66	0.08	0.18
Stable pine forest	0.46	0.69	1.27	1.74
Forest and shrubs to pine forest	0.47	1.12	1.63	3.22
Mangrove to pine forest	0.00	0.00	0.00	0.00
Soil without vegetation cover to pine forest	0.01	0.00	0.00	0.00
Wetlands to pine forest	0.00	0.00	0.00	0.00
Agriculture to pine forest	0.30	0.26	0.55	0.50
Water bodies to pine forest	0.00	0.00	0.01	0.01
Stable agriculture	32.12	4.02	10.68	3.19
Forest and shrubs to agriculture	3.15	1.31	3.24	2.23
Mangrove to agriculture	0.05	0.13	0.04	0.07
Soil without vegetation cover to agriculture	3.01	0.47	0.93	0.32
Wetlands to agriculture	0.40	0.66	0.19	0.28
Pine forest to agriculture	0.35	0.26	0.65	0.59
Water bodies to agriculture	0.04	0.02	0.02	0.01
Stable water bodies	7.78	20.01	2.72	4.21
Forest and shrubs to water bodies	0.12	0.08	0.08	0.02
Mangrove to water bodies	0.11	0.36	0.04	0.08
Soil without vegetation cover to water bodies	0.20	0.71	0.14	0.35
Wetlands to water bodies	0.16	0.47	0.08	0.15
Pine forest to water bodies	0.05	0.00	0.03	0.00
Agriculture to water bodies	0.24	0.02	0.11	0.00

175 Figure 2. Forest and shrubs change. *Omphalea* includes all the species of *Omphalea* genus

176 distributed in Cuba combined and the areas of potential distribution of *Omphalea* is referring

177 to these species' potential distribution in Cuba.

Discussion

The general global trend in forest cover is towards loss [36]. However, we found that

in Cuba, forest and shrubs (i.e., suitable vegetation cover for *Omphalea*) have increased

considerably in the past 35 years, mostly replacing agricultural lands (Fig 1, Table 1). This

result may be due to the fact that in Cuba, before 1990, extensive areas were dedicated to

growing sugarcane, especially in the center of the island (S1 Appendix 1), but after 1990, the

intensity of sugarcane production decreased [37,38]. Our results concur with [39] who

reported that over half of agricultural land in Cuba is currently unused. Although the trend we

report is opposite to the general global pattern, it is an example of how land use change is a

complex phenomenon resulting from multiple causes, and difficult to predict [40].

Forest and shrub cover increases within areas of potential distribution of *Omphalea*

(Table 1; S1 Figure) may in fact represent an advantage for populations of these plants as

these areas are known to be climatically suitable [22]. Given the obligated relationship

between *Urania boisduvalii* and *Omphalea* plants, which also drives the moth's dispersal

patterns in the country (Nuñez-Penichet et al., in prep), these gained areas could indirectly

benefit the moth. This is because the stability and increment of this type of land cover could

be increasing connectivity among fragmented patches with plant populations (Fig 2; S1

Figure). In fact, several of the stable and gained shrub and forest patches are located in the

potential migratory paths of *U. boisduvalii* identified in a previous study [41].

On the other hand, the increment of forest and shrubs in the Cuban territory does not

necessarily imply an increase of *Omphalea* plants or an increment in favorable conditions for

Urania moths. Species distributions depend on the presence of suitable abiotic conditions,

accessibility, and biotic interactions [42]. In Cuba, a significant portion of unused agricultural

lands have been reported to be covered with invasive species (e.g., *Dichrostachys cinerea,*

[39,43]; and *Vachellia farnesiana*, [44], which causes a displacement of native plant

communities [45]. In the last decades, several studies have focused on testing the potential of

D. cinerea high-quality biomass to produce a sustainable biofuel [39,46] as an alternative

source of energy, and therefore little an effort is targeted to control this invasive plant. For

this reason, performing fieldwork in the gained forest and shrubs areas will be necessary to

assess whether they can actually benefit the conservation of *Omphalea* plants and *Urania*

moths.

When we focused on potential distribution areas for *Omphalea* inside protected areas,

we detected that stability was high for all cover types explored, except for soils without

vegetation cover (S2 Figure, S3 Figure). This concurs with what has been previously reported

for 10 National Parks in Cuba [47]. However, interpretations about the conservation status of

natural vegetation inside protected areas should be done cautiously, as given the goal of our

study, we included all types of forests and shrubs in a single category. Therefore, we are not

accounting for changes at a finer detail, and forest and shrub gains do not necessarily

represent positive rates of change for primary vegetation.

Here we are presenting results that suggest the distribution of *Omphalea* plants and

for *Urania* moths in Cuba may be increasing, since we find that the areas with suitable

conditions for these species have been increasing considerably. Moreover, our results

suggest that land-use changes in Cuba may favor not only *Urania* and *Omphalea* but other

species that live in forests and shrubs. However, the presence of invasive species in those

gained forest and shrubs areas should be monitored in the field as they may represent a

serious threat to native biodiversity.

Changes in land use in Cuba during the last 35 years may represent a significant

opportunity for biodiversity conservation and sustainable management of its natural

vegetation. However, further studies that allow understanding the community composition

and structure in such forests and shrubs are needed to assess the value of these lands for

biodiversity conservation in the country.

Acknowledgments

We thank Marlon E. Cobos for his useful comments on the manuscript.

References

1.   Leemans R, Groot RS de. Millennium ecosystem assessment: ecosystems and human

well-being: a framework for assessment. Washington/Covelo/London: Island Press;

2003. p. Available: https://library.wur.nl/WebQuery/wurpubs/326575

2.   Devictor V, Julliard R, Jiguet F. Distribution of specialist and generalist species along

spatial gradients of habitat disturbance and fragmentation. Oikos. 2008;117: 507–514.

doi:10.1111/j.0030-1299.2008.16215.x

3.   Brooks TM, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Rylands AB,

Konstant WR, et al. Habitat loss and extinction in the hotspots of biodiversity. Conserv

Biol. 2002;16: 909–923. doi:10.1046/j.1523-1739.2002.00530.x

4.   de Lima RF, Dallimer M, Atkinson PW, Barlow J. Biodiversity and land-use change:

understanding the complex responses of an endemic-rich bird assemblage. Divers

Distrib. 2013;19: 411–422. doi:10.1111/ddi.12015

5.   Powers RP, Jetz W. Global habitat loss and extinction risk of terrestrial vertebrates

under future land-use-change scenarios. Nat Clim Change. 2019;9: 323–329.

doi:10.1038/s41558-019-0406-z

6.   Zechmeister HG, Schmitzberger I, Steurer B, Peterseil J, Wrbka T. The influence of

land-use practices and economics on plant species richness in meadows. Biol Conserv.

2003;114: 165–177. doi:10.1016/S0006-3207(03)00020-X

7.   Basset Y, Lamarre GPA. Toward a world that values insects. Science. 2019.

doi:10.1126/science.aaw7071

8.   Dunn RR. Modern insect extinctions, the neglected majority. Conserv Biol. 2005;19:

1030–1036. doi:10.1111/j.1523-1739.2005.00078.x

9.   Lees DC, Smith NG. Foodplant associations of the Uraniinae (Uraniidae) and their

systematic, evolutionary, and ecological significance. J Lepidopterists Soc. 1991;45:

296–347.

10. Nazari V, Schmidt BC, Prosser S, Hebert PDN. Century-old DNA barcodes reveal

phylogenetic placement of the extinct Jamaican Sunset Moth, *Urania sloanus* Cramer

(Lepidoptera: Uraniidae). PLoS ONE. 2016;11: e0164405.

doi:10.1371/journal.pone.0164405

11. Nuñez-Penichet C, Barro A. Caterpillars of *Urania boisduvalii* (Uraniidae) feed on

Omphalea trichotoma (Euphorbiaceae) fruits in western Cuba. J Lepidopterists Soc.

2020;74: 124–126. doi:10.18473/lepi.74i2.a7

12.   Smith NG. *Urania fulgens* (Colipato verde, Green Urania). 631st–633rd ed. In: Janzen

DH, editor. Historia Natural de Costa Rica. 631st–633rd ed. Chicago, USA: The

University of Chicago Press; 1991. p. 822.

13.   Smith NG. Reproductive behavior and ecology of *Urania* (Lepidoptera: Uraniidae)

moths and of their larval food plants, *Omphalea* spp. (Euphorbiaceae). In: Quintero D,

Aiello A, editors. Insects of Panama and Mesoamerica Selected studies. Oxford: Oxford

University Press; 1992. pp. 576–593.

14.   Smith NG. Host plant toxicity and migration in the dayflying moth *Urania*. Fla

Entomol. 1983;66: 76–85. doi:10.2307/3494552

15.   Barro A. Historia natural y bioacústica de *Urania boisduvalii* (Uraniidae) y

Phoenicoprocta capistrata (Arctiidae). Tesis en opción al grado científico de Doctor en

Ciencias Biológicas, Universidad de La Habana. 2006.

16.   Greuter W, Rankin R. Espermatófitos de Cuba: inventario preliminar. Parte II:

Inventario. La Habana, Cuba: Botanischer Garten & Botanisches Museum Berlin-

Dahlem; 2016. doi:10.3372/cubalist.2016.2

17.   Camacho JA, Baena GSG, Leyva GP. Fortalecimiento de la gestión del desarrollo

integral y sostenible de la Península de Guanahacabibes, Reserva de la Biosfera, Pinar

del Río, Cuba. La Habana, Cuba: Editorial Científico-Técnica; 2010.

18.   Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. Google Earth

Engine: Planetary-scale geospatial analysis for everyone. Remote Sens Environ.

2017;202: 18–27. doi:10.1016/j.rse.2017.06.031

19. Vermote E, Justice C, Claverie M, Franch B. Preliminary analysis of the performance of

the Landsat 8/OLI land surface reflectance product. Remote Sens Environ. 2016;185:

46–56. doi:10.1016/j.rse.2016.04.008

20. Foga S, Scaramuzza PL, Guo S, Zhu Z, Dilley RD, Beckmann T, et al. Cloud detection

algorithm comparison and validation for operational Landsat data products. Remote

Sens Environ. 2017;194: 379–390. doi:10.1016/j.rse.2017.03.026

21. Zhu Z, Woodcock CE. Object-based cloud and cloud shadow detection in Landsat

imagery. Remote Sens Environ. 2012;118: 83–94. doi:10.1016/j.rse.2011.10.028

22. Nuñez-Penichet C, Cobos ME, Amaro JG, Barro A. Distribución potencial del género

Omphalea (Euphorbiaceae) en Cuba: aproximación a su distribución real. Rev Jardí-n

Botánico Nac. 2016;37: 165–175.

23. Jin Y, Liu X, Chen Y, Liang X. Land-cover mapping using random forest classification

and incorporating NDVI time-series and texture: a case study of central Shandong. Int J

Remote Sens. 2018;39: 8703–8723. doi:10.1080/01431161.2018.1490976

24. Belgiu M, Drăguţ L. Random forest in remote sensing: A review of applications and

future directions. ISPRS J Photogramm Remote Sens. 2016;114: 24–31.

doi:10.1016/j.isprsjprs.2016.01.011

25. Sheykhmousa M, Mahdianpari M, Ghanbari H, Mohammadimanesh F, Ghamisi P,

Homayouni S. Support vector machine versus random forest for remote sensing image

classification: A Meta-Analysis and Systematic Review. IEEE J Sel Top Appl Earth Obs

Remote Sens. 2020;13: 6308–6325. doi:10.1109/JSTARS.2020.3026724

26. Talukdar S, Singha P, Mahato S, Shahfahad, Pal S, Liou Y-A, et al. Land-use land-cover

classification by machine learning classifiers for satellite observations—A review.

Remote Sens. 2020;12: 1135. doi:10.3390/rs12071135

27. QGIS.org. Geographic information system. 2022. Available: https://qgis.org/es/site/

28. Congalton RG. A review of assessing the accuracy of classifications of remotely sensed

data. Remote Sens Environ. 1991;37: 35–46. doi:10.1016/0034-4257(91)90048-B

29. Story M, Congalton RG. Accuracy assessment: A user's perspective. Photogramm Eng

Remote Sens. 1986;52: 397–399.

30. Fielding AH, Bell JF. A review of methods for the assessment of prediction errors in

conservation presence/absence models. Environ Conserv. 1997;24: 38–49.

doi:10.1017/S0376892997000088

31. Fitzgerald RW, Lees BG. Assessing the classification accuracy of multisource remote

sensing data. Remote Sens Environ. 1994;47: 362–368. doi:10.1016/0034-

4257(94)90103-1

32. CNAP. Plan del sistema nacional de áreas protegidas 2014-2020. Ministerio de Ciencias

Tecnología y Medio Ambiente; 2013.

33. R Core Team. R: A language and environment for statistical computing. Vienna,

Austria.: R Foundation for Statistical Computing; 2020. Available: https://www.R-

project.org/

34. Bivand R, Keitt T, Rowlingson B. rgdal: Bindings for the "Geospatial" Data

Abstraction Library. 2020. Available: https://CRAN.R-project.org/package=rgdal

35.   Hijmans RJ. raster: Geographic Data Analysis and Modeling. 2020. Available:

https://CRAN.R-project.org/package=raster

36.   Keenan RJ, Reams GA, Achard F, de Freitas JV, Grainger A, Lindquist E. Dynamics of

global forest area: Results from the FAO Global Forest Resources Assessment 2015.

For Ecol Manag. 2015;352: 9–20. doi:10.1016/j.foreco.2015.06.014

37.   Machado MR. What's going on with land-use in Cuba?: Disparate data sets and the

Cuban agricultural transition. J Land Use Sci. 2018;13: 439–446.

doi:10.1080/1747423X.2018.1533044

38.   Suárez JA, Beatóna PA, Faxas RE, Pérez OM. Energy, environment and development in

Cuba. Renew Sustain Energy Rev. 2012;16: 2724–2731.

doi:https://doi.org/10.1016/j.rser.2012.02.023

39.   Sagastume AG, Cabello JJE, Huisingh D, Vandecasteele C, Hens L. The current

potential of low-carbon economy and biomass-based electricity in Cuba. The case of

sugarcane, energy cane and marabu (Dichrostachys cinerea) as biomass sources. J Clean

Prod. 2018;172: 2108–2122. doi:10.1016/j.jclepro.2017.11.209

40.   Lambin EF, Turner BL, Geist HJ, Agbola SB, Angelsen A, Bruce JW, et al. The causes

of land-use and land-cover change: moving beyond the myths. Glob Environ Change.

2001;11: 261–269. doi:10.1016/S0959-3780(01)00007-3

41.   Nuñez-Penichet C, Cobos ME, Barro A, Soberón J. Potential migratory routes of *Urania*

boisduvalii (Lepidoptera: Uraniidae) among host plant populations. Divers Distrib.

2019;25: 478–488. doi:10.1111/ddi.12881

42. Soberón J, Peterson AT. Interpretation of models of fundamental ecological niches and

species' distributional areas. Biodivers Inform. 2005;2. doi:10.17161/bi.v2i0.4

43. Sagastume AG, Cabello JJE, Hens L, Vandecasteele C. The biomass-based electricity

generation potential of the province of Cienfuegos, Cuba. Waste Biomass Valorization.

2016;8: 2075–2085.

44. Fernández M, Williams J, Figueroa G, Graddy-Lovelace G, Machado M, Vázquez L, et

al. New opportunities, new challenges: Harnessing Cuba's advances in agroecology and

sustainable agriculture in the context of changing relations with the United States.

Kapuscinski AR, Iles A, Locke KA, Nelson E, editors. Elem Sci Anthr. 2018;6: 76.

doi:10.1525/elementa.337

45. Ruiz JDS, Remond RN, Fernandez D. An analysis of the spatial colonization of

scrubland intrusive species in the Itabo and Guanabo watershed, Cuba. Remote Sens.

2010;2: 740–757.

46. Reyes YA, Pérez M, Barrera EL, Martínez Y, Cheng KK. Thermochemical conversion

processes of Dichrostachys cinerea as a biofuel: A review of the Cuban case. Renew

Sustain Energy Rev. 2022;160: 112322. doi:10.1016/j.rser.2022.112322

47. Hernández MR, Cruz DF. Natural vegetation cover in National Parks of Cuba:

multitemporal analysis and future variation of bioclimatic conditions. Rev Jardín

Botánico Nac. 2016;37: 93–102.

Supporting information captions

S1 Table. Values of the overall accuracy of the classification, for the years 1985 and 2020, for each land cover type considered in this study.

S1 Figure. Vegetation types in the areas of potential distribution of *Omphalea* plants in Cuba for the years 1985 and 2020. *Omphalea* includes all the species of *Omphalea* genus distributed in Cuba combined.

S2 Figure. Vegetation types in the terrestrial protected areas of Cuba for the years 1985 and 2020.

S3 Figure. Cover types in the areas of potential distribution of *Omphalea* plants in Cuba that are inside the terrestrial protected areas. *Omphalea* includes all the species of *Omphalea* genus distributed in Cuba combined and the areas of potential distribution of *Omphalea* is referring to these species' potential distribution in Cuba.

S1 Appendix 1. Description of the area of study.

S1 Appendix 2. Elevational considerations used for post-processing the classified images to correct misclassified pixels.

1985

Wetlands
Agriculture
Soil without vegetation cover
Forest and shrubs
Pine forest
Mangrove
Water bodies

2020

N

0   100 km

Figure 1

Cuba

Stable forest and shrubs
Mangrove to forest and shrubs
Soil without vegetation cover to forest and shrubs
Wetlands to forest and shrubs
Pine forest to forest and shrubs
Agriculture to forest and shrubs
Water bodies to forest and shrubs

Areas of potential distrubution of *Omphalea*

0   100 km

Figure 2