A novel order-level lineage of ammonia-oxidizing *Thaumarchaeota* is widespread in marine and terrestrial environments

Yue Zheng[1†], Baozhan Wang[2†], Ping Gao[2], Yiyan Yang[3], Xiaoquan Su[4], Daliang Ning[5], Qing Tao[5], Feng Zhao[6], Dazhi Wang[1], Yao Zhang[7], Meng Li[8], Mari-K.H. Winkler[9], Anitra E. Ingalls[10], Jizhong Zhou[5,11,12], Chuanlun Zhang[13,14], David A. Stahl[10], Jiandong Jiang[2]*, Willm Martens-Habbena[15]*, Wei Qin[5]*

[1]State Key Laboratory of Marine Environmental Science and College of the Environment and Ecology, Xiamen University, Xiamen, China;

[2]Department of Microbiology, College of Life Sciences, Nanjing Agricultural University, Key Laboratory of Agricultural and Environmental Microbiology, Ministry of Agriculture and Rural Affairs, Nanjing, China.

[3]National Library of Medicine, National Institutes of Health, Bethesda, MD;

[4]College of Computer Science and Technology, Qingdao University, Qingdao, China;

[5]Department of Microbiology and Plant Biology and Institute for Environmental Genomics, University of Oklahoma, Norman, OK, USA;

[6]CAS Key Laboratory of Urban Pollutant Conversion, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, China;

[7]State Key Laboratory of Marine Environmental Science and College of Ocean and Earth Sciences, Xiamen University, Xiamen, China;

[8]Archaeal Biology Center, Institute for Advanced Study, Shenzhen University, Shenzhen, China;

[9]Department of Civil and Environmental Engineering, University of Washington, Seattle, WA, USA

[10]School of Oceanography, University of Washington, Seattle, WA, USA;

[11]School of Civil Engineering and Environmental Sciences, University of Oklahoma, Norman, OK, USA;

[12]Earth and Environmental Sciences, Lawrence Berkeley National Laboratory, Berkeley, CA, USA;

[13]Shenzhen Key Laboratory of Marine Archaea Geo-Omics, Department of Ocean Science and Engineering, Southern University of Science and Technology, Shenzhen, China;

[14]Southern Marine Science and Engineering Guangdong Laboratory (Guangzhou), Guangzhou, 511458, China;

[15]Department of Microbiology and Cell Science & Fort Lauderdale Research and Education Center, University of Florida, Davie, FL, USA.

[†]These authors contributed equally to this work: Yue Zheng, Baozhan Wang.

[*]To whom correspondence should be addressed: Jiandong Jiang (jiang_jjd@njau.edu.cn), Willm Martens-Habbena (w.martenshabbena@ufl.edu), Wei Qin (weiqin@ou.edu).

Abstract

Ammonia-oxidizing archaea (AOA) are among the most ubiquitous and abundant groups of

Archaea on Earth, widely distributed in marine, terrestrial, and geothermal ecosystems.

However, the genomic diversity, biogeography, and evolutionary process of AOA populations in

subsurface environments are vastly understudied compared to those of marine and soil AOA.

We here report a novel AOA order *Candidatus* Nitrosomirales that forms a deeply branching

basal sister lineage to the thermophilic *Ca.* Nitrosocaldales. Metagenomic and 16S rRNA gene

read mapping demonstrates the dominant presence of *Nitrosomirales* AOA in various

groundwater environments and their widespread distribution across a range of geothermal,

terrestrial, and marine habitats. Notably, terrestrial *Nitrosomirales* AOA show the genetic

capacity of using formate as an alternative source of reductant and appear to have acquired key

metabolic genes and operons from other mesophilic populations via horizontal gene transfer,

including the genes encoding urease, nitrite reductase, and V-type ATPase. Potential metabolic

versatility and acquired functions may facilitate their radiation into a variety of subsurface,

marine, and soil environments. Molecular thermometer-based evolutionary analysis suggests

that *Nitrosomirales* originated from thermophilic environments and transitioned into temperate

habitats in parallel with *Nitrososphaerales* and *Nitrosopumilales*. We also provide evidence that

terrestrial-marine habitat transitions occurred within each one of the four AOA orders, which

reveals a more complex evolutionary trajectory of major AOA lineages than previously proposed.

Together, these findings establish a robust taxonomic and evolutionary framework of AOA and

provide new insights into the ecology and evolution of this globally abundant functional guild.

Introduction

The ammonia-oxidizing archaea (AOA) represent one of the most abundant and ubiquitous

archaeal groups in the global biosphere [1, 2, 3]. They account for nearly 30% of microbial

plankton in the oceans and up to 5% of microbial populations in soils [4, 5]. Apart from their

generally high abundance in a vast range of marine and terrestrial environments, they also play

an important role in the global nitrogen cycle. They are almost exclusively responsible for

ammonia oxidation in oligotrophic marine environments and contribute as much as 80% of the

emission of ozone-depleting potent greenhouse gas nitrous oxide (N_2O) from the oceans [6, 7].

This globally abundant and ecologically significant group of archaea was assigned to a major

archaeal phylum *Thaumarchaeota* (also named as *Nitrososphaerota*) [8, 9, 10].

Although all AOA are united by a common physiology of chemoautotrophic growth on ammonia

oxidation and carbon fixation, the extensive genetic repertoire of the pan-genome suggests many

additional adaptive features are associated with the remarkable ecological success of this

functional guild as reflected by their high global abundance and wide niche breadth [11, 12, 13,

14, 15, 16, 17]. Previous AOA phylogenetic, ecological, and evolutionary analyses were based

on a standardized taxonomic framework that divided ammonia-oxidizing *Thaumarchaeota* into

four major lineages, the *Nitrosopumilales* (Group 1.1 a) [18], *Candidatus* (*Ca.*) Nitrosotaleales

(Group 1.1 a-associated, now reclassified as a family within the *Nitrosopumilales*) [19],

Nitrososphaerales (Group 1.1 b) [20], and *Ca.* Nitrosocaldales (Thermophilic AOA, ThAOA)

[21], that appear somewhat specialized, respectively, to aquatic (marine or freshwater), acidic

soil, neutral or alkaline soil, and geothermal environments [12, 13, 16, 17, 22, 23]. Intriguingly,

a recent metagenomic study of cold deep seawater (~ 2.3 °C) recovered from the Mariana Trench

yielded a metagenome-assembled genome (MAG) that appeared to be phylogenetically closely

associated with the deeply branching thermophilic AOA [24]. This marine AOA MAG, along

with two other closely related MAGs recovered from deep-sea waters (UBA213 and SAT137),

had been assigned to *Ca*. Nitrosocaldales based on the previously established phylogenetic

backbone of AOA [11, 12]. However, further phylogenomic analyses with additional genomes

that represent a broader range of genotypes of this early branching lineage are needed to resolve

and substantiate the uncertain taxonomic affiliation and evolutionary trajectory of basal AOA

clades. In addition, how widespread these early branching mesophilic AOA are in marine and

terrestrial environments, and their diversity, metabolic adaptation, and ecological significance,

remain unknown.

Here we conduct phylogenomic and comparative genomic analyses of 118 AOA and non-

ammonia-oxidizing *Thaumarchaeota* genomes, including 23 MAGs and single amplified

genomes (SAGs) of this under-studied group that were obtained from a variety of subsurface,

geothermal, soil, and marine environments. We show that these 23 MAGs and SAGs form a

highly supported monophyletic order-level lineage within the *Nitrososphaeria*, which we here

designate a new AOA order *Ca*. Nitrosomirales. The global distribution of *Ca*. Nitrosomirales

AOA was investigated by extensively searching for their 16S rRNA and *amoA* gene sequences in

marine and groundwater metagenome datasets as well as in the Microbiome Search Engine 2

database that contains over 300,000 samples sequenced from a vast range of terrestrial and

aquatic environments [25]. Our findings provide new understanding of the metabolic potential,

evolution, and biogeography of this basal branching lineage that appears to represent a

predominant AOA genotype in previously under-sampled habitats, such as many terrestrial

subsurface environments and deep-sea sponges.

Materials and Methods

The distribution of *Ca.* Nitrosomirales AOA at global scale

The 16S rRNA gene sequences of all AOA were extracted by the ssu_finder of CheckM (version

1.0.12) [26]. The extracted AOA sequences were subsequently mapped into the database of

Microbiome Search Engine 2 (MSE2, http://mse.ac.cn) as previously described [25]. The MSE2

database contains more than 300,000 16S rRNA gene amplicon and metagenomic sequencing

samples collected from marine and terrestrial habitats as well as human, animal and plant

associated microbiomes. AOA-containing microbiomes in MSE2 were identified via

VSEARCH and using a 97% sequence similarity cutoff, by comparing the amplicon and

metagenomic sequences to 16S rRNA sequences extracted from 106 AOA genomes (Table S1)

[27]. The relative abundance of AOA amplicon sequences in each sample was normalized using

the Meta-Storms algorithm [28] to reduce the 16S rRNA gene amplification bias.

In addition to searching against the 16S rRNA gene amplicon database, the metagenomic data of

seawater and groundwater samples were searched against the *amoA* genes (encoding the alpha

subunit of ammonia monooxygenase). To explore the distribution of *Nitrosomirales* AOA in the

global ocean, we searched *amoA* gene reads from global ocean metagenomic databases,

including *Tara*-Oceans (2009-2013) [29] and *Malaspina*-2010 metagenomic data [30], Hawaii

Ocean time series station metagenomic data [31], and the Mariana Trench metagenomic data

[32]. To confirm the prevalence of *Nitrosomirales* AOA in the contaminated groundwaters, we

also searched *amoA* gene reads from the contaminated groundwater metagenomic samples

collected from the Oak Ridge Integrated Field Research Challenge (OR-IFRC) experimental

sites (FW300, FW301, FW602, and GW715) [33]. Briefly, we first trimmed the raw sequencing

reads of metagenomic samples by Trimmomatic (version 0.36) [34]. The trimmed reads were

then mapped via Diamond (version 0.9.24.125) [35] with a threshold of 80% similarity and 100

bp coverage, as well as an E-value cut-off of $1 \times e^{-10}$, to an in-house AOA species genome

database containing *amoA* genes found in available AOA species, MAGs, and SAGs. The

recruited reads were further classified into different AOA lineages. In addition, we searched for

16S rRNA and *amoA* gene sequences of *Nitrosomirales* in the NCBI database with BLASTN

using a sequence similarity cutoff of 95% and 90%, respectively, and an E-value cut-off of $1 \times e^{-}$

^{10}. The phylogenetic affiliation of the identified sequences was further confirmed in the 16S

rRNA and *amoA* gene trees. Only the sequences affiliated by a bootstrap value of more than 85%

(out of 1000 replications) within the *Nitrosomirales* clade were identified as the members of

Nitrosomirales.

Comparative genomic analysis and pathway construction

Average nucleotide identity (ANI) was calculated by pyani

(https://github.com/widdowquinn/pyani) using the model of BLASTN. The percentage identity

and alignment coverage between each of two genomes were displayed in heatmap format. The

corresponding genome size and GC content were marked for each AOA genome. AOA proteins

were clustered into homolog cluster groups (HCGs) by OrthoMCL (version 2.0.9) [36]. The

thresholds of HCGs were set to achieve a pairwise coverage of 50% and identity of 50% based

on all-against-all BLASTP. Orthologs and paralogs were identified as the reciprocal best

similarity pairs that were found between and within species, respectively [37]. Core genome

represents the HCG genes shared by all AOA species genomes, MAGs, and SAGs, and pan-

genome represents the collective set of genes present in at least one AOA genome. For *m*

genomes selected out of n strains, a total of $n!/(m-1)!\cdot(n-m)!$ combinations were calculated to

determine the sizes of the core and pan- genomes. To compare the pan-genome openness of

different AOA orders, the average number of new unique genes per Mbp genome was calculated

with the sequential addition of each AOA genome. Up to 5,000 random combinations were

sampled for core genome and pan-genome analyses. Finally, the pan-genomes of *Nitrosomirales*

and other AOA orders were visualized using the Anvi'o software (version 2.2.2) [38].

The functions of HCGs were subsequently annotated according to the reference AOA genomes

(*Ca.* Nitrosocaldus cavascurensis SCU2 [39] and *Nitrosopumilus maritimus* SCM1 [40]). The

putative metabolic pathways were classified into 16 groups, including ammonia

oxidation/assimilation/nitrate reduction, urea utilization, carbon fixation/metabolism, sulfur

assimilation/metabolism, phosphorus utilization, stress response, CPISPR system, thermo-

adaptation, amino acids/vitamins/cofactors, information processes, S-layer synthesis, lipid

biosynthesis, glycosyl transferase, hydrogen oxidation, ectoine synthesis, and transporters.

Putative transporters were further identified by screening against the Transporter Classification

Database [41].

Optimal growth temperature estimates for *Nitrosomirales* AOA

The stem regions of *Nitrosomirales* AOA 16S rRNA genes were predicted based on

compensatory base-pair changes using the RNAfold package [42], and the GC contents of the

predicted stem were calculated for the *Nitrosomirales* genome-retrieved 16S rRNA gene

sequences. The evolutionary model of AOA 16S rRNA sequences along the reference tree was

estimated by a nonhomogeneous model using bppml, and a total of 100 replicates of ancestral

sequences were constructed using the program BppAncestor (BppSuite version 2.4.1) as

previously described [13, 43]. The optimal growth temperatures (OGTs) of the ancestors of

Nitrosomirales AOA were estimated using the previously calibrated linear regression between

the GC contents of AOA 16S rRNA gene stem regions and their OGTs [13]. The OGTs with

confidence intervals of corresponding nodes (ancestor nodes of all AOA and the four orders)

were inferred against the linear regression.

Phylogenomic tree construction

Phylogenomic tree construction was based on concatenated alignments of 70 conserved marker

genes (Table S2) from 106 AOA genomes (Table S1) and outgroup archaeal species. The

genomes (*Nitrosomirales sp*. UBA213 and *Nitrosomirales sp*. WS1) that had not been annotated

in NCBI and JGI were annotated using GeneMarks in this study [44] (Supplementary Datasets

S1 and S2). The outgroup species for phylogenomic analysis included non-ammonia-oxidizing

Thaumarchaeota, *Aigarchaeota*, *Bathyarchaeota*, *Crenarchaeota*, *Korarchaeota*, *Lokiarchaeota*,

Euryarchaeota, and *DPANN* archaeal species. The 70 marker genes were identified by BLASTP,

and they were individually aligned by MAFFT (version 7.221) [45]. Subsequently, the

conserved regions of alignments were extracted by Gblocks (version 0.91b) [46]. Conserved

regions were concatenated as a single evolutionary unit. For phylogenomic analysis, the best-fit

model of amino acid substitution was estimated using ProtTest (version 3.4) [47]. Maximum

likelihood phylogenomic trees were built with concatenated sequences via RAxML (version

8.0.26) using the LG+I+G+X model on the basis of 100 bootstrap replications [48].

Phylogenetic tree of key genes

The sequences of 16S rRNA, *amoA*, *atpA* and *atpC* (encoding the alpha and epsilon subunits of

ATPase, respectively), *nirK* (encoding putative nitrite reductase), and *ureC* genes (encoding the

alpha subunit of urease) were extracted from the collected 106 AOA genomes (Table S1).

Additional *Nitrosomirales* 16S rRNA genes and 33,387 AOA *amoA* genes were collected from

NCBI and extracted from a previous *amoA* gene-based phylogenetic analysis [22], respectively,

for comparative phylogenetic analysis.

The phylogenetic trees of the 16S rRNA and other key functional genes of AOA were built using

IQ-TREE (v. 2.1.2 COVID-edition) with best-fit model selection [49]. Gene sequence alignment

was carried out using MAFFT (v7.407) [45] and was then edited with Gblocks (v0.91b) [50] to

identify conserved regions. The best-fit model of evolution for each gene set was selected by

ModelFinder [51] with "-m MF –T AUTO" flag as follows: GTR+F+R4 for 16S rRNA genes,

GTR+F+I+Γ4 for *amoA* genes, LG+R5 for *atpA*, LG+R4 for *atpC* genes, WAG+F+G4 for *nirK*

genes, and LG+R3 for *ureC* genes. Subsequently, the maximum likelihood phylogenetic trees

were built using IQ-TREE with best model mentioned above and 1,000 times ultrafast

bootstrapping [52]. The non-ammonia-oxidizing *Thaumarchaeota* were used as the outgroup for

the 16S rRNA gene tree. *Thermoproteales*, *Sulfolobus*, *Desulfurococcales Ignicoccus hospitalis*

KIN4/I, *Ca*. Bathyarchaeota archaeon ex4484 231, and *Methanosuratus*/*Methanomethylicus*

were used as the outgroups for the phylogenetic trees of the subunits A and C of the A-type

ATPases, and *Thermoplasmatales* as well as *Enterococcus hirae* ATCC 9790 were used as the

outgroups for the trees of the subunits A and C of the V-type ATPases.

Availability of data and materials

AOA genomes used in this study are available in the NCBI and JGI databases. The

corresponding accession numbers are listed in Table S1. The annotation file of *Nitrosomirales*

MAGs UBA213 and WS1 are provided in GFF format (Supplementary Datasets S1 and S2).

Completeness, contamination, and coding density of AOA genomes were assessed by CheckM

(version 1.0.12) [53]. All other data products associated with this study are available from the

corresponding authors upon request.

Results and Discussion

Nitrosomirales **order represents a basal lineage of AOA**

We compiled a 118-genome dataset that comprises 106 AOA cultured genomes, MAGs, and

SAGs as well as 12 representative genomes of non-ammonia-oxidizing *Thaumarchaeota* to serve

as outgroups (Table S1). To infer the most probable evolutionary relationship among AOA taxa

and resolve the uncertain taxonomic affiliation of mesophilic members in the deepest branching

AOA lineages, we constructed a maximum likelihood phylogenomic tree from the concatenation

of 70 conserved marker genes (Table S2) present in AOA genomes and outgroup

thaumarchaeotal MAGs. We obtained 20 additional MAGs and SAGs from various terrestrial

subsurface, deep soil, and marine habitats that were clustered with three previously reported

marine AOA MAGs MTA5, UBA213, and SAT137 [11, 12, 24], and they together formed a

well-supported monophyletic group branching as a sister clade to the thermophilic *Ca.*

Nitrosocaldales AOA (Fig. 1). The monophyletic phylogeny of this deeply branching lineage

did not emerge in previous analyses using single-gene markers or only with the three reported

marine AOA MAGs [11, 12, 16, 24]. Among these newly obtained MAGs and SAGs, nine were

retrieved from **w**arm/thermal **s**pring (WS1–WS3) and **c**arbonate **s**pring (CS1–CS6) waters and

sediments, six originated from groundwaters (GW1–GW6), two from **b**asaltic **l**ava **c**aves (BLC1

and BLC2), two from **d**eep **s**andy **s**oils in switchgrass fields (DSS1 and DSS2), and one from

deep **o**cean waters (OD1) (Table S3). Eleven of these genomes showed a generally high level of

completeness ($> 78\%$) and low contamination ($< 5\%$) (Table S3). MAGs DSS1 and WS1 are

complete or nearly complete genomes and estimated to be 100% and 99.93% complete with

minimal contamination (0.97%), respectively. These MAGs and SAGs span a large size range

from 0.99–3.68 Mbp, and similar to other AOA genomes, the GC content of marine

Nitrosomirales genomes (34.8–36.1%) was lower than that of terrestrial *Nitrosomirales* genomes

(40.3–47.7%) (Table S3). The average nucleotide identity (ANI) values among these 23

genomes were 67.6%–99.6% (Table S4), reflecting generally high genomic diversity within this

group. The characterized *Nitrosomirales* genomes shared low genomic homology (<68.2% ANI)

and alignment fraction (4.8%–29.4%.) with other marine and terrestrial AOA genomes (Table

S4).

Additional whole-genome based taxonomic analysis using the Genome Taxonomy Database

Toolkit (GTDB-Tk) also showed these 23 MAGs and SAGs to be members of a distinct basal

lineage along with the other three formally described major AOA lineages (Table S3) [18, 20,

21]. We here propose the name *Ca.* Nitrosomirales to represent this deeply branching AOA

order within the class *Nitrososphaeria* (Fig. 1), branching between the orders *Nitrososphaerales*

and thermophilic *Ca*. Nitrosocaldales with the complete MAG DSS1 from deep sandy soil

representing the candidate family *Nitrosomiraceae* and genus *Nitrosomirus*. The name

Nitrosomirus refers to the organisms as ammonia oxidizers with the capacity of oxidizing

ammonia to nitrite (from the Latin "*nitrosus*", full of natron; here intended to mean nitrous) and

its surprisingly wide distribution range in various marine, soil, geothermal, and subsurface

habitats (from the Latin adjective "*mirus*", meaning surprising and amazing), spanning a range of

temperature, salinity, pressure, and nutrient availability (see the description below).

Nitrosomirales **AOA are widely distributed in diverse terrestrial, marine, and geothermal**

habitats

The recovery of many *Nitrosomirales* MAGs from subsurface, deep soil, deep ocean, and

geothermal environments suggest that *Nitrosomirales*-AOA are ubiquitous in diverse terrestrial

and marine habitats. To further explore the distribution of this under studied group in the global

biosphere, we searched the *Nitrosomirales* MAG-derived 16S rRNA sequences against the

Microbiome Search Engine 2 16S rRNA gene database that contains over 300,000 16S rRNA

gene amplicon and metagenomic sequencing samples obtained from a broader range of natural

habitats and engineered systems, as well as human, animal and plant associated microbiomes

(see Materials and Methods for details) [25]. *Nitrosomirales*-derived 16S rRNA gene sequences

were found in various terrestrial and marine habitats, including groundwaters, farmland soils,

forest soils, wetlands, and lake sediments, marine sponges, and marine aquaculture waters (Figs.

2 and Table S5). Additional extensive searches of 16S rRNA (similarity cutoff of 95% and E-

value cutoff of $1 \times e^{-10}$) in NCBI database further identified *Nitrosomirales* sequences in

(moderate)thermophilic habitats, such as hydrothermal vents (>52.0 °C and 72–103 °C), hot

springs (55.0 °C), and thermal karst well waters (73.7 °C) (Fig. 2). Both the 16S rRNA and

amoA gene-based phylogenies support the monophyletic grouping of *Nitrosomirales*

metagenomic, amplicon, and clone sequences (Figs. 2 and S1). Together, our results indicate

that the habitats of *Nitrosomirales* AOA include a wide variety of terrestrial, marine, subsurface,

and geothermal environments (Fig. 3a). Thus, the extraordinary habitat range of *Nitrosomirales*

exceeds that of any previously described AOA order-level lineages.

***Nitrosomirales* AOA represent the dominant AOA population in subsurface habitats and**

deep-sea sponges

Although the first three *Nitrosomirales* MAGs were recovered from the dark ocean,

Nitrosomirales AOA were rarely detected in global ocean metagenomes. We assessed the

number of metagenomic reads mapping to the *amoA* genes of *Nitrosomirales* AOA and other

marine AOA genotypes in global ocean metagenome databases across four oceans and two seas,

spanning from epipelagic to hadopelagic zones [29, 30, 31, 32], and found that *Nitrosomirales*

AOA only comprised at most 0.5% of the total AOA community in well-studied ocean waters

(Table S5). In contrast, the 16S rRNA and *amoA* gene read recruitment showed that

Nitrosomirales AOA can represent an abundant AOA genotype or even exclusively represent the

whole AOA community in some previously under-sampled habitats, such as deep-sea sponges

and groundwaters (Fig. 3b and Table S5).

Marine AOA have been frequently reported to associate with sponges [54, 55, 56, 57, 58] and

often dominate the whole archaeal communities of sponge microbiomes [59]. Since the

sequencing of the first sponge AOA symbiont *Ca.* Cenarchaeum symbiosum hosted by the

demosponge *Axinella mexicana* [54], additional AOA MAGs were obtained from various

shallow-water and deep-sea sponge-associated microbiomes. Marine sponges are a highly

diverse clade of metazoans that contains 125 families, 680 genera, and 11,000 species [60]. The

molecular surveys of sponge microbiota indicated that microbial communities were mostly

specific to sponge species [61], and the dominant AOA populations were even specific to

individual sponges [58]. Previous studies of thaumarchaeotal sponge symbionts only focused on

several limited sponge families of the classes *Demospongiae* and *Hexactinellida*. These sponge

symbionts were affiliated with five different AOA genera, *Nitrosopumilus* [56], *Ca.*

Cenarchaeum [54], *Ca.* Nitrosopelagicus [58], *Ca.* Nitrosospongia [55], and *Ca.*

Cenporiarchaeum [56], all of which were within the order *Nitrosopumilales*.

We found that, apart from *Nitrosopumilales*, the members belonging to *Nitrosomirales* can also

constitute a significant fraction (up to 96.05%) of the total AOA populations in deep-sea sponges

(Fig. 3b). *Nitrosomirales* AOA were specifically hosted by the deep-sea (~200–550 m)

Aciculites sponge species within the *Scleritodermidae* family and the *Porifera* phylum (Table

S5), and this lineage of sponges was poorly represented in the previous 16S rRNA gene surveys

and the global sponge microbiome metagenomic database. Intriguingly, the 16S rRNA gene

sequences that are closely related to *Ca.* Nitrosocosmicus AOA within the *Nitrososphaerales*

were found in marine sponges *Spirastrella panis* [62], *Astrosclera willeyana* [63], *Theonella*

swinhoei [64], *Pseudoceratina purpurea* (NCBI accession No.: KU064739), and *Halichondria*

oshoro (HM101091) associated microbiomes (Fig. S2). These findings significantly expand the

genetic diversity of the sponge-associated marine AOA beyond the order *Nitrosopumilales*. It is

very likely that *Nitrosomirales* AOA play an important role in the nitrogen metabolism and

nitrogenous waste removal of deep-sea *Aciculites* sponges, similar to other characterized sponge-

associated *Nitrosopumilales* AOA [54, 55, 56].

Another relatively under-sampled AOA habitat is terrestrial subsurface environment. The

biogeography of AOA in groundwater and cave ecosystems and associated environmental

variables that control the abundance and composition of AOA communities in these systems are

poorly documented [65]. Given that several *Nitrosomirales* MAGs were retrieved from the

Death Valley Regional Flow System (DVRFS), we leveraged the available 16S rRNA amplicon

sequencing data [66] collected from the DVRFS region to assess the relative abundance of

Nitrosomirales AOA across three major groundwater basins (Table S5). *Nitrosomirales* AOA

16S rRNA genes were detected in nine DVRFS groundwater sites out of the 36 total sampling

sites. Based on 16S rRNA gene read recruitment, we found that *Nitrosomirales* AOA can

constitute a substantial proportion (30.94–100.00%) of the total AOA populations in six of these

groundwater sites (Fig. 3b and Table S5). *Nitrosomirales* AOA were detected in both shallow

(0–25 m sampling depth) and deep (474–700 m sampling depth) aquifers with distinct aquatic

geochemistry, including Ca-Mg-HCO$_3$, Na-HCO$_3$, and NaCl dominated groundwaters (Table S5).

In addition, they were also found in basaltic lava caves (Table S3), together indicating their wide

distribution in terrestrial subsurface environments.

Interestingly, we also identified this lineage in the 16S rRNA gene and metagenome datasets

collected from the groundwaters at the Hanford site and the Oak Ridge Integrated Field Research

Challenge (OR-IFRC) site, respectively (Fig. 3b and Table S5), legacies of the Manhattan

Project contaminated with mixed waste, including radionuclides and nitrate [67]. *Nitrosomirales*

AOA were not detected in heavily uranium-contaminated groundwaters but can be enriched in

groundwaters with relatively high nitrate concentrations (as high as 816.56 mg/L). It is

conceivable that respiratory ammonifiers with the capacity of the dissimilatory nitrate reduction

to ammonium (DNRA) could supply ammonia to *Nitrosomirales* AOA and other archaeal and

bacterial nitrifiers in these contaminated groundwater sites [68], and thus together these

microbial communities may contribute to nitrogen transformation in terrestrial subsurface

ecosystems.

Genomic features and metabolic potential of *Nitrosomirales* AOA

We calculated the core genome and pan-genome of *Nitrosomirales* and other AOA orders to get

quantitative insights into the conserved and flexible gene pools of ammonia-oxidizing

Thaumarchaeota (Fig. S3). Comparative genomic analysis showed that all terrestrial and marine

Nitrosomirales AOA MAGs and SAGs harbored the conserved pathway genes that are involved

in the characterized central metabolism of AOA, including the copper-dependent respiration and

electron transfer systems, 3-hydroxypropionate/4-hydroxybutyrate autotrophic carbon fixation

cycle, and the biosynthesis of the B vitamin cofactors thiamin (B_1), riboflavin (B_2), pantothenate

(B_5), pyridoxine (B_6), biotin (B_7), and cobalamin (B_{12}) (Fig. 4 and Table S6). The mesophilic

Nitrosomirales AOA also contained many genes that were thought to be present only in

thermophilic *Ca*. Nitrosocaldus/Nitrosothermus AOA and associated with compatible solutes

production for thermoprotection [39, 69], such as the genes encoding mannosyl-3-

phosphoglycerate synthase and cyclic 2, 3-diphosphoglycerate synthetase (Table S6). This may

be due to the retention of ancestral features prior to the adaptive divergence of AOA into

mesophilic habitats or may suggest completely different functions of these compatible solutes in

Nitrosomirales and *Ca*. Nitrosocaldales AOA.

The pan-genome openness of *Nitrososphaerales* was the highest among the four AOA orders

(Fig. S3b), and the high genomic diversity was driven by the extensive lateral gene transfer and

gene duplication events during *Nitrososphaerales* evolution [12]. The *Nitrosomirales* AOA pan-

genome within the available dataset contains a total of 11,334 genes, and the pan-genome graph

shows that the sampling of their pan-genome is far from reaching saturation (Fig. S3b). An

average of ~160 novel unique genes per Mbp genome are expected to be identified with each

new *Nitrosomirales* species sequenced, which is comparable to the number of new unique genes

estimated for *Nitrosopumilales* AOA (Fig. S3c). We further performed Anvi'o pan-genome

analysis to assess and visualize *Nitrosomirales*-specific gene contents (Fig. 4). A total of

211,167 coding sequences of the collected AOA genomes (over 50% completeness) were

clustered into 13,495 protein clusters. Of these, 1,260 protein clusters were specifically enriched

in *Nitrosomirales* genomes but absent or scarce in other AOA orders, including the accessory

and unique genes that were assigned to certain Clusters of Orthologous Groups (COGs)

functional categories, such as transcription, replication, recombination and repair, energy

production and conversion, and transporters as well as many hypothetical genes with unknown

functions (Table S7).

Different from the members of their sister order *Ca*. Nitrosocaldales [39, 69, 70], *Nitrosomirales*

AOA lacked identifiable hydrogenase genes involved in potential hydrogen oxidation for energy

conservation (Table S6). Among the *Nitrosomirales*-specific genes, the genes that encode the

major (alpha, containing the active site) subunit of formate dehydrogenase, were found only in

Nitrosomirales genomes, including those obtained from carbonate and warm springs and deep

sandy soils (Fig. 4 and Table S6). *Nitrosomirales* genomes also encodes the putative formate

dehydrogenase beta and gamma subunits, which may be involved in electron transfer processes

and regulatory functions. In addition, the putative formate transporter genes were identified only

in terrestrial *Nitrosomirales* genomes (Fig. 4 and Table S6), strongly suggesting that formate

may serve as an alternative electron donor and source of reductant for terrestrial *Nitrosomirales*

AOA. It has been shown that *Nitrospira* nitrite-oxidizing bacteria (NOB) species were able to

grow using formate as sole substrate with O_2 or nitrate as terminal electron acceptors [71].

Interestingly, we identified genes that encode the alpha and beta subunits of the putative nitrate

reductase in ten terrestrial *Nitrosomirales* genomes (Fig. 4 and Table S6), suggesting that the

formate oxidation with nitrate reduction may be directly coupled in *Nitrosomirales* under anoxic

conditions, as observed in NOB. Future cultivation and isolation of *Nitrosomirales* AOA are

required to confirm their metabolic capacity of using formate as a source of reductant, and such

metabolic versatility may enable *Nitrosomirales* AOA to survive during periods of ammonia

deprivation.

Other unique genes that are conserved in *Nitrosomirales* AOA include two small blue copper

proteins that may function as electron shuttles in the respiratory system, putative CopA and

CopD proteins for copper sequestration and uptake, and various regulation proteins that may be

associated with stress response and adaptation, such as *Nitrosomirales*-specific transcriptional

regulatory and two-component regulatory proteins, DNA repair proteins, and Heat shock

proteins (Hsp20 family) (Fig. 4 and Table S6). Putative surface layer (S-layer) protein genes

were identified in *Nitrosomirales* genomes (Fig. 4 and Table S6). Moreover, *Nitrosomirales*

AOA were found to possess up to three additional putative PKD (polycystic kidney disease)

domain protein homologs (Fig. 4 and Table S6), which have been found in extracellular parts of

archaeal S-layer proteins that protect cells from extreme environments and assist in cell adhesion

and intercellular interaction [72, 73].

***Nitrosomirales* AOA acquire key metabolic genes via horizontal gene transfer**

Similar to other AOA orders, *Nitrosomirales* AOA genomes contain the genes coding for the

known ABC subunits of ammonia monooxygenase (AMO), the predicted AMOX, and newly

identified AMOY and AMOZ subunits [74] (Fig. 4 and Table S6). The electrons from ammonia

oxidation are transferred to oxygen via the copper-centric electron transfer chain, which would

lead to the generation of a proton motive force for ATP synthesis via energy-yielding ATPase.

The phylogenetic trees of AOA ATPase alpha (A) and epsilon (C) subunits showed clear

bifurcating topologies of archaeal-type (A-type) and vacuolar-like (V-type-like) ATPase

subgroups (Fig. S4). The A-type ATPase gene clusters were found in both marine and terrestrial

Nitrosomirales AOA genomes, and their phylogeny tracked organismal phylogeny rather than

ecological habitat (Fig. S4). Likewise, the A-type *atp* operon of marine *Nitrosomirales* AOA

shared the conserved organization and orientation with those of the terrestrial *Ca*.

Nitrosocaldales and *Nitrososphaerales* AOA, but were distinct from marine *Nitrosopumilales*

species (Fig. S5). Interestingly, four terrestrial *Nitrosomirales* MAGs encoded the entire gene

clusters for both A-type and V-type-like ATPases (Fig. S5). Partial gene clusters of V-type

ATPase were also found in the deep-sea MAG MTA5, two additional terrestrial MAGs, and a

SAG obtained from groundwaters (Fig. S5). Missing V-ATPase subunits in these *Nitrosomirales*

genomes may reflect genome incompleteness.

The V-type ATPase has been proposed to serve to maintain the cytosolic pH homeostasis in

acidophilic and acid tolerant AOA as well as deep marine AOA by pumping out excessive

cytoplasmic protons under acidic or high-pressure conditions [75]. Recent AOA comparative

population genomics and phylogenomic analyses have shown that V-type ATPase genes were

widely distributed among deep-sea water column [11] and sedimentary [76] *Nitrosopumilales*

AOA and desert *Nitrososphaerales* AOA populations [77]. Apart from the experimentally tested

function in low pH adaptation, AOA V-type ATPase may be also coupled to sodium (Na^+)

motive force at high pH levels and protect cells from high-salt stress [24, 77]. The ATPase

subunit phylogenetic trees show that the AOA V-type ATPase subgroup rooted with

Nitrosomirales AOA variants (Fig. S4). Thus, it is most likely that *Nitrosomirales* species also

acquired the V-ATPase by horizontal gene transfer, as observed for other AOA order species

[75]. It is interesting to note that the V-type *atp* operon was located next to the A-type *atp*

operon in *Nitrosomirales* MAGs (Fig. S5). In contrast, although the hadopelagic

Nitrosopumilales AOA species also contain both types of ATPases, the V-type *atp* operon was

distantly located from the A-type *atp* operon (Fig. S5), suggesting the highly mobile V-type *atp*

operon may have been reshuffled by several successive events along the evolutionary trajectory

of AOA.

Nitric oxide (NO) has been shown as a central intermediate in the archaeal ammonia oxidation

pathway [6, 78], and the putative NO-forming nitrite reductase (NirK) is conserved among

Nitrosopumilales and *Nitrososphaerales* AOA species [11]. However, no gene encoding NirK

proteins has yet been identified in *Ca.* Nitrosocaldus AOA species [39, 70]. In contrast, it was

identified in thermophilic *Ca.* Nitrosothermus AOA MAGs [69]. NirK homologs were also

identified in both marine and terrestrial *Nitrosomirales* genomes, and many *Nitrosomirales*

species encode two NirK paralogs in the genomes (Fig. S6). Intriguingly, the *nirK* genes of the

deep-sea *Nitrosomirales* MAG MTA5 and warm spring MAG WS3 did not cluster with those

encoded by the relatively closely related *Ca.* Nitrosothermus AOA, but rather grouped with those

of the distinctly related deep-sea water column B and terrestrial *Nitrosotenuis* AOA, respectively

(Fig. S6). This suggests that the *nirK* genes in these *Nitrosomirales* MAGs have been acquired

via lateral gene transfer from *Nitrosopumilales* AOA that share similar habitats. In addition,

ureC, the gene encoding the alpha subunit of urease, was also found in terrestrial *Nitrosomirales*

MAGs (Fig. S7) and putatively acquired laterally from other mesophilic terrestrial AOA lineages.

The *ureC* genes of *Nitrosomirales* MAGs were phylogenetically distinct from those of

thermophilic *Ca.* Nitrosocaldales AOA but nested among *Nitrososphaerales* AOA lineages (Fig.

S7). Thus, *Nitrosomirales* AOA may have acquired urea utilization genes to enhance metabolic

versatility during their evolution. Taken together, our results highlight the extensive lateral

transfer of key genes and operons involved in energy conservation in *Nitrosomirales* AOA, and

the acquisition of these essential metabolic genes may have facilitated their radiation into the

diversity of subsurface, marine, and geothermal environments they inhabit as now revealed by

our metagenomic-based biogeography analysis.

Hot origin of *Nitrosomirales* and terrestrial-marine habitat transitions within each AOA

order

The presence of *Nitrosomirales* sequences in geothermal habitats indicates their broad

occurrence in habitats that span a wide range of temperatures (Fig. 2). To estimate the optimal

growth temperatures (OGTs) of the ancestors of *Nitrosomirales* AOA and other AOA lineages,

we used molecular thermometer-based evolutionary analysis, which involved predicting

ancestral 16S rRNA stem composition along the reference tree [43] and using the established

linear relationship between the OGTs of cultured AOA species and their 16S rRNA stem GC

contents [13, 79]. The predicted ancestral OGT for *Nitrosomirales* AOA was 69 °C (Fig. S8),

suggesting that this deeply branching AOA lineage likely originated from hot environments. In

addition, the estimated OGTs of the ancestors of *Nitrosopumilales*, *Nitrososphaerales*, and *Ca.*

Nitrosocaldales, as well as the last common ancestor of AOA were broadly consistent with the

previously reported values by Abby et al. [13] (Fig. S8). Taken together, these results provide

additional support for the evolutionary model proposing the hot origin of AOA [13, 80].

Our comprehensive single-gene and whole-genome based phylogenetic analyses with additional

Nitrosomirales genomes and marine AOA 16S rRNA sequences provide a better resolved

framework of the AOA phylogeny and reconstruct a more complex evolutionary trajectory of

major AOA lineages than previously proposed. We found that after the initial expansion from

hot to moderate temperature environments, AOA diverged via two primary paths of adaptive

evolution that ultimately gave rise to the two major contemporary branches: the newly defined

Nitrosomirales AOA spanning both terrestrial and marine environments, and the previously

described mesophilic AOA including *Nitrososphaerales* mostly in terrestrial settings and

Nitrosopumilales mostly in marine settings (Fig. 1). Previous molecular dating analyses

suggested that AOA first transitioned into the temperate terrestrial environments before

expanding to marine environments, and further transition from shallow marine sea into deep-sea

waters awaited the oxygenation of the deep ocean during the Neoproterozoic [16, 81]. However,

we found the presence of thermophilic *Nitrosocaldales* 16S rRNA gene sequences in various

high-temperature marine environments, including shallow-sea hydrothermal vents [82], a coastal

hot springs (NCBI accession No.: JX047158), deep-sea hydrothermal fields of the Marina

Trough [83], and the walls of an active deep-sea sulfide chimney with 302 ℃ venting liquid [84]

(Fig. S9). This strongly suggests the expansion of AOA into both shallow and deep-sea habitats

prior to their transition to temperate terrestrial environments and likely places the origin of

marine AOA further back in evolutionary history than previously thought. In addition,

interestingly, we found *Nitrososphaerales* 16S rRNA sequences were present in many sponge

and coral reef samples [62, 63, 64, 85] as well as deep-sea sediments [86] (Fig. S9), revealing the

broader habitat range of *Nitrososphaerales* AOA that have been thought mainly restricted to

terrestrial environments [22]. Taken together, these results indicate that the terrestrial-marine

habitat transitions occurred in not only the order *Nitrosopumilales*, but also each one of the four

major orders throughout the evolutionary history of ammonia-oxidizing *Thaumarchaeota* (Fig.

S9), depicting a highly dynamic evolutionary process of this globally widespread functional

guild.

Conclusions

Our comparative genomic and phylogenomic analyses of 118 AOA and non-ammonia-oxidizing

Thaumarchaeota genomes revealed a new AOA cluster, *Ca*. Nitrosomirales, that forms a deeply

branching order-level lineage within the class *Nitrososphaeria*. Apart from containing expected

gene inventories for ammonia oxidation, carbon dioxide fixation, and B-vitamin biosynthesis,

Nitrosomirales AOA have the genetic capacity of using formate as an alternative energy source,

which may provide metabolic versatility under ammonia starvation. Biogeographic analysis of

16S rRNA and *amoA* genes indicates that *Nitrosomirales* AOA are widely distributed in

geothermal, terrestrial, and marine environments, and they are particularly abundant in

subsurface terrestrial environments, and some deep-sea sponges. Evidence for independent

terrestrial-marine transitions within each AOA order will foster a more detailed understanding of

the early evolution and adaptive radiation of archaeal ammonia oxidation during the Proterozoic

era.

Acknowledgement

We thank the DOE Joint Genome Institute for sharing *Nitrosomirales* MAGs and SAGs. This

work was supported by the Fundamental Research Funds for the Central Universities of China

(to JJ and BW), the National Natural Science Foundation of China grants 41977056 (to BW),

41907027 (to YZ), and the 333 high-level talent project of Jiangsu (to JJ). ML was supported by

the National Natural Science Foundation of China grant (32225003, 92251306) and the

Innovation Team Project of Universities in Guangdong Province (No. 2020KCXTD023). CZ

was supported by the Shenzhen Key Laboratory of Marine Archaea Geo-Omics, Southern

University of Science and Technology (ZDSYS201802081843490), and the Southern Marine

Science and Engineering Guangdong Laboratory (Guangzhou) (No. K19313901). YZ was

supported by the National Natural Science Foundation of China grant 92051114. WMH was

supported by Florida Agricultural Experiment Station Hatch project FLA-FTL-005680. WQ was

supported by Simons Postdoctoral Fellowship in Marine Microbial Ecology (548565) and the

startup funding of the University of Oklahoma.

Competing interests

The authors declare that they have no competing interests.

576
Figures

Figure 1. Phylogeny of *Ca*. Nitrosomirales and other AOA orders. Phylogenomic inference of

AOA species affiliated to the orders *Nitrosopumilales* (blue, basal lineage #1), *Nitrososphaerales*

(green, basal lineage #2), *Ca*. Nitrosomirales (yellow, basal lineage #3), and *Ca*. Nitrosocaldales

(orange, basal lineage #4) based on concatenated sequences of 70 conserved single-copy core

genes. The non-ammonia-oxidizing thaumarchaeotal genomes were used as outgroups (grey).

Please note that, based on the recent comparative phylogenetic analysis of AOA genomes [12],

the previously defined acidophilic AOA group *Ca.* Nitrosotaleales has been merged with

Nitrosopumilales. Confidence values were provided based on 100 bootstrap replications.

Figure 2. Phylogenic distribution of *Ca*. Nitrosomirales 16S rRNA genes according to habitat

types (squares, inner ring) and temperature ranges (circles, outer ring). Confidence values are on

the basis of 1000 bootstrap replications. The scale bar represents 2% estimated sequence

divergence. The 16S rRNA sequences that retrieved from *Ca*. Nitrosomirales genomes were

indicated with stars. The 16S rRNA genes of other thaumarchaeotal lineages were collapsed as

triangles.

Figure 3. The global distribution of *Nitrosomirales* AOA in various terrestrial and marine

environments (**a**). *Nitrosomirales* AOA were enriched in sponge-associated microbiomes and

subsurface environments (**b**). Relative abundance of *Nitrosomirales* AOA to total AOA were

estimated by comparing the number of amplicon sequencing and metagenomic reads mapping to

the 16S rRNA and *amoA* genes of *Nitrosomirales* AOA relative to those of total AOA in the

Microbiome Search Engine 2 global 16S rRNA database, groundwater metagenomes and global

ocean metagenome databases.

Figure 4. Anvi'o representation of the pangenome of *Nitrosomirales* and other AOA genomes.

Each radial layer represents an AOA genome, which was arranged by the order of phylogenomic

tree shown in Fig. 1. In each layer, dark and light colors represent the presence and absence of

protein clusters, respectively. The heatmap in the top right corner represents the conserved

functions present in the core genome of all AOA orders and highlights key unique gene contents

retrieved from the protein clusters that are specifically enriched in *Nitrosomirales* (indicated by

the black frame on the genome map rails).

References

1   Francis CA, Roberts KJ, Beman JM, Santoro AE, Oakley BB (2005). Ubiquity and diversity
of ammonia-oxidizing archaea in water columns and sediments of the ocean. *P Natl Acad Sci*
USA **102:** 14683-14688.

2   Leininger S, Urich T, Schloter M, Schwark L, Qi J, Nicol GW *et al* (2006). Archaea
predominate among ammonia-oxidizing prokaryotes in soils. *Nature* **442:** 806-809.

3   Gubry-Rangin C, Hai B, Quince C, Engel M, Thomson BC, James P *et al* (2011). Niche
specialization of terrestrial archaeal ammonia oxidizers. *P Natl Acad Sci USA* **108:** 21206-21211.

4   Karner MB, DeLong EF, Karl DM (2001). Archaeal dominance in the mesopelagic zone of
the Pacific Ocean. *Nature* **409:** 507-510.

5   Bates ST, Berg-Lyons D, Caporaso JG, Walters WA, Knight R, Fierer N (2011). Examining
the global distribution of dominant archaeal populations in soil. *ISME J* **5:** 908-917.

6   Martens-Habbena W, Qin W, Horak REA, Urakawa H, Schauer AJ, Moffett JW *et al* (2015).
The production of nitric oxide by marine ammonia-oxidizing archaea and inhibition of archaeal
ammonia oxidation by a nitric oxide scavenger. *Environ Microbiol* **17:** 2261-2274.

7   Ji Q, Buitenhuis E, Suntharalingam P, Sarmiento JL, Ward BB (2018). Global nitrous oxide
production determined by oxygen sensitivity of nitrification and denitrification. *Glob*
Biogeochem Cycles **32:** 1790-1802.

8   Oren A, Garrity GM (2021). Valid publication of the names of forty-two phyla of
prokaryotes. *Int J Syst Evol Microbiol* **71**.

9   Brochier-Armanet C, Boussau B, Gribaldo S, Forterre P (2008). Mesophilic crenarchaeota:
proposal for a third archaeal phylum, the Thaumarchaeota. *Nat Rev Microbiol* **6:** 245-252.

10   Spang A, Hatzenpichler R, Brochier-Armanet C, Rattei T, Tischler P, Spieck E *et al* (2010).
Distinct gene set in two different lineages of ammonia-oxidizing archaea supports the phylum
Thaumarchaeota. *Trends Microbiol* **18:** 331-340.

11   Qin W, Zheng Y, Zhao F, Wang YL, Urakawa H, Martens-Habbena W *et al* (2020).
Alternative strategies of nutrient acquisition and energy conservation map to the biogeography
of marine ammonia-oxidizing archaea. *ISME J* **14:** 2595-2609.

12   Sheridan PO, Raguideau S, Quince C, Holden J, Zhang LH, Williams TA *et al* (2020). Gene
duplication drives genome expansion in a major lineage of Thaumarchaeota. *Nat Commun* **11:**
5494.

13   Abby SS, Kerou M, Schleper C (2020). Ancestral reconstructions decipher major
adaptations of ammonia-oxidizing archaea upon radiation into moderate terrestrial and marine
environments. *Mbio* **11:** e02371-02320.

14   Ren M, Wang J (2022). Phylogenetic divergence and adaptation of Nitrososphaeria
across lake depths and freshwater ecosystems. *ISME J* **16:** 1491-1501.

15   Ngugi DK, Salcher MM, Andrei A-S, Ghai R, Klotz F, Chiriac M-C *et al* (2023). Postglacial
adaptations enabled colonization and quasi-clonal dispersal of ammonia-oxidizing archaea in
modern European large lakes. *Sci Adv* **9:** eadc9392.

16   Ren ML, Feng XY, Huang YJ, Wang H, Hu Z, Clingenpeel S *et al* (2019). Phylogenomics
suggests oxygen availability as a driving force in Thaumarchaeota evolution. *ISME J* **13:** 2150-
2161.

17   Herbold CW, Lehtovirta-Morley LE, Jung M-Y, Jehmlich N, Hausmann B, Han P *et al*
(2017). Ammonia-oxidising archaea living at low pH: Insights from comparative genomics.
Environ Microbiol **19:** 4939-4952.

18   Qin W, Heal KR, Ramdasi R, Kobelt JN, Martens-Habbena W, Bertagnolli AD *et al* (2017).
Nitrosopumilus maritimus gen. nov., sp nov., Nitrosopumilus cobalaminigenes sp nov.,

675 Nitrosopumilus oxyclinae sp nov., and Nitrosopumilus ureiphilus sp nov., four marine ammonia-
676 oxidizing archaea of the phylum Thaumarchaeota. *Int J Syst Evol Microbiol* **67:** 5067-5079.
677

678 19 Prosser JI, Nicol GW (2016). Candidatus Nitrosotaleales. In: Trujillo ME, Dedysh S, DeVos
679 P, Hedlund B, Kämpfer P, Rainey FA *et al* (eds). *Bergey's Manual of Systematics of Archaea and*
680 *Bacteria*. pp 1-1.
681

682 20 Stieglmeier M, Klingl A, Alves RJE, Rittmann SKMR, Melcher M, Leisch N *et al* (2014).
683 Nitrososphaera viennensis gen. nov., sp nov., an aerobic and mesophilic, ammonia-oxidizing
684 archaeon from soil and a member of the archaeal phylum Thaumarchaeota. *Int J Syst Evol*
685 *Microbiol* **64:** 2738-2752.
686

687 21 Qin W, Jewell TNM, Russell VV, Hedlund BP, de la Torre JR, Stahl DA (2017). Candidatus
688 Nitrosocaldales. In: Trujillo ME, Dedysh S, DeVos P, Hedlund B, Kämpfer P, Rainey FA *et al* (eds).
689 *Bergey's Manual of Systematics of Archaea and Bacteria*. pp 1-2.
690

691 22 Alves RJE, Minh BQ, Urich T, von Haeseler A, Schleper C (2018). Unifying the global
692 phylogeny and environmental distribution of ammonia-oxidising archaea based on *amoA* genes.
693 *Nat Commun* **9:** 1517.
694

695 23 Gubry-Rangin C, Kratsch C, Williams TA, McHardy AC, Embley TM, Prosser JI *et al* (2015).
696 Coupling of diversification and pH adaptation during the evolution of terrestrial
697 Thaumarchaeota. *P Natl Acad Sci USA* **112:** 9370-9375.
698

699 24 Zhong HH, Lehtovirta-Morley L, Liu JW, Zheng YF, Lin HY, Song DL *et al* (2020). Novel
700 insights into the Thaumarchaeota in the deepest oceans: their metabolism and potential
701 adaptation mechanisms. *Microbiome* **8:** 78.
702

703 25 Jing G, Liu L, Wang Z, Zhang Y, Qian L, Gao C *et al* (2021). Microbiome Search Engine 2: a
704 platform for taxonomic and functional search of global microbiomes on the whole-microbiome
705 level. *Msystems* **6:** e00943-00920.
706

707 26 Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW (2015). CheckM:
708 assessing the quality of microbial genomes recovered from isolates, single cells, and
709 metagenomes. *Genome Res* **25:** 1043-1055.
710

711 27 Rognes T, Flouri T, Nichols B, Quince C, Mahé F (2016). VSEARCH: a versatile open
712 source tool for metagenomics. *PeerJ* **4:** e2584.
713

714 28 Su X, Jing G, McDonald D, Wang H, Wang Z, Gonzalez A *et al* (2018). Identifying and
715 predicting novelty in microbiome studies. *Mbio* **9:** e02099-02018.
716

717 29 Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G *et al* (2015).
718 Structure and function of the global ocean microbiome. *Science* **348:** 1261359.

720   30   Duarte CM (2015). Seafaring in the 21St Century: The Malaspina 2010 Circumnavigation
721   Expedition. *Limnology and Oceanography Bulletin* **24**: 11-14.
722

723   31   Mende DR, Bryant JA, Aylward FO, Eppley JM, Nielsen T, Karl DM *et al* (2017).
724   Environmental drivers of a microbial genomic transition zone in the ocean's interior. *Nat*
725   *Microbiol* **2**: 1367-1373.
726

727   32   Liu J, Zheng Y, Lin H, Wang X, Li M, Liu Y *et al* (2019). Proliferation of hydrocarbon-
728   degrading microbes at the bottom of the Mariana Trench. *Microbiome* **7**: 47.
729

730   33   Tian R, Ning D, He Z, Zhang P, Spencer SJ, Gao S *et al* (2020). Small and mighty:
731   adaptation of superphylum Patescibacteria to groundwater environment drives their genome
732   simplicity. *Microbiome* **8**: 51.
733

734   34   Bolger AM, Lohse M, Usadel B (2014). Trimmomatic: a flexible trimmer for Illumina
735   sequence data. *Bioinformatics* **30**: 2114-2120.
736

737   35   Buchfink B, Xie C, Huson DH (2015). Fast and sensitive protein alignment using
738   DIAMOND. *Nat Methods* **12**: 59-60.
739

740   36   Li L, Stoeckert CJ, Jr., Roos DS (2003). OrthoMCL: identification of ortholog groups for
741   eukaryotic genomes. *Genome Res* **13**: 2178-2189.
742

743   37   Enright AJ, Van Dongen S, Ouzounis CA (2002). An efficient algorithm for large-scale
744   detection of protein families. *Nucleic Acids Res* **30**: 1575-1584.
745

746   38   Eren AM, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS *et al* (2021). Community-led,
747   integrated, reproducible multi-omics with anvi'o. *Nat Microbiol* **6**: 3-6.
748

749   39   Abby SS, Melcher M, Kerou M, Krupovic M, Stieglmeier M, Rossel C *et al* (2018).
750   Candidatus Nitrosocaldus cavascurensis, an ammonia oxidizing, extremely thermophilic
751   archaeon with a highly mobile genome. *Front Microbiol* **9**: 28.
752

753   40   Walker CB, de la Torre JR, Klotz MG, Urakawa H, Pinel N, Arp DJ *et al* (2010).
754   *Nitrosopumilus maritimus* genome reveals unique mechanisms for nitrification and autotrophy
755   in globally distributed marine crenarchaea. *P Natl Acad Sci USA* **107**: 8818-8823.
756

757   41   Saier MH, Reddy VS, Moreno-Hagelsieb G, Hendargo KJ, Zhang Y, Iddamsetty V *et al*
758   (2021). The Transporter Classification Database (TCDB): 2021 update. *Nucleic Acids Res* **49**:
759   D461-D467.
760

761   42   Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C, Stadler PF *et al* (2011).
762   ViennaRNA Package 2.0. *Algorithms for Molecular Biology* **6**: 26.

764 43   Guéguen L, Gaillard S, Boussau B, Gouy M, Groussin M, Rochette NC *et al* (2013). Bio++:
765 efficient extensible libraries and tools for computational molecular evolution. *Mol Biol Evol* **30:**
766 1745-1750.
767

768 44   Besemer J, Lomsadze A, Borodovsky M (2001). GeneMarkS: a self-training method for
769 prediction of gene starts in microbial genomes. Implications for finding sequence motifs in
770 regulatory regions. *Nucleic Acids Res* **29:** 2607-2618.
771

772 45   Katoh K, Standley DM (2013). MAFFT Multiple Sequence Alignment Software Version 7:
773 improvements in performance and usability. *Mol Biol Evol* **30:** 772-780.
774

775 46   Talavera G, Castresana J (2007). Improvement of phylogenies after removing divergent
776 and ambiguously aligned blocks from protein sequence alignments. *Syst Biol* **56:** 564-577.
777

778 47   Darriba D, Taboada GL, Doallo R, Posada D (2011). ProtTest 3: fast selection of best-fit
779 models of protein evolution. *Bioinformatics* **27:** 1164-1165.
780

781 48   Stamatakis A (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis
782 of large phylogenies. *Bioinformatics* **30:** 1312-1313.
783

784 49   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A *et al*
785 (2020). IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic
786 era. *Mol Biol Evol* **37:** 1530-1534.
787

788 50   Talavera G, Castresana J (2007). Improvement of phylogenies after removing divergent
789 and ambiguously aligned blocks from protein sequence alignments. *Syst Biol* **56:** 564-577.
790

791 51   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017).
792 ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14:** 587-
793 589.
794

795 52   Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS (2017). UFBoot2: improving
796 the ultrafast bootstrap approximation. *Mol Biol Evol* **35:** 518-522.
797

798 53   Pankratova G, Hederstedt L, Gorton L (2019). Extracellular electron transfer features of
799 Gram-positive bacteria. *Anal Chim Acta* **1076:** 32-47.
800

801 54   Hallam SJ, Konstantinidis KT, Putnam N, Schleper C, Watanabe Y, Sugahara J *et al* (2006).
802 Genomic analysis of the uncultivated marine crenarchaeote Cenarchaeum symbiosum. *P Natl*
803 *Acad Sci USA* **103:** 18296-18301.
804

805   55   Moeller FU, Webster NS, Herbold CW, Behnam F, Domman D, Albertsen M *et al* (2019).
806 Characterization of a thaumarchaeal symbiont that drives incomplete nitrification in the
807 tropical sponge Ianthella basta. *Environ Microbiol* **21**: 3831-3854.
808

809   56   Zhang S, Song WZ, Wemheuer B, Reveillaud J, Webster N, Thomas T (2019).
810 Comparative genomics reveals ecological and evolutionary insights into sponge-associated
811 Thaumarchaeota. *Msystems* **4**: e00288-00219.
812

813   57   Turon M, Uriz MJ (2020). New insights into the archaeal consortium of tropical sponges.
814 *Front Mar Sci* **6**: 789.
815

816   58   Steinert G, Busch K, Bayer K, Kodami S, Arbizu PM, Kelly M *et al* (2020). Compositional
817 and quantitative insights into bacterial and archaeal communities of South Pacific deep-sea
818 sponges (Demospongiae and Hexactinellida). *Front Microbiol* **11**: 716.
819

820   59   Haber M, Burgsdorf I, Handley KM, Rubin-Blum M, Steindler L (2021). Genomic insights
821 into the lifestyles of Thaumarchaeota inside sponges. *Front Microbiol* **11**: 622824.
822

823   60   Van Soest RWM, Boury-Esnault N, Vacelet J, Dohrmann M, Erpenbeck D, De Voogd NJ *et*
824 *al* (2012). Global diversity of sponges (Porifera). *PLoS One* **7**: e35105.
825

826   61   Lee OO, Wang Y, Yang JK, Lafi FF, Al-Suwailem A, Qian PY (2011). Pyrosequencing reveals
827 highly diverse and species-specific microbial communities in sponges from the Red Sea. *ISME J* **5**:
828 650-664.
829

830   62   Lee E-Y, Lee HK, Lee YK, Sim CJ, Lee J-H (2003). Diversity of symbiotic archaeal
831 communities in marine sponges from Korea. *Biomed Eng* **20**: 299-304.
832

833   63   Yang Z, Li Z (2012). Spatial distribution of prokaryotic symbionts and ammoxidation,
834 denitrifier bacteria in marine sponge Astrosclera willeyana. *Scientific Reports* **2**: 528.
835

836   64   Liu F, Han M, Zhang F, Zhang B, Li Z (2011). Distribution and abundance of Archaea in
837 South China Sea sponge Holoxea sp. and the presence of ammonia-oxidizing archaea in sponge
838 cells. *Evid Based Complementary Altern Med* **2011**: 723696.
839

840   65   Reji L, Cardarelli EL, Boye K, Bargar JR, Francis CA (2022). Diverse ecophysiological
841 adaptations of subsurface Thaumarchaeota in floodplain sediments revealed through genome-
842 resolved metagenomics. *ISME J* **16**: 1140-1152.
843

844   66   Merino N, Jackson TR, Campbell JH, Kersting AB, Sackett J, Fisher JC *et al* (2022).
845 Subsurface microbial communities as a tool for characterizing regional-scale groundwater flow.
846 *Sci Total Environ* **842**: 156768.
847

67   Watson DB, Kostka JE, Fields MW, Jardine PM (2004). The Oak Ridge Field Research
Center conceptual model.

68   Hemme CL, Tu QC, Shi Z, Qin YJ, Gao WM, Deng Y *et al* (2015). Comparative
metagenomics reveals impact of contaminants on groundwater microbiomes. *Front Microbiol* **6:**
1205.

69   Luo ZH, Rao MPN, Chen H, Hua ZS, Li Q, Hedlund BP *et al* (2021). Genomic insights of
"Candidatus Nitrosocaldaceae" based on nine new metagenome-assembled genomes, including
"Candidatus Nitrosothermus" gen nov. and two new species of "Candidatus Nitrosocaldus".
Front Microbiol **11:** 608832.

70   Daebeler A, Herbold CW, Vierheilig J, Sedlacek CJ, Pjevac P, Albertsen M *et al* (2018).
Cultivation and genomic analysis of "Candidatus Nitrosocaldus islandicus," an obligately
thermophilic, ammonia-oxidizing Thaumarchaeon from a hot spring biofilm in Graendalur
Valley, Iceland. *Front Microbiol* **9:** 193.

71   Koch H, Lücker S, Albertsen M, Kitzinger K, Herbold C, Spieck E *et al* (2015). Expanded
metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the genus *Nitrospira*. *P Natl*
Acad Sci USA **112:** 11371-11376.

72   Jing H, Takagi J, Liu J-h, Lindgren S, Zhang R-g, Joachimiak A *et al* (2002). Archaeal
surface layer proteins contain β propeller, PKD, and β helix domains and are related to
metazoan cell surface proteins. *Structure* **10:** 1453-1464.

73   Tschitschko B, Williams TJ, Allen MA, Páez-Espino D, Kyrpides N, Zhong L *et al* (2015).
Antarctic archaea–virus interactions: metaproteome-led analysis of invasion, evasion and
adaptation. *ISME J* **9:** 2094-2107.

74   Hodgskiss LH, Melcher M, Kerou M, Chen W, Ponce-Toledo RI, Savvides SN *et al* (2023).
Unexpected complexity of the ammonia monooxygenase in archaea. *ISME J.*

75   Wang BZ, Qin W, Ren Y, Zhou X, Jung MY, Han P *et al* (2019). Expansion of
Thaumarchaeota habitat range is correlated with horizontal transfer of ATPase operons. *ISME J*
13: 3067-3079.

76   Kerou M, Ponce-Toledo RI, Zhao R, Abby SS, Hirai M, Nomaki H *et al* (2021). Genomes of
Thaumarchaeota from deep sea sediments reveal specific adaptations of three independently
evolved lineages. *ISME J* **15:** 2792-2808.

77   Hwang Y, Schulze-Makuch D, Arens FL, Saenz JS, Adam PS, Sager C *et al* (2021). Leave no
stone unturned: individually adapted xerotolerant Thaumarchaeota sheltered below the
boulders of the Atacama Desert hyperarid core. *Microbiome* **9:** 234.

78   Kozlowski JA, Stieglmeier M, Schleper C, Klotz MG, Stein LY (2016). Pathways and key
intermediates required for obligate aerobic ammonia-dependent chemolithotrophy in bacteria
and Thaumarchaeota. *ISME J* **10:** 1836-1845.
895
896 79   Groussin M, Gouy M (2011). Adaptation to environmental temperature is a major
897 determinant of molecular evolutionary rates in archaea. *Mol Biol Evol* **28:** 2661-2674.
898
899 80   de La Torre JR, Walker CB, Ingalls AE, Könneke M, Stahl DA (2008). Cultivation of a
900 thermophilic ammonia oxidizing archaeon synthesizing crenarchaeol. *Environ Microbiol* **10:** 810-
901 818.
902
903 81   Stolper DA, Bucholz CE (2019). Neoproterozoic to early Phanerozoic rise in island arc
904 redox state due to deep ocean oxygenation and increased marine sulfate levels. *P Natl Acad Sci*
905 *USA* **116:** 8746-8755.
906
907 82   Price R, Lesniewski R, Nitzsche K, Meyerdierks A, Saltikov C, Pichler T *et al* (2013).
908 Archaeal and bacterial diversity in an arsenic-rich shallow-sea hydrothermal system undergoing
909 phase separation. *Front Microbiol* **4**.
910
911 83   Kato S, Takano Y, Kakegawa T, Oba H, Inoue K, Kobayashi C *et al* (2010). Biogeography
912 and biodiversity in sulfide structures of active and inactive vents at deep-sea hydrothermal
913 fields of the southern Mariana Trough. *Appl Environ Microb* **76:** 2968-2979.
914
915 84   Schrenk MO, Kelley DS, Delaney JR, Baross JA (2003). Incidence and Diversity of
916 Microorganisms within the Walls of an Active Deep-Sea Sulfide Chimney. *Appl Environ Microb*
917 **69:** 3580-3592.
918
919 85   Lins-de-Barros MM, Vieira RP, Cardoso AM, Monteiro VA, Turque AS, Silveira CB *et al*
920 (2010). Archaea, Bacteria, and Algal Plastids Associated with the Reef-Building Corals
921 Siderastrea stellata and Mussismilia hispida from Búzios, South Atlantic Ocean, Brazil. *Microb*
922 *Ecol* **59:** 523-532.
923
924 86   Takano Y, Chikaraishi Y, Ogawa NO, Nomaki H, Morono Y, Inagaki F *et al* (2010).
925 Sedimentary membrane lipids recycled by deep-sea benthic archaea. *Nat Geosci* **3:** 858-861.
926
927

Major lineages

- *Nitrosopumilales*
- *Nitrososphaerales*
- *Ca.* Nitrosomirales
- *Ca.* Nitrosocaldales
- Non-ammonia-oxidizing Thaumarchaeota

Habitats (outer ring)

- Subsurface environments
- Soil
- Geothermal environments
- Marine water/sediments
- Sponge

Habitat temperature range (inner ring)

- <20°C
- 20-45°C
- 45-60°C
- >60°C
- Not available

★ Genome

a

- ● Subsurface environments
- ● Soil
- ● Geothermal environments
- ● Marine water/sediments
- ● Sponge

b

0% 50% 100%

sponge metagenome (Porifera)
sponge tissue (Aciculites cribrophora)
sponge tissue (Aciculites cribrophora)
sponge tissue (Aciculites cribrophora)
Sponge tissue (unknown deep sea sponge)
sponge tissue
sponge tissue
Sponge tissue (unknown deep sea sponge)
Sponge tissue (unknown deep sea sponge)
sponge tissue
Sponge tissue (unknown deep sea sponge)
sponge tissue
Sponge tissue (Leiodermatium spp)

Well (Yucca Flat)
Well (Oasis Valley)
Well (Pahute Mesa)
Spring (Ash Meadows)
Spring (Ash Meadows)
Spring (Ash Meadows)
Groundwater (Hanford site)
Groundwater (Hanford site)
Well (Pahute Mesa)
Groundwater (Hanford site)
Spring (Ash Meadows)
Groundwater (Hanford site)
Well (Spring Mountains)

Sponge

Subsurface environments

- ● *Ca.* Nitrosomirales AOA
- ● *Nitrosopumilales* AOA
- ● *Nitrososphaerales* AOA
- ● *Ca.* Nitrosocaldales AOA

Nitrosopumilales *Nitrososphaerales* *Ca.* Nitrosomirales *Ca.* Nitrosocaldales

Ammonia oxidation
HP/HB carbon fixation
Vitamin B12 bisynthesis
Formate transporter
Formate dehydrogenase, major subunit
Nitrate reductase, alpha subunit
Nitrate reductase, beta subunit
Nitrosomirales-specific cupredoxin blue copper protein
Nitrosomirales-specific cupredoxin super family
copA
copD
Nitrosomirales-specific putative S-layer protein
Nitrosomirales-specific putative S-layer protein
Putative PKD domain protein
Putative PKD domain protein
Putative PKD domain protein