Words in Title: 12
Words in abstract: 146
Words in main text: 5990
Words in methods: 2015
Number of figures: 6
Number of supplementary figures: 4

Somatodendritic orientation determines tDCS-induced neuromodulation of Purkinje cell activity in awake mice.

Carlos A Sánchez-León[1,2], Guillermo Sánchez-Garrido Campos[1], Marta Fernández[3,4], Alvaro Sánchez-López[5], Javier F Medina[5] and Javier Márquez-Ruiz [1*]

[1]Department of Physiology, Anatomy and Cell Biology, Pablo de Olavide University, Ctra. de Utrera, km. 1, 41013, Seville, Spain. [2]Department of Neurology and Neurobiology, University of California Los Angeles, Los Angeles 90095, USA. [3] Department of Psychiatry, University of California Los Angeles, Los Angeles 90095, USA. [4] Department of Pharmacology, University of the Basque Country (UPV/EHU), Leioa 48940, Spain. [5]Department of Neuroscience, Baylor College of Medicine, Houston, TX, USA

Running title: tDCS neuromodulation of Purkinje cells *in vivo*

Acknowledgments: This work was supported by grants from the Spanish MINECO-FEDER (BFU2014-53820-P and BFU2017-89615-P) and FET European Union's Horizon 2020 research and innovation program (grant agreement No 101017716) to J.M-R and from the US National Institutes of Health (R01MH093727 & R01NS112917) to J.F.M and (RF1MH114269) to J.F.M and J.M-R. C.A.S-L was in receipt of an FPU grant from the Spanish Government (FPU13/04858).

Keywords: cerebellum, non-invasive brain stimulation, transcranial electrical stimulation, transcranial direct-current stimulation, high-density recordings, juxtacellular

Corresponding author:
Javier Márquez-Ruiz, Ph.D.
Brain Stimulation translational Laboratory
Universidad Pablo de Olavide
Ctra. de Utrera, km. 1
41013-Sevilla, Spain
E-mail: jmarquez@upo.es
Phone: +34-954-978054

Declaration of interests: The authors declare no competing financial interests.

Abstract

Transcranial direct-current stimulation (tDCS) of the cerebellum is a promising non-invasive neuromodulatory technique being proposed for the treatment of neurological and neuropsychiatric disorders. However, there is a lack of knowledge about how externally applied currents affect neuronal spiking activity in cerebellar circuits *in vivo*. In this study, we observe that tDCS induces a heterogeneous polarity-dependent modulation of the firing rate of Purkinje cells (PC) and non-PC in the mouse cerebellar cortex. Using a combination of juxtacellular labeling and high-density Neuropixels recordings, we demonstrate that the apparently heterogeneous effects of tDCS on PC activity can be explained by taking into account the somatodendritic orientation relative to the electric field. Our findings emphasize the importance of considering neuronal orientation and morphological aspects to increase the predictive power of tDCS computational models, enhance the reliability of current stimulation protocols and optimize desired effects in basic and clinical human applications.

Introduction

Transcranial direct-current stimulation (tDCS) is a non-invasive brain stimulation technique consisting in the application of constant weak electric currents over the scalp for several minutes through strategically positioned electrodes[1,2]. In the last two decades, tDCS has shown promising results in modulating cognitive, behavioral, and clinical traits by applying stimulation on various brain regions[3]. Cerebellar tDCS has been found to modulate both motor and also cognitive and emotional processing. This has led to its proposal as a noninvasive neuromodulatory therapy for treating cerebellum-related disorders[4,5]. As the popularity of tDCS grows, there are increasing debates regarding the variability in results among studies[6–9]. This unreliability can be attributed to methodological differences and the lack of knowledge about the physiological mechanisms of action of tDCS[10–12]. To optimize the effects of tDCS in humans, it is essential to understand the neural basis underlying this variability and develop new strategies accordingly.

Animal models have been instrumental in advancing our understanding of the mechanisms underlying tDCS effects[13], defining safety parameters[14,15], and inspiring new stimulation protocols[16]. *In vitro* studies have highlighted the importance of neuronal orientation and morphology[17–19] in modulating individual neurons' excitability. In addition, *in vivo* experiments also enable the study of the impact of tDCS on behaving brains and provide insights into the underlying neuronal mechanisms at both behavioral and physiological levels. For example, *in vivo* animal models have demonstrated the physiological effects of transcranial electrical stimulation (tES) on spike timing[20,21], local field potential oscillations[22,23], memory and learning processes[23,24], as well as the involvement of non-neuronal elements like astrocytes and microglial cells[25,26]. *In vivo* experiments also allow for the measurement of the actual intracranial electric field induced by tES[22,27] and for testing the efficacy of tES protocols at both behavioral and physiological levels[28,29]. However, *in vivo* animal models often have limited control over the applied electric fields compared to *in vitro* experiments and may lack the ability to identify which specific cell types are impacted by the stimulation. Here, we overcome these obstacles by examining the impact of tDCS on the activity of identifiable neurons in the cerebellum of mice.

Purkinje cells (PCs) are the output neurons of the cerebellar cortex and can be electrophysiologically identified by their ability to fire both complex and simple spikes[30]. Unlike the cerebral cortex, the cerebellar cortex of rodents is highly convoluted, similar to that of humans[31]. In this study, the highly-folded mouse cerebellum was used to determine the impact of the PC orientation on the final neuronal modulation induced by tDCS in awake mice. Single-neuron extracellular recording of PCs in awake mice was combined with juxtacellular recordings and subsequent staining of PC with neurobiotin under anesthesia to investigate the effects of cerebellar tDCS on the cerebellar network output. The morphological reconstruction of the recorded PCs allowed the correlation of their neuronal orientation with their response to tDCS. Finally, a high-density Neuropixels recording system was used to demonstrate the relevance of neuronal orientation in awake mice by simultaneously recording PCs with opposing orientations during the application of tDCS.

Our work offers *in vivo* evidence that points to neuronal orientation as a crucial factor in determining the impact of tDCS on neural activity, and can explain the reason why the effects are heterogeneous (or even opposite) across different layers of the same brain area. This result is essential for developing accurate computational models and emphasizes the need to consider neuronal orientation when predicting tDCS effects. Our findings suggest that high-definition tDCS electrodes in which the direction of the electric field can be flexibly controlled may be useful for enhancing the reliability of stimulation protocols and optimizing the desired tDCS effects in various cerebellum-related disorders[4,5].

Methods

Animal preparation.

Experiments were carried out on adult C57 male mice (n = 77) (University of Seville, Spain) weighting 28–35 g. All experimental procedures were carried out in accordance with European Union guidelines (2010/63/CE) and Spanish regulations (RD 53/2013) for the use of laboratory animals in chronic experiments. In addition, these experiments were submitted to and approved by the local Ethics Committee of the Pablo de Olavide University (Seville, Spain). Mice were prepared for simultaneous tES administration and electrophysiological recordings

in the lateral (left) or vermis region of the cerebellar cortex in the head-restrained awake animal, following surgical procedures described previously[22]. In brief, animals were anesthetized with a ketamine–xylazine mixture (Ketaset, 100 mg/ml, Zoetis, NJ., USA; Rompun, 20 mg/ml, Bayer, Leverkusen, Germany), and a custom-made chlorinated silver ring electrode (2.5 mm inner ø, 3.5 mm outer ø) was placed over the skull centered on left crus I-II (AP = − 6 mm; L = +2 mm; relative to bregma[32]) (Fig. 1a) or on the cerebellar vermis (AP = − 6 mm; L = 0 mm; relative to bregma) and fixed with dental cement (DuraLay, Ill., USA). A 2 mm ø craniotomy was made centered in the ring and exposing the cerebellar cortex. The dura was left intact and protected with wax bone (Ethicon, Johnson & Johnson) until recordings begun. In addition, a silver wire electrode (ø: 381 µm, A-M Systems) was also implanted over the dura surface under the left parietal bone (AP = − 0.9 mm; L = + 3 mm; relative to bregma) as electrical reference for the electrophysiological recordings. Finally, a head-holding system was implanted, consisting of three bolts screwed to the skull and a bolt placed over the skull upside down and perpendicular to the frontal plane to allow for head fixation during the experiments. The holding system was cemented to the skull.

Single unit recordings.

Recording sessions began at least two days after surgery. The animals were placed over a treadmill with an infrared sensor for locomotion activity monitoring and the head was fixed to the recording table by means of the implanted head-holding system. Bone wax was removed with the aid of a surgical microscope (SMZ-140, Motic, Barcelona, Spain) and the cortical surface was carefully cleaned with super fine forceps (Dumont #5, FST, Heidelberg, Germany) and cotton swab without damaging the dura mater.

All single-cell recordings were carried out with an amplifier (BVC-700A, Dagan corporation, MN., USA) connected to a dual extracellular-intracellular headstage (8024, Dagan corporation, MN., USA; gain error ± 1%, noise 4.5 µV root mean square). The single-cell recordings were performed with a glass micropipette (impedance 1–10 MΩ) filled with 3M NaCl, mounted on a micromanipulator (MO-10, Narishige, Tokyo, Japan). The electrode was slowly lowered at ~2 µm/s and spikes were detected based on visual (2002C and 2004C, Tektronix, OR., USA) and auditory (Audio monitor 3300, A-M Systems, WA., USA) cues. Once the spiking activity was detected, the micropipette tip was advanced slowly to properly isolate and identify single neuron activity in the recorded signal.

Juxtacellular labeling.

The procedure was similar to that of single-cell recordings except that the micropipette was filled with 2% Neurobiotin (SP-1120, Vector Laboratories, CA., USA) in 0.5 M NaCl, the tip was immersed in DiI (Vybrant DiI cell-labeling, V22885, Thermo Fisher Scientific, Mass., USA) and the impedance was periodically checked to assure that it was between 4–12 MΩ. With the headstage in extracellular mode and after single-cell activity was isolated, the micropipette tip was advanced until the negative spikes (extracellular recording) became positive spikes (juxtacellular recording) with an amplitude of at least 600 μV. Then, the headstage was switched to intracellular mode to juxtacellularly label the neuron following the method described by Pinault[33]. The firing rate of recorded neurons were modulated by passing positive current pulses (200 ms ON/OFF) at increasing intensities (1-10 nA) through the micropipette tip. After a delay of a few seconds, the electrical properties of the recorded neuron suddenly and significantly changed, increasing its firing rate and broadening the spike waveform. From this critical moment, pulse intensity was lowered to prevent cellular damage and the modulation was maintained from several seconds to minutes in order to fill the neuron with neurobiotin.

Neuropixels recording.

All Neuropixels recordings were performed using SpikeGLX (http://billkarsh.github.io/SpikeGLX/) on a computer connected to the PXIe acquisition module. Action potentials were band filtered between 0.3 and 10 kHz and sampled at 30 kHz whereas simultaneous LFPs were band filtered between 0.5 and 500 Hz and sampled at 2.5 kHz. Neuropixels's probe was coated with DiI lipophilic dye before insertion in the brain so a precise mark of the recording tract would be visible at confocal microscope. The probe was lowered in the coronal plane at 90 degrees from horizontal plane at ~2 μm/s until 4000 μm below cerebellar cortex surface. Neuropixels's probe was left to settle for 10 min to avoid drift during the recording.

Transcranial electrical stimulation.

The different protocols for transcranial currents were designed in Spike2 (Cambridge Electronic Design, CED, Cambridge, U.K.) and sent to a battery-driven linear stimulus isolator (WPI A395, Fl., USA) through an analog output from the acquisition board (CED micro1401-

3). tES was applied between the ring-electrode and a reference electrode consisting of a rubber rectangular pad (6 cm^2) attached to the back of the mice and moisten with electrogel (Electro-Cap International, OH., USA). To measure the actual voltage changes elicited intracranially, sinusoid alternating current waves were delivered at amplitudes of ± 2, ± 20 and ± 200 µA (± 0.0426, ± 0.426 and ± 4.26 mA/cm^2) at 1 Hz and recorded in steps of 1 mm from cortical surface to 4 mm depth. To characterize the effects induced by tDCS, trials of 15 or 20 s pulses at 100, 200 and 300 µA anodal and cathodal tDCS (including 5 s ramp-in and 5 s ramp-out) were applied separated by 10 s of non-stimulation.

Histology.

To reconstruct the neurobiotin-labeled neurons, mice were deeply anesthetized with ketamine–xylazine mixture (Ketaset, 100 mg/ml; Rompun, 20 mg/ml) 15 min after juxtacellular labeling and perfused transcardially with 0.9% saline followed by 4% paraformaldehyde (Panreac, Barcelona, Spain) in PBS (0.1 M, pH ~7,4). The brains were removed and stored in 4% paraformaldehyde for 24 hours, cryoprotected in 30% sucrose in PBS the next 48 hours, and then cut into 50 µm coronal slices with a freezing microtome (CM1520, Leica, Wetzlar, Germany). After three washes with PBS-Triton X-100 1% (PBS-Tx, Sigma-Aldrich, Mo., USA), sections containing neurobiotin-labelled neurons were blocked with 10% Normal Donkey Serum (NDS, 566460, Merck, Darmstadt, Germany) in PBS-Tx and then incubated overnight at room temperature in darkness with Streptavidin 1:200 (Streptavidin DyLight 488 conjugated, Thermo Fisher Scientific) in PBS-Tx. After three washes with PBS, sections were mounted on glass slides and coverslipped using Dako fluorescence mounting medium (Agilent Technologies, Santa Clara, CA, USA). To determine recording regions across cerebellar tissue in Neuropixels recordings, the same process was carried out with the exception that after the three washes in PBS-Triton X-100 1%, slices were incubated for 3 min with Hoechst 33342 dye (Merck Millipore, Billerica, MA, USA) (2µg/ml) in PBS with 0.25% Triton X-100. For confocal imaging, an *in vivo* confocal microscope (A1R HD25, Nikon, Tokyo, Japan) was used. Z-series of optical sections (0.5 µm apart) were obtained using the sequential scanning mode.

Data analysis.

Data collection. Spike activity was recorded with a glass micropipette or a Neuropixels probe

and sampled at 25 (CED micro1401-3) or 30 kHz (IMEC-PXIe) with an amplitude resolution of 12 and 10 bits, respectively. When necessary, LFP were sampled at 2.5 kHz and the remaining non-neuronal activities (tES, juxtacelullar injected currents and wheel movement) were sampled at 5 kHz.

Intracranial electric field analysis. The peak-to-peak amplitude (electric potential) of the LFP oscillations induced by tACS, were averaged for a given intensity and depth. For every intensity, the electric field strength (differences between potentials) was calculated by computing the difference in peak-to-peak values between two consecutive depths (1 mm in distance).

Single-cell activity. For glass micropipette recordings, only well isolated neurons, with high signal-to-noise ratio (at least 4 times the standard deviation of background noise) were considered for analysis. Spikes were detected offline in Spike2 (CED) and exported to Matlab 2015a (MathWorks Inc., MA., USA) software for analysis. Trials where the mouse was running were removed from analysis. For spike detection, a "DC remove" process (time constant (s): 0.001-0.0004) was applied to reduce DC level drifts, and spikes were detected based on threshold-crossing algorithm of Spike2 software. After that, the DC remove process was carried out, and all spikes were visually confirmed and PC identified as such if complex spikes (CS) were observed and had at least a 10-40 ms pause in simple spikes (SS) after CS occurrence. For Neuropixels recordings, channels showing Purkinje cells, with CS followed by a SS silence, were manually selected on SpikeGLX (http://billkarsh.github.io/SpikeGLX/) and exported to analyze on Spike2 (CED, Cambridge, UK). Spike sorting was carried out using the Spike2 software and spikes with similar waveforms were grouped together in the same templates. Putative Purkinje cell templates were subsequently curated to exclude contamination produced by other units. For this purpose, only periods where the template amplitude was stable were used, and events (spikes) with an amplitude deviation greater than one third of the average template amplitude were excluded. Additionally, the autocorrelogram was checked to discard contaminated templates with violations of the refractory period. Clusters corresponding to putative SS and CS were identified due to their waveform and their firing frequency (<3 Hz for CS and >50Hz for SS) and regularity, producing characteristic shoulders in the SS autocorrelogram. Finally, only those SS and CS from Purkinje cells unambiguously identified by the pause in their crosscorrelogram, were used for the

analysis. Subsequently, each neuron was analyzed in Matlab custom-made script. The 5 s window immediately before a stimulation ramp-in and immediately after a ramp-out were used for control and post-stimulation conditions, while the 5 s window immediately after the stimulation ramp reached the peak intensity was considered for tDCS condition. Averaged SS and CS waveforms, SS frequency, CS frequency and latency of the SS pause after CS were computed and analyzed for each condition. For SS firing rate analysis, all the trials with a given tDCS intensity and duration were averaged and then binned in 100 ms epochs in the five second windows computed for statistical analysis. For CS firing rate and latency of the SS pause after a CS analysis, the procedure was the same as before but instead of averaging between trials, we computed the different parameters for the 5 s windows (before, during and after tDCS) for every trial and the statistical comparisons were made between all the trials with a given tDCS intensity and time. Peristimulus time histograms showing the number of spikes per bin (bin size: 0.1 s for SS and 1 s for CS) were aligned with tDCS ramp-in, normalized and standardized (Z-score=$X-\mu/\sigma$, , where X is the firing rate at each bin, and μ and σ are the average and standard deviation, respectively, of the 5s control window) with respect to the average frequency of the five seconds before anodal and cathodal tDCS ramp-in. To compare the strength of the modulation for the same neuron with different tDCS intensities and between neurons, the different parameters during tDCS were normalized by their values during control condition.

Neurobiotin-labeled neurons. Confocal images were processed in ImageJ (https://imagej.nih.gov/ij/) with the image processing package Fiji (http://fiji.sc/Fiji) to obtain a z-stack reconstruction of the neurobiotin-labelled neurons. The deviation of the somatodendritic axis with respect to the active electrode was calculated by measuring the angle between the neuronal axis and an imaginary line perpendicular to the active electrode.

Statistical analysis. Statistical comparison was inferred by repeated measures ANOVA (RM-ANOVA) using Matlab 2015a. Neural activity parameters from each neuron were compared between the control, the tDCS and the post-stimulation periods. The non-parametric Friedman test was applied for comparisons when data did not permit normality assumption. The results are shown as mean ± SEM. Statistical significance was set at $p < 0.05$ in all cases.

Results

Electric field measurement in the cerebellar cortex

tDCS effects critically depend on the strength of the electric field imposed in the brain. To assess the actual electric field gradient imposed by tDCS in the cerebellar cortex in our experimental design, a group of mice (n = 9) was prepared for the chronic recording of LFPs in awake condition during simultaneous application of low-frequency transcranial alternating-current stimulation (tACS) (1 Hz) at different intensities (± 2, ± 20, and ± 200 µA) (Fig. 1a). LFPs were sequentially recorded every 1 mm from the cortical surface to 4 mm depth (15º rostro-caudal insertion angle) (Fig. 1b). Figure 1 (c) shows the LFP recordings from a representative animal during simultaneous tACS (1 Hz, 10 s, ± 20 µA) at different depths, showing a decrease in voltage for deeper recordings. The estimation of the electric field strength calculated at different depths (1, 2, 3 and 4 mm) and tACS intensities (± 2, circles; ± 20, squares; ± 200 µA, triangles) is represented in Fig. 1d (n = 9). Under the active electrode, the magnitude of the electric field decreased with depth in a logarithmic manner for the three tested intensities (R = 0.98, 0.96 and 0.95 for ± 2, ± 20, and ± 200 µA, respectively; data is presented with logarithmic abscissa axis for visual facilitation, Fig. 1d). With this data, the electric field imposed by tACS at different depths and intensities tested in our experiments was interpolated. A polynomial surface (degree 2 for depth and degree 2 for intensity axis) was fitted on the electric field values and then the coefficients (with 95% confidence bounds) were extracted from the linear model. Considering the most superficial (0.3 mm) and the deepest (2.3 mm) recorded neurons, we could expect electric field values between 60.1, 92.9 and 125.7 V/m (at 0.3 mm for 100, 200 and 300 µA, respectively) and 5.9, 20.2 and 34.6 V/m (at 2.3 mm for 100, 200 and 300 µA, respectively) at the recording places. These values are in line with *in vitro* and other *in vivo* animal studies showing modulation of neuronal firing rate under similar current densities[13].

Figure 1

Fig. 1 Intracranial electric fields induced by Cb-tACS. **a** Experimental design for combined *in vivo* electrophysiology and transcranial alternating current stimulation (tACS) in crus I-II of cerebellar cortex in awake mice, showing silver-ring active and reference (Ref.) electrode locations. Inset (*i.*) shows a schematic sagittal view of the recording chamber design. **b** Schematic representation of a sagittal section of the brain showing the reference (Ref.) and active electrodes location (gray bar) and a representative track in the lateral cerebellum highlighting the depths where the electric field was measured (color dots). **c** tACS stimulation (top trace) applied over the scalp and exemplary recording of the actual field potentials generated at different depths (from 0 to 4 mm) in a representative animal. The traces were overlapped to facilitate amplitude comparison. **d** Average (filled symbols) and individual (empty symbols) electric field strength recorded at different depths for ± 2 (circles), ± 20 (squares) and ± 200 µA (triangles) tACS.

tDCS modulates Purkinje cell activity in awake mice in a heterogeneous manner

To understand how tDCS modulates neuronal firing behavior at a single-cell level, we performed single-cell recordings in awake mice using glass micropipettes. We first focused on the potential impact of exogenous electric fields on PC firing rate. Unlike other cerebellar neuronal types, PCs can be electrophysiologically identified by the presence of high-frequency simple spikes (SS) and less frequent complex spikes (CS) followed by a brief SS silence[30,31] (Fig. 2a,b and Supplementary Fig. 1a-d). To determine the online effects of exogenous electric field

application on PC firing rate and avoid long-term plasticity mechanisms, short pulses of tDCS (15 or 20 s including 5 s ramp-in and 5 s ramp-out, ± 200 μA) separated by non-stimulating periods (10 s) were applied. Figure 2a-f shows two representative PCs and their firing behavior during tDCS (Fig. 2c-f). As observed in the Z-score-transformed average firing rate, the SS firing rate of some recorded PCs significantly increased during anodal and decreased during cathodal tDCS (Fig. 2c,e) whereas the opposite effects were observed in other recorded PCs; decreasing the SS firing rate during anodal and increasing during cathodal (Fig. 2d,f) (Paired Student's t-test, $p < 0.01$). Furthermore, we could observe a significant rebound effect when tDCS was switched off for some of the neurons (Fig. 2d,f, 'After'). A total of 24 identified PCs were recorded in crus I-II region (Fig. 2g,h) in awake mice (n = 24 animals). The impact of anodal and cathodal tDCS on the SS firing rate showed significant differences (filled circles) for 11 out of 24 individual PCs recorded (n = 24, RM-ANOVA or Friedman tests, $p < 0.05$) (Fig. 2g and Supplementary Fig. 2a,b). The data distribution was fitted by a linear model (R = 0.46, $p < 0.0003$), which indicates that most of PCs increased their firing rate with anodal stimulation and decreased with cathodal stimulation (cells in 3rd quadrant of Fig. 2g) or vice versa (cells in 1st quadrant of Fig. 2g).

No differences were observed in the waveform of recorded SS nor CS during anodal nor cathodal tDCS with respect to control condition (Supplementary Fig. 1a-d). No significant changes were observed in the CS firing rate during tDCS (Supplementary Fig. 1e-h) (n = 25, Paired Student's t-test) except for one of the recorded PCs. The impact of anodal and cathodal tDCS in CS firing rate and the CS-driven SS silence of the individual PCs recorded in crus I-II of awake mice was not significant for most cells (Supplementary Fig. 1i,j).

From these results, we can conclude that PCs, constituting the only output from cerebellar cortex, 1) are modulated by tDCS based on the electric field gradient applied in this experiment and 2) this modulatory effect is heterogeneous across the PC population.

Fig. 2 tDCS modulation of PC activity in crus I/II of awake mice. a, b Recording of spontaneous firing activity of two PCs showing the presence of SS and CS. **c,d** Z-score-transformed average PSTH (bin size: 0.1 s) of the SS activity of the 2 PCs shown in a,b before, during and after anodal (red trace) or cathodal (blue trace) tDCS. **e,f** Statistical comparison of the SS firing rate of the 2 PCs shown in a,b, measured in 5 s windows before, during and after tDCS (RM-ANOVA or Friedman tests, $p < 0.05$). Error bars represent SEM. **p < 0.01; ***p < 0.001. **g** Modulation of SS (p) of individual PCs (circles) during anodal (red) and cathodal (blue) tDCS. Filled circles represent statistically significant modulation during tDCS (n = 24, RM-ANOVA or Friedman tests, $p < 0.05$). **h** Schematic representation of the recording sites and active electrode location (gray bar) during tDCS.

Non-Purkinje cell activity is also modulated by tDCS in awake mice

Beyond the inhibitory PCs, the cerebellar cortex neuronal network also contains excitatory

(granule cell) and inhibitory (Golgi cell, Lugaro cells, basket cells and stellate cells) neurons responsible for determining the spatio-temporal PC output[34]. The modulatory effects of tDCS on the activity of these non-PC neurons could be important in feed-forward, feedback, and lateral inhibition processes underlying cerebellar function. Unlike PCs, these neurons do not show typical SS and CS in the neuronal recording and cannot be electrophysiologically identified. For analysis purposes, we decided to include these neurons in "non-PC" group. The same tDCS protocol previously used with PCs was applied to determine the online effects of exogenous electric field application on non-PC firing rate.

Fig. 3a-f shows two representative non-PCs and their firing behavior during tDCS. As observed in the Z-score-transformed average, the firing rate of some recorded non-PCs significantly increased during anodal and decreased during cathodal tDCS (Fig. 3c,d) whereas different effects were observed in other recorded neurons (Fig. 3e,f) (Paired Student's t-test, $p < 0.05$). Furthermore, similar to what we previously observed in PCs, we could observe a significant rebound effect when tDCS was switched off for some of the neurons (Fig. 3c-f). A total of 50 non-PCs were recorded in crus I-II region (Fig. 3g,h) in awake mice (n = 24 animals). The impact of anodal and cathodal tDCS in the firing rate showed significant differences (filled circles) for 35 out of 50 individual non-PCs recorded (n = 50, RM-ANOVA or Friedman tests, $p < 0.05$) (Fig. 3g and Supplementary Fig. 2c,d). The data distribution was fitted by a linear model (R = 0.56, $p < 0.0001$) and was found to have higher dispersion than the PC data (Fig. 2g). No differences were observed in the waveform of recorded spikes during anodal nor cathodal tDCS with respect to control condition (data not shown).

These results allow us to conclude that not only PCs but also non-PCs implicated in the spatio-temporal response of PCs are modulated during tDCS in a heterogeneous way.

Fig. 3 tDCS modulation of non-PC activity in crus I/II of awake mice. a, b Recording of spontaneous firing activity of two non-PCs. **c, e** Z-score-transformed average PSTH (bin size: 0.1 s) of firing rate for the 2 neurons in a,b, before, during and after anodal (red trace) or cathodal (blue trace) tDCS. **d, f** Statistical comparison of the firing rate of the 2 neurons in a,b, measured in 5 s windows before, during and after tDCS (RM-ANOVA or Friedman tests, p < 0.05). Error bars represent SEM. *p < 0.05; **p < 0.01; ***p < 0.001. **g** Modulation of firing rate of individual neurons (circles) during anodal (red) and cathodal (blue) tDCS. Filled circles represent statistically significant modulation during tDCS (n = 50, RM-ANOVA or Friedman tests, p < 0.05). **h** Schematic representation of the recording sites and active electrode (gray bar) location during tDCS.

Purkinje cell orientation explains heterogeneous tDCS modulation in anesthetized mice.

Given the large heterogeneity observed in the responses of the recorded cerebellar neurons to tDCS and considering the anatomical complexity of the highly convoluted cerebellar cortex, we wondered if the somatodendritic axis orientation of the cerebellar neurons could partially explain this variability, as suggested by previous *in vitro* studies for neurons in other brain regions[18,19,35]. For this purpose, we decided 1) to record in the cerebellar vermis region which is characterized by having PCs oppositely oriented in adjacent cortical layers (Fig. 4a,b), and 2) label some of the recorded PC with neurobiotin after electrophysiological characterization

to examine their orientation.

A total of 56 neurons (31 identified PCs and 25 non-PCs) were recorded in the vermis of 27 anesthetized mice (Fig. 4a,b) during anodal and cathodal tDCS (±200 µA). The impact of anodal and cathodal tDCS on the firing rate showed significant differences (filled circles) in 27 out of 31 PCs (Fig. 4c) and in 17 out 25 recorded non-PCs (Fig. 4d) (n = 31 PCs and 25 non-PCs, RM-ANOVA or Friedman tests, p < 0.05). The effects of tDCS on PC firing rate adjusts to a linear regression (slope -0.57, R = 0.7, p < 0.0001) (Fig. 4c), whereas the effects were scattered on non-PCs (slope -0.19, R = 0.35, p < 0.0875) (Fig. 4d).

Fig. 4 tDCS modulation of PC and non-PC activity in the vermis of anesthetized mice. a Schematic representation of the recording sites and active electrode (gray bar) location during tDCS. **b** A representative coronal section of the vermis immunofluorescently stained with Calbindin (green). The magnification inset highlights the distinct orientation of PCs in different layers, which is indicated by the drawings of PC somas and dendrites (with dendrites always extending into the molecular layer, shown in green). **c,d** Modulation of SS firing rate of individual PCs (c) and firing rate of individual non-PCs (d) during anodal (red) and cathodal (blue) tDCS over cerebellar vermis. Filled circles represent statistically significant modulation during tDCS (n = 31 PCs and 25 non-PCs, RM-ANOVA or Friedman tests, p < 0.05). GCL: granular cell layer, ML: molecular layer, PCL: Purkinje cell layer.

This result suggests that in the vermis there is a consistent polarity-dependent modulation for PCs, where anodal and cathodal modulate firing rate in opposite ways. However, this

effect is not observed in the non-PC group. The findings remained consistent even when we tested the same neurons with greater (±300 μA) and lesser (±100 μA) current intensities (Supplementary Fig. 3). From these experiments we can conclude that 1) tDCS in vermis of anesthetized mice modulates PCs and non-PCs in a heterogeneous way, 2) tDCS in vermis modulates more PCs than in Crus I-II and 3) most PCs in vermis modulate in opposite directions for anodal vs cathodal tDCS.

To test if the opposite modulation of PCs by anodal and cathodal stimulation could be related to the opposite somatodendritic orientation of these neurons with respect to the electric field, a total of 8 recorded PCs in 8 animals were successfully stained with neurobiotin using juxtacellular microinjection in the anesthetized mice either in crus I-II or vermis (see Methods). Labeled PCs were reconstructed with confocal microscopy and the deviation of the somatodendritic axis from the imaginary line perpendicular to the skull surface under the active electrode was calculated (Fig. 5a; θ angle). Figure 5a-d shows representative PCs with different θ angle together with Z-score-transformed average PSTH (bin size: 0.1 s) of the spiking activity and statistical comparison of the firing rate before, during and after anodal (red) or cathodal (blue) tDCS (Friedman test, p < 0.05). We observed that when θ was close to 0º (the somatodendritic axis was pointing toward the active electrode, Fig. 5a) anodal tDCS tended to increase the firing rate and cathodal reliably decreased it, whereas when θ was close to 180º (the somatodendritic axis was pointing away from the electrode) (Fig. 5b), the opposite modulation was observed, with anodal decreasing and cathodal increasing the firing rate. However, when θ value was close to 90º (Fig. 5c) or to 270º (Fig. 5d) (the axis was more perpendicular to the electric field) the modulatory effect was absent. Figure 5e summarizes the relationship between the θ angle and tDCS modulation of all labeled neurons. The figure represents the normalized firing rate modulation ([firing rate during tDCS/firing rate before tDCS]*100; length of the arrow) and θ angle value for each recorded neuron during anodal (Fig. 5e, at left) and cathodal (Fig. 5e, at right) tDCS. Figure 5f represents the relationship between the normalized firing rate modulation of each individual neuron with respect to its corresponding θ angle for anodal (red line) or cathodal (blue line) tDCS. The impact of tDCS in the firing rate of PC was higher for those neurons with θ values close to 0º and 180º for anodal and cathodal currents acting in an opposite polarity-dependent manner. These results corroborate that the somatodendritic axis orientation plays a critical role in explaining tDCS heterogeneous modulation of individual PCs in anesthetized mice.

Figure 5

Fig. 5 Relationship between tDCS-driven modulation of PC firing rate and somatodendritic axis orientation in anesthetized mice. a-d (Left) Confocal images of labeled neurons with different somatodendritic angles relative to the electric field (dotted white vertical line), (Right) z-score of their firing rate modulation during anodal (red) or cathodal (blue) tDCS and statistical analysis (RM-ANOVA or Friedman tests, $p < 0.05$). Error bars represent SEM. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$. **e** Relationship between firing rate modulation and somatodendritic angle for all juxtacellularly-labelled PCs (n = 8). Arrow length represents firing rate modulation during anodal (red arrows, at left) or cathodal (blue arrows, at right) tDCS at 200 μA, relative to the firing rate during control condition (represented by 100% solid circle). **f** Average change in firing rate during anodal (red) and cathodal (blue) tDCS for individual PCs with different somatodendritic orientations.

Purkinje cell orientation determines polarity of tDCS-driving firing rate modulation in awake mice.

To examine whether the effects of PC orientation that we observed in anesthetized mice can be extended to the more clinically-relevant awake state, we took advantage of Neuropixels technology and prepared 6 additional mice for chronic multiunitary neuronal recording during simultaneous tDCS. Neuropixels probes were coated with lipophilic dye (DiI) for subsequent histological reconstruction of the recording track (see Methods). A total of 9 Purkinje cells were selected for analysis because they were particularly stable and well isolated during the entire tDCS session, and because they were unequivocally identified as Purkinje cells by confirming that they fired simple and complex spikes. Figure 6a illustrates probe location marked with DiI in the cerebellar vermis of a representative animal where two different PCs with the somatodendritic axis pointing toward and away from the active electrode recorded at Ch#55 and Ch#42, respectively (Fig. 6b); note dendrites always extend into the molecular layer which is shown in green). The Z-score-transformed average PSTH (bin size: 0.1 s for SS) of the SS activity before, during and after anodal (red trace) and cathodal (blue trace) tDCS is shown for these two simultaneously recorded PCs in Figure 6c. Remarkably, we found that the two PCs, which have oppositely oriented dendrites, exhibit opposite effects in presence of anodal or cathodal tDCS (Fig. 6c). The same relationship was found for all the other PCs (Fig. 6d,e and Supplementary Fig. 4), where PCs in which the dendrites pointed toward the active tDCS electrode had increased firing rate during anodal tDCS and reduced firing rate during cathodal tDCS (Fig. 6d,e, red triangles pointing up) while PCs with dendrites pointing away from the active tDCS electrode had the opposite modulation (Fig. 6d,e, blue triangles pointing down). This experiment corroborates in the awake animal and in simultaneously recorded PCs that 1) a given tDCS polarity (either anodal or cathodal) can modulate individual PC neurons in opposite ways at the same time, increasing and decreasing their firing rates, and 2) the somatodendritic axis orientation of PCs is a key factor in determining the tDCS-driven modulation of firing rate.

Fig. 6 Impact of tDCS on PCs with opposite axodendritic orientations simultaneously recorded in the awake mice. a Probe location marked with DiI in the cerebellar vermis stained with Hoechst 33342 dye. **b** Magnification of square area in "*a*" showing the location of two oppositely oriented PCs recorded at Ch#55 and Ch#42. The orientation of the PCs in each of the 2 layers is indicated with drawings of PC soma and dendrites (yellow), in which the soma appear at the interface between the granule cell layer (GCL, shown in blue) and the molecular layer (ML, shown in green). **c** Z-score-transformed average PSTH (bin size: 0.1 s for SS) of the SS activity before, during and after anodal and cathodal tDCS pulses for each of the 2 simultaneously recorded PCs shown in b. **d** Anatomical localization of the different PCs recorded. The inset shows the recorded places marked with DiI (red) and stained with anti-Calbindin antibody (green, molecular layer) and Hoechst 33342 (blue, granule layer) (scale bar: 300 μm). PCs in which dendrites are pointed toward or away from the active tDCS electrode are denoted with triangles pointing upward or downward respectively. The color of the triangle indicates whether the modulation was an increase (red) or a decrease (blue) in firing rate. **e** Modulation of SS firing rate of individual PCs during anodal and cathodal tDCS. Filled symbols represent statistically significant modulation during tDCS, with the meaning of color and shape as in d (n = 9, RM-ANOVA or Friedman tests, p < 0.05). ML: molecular layer, PCL: Purkinje cell layer, GCL: granular cell layer.

Discussion

In the present investigation, we use *in vivo* unitary extracellular recordings to show tDCS capability to modulate neuronal activity in the cerebellar cortex of awake mice. We find that both anodal and cathodal tDCS modulates the activity of many cerebellar neurons, but the effects are extremely heterogeneous. For Purkinje cells (PCs), which are the neurons responsible for sending the output of the entire cerebellar cortex, we show that the diverse effects observed during tDCS are largely explained by the somatodendritic axis orientation with respect to the active electrode. This is shown by matching *in vivo* electrophysiological recordings with neurobiotin-labeling of individual PC neurons in anesthetized mice, as well as Neuropixels high-density recordings in awake mice under tDCS. The firing rate of PCs whose dendrites are pointed toward the stimulation electrode increases (or decreases) during anodal (or cathodal) tDCS, respectively, while the modulation of firing rate is exactly the opposite for PCs whose dendrites are pointed away from the stimulation electrode.

We observe a robust firing rate modulation during tDCS, with over 64% of the recorded neurons being modulated. The neuromodulatory effects of tDCS are highly dependent on the magnitude of the electric fields applied to the skull and the distance to the site of action in the brain[36]. As expected from cerebellar modelling studies[37,38] and previously observed *in vitro*[39] and *in vivo*[21], we find that the current flow in the cerebellum is largely uniform in direction, with the highest electric field values observed in the first millimeter of the cortex (64.8, 6.9 and 0.97 V/m for 200, 20 and 2 µA, respectively), and that the electric field strength decays logarithmically with distance. This electric field magnitude is directly related to the intensity of the electric current and the size of the electrodes, which determine the current density, as well as the shunting effect caused by the tissue between the electrodes and the brain[40,41]. Approximately 60% of the applied current is shunted by the skin, while around 20% is attenuated by the skull[28,42]. Based on previous studies that rely on computational analysis with realistic head models to estimate how electric currents propagate across the brain in different tDCS montages[37,43,44], we estimate that the electric fields applied in this study are generally higher than those typically used in human protocols (~1-1.5 V/m), although they are similar to those used in other animal studies[13,45,46]. Importantly, we observe clear firing rate modulation of PCs and non-PCs at depths of 2.3 mm and tDCS intensity of 100 µA, where the measured electric field is as low as 5.9 V/m.

The tDCS-driven modulation of firing rate is polarity-dependent for the majority of cerebellar neurons, with anodal and cathodal tDCS modulating the firing rate in opposite directions. Similar polarity-dependent modulation has been reported in previous *in vivo* studies in anaesthetized animals during application of tDCS over cerebral[28,47] and cerebellar[21] cortices. The mechanism behind the firing rate changes observed during tDCS is likely due to a polarizing effect, where a slight alteration in the resting membrane potential of neurons (depolarization or hyperpolarization, depending on tDCS polarity) will lead to opposite changes in postsynaptic somatic spiking, as demonstrated *in vitro*[17,48] and *in vivo*[28]. Consistent with this explanation, we observed that the firing rate is modulated over time in accordance with transcranial current dynamics (i.e., variable in ramp-in and ramp-out periods and stable during the 5 or 10 s at maximum intensity), except for a rebound effect observed after current termination (also observed by Asan et al.[21]). Nonetheless, we cannot rule out the possibility of indirect synaptic effects. Indeed, the electric field gradient imposed by tDCS could directly modulate a specific neuron firing rate by increasing (or decreasing) its pre-synaptic activity, i.e. by modulating the firing rate of other neurons that synapse onto it. Indeed, these synaptic changes could explain the rebound effect observed after tDCS termination. The synapses involved in the modulation of firing rate may undergo a short-term plasticity process[48–51], which can continue to affect the firing rate even after the external currents have been turned off and no polarization is exerted on the neuron.

Previous *in vitro, ex vivo* and modelling studies have highlighted the importance of various neuronal features that underly tDCS effects, including the orientation of somatodendritic axis with respect to the electrical field[38], the neuronal morphology[18,39] or the axonal orientation[19]. In this study, we demonstrate that in both anesthetized and awake animals, the orientation of the PC axodendritic axis with respect to the electric field induced by tDCS explains the observed polarity-dependent heterogeneity in firing rate modulation. The importance of PC axodendritic orientation in determining the effect of tDCS on firing rate modulation is further highlighted by our observation that pre-synaptic non-PC neurons providing inputs to PCs modulate their activity in a very heterogeneous way. In other words, our findings reveal that regardless of how tDCS impacts the activity of pre-synaptic inputs to PCs, the tDCS-driven firing rate modulation of each individual PC can be predicted by simply taking into account the orientation of the PC's dendrites relative to the electric field.

Our findings may provide some biological insight as to why the effects of cerebellar tDCS on motor control and learning have been difficult to replicate and are often reported to be fickle and unreliable[8,9]. First, we find that in most of the analyzed PCs, tDCS induces a highly variable and inconsistent modulation of two parameters that play a key role in cerebellar learning function[52]: the firing rate of PC complex spikes (CS) and the duration of the simple spike (SS) silence following the CS. Second, we show that in the same animal and at the same time, tDCS can drive completely opposite changes in the firing rate of PCs with opposite somatodendritic orientations. Groups of PCs in different layers and areas of the cerebellar cortex are linked together into functional modules, based on their projecting to the same subdivision of the deep cerebellar nuclei[53]. As a result, the final macroscopic effect of cerebellar tDCS will depend on the net modulation of all the PCs in each module within the stimulated folia, which will be strongly influenced by their (likely heterogeneous) orientation. In the future, new technological advances in high-definition tDCS may be combined with flexible control of electric field direction to target PCs in specific modules and boost reliability.

In conclusion, we found that tDCS modulates the firing rate of PCs in mice in a polarity-dependent manner, and this modulation is highly dependent on the orientation of the PC somatodendritic axis relative to the electric field. Our findings emphasize the importance of considering the neuronal orientation and morphology of target neurons when applying transcranial stimulation, at least in the cerebellum. Taking into account these neuronal features is crucial for increasing the predictive power of computational models and optimizing the desired effects of tDCS in basic and clinical human applications.

Author contributions

C.A.S-L and J.M-R conceived the original idea and designed the experiments. C.A.S-L, G.S-G.C, M.F and J.M-R performed the experiments and the data analysis. C.A.S-L, J.M-R wrote the paper. A.S-L and J.F.M assisted in the experimental design and in the interpretation of the results. All the authors contributed to the final edition of the manuscript.

References

1. Nitsche, M. A. & Paulus, W. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *J Physiol* **527 Pt 3**, 633–639 (2000).

2. Woods, A. J. *et al.* A technical guide to tDCS, and related non-invasive brain stimulation tools. *Clin Neurophysiol* **127**, 1031–1048 (2016).

3. Stagg, C. J., Antal, A. & Nitsche, M. A. Physiology of Transcranial Direct Current Stimulation. *J ECT* **34**, 144–152 (2018).

4. Manto, M. *et al.* Consensus Paper: Novel Directions and Next Steps of Non-invasive Brain Stimulation of the Cerebellum in Health and Disease. *Cerebellum* **21**, 1092–1122 (2022).

5. Grimaldi, G. *et al.* Cerebellar Transcranial Direct Current Stimulation (ctDCS): A Novel Approach to Understanding Cerebellar Function in Health and Disease. *Neuroscientist* **22**, 83–97 (2016).

6. Antal, A., Keeser, D., Priori, A., Padberg, F. & Nitsche, M. A. Conceptual and Procedural Shortcomings of the Systematic Review 'Evidence That Transcranial Direct Current Stimulation (tDCS) Generates Little-to-no Reliable Neurophysiologic Effect Beyond MEP Amplitude Modulation in Healthy Human Subjects: A Systematic Review' by Horvath and Co-workers. *Brain Stimul* **8**, 846–849 (2015).

7. Horvath, J. C., Forte, J. D. & Carter, O. Evidence that transcranial direct current stimulation (tDCS) generates little-to-no reliable neurophysiologic effect beyond MEP amplitude modulation in healthy human subjects: A systematic review. *Neuropsychologia* **66**, 213–236 (2015).

8. Jalali, R., Miall, R. C. & Galea, J. M. No consistent effect of cerebellar transcranial direct current stimulation on visuomotor adaptation. *J Neurophysiol* **118**, 655–665 (2017).

9. Beyer, L., Batsikadze, G., Timmann, D. & Gerwig, M. Cerebellar tDCS Effects on Conditioned Eyeblinks using Different Electrode Placements and Stimulation Protocols. *Front Hum Neurosci* **11**, 23 (2017).

10. Horvath, J. C., Carter, O. & Forte, J. D. No significant effect of transcranial direct current stimulation (tDCS) found on simple motor reaction time comparing 15 different simulation protocols. *Neuropsychologia* **91**, 544–552 (2016).

11. Liu, A. *et al.* Immediate neurophysiological effects of transcranial electrical stimulation. *Nature Communications* **9**, 5092 (2018).

12. Polanía, R., Nitsche, M. A. & Ruff, C. C. Studying and modifying brain function with non-invasive brain stimulation. *Nature Neuroscience* **21**, 174–187 (2018).

13. Jackson, M. P. *et al.* Animal models of transcranial direct current stimulation: Methods and mechanisms. *Clin Neurophysiol* **127**, 3425–3454 (2016).

14. Liebetanz, D. *et al.* Safety limits of cathodal transcranial direct current stimulation in rats. *Clin Neurophysiol* **120**, 1161–1167 (2009).

15. Jackson, M. P. *et al.* Safety parameter considerations of anodal transcranial Direct Current Stimulation in rats. *Brain Behav Immun* **64**, 152–161 (2017).

16. Sánchez-León, C. A., Ammann, C., Medina, J. F. & Márquez-Ruiz, J. Using animal models to improve the design and application of transcranial electrical stimulation in humans. *Curr Behav Neurosci Rep* **5**, 125–135 (2018).

17. Bikson, M. *et al.* Effects of uniform extracellular DC electric fields on excitability in rat hippocampal slices in vitro. *J Physiol* **557**, 175–190 (2004).

18. Radman, T., Ramos, R. L., Brumberg, J. C. & Bikson, M. Role of cortical cell type and morphology in subthreshold and suprathreshold uniform electric field stimulation in vitro. *Brain Stimul* **2**, 215–228, 228.e1–3 (2009).

19. Kabakov, A. Y., Muller, P. A., Pascual-Leone, A., Jensen, F. E. & Rotenberg, A. Contribution of axonal orientation to pathway-dependent modulation of excitatory transmission by direct current stimulation in isolated rat hippocampus. *J Neurophysiol* **107**, 1881–1889 (2012).

20. Huang, W. A. *et al.* Transcranial alternating current stimulation entrains alpha oscillations by preferential phase synchronization of fast-spiking cortical neurons to stimulation waveform. *Nat Commun* **12**, 3151 (2021).

21. Asan, A. S., Lang, E. J. & Sahin, M. Entrainment of cerebellar purkinje cells with directional AC electric fields in anesthetized rats. *Brain Stimul* **13**, 1548–1558 (2020).

22. Sánchez-León, C. A. *et al.* Immediate and after effects of transcranial direct-current stimulation in the mouse primary somatosensory cortex. *Sci Rep* **11**, 3123 (2021).

23. Krause, M. R. *et al.* Transcranial Direct Current Stimulation Facilitates Associative Learning and Alters Functional Connectivity in the Primate Brain. *Curr Biol* **27**, 3086-3096.e3 (2017).

24. Márquez-Ruiz, J. *et al.* Transcranial direct-current stimulation modulates synaptic mechanisms involved in associative learning in behaving rabbits. *Proc Natl Acad Sci U S A* **109**, 6710–6715 (2012).

25. Monai, H. *et al.* Calcium imaging reveals glial involvement in transcranial direct current stimulation-induced plasticity in mouse brain. *Nat Commun* **7**, 11100 (2016).

26. Mishima, T. *et al.* Transcranial Direct Current Stimulation (tDCS) Induces Adrenergic Receptor-Dependent Microglial Morphological Changes in Mice. *eNeuro* **6**, ENEURO.0204-19.2019 (2019).

27. Opitz, A. *et al.* Spatiotemporal structure of intracranial electric fields induced by transcranial electric stimulation in humans and nonhuman primates. *Sci Rep* **6**, 31236 (2016).

28. Vöröslakos, M. *et al.* Direct effects of transcranial electric stimulation on brain circuits in rats and humans. *Nat Commun* **9**, 483 (2018).

29. Grossman, N. *et al.* Noninvasive Deep Brain Stimulation via Temporally Interfering Electric Fields. *Cell* **169**, 1029-1041.e16 (2017).

30. Thach, W. T. Somatosensory receptive fields of single units in cat cerebellar cortex. *J Neurophysiol* **30**, 675–696 (1967).

31. Ito, M. *The Cerebellum and Neural Control*. (Raven Press, 1984).

32. Paxinos, G. & Franklin, K. B. J. *The Mouse Brain in Stereotaxic Coordinates*. (Academic Press, San Diego, 2013).

33. Pinault, D. A novel single-cell staining procedure performed in vivo under electrophysiological control: morpho-functional features of juxtacellularly labeled thalamic cells and other central neurons with biocytin or Neurobiotin. *J Neurosci Methods* **65**, 113–136 (1996).

34. Prestori, F., Mapelli, L. & D'Angelo, E. Diverse Neuron Properties and Complex Network Dynamics in the Cerebellar Cortical Inhibitory Circuit. *Front Mol Neurosci* **12**, 267 (2019).

35. Bikson, M., Reato, D. & Rahman, A. Cellular and network effects of transcranial direct current stimulation: insights from animal models and brain slice. *Transcranial brain stimulation* **1**, 456 (2012).

36. Opitz, A., Paulus, W., Will, S., Antunes, A. & Thielscher, A. Determinants of the electric field during transcranial direct current stimulation. *NeuroImage* **109**, 140–150 (2015).

37. Parazzini, M. *et al.* Modelling the electric field and the current density generated by cerebellar transcranial DC stimulation in humans. *Clin Neurophysiol* **125**, 577–584 (2014).

38. Rahman, A., Toshev, P. K. & Bikson, M. Polarizing cerebellar neurons with transcranial Direct Current Stimulation. *Clin Neurophysiol* **125**, 435–438 (2014).

39. Chan, C. Y. & Nicholson, C. Modulation by applied electric fields of Purkinje and stellate cell activity in the isolated turtle cerebellum. *J Physiol* **371**, 89–114 (1986).

40. Opitz, A., Falchier, A., Linn, G. S., Milham, M. P. & Schroeder, C. E. Limitations of ex vivo measurements for in vivo neuroscience. *Proc Natl Acad Sci U S A* **114**, 5243–5246 (2017).

41. Gomez-Tames, J. *et al.* Group-level and functional-region analysis of electric-field shape during cerebellar transcranial direct current stimulation with different electrode montages. *J Neural Eng* **16**, 036001 (2019).

42. Liu, A. *et al.* Immediate neurophysiological effects of transcranial electrical stimulation. *Nat Commun* **9**, 5092 (2018).

43. Ruffini, G., Fox, M. D., Ripolles, O., Miranda, P. C. & Pascual-Leone, A. Optimization of multifocal transcranial current stimulation for weighted cortical pattern targeting from realistic modeling of electric fields. *Neuroimage* **89**, 216–225 (2014).

44. Huang, Y., Datta, A., Bikson, M. & Parra, L. C. Realistic volumetric-approach to simulate transcranial electric stimulation-ROAST-a fully automated open-source pipeline. *J Neural Eng* **16**, 056006 (2019).

45. Alekseichuk, I., Mantell, K., Shirinpour, S. & Opitz, A. Comparative modeling of transcranial magnetic and electric stimulation in mouse, monkey, and human. *Neuroimage* **194**, 136–148 (2019).

46. Asan, A. S., Gok, S. & Sahin, M. Electrical fields induced inside the rat brain with skin, skull, and dural placements of the current injection electrode. *PLoS One* **14**, e0203727 (2019).

47. Bindman, L. J., Lippold, O. C. & Redfearn, J. W. THE ACTION OF BRIEF POLARIZING CURRENTS ON THE CEREBRAL CORTEX OF THE RAT (1) DURING CURRENT FLOW AND (2) IN THE PRODUCTION OF LONG-LASTING AFTER-EFFECTS. *J Physiol* **172**, 369–382 (1964).

48. Farahani, F., Kronberg, G., FallahRad, M., Oviedo, H. V. & Parra, L. C. Effects of direct current stimulation on synaptic plasticity in a single neuron. *Brain Stimul* **14**, 588–597 (2021).

49. Cirillo, G. *et al.* Neurobiological after-effects of non-invasive brain stimulation. *Brain Stimul* **10**, 1–18 (2017).

50. Kronberg, G., Bridi, M., Abel, T., Bikson, M. & Parra, L. C. Direct Current Stimulation Modulates LTP and LTD: Activity Dependence and Dendritic Effects. *Brain Stimul* **10**, 51–58 (2017).

51. Kronberg, G., Rahman, A., Sharma, M., Bikson, M. & Parra, L. C. Direct current stimulation boosts hebbian plasticity in vitro. *Brain Stimul* **13**, 287–301 (2020).

52. Medina, J. F. & Lisberger, S. G. Links from complex spikes to local plasticity and motor learning in the cerebellum of awake-behaving monkeys. *Nat Neurosci* **11**, 1185–1192 (2008).

53. Apps, R. & Hawkes, R. Cerebellar cortical organization: a one-map hypothesis. *Nat Rev Neurosci* **10**, 670–681 (2009).